Homotopy invariants of covers and KKM type lemmas

Oleg R. Musin^{*}

Abstract

Given any (open or closed) cover of a space T we associate certain homotopy classes of maps from T to n-spheres. These homotopy invariants can then be considered as obstructions for extending covers of a subspace $A \subset X$ to a cover of all of X. We are using these obstructions to obtain generalizations of the classic KKM (Knaster-Kuratowski-Mazurkiewicz) and Sperner lemmas. In particular, we show that in the case when A is a k-sphere and X is a (k + 1)-disk there exist KKM type lemmas for covers by n + 2 sets if and only if the homotopy group $\pi_k(\mathbb{S}^n) \neq 0$.

Keywords: KKM lemma, Sperner lemma, homotopy class, degree of mappings

Throughout this paper we will consider only normal topological spaces, all simplicial complexes will be finite, all manifolds with be both compact and PL, \mathbb{S}^n will denote the *n*-dimensional unit sphere, and \mathbb{B}^n will denote the *n*-dimensional unit disk. We shall denote the set of homotopy classes of continuous maps from X to Y by [X, Y].

1 Homotopy invariants of covers

First we consider labelings (colorings) of simplicial complexes. Denote by Vert(K) the vertex set of a simplicial complex K. (It is also referred to as the 0-skeleton, K^0 .) Let

$$L: \operatorname{Vert}(K) \to \{0, 1, \dots, m\}$$

be a labeling of vertices of K.

Denote by Δ^m an *m*-dimensional simplex with vertices v_0, \ldots, v_m . Let

 $f_L(u) := v_\ell$, where $u \in \operatorname{Vert}(K)$, and $\ell = L(u)$.

Since f_L is defined for all of the vertices of K, it induces a simplicial mapping $f_L : |K| \to |\Delta^m|$. This map is unique up to homeomorphism.

Note that if any simplex in K has at most m distinct labels, then f_L is a map from |K| to $\partial |\Delta^m| \cong \mathbb{S}^{m-1}$.

^{*}The research was carried out at the IITP RAS at the expense of the Russian Foundation for Sciences (project 14-50-00150).

Definition 1.1. For a simplicial complex K and a labeling L: Vert $(K) \rightarrow \{0, 1, ..., m\}$ such that K has no simplices with m + 1 distinct labels we denote by [L] the homotopy class $[f_L]$ in $[|K|, \mathbb{S}^{m-1}]$.

Example 1.1. Let K be a triangulation of \mathbb{S}^k and $L : \operatorname{Vert}(K) \to \{0, 1, \dots, m\}$ be a labeling such that K has no simplices with m + 1 distinct labels. Then $[L] \in \pi_k(\mathbb{S}^{m-1})$.

In the case k = m - 1 we have $\pi_k(\mathbb{S}^{m-1}) = \mathbb{Z}$ and

$$[L] = \deg(f_L) \in \mathbb{Z}.$$

(Here by $\deg(f)$ we denote the degree of a continuous map f from \mathbb{S}^n to itself.)

For instance, let L be a Sperner labeling of a triangulation K of $\partial \Delta^m = u_0 u_1 \dots u_m$. The rules of this labeling are:

(i) The vertices of Δ^m are colored with different colors, i. e. $L(u_i) = i$ for $0 \le i \le m$.

(ii) Vertices of K located on any n-dimensional subface of the large simplex $u_{i_0}u_{i_1}\ldots u_{i_n}$ are colored only with the colors i_0, i_1, \ldots, i_n .

Then $[L] = \deg(f_L) = 1$ in $[\mathbb{S}^{m-1}, \mathbb{S}^{m-1}] = \mathbb{Z}.$

Example 1.2. Madahar and Sarkaria [8] considered a simplicial map $\tau : \mathbb{S}_{12}^3 \to \mathbb{S}_4^2$ from a 12-vertex 3-sphere \mathbb{S}_{12}^3 onto the 4-vertex 2-sphere \mathbb{S}_4^2 (tetrahedron) with vertices $v_0v_1v_2v_3$. Actually, τ is a minimal vertices simplicial map of the Hopf invariant one.

For $u \in \operatorname{Vert}(\mathbb{S}^3_{12})$, let $L_{\tau}(u) := i$, where $\tau(u) = v_i$. Then $[L_{\tau}] = 1 \in \pi_3(\mathbb{S}^2) = \mathbb{Z}$.

Let K be a simplicial complex. Denote by St(u) the open star of a vertex $u \in Vert(K)$. In other words, St(u) is $|S| \setminus |B|$, where S is the set of all simplices in K that contain u, and B is the set of all simplices in S that does contain u.

Let

$$L: \operatorname{Vert}(K) \to \{0, 1, \dots, m\}$$

be a labeling of vertices of K. There is a natural open cover of |K|

$$\mathcal{U}_L(K) = \{U_0(K), \dots, U_m(K)\},\$$

where

$$U_{\ell}(K) := \bigcup_{u \in W_{\ell}} \operatorname{St}(u), \text{ and } W_{\ell} := \{ u \in \operatorname{Vert}(K) : L(u) = \ell \}.$$

So with any labeling L we associate a cover $\mathcal{U}_L(K)$. Now we extend Definition 1.1 to covers.

Let $\mathcal{U} = \{U_0, \ldots, U_m\}$ be an open finite cover of a space T. If $N(\mathcal{U})$ is its nerve, then there is a one-to-one correspondence between canonical maps $c: T \to |N(\mathcal{U})|$ and partitions of unity Φ subordinate to \mathcal{U} . Moreover, any two canonical maps $T \to |N(\mathcal{U})|$ are homotopic.

Since the nerve $N(\mathcal{U})$ is a subcomplex of the simplex Δ^m , we have an embedding α : $|N(\mathcal{U})| \to |\Delta^m|$. In the case when the intersection of all of the U_i is empty, i. e. $N(\mathcal{U})$ does not contain an *m*-cell, we have

$$\alpha: |N(\mathcal{U})| \to \partial |\Delta^m| \cong \mathbb{S}^{m-1}.$$

If

$$\rho_{\mathcal{U},c} := \alpha \circ c,$$

then a homotopy class $[\rho_{\mathcal{U},c}]$ in $[T, \mathbb{S}^{m-1}]$ does not depend on the canonical map $c: T \to |N(\mathcal{U})|$.

Definition 1.2. Let $\mathcal{U} = \{U_0, \ldots, U_m\}$ be an open finite cover of a space T such that the intersection of all of the U_i is empty. Denote by $[\mathcal{U}]$ the homotopy class $[\rho_{\mathcal{U},c}]$ in $[T, \mathbb{S}^{m-1}]$.

Remark. It is clear that

$$[\mathcal{U}_L(K)] = [L] \text{ in } [|K|, \mathbb{S}^{m-1}].$$

Theorem 1.1. Let T be a space and h be a homotopy class in $[T, \mathbb{S}^{m-1}]$. Then there is an open cover $\mathcal{U} = \{U_0, \ldots, U_m\}$ such that $[\mathcal{U}] = h$.

If T is a simplicial complex, then there is a triangulation K of T and a labeling L : $\operatorname{Vert}(K) \to \{0, 1, \dots, m\}$ with $[\mathcal{U}_L(K)] = h$.

Proof. Let Λ : Vert $(\Delta^m) \to \{0, 1, \ldots, m\}$ be a labeling of Δ^m with vertices v_0, \ldots, v_m such that $\Lambda(v_\ell) = \ell$ for all ℓ . Then we have a cover $\mathcal{U}_{\Lambda}(\Delta^m)$.

Let $f: T \to \mathbb{S}^{m-1}$ be a continuous map with [f] = h and

$$U_{\ell} := f^{-1}(U_{\ell}(\Delta^m)), \ \ell = 0, \dots, m.$$

It is easy to see that $[\mathcal{U}] = h$.

If T is a simplicial complex, then by the simplicial approximation theorem there is a simplicial subdivision (triangulation) K and a simplicial map $g: K \to \Delta^m$ such that g is homotopic to f. For all $v \in \operatorname{Vert}(K)$, let

$$L(v) := \Lambda(g(v)).$$

Then $[\mathcal{U}_L(K)] = h.$

Let us define the class $[\mathcal{U}]$ more explicitly. Let $\Phi = \{\varphi_0, \ldots, \varphi_m\}$ be a partition of unity subordinate to \mathcal{U} and for all $x \in T$

$$\rho_{\mathcal{U},\Phi}(x) := \sum_{i=0}^{m} \varphi_i(x) v_i,$$

where v_0, \ldots, v_m are vertices of an *m*-dimensional simplex *V* considered as vectors in \mathbb{R}^m . Then $\rho_{\mathcal{U},\Phi}$ is a continuous map from *T* to $\partial V = \mathbb{S}^{m-1}$. It is clear that

$$[\rho_{\mathcal{U},\Phi}] = [\mathcal{U}] \text{ in } [T, \mathbb{S}^{m-1}].$$

Now we extend this definition. Let $V := \{v_0, \ldots, v_m\}$ be any set of points (vectors) in \mathbb{R}^{n+1} . As above

$$\rho_{\mathcal{U},\Phi,V}(x) := \sum_{i=0}^{m} \varphi_i(x) v_i.$$

Suppose a point $p \in \mathbb{R}^{n+1}$ lies outside of the image $\rho_{\mathcal{U},\Phi,V}(T)$. For all $x \in T$, let

$$f_{\mathcal{U},\Phi,V,p}(x) := \frac{\rho_{\mathcal{U},\Phi,V}(x) - p}{||\rho_{\mathcal{U},\Phi,V}(x) - p||}.$$

Then $f_{\mathcal{U},\Phi,V,p}$ is a continuous map from T to \mathbb{S}^n .

Lemma 1.1. For given \mathcal{U}, V and p any two partitions of unity subordinate to \mathcal{U} define the same homotopy class $[f_{\mathcal{U},V,p}]$ in $[T, \mathbb{S}^n]$.

Proof. A linear homotopy $\Theta(t) = (1 - t)\Phi + t\Psi$ of two partitions of unity Φ and Ψ induces a homotopy between the maps $f_{\mathcal{U},\Phi,V,p}$ and $f_{\mathcal{U},\Psi,V,p}$.

Lemma 1.2. For any two partitions of unity Φ and Ψ subordinate to \mathcal{U} , the image $\rho_{\mathcal{U},\Phi,V}(T)$ coincides with the image $\rho_{\mathcal{U},\Psi,V}(T)$ in \mathbb{R}^{n+1} .

Proof. Consider the nerve $N(\mathcal{U})$ with vertices U_i . If we set $g(U_i) := v_i$, then we have a piecewise linear map $g : |N(\mathcal{U})| \to H$, where $H := \operatorname{conv}(V)$ is the convex hull of V in \mathbb{R}^{n+1} . Then for any partition of unity Φ we have $\rho_{\mathcal{U},\phi,V} := g \circ c$, where $c : T \to |N(\mathcal{U})|$ is the canonical map corresponding to Φ . Thus, $\rho_{\mathcal{U},\Phi,V}(T) = g(|N(\mathcal{U})|)$ and does not depend on Φ . \Box

Notation: $P_{\mathcal{U},V}(T) := \mathbb{R}^{n+1} \setminus \rho_{\mathcal{U},\Phi,V}(T).$

Note that the map $f_{\mathcal{U},\Phi,V,p}: T \to \mathbb{S}^n$ is well defined only if $p \in P_{\mathcal{U},V}(T)$.

Lemma 1.3. Let points p and q lie in the same connected component Q of $P_{\mathcal{U},V}(T)$. Then $[f_{\mathcal{U},V,p}] = [f_{\mathcal{U},V,q}]$ in $[T, \mathbb{S}^n]$.

Proof. Let s(t) be a path in Q connecting the points p and q. Then s induces a homotopy between the maps $f_{\mathcal{U},\Phi,V,p}$ and $f_{\mathcal{U},\Phi,V,q}$.

Definition 1.3. For a cover $\mathcal{U} = \{U_1, \ldots, U_m\}$ of a space T, a set V of m points in \mathbb{R}^{n+1} , and $p \in P_{\mathcal{U},V}(T)$ denote the homotopy class $[f_{\mathcal{U},V,p}]$ in $[T, \mathbb{S}^n]$ by $h(\mathcal{U}, V, p)$.

For a labeling L: Vert $(K) \rightarrow \{0, 1, ..., m\}$ of a simplicial complex K we denote by h(K, L, V, p) the homotopy class $h(\mathcal{U}_L(K), V, p)$ in $[|K|, \mathbb{S}^n]$.

Example 1.3. Let K be a heptagon with seven consecutive vertices labelled as 0, 1, 2, 3, 2, 1, 3. Let $V = \{v_0, v_1, v_2, v_3\}$ be the set of vertices of a planar square. Then h(K, L, V, p) = 1 if p lies in the triangle $v_0v_1v_3$ and h(K, L, V, p) = 0 otherwise.

Now we consider homotopy classes of covers of closed sets. Let $\mathcal{C} = \{C_0, \ldots, C_m\}$ be a closed cover of a space T. Let $\mathcal{U} = \{U_0, \ldots, U_m\}$ be an open cover of T such that U_i contains C_i for all i. We say that \mathcal{U} contains \mathcal{C} .

We may assume that the nerves $N(\mathcal{U})$ and $N(\mathcal{C})$ are isomorphic. Otherwise, if there is a subset of indices $J \subset \{0, \ldots, m\}$ such that the intersection of those U_i whose subindices are

in J is non-empty and the intersection of those C_i whose subindices are in J is empty we consider an open cover \mathcal{U}' with

$$U'_i := U_i \setminus K_J$$
, and $K_j := \bigcap_{j \in J} \overline{U}_j$.

Since $C_i \cap K_J = \emptyset$, we have that U'_i contains C_i .

Suppose two open covers \mathcal{U}^1 and \mathcal{U}^2 which both contain \mathcal{C} and that $N(\mathcal{U}^1)$, $N(\mathcal{U}^2)$ and $N(\mathcal{C})$ are isomorphic. Then the cover \mathcal{U}^3 such that each $U_i^3 := U_i^1 \cap U_i^2$ also contains \mathcal{C} moreover $N(\mathcal{U}^3)$ is isomorphic to $N(\mathcal{C})$. Since both \mathcal{U}^1 and \mathcal{U}^2 contain \mathcal{U}^3 , we have the equalities $h(\mathcal{U}^1, V, p) = h(\mathcal{U}^3, V, p) = h(\mathcal{U}^2, V, p)$.

This observation proves the following statement.

Lemma 1.4. Let C be a closed cover of a normal space T. Then there exist an open cover, U, of T which contains C and such that the nerves N(U) and N(C) are isomorphic.

If open covers \mathcal{U}^1 and \mathcal{U}^2 both contain \mathcal{C} and the nerves $N(\mathcal{U}^1)$, $N(\mathcal{U}^2)$ and $N(\mathcal{C})$ are isomorphic, then $h(\mathcal{U}^1, V, p) = h(\mathcal{U}^2, V, p)$.

This lemma shows that the homotopy class $h(\mathcal{C}, V, p) := h(\mathcal{U}, V, p)$ in $[T, \mathbb{S}^n]$ where $N(\mathcal{U}) = N(\mathcal{C})$ and such that \mathcal{U} contains \mathcal{C} is well defined.

2 Extension of covers

In this section we consider extensions of covers of a subspace A to a space X.

We call a family of sets a *cover* of a space if it is either an open or closed cover.

Definition 2.1. Let A be a subspace of a space X. Let $S = \{S_0, \ldots, S_m\}$ be a cover of A and $\mathcal{F} = \{F_0, \ldots, F_m\}$ be a cover of X. We assume that \mathcal{F} is open if S is open and closed if S is closed. We say that \mathcal{F} is an extension of S if

$$S_i = F_i \cap A$$
 for all *i*.

We start from the classic case: $A = \mathbb{S}^k$ and $X = \mathbb{B}^{k+1}$.

Theorem 2.1. Let $S = \{S_0, \ldots, S_{n+1}\}$ be a cover of \mathbb{S}^k . Suppose the intersection of all the S_i is empty. Then S can be extended to a cover \mathcal{F} of \mathbb{B}^{k+1} such that the intersection of all the F_i is empty if and only if [S] = 0 in $\pi_k(\mathbb{S}^n)$.

Proof. If S can be extended to \mathcal{F} , then we have $\rho_{S} : \mathbb{S}^{k} \to \mathbb{S}^{n}$ and $\rho_{\mathcal{F}} : \mathbb{B}^{k+1} \to \mathbb{S}^{n}$. Since $\rho_{S} = \rho_{\mathcal{F}} \circ i$ and $i : \mathbb{S}^{k} \to \mathbb{B}^{k+1}$ is null-homotopic, we have $[S] = [\rho_{S}] = 0$.

If [S] = 0, then we will show that S can be extended to a cover \mathcal{F} . From Lemma 1.4 it suffices to prove the theorem for open covers. Let $\Phi = \{\varphi_0, \ldots, \varphi_{n+1}\}$ be a partition of unity subordinate to S. Then we have a continuous map

$$\rho_{\mathcal{S},\Phi}: \mathbb{S}^k \to \partial \Delta^{n+1} = \mathbb{S}^n,$$

where $\rho_{\mathcal{S},\Phi} := \rho_{\mathcal{S},\Phi,V}$ (see Section 1) and V is the set of vertices of an (n+1)-simplex Δ^{n+1} . Since $[\rho_{\mathcal{S},\Phi}] = 0$ in $[\mathbb{S}^k, \mathbb{S}^n]$, there is a homotopy

$$H: \mathbb{S}^k \times [0,1] \to \mathbb{S}^n,$$

where $H(x,0) = \rho_{\mathcal{S},\Phi}(x)$, $H(x,1) = v_0$ for all x, and v_0 is a vertex of Δ^{n+1} .

Let L be a labeling on $\operatorname{Vert}(\Delta^{n+1})$ such that $L(v_i) = i$. Denote

$$U_{\ell}(\Phi, D) := H^{-1}(U_{\ell}(\Delta^{n+1}))), \ D := \mathbb{S}^k \times [0, 1],$$

where $U_L(\Delta^{n+1}) = \{U_\ell(\Delta^{n+1}), \ell = 1, \dots, n+2\}$ (see Section 3). It is clear that $\mathcal{U}_\ell(\Phi, D) := \{U_\ell(\Phi, D)\}$ is an open cover of $D, \mathcal{U}(\Phi, \mathbb{S}^k) := \mathcal{U}(\Phi, D)|_{\mathbb{S}^k}$ is a cover of \mathbb{S}^k , and

$$U_{\ell}(\Phi, \mathbb{S}^k) = \{ x \in \mathbb{S}^k : \varphi_{\ell}(x) > 0 \} \subset S_{\ell} \text{ for all } \ell.$$

Denote by $\Pi(\mathcal{S})$ the set of all partitions of unity subordinate to \mathcal{S} . Then for all ℓ

$$S_{\ell} = \bigcup_{\Phi \in \Pi(\mathcal{S})} U_{\ell}(\Phi, \mathbb{S}^k).$$

Let

$$W_{\ell} = \bigcup_{\Phi \in \Pi(\mathcal{S})} U_{\ell}(\Phi, D).$$

Then $\mathcal{W} := \{W_\ell\}$ is an open cover of D that extends \mathcal{S} .

The boundary of D consists of two components $D_0 := \mathbb{S}^k \times \{0\}$ and $D_1 := \mathbb{S}^k \times \{1\}$. Actually, \mathcal{W} on D_0 is \mathcal{S} and D_1 is covered only by one set, namely $D_1 \subset W_0$. Let Z be a (k+1)-disk with boundary D_1 and let

$$B := D \cup Z, \quad D \cap Z = D_1.$$

It is clear that B is homeomorphic to \mathbb{B}^{k+1} . Let $F_0 := W_0 \cup Z$ in B and let $\mathcal{F} := \{F_0, W_1, \ldots, W_{n+1}\}$, then \mathcal{F} is a cover of B that extends \mathcal{S} .

Next consider the case when A is the boundary of a manifold X.

Definition 2.2. Let $S = \{S_0, \ldots, S_{n+1}\}$ be a cover of an oriented *n*-dimensional manifold N without boundary. If the intersection of all S_i is empty, then $[S] \in \mathbb{Z} = [N, \mathbb{S}^n]$. We call [S] the degree of S and denote it by deg(S).

Theorem 2.2. Let M be an oriented (n + 1)-dimensional manifold with boundary, and let $S = \{S_0, \ldots, S_{n+1}\}$ be a cover of ∂M such that the intersection of all S_i is empty. Then S can be extended to a cover \mathcal{F} of M, such that all covers F_i have no a common point, if and only if the degree of S is zero.

Proof. From the Hopf Extension (Degree) Theorem it follows that a continuous map $f : \partial M \to \mathbb{S}^n$ can be extended to a globally defined continuous map $F : M \to \mathbb{S}^n$, with $\partial F = f$, if and only if the degree of f is zero. It implies that if S can be extended, then $\deg(\rho_S) = \deg(S) = 0$.

If deg(\mathcal{S}) = 0, then the proof that \mathcal{S} can be extended is almost the same as the proof in Theorem 2.1. In the last step we can use the *Collar Neighborhood Theorem* that ∂M has a neighborhood C in M which is homeomorphic to the product $D = \partial M \times [0, 1]$. Let $F_0 := W_0 \cup (M \setminus D)$. Then $\mathcal{F} := \{F_0, W_1, \ldots, W_{n+1}\}$ is a cover of M that extends \mathcal{S} . \Box

It is an interesting problem to find extensions of Theorems 2.1 and 2.2 for general X, A and V.

For extensions of the KKM and Sperner type lemmas we need pairs of spaces (X, A) such that covers of A, which are not null-homotopic, cannot be extended to X. So we need only the "necessary" parts of Theorems 2.1 and 2.2. Note that pairs of spaces (X, A) in these theorems satisfy the property that any continuous map $f : A \to \mathbb{S}^n$ with $[f] \neq 0$ cannot be extended to a continuous $F : M \to \mathbb{S}^n$.

Let $S = \{S_0, \ldots, S_{n+1}\}$ be a cover of A and $\mathcal{F} = \{F_0, \ldots, F_{n+1}\}$ be a cover of X. Suppose that the intersection of all of the S_i is empty. If \mathcal{F} is an extension of S and the intersection of all of the F_i is empty, then we have maps $\rho_S : A \to \mathbb{S}^n$ and $\rho_{\mathcal{F}} : X \to \mathbb{S}^n$ such that $\rho_{\mathcal{F}}|_A = \rho_S$. This fact motivates the following definition.

Definition 2.3. We say that a pair of spaces (X, A), where $A \subset X$, belongs to EP_n and write $(X, A) \in \text{EP}_n$ if there is a continuous map $f : A \to \mathbb{S}^n$ with $[f] \neq 0$ in $[A, \mathbb{S}^n]$ that cannot be extended to a continuous map $F : X \to \mathbb{S}^n$ with $F|_A = f$.

We denoted this class of pairs by EP after S. Eilenberg and L. S. Pontryagin who initiated obstruction theory in the late 1930s. Obstruction theory (see [6, 17]) considers homotopy invariants that equals zero if a map can be extended from the k-skeleton of X to the (k+1)-skeleton and is non-zero otherwise.

We conclude this section with a theorem that is a simple consequence of obstruction theory.

Theorem 2.3. Let (X, A) be a pair of spaces.

- 1. If the embedding $i : A \to X$ is null-homotopic and there are non null-homotopic maps $f : A \to \mathbb{S}^n$, then $(X, A) \in EP_n$. In particular, if $\pi_k(\mathbb{S}^n) \neq 0$, then $(\mathbb{B}^{k+1}, \mathbb{S}^k) \in EP_n$.
- 2. If X is an oriented (n+1)-dimensional manifold and $A = \partial X$, then $(X, A) \in EP_n$.

Proof. 1. Assume the converse. Then $f : A \to \mathbb{S}^n$, with $[f] \neq 0$, can be extended to $F : X \to \mathbb{S}^n$. Since $f = F \circ i$ and [i] = 0 in [A, X], we have that [f] = 0 in $[A, \mathbb{S}^n]$ which is a contradiction.

2. From the Hopf theorem $f: A \to \mathbb{S}^n$ can be extended if and only if [f] = 0.

3 KKM and Sperner type lemmas

The KKM (Knaster–Kuratowski–Mazurkiewicz) lemma states:

If a simplex Δ^m is covered by closed sets C_i for $i \in I_m := \{0, \ldots, m\}$ and that for all $J \subset I_m$ the face of Δ^m that is spanned by the vertices v_i with $i \in J$ is covered by C_i then all the C_i have a common intersection point.

This lemma was published in 1929 [7]. Actually, the KKM lemma is an extension of Sperner's lemma published one year before in 1928 [18].

Let T be a triangulation of a simplex Δ^m . Suppose that each vertex of T is assigned a unique label from I_m . A labeling L is called a *Sperner labeling* if the vertices are labelled in such a way that a vertex u of T belonging to a face that is spanned by vertices v_i from $\operatorname{Vert}(\Delta^m)$ for $i \in J \subset I_m$, can only be labelled by k from J. Sperner's lemma states:

Every Sperner labeling of a triangulation of Δ^m contains a cell labelled with a complete set of labels: $\{0, 1, \ldots, m\}$.

We consider extensions of the KKM and Sperner lemmas.

Theorem 3.1. Let $(X, A) \in EP_{m-1}$ and let $S = \{S_0, \ldots, S_m\}$ be a cover of A such that the intersection of all S_i is empty and $[S] \neq 0$ in $[A, S^{m-1}]$. If $\mathcal{F} = \{F_0, \ldots, F_m\}$ is a cover of X that extends S, then all the F_i have a common intersection point.

Proof. Assume the converse. Then $\rho_{\mathcal{S}} : A \to \mathbb{S}^{m-1}$ can be extended to $\rho_{\mathcal{F}} : X \to \mathbb{S}^{m-1}$ which is a contradiction.

Theorem 2.3 implies that if $\pi_k(\mathbb{S}^n) \neq 0$, then $(\mathbb{B}^{k+1}, \mathbb{S}^k) \in EP_n$.

Corollary 3.1. Let $\mathcal{F} = \{F_0, \ldots, F_m\}$ be a cover of \mathbb{B}^{k+1} that extends a cover \mathcal{S} of $\partial \mathbb{B}^{k+1} = \mathbb{S}^k$. If the intersection of all S_i is empty and $[\mathcal{S}] \neq 0$ in $\pi_k(\mathbb{S}^{m-1})$, then all the F_i have a common intersection point.

Note that for k = m-1 we have the KKM lemma. Indeed, the assumptions in this lemma yield that $[S] = \deg(S) = 1 \in \pi_k(\mathbb{S}^{m-1}) = \mathbb{Z}$.

It is interesting that this corollary can be non-trivial for k > m - 1. Consider the cover $\mathcal{U} := \mathcal{U}_{L_{\tau}}(K) = \{U_0(K), U_1(K), U_2(K), U_3(K)\}, K = \mathbb{S}^3_{12}$, from Example 1.2. Since $[\mathcal{U}] = 1 \in \pi_3(\mathbb{S}^2) = \mathbb{Z}$ Corollary 3.1 implies that

If a cover $\mathcal{F} = \{F_0, F_1, F_2, F_3\}$ of \mathbb{B}^4 is such that $\mathcal{F}|_{\partial \mathbb{B}^4} = \mathcal{U}$, then the intersection of all F_i is not empty.

However, for k = m = 2 any cover $S = \{S_0, S_1, S_2\}$ of \mathbb{S}^2 , where the S_i have no a common point, can be extended to a cover \mathcal{F} of \mathbb{B}^3 such that the intersection of all F_i is empty. Actually, it follows from the fact that $\pi_2(\mathbb{S}^1) = 0$.

Theorems 1.1 and 2.1 imply a condition for the existence of KKM type lemmas for arbitrary positive integers k and m.

Corollary 3.2. For given k and m there is a cover $S = \{S_0, \ldots, S_m\}$ of \mathbb{S}^k such that the intersection of all S_i is empty and for any cover \mathcal{F} of \mathbb{B}^{k+1} that extends S all the F_i have a common intersection point if and only if $\pi_k(\mathbb{S}^{m-1}) \neq 0$.

Now we extend Theorem 3.1 for homotopy classes $h(\mathcal{S}, V, p)$.

Definition 3.1. Let V be a set of points v_0, \ldots, v_m in \mathbb{R}^d . Consider a point $p \in \mathbb{R}^d$. Denote by $\operatorname{cov}_V(p)$ the collection of all subsets J in I_m such that simplices (convex hulls) in \mathbb{R}^d spanned by vertices $\{v_j, j \in J\}$ contain p.

It is clear that we have the following

Proposition 3.1. Let $S = \{S_0, \ldots, S_m\}$ be a cover of a space T. Let $V := \{v_0, \ldots, v_m\}$ and p be points in \mathbb{R}^d . Then $p \in P_{\mathcal{U},V}(T)$ if and only if for any $J \in \operatorname{cov}_V(p)$ the intersection of the S_i whose subindices i are in J is empty.

Theorem 3.2. Let $(X, A) \in EP_n$. Let $S = \{S_0, \ldots, S_m\}$ and $\mathcal{F} = \{F_0, \ldots, F_m\}$ be covers of A and X respectively. Let $V := \{v_0, \ldots, v_m\}$ and p be points in \mathbb{R}^{n+1} . Suppose \mathcal{F} extends S, for all $J \in cov_V(p)$ the intersection of the S_j whose subindices are in J is empty, and

$$h(\mathcal{S}, V, p) \neq 0 \text{ in } [A, \mathbb{S}^n].$$

Then there is $J \in \operatorname{cov}_V(p)$ such that

$$\bigcap_{j\in J} F_j \neq \emptyset.$$

Proof. Assume the converse. Then $p \in \mathbb{R}^{n+1} \setminus \rho_{\mathcal{F},V}(X)$. Therefore, $f_{\mathcal{F},V,p} : X \to \mathbb{S}^n$ is well defined. On the other side, it is an extension of the map $f_{\mathcal{S},V,p} : A \to \mathbb{S}^n$ with $[f_{\mathcal{S},V,p}] \neq 0$, which is a contradiction.

Theorem 3.2 implies a generalization of Sperner's lemma:

Theorem 3.3. Let X = |K| and A = |Q|, where K is a simplicial complex and Q is a subcomplex of K. Suppose $(X, A) \in EP_n$. Let $L : Vert(K) \to \{0, 1, ..., m\}$ be a labeling of K. Let $V := \{v_0, \ldots, v_m\}$ and p be points in \mathbb{R}^{n+1} . Suppose there are no simplices in Q whose vertices are labeled by $J \in cov_V(p)$. Let

$$h(Q, L, V, p) \neq 0$$
 in $[|Q|, \mathbb{S}^n]$.

Then there is a simplex s in K and there is $J \in cov_V(p)$ such that vertices of s have labels J.

If m = n + 1 and $[L] \neq 0$ in $[|Q|, \mathbb{S}^n]$, then there is a simplex in K that has all the labels in $0, \ldots, n + 1$.

There are many generalizations of the KKM and Sperner lemmas, see [1, 4, 5, 9, 11, 12, 13, 14, 16, 19]. Part of them follow from Theorems 3.2 and 3.3. As example, we consider here an extension of Tucker–Bacon lemma [1, 19] and [16].

Corollary 3.3. Let $(X, A) \in EP_n$. Let $\mathcal{F} = \{F_1, F_{-1}, \dots, F_n, F_{-n}\}$ be a cover of X that extends a cover S of A. Suppose $S_i \cap S_{-i} = \emptyset$ for all i and $h(S, V, O) \neq 0$ in $[A, \mathbb{S}^{n-1}]$, where $V := \{\pm e_1, \dots, \pm e_n\}, e_1, \dots, e_n$ is an orthonormal basis and O is the origin in \mathbb{R}^n . Then there is i such that the intersection of F_i and F_{-i} is not empty.

Proof. Note that $cov_V(O)$ consists of edges that join e_i and $(-e_i)$. Then Theorem 3.2 yields a proof.

Consider the case when X = M is an oriented manifold of dimension n+1 and $A = \partial M$. Then $[A, \mathbb{S}^n] = \mathbb{Z}$ and for any continuous $f : A \to \mathbb{S}^n$ we have $[f] = \deg f$. Now we show that it can improve Theorem 3.1.

Let s be a d-simplex. We say that s is fully labelled (or colored) if vertices of s are labeled (colored) by distinct labels ℓ_0, \ldots, ℓ_d .

Let T be a triangulation of M. Let $L : \operatorname{Vert}(M) \to \{0, 1, \dots, n+1\}$ be a labeling of vertices. Let ∂T denote the triangulation T on ∂M . We denote by $\deg(L, \partial T)$ the class $[\partial T, L]$ in $[\partial M, \mathbb{S}^n]$.

Theorem 3.4. Let T be a triangulation of a manifold M of dimension n with boundary. Then for a labeling $L : \operatorname{Vert}(T) \to \{0, 1, \ldots, n\}$ the triangulation T must contain at least $|\deg(L, \partial T)|$ fully colored simplices.

Proof. Actually, L implies a piecewise linear map $f_L : T \to \Delta^n$, where $f_L = \rho_{\mathcal{U}_L(T)}$ and deg $f_L = \deg(L, \partial T)$. Then any internal point y in Δ^n is regular for f_L , the set of preimages $f_L^{-1}(y)$ consists of points $u_k \in M$ such that every u_k lies inside of some fully labelled (n)-simplex $t_k \in T$, and the sum of the signs of u_k is equal to deg f_L . It proves the theorem. \Box

Let P be a convex polytope in \mathbb{R}^d with vertices $\{v_1, \ldots, v_m\}$. Let T be a triangulation of a manifold M of dimension d. Let $L : \operatorname{Vert}(T) \to \{1, 2, \ldots, m\}$ be a labeling of T. If, for $u \in \operatorname{Vert}(T)$, we have L(u) = i, then we set $f_{L,P}(u) := v_i$. Therefore, $f_{L,P}$ is defined for all vertices of T, and it uniquely defines a simplicial (piecewise linear) map $f_{L,P} : M \to \mathbb{R}^d$.

The following theorem extends Theorems 3.3, 3.4 and the De Loera - Petersen - Su theorem [4].

Theorem 3.5. Let P be a convex polytope in \mathbb{R}^d with m vertices. Let T be a triangulation of an oriented manifold M of dimension d with boundary. Let L: $Vert(T) \rightarrow \{1, 2, ..., m\}$ be a labeling such that $f_{L,P}(\partial M) \subseteq \partial P$. Then T contains at least $(m - d) |\deg(L, \partial T)|$ fully labelled d-simplices.

Proof. Consider a set of points S in the interior of P so that the interior of every simplex determined by d + 1 vertices in V := Vert(P) contains a unique point from S. In other words, for any two distinct points x and y in S the intersection of the sets $\text{cov}_V(x)$ and

 $\operatorname{cov}_V(y)$ is empty. Such sets have been called *pebble sets* by De Loera, Peterson, and Su. In [4] they proved that in P there is a pebble set of cardinality at least m - d. Note that $\operatorname{deg}(f_{L,P}) = h(\partial T, L, V, p)$ for any internal point p in P. Let us apply Theorem 3.3 for all points p in S. Using the same argument about the number of preimages of $f_{L,P}^{-1}(p)$ as in Theorem 3.4 we prove the theorem.

We conclude this paper by an extension of Theorem 3.5 for simplicial complexes. Let K be a d-dimensional simplicial complex. E. D. Bloch [2] defines the "boundary" of K, denoted Bd K, as the collection of all (d-1)-simplices of K that are contained in an odd number of d-simplices, together with all the faces of these (d-1)-simplices.

Let P be a convex polytope in \mathbb{R}^d with m vertices. Any labeling L : $\operatorname{Vert}(K) \to \{1, 2, \ldots, m\}$ defines a simplicial map $f_{L,P} : |K| \to P \subset \mathbb{R}^d$. So we have a map $f_{L,P}|_{|\operatorname{Bd} K|} : \operatorname{Bd}(X) \to \partial P \simeq \mathbb{S}^{d-1}$. Let us denote the degree of this map modulo 2 by $\operatorname{deg}_2(L, \operatorname{Bd} K)$. From [2, Theorem 1.5] it follows that the cardinality of $f_{L,P}^{-1}(p)$, where p lies inside of P, is equal to $\operatorname{deg}_2(L, \operatorname{Bd} K)$ modulo 2. Then the Pebble Set Theorem [4] yields the following theorem.

Theorem 3.6. Let P be a convex polytope in \mathbb{R}^d with m vertices. Let T be a triangulation of a simplicial complex X of dimension d. Let L: Vert $(T) \rightarrow \{1, 2, ..., m\}$ be a labeling such that $f_{L,P}(|\operatorname{Bd} T|) \subseteq \partial P$. If deg₂ $(L, \operatorname{Bd} T)$ is odd, then T contains at least (m - d) fully labelled d-simplices.

Corollary 3.4. Let T be a triangulation of a simplicial complex X of dimension d. If $\deg_2(L, \operatorname{Bd} T)$ for a labeling $L : \operatorname{Vert}(T) \to \{1, 2, \ldots, d+1\}$ is odd, then T must contain at least one fully colored d-simplex.

The KKM lemma and its relatives have many applications in several fields of pure and applied mathematics. In [15] we consider some extensions of results of this paper that can be applied in game theory and mathematical economics.

Acknowledgement. I wish to thank Alexander Dranishnikov, James Maissen and Alexey Volovikov for helpful discussions and comments.

References

- P. Bacon, Equivalent formulations of the Borsuk-Ulam theorem, Canad. J. Math., 18 (1966), 492–502.
- [2] E. D. Bloch. Mod 2 degree and a generalized no retraction theorem. Math. Nachr., 279 (2006), 490–494.
- [3] J. L. Bryant, Piecewise linear topology, Handbook of geometric topology, 219-259, North-Holland, Amsterdam, 2002.

- [4] J. A. De Loera, E. Peterson, and F. E. Su, A Polytopal Generalization of Sperner's Lemma, J. of Combin. Theory Ser. A, 100 (2002), 1-26.
- [5] K. Fan, A generalization of Tucker's combinatorial lemma with topological applications. Ann. of Math., 56 (1952), 431-437.
- [6] S.-T. Hu, Homotopy theory, Academic Press, 1959.
- [7] B. Knaster, C. Kuratowski, S. Mazurkiewicz, Ein Beweis des Fixpunktsatzes f
 ür ndimensionale Simplexe, Fundamenta Mathematicae 14 (1929): 132–137.
- [8] K. V. Madahar, K. S. Sarkaria, A minimal triangulation of the Hopf map and its application, Geom. Dedicata, 82 (2000), 105–114.
- [9] F. Meunier, Sperner labellings: a combinatorial approach, J. of Combin. Theory Ser. A, 113 (2006), 1462-1475.
- [10] J. W. Milnor, Topology from the differentiable viewpoint, The University Press of Virginia, Charlottesville, Virginia, 1969.
- [11] O. R. Musin, Borsuk-Ulam type theorems for manifolds, Proc. Amer. Math. Soc. 140 (2012), 2551-2560.
- [12] O. R. Musin, Extensions of Sperner and Tucker's lemma for manifolds, J. of Combin. Theory Ser. A, 132 (2015), 172–187.
- [13] O. R. Musin, Sperner type lemma for quadrangulations, Moscow Journal of Combinatorics and Number theory, 5 (2015), arXiv:1406.5082.
- [14] O. R. Musin, Generalizations of Tucker–Fan–Shashkin lemmas, arXiv:1409.8637, appear in Arnold Math. J.
- [15] O. R. Musin, KKMS type theorems with boundary conditions, in preparation
- [16] O. R. Musin and A. Yu. Volovikov, Borsuk–Ulam type spaces, arXiv:1507.08872, appear in Moscow Math. J.
- [17] E. H. Spanier, Algebraic topology, McGraw-Hill, 1966.
- [18] E. Sperner, Neuer Beweis für die Invarianz der Dimensionszahl und des Gebietes, Abh. Math. Sem. Univ. Hamburg 6 (1928), 265-272.
- [19] A. W. Tucker, Some topological properties of the disk and sphere. In: Proc. of the First Canadian Math. Congress, Montreal, 285-309, 1945.

Oleg R. Musin University of Texas Rio Grande Valley, One West University Boulevard, Brownsville, TX, 78520 and IITP RAS, Bolshoy Karetny per. 19, Moscow, 127994, Russia

 $E\text{-}mail\ address:$ oleg.musin@utrgv.edu