

Finitary Corecursion for the Infinitary Lambda Calculus*

Stefan Milius¹ and Thorsten Wißmann¹

1 Lehrstuhl für Theoretische Informatik, FAU Erlangen-Nürnberg, Germany

Abstract

Kurz et al. have recently shown that infinite λ -trees with finitely many free variables modulo α -equivalence form a final coalgebra for a functor on the category of nominal sets. Here we investigate the rational fixpoint of that functor. We prove that it is formed by all rational λ -trees, i.e. those λ -trees which have only finitely many subtrees (up to isomorphism). This yields a corecursion principle that allows the definition of operations such as substitution on rational λ -trees.

1998 ACM Subject Classification F.3.2 Semantics of Programming Languages, F.4.1 Mathematical Logic, D.3.1 Formal Definitions and Theory

Keywords and phrases rational trees, infinitary lambda calculus, coinduction

1 Introduction

One of the most important concepts in computer science is the λ -calculus. It is a very simple notion of computation because its syntax consists only of three constructs: variables, λ -abstraction and function application, and its semantics consists of only two concepts α -conversion for renaming of bound variables and β -conversion for executing function applications. Yet it is very powerful since it is Turing complete and allows to define many notions of higher level programming languages such as booleans, if-then-else, natural numbers, arithmetic operations, lists including mapping and folding, recursion etc.¹

However, whenever one wants to deal with inductive and coinductive definitions in the presence of variable binding subtle issues arise and one has to be careful not to mess up the variable binding. One solution to these problems has been proposed by Gabbay and Pitts [12]. They use *nominal sets* as a framework for dealing with binding operators, abstraction and structural induction. Nominal sets go back to Fraenkel's and Mostowski's permutation model for set theory devised in the 1920s and 1930s. They are sets equipped with an action of the group of finite permutations on a given fixed set \mathcal{V} of atoms (here these play the role of variables). For an arbitrary nominal set one can then define the notions of “free” and “bound” variables using the notion of support (we recall this in Section 2.2). Gabbay and Pitts then consider the functor

$$L_\alpha X = \mathcal{V} + [\mathcal{V}]X + X \times X$$

expressing the type of the term constructors of the λ -calculus (note that the abstraction functor $[\mathcal{V}]X$ is a quotient of $\mathcal{V} \times X$ modulo renaming “bound” variables). And they prove that the initial algebra for L_α is formed by all λ -terms modulo α -equivalence.

* This work is supported by the Deutsche Forschungsgemeinschaft (DFG) under project MI 717/5-1

¹ Depending on the application a third semantic concept, η conversion, may be of interest. But this is neither needed for Turing completeness nor for our work.



Recently, Kurz et al. [18] have characterized the final coalgebra for L_α (and more generally, for functors arising from so-called binding signatures): it is carried by the set of all infinitary λ -terms (i.e. finite or infinite λ -trees) with finitely many free variables modulo α -equivalence. This then allows to define operations on infinitary λ -terms by coinduction, for example substitution and operations that assign to an infinitary λ -term its normal form computations (e.g. the Böhm, Levy-Longo, and Berarducci trees of a given infinitary λ -term).

Our contribution in this paper is to give a characterization of the *rational fixpoint* of the functor L_α . In general, the rational fixpoint for a functor F lies between the initial algebra and the final coalgebra for F . If one thinks of it as a coalgebra, it is characterized as the final locally finitely presentable F -coalgebra. Intuitively, one may think of it as collecting all behaviours of “finite” (more technically, finitely presentable carried) F -coalgebras. Examples include regular languages, eventually periodic and rational streams, rational formal power-series etc. For a polynomial endofunctor F_Σ on sets associated to the signature Σ , the rational fixpoint consists of regular Σ -trees of Elgot [10], i.e. those (finite and infinite) Σ -trees having only finitely many different subtrees (up to isomorphism). We will prove in Section 3 that the rational fixpoint for L_α on \mathbf{Nom} is carried by all rational λ -trees modulo α -equivalence. Before that we recall in Section 2 preliminaries on the infinitary λ -calculus, nominal sets and the rational fixpoint. The finality principle of the rational fixpoint may be understood as a finitary corecursion principle. In Section 4 we show applications of our main result, in particular, that the coinductive definition of substitution given in [18] restricts to rational trees. We also discuss coinductive definitions concerning normal form computations. We conclude in Section 5.

Related work. The work presented here is based on the second author’s student project reported in [27].

A related approach to variable binding operations which uses presheaves over finite sets was proposed by Fiore, Plotkin and Turi [11]. By now this has developed into a respectable body of work by these and other authors. Most related to our work here is the coinductive approach to infinitary and rational λ -terms studied by Adámek, Milius and Velebil [3]. This work considers an endofunctor very similar to L_α but on the category of presheaves on finite sets. Its final coalgebra is shown to be the presheaf of all infinite λ -trees and the rational fixpoint the presheaf of all rational trees – each of them modulo α -equivalence.

2 Preliminaries

We assume that readers are familiar with basic notions of category theory and with algebras and coalgebras for an endofunctor. For a given endofunctor F on the category \mathcal{C} we will write $t : \nu F \rightarrow F(\nu F)$ for the final coalgebra (assuming that it exists). Given an F -coalgebra (C, c) we write $c^\dagger : (C, c) \rightarrow (\nu F, t)$ for the unique F -coalgebra homomorphism from C to νF . The category of coalgebras for an endofunctor F is denoted by $\mathbf{Coalg} F$. For introductory texts on coalgebras see [25, 16, 1].

We will now give some background on the (infinitary) λ -calculus, on nominal sets and on the rational fixpoint of a functor as needed in the present paper.

2.1 Infinitary λ -Calculus and Rational Trees

Before we talk about infinitary λ -terms (aka λ -trees) first recall that ordinary λ -terms are defined starting from a fixed countable set of variables \mathcal{V} by the grammar

$$T ::= x \mid \lambda x.T \mid TT,$$

where x ranges over \mathcal{V} . We denote the set of all λ -terms by Λ . Free and bound variables and substitution are defined as usual with the operator $\lambda x.(-)$ binding x in its argument. Often one considers λ -terms modulo α -equivalence, i.e., the least equivalence relation on λ -terms identifying two terms that arise by consistently renaming bound variables. One can think of a term $\lambda x.T$ as representing a computation that takes a parameter P that is used in all free occurrences of x in T . Hence, the main computation rule of the λ -calculus is β -reduction, i.e. the rule

$$(\lambda x.T)P \rightarrow_{\beta} T[x \mapsto P].$$

For example we have $(\lambda x.\lambda y.x) a b \rightarrow_{\beta} (\lambda y.a) b \rightarrow_{\beta} a$, where a cannot be reduced further. However, terms may have infinite reduction sequences; a prominent example is Yf for the Y -combinator defined as $Y := \lambda g.(\lambda x.g(xx)) (\lambda x.g(xx))$ we have:

$$\begin{aligned} Yf &= (\lambda g.(\lambda x.g(xx)) (\lambda x.g(xx)))f \\ &\rightarrow_{\beta} (\lambda x.f(xx)) (\lambda x.f(xx)) \rightarrow_{\beta} f((\lambda x.f(xx)) (\lambda x.f(xx))) \\ &\rightarrow_{\beta} f(f((\lambda x.f(xx)) (\lambda x.f(xx)))) \rightarrow_{\beta} \dots \end{aligned}$$

Informally speaking, this “converges” to the infinite term $f(f(f(\dots)))$. If one takes such infinite terms as legal objects of the λ -calculus one is led to *infinitary* λ -calculus. There one replaces λ -terms by (finite and infinite) λ -trees. A λ -tree is a rooted and ordered tree with leaves labelled by variables in \mathcal{V} and with two sorts of inner nodes: nodes with one successor labelled by λx for some variable $x \in \mathcal{V}$ and nodes with two successors labelled by $@$. For example, we have the λ -trees

$$(2.1)$$

representing the λ -term $(\lambda x.xx)(\lambda x.xx)$ and the infinite term $f(f(f(\dots)))$, respectively. Let Λ^{∞} be the set of all λ -trees. The notions of free and bound variables of a λ -tree are clear: a variable x is bound in a λ -tree t if there is a path from a leaf labelled by x to the root of t that contains a node labelled by λx , and x is free in t if there is a path from an x -labelled leaf to the root of t that does not contain any node labelled by λx .

The classic approach to defining operations such a substitution on λ -trees uses that Λ^{∞} is the metric completion of Λ under a natural metric; this idea of using a metric approach to dealing with infinite trees goes at least back to Arnold and Nivat [5]. Thus, every infinite λ -tree is regarded as the limit of the Cauchy sequence of its truncations at level n . Notions such as α -equivalence and substitution of λ -trees are then defined by extending the corresponding notions on finite λ -trees (i.e. λ -terms) continuously. More concretely, two λ -trees s and t are α -equivalent iff for every natural number n the pair of truncations at level n of s and t are α -equivalent λ -terms (see [18, Definition 5.17]).

Our aim in this paper is to give a coalgebraic characterization of an important subclass of all λ -trees, the so called *rational* λ -trees. The following definition follows Ginali’s characterization [15] of regular Σ -trees for a signature Σ :

► **Definition 2.1.** A λ -tree having only finitely many subtrees (up to isomorphism) is called *rational*. A λ -tree modulo α -equivalence, i.e. an α -equivalence class of λ -trees, is called *rational* if it contains at least one rational λ -tree.

containing any variable, i.e. the empty set supports x .

3. The set Λ of all λ -terms forms a nominal set with the group action given by renaming of free variables. Every λ -term is supported by the set of its free variables. In contrast the set Λ^∞ of all λ -trees is not nominal since λ -trees with infinitely many free variables do not have finite support. However, the set $\Lambda_{\text{ffv}}^\infty$ of all λ -trees with finitely many free variables is nominal.

Notice that if $S \subseteq V$ supports $x \in X$, then $S' \supseteq S$ also supports $x \in X$. So S supporting x only means that by not touching the members of S one does not modify the term x . But it is more interesting to talk about the variables actually occurring in x . This is achieved by considering the smallest set supporting x , which is denoted by $\text{supp}(x)$. If $v \in \mathcal{V} \setminus \text{supp}(x)$, we say that v is *fresh for* x , denoted by $v \# x$.

► **Example 2.3.** The set $\mathcal{P}_f(\mathcal{V})$ of finite subsets of \mathcal{V} , together with the point-wise action is a nominal set. The support of each $u \in \mathcal{P}_f(\mathcal{V})$ is u itself: $\pi \cdot u = \{\pi \cdot x \mid x \in u\}$ and $\text{supp}(u) = u$. Note that $\mathcal{P}(\mathcal{V})$ with the point-wise action is not a nominal set because the infinite $\{v_0, v_2, v_4, \dots\}$ does not have any finite support.

The morphisms of nominal sets are those maps which are equivariant: an *equivariant map* $f : (X, \cdot) \rightarrow (Y, \star)$ is a map $f : X \rightarrow Y$ with $f(\pi \cdot x) = \pi \star f(x)$ for all $\pi \in \mathfrak{S}(\mathcal{V}), x \in X$.

For example, the function $\text{supp} : X \rightarrow \mathcal{P}_f(\mathcal{V})$ mapping each element to its (finite) support is an equivariant map.

► **Remark 2.4.** For any equivariant $f : (X, \cdot) \rightarrow (Y, \star)$, we have $\text{supp}(f(x)) \subseteq \text{supp}(x)$ for any $x \in X$.

The nominal sets – together with the equivariants as morphisms – form a category, denoted by **Nom**. As shown in [14], this category is (equivalent to) a Grothendieck topos (the so-called Shanel topos), and so it has rich categorical structure. We only mention some facts needed for the current paper.

Monomorphisms and epimorphisms in **Nom** are precisely the injective and surjective equivariant maps, respectively. It is not difficult to see that every epimorphism in **Nom** is strong, i.e., it has the unique diagonalization property w.r.t. any monomorphism: given an epimorphism $e : A \twoheadrightarrow B$, a monomorphism $m : C \hookrightarrow D$ and $f : A \rightarrow C, g : B \rightarrow D$ with $g \cdot e = m \cdot f$, there exists a unique diagonal $d : B \rightarrow C$ with $d \cdot e = f$ and $m \cdot d = g$.

Furthermore, **Nom** has image-factorizations; that means that every equivariant map $f : A \rightarrow C$ factorizes as $f = m \cdot e$ for an epimorphism $e : A \twoheadrightarrow B$ and a monomorphism $m : B \hookrightarrow C$. Note that the intermediate object B is (isomorphic to) the image $f[A]$ in B with the restricted action. For an endofunctor F on **Nom** preserving monos this factorization systems lifts to **Coalg** F : every F -coalgebra homomorphism f has a factorization $f = m \cdot e$ where e and m are F -coalgebra homomorphisms that are epimorphic and monomorphic in **Nom**, respectively.

Recall from [23, Section 2.2] that **Nom** is complete and cocomplete with colimits and finite limits formed as in **Set**. In fact, **Nom** is a locally finitely presentable category in the sense of Gabriel and Ulmer [13] (see also Adámek and Rosický [4]). We shall not recall that notion here as it is not needed in the current paper; intuitively, a locally finitely presentable category is a category with a well behaved “finite” objects (called *finitely presentable* objects) such that every object can be build (as a filtered colimit) from these. Petrişan [22, Proposition 2.3.7] has shown that the finitely presentable objects of **Nom** are precisely the orbit-finite nominal sets.

► **Definition 2.5.** For a nominal set (X, \cdot) and $x \in X$ the set $\{\pi \cdot x \mid \pi \in \mathfrak{S}(\mathcal{V})\}$ is called the *orbit* of x . A nominal set (X, \cdot) is said to be *orbit-finite* if it has only finitely many orbits.

The notion of orbit-finiteness plays a central role in our paper since the rational λ -trees modulo α -equivalence are described by precisely all the coalgebras with an orbit-finite carrier for the functor L_α further below (cf. Proposition 2.11 and Theorem 3.4).

We now collect a few easy properties of orbit-finite sets that we are going to need. (The proofs can be found in the appendix.)

► **Lemma 2.6.** *For any $x_1, x_2 \in X$ in the same orbit, we have $|\text{supp}(x_1)| = |\text{supp}(x_2)|$.*

► **Lemma 2.7.** *For an element x of a nominal set X , there are at most $|\text{supp}(x)|!$ many elements with support $\text{supp}(x)$ in the orbit of x .*

► **Lemma 2.8.** *For a finite set $W \subseteq \mathcal{V}$ and an orbit \mathcal{O} of the nominal set X there are only finitely many elements in \mathcal{O} whose support is contained in W .*

Let us now recall from Kurz et al. [18] how all λ -trees form a final coalgebra in **Nom**. First consider the following endofunctor on **Nom**:

$$LX = \mathcal{V} + \mathcal{V} \times X + X \times X;$$

its coproduct components describe the type of the term constructors of the λ -calculus (variables, λ -abstraction and application, respectively). As shown in [18], the final coalgebra for this functor is carried by the set of all λ -trees containing finitely many (free and bound) variables.² Its coalgebra structure is the obvious map decomposing a λ -tree at the root: a single node λ -tree is mapped to its node label in \mathcal{V} , a λ -tree whose root is labeled by λx to (x, t) , where t is the λ -tree defined by the successor of the root and a λ -tree with root label $@$ to the pair of λ -trees defined by the successors of the root.

Since this final coalgebra completely disregards α -equivalence it is not possible to define substitution as a total operation on it. The solution is to replace the second component $\mathcal{V} \times X$ of L by Gabbay and Pitts abstraction functor [12, Lemma 5.1] that takes α -equivalence into account:

► **Definition 2.9.** Let (X, \cdot) be a nominal set. We define α -equivalence \sim_α as the relation on $\mathcal{V} \times X$ as

$$(v_1, x_1) \sim_\alpha (v_2, x_2) \text{ if there exists } z \# \{v_1, v_2\}, z \# x_1, z \# x_2 \text{ with } (v_1 z)x_1 = (v_2 z)x_2.$$

The \sim_α -equivalence class of (v, x) is denoted by $\langle v \rangle x$. The abstraction $[\mathcal{V}]X$ of the nominal X is the quotient $(\mathcal{V} \times X)/\sim_\alpha$ with the group action defined by

$$\pi \cdot \langle v \rangle x = \langle \pi(v) \rangle (\pi \cdot x).$$

For an equivariant map $f : X \rightarrow Y$, $[\mathcal{V}]f : [\mathcal{V}]X \rightarrow [\mathcal{V}]Y$ is defined by $\langle v \rangle x \mapsto \langle v \rangle (f(x))$.

Note that the abstraction functor $[\mathcal{V}](-)$ is *strong*, i.e., we have a natural transformation τ with components $\tau_{X,Y} : [\mathcal{V}]X \times Y \rightarrow [\mathcal{V}](X \times Y)$ given by $\tau_{X,Y}(\langle v \rangle x, y) = \langle v \rangle (x, y)$; we will need the strength τ in Section 4.1. Now one considers the endofunctor L_α on **Nom** given by

$$L_\alpha X = \mathcal{V} + [\mathcal{V}]X + X \times X.$$

Gabbay and Pitts [12] showed that its initial algebra consists of all λ -terms modulo α -equivalence, and the main result of Kurz et al. [18] is that the final coalgebra νL_α is carried by the set $\Lambda_{\text{ffv}}^\infty$ of all λ -trees with finitely many free variables quotiented by α -equivalence. The coalgebra structure is the same as on the final coalgebra for L – one can show that this is well-defined on equivalence classes modulo α -equivalence.

² Note that this is different from the set $\Lambda_{\text{ffv}}^\infty$ mentioned in Example 2.10.3; λ -trees in the latter may have infinitely many bound variables.

2.3 The Rational Fixpoint

Recall that by Lambek's Lemma [19], the structure of an initial algebra and a final coalgebra for a functor F are isomorphisms, so both yield *fixpoints* of F . Here we shall be interested in a third fixpoint that lies in between initial algebra and final coalgebra called the *rational fixpoint* of F . This can on the one hand be characterized as the initial iterative algebra for F (see [2]) or as the final locally finitely presentable coalgebra for F (see [20]). We will only recall the latter description since the former will not be needed in this paper.

The rational fixpoint can be defined for any *finitary* endofunctor F on a locally finitely presentable (lfp, for short) category \mathcal{C} , i.e. F is an endofunctor on \mathcal{C} that preserves filtered colimits. Examples of lfp categories are **Set**, the categories of posets and of graphs, every finitary variety of algebras (such as groups, rings, and vector spaces) and every Grothendieck topos (such as **Nom**). The finitely presentable objects in these categories are: all finite sets, posets or graphs, those algebras presented by finitely many generators and relations, and, as we mentioned before, the orbit-finite nominal sets.

Now let $F : \mathcal{C} \rightarrow \mathcal{C}$ be finitary on the locally finitely presentable category \mathcal{C} and consider the full subcategory $\text{Coalg}_f F$ of $\text{Coalg } F$ given by all F -coalgebras with a finitely presentable carrier. In [20] the *locally finitely presentable* F -coalgebras were characterized as precisely those coalgebras that arise as a colimit of a filtered diagram of coalgebras from $\text{Coalg}_f F$. It follows that the *final* locally finitely presentable coalgebra can be constructed as the colimit of *all* coalgebras from $\text{Coalg}_f F$. More precisely, one defines a coalgebra $r : \varrho F \rightarrow F(\varrho F)$ as the colimit of the inclusion functor of $\text{Coalg}_f F$: $(\varrho F, r) := \text{colim}(\text{Coalg}_f F \hookrightarrow \text{Coalg } F)$. Note that since the forgetful functor $\text{Coalg } F \rightarrow \mathcal{C}$ creates all colimits this colimit is actually formed on the level of \mathcal{C} . The colimit ϱF then carries a uniquely determined coalgebra structure r making it the colimit above.

As shown in [2], ϱF is a fixpoint for F , i.e. its coalgebra structure r is an isomorphism. From [20] we obtain that local finite presentability of a coalgebra (C, c) has the following concrete characterizations: (1) for $\mathcal{C} = \text{Set}$ local finiteness, i.e. every element of C is contained in a finite subcoalgebra of C ; (2) for $\mathcal{C} = \text{Nom}$, local orbit-finiteness, i.e. every element of C is contained in an orbit-finite subcoalgebra of C ; (3) for \mathcal{C} the category of vector spaces over a field K , local finite dimensionality, i.e., every element of C is contained in a subcoalgebra of C carried by a finite dimensional subspace of C .

► **Example 2.10.** We list only a few examples of rational fixpoints; for more see [2, 20, 8].

1. Consider the functor $FX = 2 \times X^A$ on **Set** where A is an input alphabet and $2 = \{0, 1\}$. The F -coalgebras are precisely the deterministic automata over A (without initial states). The final coalgebra is carried by the set $\mathcal{P}(A^*)$ of all formal languages and the rational fixpoint is its subcoalgebra of regular languages over A .
2. For $FX = \mathbb{R} \times X$ on **Set** the final coalgebra is carried by the set \mathbb{R}^ω of all real streams and the rational fixpoint is its subcoalgebra of all eventually periodic streams, i.e. streams $uvvv \cdots$ with $u, v \in \mathbb{R}^*$. Taking the same functor on the category of real vector spaces we get the same final coalgebra \mathbb{R}^ω with the componentwise vector space structure, but this time the rational fixpoint is formed by all rational streams (see [26, 20]).
3. Let Σ be a signature of operation symbols with prescribed arity, i.e. a sequence $(\Sigma_n)_{n < \omega}$ of sets. This give rise to an associated polynomial endofunctor F_Σ on **Set** given by $F_\Sigma X = \coprod_{n < \omega} \Sigma_n \times X^n$. Its initial algebra is formed by all Σ -terms and its final coalgebra by all (finite and infinite) Σ -trees, i.e. rooted and ordered trees such that every node with n children is labelled by an n -ary operation symbol. And the rational fixpoint consists precisely of all regular Σ -trees of Elgot [10] (see also Courcelle [9]), i.e. those Σ -trees having only finitely many different subtrees up to isomorphism (see Ginali [15]).

Note that in all the above examples the rational fixpoint ϱF always occurs as a sub-coalgebra of the final coalgebra νF . But this need not be the case in general (see [8, Example 3.15] for a counterexample). However, we have the following result:

► **Proposition 2.11** ([8, Proposition 3.12]). *Suppose that in \mathcal{C} finitely presentable objects are closed under strong quotients and that F is finitary and preserves monomorphisms. Then the rational fixpoint ϱF is the subcoalgebra of νF given by the union of images of all coalgebra homomorphisms $c^\dagger : (C, c) \rightarrow (\nu F, t)$ where (C, c) ranges over $\text{Coalg}_f F$.³*

In particular, for a finitary functor F on Set (or Nom resp.) preserving monomorphisms the rational fixpoint is the union in νF of images of all finite (or orbit-finite resp.) coalgebras; in symbols:

$$\varrho F = \bigcup_{(C, c) \text{ in } \text{Coalg}_f F} c^\dagger[C] \subseteq \nu F.$$

Note that it is sufficient to let (C, c) range over those coalgebras in $\text{Coalg}_f F$ where c^\dagger is injective (or an inclusion map) because for an arbitrary (orbit-)finite (C, c) in $\text{Coalg}_f F$ its image $c^\dagger[C]$ is an (orbit-)finite F -coalgebra, too.

3 The Rational Fixpoint in Nominal Sets

In this section we are going to prove the main result of our paper, a characterization of the rational fixpoint for the functor L_α as the rational λ -trees modulo α -equivalence.

But we start with the rational fixpoint of the functor $LX = \mathcal{V} + \mathcal{V} \times X + X \times X$. Note that both functors L and L_α are finitary and preserve monomorphisms (to see this use [18, Proposition 5.6] and the fact the forgetful functor from Nom to Set creates colimits).

► **Proposition 3.1.** *The rational fixpoint of the functor $L : \text{Nom} \rightarrow \text{Nom}$ is formed by all rational λ -trees.*

In the proof of the following theorem we will slightly abuse notation and consider $LX = \mathcal{V} + \mathcal{V} \times X + X \times X$ as an endofunctor on Set . Note that its final coalgebra is formed by the set Λ^∞ of all λ -trees and its rational fixpoint by all rational λ -trees (this follows from Example 2.10.3).

► **Theorem 3.2.** *Let $X \xrightarrow{a} L_\alpha X$ be an orbit-finite coalgebra. Then for all $\text{root} \in X$, $a^\dagger(\text{root}) \in \varrho L_\alpha$ is a rational λ -tree.*

Proofsketch. (for a full proof see the appendix) Let $m := \max_{x \in X} |\text{supp}(x)|$ be the maximal number of free variables in any element of X . This exists by Lemma 2.6 since X is orbit-finite. Let $W \subseteq \mathcal{V}$ be some set of $m+1$ variables containing $\text{supp}(\text{root})$. Hence for all $x \in X$ there exists a $w \in W$ with $w \# x$.

In the following, one constructs a rational λ -tree in the α -equivalence class of $a^\dagger(\text{root})$. First, define an L -coalgebra $C \xrightarrow{c} LC = \mathcal{V} + \mathcal{V} \times C + C \times C$ in Set with $C := \{x \in X \mid \text{supp}(x) \subseteq W\}$ and

$$c(x) = \begin{cases} w & \text{if } a(x) = w \in W \subseteq \mathcal{V} \\ (\ell, r) & \text{if } a(x) = (\ell, r) \in X \times X \\ (w, y) & \text{if } a(x) = \langle v \rangle y' \text{ and } y = (v \ w)y' \text{ for some } w \in W \setminus \text{supp}(x) \end{cases}$$

³ In a general lfp category the *image* of c^\dagger is obtained by taking a strong epi-mono factorization of c^\dagger , and the *union* is then obtained as a directed colimit of the resulting subobjects of $(\nu F, t)$.

Next one readily verifies that c is well-defined, i.e., its image really lies in LC .

Furthermore, C is finite because X is orbit-finite and within any orbit there are only finitely many elements with a support contained in W by Lemma 2.8. Let c^\dagger denote the unique L -coalgebra homomorphism into the final L -coalgebra in **Set**. Since C is finite, we know that $c^\dagger : C \rightarrow \nu L$ factors through the rational fixpoint, i.e. for every $x \in C$, $c^\dagger(x)$ is a rational λ -tree. One then proves that $[c^\dagger(x)]_\alpha = a^\dagger(x)$ for all $x \in C \subseteq X$, where $[-]_\alpha$ denotes α -equivalence classes. This involves a non-trivial induction argument using the *final chains* of the set functor L and the functor L_α on **Nom** as well as technical details from [18] (see the appendix). It follows that $a^\dagger(\text{root})$ is rational. \blacktriangleleft

For the L -coalgebra (C, c) in **Set** from the previous proof, we know that for any $x \in C$, the resulting tree $c^\dagger(x)$ has at most $|C|$ subtrees. This does not hold for an L_α -coalgebra (X, a) in **Nom**: if X has a non-trivial action, then the cardinality of X is at least infinite, i.e. the cardinality does not give a reasonable bound for the number of subtrees. And the number of orbits n is not a bound either. The problem is that multiple elements from the same orbit may represent different subtrees. For example, consider the rational tree

$$t := \begin{array}{c} \textcircled{\alpha} \\ / \quad \backslash \\ v_0 \quad v_1 \end{array},$$

and let $X := \mathcal{V} + \{(\ell, r) \in \mathcal{V} \times \mathcal{V} \mid \ell \neq r\}$ be equipped with the coalgebra structure

$$X \xrightarrow{a} L_\alpha X, \quad a(x) = \begin{cases} v & \text{if } x = v \in \mathcal{V} \\ (\ell, r) & \text{if } x = (\ell, r) \in \mathcal{V} \times \mathcal{V}. \end{cases}$$

X is constructed to have two orbits: one consisting of single variables and one consisting of unequal ordered pairs of variables. We have $(v_0, v_1) \in X$ and $a^\dagger((v_0, v_1)) = [t]_\alpha$. But t has three subtrees, namely t itself, v_0 , and v_1 . This example is expanded later in Example 3.5.

But when looking closer at the construction of C in the previous proof, we can give a bound on the number of its elements, i.e. the number of (up to isomorphism) different subtrees of the rational tree $c^\dagger(\text{root})$.

► Proposition 3.3. *Let (X, a) be an orbit-finite L_α -coalgebra with n orbits and let $m = \max_{x \in X} |\text{supp}(x)|$. Then the number of elements of the coalgebra C (as constructed in the previous proof) is bounded by $n \cdot (m + 1)!$.*

Proof. Recall from the proof of Theorem 3.2 that $C := \{x \in X \mid \text{supp}(x) \subseteq W\}$, where W is a set of $m + 1$ variables. Consider a fixed orbit O whose elements have a support of cardinality k : there are at most $\binom{m+1}{k}$ possibilities of choosing a k -element subset S of W and for any such S there are at most $k!$ elements in O with support S , by Lemma 2.7. Thus, the number of elements of O in C is at most

$$\binom{m+1}{k} \cdot k! = \frac{(m+1)!}{k! \cdot (m+1-k)!} \cdot k! = \frac{(m+1)!}{(m+1-k)!} \stackrel{k \leq m}{\leq} \frac{(m+1)!}{(m+1-m)!} = (m+1)!.$$

In total, the cardinality of C is bounded by $n \cdot (m + 1)!$. \blacktriangleleft

That the number of orbits n occurs linearly is not surprising, because if we have some finite carrier set X with the trivial action that “uses” all its elements for the coalgebra structure, we have exactly one subtree per element of X . In Example 3.5 we shall see that the factor is $(m + 1)!$ is necessary. But before that we state and prove our main result:

► **Theorem 3.4.** *The rational fixed point ϱL_α contains precisely the rational λ -trees modulo α -equivalence.*

Proof. After Theorem 3.2 it only remains to show that all α -equivalence classes of rational λ -trees are in ϱL_α . Let $u_\alpha \in \nu L_\alpha$ be rational, witnessed by some rational representative $u \in u_\alpha$ with only finitely many subtrees (up to isomorphism). Let C be the finite set of all subtrees of u and define the nominal set X as

$$X := \bigcup_{s \in C} O([s]_\alpha) \subseteq \nu L_\alpha,$$

where $[s]_\alpha \in \nu L_\alpha$ is the α -equivalence class of the subtree s and $O(y) \subseteq \nu L_\alpha$ denotes the orbit of a given element $y \in \nu L_\alpha$. Note that the group action of νL_α restricts to X since it is a union of orbits.

Next we define a coalgebra structure $a : X \rightarrow L_\alpha X$ by restriction of the structure $t : \nu L_\alpha \rightarrow L_\alpha(\nu L_\alpha)$: set $a(x) := t(x)$ for all $x \in X$. We need to check that this is well-defined, i.e. that $t(x)$ really lies in $L_\alpha X$ for every $x \in X$. For this we consider three cases:

1. The case $t(x) \in \mathcal{V}$ is clear;
2. Suppose that $t(x) = \langle v \rangle y \in [\mathcal{V}](\nu L_\alpha)$ where $x = \pi \cdot [\lambda w.s]_\alpha$ for some $\pi \in \mathfrak{S}(\mathcal{V})$ and some subtree $\lambda w.s$ of u . Then we have

$$\langle w \rangle [s]_\alpha = t([\lambda w.s]_\alpha) = t(\pi^{-1} \cdot x) = \pi^{-1} \cdot t(x) = \pi^{-1} \cdot \langle v \rangle y = \langle \pi^{-1}(v) \rangle (\pi^{-1} \cdot y).$$

By the definition of abstraction, we have some $z \in \mathcal{V}$ with

$$(w z) \cdot [s]_\alpha = (\pi^{-1}(v) z) \cdot \pi^{-1} \cdot y.$$

Hence, y is in the orbit of $[s]_\alpha$ and therefore $t(x)$ is in $[\mathcal{V}]X$.

3. For $t(x) = (\ell, r) \in \nu L_\alpha \times \nu L_\alpha$, let $x \in X$ be $\pi \cdot [(s_\ell, s_r)]_\alpha$ for some $\pi \in \mathfrak{S}(\mathcal{V})$ and subtrees s_ℓ, s_r of u . Analogously to the previous case, we have


$$(\ell, r) = t(x) = t(\pi \cdot [(s_\ell, s_r)]_\alpha) = \pi \cdot ([s_\ell]_\alpha, [s_r]_\alpha) = (\pi \cdot [s_\ell]_\alpha, \pi \cdot [s_r]_\alpha) \in X \times X.$$

By construction, $u_\alpha \in X$ and $a^\dagger(x) = x$ holds for all $x \in X$. By the finiteness of C , X is orbit-finite and thus $a^\dagger[X] \subseteq \varrho L_\alpha$, and in particular $u_\alpha \in \varrho L_\alpha$. ◀

The following example shows that the bound in Proposition 3.3 on the number of elements of the L -coalgebra C from the proof of Theorem 3.2 can essentially not be improved even if we omit the usage of λ -abstraction.

► **Example 3.5.** Let $\ell \geq 1$ be a fixed natural number and let $m = 2^{\ell-1}$. Further let $V = \{v_1, \dots, v_m\}$ be a set of m variables and consider the rational λ -trees parametrized by permutations $\sigma \in \mathfrak{S}(\mathcal{V})$ shown in Figure 2. With r_σ and h_σ we denote the corresponding subtrees rooted at the indicated nodes. Note that the α -equivalence classes in ϱL_α of each r_σ and of any of its subtrees are singletons since r_σ does not contain any λ -abstraction. For this reason we shall henceforth abuse notation and denote those equivalence classes by their representatives. Observe further that the group action on ϱL_α satisfies $\tau \cdot r_\sigma = r_{\tau\sigma}$ for any $\tau, \sigma \in \mathfrak{S}(\mathcal{V})$. This implies that all r_σ and h_σ , respectively, lie in the same orbit. Similarly, one can see that all nodes on the same level in the right-hand maximal subtrees of every r_σ (indicated by the dotted rectangles) lie in the same orbit.

Now consider r_{id} and the corresponding orbit-finite subcoalgebra X of ϱL_α from the proof of Theorem 3.4. The elements of X are all subtrees of r_{id} with the group action



■ **Figure 2**

inherited from ϱL_α . By the above reasoning we see that X has precisely $\ell + 2$ orbits. Hence, the number of orbits of X is logarithmic in m . But the number of subtrees of r_{id} grows faster than $m!$. To see this, notice first that the permutations $(1\ 2)$ and $(1\ 2\ \dots\ m)$ generate the group of all permutations of m elements. Thus, we see that r_{id} has all r_σ as subtrees where σ is any permutation that fixes $\mathcal{V} \setminus V$. For different σ and τ we have that r_σ , h_σ , r_τ and h_τ are pairwise non-isomorphic. Thus, we see that r_{id} has at least $2 \cdot m!$ subtrees. In addition, consider the i -th level (from 1 at the bottom to ℓ at the top) on the right in Figure 2. Each node on that level covers $k = 2^{i-1}$ variables. Then for each permutation of those k variables there exists a subtree of the right-hand successor of some subtree r_σ of r_{id} which is a complete binary tree of height i with the front given by the permutation. Thus, the total number of subtrees of r_{id} is precisely

$$2 \cdot m! + \sum_{i=1}^{\ell} \left(\binom{m}{2^{i-1}} \cdot (2^{i-1})! \right) = 2 \cdot m! + \sum_{i=1}^{\ell} \frac{m!}{(m - 2^{i-1})!}.$$

4 Application: Corecursive Definitions on Rational λ -trees

Our result in Theorem 3.4 that rational λ -trees modulo α -equivalence form the final locally orbit-finite L_α -coalgebra yields a corecursion principle. In this section we shall demonstrate this principle by considering two easy applications. First we show that substitution as defined corecursively for all λ -trees in [18] restricts to rational λ -trees. Secondly, we discuss the corecursive definition of the computation of the Böhm tree of a given rational λ -tree.

4.1 Substitution on Rational λ -trees

When performing operations known from (infinitary) λ -calculus on rational λ -trees, it is not clear whether the resulting λ -tree still is rational in general. One such operation is the substitution function $\text{subs} : \nu L_\alpha \times \mathcal{V} \times \nu L_\alpha \rightarrow \nu L_\alpha$, which for a given triple (t, v, s) replaces each occurrence of the variable v in t by s . Kurz et al. [18] show how to define subs by coinduction: to do this they define an L_α -coalgebra whose unique homomorphism into the final coalgebra yields subs . It is possible to adapt this to the orbit-finite case as follows.

For arbitrary coalgebras $A \xrightarrow{a} L_\alpha A$ and $B \xrightarrow{b} L_\alpha B$ with orbit-finite carriers, one defines $\text{subs}_{A,B} : A \times \mathcal{V} \times B \rightarrow \varrho L_\alpha$. This map $\text{subs}_{A,B}$ describes the substitution of a variable within an element of the coalgebra A by some element of the coalgebra B . In the following

$$B \xrightarrow{\text{inl}} B + A \times \mathcal{V} \times B \xleftarrow{\text{inr}} A \times \mathcal{V} \times B \quad \text{and} \quad \mathcal{V} \xrightarrow{\text{in}_1} L_\alpha X \xleftarrow{\text{in}_2} [\mathcal{V}]X$$

denote coproduct injections. Now define $B + A \times \mathcal{V} \times B \xrightarrow{[g,h]} L_\alpha(B + A \times \mathcal{V} \times B)$ with $g = L_\alpha(\text{inl}) \circ b$ and $h = [h_{\text{Var}}, h_{\text{Abs}}, h_{\text{App}}] \circ (a \times \text{id} \times \text{id})$ using distributivity and

- $h_{\text{Var}} : \mathcal{V} \times \mathcal{V} \times B \rightarrow L_\alpha(B + A \times \mathcal{V} \times B)$, $h_{\text{Var}}(v, w, x) = \begin{cases} \text{in}_1 v & \text{if } v \neq w \\ L_\alpha(\text{inl})(\underbrace{b(x)}_{\in L_\alpha(B)}) & \text{if } v = w \end{cases}$
- $h_{\text{Abs}} : ([\mathcal{V}]A) \times \mathcal{V} \times B \rightarrow L_\alpha(B + A \times \mathcal{V} \times B)$, $h_{\text{Abs}} := \text{in}_2 \circ \tau_{A, \mathcal{V} \times B}$, using the strength τ of the functor $[\mathcal{V}]$ with $\tau_{A, \mathcal{V} \times B} : ([\mathcal{V}]A) \times \mathcal{V} \times B \rightarrow [\mathcal{V}](A \times \mathcal{V} \times B)$.
- $h_{\text{App}} : A \times A \times \mathcal{V} \times B \rightarrow L_\alpha(B + A \times \mathcal{V} \times B)$ is the composition of the obvious steps $A \times A \times \mathcal{V} \times B \rightarrow (A \times \mathcal{V} \times B) \times (A \times \mathcal{V} \times B) \rightarrow L_\alpha(B + A \times \mathcal{V} \times B)$.

$B + A \times \mathcal{V} \times B$ is orbit-finite, so we get a unique L_α -coalgebra homomorphism $[g', h']$ into ϱL_α , making the following diagram commute:

$$\begin{array}{ccc} B + A \times \mathcal{V} \times B & \xrightarrow{[g,h]} & L_\alpha(B + A \times \mathcal{V} \times B) \\ \downarrow [g', h'] & & \downarrow L_\alpha([g', h']) \\ \varrho L_\alpha & \xrightarrow{f_\alpha} & L_\alpha(\varrho L_\alpha) \end{array}$$

Now we define $\text{subs}_{A,B} := h'$. It remains to extend this to the desired domain $\varrho L_\alpha \times \mathcal{V} \times \varrho L_\alpha$. Let I be the set of all orbit-finite subcoalgebras (X, a) of $(\nu L_\alpha, t_\alpha)$. Then we know from (the discussion following) Proposition 2.11 that

$$\varrho L_\alpha = \bigcup_{(X,a) \text{ in } I} X.$$

Hence, we can define substitution $\text{subs}_{\text{rat}} : \varrho L_\alpha \times \mathcal{V} \times \varrho L_\alpha \rightarrow \varrho L_\alpha$ on rational λ -trees by

$$\text{subs}_{\text{rat}}(x, v, y) = \text{subs}_{A,B}(x, v, y), \text{ for some } A, B \in I \text{ with } x \in A \text{ and } y \in B.$$

It remains to prove that the result does not depend on the choice of A and B . But if we have any other (A', a') in I with $x \in A'$ then since both A and A' are subcoalgebras we have $a(x) = t_\alpha(x) = a'(x)$. Similarly for B and $y \in B$.

Thus, since the function h is defined by pattern matching, i.e. on the alternatives indicated by a , it behaves independently from the choice of A and B as desired.

To summarize, we can say that one can define operations on ϱL_α if these n -ary operations can be defined restricted to n orbit-finite subcoalgebras of νL_α as we have just seen for the operation of substitution.

4.2 Normalization of Rational λ -trees

In the λ -calculus different kinds of normal forms play an important role. One of them is the *head normal form* (*hnf*, for short). A λ -term is in hnf if it is of the form $\lambda x_1 \dots \lambda x_n. y N_1 \dots N_m$, where y is a variable and the N_i are arbitrary terms. If one recursively requires the N_i to be in hnf as well, one gets the definition of *Böhm trees*. Adding an additional constant symbol \perp to the syntax of the λ -calculus allows the following corecursive definition of the Böhm tree $\text{BT}(M)$ of a λ -term M (see [18]):

$$\text{BT}(M) = \begin{cases} \lambda x_1 \dots \lambda x_n. y \text{BT}(N_1) \dots \text{BT}(N_m) & \text{if } M \rightarrow_\beta \lambda x_1 \dots \lambda x_n. y N_1 \dots N_m \\ \perp & \text{otherwise,} \end{cases} \quad (4.1)$$

5 Conclusion and Future Work

We have contributed to the abstract algebraic study of variable binding using nominal sets. In particular, we have extended a recent coalgebraic approach to infinitary λ -calculus due to Kurz et al. Whereas they proved in [18] that λ -trees with finitely many variables modulo α -equivalence form the final coalgebra for the functor L_α on **Nom** we have given a characterization of the rational fixpoint of that functor. It contains precisely the rational λ -trees modulo α -equivalence.

This characterization entails a corecursion principle for rational λ -trees because the rational fixpoint is the final locally orbit-finite coalgebra for L_α . In this sense we have achieved finitary corecursion for the infinitary λ -calculus. We have demonstrated the new principle and its limitations with two applications: a corecursive definition of substitution and of a normalform computation.

Our work is only a first step in the study of the coalgebraic approach to finitary coinduction for infinitary terms with variable binding operators. First, it should be clear that our results generalize from λ -terms to the rational fixpoint for endofunctors on **Nom** associated to a binding signature. Other points for future work are: (1) the extension of the coalgebraic approach to rational and infinitary λ -terms using nominal sets to treat the solutions of higher-order recursion schemes as was done in the setting of presheaves on finite sets in [3], and (2) the study of specification formats that extend our simple corecursion principle that follows from finality; more precisely, Bonsangue et al. [7, 21] have proposed bipointed specifications as an abstract format (by restricting Turi's and Plotkin's abstract GSOS rules [24]) to specify algebraic operations on the rational fixpoint of an endofunctor. It should be interesting to work out a concrete rule format corresponding to bipointed specifications for rational λ -terms and rational terms for arbitrary binding signatures. Last, but not least, the similarity of the results in [3] on the one hand and those in [18] and here on the other hand is so striking that there should be a formal connection; however, to our knowledge this has not been worked out in the literature yet.

References

- 1 Jiří Adámek. Introduction to coalgebra. *Theory Appl. Categ.*, 14:157–199, 2005.
- 2 Jiří Adámek, Stefan Milius, and Jiri Velebil. Iterative algebras at work. *Math. Structures Comput. Sci.*, 16(6):1085–1131, 2006.
- 3 Jiří Adámek, Stefan Milius, and Jiri Velebil. Semantics of higher-order recursion schemes. *Log. Methods Comput. Sci.*, 7(1), 2011.
- 4 Jiří Adámek and Jiří Rosický. *Locally presentable and accessible categories*. Cambridge University Press, 1994.
- 5 André Arnold and Maurice Nivat. The metric space of infinite trees. algebraic and topological properties. *Fundam. Inform.*, 3(4):445–476, 1980.
- 6 Hendrik Pieter Barendregt. *The Lambda calculus: Its syntax and semantics*. North-Holland, Amsterdam, 1984.
- 7 Marcello M. Bonsangue, Stefan Milius, and Jurriaan Rot. On the specification of operations on the rational behaviour of systems. In *Proc. EXPRESS/SOS'12*, volume 89 of *Electron. Proc. Theoret. Comput. Sci.*, pages 3–18, 2012.
- 8 Marcello M. Bonsangue, Stefan Milius, and Alexandra Silva. Sound and complete axiomatizations of coalgebraic language equivalence. *ACM Trans. Comput. Log.*, 14(1:7), 2013.
- 9 Bruno Courcelle. Fundamental properties of infinite trees. *Theoret. Comput. Sci.*, 25:95–169, 1983.

- 10 Calvin C. Elgot. Monadic computation and iterative algebraic theories. In H. E. Rose and J. C. Sheperdson, editors, *Logic Colloquium '73*, volume 80, pages 175–230, Amsterdam, 1975. North-Holland Publishers.
- 11 Marcelo Fiore, Gordon D. Plotkin, and Daniele Turi. Abstract syntax and variable binding. In *Proc. Logic in Computer Science 1999*, pages 193–202. IEEE Press, 1999.
- 12 Murdoch Gabbay and Andrew M. Pitts. A new approach to abstract syntax involving binders. In *Proc. LICS'99*, pages 214–224. IEEE Computer Society Press, 1999.
- 13 Peter Gabriel and Friedrich Ulmer. *Lokal präsentierbare Kategorien*, volume 221 of *Lecture Notes Math.* Springer-Verlag, 1971.
- 14 Fabio Gaducci, Marino Miculan, and Ugo Montanari. About permutation algebras, (pre)sheaved and names sets. *Higher-Order Symb. Comput.*, 19:283–304, 2006.
- 15 Susanna Ginali. Regular trees and the free iterative theory. *J. Comput. System Sci.*, 18:228–242, 1979.
- 16 Bart Jacobs and Jan Rutten. A tutorial on (co)algebras and (co)induction. *EATCS Bulletin*, 62:62–222, 1997.
- 17 Richard Kennaway, Jan Willem Klop, Ronan Sleep, and Fer-Jan de Vries. Infinitary Lambda Calculus. *Theoret. Comput. Sci.*, 175(1):93–125, 1997.
- 18 Alexander Kurz, Daniela Petrisan, Paula Severi, and Fer-Jan de Vries. Nominal coalgebraic data types with applications to lambda calculus. *Log. Methods Comput. Sci.*, 9(4), 2013.
- 19 Joachim Lambek. A fixpoint theorem for complete categories. *Math. Z.*, 103:151–161, 1968.
- 20 Stefan Milius. A sound and complete calculus for finite stream circuits. In *Proc. LICS'10*, pages 449–458. IEEE Computer Society, 2010.
- 21 Stefan Milius, Marcello M. Bonsangue, Robert S.R. Myers, and Jurriaan Rot. Rational operation models. In *Proc. MFPS XXIX*, volume 298 of *Electron. Notes Theor. Comput. Sci.*, pages 257–282, 2013.
- 22 Daniela Petrişan. *Investigations into Algebra and Topology over Nominal Sets*. dissertation, University of Leicester, 2011.
- 23 Andrew M. Pitts. *Nominal Sets: Names and Symmetry in Computer Science*, volume 57 of *Cambridge Tracts in Theoretical Computer Science*. Cambridge University Press, 2013.
- 24 Gordon D. Plotkin and Daniele Turi. Towards a mathematical operational semantics. In *Proc. Logic in Computer Science (LICS'97)*, pages 280–291, 1997.
- 25 Jan Rutten. Universal coalgebra: a theory of systems. *Theoret. Comput. Sci.*, 249(1):3–80, 2000.
- 26 Jan Rutten. Rational streams coalgebraically. *Log. Methods Comput. Sci.*, 4(3:9):22 pp., 2008.
- 27 Thorsten Wißmann. The rational fixed point in nominal sets and its application to infinitary lambda-calculus. Project report, available at <http://thorsten-wissmann.de/theses/project-wissmann.pdf>, October 2014.

A Proof of Lemma 2.6

Proof. Assuming $\pi \cdot x_1 = x_2$, the equivariance of supp provides the equation

$$\text{supp}(x_1) = \text{supp}(\pi \cdot x_2) = \pi \cdot \text{supp}(x_2) = \{\pi(v) \mid v \in \text{supp}(x_2)\}.$$

Since π is a bijective map this establishes the desired bijection between $\text{supp}(x_1)$ and $\text{supp}(x_2)$. ◀

B Proof of Lemma 2.7

Proof. Consider the set

$$N = \{\pi \cdot x \mid \pi \in \mathfrak{S}(\mathcal{V}) \text{ and } \text{supp}(\pi \cdot x) = \text{supp}(x)\}$$

All π with $\pi \cdot x \in N$ have to fulfill the property $\text{supp}(x) = \text{supp}(\pi \cdot x) = \pi \cdot \text{supp}(x)$. This means, π restricts to $\text{supp}(x)$, i.e. $\pi(v) \in \text{supp}(x)$ iff $v \in \text{supp}(x)$. Hence, each such π can be factorized into two permutations $f_\pi, g_\pi \in \mathfrak{S}(\mathcal{V})$ where

$$f_\pi(v) = \begin{cases} \pi(v) & \text{if } v \in \text{supp}(x) \\ v & \text{if } v \notin \text{supp}(x) \end{cases}, \quad g_\pi(v) = \begin{cases} v & \text{if } v \in \text{supp}(x) \\ \pi(v) & \text{if } v \notin \text{supp}(x) \end{cases}$$

and with $f_\pi \circ g_\pi = g_\pi \circ f_\pi = \pi$. Since $\text{supp}(x)$ supports x we have $g_\pi \cdot x = x$, and this implies

$$\begin{aligned} N &= \{f_\pi \cdot g_\pi \cdot x \mid \pi \in \mathfrak{S}(\mathcal{V}) \text{ and } \pi \cdot \text{supp}(x) = \text{supp}(x)\} \\ &= \{f_\pi \cdot x \mid \pi \in \mathfrak{S}(\mathcal{V}) \text{ and } \pi \cdot \text{supp}(x) = \text{supp}(x)\}. \end{aligned}$$

The set of permutations $\{f_\pi \mid \pi \cdot x \in N\}$ has cardinality $|\text{supp}(x)|!$, hence so is N . ◀

C Proof of Lemma 2.8

Proof. First there are only finitely many elements whose support is exactly W . If there is no element in \mathcal{O} with support W , then we are done. And if x in \mathcal{O} satisfies $\text{supp}(x) = W$, there are only finitely many elements in \mathcal{O} with the same support by Lemma 2.7.

The statement of the lemma now follows since W has only finitely many subsets and for each $W' \subseteq W$ there are only finitely many elements with support W' . ◀

D Proof of Proposition 3.1

Proof. Take a tree $t \in \varrho L$ and an orbit-finite subcoalgebra (S, s) of ϱL with $t \in S$. Every subtree t' of t has support

$$\text{supp}(t') \subseteq \text{supp}(t).$$

But there are only finitely many $t' \in S$ with that property; this follows from Lemma 2.8 using $W = \text{supp}(t)$ and that S is orbit-finite. Hence, t has only finitely many subtrees, i.e. t is rational. So ϱL contains only rational λ -trees. Conversely, for each rational λ -tree t we can construct an orbit-finite L -coalgebra by taking the subtrees of t as the carrier set of that coalgebra and with the coalgebra structure given by decomposing trees at the root. ◀

E Proof of Theorem 3.2

Let m be the maximal number of free variables in any element of X , i.e.

$$m := \max_{x \in X} |\text{supp}(x)|.$$

The maximum exists by Lemma 2.6 since X is orbit-finite.

Let $W \subseteq \mathcal{V}$ be some set of $m + 1$ variables containing $\text{supp}(\text{root})$. Hence

$$\text{for all } x \in X \text{ there exists a } w \in W \text{ with } w \# x. \quad (\text{E.1})$$

In the following, we will construct a rational λ -tree in the α -equivalence class of $a^\dagger(\text{root})$. First, define an L -coalgebra $C \xrightarrow{c} LC = \mathcal{V} + \mathcal{V} \times C + C \times C$ in **Set**:

$$C := \{x \in X \mid \text{supp}(x) \subseteq W\}, \quad c(x) = \begin{cases} w & \text{if } a(x) = w \in W \subseteq \mathcal{V} \\ (\ell, r) & \text{if } a(x) = (\ell, r) \in X \times X \\ (w, y) & \text{if } a(x) = \langle v \rangle y' \text{ and } y = (v w) y' \\ & \text{for some } w \in W \setminus \text{supp}(x) \end{cases}$$

We verify that c is well-defined, i.e., its image lies in LC :

- For the case $a(x)$ in \mathcal{V} , $a(x)$ is also in W because $x \in C$ and thus

$$\text{supp}(a(x)) \subseteq \text{supp}(x) \subseteq W.$$

- For the case $a(x) = (\ell, r)$ in $X \times X$,

$$\text{supp}(\ell) \cup \text{supp}(r) \subseteq \text{supp}(x) \subseteq W.$$

So $\ell, r \in C$.

- For $a(x) = \langle v \rangle y' \in [\mathcal{V}]X$, we know by (E.1), that there is such a fresh $w \in W$. So $\langle w \rangle y = \langle v \rangle y'$. In particular, $y \in C$, because

$$\text{supp}(y) \subseteq \{w\} \cup \text{supp}(a(x)) \subseteq \{w\} \cup \text{supp}(x) \subseteq \{w\} \cup W \subseteq W.$$

The last item gives that $\langle w \rangle y = \langle v \rangle y'$, hence the following diagram commutes in **Set**:

$$\begin{array}{ccc} C & \xrightarrow{c} & LC \xleftarrow{Li} LX \\ \downarrow i & & \downarrow qx \\ X & \xrightarrow{a} & L_\alpha X \end{array} \quad (\text{E.2})$$

Observe that C is finite, because X is orbit-finite and within an orbit there are only finitely many elements with a support contained in W by Lemma 2.8. Let c^\dagger denote the unique L -coalgebra morphism into the final L -coalgebra in **Set**. Since C is finite, we know that $c^\dagger : C \rightarrow \nu L$ factors through the rational fixpoint, i.e. for every $x \in C$, $c^\dagger(x)$ is a rational λ -tree. In particular, $c^\dagger(\text{root})$ is a rational λ -tree. Using (E.2) and the *final chains* of the set functor L and the functor L_α on **Nom** we shall prove below that

$$[c^\dagger(x)]_\alpha = a^\dagger(x) \quad \text{for all } x \in C \subseteq X, \quad (\text{E.3})$$

where $[-]_\alpha$ denotes α -equivalence classes. For $x = \text{root}$ this gives that $c^\dagger(\text{root})$ is the desired rational λ -tree in $a^\dagger(\text{root})$ and thus the α -equivalence class $a^\dagger(\text{root})$ is rational.

We will now prove Equation (E.3). Before we can proceed to the proof we need to recall some technical background (cf. [18]). First recall that every endofunctor F on a category \mathcal{C} with a final object 1 induces its final chain

$$1 \leftarrow F1 \leftarrow FF1 \leftarrow \dots$$

Whenever the limit of this chain exists in \mathcal{C} and F preserves this limit, then it carries the final F -coalgebra νF . Moreover, every F -coalgebra $c : C \rightarrow FC$ induces its *canonical cone* $c^n : C \rightarrow F^n 1$, $n < \omega$, on the final chain defined inductively as: $c^0 : C \rightarrow 1$ is the unique map, and given c^n we define

$$c^{n+1} = C \xrightarrow{c} FC \xrightarrow{Fc^n} FF^n 1 = F^{n+1} 1 \quad \text{for all } n > 0.$$

The unique F -coalgebra homomorphism $c^\dagger : C \rightarrow \nu F$ is then induced by the universal property of the limit – c^\dagger is the unique morphism such that

$$c^n = (C \xrightarrow{c^\dagger} \nu F \xrightarrow{p_n} F^n 1) \quad \text{for every } n < \omega, \quad (\text{E.4})$$

where $p_n : \nu F \rightarrow F^n 1$ denote the limit projections.

Now let us turn to our concrete functors of interest. The terminal coalgebra for the functor $L : \mathbf{Set} \rightarrow \mathbf{Set}$ consists of all finite and infinite λ -trees. Since L preserves all limits, νL is the limit of its final chain $L^n 1$ and we denote the limit projections by

$$p_n : \nu L \rightarrow L^n 1$$

and the canonical cone induced by $c : C \rightarrow LC$ by

$$c^n : C \rightarrow L^n 1.$$

We also consider the final chain of the functor L_α on \mathbf{Nom} , and we take its limit in $\lim L_\alpha^n 1$ in \mathbf{Set} with the projections

$$p_{\alpha,n} : \lim L_\alpha^n 1 \rightarrow L_\alpha^n 1, \quad \text{for } n < \omega.$$

Notice that this limit does *not* carry the final L_α -coalgebra. But νL_α is a subset of the limit via the unique map $\iota_\alpha : \nu L_\alpha \rightarrow \lim L_\alpha^n 1$ induced by the canonical cone $(t_\alpha^n)_{n < \omega}$ induced by the structure $t_\alpha : \nu L_\alpha \rightarrow L_\alpha(\nu L_\alpha)$ of the final coalgebra, i.e., ι_α is the unique map such that

$$t_\alpha^n = (\nu L_\alpha \xrightarrow{\iota_\alpha} \lim L_\alpha^n 1 \xrightarrow{p_{\alpha,n}} L_\alpha^n 1) \quad \text{for all } n < \omega. \quad (\text{E.5})$$

We will also use the canonical cone

$$a^n : X \rightarrow L_\alpha^n 1, \quad n < \omega,$$

of our given L_α -coalgebra (X, a) .

As a final ingredient we need to relate the final chains of L and L_α . To this end observe that L can be construed as an endofunctor on \mathbf{Nom} (more precisely, L lifts to \mathbf{Nom}), and then we have a quotient natural transformation $q : L \rightarrow L_\alpha$ with components

$$q_X = (LX = \mathcal{V} + \mathcal{V} \times X + X \times X \xrightarrow{\text{id} + \theta_X + \text{id} \times \text{id}} \mathcal{V} + [\mathcal{V}]X + X = L_\alpha X,$$

where $\theta_X : \mathcal{V} \times X \rightarrow [\mathcal{V}]X$ is the natural transformation given by the canonical quotient maps (cf. Definition 2.9). Using q one can define a natural transformation $[-]_\alpha^n : L^n 1 \rightarrow L_\alpha^n 1$, $n < \omega$, between the respective final chains by induction: $[-]_\alpha^0$ is the identity on 1 and

$$[-]_\alpha^{n+1} = L^{n+1} 1 = LL^n 1 \xrightarrow{L[-]_\alpha^n} LL_\alpha^n 1 \xrightarrow{qL_\alpha^n} L_\alpha L_\alpha^n 1 = L_\alpha^{n+1} 1.$$

Now we regard the two final chains and the natural transformation given by the $[-]_\alpha^n$ in Set and use the universal property of the limit of the $L_\alpha^n 1$ to obtain a unique map $[-]_\alpha : \nu L \rightarrow \lim L_\alpha^n 1$ such that the following squares commute for all $n < \omega$:

$$\begin{array}{ccc} L^n 1 & \xleftarrow{p_n} & \nu L \\ [-]_\alpha^n \downarrow & & \downarrow [-]_\alpha \\ L_\alpha^n 1 & \xleftarrow{p_{\alpha,n}} & \lim L_\alpha^n 1 \end{array} \quad (\text{E.6})$$

We denote by $i : C \hookrightarrow X$ the inclusion map and consider the following diagram:

$$\begin{array}{ccccc} & & c^\dagger & & \\ & \swarrow & \longleftarrow & \longleftarrow & \longleftarrow \\ \nu L & \xleftarrow{\quad} & \Lambda_{\text{ffv}}^\infty & \xleftarrow{\quad} & C \\ [-]_\alpha \downarrow & & \downarrow & & \downarrow i \\ \lim L_\alpha^n 1 & \xleftarrow{\nu_\alpha} & \nu L_\alpha & \xleftarrow{a^\dagger} & X \end{array} \quad (\text{E.7})$$

The set $\Lambda_{\text{ffv}}^\infty$ consists of all λ -trees with finitely many free variables, and this makes the left-hand square above the pullback of $[-]_\alpha$ along ν_α as proved in [18, Proposition 5.33]. There it was also shown that the map opposite $[-]_\alpha$ is surjective (see [18, Theorem 5.34]), and it follows that νL_α consists of all equivalence classes modulo α -equivalence of λ -trees in $\Lambda_{\text{ffv}}^\infty$.

Now note that once we prove that the outside of Diagram (E.7) commutes we obtain the dashed map making the diagram commutative. This finishes our proof since the right-hand square of (E.7) is precisely our desired equation (E.3). We now establish the commutativity of the outside of the diagram by proving inductively that

$$p_{\alpha,n} \cdot \nu_\alpha \cdot a^\dagger \cdot i = p_{\alpha,n} \cdot [-]_\alpha \cdot c^\dagger, \quad \text{for all limit projections } p_{\alpha,n}, n < \omega.$$

The base case is clear and for the induction step we compute:

$$\begin{aligned} p_{\alpha,n+1} \cdot \nu_\alpha \cdot a^\dagger \cdot i &= t_\alpha^{n+1} \cdot a^\dagger \cdot i && \text{by (E.5)} \\ &= L_\alpha t_\alpha^n \cdot t_\alpha \cdot a^\dagger \cdot i && \text{definition of } t_\alpha^{n+1} \\ &= L_\alpha t_\alpha^n \cdot L_\alpha a^\dagger \cdot a \cdot i && a^\dagger \text{ coalgebra homomorphism} \\ &= L_\alpha t_\alpha^n \cdot L_\alpha a^\dagger \cdot q_X \cdot Li \cdot c && \text{see (E.2)} \\ &= q_{L_\alpha^n 1} \cdot Lt_\alpha^n \cdot La^\dagger \cdot Li \cdot c && q \text{ natural and } t_\alpha^n, a^\dagger \text{ equivariant} \\ &= q_{L_\alpha^n 1} \cdot Lp_{\alpha,n} \cdot L\nu_\alpha \cdot La^\dagger \cdot Li \cdot c && \text{by (E.5)} \\ &= q_{L_\alpha^n 1} \cdot L(p_{\alpha,n} \cdot \nu_\alpha \cdot a^\dagger \cdot i) \cdot c && \text{functoriality of } L \\ &= q_{L_\alpha^n 1} \cdot L(p_{\alpha,n} \cdot [-]_\alpha \cdot c^\dagger) \cdot c && \text{induction hypothesis} \\ &= q_{L_\alpha^n 1} \cdot Lp_{\alpha,n} \cdot L[-]_\alpha \cdot Lc^\dagger \cdot c && \text{functoriality of } L \\ &= q_{L_\alpha^n 1} \cdot L[-]_\alpha^n \cdot Lp_n \cdot Lc^\dagger \cdot c && \text{by (E.6)} \\ &= q_{L_\alpha^n 1} \cdot L[-]_\alpha^n \cdot Lc^n \cdot c && \text{by (E.4)} \\ &= [-]_\alpha^{n+1} \cdot Lc^n \cdot c && \text{definition of } [-]_\alpha^{n+1} \\ &= [-]_\alpha^{n+1} \cdot c^{n+1} && \text{definition of } c^{n+1} \\ &= [-]_\alpha^{n+1} \cdot p_{n+1} \cdot c^\dagger && \text{by (E.4)} \\ &= p_{\alpha,n+1} \cdot [-]_\alpha \cdot c^\dagger && \text{by (E.6)} \end{aligned}$$

This completes the proof of the desired equation (E.3). ◀