# Estimation of the parameter of a dynamically selected population for two subclasses of the exponential family

Morteza Amini†∗and Nader Nematollahi ‡

† Department of Statistics, School of Mathematics, Statistics and computer Science,

College of Science, University of Tehran, P.O. Box 14155-6455, Tehran, Iran

‡ Department of Statistics, Allameh Tabataba'i University, Tehran, Iran

September 21, 2021

#### Abstract

We introduce the problem of estimation of the parameters of a dynamically selected population in an infinite sequence of random variables and provide its application in the statistical inference based on record values from a non-stationary scheme. We develop unbiased estimation of the parameters of the dynamically selected population and evaluate the risk of the estimators. We provide comparisons with natural estimators and obtain asymptotic results. Finally, we illustrate the applicability of the results using real data.

Keywords: Extreme value theory, General record models, Partial maxima, Pfeifer model, Selected population, Uniformly minimum variance unbiased estimator.

## 1 Introduction

The problem of estimating parameters of selected populations has wide practical applications in estimation of experimental data in agriculture, industry and medicine. Some of the real world applications of this theory are the problem of estimating the average yield of a selected variety of plant with maximum yield (Kumar and Kar, 2001), estimating the average fuel efficiency of the vehicle with minimum fuel consumption (Kumar and Gangopadhyay, 2005) and selecting the regimen with maximal efficacy or minimal toxicity from a set of regimens and estimating a treatment effect for the selected regimen (Sill and Sampson, 2007).

<sup>∗</sup>Corresponding Author, E-mail address: morteza.amini@ut.ac.ir (Morteza Amini) nematollahi@atu.ac.ir (Nader Nematollahi)

The problem of estimation after selection has received considerable attention by many researches in the past three decades. Interested readers are referred to, for example, Gibbons et al. (1977) for more details. Some other contributions in this area include Sarkadi (1967), Dahiya (1974), Kumar and Kar (2001), Misra et al. (2006a,b), Kumar et al. (2009) and Nematollahi and Motammed-Shariati (2012). For a summary of results, as well as a list of references until 2006, see Misra et al. (2006 a,b).

In this paper, we introduce and develop the problem of estimation of the parameters of a dynamically selected population from a sequence of infinite populations which is not studied in the literature, according to the best of our knowledge. Let  $X_1, X_2, \cdots$  be a sequence of random variables where  $X_i$  is drawn from population  $\Pi_i$  with corresponding cumulative distribution function (cdf)  $F_{\theta_i}$ . and probability density function (pdf)  $f_{\theta_i}$ . The traffic volume trend, daily temperatures, sequences of stock quotes, or sequences of estimators of interior water volume in a dam reservoir are examples of such sequences.

Suppose we want to estimate the parameter of the population corresponding to the largest value of the sequence  $X_1, X_2, \cdots$  yet seen, that is

$$
\theta^U_{[n]} = \theta_{T_n},
$$

where  $T_1 = 1$ , with probability one, and for  $n > 1$ 

$$
T_n = \min\{j; j > T_{n-1}; X_j > X_{T_{n-1}}\},\
$$

or similarly the parameter of the population corresponding to the smallest value of the sequence  $X_1, X_2, \cdots$  yet seen, that is

$$
\theta^L_{[n]} = \theta_{T'_n},
$$

where  $T_1' = 1$ , with probability one, and for  $n > 1$ 

$$
T'_{n} = \min\{j; j > T_{n-1}; X_{j} < X_{T_{n-1}}\}.
$$

We want to estimate  $\theta_{[n]}^U$ , and similarly the lower ones  $\theta_{[n]}^L$ . This happens for example, when we want to estimate the largest value of traffic volume or stock quotes yet seen, the temperature of the coldest day or the largest volume of the coming water into the dam reservoir, up to now.

For simplicity, we denote  $\theta_{[n]}^U$  by  $\theta_{[n]}$  hereafter. We may write

<span id="page-1-0"></span>
$$
\theta_{[n]} = \sum_{j=n}^{\infty} \theta_j I_j(X_1, X_2, \ldots),
$$
\n(1)

where

<span id="page-1-1"></span>
$$
I_j = I_j(X_1, X_2, \ldots) = \begin{cases} 1, & \max X_k < X_{T_{n-1}} < X_j \\ 1, & \max X_k < j-1 \\ 0, & \text{o.w.} \end{cases}
$$
\n
$$
= I(\max\{X_k; \ T_{n-1} + 1 \le k \le j-1\} < X_{T_{n-1}} < X_j). \tag{2}
$$

The statistics  $U_n = X_{T_n}$  and  $L_n = X_{T'_n}$  are called upper and lower records, respectively. In the sequence  $X_1, X_2, \ldots$ , the sequences of partial maxima and upper record statistics are defined by  $M_n = \max\{X_1, X_2, \ldots, X_n\}$  and  $U_n = X_{T_n} = M_{T_n}$ , respectively, where  $T_1 = 1$ with probability 1, and  $T_{n+1} = \min\{j; M_j > M_{T_n}\}\$ , for  $n \geq 1$ . The record statistics  $U_n$ could be viewed as the dynamic maxima of the original random variables. So, we call the problem of estimating  $\theta_{[n]}$  as the estimation of the parameter of a dynamically selected population.

There is a vast literature on records for iid as well as non-stationary random variables. A thorough survey of available results, until 1998, is given in the book of Arnold et al. (1998). More recent articles on record values include, among others, Amini and Balakrishnan (2013, 2015), Doostparast and Emadi (2013), Salehi et al. (2013), Ahmadi and Balakrishnan (2013, 2010), Psarrakos and Navarro (2013), Raqab and Ahmadi (2012), Zarezadeh and Asadi (2010), Kundu et al. (2009) and Baklizi (2008).

This problem is related to the so-called *general record model*. The geometrically increasing populations, the Pfeifer, the linear drift and the  $F^{\alpha}$  record models are some of the generally used record models. The basics of non-stationary schemes for the record values are due to Nevzorov (1985, 1986) and Pfeifer (1989, 1991), who considered the socalled  $F^{\alpha}$ -scheme, that is the sequences of independent random variables with distribution  $F_k(x) = (F(x))^{\theta_k}, k = 1, 2, \ldots$ , where F is a continuous cdf and  $\theta_k$ 's are positive parameters. Further generalization of the  $F^{\alpha}$ -scheme was suggested by Ballerini and Resnick (1987). Although non-stationary schemes could be employed in the most general setting, the special case of improving populations is usually of special interest. Alternative nonstationary schemes include geometrically increasing populations, linear trend and Pfeifer models.

In all the above models, strict assumptions are made on the sequence of parameters  $\{\theta_i\}_{i\geq 1}$ . For instance, in  $F^{\alpha}$  record model, the sequence of the parameters is assumed to be known or depend on a fixed unknown parameter. In the linear drift model, a linearly increasing population is assumed as the underlying population. However, certain natural phenomena may behave otherwise. For example, an earthquake is produced by a natural phenomenon which has a pivotal parameter that varies based on an unknown model. In order to predict extremely destructive earthquakes, a very important question is on the value of the parameters which cause a new record in the sequence of earthquakes? This motivates us to study the problem of dynamic after-selection estimation.

The rest of this paper is organized as follows. The theoretical results of the dynamic after-selection problem, consisting unbiased estimation of the parameters of the model as well as unbiased estimation of the risk of the estimators are presented in Sections 2 and 3. In Section 4, we compare the proposed estimators with some natural estimators. Asymptotic distributional results for studying the limiting behavior of the risks of the estimators are studied in Section 5. Finally, a real data example is considered in section 6 to illustrate the applicability of the results.

## 2 Minimum variance unbiased estimation

Let  $\boldsymbol{\theta} = (\theta_1, \theta_2, \ldots), \mathbf{X} = (X_1, X_2, \ldots)$  and  $h_{\mathbf{X}}(\boldsymbol{\theta})$  be a random parameter (a function of **X** and  $\theta$ ). Suppose that  $h_{\mathbf{X}}(\theta)$  is estimated by  $\delta(\mathbf{X})$ . Following Lehmann (1951), the estimator  $\delta(\mathbf{X})$  is said to be risk unbiased for  $h_{\mathbf{X}}(\theta)$  under the loss function  $L(h_{\mathbf{X}}(\theta), \delta(\mathbf{X}))$ , if it satisfies

<span id="page-3-0"></span>
$$
E_{\theta}(L(h_{\mathbf{X}}(\theta), \delta(\mathbf{X}))) \le E_{\theta}(L(h_{\mathbf{X}}(\theta'), \delta(\mathbf{X}))), \ \forall \theta' \ne \theta.
$$
 (3)

Under the squared error loss (SEL) function

<span id="page-3-3"></span>
$$
L(h_{\mathbf{X}}(\boldsymbol{\theta}), \delta(\mathbf{X})) = (h_{\mathbf{X}}(\boldsymbol{\theta}) - \delta(\mathbf{X}))^{2},
$$

the condition [\(3\)](#page-3-0) reduces to

$$
E_{\theta}(\delta(\mathbf{X})) = E_{\theta}(h_{\mathbf{X}}(\theta)).
$$
\n(4)

In this section, we use the U-V method of Robbins (1988), to find Uniformly Minimum Variance Unbiased (UMVU) estimator of  $\theta_{[n]}$  under the two models 1 and 2, presented below.

**Model 1:** Let  $X_1, X_2, \cdots$  be a sequence of independent absolutely continuous random variables with pdf

<span id="page-3-1"></span>
$$
f(x_i; \theta_i) = c(x_i)\theta_i^{-p} e^{-S(x_i)/\theta_i}, \qquad (5)
$$

where  $S(X_i)$  is a complete sufficient statistic with the Gamma $(p, \theta_i)$ -distribution. Some well-known members of the above family are:

- 1. Exponential( $\theta_i$ ), with  $p = 1$ ,  $S(x_i) = x_i$  and  $c(x_i) = 1$ ;
- 2. Gamma $(p, \theta_i)$ , with  $S(x_i) = x_i$  and  $c(x_i) = x_i^{p-1}$  $i^{p-1}/\Gamma(p);$
- 3. Normal $(0, \sigma_i^2)$ , with  $\theta_i = \sigma_i^2$ ,  $p = 1/2$ ,  $S(x_i) = x_i^2/2$  and  $c(x_i) = (2\pi)^{-1/2}$ ;
- 4. Inverse Gaussian( $\infty, \lambda_i$ ), with  $\theta_i = 1/\lambda_i$ ,  $p = 1/2$ ,  $S(x_i) = 1/(2x_i)$  and  $c(x_i) =$  $(2x_i^3)^{-1/2};$
- 5. Weibull $(\eta_i, \beta)$ , with known  $\beta, \theta_i = \eta_i^{\beta}$  $i^{\beta}, p = 1, S(x_i) = x_i^{\beta}$  $i \atop i$  and  $c(x_i) = \beta x_i^{\beta - 1};$
- 6. Rayleigh $(\beta_i)$ , with  $\theta_i = \beta_i^2$ ,  $p = 1$ ,  $S(x_i) = x_i^2/2$  and  $c(x_i) = x_i$ .

To estimate  $\theta_{[n]}$  in the family of distributions [\(5\)](#page-3-1), we first consider the estimation of  $\theta_{[n]}$  under the Gamma $(p, \theta_i)$ -distribution with pdf

<span id="page-3-2"></span>
$$
f(x_i|\theta_i) = \frac{1}{\theta_i^p \Gamma(p)} x_i^{p-1} \exp\{x_i/\theta_i\}, \quad i = 1, 2, \cdots
$$
 (6)

By using the U-V method of Robbins (1988), we have the following lemma (see also Vellaisamy and Sharma, 1989).

**Lemma 1** Let  $X_1, X_2, \cdots$  be a sequence of independent random variables with densities *defined in* [\(6\)](#page-3-2). Let  $u_j(\mathbf{x})$  be a real-valued function such that for  $j = 1, 2, \cdots$ ,

*(i)*  $E_{\theta}[|u_j(\mathbf{X})|] < \infty, \quad \forall \theta$ 

(*ii*) 
$$
\int_0^{x_j} u_j(x_1, \dots, x_{j-1}, t, x_{j+1}, \dots) t^{p-1} dt < \infty
$$
,  $\forall x_j > 0$ .

*Then the functions*

$$
\nu_j(\mathbf{X}) = \frac{1}{X_j^{p-1}} \int_0^{X_j} u_j(X_1, \cdots, X_{j-1}, t, X_{j+1}, \cdots) t^{p-1} dt, \ j = 1, 2, \cdots,
$$

*satisfy*

<span id="page-4-0"></span>
$$
E_{\theta} \left[ \nu_j(\mathbf{X}) \right] = E_{\theta} \left[ \theta_j u_j(\mathbf{X}) \right], \ j = 1, 2, \cdots.
$$

The next result obtains the unbiased estimator of  $\theta_{[n]}$ , under the SEL function, for the Gamma $(p, \theta_i)$  distribution with the pdf of  $X_i$  as in [\(6\)](#page-3-2).

**Theorem 1** For the Gamma $(p, \theta_i)$  distribution with the pdf of  $X_i$  as in [\(6\)](#page-3-2), an unbiased *estimator of*  $\theta_{[n]}$ *, under SEL function, which satisfies* [\(4\)](#page-3-3) *with*  $h_{\mathbf{X}}(\boldsymbol{\theta}) = \theta_{[n]}$ *, is* 

$$
V_1(\mathbf{X}) = \frac{U_n}{p} \left( 1 - \left( \frac{U_{n-1}}{U_n} \right)^p \right),\tag{7}
$$

where  $U_n$  is the  $n^{\text{th}}$  upper record value of the sequence  $X_1, X_2, \ldots$ 

**Proof** From [\(1\)](#page-1-0), [\(2\)](#page-1-1) and Lemma 1, an unbiased estimator of  $\theta_{[n]}$ , under SEL function, based on  $X_1, X_2, \ldots$  is given by

$$
V_1(\mathbf{X}) = \sum_{j=n}^{\infty} \nu_j(\mathbf{X}) = \sum_{j=n}^{\infty} \frac{1}{X_j^{p-1}} \int_0^{X_j} t^{p-1} I_j(X_1, X_2, \dots, X_{j-1}, t, X_{j+1}, \dots) dt
$$

where  $I_i(X_1, X_2, \ldots)$  is defined in [\(2\)](#page-1-1). Thus,

$$
V_1(\mathbf{X}) = \sum_{j=n}^{\infty} \frac{I(max\{X_k; X_{T_{n-1}+1} \le k \le j-1\} < U_{n-1} < X_j)}{X_j^{p-1}} \\
\times \left\{ \int_{U_{n-1}}^{X_j} t^{p-1} dt \right\} \\
= \frac{U_n^p - U_{n-1}^p}{p U_n^{p-1}} = \frac{U_n}{p} \left( 1 - \left( \frac{U_{n-1}}{U_n} \right)^p \right).
$$

To find an unbiased estimator of  $\theta_{[n]}$  under the Model 1 with the pdf of  $X_i$  as in [\(5\)](#page-3-1), let  $Y_i = S(X_i) \sim \text{Gamma}(p, \theta_i), i = 1, 2, ..., Y = (Y_1, Y_2, ...)$  and  $y = (y_1, y_2, ...)$ . Then,

 $\Box$ 

by replacing  $X_i$  with  $Y_i = S(X_i)$  in Theorem [1,](#page-4-0) an unbiased estimator of  $\theta_{[n]}$ , under the SEL function, for the general family [\(5\)](#page-3-1), can be obtained as

$$
V_2(\mathbf{X}) = \frac{U_n^S}{p} \left( 1 - \left( \frac{U_{n-1}^S}{U_n^S} \right)^p \right),\tag{8}
$$

where  $U_n^S$  is the n<sup>th</sup> upper record value of the sequence  $Y_1, Y_2, \ldots$ 

For a monotone function  $S(.)$  (available in all of the above examples, except in the normal distribution),  $U_n^S$  can be obtained simply as  $S(U_n)$  for an increasing S and as  $S(L_n)$  for a decreasing S. For example, for the Rayleigh $(\beta_i)$ -distribution, an unbiased estimator for  $\beta_{[n]}$  is

$$
\hat{\beta}_{[n]} = \frac{U_n^2/2}{1} \left( 1 - \left( \frac{U_{n-1}^2/2}{U_n^2/2} \right)^1 \right) = \frac{U_n^2}{2} \left( 1 - \left( \frac{U_{n-1}}{U_n} \right)^2 \right) = \frac{U_n^2 - U_{n-1}^2}{2}.
$$

**Model 2:** For  $X_i$ ,  $i = 1, 2, \dots$ , consider two families of distributions, the first with  $X_i$ having the survival function

<span id="page-5-0"></span>
$$
\bar{F}_{\theta_i}(x) = 1 - F_{\theta_i}(x) = (\bar{G}(x))^{\theta_i^{-1}},
$$
\n(9)

and the second with  $X_i$  having the cdf

<span id="page-5-1"></span>
$$
F_{\theta_i}(x) = (G(x))^{\theta_i^{-1}},\tag{10}
$$

in which  $G(x)$  is a cdf, free of  $\theta_i$ , and  $\overline{G}(x) = 1 - G(x)$ . We assume G to be known. These are called *proportional hazard rate* and *proportional reversed hazard rate* families, or simply  $F^{\alpha}$  models in the context of record values. Some well-known members of the above families are:

- 1. Exponential( $\theta_i$ ), a member of [\(9\)](#page-5-0) with  $\bar{G}(x) = e^{-x}, x > 0;$
- 2. Rayleigh $(\theta_i)$ , a member of [\(9\)](#page-5-0) with  $\bar{G}(x) = e^{-x^2/2}$ ,  $x > 0$ ;
- 3. Beta $(\theta_i^{-1}, 1)$ , a member of [\(10\)](#page-5-1) with  $G(x) = x, 0 < x < 1$ ;
- 4. Pareto $(\theta_i^{-1}, \beta)$ , a member of [\(9\)](#page-5-0) with  $\bar{G}(x) = \beta/x$ ,  $x > \beta$ , and
- 5. Burr $(\alpha, \theta_i^{-1})$ , a member of [\(9\)](#page-5-0) with  $\bar{G}(x) = (1 + x^{\alpha})^{-1}$ ,  $x > 0$ .

By making use of U-V method of Robbins (1988) for the family [\(9\)](#page-5-0), we have the following lemma.

**Lemma 2** Let  $X_1, X_2, \cdots$  be a sequence of independent random variables with survival *function defined in [\(9\)](#page-5-0). Let*  $u_j(\mathbf{x})$  *be a real-valued function such that for*  $j = 1, 2, \cdots$ ,

*(i)*  $E_{\theta}[|u_j(\mathbf{X})|] < \infty$ ,  $\forall \theta$ 

$$
(ii) \int_{-\infty}^{x_j} u_j(x_1, \cdots, x_{j-1}, t, x_{j+1}, \cdots) h(t) dt < \infty, \quad \forall x_j > 0,
$$

*in which*  $h = g/\bar{G}$  *is the hazard function of* G *and* g *is the corresponding pdf of* G. Then *the functions*

$$
\nu_j(\mathbf{X}) = \int_{-\infty}^{X_j} u_j(X_1, \cdots, X_{j-1}, t, X_{j+1}, \cdots) h(t) dt, \ j = 1, 2, \cdots,
$$

*satisfy*

$$
E_{\theta}[\nu_j(\mathbf{X})] = E_{\theta}[\theta_j u_j(\mathbf{X})], \ j = 1, 2, \cdots.
$$

**Proof** For one component problem (i.e., a single random variable  $X_j$ ,  $j \ge 1$ ), let  $\nu(x) =$  $\int_{-\infty}^{x} u(t)h(t) dt$ . Then, we have

$$
\theta_j \mathcal{E}(u(X_j)) = \int_{-\infty}^{+\infty} u(x) [\bar{G}(x)]^{\theta_j^{-1} - 1} g(x) dx
$$
  
\n
$$
= \int_{-\infty}^{+\infty} u(x) \bar{F}_{\theta_j}(x) h(x) dx
$$
  
\n
$$
= \int_{-\infty}^{+\infty} u(x) h(x) \left\{ \int_x^{+\infty} dF_{\theta_j}(y) \right\} dx
$$
  
\n
$$
= \int_{-\infty}^{+\infty} \int_{-\infty}^y u(x) h(x) dx dF_{\theta_j}(y) = \int_{-\infty}^{+\infty} \nu(x) dF_{\theta_j}(x).
$$

For the sequence  $X_1, X_2, \ldots$ , the result follows by a similar calculation.  $\Box$ 

The next result gives the unbiased estimator of  $\theta_{[n]}$ , under SEL function, for the general family [\(9\)](#page-5-0).

**Theorem 2** *Assume* G to be known and let  $H = -\log \bar{G}$  be the cumulative hazard function *of* G. For the general family [\(9\)](#page-5-0), an unbiased estimator of  $\theta_{[n]}$ , under the SEL function, is

$$
V_3(\mathbf{X}) = H(U_n) - H(U_{n-1}).
$$
\n(11)

**Proof** From [\(1\)](#page-1-0), [\(2\)](#page-1-1) and Lemma 2, an unbiased estimator of  $\theta_{[n]}$  is given by

$$
V_3(\mathbf{X}) = \sum_{j=n}^{\infty} \nu_j(\mathbf{X}) = \sum_{j=n}^{\infty} \int_0^{X_j} h(t) I_j(X_1, X_2, \cdots, X_{j-1}, t, X_{j+1}, \cdots) dt
$$
  
= 
$$
\sum_{j=n}^{\infty} \left\{ \int_{U_{n-1}}^{X_j} h(t) dt \right\}
$$
  

$$
\times I(max\{X_k; X_{T_{n-1}+1} \le k \le j-1\} < U_{n-1} < X_j)
$$
  
= 
$$
H(U_n) - H(U_{n-1}).
$$

 $\Box$ 

**Remark 1** Similarly, for the family [\(10\)](#page-5-1), an unbiased estimator for  $\theta_{[n]}$ , under the SEL function, is

$$
V_4(\mathbf{x}) = R(U_n) - R(U_{n-1}),
$$

where  $R = \log G$  is the cumulative reversed hazard function of the known cdf G.

**Remark 2** Note that  $(X_1, X_2, \dots)$  is a complete sufficient statistic for  $(\theta_1, \theta_2, \dots)$ . Hence, the above unbiased estimators of  $\theta_{[n]}$  are indeed UMVU estimators of  $\theta_{[n]}$ .

## 3 Estimation of the Risks

To compare the UMVU estimator with other estimators, we need to compute the risk function of the proposed estimators.

Under the SEL function, the risk of an estimator  $V$  is

$$
\mathcal{R}(V,\theta_{[n]})=\mathcal{E}(V^2)+\mathcal{E}(\theta_{[n]}^2)-2\mathcal{E}(V\theta_{[n]}).
$$

The UMVU estimators obtained in Section 3 are functions of  $(U_n, U_{n-1})$ . Suppose we want to estimate the risk of an estimator of  $\theta_{[n]}$  which depend on **X** only through  $U_n$  and  $U_{n-1}$ , i.e.  $V = V(U_n, U_{n-1})$ . Then, we have the following results, under Models 1 and 2, respectively.

<span id="page-7-0"></span>Theorem 3 *Under the Model 1 and the SEL function, an unbiased estimator of the risk of an estimator*  $V = V(U_n^S, U_{n-1}^S)$  *of*  $\theta_{[n]}$  *is* 

$$
W(U_n^S, U_{n-1}^S) = V^2(U_n^S, U_{n-1}^S) - 2 \frac{\int_{U_{n-1}^S}^{U_n^S} t^{p-1} V(t, U_{n-1}^S) dt}{(U_n^S)^{p-1}} + \frac{(U_n^S)^{p+1} - (U_{n-1}^S)^{p+1} - (p+1) (U_{n-1}^S)^p (U_n^S - U_{n-1}^S)}{p(p+1) (U_n^S)^{p-1}}.
$$

**Proof** From Lemma 1 with  $Y_i = S(X_i)$ , we have

$$
E(\theta_{[n]}^2) = \sum_{j=n}^{\infty} \theta_j^2 E(I_j(\mathbf{Y})) = \sum_{j=n}^{\infty} \theta_j E[\nu_j(\mathbf{Y})]
$$
  
= 
$$
\sum_{j=n}^{\infty} E[\nu_j^*(\mathbf{Y})],
$$

where

$$
\nu_j^*(\mathbf{Y}) = \frac{1}{Y_j^{p-1}} \int_0^{Y_j} s^{p-1} \nu_j(Y_1, \dots, Y_{j-1}, s, Y_{j+1}, \dots) ds
$$
  
= 
$$
\frac{1}{Y_j^{p-1}} \int_0^{Y_j} s^{p-1} \left\{ \frac{1}{s^{p-1}} \int_0^s t^{p-1} I_j(Y_1, \dots, Y_{j-1}, t, Y_{j+1}, \dots) dt \right\} ds.
$$

Therefore

$$
E(\theta_{[n]}^2) = E\left[\sum_{j=n}^{\infty} \frac{I_j(\mathbf{Y})}{Y_j^{p-1}} \int_{U_{n-1}^S}^{Y_j} \int_{U_{n-1}^S}^s t^{p-1} dt ds\right]
$$
  
= 
$$
E\left[\frac{1}{(U_n^S)^{p-1}} \int_{U_{n-1}^S}^{U_n^S} \int_{U_{n-1}^S}^s t^{p-1} dt ds\right]
$$
  
= 
$$
E\left[\frac{(U_n^S)^{p+1} - (U_{n-1}^S)^{p+1} - (p+1) (U_{n-1}^S)^p (U_n^S - U_{n-1}^S)}{p(p+1) (U_n^S)^{p-1}}\right].
$$

Furthermore

$$
E(\theta_{[n]}V(U_n^S, U_{n-1}^S)) = \sum_{j=n}^{\infty} \theta_j E(I_j(\mathbf{Y})V(Y_j, U_{n-1}^S))
$$
  
= 
$$
\sum_{j=n}^{\infty} E\left[\frac{1}{Y_j^{p-1}} \int_0^{Y_j} t^{p-1} V(t, U_{n-1}^S) \times I_j(Y_1, \dots, Y_{j-1}, t, Y_{j+1}, \dots) dt\right]
$$
  
= 
$$
E\left[\frac{1}{(U_n^S)^{p-1}} \int_{U_{n-1}^S}^{U_n^S} t^{p-1} V(t, U_{n-1}^S) dt\right].
$$

Which completes the proof.  $\hfill \Box$ 

An immediate corollary of Theorem [3](#page-7-0) is as follows.

Corollary 1 *Under the Model 1 and the SEL function, an unbiased estimator of the risk of*

$$
V_2 = \frac{U_n^S}{p} \left( 1 - \left( \frac{U_{n-1}^S}{U_n^S} \right)^p \right)
$$

*is*

<span id="page-8-0"></span>
$$
W_2(U_n^S, U_{n-1}^S) = \frac{\left(U_n^S\right)^2}{p^2} \left(1 - \left(\frac{U_{n-1}^S}{U_n^S}\right)^p\right)^2 - \frac{\left(U_n^S\right)^{p+1} - \left(U_{n-1}^S\right)^{p+1} - (p+1)\left(U_{n-1}^S\right)^p \left(U_n^S - U_{n-1}^S\right)}{p(p+1)\left(U_n^S\right)^{p-1}}.
$$

Theorem 4 *For the general family* [\(9\)](#page-5-0)*, and under the SEL function, an unbiased estimator of the risk of an estimator*  $V = V(U_n, U_{n-1})$  *of*  $\theta_{[n]}$  *is* 

$$
W(U_n, U_{n-1}) = V^2(U_n, U_{n-1}) + \frac{(H(U_n) - H(U_{n-1}))^2}{2}
$$

$$
-2 \int_{U_{n-1}}^{U_n} h(t) V(t, U_{n-1})) dt.
$$

Proof From Lemma 2 and using similar argument as in the proof of Theorem [3,](#page-7-0) we have

$$
E(\theta_{[n]}^{2}) = \sum_{j=n}^{\infty} \theta_{j}^{2} E(I_{j}((X)))
$$
  
\n
$$
= \sum_{j=n}^{\infty} \theta_{j} E\left[\int_{-\infty}^{X_{j}} h(t) I_{j}(X_{1},...,X_{j-1},t,X_{j+1},...) dt\right]
$$
  
\n
$$
= \sum_{j=n}^{\infty} E\left[\int_{-\infty}^{X_{j}} h(s) \int_{-\infty}^{s} h(t) I_{j}(X_{1},...,X_{j-1},t,X_{j+1},...) dt ds\right]
$$
  
\n
$$
= E\left[\int_{U_{n-1}}^{U_{n}} h(s) \int_{U_{n-1}}^{s} h(t) dt ds\right]
$$
  
\n
$$
= E\left[\frac{H^{2}(U_{n}) - H^{2}(U_{n-1})}{2} - H(U_{n-1})(H(U_{n}) - H(U_{n-1}))\right]
$$
  
\n
$$
= E\left[\frac{(H(U_{n}) - H(U_{n-1}))^{2}}{2}\right].
$$

Furthermore

$$
E(\theta_{[n]}V(U_n, U_{n-1})) = \sum_{j=n}^{\infty} \theta_j E(I_j(\mathbf{X})V(X_j, U_{n-1}))
$$
  
= 
$$
\sum_{j=n}^{\infty} E\left(\int_0^{X_j} h(t)V(t, U_{n-1}) \times I_j(X_1, \dots, X_{j-1}, t, X_{j+1}, \dots) dt\right)
$$
  
= 
$$
E\left(\int_{U_{n-1}}^{U_n} h(t)V(t, U_{n-1}) dt\right).
$$

This completes the proof.  $\hfill \Box$ 

An immediate corollary of Theorem [4](#page-8-0) is as follows.

Corollary 2 *For the general family* [\(9\)](#page-5-0) *and under the SEL function, (i) an unbiased estimator of the risk of*

<span id="page-9-0"></span>
$$
V_3 = H(U_n) - H(U_{n-1})
$$

*is*

$$
W_3(U_n, U_{n-1}) = \frac{1}{2}(H(U_n) - H(U_{n-1}))^2;
$$

*(ii)* the risk of  $V_3$  *is* 

$$
R(H(U_n) - H(U_{n-1}), \theta_{[n]}) = E(\theta_{[n]}^2).
$$

**Remark 3** The results for the general family [\(10\)](#page-5-1) can be obtained by replacing  $H(\cdot)$  with  $R(\cdot) = \log G(\cdot)$  in Theorem [4](#page-8-0) and Corollary [2.](#page-9-0)

**Remark 4** Since  $(X_1, X_2, \dots)$  is a complete sufficient statistic for  $(\theta_1, \theta_2, \dots)$ , the above unbiased estimators of  $R(V, \theta_{[n]})$  are indeed, UMVU estimators of  $R(V, \theta_{[n]})$ .

The following result presents the distribution of the unbiased estimator in the family [\(9\)](#page-5-0).

Lemma 3 *In the general family* [\(9\)](#page-5-0)*, the following identities hold:*

*(i)* For every  $n \geq 1$  and  $y > 0$ ,

$$
\Pr(H(U_n) - H(U_{n-1}) > y) = \sum_{j=1}^{\infty} e^{-y/\theta_j} \Pr(T_n = j);
$$

*(ii)* For every  $k \geq 2$ ,  $n_1 > n_2 > \cdots$ ,  $n_k \geq 1$  and  $y_1, \ldots, y_k > 0$ ,

$$
\Pr(\bigcap_{i=1}^k \{H(U_{n_i}) - H(U_{n_i-1}) > y_i\}) = \sum_{j_1 < \cdots < j_k} \prod_{i=1}^k e^{-y_i/\theta_{j_i}} \Pr(\bigcap_{i=1}^k \{T_{n_i} = j_i\}).
$$

**Proof** Let  $U_n^* = H(U_n)$  and  $X_n^* = H(X_n)$ ,  $n \ge 1$ . We only prove part (i). Part (ii) is proved in a simillar way. Using the fact that  $X_i^* \sim \text{Exponential}(\theta_i)$  and the lack of memory property of the exponential distribution,

$$
\Pr(U_n^* - U_{n-1}^* > y) = \Pr(X_{T_n}^* - X_{T_{n-1}}^* > y)
$$
  
\n
$$
= \sum_{i < j} \Pr(X_j^* - X_i^* > y | T_n = j, T_{n-1} = i)
$$
  
\n
$$
\times \Pr(T_n = j, T_{n-1} = i)
$$
  
\n
$$
= \sum_{i < j} \Pr(X_j^* - X_i^* > y | X_j^* > X_i^*) \Pr(T_n = j, T_{n-1} = i)
$$
  
\n
$$
= \sum_{i < j} \int \Pr(X_j^* - x > y | X_j^* > x) \Pr(T_n = j, T_{n-1} = i)
$$
  
\n
$$
\times f_{X_i^*}(x) dx
$$
  
\n
$$
= \sum_{i < j} \int \Pr(X_j^* > y) \Pr(T_n = j, T_{n-1} = i) f_{X_i^*}(x) dx
$$
  
\n
$$
= \sum_{j=1}^{\infty} e^{-y/\theta_j} \Pr(T_n = j),
$$

which is the required result.  $\Box$ 

## 4 Inadmissibility of the natural estimator of  $\theta_{[n]}$

For the general family with pdf [\(5\)](#page-3-1), we have

$$
E(S(X_i)/p) = \theta_i.
$$

Thus, a natural estimator for  $\theta_{[n]}$ , for this family of distributions is  $U_n^S/p$ . For the general family with the survival function [\(9\)](#page-5-0), we have

$$
E(H(X_i)) = \theta_i,
$$

which candidates  $H(U_n)$  as a natural estimator of  $\theta_{[n]}$ . So a risk comparison of the natural estimators with UMVUEs of  $\theta_{[n]}$ , for both families of distributions is considered.

The following Corollary of Theorem [4](#page-8-0) states that, under Model 2, the UMVUE dominates the natural estimator.

Corollary 3 *For the general family* [\(9\)](#page-5-0) *and under the SEL function, we have*

$$
R(H(U_n), \theta_{[n]}) > R(H(U_n) - H(U_{n-1}), \theta_{[n]}).
$$

Proof First, we have

$$
E(H(U_{n-1})\theta_{[n]}) = \sum_{j=n}^{\infty} \theta_j E(I_j(\mathbf{X})H(U_{n-1}))
$$
  
= 
$$
E\left(H(U_{n-1})\sum_{j=n}^{\infty} \int_0^{X_j} h(t)I_j(X_1,\ldots,X_{j-1},t,X_{j+1},\ldots) dt\right)
$$
  
= 
$$
E\left(H(U_{n-1})\int_{U_{n-1}}^{U_n} h(t) dt\right)
$$
  
= 
$$
E(H(U_{n-1})(H(U_n) - H(U_{n-1}))).
$$

Consequently,

$$
R(H(U_n), \theta_{[n]}) - R(H(U_n) - H(U_{n-1}), \theta_{[n]})
$$
  
= 2E(H(U\_n)H(U\_{n-1})) - E(H<sup>2</sup>(U\_{n-1}))  
- 2E(H(U\_{n-1})\theta\_{[n]})  
= 2E(H(U\_n)H(U\_{n-1})) - E(H<sup>2</sup>(U\_{n-1}))  
- 2E(H(U\_{n-1})(H(U\_n) - H(U\_{n-1})))  
= E(H<sup>2</sup>(U\_{n-1})) > 0.

This completes the proof.  $\Box$ 

However, under Model 1, no explicit results can be obtained for domination of the UMVUE or the natural estimator with respect to the other, since we have similarly

$$
R(V_2(\mathbf{X}), \theta_{[n]}) - R(U_n^S/p, \theta_{[n]})
$$
  
= 
$$
E\left(\frac{(U_{n-1}^S)^{2p} - 2(U_{n-1}^S)^p (U_n^S)^p + 2p (U_n^S)^{p-1} (U_{n-1}^S)^p (U_n^S - U_{n-1}^S)}{p^2 (U_n^S)^{2p-2}}\right).
$$

To compare the UMVUE and the natural estimator under Model 1, we run a simulation study, which is described in the following section.

#### 4.1 Simulation study

We assume  $X_i \sim \text{Gamma}(p, \theta_i), i = 1, 2, \ldots$  To compare the risks of the UMVUE  $\hat{\theta}^1_{[n]} \,=\, \frac{U_n}{p} \left( 1 - \left( \frac{U_{n-1}}{U_n} \right) \right.$  $\left(\frac{U_{n-1}}{U_n}\right)^p$ , with that of the natural estimator  $\hat{\theta}_{[n]}^2 = \frac{U_n}{p}$ , for  $n = 2, 3, 4$ ,  $p = 0.5, 2$ , we consider three different models for the sequence of parameters as follows:

Model 1 (An stochastic, positive error auto-regressive model):

$$
\theta_i = Z_i \theta_{i-1} + \epsilon_i, \quad \epsilon_i \stackrel{\text{i.i.d}}{\sim} \exp(1), \quad Z_i \stackrel{\text{i.i.d}}{\sim} U(0, 1), \quad i \ge 1, \quad \theta_0 = 0;
$$

Model 2 (An stochastic Geometrically increasing population):

$$
\theta_i = C_i (1 + D_i/10)^{i-1}, \quad C_i, D_i \stackrel{\text{i.i.d}}{\sim} U(0, 1);
$$

Model 3 (White noise model):

$$
\theta_i = 10 + \varepsilon_i, \quad \varepsilon_i \stackrel{\text{i.i.d}}{\sim} N(0, 1).
$$

The simulated bias and risks of the estimators are tabulated in Table [1.](#page-13-0) As one can observe from Table [1,](#page-13-0) the simulated risks of  $\hat{\theta}_{[n]}^1$  are less than those of  $\hat{\theta}_{[n]}^2$ . Also, biases and risks are increasing in n, except the risks of  $\hat{\theta}_{[n]}^1$ , under the white noise Model 3.

### 5 Asymptotic results

From Corollary [2,](#page-9-0) the risk of the UMVUE of  $\theta_{[n]}$  for the general family [\(9\)](#page-5-0),  $V_3 = H(U_n) H(U_{n-1}),$  is

$$
R(V_3, \theta_{[n]}) = \frac{1}{2}E((H(U_n) - H(U_{n-1}))^2)
$$
  
= 
$$
\frac{1}{2}E((U_n^H - U_{n-1}^H)^2),
$$

<span id="page-13-0"></span>

			Model 1		
$\boldsymbol{p}$		$\boldsymbol{n}$	$\overline{2}$	3	4
0.5	$\hat{\theta}^1_{[n]}$	Risk	9.440638	14.75326	18.54895
	$\left  n\right $	<b>Bias</b>	1.524951	4.747217	9.160673
		Risk	23.1851	84.08421	209.7748
$\overline{2}$	$\boldsymbol{n}$	<b>Risk</b>	3.224838	6.856674	10.66222
	n	<b>Bias</b>	0.5978639	1.782032	3.29696
		Risk	3.886525	12.33907	27.96078
			Model 2		
$\,p\,$		$\overline{n}$	$\overline{2}$	3	4
0.5	$\widetilde{\theta^1_{[n]}}$	<b>Risk</b>	2.224561	53.26235	1785.95
	n	<b>Bias</b>	0.7864656	2.342428	6.334353
		Risk	5.501025	94.40079	2499.64
$\overline{2}$	n	Risk	0.5376576	2.314486	19.68881
	n	<b>Bias</b>	0.3038626	0.72345	1.335166
		<b>Risk</b>	0.6209572	2.658157	19.79643
			Model 3		
$\boldsymbol{p}$		$\overline{n}$	$\overline{2}$	3	$\overline{4}$
0.5	$\hat{\theta}^1$ $\lceil n \rceil$	<b>Risk</b>	161.3311	146.8202	125.2359
	$\left  n\right $	<b>Bias</b>	13.682	30.34559	47.98977
		Risk	685.7074	1851.813	3543.839
$\overline{2}$	[n]	Risk	64.93679	74.52687	82.06017
	$\frac{2}{\lvert n \rvert}$	<b>Bias</b>	7.023687	13.47608	19.60645
		Risk	131.9781	297.2568	537.5641

Table 1: Simulated bias and risk of the UMVUE and the natural estimator of  $\theta_{[n]}$  under three different models from gamma distribution for different values of  $n$  and  $p$ .

where  $U_n^H$  is the n<sup>th</sup> upper record value form the sequence  $Y_1, Y_2, \ldots$ , with  $Y_i \sim \text{Exp}(\theta_i)$ .

Hence, asymptotic joint distribution of  $U_n^H$  and  $U_{n-1}^H$  would be useful for computing the risks of the estimators. The following theorem proposes the required asymptotic distribution.

**Theorem 5** Let  $a(n)$  and  $b(n)$  be such that

<span id="page-14-0"></span>
$$
G^n(a(n) + b(n)x) \to \Psi(x),
$$

 $as n \to \infty$  *for all real x, where*  $\Psi$  *is one of the three extreme value cdfs (see Resnick, 1987, p.* 38). Then, for the family [\(9\)](#page-5-0) with  $\sum_{i=1}^{\infty} \theta_i^{-1} = \infty$ , and letting  $U_n^* = \frac{U_n - a(\sum_{i=1}^{T_n} \theta_i^{-1})}{b(\sum_{i=1}^{T_n} \theta_i^{-1})}$  $\frac{b(\sum_{i=1}^{T_n} \theta_i^{-1})}{b(\sum_{i=1}^{T_n} \theta_i^{-1})}$  and  $U_{n-1}^* = \frac{U_{n-1}-a(\sum_{i=1}^{T_n} \theta_i^{-1})}{b(\sum_{i=1}^{T_n} \theta_i^{-1})}$  $\frac{b(\sum_{i=1}^{T_n} \theta_i^{-1})}{b(\sum_{i=1}^{T_n} \theta_i^{-1})}$ , we have, for all  $y > z$ ,  $f_{U_n^*,U_{n-1}^*}(y,z) \to \frac{\psi(y)\psi(z)}{\Psi(y)}$  $\frac{y}{\Psi(y)}, \quad y > z,$ 

 $as n \to \infty$ *, where*  $\psi$  *is the corresponding pdf of*  $\Psi$ *.* 

**Proof.** Letting  $S(i) = \sum_{j=1}^{i} \theta_j^{-1}$ ,  $S^{(2)}(i) = \sum_{j=1}^{i} \theta_j^{-2}$  and  $X_{i:k}$  is the *i*<sup>the</sup> order statistic of  $X_1, \ldots, X_k$ . Using the independence of  $(X_{i-1:i}, X_{i:i})$  and  $T_n$  under the  $F^{\alpha}$  model (Ballerini and Resnick, 1987), we have

$$
f_{U_n, U_{n-1}}(y, z) = \sum_{i=n}^{\infty} f_{X_{i:i}, X_{i-1:i}}(y, z | T_n = i) P(T_n = i)
$$
  
\n
$$
= \sum_{i=n}^{\infty} f_{X_{i:i}, X_{i-1:i}}(y, z) P(T_n = i)
$$
  
\n
$$
= \sum_{i=n}^{\infty} P(T_n = i) \sum_{i_1, i_2 \in \{1, \cdots, i\}; i_1 \neq i_2} [G(z)]^{S(i)-2} \theta_{i_1}^{-1} \theta_{i_2}^{-1} g(y) g(z)
$$
  
\n
$$
= \sum_{i=n}^{\infty} P(T_n = i) [G(z)]^{S(i)-2} (S(i))^2 g(y) g(z) \left[ 1 - \frac{S^{(2)}(i)}{(S(i))^2} \right]
$$
  
\n
$$
= \mathbf{E} \left[ [G(z)]^{S(T_n)-2} (S(T_n))^2 g(y) g(z) \left[ 1 - \frac{S^{(2)}(T_n)}{(S(T_n))^2} \right] \right].
$$

Consequently, since g satisfies the Von-Mises conditions (see Resnick, 1987) and  $\left[1 - \frac{S^{(2)}(n)}{(S(n))^2}\right]$  $rac{S^{(2)}(n)}{(S(n))^2}$   $\rightarrow$ 1, as  $n \to \infty$ , we have

$$
f_{U_n^*, U_{n-1}^*}(y, z) = \mathbb{E}\left[ [G(b(S(Tn))z + a(S(Tn)))]^{S(T_n)-2}(S(T_n)b(S(Tn)))^2 \times g(b(S(Tn))y + a(S(Tn))) \right] \times g(b(S(Tn))z + a(S(Tn)))\left[1 - \frac{S^{(2)}(T_n)}{(S(T_n))^2}\right] \longrightarrow \Psi(z)\frac{\psi(y)}{\Psi(y)}\frac{\psi(z)}{\Psi(z)}.
$$

Thus, the proof is complete.  $\Box$ 

When G is standard exponential distribution, we have  $a(n) = \log n$ ,  $b(n) = 1$  and  $\Psi(x) = \exp\{-\exp(-x)\}.$  Therefore, letting  $U_n^* = U_n - \log(\sum_{i=1}^{T_n} \theta_i^{-1})$  and  $U_{n-1}^* = U_{n-1}$  $log(\sum_{i=1}^{T_n} \theta_i^{-1}),$  as  $n \to \infty$ , we have

$$
f_{U_n^*, U_{n-1}^*}(y, z) \to \exp(-(z + y)) \exp\{-\exp(-z)\}, \quad y > z,
$$

and consequently for each y and z, as  $n \to \infty$ , we have

 $F_{U_n^*, U_{n-1}^*}(y, z) \to \exp\{-e^{-\min(y, z)}\} [1 + I(y > z)(e^{-z} - e^{-y})].$ 

However,  $U_n^*$  and  $U_{n-1}^*$  depend on the unknown  $\theta$ . The following result solves this problem using the fact that under the  $F^{\alpha}$  model,  $n^{-1/2}(\log(S(T_n)) - n)$  converges in law to the standard normal distribution (see Nevzerov, 1995).

<span id="page-15-0"></span>**Theorem 6** *Under the family* [\(9\)](#page-5-0) *with*  $G(x) = 1 - \exp(-x)$ ,  $x > 0$ , *with the assumptions of Theorem [5,](#page-14-0) and letting*  $T_n^* = n^{-1/2}(\log(S(T_n)) - n)$ *, as*  $n \to \infty$ *, for fixed* y*,* z and t*,* we *have*

$$
F_{U_n^*, U_{n-1}^*, T_n^*}(y, z, t) \to \Phi(t) \exp\{-e^{-\min(y, z)}\}[1 + I(y > z)(e^{-z} - e^{-y})],
$$

*where* Φ *is the cdf of the standard normal distribution.*

Proof As in the proof of Theorem [5,](#page-14-0) we have

$$
F_{U_n^*, U_{n-1}^*, T_n^*}(y, z, t) = \sum_{i=n}^{\infty} F_{X_{i:i} - \log(S(i)), X_{i-1:i} - \log(S(i))}(y, z)
$$
  
 
$$
\times I(n^{-1/2}(\log(S(i)) - n) < t)P(T_n = i)
$$
  
\n
$$
= \sum_{i=n}^{\infty} \exp\{-e^{-\min(y, z)} + O(1/S(i))\}
$$
  
\n
$$
\times [1 + I(y > z)(e^{-z} - e^{-y})]
$$
  
\n
$$
\times I(n^{-1/2}(\log(S(i)) - n) < t)P(T_n = i)
$$
  
\n
$$
\longrightarrow \exp\{-e^{-\min(y, z)}\}[1 + I(y > z)(e^{-z} - e^{-y})]\Phi(t),
$$

as  $n \to \infty$ , which is the required result.  $\Box$ 

By Theorem [6,](#page-15-0) we have

$$
P\left(\frac{U_n - n}{\sqrt{n}} \le x, \frac{U_{n-1} - n}{\sqrt{n}} \le y, \right)
$$
  
= 
$$
P\left(\frac{U_n - \log S(T_n) + \log S(T_n) - n}{\sqrt{n}} \le x, \frac{U_{n-1} - \log S(T_n) + \log S(T_n) - n}{\sqrt{n}} \le y\right)
$$
  

$$
\longrightarrow \Phi(\min\{x, y\}) = \min\{\Phi(x), \Phi(y)\},
$$
 (12)

<span id="page-15-1"></span>

as  $n \to \infty$ , which is the upper Fréchet Höeffding bound; see, e.g., Fréchet (1951) or Nelsen (1999, p. 9). The following Corollary, presents an approximate formula for the risk of UMVUE of  $\theta_{[n]}$ , for the family [\(9\)](#page-5-0).

Corollary 4 *For the family* [\(9\)](#page-5-0)*, under the assumptions of Theorem [5,](#page-14-0) we have*

$$
R(H(U_n) - H(U_{n-1}), \theta_{[n]}) = o(n), \quad \text{as} \quad n \to \infty.
$$

Proof From  $(12)$  and by Höeffding's theorem,

$$
\lim_{n \to \infty} \text{Cor}\left(\frac{U_{n-1}^H - n}{\sqrt{n}}, \frac{U_n^H - n}{\sqrt{n}}\right) = \lim_{n \to \infty} \text{Cov}\left(\frac{U_{n-1}^H - n}{\sqrt{n}}, \frac{U_n^H - n}{\sqrt{n}}\right)
$$

$$
= \int \int \min(\Phi(x), \Phi(y)) - \Phi(x)\Phi(y) \, dx \, dy.
$$

The above double integral can be simplified by algebraic manipulations as

$$
1 + \int x \phi^{2}(x) dx - \int \phi(x) (1 - 2\Phi(x)) dx = 1,
$$

in which  $\phi$  is the pdf of the standard normal distribution. Thus, we have

$$
\frac{1}{n}R(U_n^H - U_{n-1}^H, \theta_{[n]}) = \frac{1}{2}E\left(\frac{U_n^H - n - (U_{n-1}^H - n)}{\sqrt{n}}\right)^2 \to 0,
$$
  
as  $n \to \infty$ .

## 6 Rainfall data: an illustrative example

In this section, we utilize the data set which represents the records of the amount of annual (January 1-December 31) rainfall in inches recorded at Los Angeles Civic Center LACC during the 100-year period from 1890 until 1989, presented by Arnold et al. [1998, p. 180].

A member of the  $F^{\alpha}$  model (Model 2) with survival function as in [\(9\)](#page-5-0), that is the Rayleigh distribution with cdf

$$
F(x) = 1 - \exp\left\{ \frac{-(x-4)^{1.9}}{113.23} \right\}, \ x > 4,\tag{13}
$$

is well-fitted to the data. The p-value for two-sample Kolmogorov-Smirnov test is 0.3333. Figure [1](#page-17-0) shows the empirical distribution function of the rainfall data and the cdf in [\(13\)](#page-16-0). Thus, we take

<span id="page-16-0"></span>
$$
H(x) = (x - 4)^{1.9},
$$

to be the known cumulative hazard rate function of the base distribution  $G(x) = 1 \exp\left\{-(x-4)^{1.9}\right\}, x > 4.$ 

Suppose that the only observations are the sequence of upper record values as follows:



Figure 1: Empirical cdf of the rainfall data.

<span id="page-17-0"></span>

We consider two hypotheses:

 $H_0$ : (Stationary model)  $X_1, X_2, \ldots \stackrel{\text{iid}}{\sim} F_\theta(x) = 1 - \exp\left\{ \frac{-(x-4)^{1.9}}{\theta} \right\}$  $\frac{(-4)^{1.9}}{\theta}$ ,  $x > 4$ ;  $H_1:$  (Non-stationary model)  $X_i \sim F_{\theta_i}(x) = 1 - \exp\left\{\frac{-(x-4)^{1.9}}{\theta_i}\right\}$  $\left\{\frac{-4}{\theta_i}\right\}, x > 4, i = 1, 2, \dots$  and Xs are independent.

Under  $H_0, \ \theta_{[n]} = \theta, \ n = 1, 2, \ldots$ , with probability 1. Hence,  $\hat{\theta}_{[n]} = \frac{H(U_n)}{n} = \frac{(U_n - 4)^{1.9}}{n}$  $\frac{-4)^{n}}{n}$  is the UMVUE of  $\theta_{[n]} = \theta$ . Also,  $R(\hat{\theta}_{[n]}, \theta_{[n]}) = \text{Var}\left(\frac{H(U_n)}{n}\right)$  $\left(\frac{U_n}{n}\right) = \frac{\theta^2}{n}$  $\frac{\partial^2}{\partial n}$ , with unbiased estimator  $\hat{\mathcal{R}}(\hat{\theta}_{[n]}, \theta_{[n]}) = \frac{[H(U_n)]^2}{n^2(n+1)} = \frac{(U_n-4)^{3.8}}{n^2(n+1)}.$ 

Under  $H_1$ ,  $\hat{\theta}_{[n]} = H(U_n) - H(U_{n-1}) = (U_n - 4)^{1.9} - (U_{n-1} - 4)^{1.9}$  and the unbiased estimator of its risk is  $\hat{R}(\hat{\theta}_{[n]}, \theta_{[n]}) = \frac{(H(U_n) - H(U_{n-1}))^2}{2} = \frac{((U_n - 4)^{1.9} - (U_{n-1} - 4)^{1.9})^2}{2}$  $\frac{(U_{n-1}-4)^{-\alpha}$ )<sup>-</sup><br>2

Figure [2](#page-18-0) shows the values of  $\hat{\theta}_{[n]}$  and their corresponding 3- $\sigma$  region

$$
\left(\max\left\{0,\hat{\theta}_{[n]}-1.5\sqrt{\hat{\mathcal{R}}(\hat{\theta}_{[n]},\theta_{[n]})}\right\},\hat{\theta}_{[n]}+1.5\sqrt{\hat{\mathcal{R}}(\hat{\theta}_{[n]},\theta_{[n]})}\right),\right)
$$

under  $H_0$  and  $H_1$ .



<span id="page-18-0"></span>Figure 2: Estimates path (solid line) and 3- $\sigma$  regions (upper and lower dashed lines) of  $\theta_{[n]}$ , under the stationary (straight lines) and non-stationary (zigzag lines) assumptions, for the rainfall data.

To test  $H_0$  against  $H_1$  using the record sequence we propose the scale invariant test statistic

$$
T = \frac{1}{n-1} \sum_{i=2}^{n} \left( \frac{\hat{\theta}_{[i]}}{\hat{\theta}_{[i-1]}} - 1 \right)^2.
$$
 (14)

Since, under  $H_0$ , all  $\hat{\theta}_{[i]}$ s are equal, the null hypothesis is rejected for large values of T.

We use the fact that under  $H_0$ , the random variables  $H(U_n) - H(U_{n-1}), n \geq 2$  are iid exponential, to deduce that under  $H_0$ ,

<span id="page-18-1"></span>
$$
T \stackrel{\text{d}}{=} \frac{1}{n-1} \sum_{i=2}^{n} \left( \frac{Z_i}{Z_{i-1}} - 1 \right)^2,\tag{15}
$$

where  $\stackrel{\text{d}}{=}$  stands for the identically distributed and  $Z_1, \ldots, Z_n \stackrel{\text{iid}}{\sim} Exp(1)$ .

Deriving the exact distribution of  $T$  is far from reach. However, one can estimate the distribution quantiles of  $T$  using a Monté Carlo simulation study.

To generate random variables identically distributed as  $T$ , one may generate an iid sample form standard exponential, namely,  $Z_1, \ldots, Z_n$ , and return  $T = \frac{1}{n-1}$  $\frac{1}{n-1}\sum_{i=2}^n\left(\frac{Z_i}{Z_i}\right)$  $\frac{Z_i}{Z_{i-1}}-1\Big)^2.$ 

Table [2](#page-19-0) presents the simulated values of  $\alpha$ -critical values of T,  $t_n(\alpha)$ , for  $n = 2, \ldots, 10$ , and  $\alpha = 0.01, 0.025, 0.05, 0.1$ , which are generated using R.14.1 package with  $10^5$  iterations.

<span id="page-19-0"></span>

		$\alpha$		
n	0.01	0.025	0.05	0.1
$\overline{2}$	8645.63	1368.24	326.02	64.61
3	19003.73	3113.96	723.25	164.76
$\overline{4}$	27929.12	4681.26	1093.01	264.36
5	37018.72	6343.56	1529.97	355.73
6	49769.98	7707.69	2007.57	456.78
7	64315.21	9211.87	2388.29	563.19
8	70630.56	10801.06	2698.59	655.51
9	73372.31	11655.77	3131.44	747.15
10	92847.93	13727.53	3500.69	883.22

Table 2: The critical values of the test statistic [\(15\)](#page-18-1)

The hypothesis  $H_0$  is rejected at level  $\alpha$  as

$$
T>t_n(\alpha).
$$

For the rainfall data we obtain  $T = 279.14$ , which is less than  $t_8(0.05) = 2698.59$ . Therefore,  $H_0$  is not rejected in favor of  $H_1$  at level  $\alpha = 0.05$ .

## 7 Concluding remarks

The problem of estimating parameters of the dynamically selected populations can be extended to the Bayesian context. Moreover, the problem of unbiased estimation of the selected parameters under other loss functions is of interest. The distributional models which are not members of studied families can be studied separately, specially the discrete distribution. Another problem is to find the two stage (conditionally) unbiased estimators of the parameters of the dynamically selected populations. These problems are treated in an upcoming work, to appear in subsequent papers.

## Acknowledgements

The authors thank the anonymous referee for his/her useful comments and suggestions on an earlier version of this manuscript which resulted in this improved version.

## References

[1] Ahmadi J. and Balakrishnan N. (2010). Prediction of order statistics and record values from two independent sequences, *Statistics*, 44, 417 – 430.

- [2] Ahmadi J. and Balakrishnan N. (2013). On the nearness of record values to order statistics from Pitman's measure of closeness, *Metrika*, 76, 521 – 541.
- [3] Amini M. and Balakrishnan N. (2013). Nonparametric Meta-Analysis of Independent Samples of Records. *Computational Statistics & Data Analysis*, 66, 70 – 81.
- [4] Amini M. and Balakrishnan N. (2015). Pooled Parametric Inference for minimal repair systems. *Computational Statistics*, DOI: 10.1007/s00180-014-0552-8.
- [5] Arnold B. C., Balakrishnan N. and Nagaraja H. N. (1998). *Records*, John Wiley & Sons, New York.
- [6] Baklizi A. (2008). Likelihood and Bayesian estimation of using lower record values from the generalized exponential distribution, *Computational Statistics & Data Analysis*, 52, 3468 – 3473.
- [7] Ballerini R. and Resnick S.I. (1987). Embedding sequences of successive maxima in extremal processes with applications. *Journal of Applied Probability*, 24, 827 – 837.
- [8] Dahiya, R. C. (1974). Estimation of the Mean of the Selected Population, *Journal of the American Statistical Association*, 69, 226 – 230.
- [9] Doostparast, M. and Emadi M. (2013). Evidential inference and optimal sample size determination on the basis of record values and record times under random sampling scheme, *Statistical Methods & Applications*, doi:10.1007/s10260-012-0228-x.
- [10] Fréchet, M. (1951). Sur les tableaux de corrélation dont les marges sont données. *Annales de l'Universit´e de Lyon Section A.* (3), 14, 53 – 77.
- [11] Gibbons, J. D., Olkin, I. and Sobel, M. (1977). *Selecting and Ordering Populations. A New Statistical Methodology.* New York: John Wiley and Sons.
- [12] Kumar S. and Kar A. (2001). Estimation quantiles of a selected exponential population. *Statistics & Probability Letters*, 52, 9 - 19.
- [13] Kumar S. and Gangopadhyay A.K. (2005). Estimation parameters of a selected Pareto population. *Statistical Methodology*, 2, 121 - 130.
- [14] Kumar S., Mahapatra A.K. and Vellaisamy P. (2009). Reliability estimation of the selected exponential populations. *Statistics & Probability Letters*, 79, 1372 - 1377.
- [15] Kundu C., Nanda A. K. and Hu T. (2009). A note on reversed hazard rate of order statistics and record values, *Journal of Statistical Planning and Inference*, 139, 1257 – 1265.
- [16] Lehmann, E.L. (1951). A general concept of unbiasedness. *Annals of Mathematical Statistics*, 22, 578-592.
- [17] Misra N., Vander Meulen E.C. and Branden K.V. (2006a). On estimating the scale parameter of the selected gamma population under the scale invariant squared error loss function. *Journal of Computational and Applied Mathematics*, 186, 268 - 282.
- [18] Misra N. Vander Meulen E.C. and Brandan K.V. (2006b). On some inadmissibility results for the scale parameters of selected gamma populations. *Journal of Statistical Planning and Inference*, 136, 2340 - 2351.
- [19] Nelsen, R. B. (1999). *An Introduction to Copulas.* Lecture Notes in Statistics. Springer, New York.
- [20] Nematollahi, N. and Motammed-Shariati, F. (2012). Estimation of the parameter of the selected uniform population under the entropy loss function, *Journal of Statistical Planning and Inference*, 142, 2190 – 2202.
- [21] Nevzorov V.B. (1985). On record times and inter-record times for sequences of nonidentically distributed random variables. *Zap. Nauehn. Sere. LOMI.*, 142, 109 – 118.
- [22] Nevzorov V.B. (1986). Two characterizations using records. *Lecture Notes in Mathematics*, 1233, 79 – 85.
- [23] Nevzorov V. (1995). Asymptotic distributions of records in non-stationary schemes. *Journal of Statistical Planning and Inference*, 45, 261 – 273.
- [24] Pfeifer D. (1989). Extremal processes, secretary problems and the 1/e law, *Journal of Applied Propagability*, 27, 722 – 733.
- [25] Pfeifer D. (1991). Some remarks on Nevzorov's record model. *Advances in Applied Propagability*, 23, 823 – 834.
- [26] Psarrakos, G. and Navarro, J. (2013). Generalized cumulative residual entropy and record values, *Metrika*, 76,623 – 640.
- [27] Raqab M. Z. and Ahmadi J. (2012). Pitman closeness of record values from two sequences to population quantiles, *Journal of Statistical Planning and Inference*, 142, 855 – 862.
- [28] Resnick, S. (1987). *Extreme Values, Regular Variation, and Point Processes.*, Springer-Verlag, New York.
- [29] Robbins H. (1988). The U.V methods of estimation. In: Gupta, S.S., Berger, J.O. (Eds.), Statistical Decision Theory and Related Topics IV, vol.1. Springer - Verlag, NewYork, pp. 265 - 270.
- [30] Salehi M. Ahmadi J. and Balakrishnan, N. (2013). Prediction of order statistics and record values based on ordered ranked set sampling, *Journal of Statistical Computation and Simulation*, doi = 10.1080/00949655.2013.803194.
- [31] Sarkadi, K. (1967). Estimation after selection. *Studia Scientarium Mathematicarum Hungarica*, 2, 341–350.
- [32] Sill, M. W. and Sampson, A. R. (2007). Extension of a Two-Stage Conditionally Unbiased Estimator of the Selected Population to the Bivariate Normal Case, *Communications in Statistics - Theory and Methods*, 36, 801 – 813.
- [33] Vellaisamy P. and Sharma D. (1989). A note on the estimation of the mean of the selected gamma population. *Communications in Statistics - Theory and Methods*, 18,  $555 - 560.$
- [34] Zarezadeh S. and Asadi M. (2010). Results on residual Rényi entropy of order statistics and record values, *Information Sciences*, 180, 4195 – 4206.