Spin nematics, valence-bond solids and spin liquids in SO(N) quantum spin models on the triangular lattice

Ribhu K. Kaul

Department of Physics & Astronomy, University of Kentucky, Lexington, KY-40506-0055

We introduce a simple model of SO(N) spins with two-site interactions which is amenable to quantum Monte-Carlo studies without a sign problem on *non-bipartite* lattices. We present numerical results for this model on the two-dimensional triangular lattice where we find evidence for a spin nematic at small N, a valence-bond solid (VBS) at large N and a quantum spin liquid at intermediate N. By the introduction of a sign-free four-site interaction we uncover a rich phase diagram with evidence for both first-order and exotic continuous phase transitions.

The destruction of magnetic order by quantum fluctuations in spin systems is frequently invoked as a route to exotic condensed matter physics such as spin liquid phases and novel quantum critical points [1–3]. The most commonly studied spin Hamiltonians have symmetries of the groups SO(3) and SU(2) which describe the rotational symmetry of 3-dimensional space. Motivated both by theoretical and experimental [4] interest, spin models with larger-N symmetries have been introduced, e.g. extensions of SU(2) to SU(N) [5–8] or Sp(N) [9].

The extension of SO(3) to SO(N) is an independant large-N enlargement of symmetry, with its own physical motivations [10]. While there have been many studies of SO(N) spin models in one dimension [11–13], our understanding of their ground states and quantum phase transitions in higher dimension is in its infancy. To this end, we introduce here a simple SO(N) spin model that surprisingly is sign free on *any* non-bipartite lattice. This model provides us with a new setting in which the destruction of magnetic order can be studied in higher dimensions using *unbiased* methods. As an example of interest, we present the results of a detailed study of the phase diagram of the our SO(N) anti-ferromagnet on the two-dimensional triangular lattice.

Models. – Consider a triangular lattice, each site of which has a Hilbert state of N states, we will denote the state of site j as $|\alpha\rangle_j$ $(1 \le \alpha \le N)$. Define the N(N-1)/2 generators of SO(N) on site i as $\hat{L}_i^{\alpha\beta}$ with $\alpha < \beta$; they will be chosen in the fundamental representation on all sites: $\hat{L}_j^{\alpha\beta}|\gamma\rangle_j = i\delta_{\beta\gamma}|\alpha\rangle_j - i\delta_{\alpha\gamma}|\beta\rangle_j$. Now consider the following SO(N) [14] symmetric lattice model for $N \ge 3$,

$$\hat{H}_J = -\frac{J}{N^2 - 2N} \sum_{\langle ij \rangle} (\hat{L}_i \cdot \hat{L}_j)^2, \qquad (1)$$

where the "·" implies a summation over the N(N-1)/2generators and $\langle ij \rangle$ is the set of nearest neighbors. To see that \hat{H}_J does not suffer from the sign problem, define a "singlet" state on a bond, $|S_{ij}\rangle \equiv \frac{1}{\sqrt{N}} \sum_{\alpha} |\alpha \alpha\rangle_{ij}$ and the singlet projector $\hat{P}_{ij} = |S_{ij}\rangle\langle S_{ij}|$. Using these operators and ignoring a constant shift we find the simple form [15],

$$\hat{H}_J = -J \sum_{\langle ij \rangle} \hat{P}_{ij}.$$
 (2)

We make four observations: First, it is possible to create an SO(N) spin singlet with only two spins for all N (in contrast to SU(N) where N fundamental spins are required to create a singlet); Second Eq. (1) being a sum of projectors on this two-site singlet is the simplest SO(N) coupling, despite it being a biquadratic interaction in the generators $\hat{L}^{\alpha\beta}$; Third, since the singlet has a positive expansion, \hat{H}_J is Marshall positive on any lattice; Fourth, on bipartite lattices \hat{H}_J is equivalent to the familiar SU(N) anti-ferromagnet [6], i.e. the obvious SO(N) of Eq. (1) is enlarged to an SU(N) symmetry. Since the bipartite SU(N) case has been studied in great detail in past work on various lattices [7, 16–23], we shall concern ourselves here with the non-bipartite SO(N) case which is relatively unexplored.

Phases of \hat{H}_J : Starting at N = 3, Eq. (1) becomes $\hat{H} = -\frac{J}{3} \sum_{\langle ij \rangle} \left(\vec{S}_i \cdot \vec{S}_j \right)^2$ with \vec{S} the familiar S = 1 representation of angular momentum. Previous numerical work has shown that this triangular lattice S = 1 biquadratic model [24, 25] has an SO(3) symmetry breaking "spin nematic" magnetic ground state (we shall denote this phase by SN). The ground state of \hat{H}_J for N > 3 has not been studied in the past.

In the large-N limit, analogous to previous work for SU(N) anti-ferromagnets on bipartite lattices [26], the ground state is infinitely degenerate and consists of dimer coverings where each dimer is in $|S_{ij}\rangle$. At leading order in 1/N, \hat{H}_J introduces off-diagonal moves which re-arrange parallel dimers around a plaquette, mapping \hat{H}_J at large-N to a quantum dimer model on the triangular lattice with only a kinetic term,

$$\hat{H}_{\text{QDM}} = -t \sum_{\text{plaq}} \left\{ \left(|\overleftarrow{}\rangle_i \langle \overleftarrow{} \rangle_i + h.c. \right) \right\}$$
(3)

where the sum on plaquettes includes all closed loops of length four on the triangular lattice. The ground state of this model has been found in previous analytic [27] and numerical work [28] to be a $\sqrt{12} \times \sqrt{12}$ valence bond solid (VBS), breaking the lattice translation symmetry. We thus expect that at large but finite values of N, \hat{H}_J should restore its SO(N) symmetry and enter this same VBS state.



FIG. 1. Equal time structure factors for SN order $[S_{\rm SN}(\mathbf{k})]$, and susceptibility for VBS order $[\chi_{\rm VBS}(\mathbf{k})]$ shown for N = 10and N = 14, for the \hat{H}_J model, Eq. (1) with L = 48. The Bragg peaks for SN (VBS) weaken (sharpen) with increasing N. The cartoon of the Brillouin zones shows the location of the ordering vectors of both order parameters. Quantitative finite size scaling of these orders is shown in Fig. 2.

Since \hat{H}_J has SN order for N = 3 and is expected to have a non-magnetic VBS at large-N, it is interesting to ask what the nature of the transition at which SN magnetism is destroyed. The answer to this question is unclear based on current theoretical ideas and is best settled by unbiased numerical simulations. Exploiting that H_J has no sign problem we study it as a function of N on $L \times L$ lattices at temperture β by unbiased stochastic series expansion [29] quantum Monte Carlo simulations, with a previously described algorithm [24]. The SN state is described by the matrix order parameter $\hat{Q}_{\alpha\beta} = |\alpha\rangle\langle\beta| - \frac{1}{N}$. The static structure factor, $S_{\rm SN}(\mathbf{k}) = \frac{1}{N_{\rm site}} \sum_{ij} e^{i\mathbf{k}\cdot(\mathbf{r}_i - \mathbf{r}_j)} \langle \hat{Q}_{\alpha\alpha}(i)\hat{Q}_{\alpha\alpha}(j) \rangle$ is used to detect SN order. For the VBS order, we construct the \mathbf{k} dependent susceptibility of dimerdimer correlation functions in the usual way from imaginary time-displaced operators: $\chi_{\text{VBS}}(\mathbf{k}) = \frac{1}{N_{\text{site}}} \sum_{ij} e^{i\mathbf{k}\cdot(\mathbf{r}_i - \mathbf{r}_j)} \frac{1}{\beta} \int d\tau \langle \hat{P}_{\mathbf{r}_i,\mathbf{r}_i + \hat{x}}(\tau) \hat{P}_{\mathbf{r}_j,\mathbf{r}_j + \hat{x}}(0) \rangle.$ Throughout this paper we have fixed $\beta = L$ for our finite size scaling [15].

As shown in Fig. 1, a peak in $S_{\rm SN}(\mathbf{k})$ is found at the Γ point. Comparing the data at N = 10 and N = 14, already qualitatively it is possible to see the peak in $S_{\rm SN}(\mathbf{k})$ softens as N is increased. In contrast $\chi_{\rm VBS}(\mathbf{k})$ develops sharp peaks at the X and M points as N is increased. These are precisely the momenta at which previous numerical studies of the triangular lattice quantum dimer model Eq. (3) have observed Bragg peaks [28], validating the large-N mapping to Eq. (3) made earlier. To detect at which N, the magnetic order is destroyed and the VBS order first sets in, we study the ratio, $R_{\rm SN} = 1 - \frac{S_{\rm SN}(\Gamma + \mathbf{a}2\pi/L)}{S_{\rm SN}(\Gamma)}$ (where $\mathbf{a} \equiv \mathbf{x} - \mathbf{y}/\sqrt{3}$) as a function of L. $R_{\rm SN}$ must diverge in a phase in which the Bragg peak height scales with volume and becomes infinitely sharp. On the other hand it must go to zero in



FIG. 2. Crossing plots of the ratios $R_{\rm SN}$ and $R_{\rm VBS}$ as a function of the discrete variable N for the \hat{H}_J model. It is seen that spin nematic order is present for $N \leq 10$. VBS order on the other hand is present for N > 12. N = 12 appears to be on the verge of VBS order. Interestingly, N = 11 has no SN or VBS order. In the text, we present evidence that this phase is a QSL. The inset in the upper panel shows $R_{\rm SN}$ scales to zero at N = 11, despite non-monotonic behavior at intermediate L.

a phase in which the correlation length is finite and the height and width of the Bragg peak saturate with system size. At a critical point standard finite size scaling arguments imply that the ratio, $R_{\rm SN}$ becomes volume independent. All of these facts together imply a crossing in this quantity for different L. Fig. 2 shows the $R_{\rm SN}$ and $R_{\rm VBS}$ ratios (an analogous quantity constructed for the VBS order from $\chi_{\text{VBS}}(\mathbf{k})$ close to the **M**-point) as a function of the discrete variable N for different L. The data for $R_{\rm SN}$ shows that the magnetic order is present for $N \leq 10$. The $R_{\rm VBS}$ data shows that the long-range VBS order is present for N > 12. From Fig. 2 we find that N = 12 is on the verge of developing VBS order; from the system sizes accesible we are unable to reliably conclude whether N = 12 has long range VBS order or not from our study. However, taken together the data show definitively that N = 11 has neither VBS nor SN order. As we shall substantiate below, at N = 11, \hat{H}_J is a quantum spin-liquid (QSL).

J-Q models: In order to clarify the global phase diagram of SO(N) anti-ferromagnets and access the quantum phase transitions between the SN, VBS and QSL phases found in \hat{H}_J , it is of interest to find an interaction that can tune between these phases at *fixed N*. In order to be meaningful, the new coupling must preserve all the symmetries of \hat{H}_J . To this end, we introduce and study a generalization of the four-site Q term of SU(2) spins [30],

$$\hat{H}_Q = -Q \sum_{\langle ijkl \rangle} \left(\hat{P}_{ij} \hat{P}_{kl} + \hat{P}_{il} \hat{P}_{jk} \right) \tag{4}$$

where the sum includes elementary plaquettes of length four on the triangular lattice (with periodic boundary



FIG. 3. Phase diagram of \hat{H}_{JQ} [Eqs. (2,4)] for different values of N. The left panel shows the phase diagram for small N, where a first order SN-VBS transition is found for $6 \leq N \leq 9$, (see Fig. 4). As N is increased we find the first order transition weakens. The right panel shows how an intermediate QSL phase emerges for N = 10 and N = 11. Transitions from the QSL to both SN and VBS phases are continuous on the large systems studied, see Fig. 5.

conditions on an $L \times L$ system there are $3L^2$ such plaquettes). For a fixed-N, \hat{H}_Q provides a tuning parameter which preserve both the internal and lattice symmetries of \hat{H}_J and hence allows us to study the generic phase diagram of SO(N) magnets. A summary of the phase diagram of \hat{H}_{JQ} in the N-Q/J plane is in Fig. 3: The Qinteraction destroys the SN order and gives way to VBS order only for $N \ge 6$. We have found evidence for direct first-order SN-VBS transitions for $6 \le N < 10$ and exotic continuous SN-VBS transitions for N = 10 and N = 11.

As an example of our observed first-order behavior we present in Fig. 4, our study of the N = 7 QMC data for the spin stiffness $\rho_s \equiv \langle W_x^2 \rangle / L$ (where W_x is the winding number of the spin world lines), which acts as a sensitive order parameter for the SN phase, and the VBS order parameter $O_{\rm VBS}^2 \equiv \chi_{\rm VBS}(\mathbf{M})/N_{\rm site}$. Clear evidence for a direct first order SN-VBS transition at N = 7 is found.

The nature of the transition changes at N = 10, where evidence for two phase transitions is found. As shown in Fig. 5 the SN order vanishes at a Q/J smaller than the value at which VBS order develops. Although the difference is small for N = 10, it is significant. The data for N = 11 in Fig. 5 shows that the SN and VBS orders do not vanish at the same point. In fact $R_{\rm SN}$ indicates that the SN order has vanished already at Q/J = 0, consistent with our previous analysis of \hat{H}_J . As illustrated by the dashed and solid lines in Fig. 3, the appearance of the QSL phase is consistent with a global phase diagram for the SO(N) magnets.

QSL phase and criticality: We have identified the ground state between SN and VBS as a QSL, since it does not show evidence for any Landau-order. Were the intermediate phase characterized by a conventional order parameter, we would have expected strong first order



FIG. 4. First-order SN to VBS transition in H_{JQ} at N = 7. The upper panel shows the VBS order parameter and the stiffness as a function for Q/J for different L indicating a direct SN-VBS transition. The lower panel shows MC histories (and histograms in the inset) at Q/J = 1.26, providing clear evidence that the SN-VBS transition at N = 7 is direct and first-order.

transitions of the kind between SN and VBS (see Fig. 4), instead we find continuous transitions.

There are field theoretic reasons to expect a QSL on quantum disordering a spin nematic. The long-distance description of our SO(N) models is given by a RP^{N-1} theory (in contrast to the \mathbb{CP}^{N-1} description of $\mathrm{SU}(N)$ models [31]), which can be described as N real matter fields coupled to a Z_2 gauge field. Such a theory is expected to host three phases [32], a symmetry breaking phase in which the matter condenses (which we identify in our spin model as the SN), a stable phase in which the matter gets a gap and the Z_2 gauge theory is deconfined (identified here as the QSL) and a phase in which matter is gapped and the Z_2 is confined (identified here as the VBS). Thus, the SN-QSL critical point should be in the universality class of $O(N)^*$ critical point [3]. The QSL-VBS phase transition should be in the same universality class as the critical point between these identical phases in the quantum dimer model since the magnetic fluctuations are gapped in both the QSL and VBS phases. A previous analysis of this phase transition has predicted an $O(4)^*$ phase transition [27], where the VBS order parameter is identified with a bilinear of the primary field.

A detailed study of the critical phenomena at N = 10and N = 11 is clearly beyond the scope of the current manuscript. We shall be satisfied here with a brief analysis: At the QSL-VBS critical point, we are able to carry out reasonable data collapses [15] at both N = 10 and



Crossings of $R_{\rm SN}$ (above) and $R_{\rm VBS}$ (below) sig-FIG. 5. naling the location of the onset of long-range SN and VBS orders at N = 10 (left) and N = 11 (right). At N = 10, $R_{\rm SN}$ and $R_{\rm VBS}$ cross at close but significantly different couplings, $Q_c = 0.100(5)$ and $Q_c = 0.117(2)$ respectively. At $N = 11, R_{\rm SN}$ appears to have crossed at Q/J < 0 (we cannot study this region because of the sign problem), whereas $R_{\rm VBS}$ crosses at $Q_c = 0.042(3)$. From the location of the crossings, for both N = 10 and N = 11, we can infer an intermediate phase which is neither SN nor VBS, as shown in Fig. 3(b). We present arguments that this phase is a QSL. No direct evidence for first order behavior is found at either of the transitions, though a weakly first order SN-QSL cannot be ruled out. The QSL-VBS transitions shows good scaling behavior with unconventional critical exponents.

N = 11 for $O_{\rm VBS}^2$ (for both X and M ordering vectors, see Fig. 1) and R_{VBS} , where we find, $\eta_{\text{VBS}} = 1.3(2)$ and $\nu_{\rm VBS} = 0.65(20)$ for the anomalous dimension of $O_{\rm VBS}$. The unusually large value of $\eta_{\rm VBS}$ is a direct consequence of fractionalization in the intermediate QSL phase and is often regarded as a smoking gun diagnostic of exotic critical points (see e.g., [33]). More quantitatively, our critical exponents are in rough agreement with the best estimate of $\eta = 1.375(5)$ of the bilinear field and $\nu = 0.7525(10)$ in the O(4) model [34]. We note that the values for $\eta_{\rm VBS}$ and $\nu_{\rm VBS}$ agree within the quoted errors for N = 10 and N = 11. Taken together, this bolsters the case that the intermediate QSL phase has Z_2 fractionalization, albeit more work is needed for a definitive identification. Unfortunately, the SN-QSL transition, observed only at N = 10, has large corrections to scaling and we are unable to reliably determine its critical exponents or determine whether it is a weakly first order transition (no direct evidence for a first-order transition has been found of the type shown for the N = 7 case).

In summary, we have introduced a new family of signfree SO(N) spin models, which can be regarded as nonbipartite generalizations of their popular SU(N) cousins. The triangular lattice model which we have studied thoroughly here hosts a spin nematic, a VBS with a large unit cell, a quantum spin liquid phase and unusual quantum critical points. The absence in the SO(N) models of a direct continuous "deconfined quantum critical point" [33] is in striking contrast to previous simulations of the related bipartite SU(N) models [8, 23]. We have offered a plausible field theoretic scenario that naturally explains this difference. It is interesting that the absence (presence) of a QSL in bipartite SU(N) (non-bipartite SO(N)) spin models seems to track the absence or presence of this phase in the kind of quantum dimer models that our model maps to at large-N [35].

While the study in this paper has focussed on the triangular lattice, our family of models, Eq. (2,4) may be constructed sign free on any two or three dimensional non-bipartite lattice. Because of the larger degree of frustration, the kagome system may provide a wider swath of the QSL phase and hence could possibly allow a more detailed study of this phase, even if the phase diagram is of the same form found here. Exploring the phase diagram and quantum phase transitions of the three dimensional pyrochlore system is an exciting open direction for future work.

The author is grateful to J. Chalker, T. Lang, M. Levin, R. Mong, G. Murthy, A. Nahum, A. Sandvik, T. Senthil and M. Zaletel for helpful discussions. This research was supported in part by NSF DMR-1056536.

- L. Balents, Nature 464, 199 (2010), URL http://dx.doi. org/10.1038/nature08917.
- [2] S. Sachdev, Quantum Phase Transitions (Cambridge University Press, 1999).
- [3] C. Xu, Int. J. Mod. Phys. B p. 1230007 (2012).
- [4] C. Wu, Physics **3**, 92 (2010).
- [5] B. Sutherland, Phys. Rev. B 12, 3795 (1975), URL http: //link.aps.org/doi/10.1103/PhysRevB.12.3795.
- [6] I. Affleck, Phys. Rev. Lett. 54, 966 (1985), URL http: //link.aps.org/doi/10.1103/PhysRevLett.54.966.
- [7] N. Read and S. Sachdev, Phys. Rev. Lett. 62, 1694 (1989), URL http://link.aps.org/doi/10.1103/PhysRevLett. 62.1694.
- [8] R. K. Kaul, R. G. Melko, and A. W. Sandvik, Annu. Rev. Cond. Matt. Phys 4, 179 (2013), URL http://www.annualreviews.org/doi/abs/10.1146/ annurev-conmatphys-030212-184215.
- [9] N. Read and S. Sachdev, Phys. Rev. Lett. 66, 1773 (1991), URL http://link.aps.org/doi/10.1103/PhysRevLett. 66.1773.
- [10] E. Demler, W. Hanke, and S.-C. Zhang, Rev. Mod. Phys. 76, 909 (2004), URL http://link.aps.org/doi/ 10.1103/RevModPhys.76.909.
- [11] H.-H. Tu, G.-M. Zhang, and T. Xiang, Phys. Rev. B 78, 094404 (2008), URL http://link.aps.org/doi/10. 1103/PhysRevB.78.094404.
- [12] F. Alet, S. Capponi, H. Nonne, P. Lecheminant, and I. P. McCulloch, Phys. Rev. B 83, 060407 (2011), URL http: //link.aps.org/doi/10.1103/PhysRevB.83.060407.

- [13] K. Okunishi and K. Harada, Phys. Rev. B 89, 134422 (2014), URL http://link.aps.org/doi/10. 1103/PhysRevB.89.134422.
- [14] Strictly speaking the symmetry of our model is an SO(N) for odd-N and an $O(N)/\mathbb{Z}_2$ for even N. This point is discussed further in the supplementary materials.
- [15] Please refer to supplementary materials for more details on the model and the numerical simulations.
- [16] G. Santoro, S. Sorella, L. Guidoni, A. Parola, and E. Tosatti, Phys. Rev. Lett. 83, 3065 (1999), URL http: //link.aps.org/doi/10.1103/PhysRevLett.83.3065.
- [17] K. Harada, N. Kawashima, and M. Troyer, Phys. Rev. Lett. 90, 117203 (2003), URL http://journals.aps. org/prl/abstract/10.1103/PhysRevLett.90.117203.
- [18] K. S. D. Beach, F. Alet, M. Mambrini, and S. Capponi, Phys. Rev. B 80, 184401 (2009), URL http://link.aps. org/doi/10.1103/PhysRevB.80.184401.
- [19] J. Lou, A. Sandvik, and N. Kawashima, Phys. Rev. B 80, 180414 (2009), URL http://link.aps.org/doi/10. 1103/PhysRevB.80.180414.
- [20] R. K. Kaul, Phys. Rev. B 85, 180411 (2012), URL http: //link.aps.org/doi/10.1103/PhysRevB.85.180411.
- [21] R. K. Kaul and A. W. Sandvik, Phys. Rev. Lett. 108, 137201 (2012), URL http://link.aps.org/doi/10. 1103/PhysRevLett.108.137201.
- [22] M. S. Block and R. K. Kaul, Phys. Rev. B 86, 134408 (2012), URL http://link.aps.org/doi/10. 1103/PhysRevB.86.134408.
- [23] M. S. Block, R. G. Melko, and R. K. Kaul, Phys. Rev. Lett. 111, 137202 (2013), URL http://link.aps.org/ doi/10.1103/PhysRevLett.111.137202.
- [24] R. K. Kaul, Phys. Rev. B 86, 104411 (2012), URL http: //link.aps.org/doi/10.1103/PhysRevB.86.104411.
- [25] A. Laeuchli, F. Mila, and K. Penc, Phys. Rev. Lett. 97 (2006).
- [26] N. Read and S. Sachdev, Nuclear Physics B 316, 609 (1989), ISSN 0550-3213, URL http://www.sciencedirect.com/science/article/ pii/0550321389900618.
- [27] R. Moessner and S. L. Sondhi, Phys. Rev. B 63, 224401 (2001), URL http://link.aps.org/doi/10. 1103/PhysRevB.63.224401.
- [28] A. Ralko, M. Ferrero, F. Becca, D. Ivanov, and F. Mila, Phys. Rev. B 74, 134301 (2006), URL http://link.aps. org/doi/10.1103/PhysRevB.74.134301.
- [29] A. W. Sandvik, AIP Conf. Proc. 1297, 135 (2010), URL http://scitation.aip.org/content/aip/proceeding/ aipcp/10.1063/1.3518900.
- [30] A. W. Sandvik, Phys. Rev. Lett. 98, 227202 (2007), URL http://link.aps.org/doi/10.1103/PhysRevLett. 98.227202.
- [31] N. Read and S. Sachdev, Phys. Rev. B 42, 4568 (1990).
- [32] P. E. Lammert, D. S. Rokhsar, and J. Toner, Phys. Rev. Lett. 70, 1650 (1993), URL http://link.aps.org/doi/ 10.1103/PhysRevLett.70.1650.
- [33] T. Senthil, A. Vishwanath, L. Balents, S. Sachdev, and M. P. A. Fisher, Science 303, 1490 (2004), URL http://www.sciencemag.org/content/303/5663/ 1490.abstract.
- [34] H. Ballesteros, L. Fernndez, V. Martn-Mayor, and A. M. Sudupe, Physics Letters B 387, 125 (1996), ISSN 0370-2693, URL http://www.sciencedirect.com/ science/article/pii/0370269396009847.
- [35] R. Moessner and S. L. Sondhi, Phys. Rev. Lett. 86,

1881 (2001), URL http://link.aps.org/doi/10.1103/ PhysRevLett.86.1881.

Model and Symmetries

Here we provide some additional details of the models introduced in Eq. (1) and Eq. (2).

Mapping between Eqs. (1) and (2)

To see the connection between the two Hamiltonians Eq. (1) and Eq. (2). We consider two SO(N) spins. We can combine them into three representations: a singlet (S), symmetric (χ) and anti-symmetric (Φ) representations of dimensions: $1, \frac{N^2}{2} + \frac{N}{2} - 1$ and $\frac{N^2}{2} - \frac{N}{2}$. Now construct projectors on these representations, P_S, P_{χ} and P_{Φ} . Clearly $P_S + P_{\chi} + P_{\Phi} = 1$ and $P_S^2 = P_S, P_{\chi}^2 = P_{\chi}, P_{\Phi}^2 = P_{\Phi}$. It is straightforward to show that, $(L_i \cdot L_j) = -(N-1)P_S - P_{\chi} + P_{\Phi}$ by explicitly acting on the symmetrized wave-functions. From this it follows that $(L_i \cdot L_j)^2 = (N-1)^2 P_S + P_{\chi} + P_{\Phi}$. From which it follows that $(L_i \cdot L_j)^2 = ((N-1)^2 - 1) P_S + 1$, which proves as claimed that for $N \geq 3$, Eq. (1) and Eq. (2) are equivalent up to a constant.

N=2

Although not studied in this manuscript, for the sake of completeness, we discuss our model at N = 2. Even though Eq. (1) is trivial for N = 2 (since squaring the only SO(2) generator is just an identity operator), Eq. (2) is a well defined non-trivial model. Identifying the two colors with \uparrow and \downarrow spins, Eq. (2) becomes $\hat{H} = \sum_{\langle ij \rangle} -J(S_i^x S_j^x + S_i^z S_j^z) + JS_i^y S_j^y$. Previous work on the triangular lattice N = 2 model has found clear evidence for SO(2) symmetry breaking superfluid order [1–3].

Symmetries

We now discuss the symmetries of the model Eq. (2). We begin by observing that this model is invariant under uniform O(N) rotations where we multiply each basis state by an orthogonal matrix (one that satisfies $O^T O =$ 1), since this leaves the singlet state invariant, *i.e.*

$$\sum_{\alpha} |\alpha\alpha\rangle \to \sum_{\alpha} O_{\alpha\gamma} O_{\alpha\eta} |\gamma\eta\rangle = \sum_{\alpha} |\alpha\alpha\rangle.$$
 (5)

However we should identify rotations that only differ by changing all the local basis states by the same phase (in this case a sign). Here it becomes necessary to distinguish between even and odd N. This is because the matrix -1 has determinant 1 for even N and -1 for odd N. Thus

size	N	Q	$\beta_{\rm QMC}$	E_{ex}	$E_{\rm QMC}$
2×2	4	0	16	-1.5	-1.49997(3)
2×2	4	1	16	-4.5	-4.5000(1)
2×2	5	0	16	-1.4	-1.40000(5)
2×2	5	1	16	-4.2	-4.2001(1)
2×3	3	2	16	-6.0657499233	-6.0656(1)
3×3	2	0	16	-1.7026987262	-1.70269(2)
3×3	2	1	16	-3.7290340614	-3.72900(3)
3×3	2	2	16	-5.7639923092	-5.76402(5)

TABLE I. Test comparisons of ground state energies from exact diagonalization and average energies from finite-T QMC studies of the SO(N) model introduced here. Note that J = 1 always. The energies reported here are per site and on triangular lattices with periodic boundary conditions such that there are always $3L^2$ bonds and $3L^2$ plaquettes in Eqs. (2,4). This causes some terms to be appear more than once for the 2×2 and 2×3 systems.

for odd-N the symmetry is simply SO(N), since the rest of O(N) is obtained from SO(N) by multiplying by -1. For even-N however SO(N) has pairs of elements that cause the same basis transformation up to a sign, *e.g.* 1 and -1. On the other hand unlike the case of odd-N, the O(N) matrices with determinant -1 are independent symmetries, so the symmetry realized for even-N is an $\frac{O(N)}{Z_2}$.

Symmetry on Bipartite Lattices

On bipartite lattices the orthogonal rotation symmetry, Eq. (5) gets extended to a unitary symmetry (with $U^{\dagger}U = 1$) so long as the singlet is defined between sites on opposite sub-lattices, and A sub-lattice spins are rotated by U and B sub-lattice spins are rotated by U^* ,

$$\sum_{\alpha} |\alpha\alpha\rangle \to \sum_{\alpha} U^*_{\alpha\gamma} U_{\alpha\eta} |\gamma\eta\rangle = \sum_{\alpha} |\alpha\alpha\rangle.$$
(6)

Since a uniform phase change of the all the states locally does not have physical consequences, the model is said to have an SU(N) symmetry, as has been discussed and extensively studied previously in such models, see *e.g.* Ref. [4] for a review.

Ground state theorems

Marshall's sign theorem guarantees that the ground state of $H_{JQ} = H_J + H_Q$ is an SO(N) singlet. In addition, on the triangular lattice, which is the focus of our study here, there is no simple translationally invariant covering of two-site singlets, leading us to suspect that a generalization of the SU(2) square lattice Lieb-Schultz



FIG. 6. Estimation of optimal β/L ratio for our simulations by comparison of the fluctuations of the temporal $(\langle W_{\tau}^2 \rangle)$ and spatial $(\langle W_x^2 \rangle)$ winding numbers . Data shown is for \hat{H}_J at N = 10.

Mattis (LSM) theorem [5] applies to H_{JQ} on this lattice, *i.e.* in the thermodynamic limit there must be a degeneracy in the ground state, so that a simple gapped paramagnet is not possible – either a symmetry is broken or the ground state is exotic. A rigorous proof of this intuitive assertion is expected to be at least as technical as the proof for the bipartite N = 2 case [5] and is beyond the scope of this work. As we saw above, on a one-dimensional chain, which is bipartite, our model is equivalent to the SU(N) model studied by Affleck [6] and is hence expected to have an LSM degeneracy.

Numerical Simulations

QMC energy tests

Here we provide the results of some QMC tests on small lattices for the total energy per spin of our models, $\hat{H}_J + H_Q$, Eqs. (2,4) on the triangular lattice, for completeness and future comparisons.

Choice of $\beta = L$

In Fig. 6 we show the dependence of the fluctuations of the temporal and spatial winding numbers on the "aspect ratio" of our simulation cell. β is a measure of the extent of the imaginary time and L is an estimate for the linear spatial extent. We study the fluctuations of the temporal and spatial winding numbers as the ratio β/L is varied for two different sizes, L = 32 and L = 48 at N = 10 in the model \hat{H}_J . We find that both quantities are balanced at a value of β/L which is of the order of one (close to

1.42) and that the crossing point does not move much with system size. Thus for simplicity we have chosen $\beta/L = 1$ throughout the paper.



FIG. 7. Finite size scaling of spin nematic order parameter and spin stiffness for the model \hat{H}_J , in Eq. (1) with $10 \leq N \leq$ 14. Shown on the left is the square of the spin nematic order parameter, $O_{\rm SN}^2$ (the height of the Bragg peak in Fig. 1), and on the right is the spin stiffness, ρ_s , plotted as a function of 1/L. The data confirms that the system is magnetically ordered for $N \leq 10$ and non-magnetic for $N \geq 11$.



FIG. 8. Finite size scaling of VBS order parameter for the model \hat{H}_J , in Eq. (1) with $10 \leq N \leq 14$. For $N \geq 13$ we encounter difficulties in equilibrating the system for sizes larger than L = 48 due to formation of long-range VBS order. Rather than extrapolating O_{VBS}^2 , we study the crossing of the ratio R_{VBS} , which provides a reliable way to detect the onset of long-range VBS order on moderate system sizes [see Fig. 2].

Extrapolation of order parameters

The simplest estimate for long range order is to study whether the height of the Bragg peak per unit volume extrapolates to a finite quantity in the thermodynamic limit. Unfortunately, this method becomes increasingly unreliable when the measured order is weak, e.g., close to a critical point. In such cases, results from extrapolations will depend on the form of the extrapolation used. A thorough discussion of these difficulties in quantum spin systems may be found in the literature [7]. It is for this reason that we prefer to work with the R ratios defined in the text. The disadvantage is that we do not know the order parameter in the thermodynamic limit, but the advantage is we can calculate the critical coupling reliably by studying the crossing of the R ratio. For completeness we present here the data required for extrapolation of both SN and VBS order parameters for H_{I} . To test quantitatively for long range order we study the scaling of the height of the peak in $S_{\rm SN}(\mathbf{k})$, $O_{\rm SN}^2 \equiv S_{\rm M}({\bf k}=0)/N_{\rm site}$ and the spin stiffness ρ_s on finite size systems with $N_{\text{site}} = L \times L$. Both quantities are expected to be finite in the M state and zero when the O(N) symmetry is restored. Fig. 7 shows finite size data for both quantities for different values of N. From these plots we conclude that the M symmetry is broken up to N=10 and is restored for $N\geq 11,$ because $O_{\rm SN}^2$ scales to zero for these N. This behavior is mirrored in ρ_s , albeit for intermediate L there is some non-monotonic behavior for N = 11. This is consistent with our conclusions in the main text made from the analysis of $R_{\rm SN}$

Finite size scaling for the VBS order parameter is shown in Fig. 8. Notice for the cases where there is VBS order (N = 13, 14) we only have data for $L \leq 48$. For system sizes larger than this we face serious equilibration issues with QMC as is expected, since the simulation gets locked into a symmetry broken VBS state. The plot serves to illustrate te ambiguity faced by making direct extrapolations. On the other hand, a study of the Rratios shown in Fig. 2 provides a more clear cut way to locate the critical point.

QSL-VBS and SN-QSL phase transitions

Here we present some details of the study of the both the QSL-VBS (N = 10, 11) and SN-QSL (N = 10) phase transitions found in our model.

We obtain critical exponents at the QSL-VBS critical point by attempting a data collapse, see Fig. 9,10. We use the standard finite size scaling ansatz for the order parameter and the crossing ratio,

$$\langle O_{\rm VBS}^2 \rangle = L^{-(1+\eta_{\rm VBS})} \mathcal{F}_O(g L^{1/\nu_{\rm VBS}}) \tag{7}$$

$$R_{\rm VBS} = \mathcal{F}_R(gL^{1/\nu_{\rm VBS}}) \tag{8}$$



FIG. 9. Examples of our data collapses at QSL-VBS critical point at N = 10 shown here with the best parameters determined individually for each observable. The parameters used are $Q_c = 0.1170, 0.1187, 0.1171, 1/\nu_{VBS} = 2.171, 2.152, 1.733$ and $\eta_{VBS} = 1.414, 1.106, -.$



FIG. 10. Examples of our data collapses at QSL-VBS critical point at N = 11 shown here with the best parameters determined individually for each observable. The parameters used are $Q_c = 0.04330, 0.04182, 0.03917, 1/\nu_{\rm VBS} = 1.902, 1.858, 1.244$ and $\eta_{\rm VBS} = 1.383, 1.133, -$.

where $g = (Q - Q_c)/J$. We continue to work with $\beta = L$ as discussed. No attempt is made to make use of corrections to this leading scaling behavior. Our main objective is to determine the universal number η_{VBS} for the QSL-VBS transition for N = 10 and N = 11. We find acceptable collapses for our data sets over a wide range of ν_{VBS} . On the other hand, the estimate for η_{VBS} is relatively stable over our various fits. The values and errors of the critical exponents quoted in the main text are based on the variation observed by using different data



FIG. 11. Crossing of L and L/2 for various dimensionless quantities for N = 10. The dashed line is a quantity ($R_{\rm VBS}$) whose crossing locates the VBS transitions and the solid lines are for two independent quantities (ρ_s and $R_{\rm SN}$) that locate the SN transition. The blue and red semi-circle shows the range of critical couplings based on the extrapolation of this data. Note that (1) there is clear evidence for an intermediate phase. (2) the errors in the SN transition are significantly larger than those for the VBS transition.

sets. A higher precision study should be possible with access to more accurate data and larger system sizes. In order to carry out the collapse numerically, we make use of a recently developed Bayesian approach to scaling [8]. We note that difficulties in obtaining accurate values of the critical exponents at exotic transitions in quantum spin models is a well-documented difficulty [9].

Another quantum phase transition takes place between SN and QSL. In our model this transition appears only at N = 10. In Fig. 11 we study the drift of various crossing

quantities at the critical points. Presumably the significant drift for the crossing at the SN-QSL transition are due to corrections to scaling. We have looked for signs of first order behavior as we found for smaller-N and not found them here, though the possibility of a very weak first order transition cannot be ruled out. The corrections to scaling hamper efforts to extract critical exponents at this phase transition. In contrast the QSL-VBS transition shows a reasonably converged crossing point with a nice scaling regime, where the crossing points do not depend significantly on L.

- R. G. Melko, A. Paramekanti, A. A. Burkov, A. Vishwanath, D. N. Sheng, and L. Balents, Phys. Rev. Lett. 95, 127207 (2005), URL http://link.aps.org/doi/10. 1103/PhysRevLett.95.127207.
- [2] D. Heidarian and K. Damle, Phys. Rev. Lett. 95, 127206 (2005), URL http://link.aps.org/doi/10. 1103/PhysRevLett.95.127206.
- [3] S. Wessel and M. Troyer, Phys. Rev. Lett. 95, 127205 (2005), URL http://link.aps.org/doi/10. 1103/PhysRevLett.95.127205.
- [4] R. K. Kaul, R. G. Melko, and A. W. Sandvik, Annu. Rev. Cond. Matt. Phys 4, 179 (2013), URL http://www.annualreviews.org/doi/abs/10.1146/ annurev-conmatphys-030212-184215.
- [5] M. B. Hastings, Phys. Rev. B 69, 104431 (2004).
- [6] I. Affleck, Phys. Rev. Lett. 54, 966 (1985), URL http: //link.aps.org/doi/10.1103/PhysRevLett.54.966.
- [7] A. W. Sandvik, Phys. Rev. B 85, 134407 (2012), URL http://link.aps.org/doi/10.1103/PhysRevB.85. 134407.
- [8] K. Harada, Phys. Rev. E 84, 056704 (2011), URL http: //link.aps.org/doi/10.1103/PhysRevE.84.056704.
- [9] K. Harada, T. Suzuki, T. Okubo, H. Matsuo, J. Lou, H. Watanabe, S. Todo, and N. Kawashima, Phys. Rev. B 88, 220408 (2013), URL http://link.aps.org/doi/10. 1103/PhysRevB.88.220408.