

The distribution of work performed on a NIS junction

Jaime E. Santos

Email: jaie@nanotech.dtu.dk

Center for Nanostructured Graphene (CNG), Department of Micro and Nanotechnology, Technical University of Denmark, DK-2800 Kongens Lyngby, Denmark

Centro de Física and Departamento de Física, Universidade do Minho, P-4710-057 Braga, Portugal

Pedro Ribeiro

Email: ribeiro.pedro@gmail.com

Russian Quantum Center, Novaya Street 100 A, Skolkovo, Moscow Area, 143025 Russia

Stefan Kirchner

Email: stefan.kirchner@correlated-matter.com

Center for Correlated Matter, Zhejiang University, Hangzhou, Zhejiang 310058, China
Max Planck Institute for the Physics of Complex Systems, Nöthnitzer Str. 38, D-01187 Dresden, Germany

Max Planck Institute for Chemical Physics of Solids, Nöthnitzer Str. 40, D-01187 Dresden, Germany

Abstract. We propose an experimental setup to measure the work performed in a normal-metal/insulator/superconducting (NIS) junction, subjected to a voltage change and in contact with a thermal bath. We compute the performed work and argue that the associated heat release can be measured experimentally. Our results are based on an equivalence between the dynamics of the NIS junction and that of an assembly of two-level systems subjected to a circularly polarised field, for which we can determine the work-characteristic function exactly. The average heat dissipated by the NIS junction, as well as its fluctuations, are determined. From the work characteristic function, we also compute the work probability-distribution and show that it does not have a Gaussian character. Our results allow for a direct experimental test of the Crooks-Tasaki fluctuation relation.

PACS numbers: 05.70.Ln,74.45.+c,74.50.+r

1. Introduction

Fluctuation relations are central to our present understanding of statistical mechanics. Their long and distinguished history goes back to at least the studies by Callen and Welton [1], Green [2] and Kubo [3], which were inspired by the works of Einstein on the Brownian movement [4] and of Johnson [5] and Nyquist [6] on noise in electrical circuits.

The derivation by Jarzynski of a rather general non-equilibrium work relation [7], linking the average of the exponential of the work being performed on a system with the equilibrium free energy difference between initial and final equilibrium states of the system, is of particular interest. Subsequently, Crooks [8, 9] obtained the Jarzynski equality from a relation between the probability of a given amount of work being performed on a system and the probability that the system performs the same amount of work on its surroundings if the time-reversed protocol is undertaken. Such heightened interest in non-equilibrium work relations has not only been fuelled by novel theoretical advances in out-of-equilibrium dynamics, but also by experimental progress in preparing and probing non-equilibrium evolution (see [10, 11, 12, 13, 14] for reviews on the subject).

In the present paper, we propose a relatively simple yet realistic experiment based on a proposal by Crooks [15], which would allow for a direct test of the Crooks-Tasaki fluctuation relation. The experimental system we consider consists of a normal-metal/insulator/superconducting (NIS) junction between a superconductor and a normal metal, which is initially short-circuited and is subjected to a given voltage protocol (see below). We establish that for the proposed protocol, the full work distribution-function has non-Gaussian character with a non-standard decay exponent. We also compute the first two moments of such distribution, which are equal to the average heat dissipated and its variance.

In order to obtain a closed expression for the characteristic function of the work distribution, we derive an equivalence between the dynamics of an NIS junction and the one of an assembly of two-level systems subjected to a circularly polarized field. Using this mapping, the work distribution is determined. For realistic parameters of experimentally available NIS junctions as *e.g.* those of Ref. [16], our findings show that at cryogenic temperatures, even the tiny amounts of heat released will result in sizeable volume changes for a probe connected to the junction.

The structure of the paper is as follows: Sect. 2 places the recent developments briefly outlined above into their proper historical context and reviews the theoretical tools necessary to study work-fluctuation relations at the quantum level. In Sect. 3, we introduce the Hamiltonian of the NIS junction and describe in detail the work protocol applied to the system. In Sect. 4, we use results that are derived in Appendix A and Appendix B to compute both the average work dissipated as well as its fluctuations and the full work distribution. Sect. 5 contains the main result of our paper, showing that the heat that is dissipated can be measured by determining the volume change of a probe that absorbs it. We provide estimates of the relative volume change, which should

be measurable using capacitive methods [17, 18]. We make contact with experimentally measurable quantities and give order-of-magnitude estimates for realistic NIS junctions. Finally, in Sect. 6, we present our conclusions. For increased readability, most technical details have been relegated to appendices.

2. Perspective and previous works

In this section, we give a brief historical overview of the evolution of the subject of non-equilibrium fluctuation theorems. Besides the early contributions already mentioned above, a statement of a general fluctuation theorem involving the free energy of a gas of hard-spheres, interacting at large distances via the Lennard-Jones potential, can be found in the work of Zwanzig [19]. A subsequent important development was the work of Bochkov and Kuzovlev [20, 21], which also concerned the relation of probabilities between a given work protocol and its time reversed version, but which adopted a different perspective to that of Jarzynski and Crooks, namely the former authors considered the generalised force that performs work on the system as being external to the dynamical description of the system, rather than internal. Another significant contribution was the work of Evans and Searles (see [22] and references therein), which considers the relation between the probability for a dynamical trajectory characterised by a certain value of a dissipation function and its time-reversed counterpart. Regarding developments concerning the validity and extension of the Jarzynski-Crooks relations at the classical level, see the experimental works [23, 24, 25, 26, 27, 28, 29, 30, 31, 32] and the theoretical ones [33, 34, 35, 36, 37, 38], the above list not being exhaustive.

A quantum generalisation of the Crooks equality was obtained by Tasaki [39] and also by Kurchan [40] (for the case of cyclic protocols). A version of the Jarzynski equality valid for quantum systems was also derived by Yukawa [41], albeit treating work at a quantum level as an observable, a view that was shown not to be correct [42]. Such a view led in the past to a debate concerning the validity of such fluctuation relations at the quantum level [43, 44, 45]. The extension of the Jarzynski equality to isolated systems, using the definition of the work distribution function of Tasaki and Kurchan, is due to Mukamel [46]. A quantum generalisation of the work of Bochkov and Kuzovlev referred above, as well as a clarification of the relation of their work to that of Jarzynski and Crooks, can be found in a paper by Campisi et al. [47]. Extensions of non-equilibrium work relations to isolated systems (micro-canonical fluctuation theorems) are given in [48, 49], whereas work relations valid for arbitrary open quantum systems were obtained in [50]. The last reference shows the validity of such theorems for systems that can arbitrarily exchange heat with the bath while a work protocol is being applied to them (see also below). Other theoretical developments concerning quantum systems can be found in the references [51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71], where again the list is not exhaustive.

At the experimental level, several proposals to demonstrate the validity of the




Figure 1. (a) NIS junction. (b) Schematic representation of the voltage protocol considered

Crooks-Tasaki relation in the quantum domain have been presented, including the use of a series of projective measurements [72], the measurement of this relation in the optical spectra of systems subjected to weak quenches [73], or the use of qubit interferometry [74, 75, 76]. This latter proposal has been successfully implemented, see [77]. An example of the experimental confirmation of a generalised version of the Jarzynski equality in the quantum domain was discussed in [70]. Finally, a discussion of possible solid state experiments performed on quantum heat engines is given in [71].

The above mentioned proposal by Crooks [15] relies on an indirect method of observation namely, the measurement of the heat dissipated by a quantum system after a cyclic work protocol has been performed on the system. A specific set-up to measure the heat released by a circuit including a resistor and a Cooper-pair box, based on the temperature increase of a local probe, was already presented in [78], where the Cooper-pair box acts as a single two-level system. The present proposal has the additional advantage that the heat released scales with the contact area between the normal-metal and the superconducting films.

The work probability distribution, $P_\beta(W, \tau)$, is defined, for a class of work protocols where the system is decoupled from the thermal bath during the application of the protocol (see below), of duration τ , as [39]

$$P_\beta(W, \tau) = \frac{1}{\mathcal{Z}(\beta)} \sum_{m,n} e^{-\beta E_m^0} |\langle n, \tau | \hat{U}(\tau, 0) | m, 0 \rangle|^2 \delta(W - E_n^\tau + E_m^0), \quad (1)$$

where $|m, 0\rangle$ are the eigenstates of the system's Hamiltonian at the beginning of the protocol, $\hat{H}(0)$ (which is given for the NIS junction by (5) at $t = 0$, see below), and E_m^0 are the corresponding eigenvalues, and where $|n, \tau\rangle$ are the eigenstates of the system's Hamiltonian at the end of the protocol, $\hat{H}(\tau)$ (which is given by the same equation at $t = \tau$), with E_n^τ being the corresponding eigenvalues, with $\beta = 1/(k_B T)$ and $\mathcal{Z}(\beta) = \text{Tr}(e^{-\beta \hat{H}(0)})$. The operator $\hat{U}(\tau, 0)$ is the time-evolution operator of the system during the protocol, i.e. in the interval $[0, \tau]$, which can be written as a time-ordered exponential of the full Hamiltonian of the system, given again for the NIS junction by equation (5).

The work characteristic-function of the NIS junction is the Fourier transform of

equation (1), and is given by

$$\mathcal{G}_\beta(v, \tau) = \text{Tr} \left[\widehat{U}^\dagger(\tau, 0) e^{i\widehat{H}(\tau)v} \widehat{U}(\tau, 0) e^{-i\widehat{H}(0)v} \widehat{\rho}(\beta) \right], \quad (2)$$

where $\widehat{\rho}(\beta) = e^{-\beta\widehat{H}(0)}/\mathcal{Z}(\beta)$ is the density matrix of the junction before the application of the protocol.

The unitary evolution of the system in the interval $[0, \tau]$ reflects the work protocol's restriction mentioned above, i.e. the NIS junction is disconnected from the thermal bath at $t = 0$ and remains adiabatically insulated while the protocol is being applied, being reconnected to the thermal bath at $t = \tau$ and undergoing equilibration through the exchange of heat with the bath afterwards. It is possible to lift such a restriction on the protocol by explicitly considering the interaction of the system with the thermal bath [50], but we will not consider such an extension here.

The knowledge of the work characteristic-function allows one to compute all work moments. If one performs the analytic continuation $v = i\beta$ in the above definition and uses the cyclic character of the trace operation, one obtains

$$\mathcal{G}_\beta(i\beta, \tau) = \mathcal{Z}_\tau(\beta)/\mathcal{Z}(\beta) = e^{-\beta\Delta F}, \quad (3)$$

where $\mathcal{Z}_\tau(\beta) = \text{Tr}(e^{-\beta\widehat{H}(\tau)})$ is the partition function for a junction in equilibrium, described by the Hamiltonian $\widehat{H}(\tau)$, ΔF being the difference in free-energies between the final equilibrium state (after thermalisation at a time $t > \tau$) and the initial equilibrium state at $t = 0$. Since in our case the Hamiltonian goes through a cycle, $\Delta F = 0$. Expressing $\mathcal{G}_\beta(i\beta, \tau)$ through its Fourier transform, we obtain

$$\langle e^{-\beta W} \rangle = \int_{-\infty}^{+\infty} dW e^{-\beta W} P_\beta(W, \tau) = e^{-\beta\Delta F} = 1, \quad (4)$$

which is a statement of the Jarzynski equality, which the work probability distribution satisfies by definition. Considering the relation between $\mathcal{G}_\beta(v, \tau)$ and $\mathcal{G}_\beta(v + i\beta, \tau)$, using again the cyclic character of the trace, it is possible to derive the identity $P_\beta(-W, -\tau) = e^{-\beta W} P_\beta(W, \tau)$ where by $P_\beta(W, -\tau)$ we denote the work probability distribution for the time-reversed protocol. This is a statement of the Crooks-Tasaki relation.

3. NIS junction model

The NIS junction is composed of a normal-metal film, a thin insulating layer and a superconducting film. We consider that such a junction can be described by an Hamiltonian consisting of three terms: a Fermi-gas describing the normal-metal film, a BCS superconductor describing the superconducting film and a tunnelling Hamiltonian coupling the two [79, 80, 81, 82, 83, 84, 85], which gives rise to induced superconductivity in the normal-metal film through the proximity effect (see [86] and references therein). We use the Hamiltonian

$$\widehat{H}(t) = \widehat{H}_n(t) + \widehat{H}_s + \widehat{X}. \quad (5)$$

The operator $\widehat{H}_n(t) = \sum_{\mathbf{k},\sigma} (\xi_{\mathbf{k}}^n - e\phi(t)) \widehat{c}_{n\mathbf{k}\sigma}^\dagger \widehat{c}_{n\mathbf{k}\sigma}$ is a free-fermion Hamiltonian describing the normal-metal, with $\phi(t)$ being the voltage difference across the junction, whose value changes in time. The operator

$$\widehat{H}_s = \sum_{\mathbf{k},\sigma} \xi_{\mathbf{k}}^s \widehat{c}_{s\mathbf{k}\sigma}^\dagger \widehat{c}_{s\mathbf{k}\sigma} + \sum_{\mathbf{k}} \left(\Delta_{\mathbf{k}} \widehat{c}_{s\mathbf{k}\uparrow}^\dagger \widehat{c}_{s-\mathbf{k}\downarrow}^\dagger + \overline{\Delta}_{\mathbf{k}} \widehat{c}_{s-\mathbf{k}\downarrow} \widehat{c}_{s\mathbf{k}\uparrow} \right), \quad (6)$$

is the Hamiltonian describing the BCS superconductor, where the pairing function $\Delta_{\mathbf{k}}$ is different from zero on a band of width $2\hbar\omega_D$ around the Fermi-level, where ω_D is the superconducting material's Debye frequency. Finally, the third term of Eq. (5) is given by

$$\widehat{X} = \sum_{\mathbf{k},\sigma} t^\sigma(\mathbf{k}) \left(\widehat{c}_{s\mathbf{k}\sigma}^\dagger \widehat{c}_{n\mathbf{k}\sigma} + \widehat{c}_{n-\mathbf{k}-\sigma}^\dagger \widehat{c}_{s-\mathbf{k}-\sigma} \right), \quad (7)$$

and represents the tunnelling process across the junction. We consider the overall tunnelling matrix element to be small with respect to the gap function $\Delta_{\mathbf{k}}$. Such matrix element is taken to be invariant under time-reversal, i.e. $\overline{t}^\sigma(\mathbf{k}) = t^{-\sigma}(-\mathbf{k})$ [83]. The fermions' momentum is directed along the plane of the films and we assume it to be conserved in a tunnelling process. We will measure the kinetic energy of electrons in the normal-metal film and in the superconducting film with respect to their common Fermi-energy μ . The super- or subscripts n and s refer to operators or quantities pertaining to the normal-metal or to the superconductor, respectively.

We consider the NIS to be initially ($t < 0$) in equilibrium with an heat bath at temperature T and that the voltage ϕ across the junction is zero. Such voltage is changed from 0 to a finite value V_{\max} and returned to zero within a finite time-interval starting at $t = 0$ and ending at $t = \tau$. During such an interval the junction is decoupled from the heat bath. For simplicity, we consider that the voltage protocol is symmetric with respect to the mid-point of the interval $t = \tau/2$. However, the calculation presented in Appendix A is valid for more general protocols. Also, $\phi(t)$ varies smoothly within a time-interval $\tau' \ll \tau$ and is returned to zero at the end of the protocol within the same time frame (see figure 1 (b)).

We will further place ourselves within the so-called adiabatic limit, i.e. we will consider that the switching-time $\tau' \gg \hbar/\Delta$, where Δ is the value of the gap-function within the superconductor. Thus, the voltage difference across the junction is held approximately constant within the microscopic time-scale of the superconductor. Therefore, the separation of time-scales in our problem is such that $\tau \gg \tau' \gg \hbar/\Delta$.

Using the Hamiltonian for an NIS junction and the voltage protocol specified, the work probability distribution can be obtained using Eq. (2), see next section. Note that although the Hamiltonian describing the system goes through a cycle and thus $P_\beta(-W, -\tau) = e^{-\beta W} P_\beta(W, \tau)$ holds, in the present case the time-reversed protocol cannot be identified with the original protocol since, due to the proximity effect, an induced superconducting order parameter will appear in the normal-metal. Initially, there will be no phase difference between the induced order parameter and the order parameter of the superconductor but, upon application of the voltage protocol, a phase

difference will appear and this will imply that the final equilibrium state of the junction will be distinct from the initial one.

4. Work dissipated due to the application of the voltage protocol to the junction

In this section, we establish that the dynamics of the NIS junction is equivalent to that of an assembly of quantum spins in a time-dependent magnetic field. This entails an equality between the work probability distributions of both systems, from which it follows that one can compute the average heat dissipated by the NIS junction, as well as its higher order fluctuations.

Starting from Eq. (5), the explicit dependence of the Hamiltonian on the time-dependent bias voltage can be eliminated by a gauge transformation at the expense of acquiring a time-dependent hopping term between the normal and superconducting films. Thus, the Hamiltonian becomes

$$\hat{H}_1(t) = \hat{H}_n(0) + \hat{H}_s + \widehat{\mathcal{X}}(t) \quad (8)$$

with

$$\widehat{\mathcal{X}}(t) = \sum_{\mathbf{k}, \sigma} t^\sigma(\mathbf{k}) \left(e^{ie\Phi(t)/\hbar} \hat{c}_{s\mathbf{k}\sigma}^\dagger \hat{c}_{n\mathbf{k}\sigma} + e^{-ie\Phi(t)/\hbar} \hat{c}_{n-\mathbf{k}-\sigma}^\dagger \hat{c}_{s-\mathbf{k}-\sigma} \right), \quad (9)$$

where $\Phi(t) = \int_0^t du \phi(u)$. Anticipating that, in general, $t^\sigma(\mathbf{k}) \ll \Delta$, one can, in the adiabatic limit discussed above, perform a generalised Schrieffer-Wolff transformation that takes into account the explicit time-dependence of $\widehat{\mathcal{X}}(t)$. After such a transformation, the metal and superconducting films effectively decouple and only the transformed metallic excitations are subjected to the time-dependent bias. The Hamiltonian, $\widehat{\mathcal{H}}_n(t)$, which describes the normal-metal film, explicitly reads

$$\begin{aligned} \widehat{\mathcal{H}}_n(t) = & \sum_{\mathbf{k}, \sigma} \tilde{\xi}_{\mathbf{k}\sigma}^n \hat{c}_{n\mathbf{k}\sigma}^\dagger \hat{c}_{n\mathbf{k}\sigma} \\ & + \sum_{\mathbf{k}} \left(\Delta_{\mathbf{k}}^n e^{-2ie\Phi(t)/\hbar} \hat{c}_{n\mathbf{k}\uparrow}^\dagger \hat{c}_{n-\mathbf{k}\downarrow}^\dagger + \overline{\Delta}_{\mathbf{k}}^n e^{2ie\Phi(t)/\hbar} \hat{c}_{n-\mathbf{k}\downarrow} \hat{c}_{n\mathbf{k}\uparrow} \right), \end{aligned} \quad (10)$$

where

$$\tilde{\xi}_{\mathbf{k}\sigma}^n = \xi_{\mathbf{k}}^n + \frac{|t^\sigma(\mathbf{k})|^2 (\xi_{\mathbf{k}}^n + \xi_{\mathbf{k}}^s)}{(\xi_{\mathbf{k}}^n)^2 - (E_{\mathbf{k}}^s)^2}, \quad (11)$$

$$\Delta_{\mathbf{k}}^n = - \frac{\Delta_{\mathbf{k}} (|t^\uparrow(\mathbf{k})|^2 + |t^\downarrow(\mathbf{k})|^2)}{2 [(\xi_{\mathbf{k}}^n)^2 - (E_{\mathbf{k}}^s)^2]}, \quad (12)$$

are, respectively, the renormalized kinetic energy of electrons and the induced superconducting parameter, in the normal metal, due to the proximity effect, $E_{\mathbf{k}}^s = \sqrt{(\xi_{\mathbf{k}}^s)^2 + |\Delta_{\mathbf{k}}|^2}$ being the energy of the superconducting excitations. This result is shown in detail in Appendix A. We thus conclude that for spin-independent tunnelling matrix elements, the Hamiltonian describing the normal-metal is equivalent, in each $((\mathbf{k}, \uparrow), (-\mathbf{k}, \downarrow))$ subspace, to the Hamiltonian of a two-level system under the action of a

circularly polarised field. Furthermore, \hat{H}_s does not contribute to the work-characteristic function, see Appendix A.

One can obtain the work characteristic function for a NIS junction simply by considering the product of individual characteristic-functions for two-level systems subjected to a circularly polarised field, see Appendix B. One obtains for the logarithm of the overall work characteristic-function, which is the generating function of the connected work-moments, the result

$$\mathcal{W}_\beta(v, \tau) \equiv \ln \mathcal{G}_\beta(v, \tau) = \sum_{\mathbf{k}} \ln \left\{ (1 - p_{\mathbf{k}}) + p_{\mathbf{k}} \cdot \frac{\cosh \left[(\beta + 2iv) \sqrt{|\Delta_{\mathbf{k}}^n|^2 + (\tilde{\xi}_{\mathbf{k}}^n)^2} \right]}{\cosh \left(\beta \sqrt{|\Delta_{\mathbf{k}}^n|^2 + (\tilde{\xi}_{\mathbf{k}}^n)^2} \right)} \right\}, \quad (13)$$

with $p_{\mathbf{k}} = \frac{|\Delta_{\mathbf{k}}^n|^2 (eV_{\max})^2}{(|\Delta_{\mathbf{k}}^n|^2 + (\tilde{\xi}_{\mathbf{k}}^n)^2)(|\Delta_{\mathbf{k}}^n|^2 + (\tilde{\xi}_{\mathbf{k}}^n - eV_{\max})^2)} \sin^2 \left(\frac{\tau}{\hbar} \sqrt{|\Delta_{\mathbf{k}}^n|^2 + (\tilde{\xi}_{\mathbf{k}}^n - eV_{\max})^2} \right)$. The detailed calculation can be found in Appendix B.

The average work dissipated per atom in the normal-metal film, \bar{w} , is obtained from the derivative of Eq. (13), evaluated at $v = 0$,

$$\bar{w} = \frac{2(eV_{\max})^2}{\mathcal{N}_{at}} \sum_{\mathbf{k}} \tanh \left(\beta \sqrt{|\Delta_{\mathbf{k}}^n|^2 + (\tilde{\xi}_{\mathbf{k}}^n)^2} \right) \times \frac{|\Delta_{\mathbf{k}}^n|^2 \sin^2 \left(\frac{\tau}{\hbar} \sqrt{|\Delta_{\mathbf{k}}^n|^2 + (\tilde{\xi}_{\mathbf{k}}^n - eV_{\max})^2} \right)}{\sqrt{|\Delta_{\mathbf{k}}^n|^2 + (\tilde{\xi}_{\mathbf{k}}^n)^2} (|\Delta_{\mathbf{k}}^n|^2 + (\tilde{\xi}_{\mathbf{k}}^n - eV_{\max})^2)}, \quad (14)$$

where \mathcal{N}_{at} is the total number of atoms in the normal-metal film. From Eq. (11), one has that for weak-coupling between the two films, $\tilde{\xi}_{\mathbf{k}}^n \approx \xi_{\mathbf{k}}^n$. Moreover, if, as in the system experimentally studied in [16], the dispersion relation is the same in both the normal-metal and the superconducting film (since the normal-metal film is composed of the same material as that of the superconducting one, but weakly doped with another metal), we have from Eq. (12) that $\Delta_{\mathbf{k}}^n = \frac{|t(\mathbf{k})|^2}{\Delta_{\mathbf{k}}^*}$. Assuming that both the order parameter $\Delta_{\mathbf{k}}$ and $|t(\mathbf{k})|^2$ only depend on \mathbf{k} through their dependence on $\xi_{\mathbf{k}}^n$ and are only non-zero in a vicinity of width $\hbar\omega_D$ around the Fermi level of the material composing the films, where ω_D is the Debye frequency of the said material, one can transform the above summation into an integral and write the above equation as

$$\bar{w} = 2(eV_{\max})^2 \int_{-\hbar\omega_D}^{\hbar\omega_D} d\xi \rho(\xi) \tanh \left(\beta \sqrt{|\Delta^n(\xi)|^2 + \xi^2} \right) \times \frac{|\Delta^n(\xi)|^2 \sin^2 \left(\frac{\tau}{\hbar} \sqrt{|\Delta^n(\xi)|^2 + (\xi - eV_{\max})^2} \right)}{\sqrt{|\Delta^n(\xi)|^2 + \xi^2} (|\Delta^n(\xi)|^2 + (\xi - eV_{\max})^2)}, \quad (15)$$

where we have introduced $\rho(\xi) = \frac{1}{\mathcal{N}_{at}} \sum_{\mathbf{k}} \delta(\xi - \xi_{\mathbf{k}}^n)$, the density of states per atom (and per spin-species) in the normal metal film.

Differentiating Eq. (13) twice with respect to v , at $v = 0$, one obtains the mean-square deviation and thus the relative deviation $\sqrt{\delta w^2}/\bar{w}$ of the work performed by the

NIS junction during the protocol. If one transforms such result into an integral using the same assumptions we have used to obtain Eq. (15), this integral can also be computed numerically. In a similar fashion, higher moments can be calculated.

The full characterisation of the work fluctuations requires the determination of the work probability distribution, through the computation of the inverse Fourier transform of $\mathcal{G}_\beta(v, \tau)$. Writing the probability distribution in terms of the logarithm of the characteristic function, as defined in Eq. (13), one has

$$P_\beta(W, \tau) = \int_{-\infty}^{+\infty} \frac{dv}{2\pi} e^{-iWv + \mathcal{W}_\beta(v, \tau)}. \quad (16)$$

Since W is an extensive quantity, *i.e.* it is proportional to \mathcal{N}_{at} , the integral can be computed using a saddle-point approximation. Minimising the argument of the exponential in Eq. (16), one obtains the following saddle-point equation

$$w = \int_{-\hbar\omega_D}^{\hbar\omega_D} d\xi \rho(\xi) \partial_x S(\xi, x), \quad (17)$$

where $w = W/\mathcal{N}_{at}$ is the work per atom of the sample and $S(\xi, x)$ is given by

$$S(\xi, x) = \ln \left\{ (1 - p(\xi)) + p(\xi) \cdot \frac{\cosh \left[(\beta + 2x) \sqrt{|\Delta^n(\xi)|^2 + \xi^2} \right]}{\cosh \left[\beta \sqrt{|\Delta^n(\xi)|^2 + \xi^2} \right]} \right\}, \quad (18)$$

where $p(\xi) = \frac{|\Delta^n(\xi)|^2 (eV_{\max})^2}{(|\Delta^n(\xi)|^2 + \xi^2)(|\Delta^n(\xi)|^2 + (\xi - eV_{\max})^2)} \sin^2 \left(\frac{\tau}{\hbar} \sqrt{|\Delta^n(\xi)|^2 + (\xi - eV_{\max})^2} \right)$. Note that the solutions, $x(w)$, of the saddle-point equation (17) correspond to the analytic continuation of the exponent of the integral given in Eq. (16) to imaginary values of $v = -ix(w)$. Numerically solving Eq. (17) for a set of values of w (with the same assumptions as in the two calculations previously performed) and substituting its solution in Eq. (16), one obtains the result plotted in figure 2 for the logarithm of $P_\beta(W, \tau)$, divided by \mathcal{N}_{at} . This calculation indicates that the work probability distribution is a stretched exponential for both positive and negative W . In the inset, the exponent associated with such a stretched exponential is determined for positive W (blue triangles).

We have also subtracted the factor βw from the logarithm of the probability distribution at negative W (green squares in the inset) and the function so obtained coincides with the logarithm of the probability distribution for positive W , thus showing that the distribution obeys the Crooks-Tasaki relation at the level of the numerics.

The above figure is a clear indication of the non-Gaussian nature of the work distribution, which may allow the reconstruction of the distribution above from experimental data, using the method that we propose in the next section.

5. Isothermal measurement of the dissipated heat

Our primary motivation for the present study was the analysis of the validity of the Crooks-Tasaki relation in an experimentally relevant setting. As shown above, this can be accomplished by measuring the heat dissipated in the NIS junction during a full




Figure 2. Logarithm of the work per atom distribution for a AlMn/AlO₂/Al junction at $T = 1$ K.

cycle of the protocol. Since such measurements require a high degree of control of the energy flows occurring in an experimental set-up, it is advantageous that the released heat scales with the contact area between the superconducting and normal metal films.

The measurement of the released heat is carried out at constant temperature by coupling the system to an external probe that absorbs a quantity of heat $\Delta Q = W$ isothermally. If the absorption occurs reversibly, it corresponds to an increase of entropy of the probe equal to $\Delta S = \Delta Q/T$. The probe will increase in volume as result of the heat absorbed, the variation of volume being given by

$$\Delta \mathcal{V} = \frac{\kappa_T}{\alpha} \Delta S, \quad (19)$$

where κ_T is the isothermal compressibility of the probe and α its thermal expansion coefficient, as follows from equilibrium thermodynamics. Therefore, the relative variation of the volume of the probe is given by

$$\frac{\Delta \mathcal{V}}{\mathcal{V}} = \frac{W}{\gamma n c_v T}, \quad (20)$$

where $\gamma = \frac{\alpha v_m}{c_v \kappa_T}$ is the Grüneisen parameter of the material that constitutes the probe, v_m its molar volume, n is the number of moles of material contained in the probe, and c_v the material's molar heat capacity. The measurement of the relative variation of the probe's volume thus gives direct experimental access to the work produced during the prescribed protocol. The film itself may act as the probe. A histogram of the distribution of data obtained for multiple realizations of the protocol will thus allow the reconstruction of the full work distribution.

We now turn to a discussion of the experimental verification. As a concrete example, we will consider the junction studied in reference [16], where manganese

doped aluminium (the Mn concentration being 0.4% in mass) acts as the normal-metal electrode in the junction and pure aluminium acts as the superconducting electrode (with aluminium-oxide being the insulating material).

In aluminium, the density of states is approximately constant in the vicinity of the Fermi surface yielding $\rho \approx 0.0117\omega_D^{-1}$ per atom (and per spin) [87, 88], in units of the Debye frequency, which is approximately $\omega_D = 10^{14}$ Hz in aluminium. This metal is a BCS superconductor with a transition temperature, $T_c = 1.20$ K [89], and the gap-function given by $\Delta = 1.764 k_B T_c$, where one uses the universal (BCS) relation [90], valid in the weak-coupling limit. For an intermediate coupling superconductor the gap is larger than predicted by the BCS relation. This is of no concern here, since we assumed $\tau \gg \tau' \gg \hbar/\Delta \simeq 3.6 \times 10^{-12}$ s.

For concreteness, we consider the example of a protocol of duration $\tau = 10^{-6}$ s with an applied voltage $V_{\max} = 8 \times 10^{-5}$ V at a temperature of $T = 1$ K. Performing the integral in (15), assuming a typical tunnelling element $t(\mathbf{k}) \simeq 0.5\Delta$, we obtain, using see Eq. (14), an average work $\bar{w} \approx 2 \times 10^{-8} \hbar\omega_D$ per atom which amounts to approximately $5 \mu\text{J}$ of dissipated heat per gram of aluminium. For these parameters, we also obtain a standard deviation of $\overline{\delta w^2} \approx 6.9 \times 10^{-11} (\hbar\omega_D)^2$ corresponding to a relative deviation, $\sqrt{\overline{\delta w^2}}/\bar{w} \approx 420$, which is a large value for such a quantity.

For aluminium, $\gamma \approx 2.1$ [91] and $c_v \approx 1.5 \text{ mJ mol}^{-1} \text{ K}^{-1}$ [18]. Assuming that the film itself will act as the probe, if we use $\langle W \rangle_\beta \approx 5 \mu\text{J/g}$, we obtain a relative deviation $\frac{\Delta\gamma}{\gamma} \approx 0.01$, on average, which should be measurable using, e.g. capacitive methods [17, 18], even for such a small heat release, provided that other sources of heat dissipation can be minimized.

6. Conclusions

Fluctuation-dissipation relations play a central role in thermodynamics and the approach to equilibrium, but the direct experimental verification of the various relations can often be difficult. In this work, we discussed a simple but realistic experimental set-up that will allow to test the Crooks-Tasaki fluctuation relation. Our proposal consists of an NIS junction of a superconductor and a normal metal where the bias voltage is altered according to a specified protocol, see Fig. 1-(b).

As shown, the released heat scales with the contact area between the superconducting and normal metal films. Our proposal is thus amenable to experimental verification, by coupling the junction to a probe that isothermally absorbs the dissipated heat and changes its volume as a result. Order of magnitude estimates of this effect for the aluminium-based junctions of Ref. [16] were presented in Sect. 5.

The calculation of the full work distribution function and its first moments clearly establishes an exponential behavior of such quantity that displays a non-Gaussian character with a non-standard decay exponent.

Our results are based on an equivalence between the dynamics of a NIS junction and the dynamics of an assembly of two-level systems subjected to a circularly polarized

field. This equivalence holds in an adiabatic limit in which the switching-on/off of the potential difference across the junction takes place in time scales much larger than the microscopic time scale of the system, determined by the value of the superconducting gap.

Acknowledgements: We acknowledge helpful discussions with O. Stockert, J. Pekola, E. Lutz, M. Kiselev, A. Muramatsu, K. Kroy, E. Lage, J. L. Santos, M. I. Vasilevskiy, N. M. R. Peres and A.-P. Jauho. J.E.S. work contract at the University of Minho, where most of the present research was conducted, was financed in the framework of the Program of Recruitment of Post Doctoral Researchers for the Portuguese Scientific and Technological System, within the Operational Program Human Potential (POPH) of the QREN, participated by the European Social Fund (ESF) and national funds of the Portuguese Ministry of Education and Science (MEC). He also acknowledges support provided to the current research project by FEDER through the COMPETE Program and by FCT in the framework of the Strategic Project PEST-C/FIS/UI607/2011. He would also like to thank the support of the MPIPkS in Dresden within the framework of their Visitors' Program at several stages of this work, as well as the support of the Center for Nanostructured Graphene (CNG) at the Technical University of Denmark at the final stage of the work. The CNG is sponsored by the Danish National Research Foundation, Project No. DNRF58. S. Kirchner acknowledges partial support by the National Natural Science Foundation of China, grant No.11474250.

Appendix A. Mapping of the work characteristic-function to that of an assembly of two-level systems

Following [92] and starting from the Hamiltonian (5), we change representation, so as to write the original time-evolution operator $\hat{U}(t, 0)$, determined by such Hamiltonian, in terms of the time-evolution operator in a new representation, $\hat{V}(t, 0)$, with $0 \leq t \leq \tau$,

$$\hat{U}(t, 0) = e^{ie\Phi(t)\hat{N}_n/\hbar} \hat{V}(t, 0), \quad (\text{A.1})$$

where $\Phi(t) = \int_0^t du \phi(u)$ and $\hat{N}_n = \sum_{\mathbf{k}, \sigma} \hat{c}_{n\mathbf{k}\sigma}^\dagger \hat{c}_{n\mathbf{k}\sigma}$ is the number of particles on the normal side of the junction. This change of representation corresponds to a gauge transformation that eliminates the dependence on $\phi(t)$ of $\hat{H}_n(t)$ and transfers such dependence to the tunnelling part of the Hamiltonian. The transformed operator, $\hat{H}_1(t) = e^{-ie\Phi(t)\hat{N}_n/\hbar} \hat{H}(t) e^{ie\Phi(t)\hat{N}_n/\hbar}$, is given by $\hat{H}_1(t) = \hat{H}_0 + \lambda \hat{\mathcal{X}}(t)$, where λ is a small dimensionless parameter that sets the scale of the tunnelling matrix element and which is explicitly written here for convenience of calculation. The operator $\hat{H}_0 = \hat{H}_n(0) + \hat{H}_s$ describes the normal-metal film and the superconducting film at $t = 0$, in the absence of the tunnelling operator (7). On the other hand, the operator $\hat{\mathcal{X}}(t)$ is given by equation (7).

The operator $\hat{V}(t, 0)$ can be expressed in terms of a time-ordered exponential involving the transformed Hamiltonian $\hat{H}_1(t)$. This exponential can be written in terms

of an ordered product of exponentials, evaluated at increasing times, as

$$\widehat{V}(t, 0) = \prod_{l=1}^N \exp \left[-\frac{it}{\hbar N} \widehat{H}_1(u_l) \right] \quad (\text{A.2})$$

where $N \rightarrow \infty$ and $u_l = l \cdot t/N$. We now introduce a second-unitary transformation, dependent on both λ and on t , by defining a transformed Hamiltonian operator as

$$\widehat{H}_2(t) = e^{i\lambda\widehat{S}(t)} \widehat{H}_1(t) e^{-i\lambda\widehat{S}(t)}, \quad (\text{A.3})$$

where the Hermitian operator $\widehat{S}(t)$ will be adequately chosen below. Substituting this result in (A.2), we obtain

$$\widehat{V}(t, 0) = e^{-i\lambda\widehat{S}(t)} \prod_{l=1}^N \left[\exp \left[-\frac{it}{\hbar N} \widehat{H}_2(u_l) \right] e^{i\lambda\widehat{S}(u_l)} e^{-i\lambda\widehat{S}(u_{l-1})} \right] e^{i\lambda\widehat{S}(0)}. \quad (\text{A.4})$$

Using the Baker-Campbell-Hausdorff formula, one can write, to order λ^2 , and to order $1/N$

$$\begin{aligned} e^{i\lambda\widehat{S}(u_l)} e^{-i\lambda\widehat{S}(u_{l-1})} &\approx e^{i\lambda(\widehat{S}(u_l) - \widehat{S}(u_{l-1})) + \frac{\lambda^2}{2} [\widehat{S}(u_l), \widehat{S}(u_{l-1})]_-} \\ &\approx e^{\frac{i\lambda}{N} \left(\frac{d\widehat{S}}{du} + \frac{i\lambda}{2} \left[\widehat{S}(u_l), \frac{d\widehat{S}}{du} \right]_- \right)}, \end{aligned} \quad (\text{A.5})$$

where $[,]_-$ is the commutator of two operators and where the derivative of $\widehat{S}(u)$ is computed at u_l . Substituting (A.5) in (A.4), we obtain to order λ^2 and order $1/N$, the result

$$\widehat{V}(t, 0) = e^{-i\lambda\widehat{S}(t)} \prod_{l=1}^N \exp \left[-\frac{it}{\hbar N} \widehat{\mathcal{H}}(u_l) \right] e^{i\lambda\widehat{S}(0)}, \quad (\text{A.6})$$

where, to order λ^2 , the operator $\widehat{\mathcal{H}}(t)$ is given by

$$\begin{aligned} \widehat{\mathcal{H}}(t) &\approx e^{i\lambda\widehat{S}(t)} \left(\widehat{H}_0 + \lambda \widehat{\mathcal{X}}(t) \right) e^{-i\lambda\widehat{S}(t)} - \lambda\hbar \frac{d\widehat{S}}{dt} - \frac{i\hbar\lambda^2}{2} \left[\widehat{S}(t), \frac{d\widehat{S}}{dt} \right]_- \\ &\approx \widehat{H}_0 + \lambda \left(-\hbar \frac{d\widehat{S}}{dt} + \widehat{\mathcal{X}}(t) + i \left[\widehat{S}(t), \widehat{H}_0 \right]_- \right) \\ &\quad + \frac{i\lambda^2}{2} \left[\widehat{S}(t), -\hbar \frac{d\widehat{S}}{dt} + \widehat{\mathcal{X}}(t) + i \left[\widehat{S}(t), \widehat{H}_0 \right]_- \right]_- \\ &\quad + \frac{i\lambda^2}{2} \left[\widehat{S}(t), \widehat{\mathcal{X}}(t) \right]_- . \end{aligned} \quad (\text{A.7})$$

We choose the hermitian operator $\widehat{S}(t)$ such that the linear term in λ vanishes in (A.7), i.e. such operator obeys the first order ordinary differential equation

$$\hbar \frac{d\widehat{S}}{dt} = \widehat{\mathcal{X}}(t) + i \left[\widehat{S}(t), \widehat{H}_0 \right]_- , \quad (\text{A.8})$$

whereas the transformed Hamiltonian reduces to

$$\widehat{\mathcal{H}}(t) = \widehat{H}_0 + \frac{i\lambda^2}{2} \left[\widehat{S}(t), \widehat{\mathcal{X}}(t) \right]_- . \quad (\text{A.9})$$

One still needs to specify an initial condition for (A.8) to be properly defined, which we will do below. The above choice of $\widehat{S}(t)$ corresponds to a time-dependent Schrieffer-Wolff transformation [93, 94], which is reminiscent of the work of [95, 96]. One can thus write the operator $\widehat{U}(t, 0)$ as $\widehat{U}(t, 0) = e^{ie\Phi(t)\widehat{N}_n/\hbar} e^{-i\lambda\widehat{S}(t)} \widehat{\mathcal{V}}(t, 0) e^{i\lambda\widehat{S}(0)}$, where

$$\widehat{\mathcal{V}}(t, 0) = T \exp \left[-\frac{i}{\hbar} \int_0^t du \widehat{\mathcal{H}}(u) \right], \quad (\text{A.10})$$

which is valid to order λ^2 and where $\widehat{\mathcal{H}}(u)$ is given by (A.9). Substituting this expression for $\widehat{U}(t, 0)$ in (2) and using the cyclic character of the trace operation, one obtains

$$\mathcal{G}_\beta(v, \tau) = Tr \left[\widehat{\mathcal{V}}^\dagger(\tau, 0) e^{i\lambda\widehat{S}(\tau)} e^{i\widehat{H}_1(\tau)v} e^{-i\lambda\widehat{S}(\tau)} \cdot \widehat{\mathcal{V}}(\tau, 0) e^{i\lambda\widehat{S}(0)} e^{-i\widehat{H}(0)v} \widehat{\rho}(\beta) e^{-i\lambda\widehat{S}(0)} \right]. \quad (\text{A.11})$$

Under the action of, respectively, $\widehat{S}(0)$ and $\widehat{S}(\tau)$, $\widehat{H}(0)$ and $\widehat{H}_1(\tau)$ do not transform into $\widehat{\mathcal{H}}(0)$ and $\widehat{\mathcal{H}}(\tau)$, as the factors that involve the derivative of $\widehat{S}(t)$ at these points, and which appear in (A.7), are not present in the transformation. However, we may still fix the derivative of $\widehat{S}(t)$ at one of these points, by the choice of an appropriate initial condition. We thus choose $\left. \frac{d\widehat{S}}{dt} \right|_{t=0} = 0$. The explicit computation of $\mathcal{G}_\beta(v, \tau)$ now requires the solution of (A.8), subjected to the condition that we have imposed at $t = 0$. In order to solve this equation, we write $\widehat{S}(t)$ in the form

$$\begin{aligned} \widehat{S}(t) = & \sum_{k\sigma} [r_{\mathbf{k}}^\sigma(t) \widehat{c}_{s\mathbf{k}\sigma}^\dagger \widehat{c}_{n\mathbf{k}\sigma} + \bar{r}_{\mathbf{k}}^\sigma(t) \widehat{c}_{n\mathbf{k}\sigma}^\dagger \widehat{c}_{s\mathbf{k}\sigma}] \\ & + \sum_{k\sigma} [\bar{s}_{\mathbf{k}}^\sigma(t) \widehat{c}_{s-\mathbf{k}-\sigma}^\dagger \widehat{c}_{n-\mathbf{k}-\sigma}^\dagger + s_{\mathbf{k}}^\sigma(t) \widehat{c}_{n-\mathbf{k}-\sigma} \widehat{c}_{s-\mathbf{k}-\sigma}], \end{aligned} \quad (\text{A.12})$$

where $r_{\mathbf{k}}^\sigma(t)$ and $s_{\mathbf{k}}^\sigma(t)$ are complex-valued functions, to be determined. Substituting (A.12) in (A.8) and equating the terms pertaining to the same pairs of operators, we obtain the coupled system of first-order ordinary differential equations

$$\begin{cases} \frac{dr_{\mathbf{k}}^\sigma}{dt} = \frac{t^\sigma(\mathbf{k})}{\hbar} e^{ie\Phi(t)/\hbar} + \frac{i}{\hbar} (\xi_{\mathbf{k}}^n - \xi_{\mathbf{k}}^s) r_{\mathbf{k}}^\sigma + \frac{i}{\hbar} \sigma \Delta_{\mathbf{k}} s_{\mathbf{k}}^\sigma \\ \frac{ds_{\mathbf{k}}^\sigma}{dt} = \frac{i}{\hbar} (\xi_{\mathbf{k}}^n + \xi_{\mathbf{k}}^s) s_{\mathbf{k}}^\sigma + \frac{i}{\hbar} \sigma \bar{\Delta}_{\mathbf{k}} r_{\mathbf{k}}^\sigma \end{cases}, \quad (\text{A.13})$$

with the initial conditions $\left. \frac{dr_{\mathbf{k}}^\sigma}{dt} \right|_{t=0} = \left. \frac{ds_{\mathbf{k}}^\sigma}{dt} \right|_{t=0} = 0$. These conditions, once substituted in (A.13) at $t = 0$, yield for $r_{\mathbf{k}}^\sigma(0)$ and $s_{\mathbf{k}}^\sigma(0)$, the result

$$\begin{cases} r_{\mathbf{k}}^\sigma(0) = \frac{it^\sigma(\mathbf{k})(\xi_{\mathbf{k}}^n + \xi_{\mathbf{k}}^s)}{(\xi_{\mathbf{k}}^n)^2 - (E_{\mathbf{k}}^s)^2} \\ s_{\mathbf{k}}^\sigma(0) = -\frac{it^\sigma(\mathbf{k})\sigma\bar{\Delta}_{\mathbf{k}}}{(\xi_{\mathbf{k}}^n)^2 - (E_{\mathbf{k}}^s)^2} \end{cases}. \quad (\text{A.14})$$

This choice of parameters corresponds to the time-independent Schrieffer-Wolff transformation introduced in [97] and such transformation can be applied to the Hamiltonian of the system at $t \leq 0$, since $\phi(t) = 0$ in this interval.

Writing the pairing function in terms of an amplitude and a phase, i.e. $\Delta_{\mathbf{k}} = |\Delta_{\mathbf{k}}| e^{i\varphi}$, where φ is independent of \mathbf{k} , we define the new variables $x_{\mathbf{k}}^\sigma(t) = u_{\mathbf{k}}^\sigma(t) e^{-\frac{i}{\hbar}\xi_{\mathbf{k}}^n t - \frac{i}{2}\varphi}$, $y_{\mathbf{k}}^\sigma(t) = v_{\mathbf{k}}^\sigma(t) e^{-\frac{i}{\hbar}\xi_{\mathbf{k}}^n t + \frac{i}{2}\varphi}$. Substituting these expressions in equation

(A.13) and differentiating the result, we obtain two decoupled second-order equations for $x_{\mathbf{k}}^\sigma(t)$ and $y_{\mathbf{k}}^\sigma(t)$, with appropriate boundary conditions. These equations can be easily solved.

The solutions of such equations, expressed in terms of the original functions $r_{\mathbf{k}}^\sigma(t)$, $s_{\mathbf{k}}^\sigma(t)$, are given by

$$\begin{aligned} r_{\mathbf{k}}^\sigma(t) = & it^\sigma(\mathbf{k}) e^{\frac{i}{\hbar}\xi_{\mathbf{k}}^n t} \left[\frac{\xi_{\mathbf{k}}^n + \xi_{\mathbf{k}}^s}{(\xi_{\mathbf{k}}^n)^2 - (E_{\mathbf{k}}^s)^2} \left(\cos(\omega_{\mathbf{k}}^s t) - \frac{i\xi_{\mathbf{k}}^n}{E_{\mathbf{k}}^s} \cdot \sin(\omega_{\mathbf{k}}^s t) \right) \right. \\ & + \frac{1}{\hbar E_{\mathbf{k}}^s} \int_0^t du (e\phi(u) - \xi_{\mathbf{k}}^n - \xi_{\mathbf{k}}^s) \\ & \left. \cdot e^{\frac{i}{\hbar}(e\Phi(u) - \xi_{\mathbf{k}}^n u)} \sin(\omega_{\mathbf{k}}^s(t-u)) \right], \end{aligned} \quad (\text{A.15})$$

and

$$\begin{aligned} s_{\mathbf{k}}^\sigma(t) = & -i\sigma \bar{\Delta}_{\mathbf{k}} t^\sigma(\mathbf{k}) e^{\frac{i}{\hbar}\xi_{\mathbf{k}}^n t} \left[\frac{1}{(\xi_{\mathbf{k}}^n)^2 - (E_{\mathbf{k}}^s)^2} \left(\cos(\omega_{\mathbf{k}}^s t) - \frac{i\xi_{\mathbf{k}}^n}{E_{\mathbf{k}}^s} \cdot \sin(\omega_{\mathbf{k}}^s t) \right) \right. \\ & \left. - \frac{1}{\hbar E_{\mathbf{k}}^s} \int_0^t du e^{\frac{i}{\hbar}(e\Phi(u) - \xi_{\mathbf{k}}^n u)} \sin(\omega_{\mathbf{k}}^s(t-u)) \right]. \end{aligned} \quad (\text{A.16})$$

Now, one can show, using the know properties of the exponential function and integration by parts, the following identity

$$\begin{aligned} \frac{1}{\hbar E_{\mathbf{k}}^s} \int_0^t du e^{\frac{i}{\hbar}(e\Phi(u) - \xi_{\mathbf{k}}^n u)} \sin(\omega_{\mathbf{k}}^s(t-u)) = & -\frac{1}{(\xi_{\mathbf{k}}^n)^2 - (E_{\mathbf{k}}^s)^2} \cdot \left[e^{\frac{i}{\hbar}(e\Phi(t) - \xi_{\mathbf{k}}^n t)} \right. \\ & - \left(\cos(\omega_{\mathbf{k}}^s t) - \frac{i\xi_{\mathbf{k}}^n}{E_{\mathbf{k}}^s} \cdot \sin(\omega_{\mathbf{k}}^s t) \right) \\ & - \frac{i}{\hbar} \int_0^t du e\phi(u) e^{\frac{i}{\hbar}(e\Phi(u) - \xi_{\mathbf{k}}^n u)} \\ & \cdot \left(\cos(\omega_{\mathbf{k}}^s(t-u)) \right. \\ & \left. \left. - \frac{i\xi_{\mathbf{k}}^n}{E_{\mathbf{k}}^s} \cdot \sin(\omega_{\mathbf{k}}^s(t-u)) \right) \right]. \end{aligned} \quad (\text{A.17})$$

Using (A.17) in (A.15) and (A.16), one can write the solution of the system given in (A.13) as

$$\begin{aligned} r_{\mathbf{k}}^\sigma(t) = & \frac{it^\sigma(\mathbf{k}) (\xi_{\mathbf{k}}^n + \xi_{\mathbf{k}}^s)}{(\xi_{\mathbf{k}}^n)^2 - (E_{\mathbf{k}}^s)^2} \cdot e^{\frac{ie}{\hbar}\Phi(t)} + \frac{t^\sigma(\mathbf{k})}{\hbar[(\xi_{\mathbf{k}}^n)^2 - (E_{\mathbf{k}}^s)^2]} \int_0^t du e\phi(u) \\ & \times e^{\frac{i}{\hbar}(e\Phi(u) + \xi_{\mathbf{k}}^n(t-u))} \cdot \left((\xi_{\mathbf{k}}^n + \xi_{\mathbf{k}}^s) \cdot \cos(\omega_{\mathbf{k}}^s(t-u)) \right. \\ & \left. - i \frac{\xi_{\mathbf{k}}^s(\xi_{\mathbf{k}}^n + \xi_{\mathbf{k}}^s) + |\Delta_{\mathbf{k}}|^2}{E_{\mathbf{k}}^s} \cdot \sin(\omega_{\mathbf{k}}^s(t-u)) \right), \end{aligned} \quad (\text{A.18})$$

and

$$\begin{aligned} s_{\mathbf{k}}^\sigma(t) = & -\frac{it^\sigma(\mathbf{k}) \sigma \bar{\Delta}_{\mathbf{k}}}{(\xi_{\mathbf{k}}^n)^2 - (E_{\mathbf{k}}^s)^2} \cdot e^{\frac{ie}{\hbar}\Phi(t)} - \frac{t^\sigma(\mathbf{k}) \sigma \bar{\Delta}_{\mathbf{k}}}{\hbar[(\xi_{\mathbf{k}}^n)^2 - (E_{\mathbf{k}}^s)^2]} \int_0^t du e\phi(u) \\ & \times e^{\frac{i}{\hbar}(e\Phi(u) + \xi_{\mathbf{k}}^n(t-u))} \cdot \left(\cos(\omega_{\mathbf{k}}^s(t-u)) - \frac{i\xi_{\mathbf{k}}^n}{E_{\mathbf{k}}^s} \cdot \sin(\omega_{\mathbf{k}}^s(t-u)) \right). \end{aligned} \quad (\text{A.19})$$

We stated above that we are considering protocols in which the applied voltage difference $\phi(t)$ changes from 0 to V_{\max} in a time interval τ' i.e. $\phi(t) = V_{\max}\psi(t/\tau')$, where $\psi(x)$

changes from 0 to 1 in an interval of length 1. The integrals which enter the second term of both (A.18) and (A.19) are all, after the change of variables $u \rightarrow u/\tau'$, of the form

$$\frac{eV_{\max}\tau'}{\hbar} \int_0^{t/\tau'} du \psi(u) e^{\frac{ieV_{\max}\tau'}{\hbar}\Psi(u)} e^{-\frac{i\tau'}{\hbar}(\xi_{\mathbf{k}}^n \pm E_{\mathbf{k}}^s)u}, \quad (\text{A.20})$$

where $\Psi(u) = \int_0^u dv \psi(v)$. The term $\psi(u) e^{\frac{ieV_{\max}\tau'}{\hbar}\Psi(u)}$ is a bounded function, whereas the exponential $e^{-\frac{i\tau'}{\hbar}(\xi_{\mathbf{k}}^n \pm E_{\mathbf{k}}^s)u}$ is at least equal to $e^{-\frac{i\tau'}{\hbar}|\Delta_{\mathbf{k}}|u}$, which in the adiabatic regime $\tau' \gg \hbar/\Delta$ oscillates rapidly. Thus, provided that $t \gg \tau'$ and $eV_{\max}\tau'/\hbar \leq 1$, then (A.20) is zero by the Riemann-Lebesgue lemma. From these conditions, we also obtain $eV_{\max} \ll \Delta$, i.e. the tunnelling voltage applied should be well within the gap. Therefore, we conclude that in the adiabatic limit and for $t \gg \tau'$, (A.18) and (A.19) reduce to

$$\begin{cases} r_{\mathbf{k}}^{\sigma}(t) \approx \frac{it^{\sigma}(\mathbf{k})(\xi_{\mathbf{k}}^n + \xi_{\mathbf{k}}^s)}{(\xi_{\mathbf{k}}^n)^2 - (E_{\mathbf{k}}^s)^2} \cdot e^{\frac{ie}{\hbar}\Phi(t)} \\ s_{\mathbf{k}}^{\sigma}(t) \approx -\frac{it^{\sigma}(\mathbf{k})\sigma\bar{\Delta}_{\mathbf{k}}}{(\xi_{\mathbf{k}}^n)^2 - (E_{\mathbf{k}}^s)^2} \cdot e^{\frac{ie}{\hbar}\Phi(t)} \end{cases}. \quad (\text{A.21})$$

From such a discussion and since $\phi(\tau) = 0$, it also follows that $\left. \frac{dr_{\mathbf{k}}^{\sigma}}{dt} \right|_{t=\tau} \approx \left. \frac{ds_{\mathbf{k}}^{\sigma}}{dt} \right|_{t=\tau} \approx 0$ and hence we obtain $\left. \frac{d\hat{S}}{dt} \right|_{t=\tau} = 0$. In such a case, we can write equation (A.11) as

$$\mathcal{G}_{\beta}(v, \tau) = Tr \left[\widehat{\mathcal{V}}^{\dagger}(\tau, 0) e^{i\widehat{\mathcal{H}}(\tau)v} \widehat{\mathcal{V}}(\tau, 0) e^{-i\widehat{\mathcal{H}}(0)v} \widehat{\rho}(\beta) \right], \quad (\text{A.22})$$

where $\widehat{\mathcal{H}}(\tau)$ is given by (A.9) (with t substituted by τ in that equation). Substituting the operator $\widehat{S}(t)$ by its expression as given in (A.12), with $r_{\mathbf{k}}^{\sigma}(t)$ and $s_{\mathbf{k}}^{\sigma}(t)$ given by (A.21), in (A.9), we obtain that $\widehat{\mathcal{H}}(t) = \widehat{\mathcal{H}}_s + \widehat{\mathcal{H}}_n(t)$, where the operators $\widehat{\mathcal{H}}_s$ and $\widehat{\mathcal{H}}_n(t)$, pertaining, respectively, to the superconductor and the normal-metal, are given by

$$\widehat{\mathcal{H}}_s = \sum_{\mathbf{k}, \sigma} \tilde{\xi}_{\mathbf{k}\sigma}^s \widehat{c}_{s\mathbf{k}\sigma}^{\dagger} \widehat{c}_{s\mathbf{k}\sigma} + \sum_{\mathbf{k}} \left(\Delta_{\mathbf{k}}^s \widehat{c}_{s\mathbf{k}\uparrow}^{\dagger} \widehat{c}_{s-\mathbf{k}\downarrow}^{\dagger} + \bar{\Delta}_{\mathbf{k}}^s \widehat{c}_{s-\mathbf{k}\downarrow} \widehat{c}_{s\mathbf{k}\uparrow} \right), \quad (\text{A.23})$$

with

$$\tilde{\xi}_{\mathbf{k}\sigma}^s = \xi_{\mathbf{k}}^s - \frac{\lambda^2 |t^{\sigma}(\mathbf{k})|^2 (\xi_{\mathbf{k}}^n + \xi_{\mathbf{k}}^s)}{(\xi_{\mathbf{k}}^n)^2 - (E_{\mathbf{k}}^s)^2}, \quad (\text{A.24})$$

$$\Delta_{\mathbf{k}}^s = \Delta_{\mathbf{k}} \cdot \left(1 - \frac{\lambda^2}{2} \cdot \frac{|t^{\uparrow}(\mathbf{k})|^2 + |t^{\downarrow}(\mathbf{k})|^2}{(\xi_{\mathbf{k}}^n)^2 - (E_{\mathbf{k}}^s)^2} \right), \quad (\text{A.25})$$

being the renormalised kinetic energy and pairing function in the superconductor, and

$$\begin{aligned} \widehat{\mathcal{H}}_n(t) &= \sum_{\mathbf{k}, \sigma} \tilde{\xi}_{\mathbf{k}\sigma}^n \widehat{c}_{n\mathbf{k}\sigma}^{\dagger} \widehat{c}_{n\mathbf{k}\sigma} \\ &+ \sum_{\mathbf{k}} \left(\Delta_{\mathbf{k}}^n e^{-2ie\Phi(t)/\hbar} \widehat{c}_{n\mathbf{k}\uparrow}^{\dagger} \widehat{c}_{n-\mathbf{k}\downarrow}^{\dagger} + \bar{\Delta}_{\mathbf{k}}^n e^{2ie\Phi(t)/\hbar} \widehat{c}_{n-\mathbf{k}\downarrow} \widehat{c}_{n\mathbf{k}\uparrow} \right), \end{aligned} \quad (\text{A.26})$$

with

$$\tilde{\xi}_{\mathbf{k}\sigma}^n = \xi_{\mathbf{k}}^n + \frac{\lambda^2 |t^{\sigma}(\mathbf{k})|^2 (\xi_{\mathbf{k}}^n + \xi_{\mathbf{k}}^s)}{(\xi_{\mathbf{k}}^n)^2 - (E_{\mathbf{k}}^s)^2}, \quad (\text{A.27})$$

$$\Delta_{\mathbf{k}}^n = -\frac{\lambda^2}{2} \cdot \frac{\Delta_{\mathbf{k}} (|t^{\uparrow}(\mathbf{k})|^2 + |t^{\downarrow}(\mathbf{k})|^2)}{(\xi_{\mathbf{k}}^n)^2 - (E_{\mathbf{k}}^s)^2}, \quad (\text{A.28})$$

being the renormalised kinetic energy and induced pairing function in the normal-metal, due to the proximity effect. However, such an induced pairing function is multiplied by a time-dependent phase, due to the applied voltage across the junction. If one assumes that the tunnelling square amplitudes are independent of spin, i.e. $|t^\uparrow(\mathbf{k})|^2 = |t^\downarrow(\mathbf{k})|^2$, the modified kinetic energies are also independent of the spin one can, introducing in each $((\mathbf{k}, \uparrow), (-\mathbf{k}, \downarrow))$ subspace the operators,

$$\hat{\sigma}_{\mathbf{k}}^z = \hat{c}_{n\mathbf{k}\uparrow}^\dagger \hat{c}_{n\mathbf{k}\uparrow} + \hat{c}_{n-\mathbf{k}\downarrow}^\dagger \hat{c}_{n-\mathbf{k}\downarrow} - 1, \quad (\text{A.29})$$

$$\hat{\sigma}_{\mathbf{k}}^+ = \hat{c}_{n\mathbf{k}\uparrow}^\dagger \hat{c}_{n-\mathbf{k}\downarrow}^\dagger, \quad (\text{A.30})$$

$$\hat{\sigma}_{\mathbf{k}}^- = \hat{c}_{n-\mathbf{k}\downarrow} \hat{c}_{n\mathbf{k}\uparrow}, \quad (\text{A.31})$$

whose algebra is isomorphic to the spin 1/2 algebra, write $\widehat{\mathcal{H}}_n(t)$, up to a constant factor, as

$$\widehat{\mathcal{H}}_n(t) = \sum_{\mathbf{k}} \left(\tilde{\xi}_{\mathbf{k}}^n \hat{\sigma}_{\mathbf{k}}^z + \Delta_{\mathbf{k}}^n e^{-2ie\Phi(t)/\hbar} \hat{\sigma}_{\mathbf{k}}^+ + \overline{\Delta}_{\mathbf{k}}^n e^{2ie\Phi(t)/\hbar} \hat{\sigma}_{\mathbf{k}}^- \right). \quad (\text{A.32})$$

We thus conclude that the Hamiltonian describing the normal-metal is equivalent, in each $((\mathbf{k}, \uparrow), (-\mathbf{k}, \downarrow))$ subspace, to the Hamiltonian of a two-level system under the action of a circularly polarised field. Moreover, since $\widehat{\mathcal{H}}_s$ is time-independent and commutes with $\widehat{\mathcal{H}}_n(t)$, it does not contribute to (A.22). We can thus write this equation as

$$\mathcal{G}_\beta(v, \tau) = \text{Tr} \left[\widehat{\mathcal{W}}^\dagger(\tau, 0) e^{i\widehat{\mathcal{H}}_n(\tau)v} \widehat{\mathcal{W}}(\tau, 0) e^{-i\widehat{\mathcal{H}}_n(0)v} \hat{\rho}_n(\beta) \right], \quad (\text{A.33})$$

where

$$\widehat{\mathcal{W}}(t, 0) = T \exp \left[-\frac{i}{\hbar} \int_0^t du \widehat{\mathcal{H}}_n(u) \right], \quad (\text{A.34})$$

and $\hat{\rho}_n(\beta) = e^{-\beta\widehat{\mathcal{H}}_n(0)}/\mathcal{Z}_n(\beta)$, with $\mathcal{Z}_n(\beta) = \text{Tr}(e^{-\beta\widehat{\mathcal{H}}_n(0)})$. Therefore, we have shown that there exists an equivalence, in what concerns the calculation of the work characteristic-function in the adiabatic limit, between the dynamics of the NIS junction and that of an assembly of independent two-level systems, subjected to a circularly polarised field. In Appendix B, we will consider the calculation of the work characteristic-function of the latter system.

Appendix B. The work characteristic-function of a two-level system in contact with a thermal bath

We now wish to compute (A.33) where $\widehat{\mathcal{H}}_n(t)$ is given by (A.32). Since the different \mathbf{k} modes are independent, we can restrict the calculation of of the said function to that of a single mode. Furthermore, we will drop the \mathbf{k} label, as no confusion can arise. Since $\tau \gg \tau'$, we can write $\Phi(t) \approx \hbar\omega t/2$, where $\omega = 2eV_{\text{max}}/\hbar$. We will change somewhat the notation with regard to the previous section, so as to keep the result obtained as general as possible, rather than identifying it solely with the dynamics of the NIS junction. In the new notation, the initial Hamiltonian is given by

$$\widehat{\mathcal{H}}_n(0) = \frac{\hbar}{2} \hat{\sigma}^z + \frac{\Gamma}{2} \hat{\sigma}^x, \quad (\text{B.1})$$

where in the NIS junction context $h = 2\tilde{\xi}^n$ and $\Gamma = 2\Delta^n$, and where, without loss of generality, Δ^n can be chosen to be a real-number, as one can always fix the arbitrary superconducting phase φ to be zero.

The quantities h and Γ can be viewed as the components of a constant (pseudo-)magnetic field applied to the two-level system. Since the system is initially in equilibrium with a thermal bath, the system's partition function is given by $\mathcal{Z}_n(\beta) = 2 \cosh(\beta\sqrt{\Gamma^2 + h^2}/2)$.

The time-dependent Hamiltonian $\widehat{\mathcal{H}}_n(t)$ can be written as

$$\widehat{\mathcal{H}}_n(t) = \frac{h}{2} \hat{\sigma}^z + \frac{\Gamma}{2} (\cos(\omega t) \hat{\sigma}^x + \sin(\omega t) \hat{\sigma}^y). \quad (\text{B.2})$$

The dynamics of the two-level system is such that the constant applied field is substituted by a circularly polarised field in $[0, \tau]$, i.e. by a Rabi dynamics during that interval. After the application of the protocol, the circularly polarised field is again replaced by a time-independent field and the system is described by the constant Hamiltonian $\widehat{\mathcal{H}}_n(\tau)$. In the process, the applied field has been rotated by an angle $\theta = \omega\tau$ around the z axis.

It is well known that for the Rabi dynamics, one can write the time-evolution operator $\widehat{\mathcal{U}}(\tau, 0)$ in (A.33) as $\widehat{\mathcal{U}}(\tau, 0) = \widehat{\mathcal{R}}_\tau \widehat{\mathcal{U}}_\tau$, with $\widehat{\mathcal{R}}_\tau = e^{-i\omega\tau\hat{\sigma}^z/2}$ representing a pure rotation around the z axis and $\widehat{\mathcal{U}}_\tau = e^{-i\widehat{\mathcal{H}}'\tau/\hbar}$, where $\widehat{\mathcal{H}}'$ is a time-independent pseudo-Hamiltonian, which depends on ω , and is given by

$$\widehat{\mathcal{H}}' = \frac{1}{2}(h - \hbar\omega) \hat{\sigma}^z + \frac{\Gamma}{2} \hat{\sigma}^x. \quad (\text{B.3})$$

Substituting this expression for $\widehat{\mathcal{U}}(\tau, 0)$ in (A.33) and noting that $\widehat{\mathcal{R}}_\tau^\dagger e^{i\widehat{\mathcal{H}}_n(\tau)u} \widehat{\mathcal{R}}_\tau = e^{i\widehat{\mathcal{R}}_\tau^\dagger \widehat{\mathcal{H}}_n(\tau) \widehat{\mathcal{R}}_\tau u} = e^{i\widehat{\mathcal{H}}_n(0)u}$, since $\widehat{\mathcal{R}}_\tau^\dagger \widehat{\mathcal{H}}_n(\tau) \widehat{\mathcal{R}}_\tau = \widehat{\mathcal{H}}_n(0)$, one obtains for the characteristic-function the result

$$\mathcal{G}_\beta(v, \tau) = \frac{1}{\mathcal{Z}_n(\beta)} \text{Tr} \left[e^{i\widehat{\mathcal{H}}'\tau/\hbar} e^{i\widehat{\mathcal{H}}_n(0)v} e^{-i\widehat{\mathcal{H}}'\tau/\hbar} e^{-i\widehat{\mathcal{H}}_n(0)v} e^{-\beta\widehat{\mathcal{H}}_n(0)} \right]. \quad (\text{B.4})$$

Performing the trace over the complete set of states that diagonalises $\widehat{\mathcal{H}}_n(0)$, we obtain for $\mathcal{G}_\beta(v, \tau)$, the expression

$$\begin{aligned} \mathcal{G}_\beta(v, \tau) &= \frac{1}{\mathcal{Z}_n(\beta)} \sum_{\sigma, \sigma'} | \langle \sigma, \hat{\mathbf{n}} | e^{i\widehat{\mathcal{H}}'\tau/\hbar} | \sigma', \hat{\mathbf{n}} \rangle |^2 \\ &\quad \times e^{i(\sigma' - \sigma)v\sqrt{\Gamma^2 + h^2}/2} e^{-\beta\sigma\sqrt{\Gamma^2 + h^2}/2}, \end{aligned} \quad (\text{B.5})$$

where the unit vector $\hat{\mathbf{n}}$ refers to the direction of the applied field at $t = 0$, its components being given by $\hat{n}_x = \frac{\Gamma}{\sqrt{\Gamma^2 + h^2}}$ and $\hat{n}_z = \frac{h}{\sqrt{\Gamma^2 + h^2}}$. Moreover, writing $\widehat{\mathcal{H}}' = \frac{1}{2}\sqrt{\Gamma^2 + (h - \hbar\omega)^2} (\hat{\mathbf{n}}' \cdot \hat{\boldsymbol{\sigma}})$, where $\hat{n}'_x = \frac{\Gamma}{\sqrt{\Gamma^2 + (h - \hbar\omega)^2}}$ and $\hat{n}'_z = \frac{h - \hbar\omega}{\sqrt{\Gamma^2 + (h - \hbar\omega)^2}}$, we can expand the exponential $e^{i\widehat{\mathcal{H}}'\tau/\hbar}$ as

$$e^{i\widehat{\mathcal{H}}'\tau/\hbar} = \cos\left(\frac{\tau\sqrt{\Gamma^2 + (h - \hbar\omega)^2}}{2\hbar}\right) \mathbb{1}$$

$$\begin{aligned}
& + i (\hat{\mathbf{n}}' \cdot \hat{\mathbf{n}}) \sin \left(\frac{\tau \sqrt{\Gamma^2 + (h - \hbar\omega)^2}}{2\hbar} \right) (\hat{\mathbf{n}} \cdot \hat{\boldsymbol{\sigma}}) \\
& + i \sin \left(\frac{\tau \sqrt{\Gamma^2 + (h - \hbar\omega)^2}}{2\hbar} \right) [\hat{\mathbf{n}} \times (\hat{\mathbf{n}}' \times \hat{\mathbf{n}})] \cdot \hat{\boldsymbol{\sigma}}. \tag{B.6}
\end{aligned}$$

The first two terms of (B.6) are diagonal in the $|\sigma, \hat{\mathbf{n}}\rangle$ basis, whereas the last term is only non-zero when evaluated between two states of the $|\sigma, \hat{\mathbf{n}}\rangle$ basis with opposite spin. We thus obtain that the square moduli of the matrix elements that appear in equation (B.5) are given by

$$\begin{aligned}
|\langle \sigma, \hat{\mathbf{n}} | e^{i\hat{\mathcal{H}}'\tau/\hbar} | \sigma', \hat{\mathbf{n}} \rangle|^2 &= \left[\cos^2 \left(\frac{\tau \sqrt{\Gamma^2 + (h - \hbar\omega)^2}}{2\hbar} \right) \right. \\
& \quad \left. + (\hat{\mathbf{n}}' \cdot \hat{\mathbf{n}})^2 \sin^2 \left(\frac{\tau \sqrt{\Gamma^2 + (h - \hbar\omega)^2}}{2\hbar} \right) \right] \delta_{\sigma, \sigma'} \\
& \quad + |\hat{\mathbf{n}} \times (\hat{\mathbf{n}}' \times \hat{\mathbf{n}})|^2 \\
& \quad \times \sin^2 \left(\frac{\tau \sqrt{\Gamma^2 + (h - \hbar\omega)^2}}{2\hbar} \right) \delta_{\sigma, -\sigma'}. \tag{B.7}
\end{aligned}$$

Taking into account that $|\hat{\mathbf{n}} \times (\hat{\mathbf{n}}' \times \hat{\mathbf{n}})|^2 = 1 - (\hat{\mathbf{n}}' \cdot \hat{\mathbf{n}})^2$ and using the expressions for the components of $\hat{\mathbf{n}}$ and $\hat{\mathbf{n}}'$ given above in (B.7), we finally obtain the expression

$$\begin{aligned}
\mathcal{G}_\beta(v, \tau) &= \left[\cos^2 \left(\frac{\tau \sqrt{\Gamma^2 + (h - \hbar\omega)^2}}{2\hbar} \right) \right. \\
& \quad \left. + \frac{(\Gamma^2 + h^2 - h\hbar\omega)^2}{(\Gamma^2 + h^2)(\Gamma^2 + (h - \hbar\omega)^2)} \cdot \sin^2 \left(\frac{\tau \sqrt{\Gamma^2 + (h - \hbar\omega)^2}}{2\hbar} \right) \right] \\
& \quad + \frac{\Gamma^2 (\hbar\omega)^2}{(\Gamma^2 + h^2)(\Gamma^2 + (h - \hbar\omega)^2)} \cdot \sin^2 \left(\frac{\tau \sqrt{\Gamma^2 + (h - \hbar\omega)^2}}{2\hbar} \right) \\
& \quad \times \frac{\cosh \left[(\beta/2 + iv) \sqrt{\Gamma^2 + h^2} \right]}{\cosh \left(\beta \sqrt{\Gamma^2 + h^2} / 2 \right)}. \tag{B.8}
\end{aligned}$$

It is trivial to check that $\mathcal{G}_\beta(v, \tau)$, as given by (B.8), fulfils both the equality $\mathcal{G}_\beta(0, \tau) = 1$ and $\mathcal{G}_\beta(i\beta, \tau) = 1$, which provides a check on the correctness of the result, since these equalities are built into the definition of $\mathcal{G}_\beta(v, \tau)$ by construction, as pointed out above.

Performing the inverse Fourier transform on $\mathcal{G}_\beta(u, \tau)$, we obtain for $P_\beta(W, \tau)$ the result

$$\begin{aligned}
P_\beta(W, \tau) &= \left[\cos^2 \left(\frac{\tau \sqrt{\Gamma^2 + (h - \hbar\omega)^2}}{2\hbar} \right) + \right. \\
& \quad \left. \frac{(\Gamma^2 + h^2 - h\hbar\omega)^2}{(\Gamma^2 + h^2)(\Gamma^2 + (h - \hbar\omega)^2)} \sin^2 \left(\frac{\tau \sqrt{\Gamma^2 + (h - \hbar\omega)^2}}{2\hbar} \right) \right] \delta(W)
\end{aligned}$$

$$\begin{aligned}
& + \frac{\Gamma^2(\hbar\omega)^2}{(\Gamma^2 + h^2)(\Gamma^2 + (h - \hbar\omega)^2)} \sin^2 \left(\frac{\tau\sqrt{\Gamma^2 + (h - \hbar\omega)^2}}{2\hbar} \right) \\
& \times \frac{e^{\beta W/2}}{2 \cosh(\beta\sqrt{\Gamma^2 + h^2}/2)} \\
& \cdot \left[\delta(W - \sqrt{\Gamma^2 + h^2}) + \delta(W + \sqrt{\Gamma^2 + h^2}) \right]. \tag{B.9}
\end{aligned}$$

The form of (B.9) is easily interpreted from the two-level structure of the problem, as either no work is performed on the system if the time-dependent field does not induce transitions between the levels (first term), or otherwise the time-dependent field induces a transition between the ground state and the excited state, involving an amount of work performed on the system equal to $W = \sqrt{\Gamma^2 + h^2}$, or the inverse transition is induced involving a negative amount of work $W = -\sqrt{\Gamma^2 + h^2}$ being performed on the system (second term). Note that if one sets $W \rightarrow -W$, $\omega \rightarrow -\omega$ and $h \rightarrow -h$, $\Gamma \rightarrow -\Gamma$ in (B.9), one can directly check that $P_\beta(-W, -\tau) = e^{-\beta W} P_\beta(W, \tau)$, i.e. the work distribution satisfies the Crooks-Tasaki relation.

The average work performed on the system during the application of the protocol can be computed either by differentiating (B.8) with respect to v , or directly from (B.9), and is given by

$$\begin{aligned}
\langle W \rangle_\beta & = \frac{\Gamma^2(\hbar\omega)^2}{\sqrt{\Gamma^2 + h^2}(\Gamma^2 + (h - \hbar\omega)^2)} \sin^2 \left(\frac{\tau\sqrt{\Gamma^2 + (h - \hbar\omega)^2}}{2\hbar} \right) \\
& \times \tanh(\beta\sqrt{\Gamma^2 + h^2}/2). \tag{B.10}
\end{aligned}$$

Since the change in the free energy of the system is zero during the transformation, this quantity is equal to the energy dissipated by the system into the thermal bath during the equilibration process occurring after $t > \tau$. One can easily check that, according to (B.10), $\langle W \rangle_\beta$ is always larger or equal to zero in agreement with the second law of thermodynamics. It has a maximum at resonance, i.e. if $\omega = (\Gamma^2 + h^2)/(\hbar h)$ and if $\omega\tau | \Gamma | / \sqrt{\Gamma^2 + h^2} = (2n+1)\pi$, with n being an integer. Note that for a single two-level system, one can, for each value of τ , choose ω such that $\langle W \rangle_\beta = 0$, i.e. one takes ω such that $\tau\sqrt{\Gamma^2 + (h - \hbar\omega)^2}/\hbar = 2n\pi$. The existence of such a minimum can be understood from that fact that for such choice of ω , $e^{i\hat{H}'\tau/\hbar} = \mathbb{1}$, and thus no transitions between the levels are induced by the unitary transformation, see also (B.5). This behaviour regarding the dissipated work in a two-level system mirrors the corresponding behaviour of the Rabi formula for the transition probabilities of such a system under the influence of a circularly polarised field.

The mean-square deviation of the work performed during the application of the protocol can be computed either by differentiating (B.8) twice with respect to v , or directly from (B.9), and is given by

$$\langle (\delta W)^2 \rangle_\beta = \frac{\Gamma^2(\hbar\omega)^2}{(\Gamma^2 + (h - \hbar\omega)^2)} \sin^2 \left(\frac{\tau\sqrt{\Gamma^2 + (h - \hbar\omega)^2}}{2\hbar} \right) \times$$

$$\left[1 - \frac{\Gamma^2(\hbar\omega)^2}{(\Gamma^2 + h^2)(\Gamma^2 + (h - \hbar\omega)^2)} \sin^2 \left(\frac{\tau\sqrt{\Gamma^2 + (h - \hbar\omega)^2}}{2\hbar} \right) \right. \\ \left. \times \tanh \left(\beta\sqrt{\Gamma^2 + h^2}/2 \right) \right]. \quad (\text{B.11})$$

In Sect. 4, we use (B.8) to compute the work dissipated by the NIS junction due to the application of the voltage protocol. Since it is also of interest, albeit not for the solution of the NIS problem, we leave to Appendix C the calculation of the work characteristic-function for an isolated two-level system due to the application of a circularly polarised field to such a system.

Appendix C. The generating function of an isolated two-level system

One can use the results obtained in Appendix B to compute the characteristic function or the work distribution function for an isolated system, described by a micro-canonical ensemble, which undergoes a transformation that is analogous to the one described in that section, i.e. the system is initially isolated and is coupled to the circularly polarised field at $t = 0$, being decoupled from it at $t = \tau$. Such characteristic function is given by an inverse Laplace transform [48] that involves $\mathcal{G}_\beta(v, \tau)$ and the partition function $\mathcal{Z}(\beta) = 2 \cosh(\beta\sqrt{\Gamma^2 + h^2}/2)$

$$\mathcal{G}_E(v, \tau) \omega_0(E) = \int_C \frac{d\beta}{2\pi i} e^{\beta E} \mathcal{G}_\beta(v, \tau) \mathcal{Z}(\beta), \quad (\text{C.1})$$

where E is the energy of the system and $\omega_0(E) = \delta(E - \sqrt{\Gamma^2 + h^2}/2) + \delta(E + \sqrt{\Gamma^2 + h^2}/2)$ is the density of states of the system at $t = 0$. The contour C is the inverse Laplace transform contour from $c - i\infty$ and $c + i\infty$, with c chosen such that all the singularities of the integrand are located to the left of c . In our case, $c = 0$.

Note that these results cannot be generalised to an assembly of two-level systems, since the work characteristic-function $\mathcal{G}_\beta(v, \tau)$ is in this case the product of characteristic-functions for the individual systems.

The integral can be readily performed and after factoring the term $\omega_0(E)$ out, we obtain for $\mathcal{G}_E(v, \tau)$ the result

$$\mathcal{G}_E(v, \tau) = \left[\cos^2 \left(\frac{\tau\sqrt{\Gamma^2 + (h - \hbar\omega)^2}}{2\hbar} \right) + \frac{(\Gamma^2 + h^2 - h\hbar\omega)^2}{(\Gamma^2 + h^2)(\Gamma^2 + (h - \hbar\omega)^2)} \sin^2 \left(\frac{\tau\sqrt{\Gamma^2 + (h - \hbar\omega)^2}}{2\hbar} \right) \right] + \frac{\Gamma^2(\hbar\omega)^2}{(\Gamma^2 + h^2)(\Gamma^2 + (h - \hbar\omega)^2)} \sin^2 \left(\frac{\tau\sqrt{\Gamma^2 + (h - \hbar\omega)^2}}{2\hbar} \right) e^{-2ivE}. \quad (\text{C.2})$$

From this expression, one can obtain, as above, by inverse Fourier transformation of $\mathcal{G}_E(v, \tau)$, the work function distribution for an isolated two-level system. This is given

by

$$\begin{aligned}
 P_E(W, \tau) = & \left[\cos^2 \left(\frac{\tau \sqrt{\Gamma^2 + (h - \hbar\omega)^2}}{2\hbar} \right) + \right. \\
 & \left. \frac{(\Gamma^2 + h^2 - h\hbar\omega)^2}{(\Gamma^2 + h^2)(\Gamma^2 + (h - \hbar\omega)^2)} \sin^2 \left(\frac{\tau \sqrt{\Gamma^2 + (h - \hbar\omega)^2}}{2\hbar} \right) \right] \delta(W) \\
 & + \frac{\Gamma^2(\hbar\omega)^2}{(\Gamma^2 + h^2)(\Gamma^2 + (h - \hbar\omega)^2)} \\
 & \times \sin^2 \left(\frac{\tau \sqrt{\Gamma^2 + (h - \hbar\omega)^2}}{2\hbar} \right) \delta(W + 2E) . \tag{C.3}
 \end{aligned}$$

This result can be easily interpreted if one again notes that either no transitions occur between the two levels and in this case the work performed is zero (first term), or otherwise the work performed is $-2E$ where E is the energy of the initial level (second term). Also, note that $P_{E+W}(-W, -\tau) = P_E(W, \tau)$, which is the version of the Crooks-Tasaki relation appropriate for isolated systems in which the expressions for the micro-canonical density of states corresponding to the initial and to the final Hamiltonian are identical, since the spectrum of the system does not change under application of the work protocol [48].

We can also compute the average work dissipated in the transformation, using the expression for $\mathcal{G}_E(v, \tau)$ or that for $P_E(W, \tau)$. This is given by

$$\langle W \rangle_E = - \frac{2E \Gamma^2(\hbar\omega)^2}{(\Gamma^2 + h^2)(\Gamma^2 + (h - \hbar\omega)^2)} \sin^2 \left(\frac{\tau \sqrt{\Gamma^2 + (h - \hbar\omega)^2}}{2\hbar} \right) . \tag{C.4}$$

Again, one can minimise or maximise this quantity by an appropriate choice of the value of ω . This result may be relevant in the context of quantum phase-shift gates [98] where the transformation can be implemented at finite frequency without generation of dissipated heat.

Finally, we can also compute the mean-square deviation of the work performed during the application of the protocol, using the expression for $\mathcal{G}_E(v, \tau)$ or that for $P_E(W, \tau)$. This quantity is given by

$$\begin{aligned}
 \langle (\delta W)^2 \rangle_E = & \frac{4E^2 \Gamma^2(\hbar\omega)^2}{(\Gamma^2 + h^2)(\Gamma^2 + (h - \hbar\omega)^2)} \sin^2 \left(\frac{\tau \sqrt{\Gamma^2 + (h - \hbar\omega)^2}}{2\hbar} \right) . \tag{C.5} \\
 & \left[1 - \frac{\Gamma^2(\hbar\omega)^2}{(\Gamma^2 + h^2)(\Gamma^2 + (h - \hbar\omega)^2)} \sin^2 \left(\frac{\tau \sqrt{\Gamma^2 + (h - \hbar\omega)^2}}{2\hbar} \right) \right] .
 \end{aligned}$$

References

- [1] Callen H B and Welton T A. Irreversibility and generalized noise. *Phys. Rev.*, 83:34–40, 1951.
- [2] Green M S. Markoff random processes and the statistical mechanics of time-dependent phenomena. *J. Chem. Phys.*, 20:1281, 1952.

- [3] Kubo R. Statistical mechanical theory of irreversible processes i. J. Phys. Soc. Jpn., 12:570586, 1957.
- [4] Einstein A. Investigations on the theory of Brownian movement. Methuen, London, UK, 1 edition, 1926.
- [5] Johnson J B. Thermal agitation of electricity in conductors. Phys. Rev., 32:97–109, 1928.
- [6] Nyquist H. Thermal agitation of electric charge in conductors. Phys. Rev., 32:110–113, 1928.
- [7] Jarzynski C. Nonequilibrium equality for free energy differences. Phys. Rev. Lett., 78:2690–2693, 1997.
- [8] Crooks G E. Nonequilibrium measurements of free energy differences for microscopically reversible markovian systems. J. Stat. Phys., 90:1481–1487, 1998.
- [9] Crooks G E. Path-ensemble averages in systems driven far from equilibrium. Phys. Rev. E, 61:2361–2366, 2000.
- [10] Ritort F. Work fluctuations and transient violations of the second law: perspectives in theory and experiments. Séminaire Poincaré, 2:63–87, 2003.
- [11] Ritort F. Fluctuations in open systems. Physics, 2:43, 2009.
- [12] Jarzynski C. Equalities and inequalities: Irreversibility and the second law of thermodynamics at the nanoscale. Séminaire Poincaré, 15:77–102, 2010.
- [13] Campisi M, Talkner P, and Hänggi P. Colloquium: Quantum fluctuation relations: Foundations and applications. Rev. Mod. Phys., 83:771–791, 2011.
- [14] Campisi M, Talkner P, and Hänggi P. Erratum: Colloquium: Quantum fluctuation relations: Foundations and applications. Rev. Mod. Phys., 83:1653, 2011.
- [15] Crooks G E. On the jarzynski relation for dissipative quantum dynamics. J. Stat. Mech.: Theor. Exp., 2008:P10023, 2008.
- [16] Lowell P J, ONeil G C, Underwood J M, and Ullom J N. Macroscale refrigeration by nanoscale electron transport. App. Phys. Lett., 102:082601, 2013.
- [17] White W K and Meeson P J. Experimental techniques in low-temperature physics. OUP, 4 edition, 2002.
- [18] Pobell F. Matter and methods at low temperatures. Springer, 2 edition, 2007.
- [19] Zwanzig R W. High temperature equation of state by a perturbation method. i. nonpolar gases. J. Chem. Phys., 22:1420–1426, 1954.
- [20] Bochkov G N and Kuzovlev Y E. General theory of thermal fluctuations in nonlinear systems. Sov. Phys. JETP, 45:125–130, 1977.
- [21] Bochkov G N and Kuzovlev Y E. Nonlinear fluctuationdissipation relations and stochastic models in nonequilibrium thermodynamics i. generalized fluctuation-dissipation theorem. Physica A, 106:443–479, 1981.
- [22] Sevick E M, Prabhakar R, Williams S R, and Searles D J. Fluctuation theorems. Annu. Rev. Phys. Chem., 59:60333, 2008.
- [23] Hummer G and Szabo A. Free energy reconstruction from nonequilibrium single-molecule pulling experiments. PNAS, 98:3658 3661, 2001.
- [24] Liphardt J, Dumont S, Smith S B, Tinoco Jr I, and Bustamante C. Equilibrium information from nonequilibrium measurements in an experimental test of jarzynskis equality. Science, 296:1832–1835, 2002.
- [25] Wang G M, Sevick E M, Mittag E, Searles D J, and Evans D J. Experimental demonstration of violations of the second law of thermodynamics for small systems and short time scales. Phys. Rev. Lett., 89:050601, 2002.
- [26] Collin D, Ritort F, Jarzynski C, Smith S B, Tinoco Jr I, and Bustamante C. Verification of the crooks fluctuation theorem and recovery of rna folding free energies. Nature, 437:231–234, 2005.
- [27] Douarche F, Ciliberto S, and Petrosyan A. Estimate of the free energy difference in mechanical systems from work fluctuations: experiments and models. J. Stat. Phys., 2005:P09011, 2005.
- [28] Schuler S, Speck T, Tietz C, Wrachtrup J, and Seifert U. Experimental test of the fluctuation theorem for a driven two-level system with time-dependent rates. Phys. Rev. Lett., 94:180602,

- 2005.
- [29] Junier I, Mossa A, Manosas M, and Ritort F. Recovery of free energy branches in single molecule experiments. Phys. Rev. Lett., 102:070602, 2009.
 - [30] Mossa A, Manosas M, Fornas N, Huguet J M, and Ritort F. Dynamic force spectroscopy of dna hairpins: I. force kinetics and free energy landscapes. J. Stat. Phys., 2009:P02060, 2009.
 - [31] Toyabe S, Sagawa T, Ueda M, Muneyuki E, and Sano M. Experimental demonstration of information-to-energy conversion and validation of the generalized jarzynski equality. Nature Phys., 6:988–992, 2010.
 - [32] Berut A, Arakelyan A, Petrosyan A, Ciliberto S, Dillenschneider R, and Lutz E. Experimental verification of landauers principle linking information and thermodynamics. Nature, 483:187–189, 2012.
 - [33] Jarzynski C. How does a system respond when driven away from thermal equilibrium? PNAS, 98:3636–3638, 2001.
 - [34] Jarzynski C. Nonequilibrium work theorem for a system strongly coupled to a thermal environment. J. Stat. Mech., 2004:P09005, 2004.
 - [35] Cleuren B, Van den Broeck C, and Kawai R. Fluctuation and dissipation of work in a joule experiment. Phys. Rev. Lett., 96:050601, 2006.
 - [36] Palmieri B and Ronis D. Jarzynski equality: Connections to thermodynamics and the second law. Phys. Rev. E, 75:011133, 2007.
 - [37] Campisi M, Talkner P, and Hänggi P. Finite bath fluctuation theorem. Phys. Rev. E, 80:031145, 2009.
 - [38] Seifert U. Stochastic thermodynamics, fluctuation theorems and molecular machines. Rep. Prog. Phys., 75:126001, 2012.
 - [39] Tasaki H. Jarzynski relations for quantum systems and some applications. cond-mat/0009244v2, page unpublished, 2000.
 - [40] Kurchan J. A quantum fluctuation theorem. cond-mat/0007360, page unpublished, 2000.
 - [41] Yukawa S. A quantum analogue of the jarzynski equality. J. Phys. Soc. Jp., 69:2367–2370, 2000.
 - [42] Talkner P, Lutz E, and Hänggi P. Fluctuation theorems: Work is not an observable. Phys. Rev. E, 75:050102(R), 2007.
 - [43] Chernyak V and Mukamel S. Effect of quantum collapse on the distribution of work in driven single molecules. Phys. Rev. Lett., 93:048302, 2004.
 - [44] Allahverdyan A E and Nieuwenhuizen Th M. Fluctuations of work from quantum subensembles: The case against quantum work-fluctuation theorems. Phys. Rev. E, 71:066102, 2005.
 - [45] A. Engel and R. Nolte. Jarzynski equation for a simple quantum system: Comparing two definitions of work. Eur. Phys. Lett., 79:10003, 2007.
 - [46] Mukamel S. Quantum extension of the jarzynski relation: Analogy with stochastic dephasing. Phys. Rev. Lett., 90:170604, 2003.
 - [47] Campisi M, Talkner P, and Hänggi P. Quantum bochkovkuzovlev work fluctuation theorems. Phil. Trans. R. Soc., 369:291–306, 2011.
 - [48] Talkner P, Hänggi P, and Morillo M. Microcanonical quantum fluctuation theorems. Phys. Rev. E, 77:051131, 2008.
 - [49] Talkner P, Morillo M, Yi J, and Hänggi P. Statistics of work and fluctuation theorems for microcanonical initial states. N. J. Phys., 15:095001, 2013.
 - [50] Campisi M, Talkner P, and Hänggi P. Fluctuation theorem for arbitrary open quantum systems. Phys. Rev. Lett., 102:21040, 2009.
 - [51] Esposito M and Mukamel S. Fluctuation theorems for quantum master equations. Phys. Rev. E, 73:046129, 2009.
 - [52] Talkner P and Hänggi P. The tasakicrooks quantum fluctuation theorem. J. Phys. A: Math. Theor., 40:F569F571, 2007.
 - [53] Deffner S and Lutz E. Nonequilibrium work distribution of a quantum harmonic oscillator. Phys. Rev. E, 77:021128, 2008.

- [54] Silva A. Statistics of the work done on a quantum critical system by quenching a control parameter. *Phys. Rev. Lett.*, 101:120603, 2008.
- [55] Talkner P, Burada P S, and Hänggi P. Statistics of work performed on a forced quantum oscillator. *Phys. Rev. E*, 78:011115, 2008.
- [56] Campisi M, Talkner P, and Hänggi P. Thermodynamics and fluctuation theorems for a strongly coupled open quantum system: an exactly solvable case. *J. Phys. A: Math. Theor.*, 42:392002, 2009.
- [57] Talkner P, Burada P S, and Hänggi P. Erratum: Statistics of work performed on a forced quantum oscillator. *Phys. Rev. E*, 79:039902, 2009.
- [58] Campisi M, Talkner P, and Hänggi P. Fluctuation theorems for continuously monitored quantum fluxes. *Phys. Rev. Lett.*, 105:140601, 2009.
- [59] Deffner S, Abah O, and Lutz E. Quantum work statistics of linear and nonlinear parametric oscillators. *Chem. Phys.*, 375:200–208, 2010.
- [60] Averin D V and Pekola J P. Statistics of the dissipated energy in driven single-electron transitions. *Eur. Phys. Lett.*, 96:67004, 2011.
- [61] Kawamoto T and Hatano N. Test of fluctuation theorems in non-markovian open quantum systems. *Phys. Rev. E*, 84:031116, 2011.
- [62] Yi J and Talkner P. Work statistics of charged noninteracting fermions in slowly changing magnetic fields. *Phys. Rev. E*, 83:041119, 2011.
- [63] Gambassi A and Silva A. Large deviations and universality in quantum quenches. *Phys. Rev. Lett.*, 109:250602, 2012.
- [64] Ngo V A and Haas S. Demonstration of jarzynskis equality in open quantum systems using a stepwise pulling protocol. *Phys. Rev. E*, 86:031127, 2012.
- [65] Smacchia P and Silva A. Universal energy distribution of quasiparticles emitted in a local time-dependent quench. *Phys. Rev. Lett.*, 109:037202, 2012.
- [66] Smacchia P and Silva A. Work distribution and edge singularities for generic time-dependent protocols in extended systems. *Phys. Rev. E*, 88:042109, 2013.
- [67] Solinas P, Averin D V, and Pekola J P. Work and its fluctuations in a driven quantum system. *Phys. Rev. B*, 87:060508(R), 2013.
- [68] Sotiriadis S, Gambassi A, and Silva A. Statistics of the work done by splitting a one-dimensional quasicondensate. *Phys. Rev. E*, 87:052129, 2013.
- [69] Marino J and Silva A. Nonequilibrium dynamics of a noisy quantum ising chain: Statistics of work and prethermalization after a sudden quench of the transverse field. *Phys. Rev. B*, 89:024303, 2014.
- [70] Khaymovich I M, Koski J V, Saira O-P, Kravtsov V E, and Pekola J P. Multifractality of random eigenfunctions and generalization of jarzynski equality. *Nat. Comm.*, 6:7010, 2015.
- [71] Campisi M, Pekola J, and Fazio R. Nonequilibrium fluctuations in quantum heat engines: theory, example, and possible solid state experiments. *N. J. Phys.*, 17:035012, 2014.
- [72] Huber G, Schmidt-Kaler F, Deffner S, and Lutz E. Employing trapped cold ions to verify the quantum jarzynski equality. *Phys. Rev. Lett.*, 101:070403, 2008.
- [73] Heyl M and Kehrein S. Crooks relation in optical spectra: universality in work distributions for weak local quenches. *Phys. Rev. Lett.*, 108:190601, 2012.
- [74] Dorner R, Clark S R, Heaney L, Fazio R, Goold J, and Vedral V. Extracting quantum work statistics and fluctuation theorems by single-qubit interferometry. *Phys. Rev. Lett.*, 110:230601, 2013.
- [75] Mazzola L, De Chiara G, and Paternostro M. Measuring the characteristic function of the work distribution. *Phys. Rev. Lett.*, 110:230602, 2013.
- [76] Campisi M, Blattmann R, Kohler S, Zueco D, and Hänggi P. Employing circuit qed to measure non-equilibrium work fluctuations. *N. J. Phys.*, 15:105028, 2013.
- [77] Batalhao T B, Souza A M, Mazzola L, Auccaise R, Sarthour R S, Oliveira I S, Goold J, De Chiara G, Paternostro M, and Serra R M. Experimental reconstruction of work distribution and study

- of fluctuation relations in a closed quantum system. Phys. Rev. Lett., 89:024303, 2014.
- [78] Pekola J P, Solinas P, Shnirman A, and Averin D V. Calorimetric measurement of work in a quantum system. N. J. Phys., 15:115006, 2013.
- [79] Bardeen J. Tunneling from a many-particle point of view. Phys. Rev. Lett., 6:57–59, 1961.
- [80] Cohen M H, Falicov L M, and Phillips J C. Superconductive tunneling. Phys. Rev. Lett., 8:316–318, 1962.
- [81] Ambegaokar V and Baratoff A. Tunneling between superconductors. Phys. Rev. Lett., 10:486–489, 1963.
- [82] Ambegaokar V and Baratoff A. Tunneling between superconductors (errata). Phys. Rev. Lett., 11:104, 1963.
- [83] McMillan W L. Tunneling model of the superconducting proximity effect. Phys. Rev., 175:537–542, 1968.
- [84] Blonder G E, Tinkham M, and Klapwijk T M. Transition from metallic to tunneling regimes in superconducting microconstrictions: Excess current, charge imbalance, and supercurrent conversion. Phys. Rev. B, 25:4515–4532, 1982.
- [85] Cuevas J C, Martín-Rodero A, and Yeyati A L. Hamiltonian approach to the transport properties of superconducting quantum point contacts. Phys. Rev. B, 54:7366–7379, 1996.
- [86] Wolf E L. Principles of electron tunneling spectroscopy. Oxford University Press, Oxford, UK, 2 edition, 2011.
- [87] Snow E C. Self-consistent band structure of aluminum by an augmented-plane-wave method. Phys. Rev., 158:683–688, 1967.
- [88] Smrca L. Energy band structure of aluminium by the augmented plane wave method. Ch. J. Phys. B, 20:291–300, 1970.
- [89] Matthias B T, Geballe T H, and Compton V B. Superconductivity. Rev. Mod. Phys., 15:1–22, 1963.
- [90] Tinkham M. Introduction to superconductivity. Dover Editions, 2 edition, 2004.
- [91] Callen H B. Thermodynamics: an introduction to the physical theories of equilibrium thermostatics and irreversible thermodynamics. Wiley, 1 edition, 1960.
- [92] Rogovin D and Scalapino D J. Fluctuation phenomena in tunnel junctions. Ann. Phys., 86:1–90, 1974.
- [93] Foldy L L and Wouthuysen S A. On the dirac theory of spin 1/2 particles and its non-relativistic limit. Phys. Rev., 78:2936, 1950.
- [94] Schrieffer J R and Wolff P A. Relation between the anderson and kondo hamiltonians. Phys. Rev., 149:491492, 1966.
- [95] Wegner F. Flow-equations for hamiltonians. Ann. Phys. (Leipzig), 3:77, 1994.
- [96] Kehrein S K and Mielke A. Theory of the anderson impurity model: The schrieffer wolff transformation reexamined. Ann. Phys., 252:1–32, 1996.
- [97] Donabidowicz A and Domański T. The in-gap charge current through the correlated quantum dot hybridized with superconductor. Act. Phys. Pol. A, 114:75–82, 2008.
- [98] Martinis J M, Nam S, Aumentado J, and Urbina C. Rabi oscillations in a large josephson-junction qubit. Phys. Rev. Lett., 89:117901, 2002.