

The thermodynamics of protein aggregation reactions may underpin the enhanced metabolic efficiency associated with heterosis, some balancing selection, and the evolution of ploidy levels

B.R. Ginn¹

Abstract

Identifying the physical basis of heterosis (or “hybrid vigor”) has remained elusive despite over a hundred years of research on the subject. The three main theories of heterosis are dominance theory, overdominance theory, and epistasis theory. Kacser and Burns (1981) identified the molecular basis of dominance, which has greatly enhanced our understanding of its importance to heterosis. This paper aims to explain how overdominance, and some features of epistasis, can similarly emerge from the molecular dynamics of proteins. Possessing multiple alleles at a gene locus results in the synthesis of different allozymes at reduced concentrations. This in turn reduces the rate at which each allozyme forms soluble oligomers, which are toxic and must be degraded, because allozymes co-aggregate at low efficiencies. The model developed in this paper can explain how heterozygosity impacts the metabolic efficiency of an organism. It can also explain why the viabilities of some inbred lines seem to decline rapidly at high inbreeding coefficients ($F > 0.5$), which may provide a physical basis for truncation selection for heterozygosity. Finally, the model has implications for the ploidy level of organisms. It can explain why polyploids are frequently found in environments where severe physical stresses promote the formation of soluble oligomers. The model can also explain why complex organisms, which need to synthesize aggregation-prone proteins that contain intrinsically unstructured regions (IURs) and multiple domains because they facilitate complex protein interaction networks (PINs), tend to be diploid while haploidy tends to be restricted to relatively simple organisms.

Keywords: heterosis, truncation selection, ploidy, protein interaction network, thermodynamics

1. Introduction

Heterosis, or “hybrid vigor”, refers to the superior performance of highly heterozygous individuals relative to less heterozygous individuals on a number of biological metrics (Lippman and Zamir 2007; Hochholdinger and Hoecker 2007; Birchler et al. 2010). Biologists have been aware of the phenomena for over a hundred years (e.g. Darwin 1876), and it has been exploited to improve crop yields substantially over the twentieth century, especially in maize (Crow 1998; Duvick 2001), but there is still debate over its origin (for review see Lippman and Zamir 2007; Hochholdinger and Hoecker 2007; Birchler et al. 2010). Three different theories are usually used to explain heterosis: dominance theory (Davenport 1908; Bruce 1910; Jones 1917), overdominance theory (Shull 1948; East 1936), and epistasis (Powers 1944; Williams

1959). Dominance theory attributes the benefits of heterozygous genotypes to the masking of recessive deleterious alleles. Proponents of overdominance theory argue that heterozygosity itself can have benefits, even in the absence of deleterious mutations. Proponents of epistasis argue that heterosis comes from positive interactions between multiple gene loci. Currently, dominance theory is the most widely accepted theory of heterosis (Crow 1998; Charlesworth and Willis 2009).

However, dominance theory is unlikely to be the sole explanation for heterosis. Five lines of evidence from the literature for overdominance and epistasis are briefly mentioned here. First, the performance of hybrid rice cannot be explained solely by dominance theory, and may require some combination of overdominant and epistatic interactions (Li et al. 2001; Zhou et al. 2012). Second, breeding experiments performed on polyploid plants appear to indicate that possessing three or more alleles at a gene locus is more beneficial

Email address: bginn3@gmail.com (B.R. Ginn)

than possessing two, which is difficult to explain with dominance theory alone (Groose et al. 1989; Riddle and Birchler 2008). Third, the rate of heterozygosity decline that occurs after multiple generations of inbreeding is slower than predicted by dominance theory (Rumball et al. 1994; Demontis et al. 2009; Chelo and Teotónio 2012). The slow decline may reflect balancing selection (selection for overdominant loci) or linkage between the measured genetic markers and deleterious alleles (associative overdominance). Fourth, haplodiploid (Henter 2003; Tortajada et al. 2009) and selfing species (Husband and Schemske 1996) show greater degrees of inbreeding depression than would be expected if deleterious recessive alleles alone were responsible (although numerous very mildly deleterious alleles may explain these findings). Finally, several authors have found evidence of heterozygosity-fitness correlations (HFC's) in wild populations, which are taken as evidence of overdominance (Lesica and Allendorf 1992; Silva et al. 2006; Ferreira and Amos 2006; Mäkinen et al. 2008; Hoffman et al. 2010).

One of the appeals of the dominance theory of heterosis is that its mechanisms have firm theoretical foundations. Population genetics theory predicts that recessive deleterious mutations should be common in populations (Charlesworth and Willis 2009). Furthermore, Kacser and Burns (1981) presented theoretical and experimental results that show why deleterious mutations are frequently recessive.

In contrast, there is no widely accepted mechanism for overdominance and selection for heterozygous genotypes (Charlesworth and Willis 2009; Zhou et al. 2012). This paper attempts to provide such a mechanism, which will provide an explanation for how heterozygosity can result in heterosis even in the absence of deleterious mutations. An interesting feature of the theory presented in this paper is that overdominant loci are inherently epistatic, which may be a reason why papers that find evidence for overdominance also tend to find evidence for epistasis (Li et al. 2001; Zhou et al. 2012).

The central hypothesis of this paper, that the specificity of protein aggregation reactions provide a physical basis for overdominance and heterozygous advantage, is supported by the findings of previous papers that relate heterozygosity to protein metabolism. The earliest papers, such as Koehn and Shumway (1982) and Hawkins et al. (1986), found that both metabolic efficiency and protein turnover increase with decreasing heterozygosity in marine bivalves. More recently, Kristensen et al. (2002) and Pedersen et al. (2005) found that inbred lines of *Drosophila*

melanogaster produce higher concentrations of molecular chaperones than outbred lines, which they hypothesized was due to higher rates of protein aggregation. Both Kristensen et al. (2002) and Goff (2011) argued that these previous findings can be explained if homozygosity for deleterious mutations leads to greater expression of unstable proteins by inbred individuals. In contrast, Ginn (2010) attempted to show that the previous findings could be explained by an overdominance model if protein aggregation is assumed to be a highly specific process (see below). Mead et al. (2003) had already shown that the specificity of prion amyloid formation resulted in balancing selection at the prion protein gene. However, Mead et al. (2003) and Ginn (2010) only considered the benefits of heterozygosity at a single gene locus. This paper will provide a biochemical explanation for how heterozygosity at multiple gene loci can lead to truncation selection, which can maintain protein polymorphisms at numerous gene loci in natural populations (King 1967; Milkman 1967; Sved et al. 1967).

One of the benefits of this paper's theoretical approach is that it can potentially explain several trends in the ploidy level of organisms. The reason why organisms have different ploidy levels is still poorly understood (Otto and Gerstein 2008; Otto and Whitton 2000; Mable 2004; Madlung 2013). Yet, the evolution of different ploidy levels is important to any theory of heterosis since heterozygosity cannot exist in strictly haploid organisms. Most theories that attempt to explain the evolution of higher ploidy levels have focused on the masking of deleterious recessive alleles or have compared the rates of evolution of organisms with different ploidy levels (see Orr and Otto 1994, Otto 2007, and Otto and Gerstein 2008 for review). The theory developed in this paper will instead focus on how higher ploidy levels can help organisms cope with protein aggregation. The theory can potentially explain four different trends: 1) the frequent occurrence of polyploid organisms in harsh environments, 2) the restriction of haploidy to relatively simple organisms, 3) the relative stress tolerances of plant gametophytes and sporophytes, and 4) why complexity (and diploidy) is associated with the production of sexual spores in plants and fungi. Thus, the theory presented in this paper attempts to provide a complete theory of heterozygous advantage that unifies our understanding of heterosis, selection for heterozygosity, and ploidy level.

2. Metabolic Heterosis

2.1. Inbreeding and Metabolic Efficiency

Numerous studies have been published on a phenomena that I will call "metabolic heterosis" in this paper. Metabolic heterosis is the observed correlation between the growth rate of organisms and their heterozygosities as measured by allozyme or microsatellite markers (Koehn and Shumway 1982; Garton et al. 1984; Mitton 1985; Danzmann et al. 1987; Mitton 1993; Hedgecock et al. 1996; Pogson and Fevolden 1998; Bayne et al. 1999; Hawkins and Day 1999; Hawkins et al. 2000; Bayne 2004; Borrell et al. 2004; Pujolar et al. 2005; Ketola and Kotiaho 2009). Some of these authors maintain that their correlations indicate a relationship between heterozygosity and metabolic efficiency (Garton et al. 1984; Mitton 1993; Mitton 1997; Pogson and Fevolden 1998; Borrell et al. 2004). While most studies have focused on the size and mass of the organism, metabolic efficiency can affect other fitness traits. For instance, Gajardo and Beardmore (1989) and Gajardo et al. (2001) have shown a positive correlation between heterozygosity and the percentage of female *Artemia* that produce energetically expensive encysted offspring rather than energetically cheaper nauplii. While many studies have concluded that heterozygosity is correlated with the metabolic efficiency of organisms, there is still no consensus view on the underlying mechanism behind this correlation. Furthermore, there is an additional debate over whether the fitness parameters are correlated with the heterozygosity of the organism across all gene loci (the "general effect" hypothesis) or only with the heterozygosity of gene loci near the measured genetic markers (the "local effect" hypothesis) (Mitton and Pierce 1980; Balloux et al. 2004; Szulkin et al. 2010).

The correlation between heterozygosity and metabolic efficiency may be explained by protein turnover (Hawkins et al. 1986; Hawkins et al. 1989; Hedgecock et al. 1996; Bayne 2004). Hawkins et al. (1986) showed that lower heterozygosity leads to higher levels of protein turnover in the blue mussel, *Mytilus edulis*, using ¹⁵N labeled food and allozyme markers. Protein turnover refers to an organism's daily degradation and synthesis of proteins, both of which are energy consuming processes. Therefore, these papers argued, an inbred organism's biomass may be more energetically expensive to sustain than an outbred organism's due to higher levels of protein turnover.

One of the processes that enhances protein turnover is protein aggregation. Most proteins consist of




Figure 1: An unfolded polypeptide chain may either refold into its native conformation or bind to another unfolded polypeptide chain. The two reactions compete against each other, and the relative rates of each reaction will determine the folding efficiency of the polypeptide chain. The shaded areas on the polypeptide chain represent binding sites that stabilize the folded protein or soluble oligomer.

polypeptide chains that must fold into a native conformation in order to be functional. Many proteins, especially meta-stable proteins containing intrinsically unstructured regions (IURs), are continuously unfolding and refolding in an organism (Olzscha et al. 2011; also see Section 3.4 below). However, as shown in Figure 1, there are two alternative pathways that unfolded polypeptide chains may take. First, the polypeptide chain may refold into its correct conformation and become a functional protein. Alternatively, the polypeptide chain may bind with another unfolded chain and form a soluble oligomer (Silow and Oliveberg 1997; Bitan et al. 2001; Idicula-Thomas and Balaji 2007; Kaye et al. 2003; Kaye et al. 2004; Cleary et al. 2005; Haass and Selkoe 2007; Vieira et al. 2007; Wei et al. 2007). Soluble oligomers can then bind with additional unfolded chains and eventually become a solid protein aggregate. The formation of soluble oligomers and solid protein aggregates is detrimental for two reasons. First, protein aggregation competes with the proper folding of a protein (Kiefhaber et al. 1991). A protein's folding efficiency decreases when more unfolded polypeptide chains bind to each other. Second, soluble oligomers and solid protein aggregates are cytotoxic species that have been associated with several disorders (Kaye et al. 2003; Haass and Selkoe 2007;

Vieira et al. 2007; Shankar et al. 2008). Therefore, the viability of organisms depends on the ability of their proteins to maintain their correct conformations and avoid aggregation.

For this reason, all organisms produce numerous molecular chaperones that prevent unfolded polypeptide chains from aggregating. Molecular chaperones can bind to unfolded polypeptide chains, thereby preventing their aggregation, or they can tag the polypeptide chains with ubiquitin, which marks the polypeptide chains for destruction by the proteasome (Hayes and Dice 1996; Kopito 2000; Maurizi 2002; McClellan et al. 2005; Kaganovich et al. 2008). The proteasomal system also degrades proteins after they have aggregated (Dougan et al. 2006; Rubinsztein 2006; Liberek et al. 2008; Tetzlaff et al. 2008). In addition, protein aggregates may also be degraded via autophagy, whereby aggregated proteins are transported to lysosomes and digested (Kopito 2000; Garcia-Mata et al. 2002; Kruse et al. 2006; Yorimitsu and Klionsky 2007). Choe and Strange (2008) observed that half of the genes up-regulated when the nematode *C. elegans* is exposed to aggregate promoting environmental stresses are associated with protein degradation. Especially up-regulated were genes associated with proteasomal and lysosomal degradation.

Both the molecular chaperones that prevent aggregation and the degradation pathways that destroy soluble oligomers (and protein aggregates) must consume ATP while functioning. In addition, new polypeptide chains will have to be synthesized in order to take the place of degraded chains, so high rates of protein aggregation will also result in greater energy expenditure for protein synthesis. Thus, an organism's efforts to maintain protein homeostasis will generally result in high levels of protein turnover and an associated energy expenditure.

The expression of molecular chaperones is correlated with the heterozygosity of organisms. Kristensen et al. (2002) and Pedersen et al. (2005), using an enzyme-linked immunosorbant assay, found that inbred fruit flies (*D. melanogaster* and *Drosophila buzzati*) synthesized more heat shock proteins (*Hsps*) than outbred fruit flies at benign and elevated temperatures. Since *Hsps* are a type of molecular chaperone, the authors of these papers concluded that inbred fruit flies contain a higher number of unfolded or misfolded polypeptide chains, and potentially higher rates of protein aggregation, than outbred fruit flies, even at benign temperatures. Thus, low levels of heterozygosity may result in higher levels of protein aggregation, which in turn can result in higher rates of protein turnover and

lower metabolic efficiencies.¹

Kristensen et al. (2002) and Kristensen et al. (2009) used the dominance theory of inbreeding depression to explain the correlation between heterozygosity and lower *Hsp* concentrations. They argued that proteins encoded by deleterious recessive alleles may be less stable, and more prone to aggregation, than the proteins encoded by normal alleles. Consequently, the increased expression of deleterious recessive alleles by inbred organisms may increase their demand for molecular chaperones. The following subsection will develop an overdominance theory to explain the correlations between heterozygosity and *Hsp* concentrations, protein turnover, and metabolic efficiency. The overdominant explanation will then be extended in subsequent sections to provide a biochemical basis for truncation selection that favors heterozygosity. Afterwards, the truncation selection model will be used to explain why higher ploidy levels are advantageous in certain circumstances.

2.2. Model

A heuristic model is developed in this subsection that shows how an organism's heterozygosity can influence its expression of molecular chaperones, protein turnover, and metabolic efficiency. This model shows how protein aggregation reactions can provide a physical basis for overdominance and metabolic heterosis, which may explain the results of some breeding experiments (Li et al. 2001; Zhou et al. 2012). The model will also show that the relationship between heterozygosity and metabolic efficiency should be linear (additive epistasis) as described in several studies (Koehn and Shumway 1982; Garton et al. 1984; Mitton and Grant 1984; Hawkins et al. 1986). This contrasts with the usual exponential relationship between heterozygosity and phenotype anticipated by multiplicative epistasis (Charlesworth and Willis 2009).

The model will assume that an organism maintains steady-state concentrations of functional proteins that are continuously unfolding and either refolding or forming soluble oligomers. Soluble oligomers must be degraded by the organism when they form because they are toxic. Also, the organism must synthesize new proteins to replace those that were removed when the organism destroyed its soluble oligomers. The result is three steady-state concentrations, $[F]_{steady}$, $[U]_{steady}$, and $[O]_{steady}$, which are the concentrations of

¹Chen et al. (2006) has supported these findings with similar experiments performed on Pacific Abalone populations.

folded protein, unfolded protein, and soluble oligomer, respectively. In this model, it will be assumed that the organism maintains soluble oligomers at a critical steady-state concentration. If the steady-state concentration of soluble oligomers rises, then the organism will increase its concentration of molecular chaperones to lower the soluble oligomers' steady-state concentration back to their critical level. This may be accomplished through a feedback mechanism, such as the unfolded protein response (UPR) that occurs inside the endoplasmic reticulum (Schröder and Kaufman 2005; Bernales et al. 2006). As a consequence of the steady-state assumption, the rates of soluble oligomer formation and degradation will primarily depend on the rate that unfolded polypeptide chains are introduced into the system (see Equation 10 below), which is similar to the *in vivo* model described in Kiefhaber et al. (1991).

Another assumption for the model is that the initial binding reactions are the rate-limiting step in the various protein aggregation pathways, and that molecular chaperones prevent the accumulation of protein aggregation products “downstream” from the initial binding reactions (see Dobson (2003) for an overview of the many aggregation pathways that can occur). This approach has been used by other researchers to successfully model protein aggregation dynamics both *in vitro* and *in vivo* (Kiefhaber et al. 1991; Hasegawa et al. 1999; Borgia et al. 2013). As a consequence of this assumption, the kinetics of soluble oligomer formation will follow second-order rate laws in the model.

The first thing to consider is the rate at which an unfolded polypeptide chain folds into its native conformation. The polypeptide chain may be unfolded because it has been recently synthesized or because a previously native protein unraveled. The latter process may be part of the protein's normal condition (perhaps because it contains intrinsically unstructured regions) or may be induced by environmental stress. Regardless, most unfolded polypeptide chains must fold into their correct conformation in order to be functional. This takes time, especially if the folding chain becomes trapped in a metastable intermediate state (Onuchic et al. 1995; Levy et al. 2005; Nevo et al. 2005). Nevertheless, folding (and refolding) proceeds according to a first-order rate law (Kiefhaber et al. 1991).

$$\frac{d[N]}{dt} = k_f[U] \quad (1)$$

where $[N]$ is the concentration of native protein, t is time

k_f is the rate constant for the folding reaction, and $[U]$ is the concentration of unfolded polypeptide chains.

Alternatively, the unfolded polypeptide chain may bind to another and form a soluble oligomer, which may serve as a seed for protein aggregation. The process of protein aggregation is highly specific in that protein aggregates are highly enriched with a single protein species, even when two or more polypeptides are aggregating simultaneously (London et al. 1974; Speed et al. 1996; Kopito 2000; Rajan et al. 2001; Morell et al. 2008). The process may be so specific that small differences in amino acid sequence can inhibit co-aggregation of different polypeptide chains. For example, Mead et al. (2003), O’Nuallain et al. (2004), and Apostol et al. (2010) have found that a single point mutation can prevent amyloid fibrils from co-aggregating. Other researchers have similarly found that amyloid formation can be inhibited in mixtures of polypeptide variants (Hasegawa et al. 1999; Rochet et al. 2000; Lashuel et al. 2003; Yagi et al. 2005; Lewis et al. 2006; Tahiri-Alaoui et al. 2006).

Both O’Nuallain et al. (2004) and Apostol et al. (2010) have argued that the specificity of amyloid formation may be due to changes in amyloid conformation brought about by point mutations. King et al. (1996), Sinha and Nussinov (2001), and Xu et al. (2013) have shown that point mutations can change the probabilities of unfolded polypeptide chains assuming particular conformations without altering the stability or conformation of the native protein. Thus, point mutations may reduce the probability of two allozymes assuming similar conformations, which would decrease the likelihood of them aligning properly in order to co-aggregate (O’Nuallain et al. 2004; Ma and Nussinov 2012). In addition, point mutations may introduce incompatibilities (e.g. steric repulsion between side chains) that prevent two polypeptide chains from co-aggregating (Apostol et al. 2010), though this would seem to be a less general mechanism. Note that some polymorphisms do co-aggregate (see Wright et al. (2005) and Krebs et al. (2004)), but they are not likely to persist in natural populations since they do not confer any heterozygous advantage. Also, the specificity of protein aggregation may only apply to proteins with certain physical properties. This paper will discuss the importance of proteins with intrinsically unstructured regions (IURS) in Sections 3.4 and 4.2. However, this is not a major limitation since balancing selection would only apply to a fraction of all protein coding gene loci.

The specificity of protein aggregation implies that the formation of soluble oligomers proceeds as a second order reaction (Kiefhaber et al. 1991; Bitan et al. 2001;

Zhdanov and Kasemo 2004; Zhu et al. 2010). For example, the rate law for the formation of a dimer will be:

$$\frac{d[O]}{dt} = k_b[U]^2 \quad (2)$$

where $[O]$ is the concentration of the soluble oligomer and k_b is the rate constant for the self-binding reaction.

A comparison of Equations 1 and 2 reveals that the rate of soluble oligomer formation is more dependent upon the concentration of unfolded polypeptide chains than is the rate of protein folding. For example, an individual that is heterozygous at a given gene locus will synthesize two different allozymes, which should have approximately half the concentration that they would have if the individual were homozygous at the gene locus. From Equation 1, it follows that each allozyme will fold half as fast, and the combined rate of folding will be about the same for heterozygous and homozygous individuals. This can be expressed generally as:

$$\frac{d[N]}{dt} = \sum r_i k_{fi} [U_i] \quad (3)$$

where r_i is the actual concentration of allozyme i in an individual divided by the concentration of i if the individual were homozygous for i (r_i will have a value of 1 in an organism that is homozygous for i , a value of 0 in an organism that does not produce i , and a value of ≈ 0.5 in an organism that is heterozygous for i).

In contrast, soluble oligomers of an allozyme will form at one-quarter the rate in an individual that is heterozygous for the allozyme than in an individual that is homozygous for the allozyme ($\text{rate}_{het} \approx 0.25 \text{rate}_{hom}$). The combined rate of soluble oligomer formation for allozymes in heterozygous individuals will be approximately half the rate for homozygous individuals ($0.25 \text{rate}_{hom} + 0.25 \text{rate}_{hom} = 0.5 \text{rate}_{hom}$), or more generally:

$$\frac{d[O]}{dt} = \sum r_i^2 k_{bi} [U_i]^2 \quad (4)$$

Equations 1–4 show that diluting the concentration of an unfolded polypeptide chain shifts the competition between protein folding and soluble oligomer formation in favor of folding. Thus, heterozygosity increases the folding efficiency of unfolded polypeptide chains simply by diluting their concentrations.

Another way to think of the influence that heterozygosity has on soluble oligomer formation is to consider the number of collisions that will occur in a given time period. In a homozygous

organism, all of the collisions will be between the same allozyme. In contrast, 50% of the collisions in a heterozygous organism will be between alternate allozymes. Each allozyme buffers the soluble oligomer formation reaction of the other. The critical assumption is that soluble oligomer formation is highly specific, which is corroborated by the research papers cited above.

Equation 4 gives the rate of dimer formation at a single gene locus. The total rate of soluble oligomer addition for all alleles at all gene loci is:

$$\begin{aligned} \frac{dA}{dt} = & \sum_j \sum_i k_{b_{1ji}} r_{U_{ji}}^2 [U_{ji}]^2 \\ & + \sum_j \sum_i k_{b_{2ji}} r_{D_{ji}} r_{U_{ji}} [D_{ji}] [U_{ji}] \\ & + \sum_j \sum_i k_{b_{3ji}} r_{T_{ji}} r_{U_{ji}} [T_{ji}] [U_{ji}] \dots \end{aligned} \quad (5)$$

where $[U_{ji}]$ is the concentration of unfolded polypeptide chain expressed by each allele i at each gene locus j , $[D]$ is the concentration of dimer, $[T]$ is the concentration of trimer, r_{ji} is the concentration of a chemical species in an organism that is heterozygous for ji divided by its concentration in an organism that is homozygous for ji , and $k_{b_{ji}}$ is the rate law constant for the binding reaction of each unfolded polypeptide chain encoded by each allele at each gene locus.

For the purpose of simplicity, it is assumed in the rest of this subsection that a single molecular chaperone is responsible for removing soluble oligomers. The rate of removal can be expressed in terms of the Michaelis-Menten Equation for the molecular chaperone (Kondepudi 2008):

$$\frac{dR}{dt} = \frac{R_{max} [O]_{steady}}{K_m + [O]_{steady}} \quad (6)$$

where R_{max} is the chaperone's maximum rate of removal, $[O]_{steady}$ is the steady-state concentration of soluble oligomer, and K_m is the Michaelis-Menten constant for the molecular chaperone. The value of R_{max} is directly proportional to the concentration of molecular chaperone:

$$R_{max} = k_2 M_t \quad (7)$$

where M_t is the total concentration of molecular chaperone and k_2 is the rate law constant for the molecular chaperone's catalyzing reaction. Combining Equations 5, 6, and 7 and solving for M_t gives:

$$M_t = \left(\frac{K_m + [O]_{steady}}{k_2[O]_{steady}} \right) \frac{dA}{dt} \quad (8)$$

Equation 8 gives the concentration of molecular chaperone necessary to maintain a particular steady-state concentration of soluble oligomers at a given heterozygosity. It predicts that inbred organisms will have higher concentrations of molecular chaperones than outbred organisms, which has been confirmed by Kristensen et al. (2002).

The degradation of soluble oligomers must be balanced by the synthesis of new proteins in order to maintain a steady-state concentration of properly functioning protein within the cytosol:

$$\frac{dP}{dt} = \nu \frac{dR}{dt} = \nu \frac{dA}{dt} \quad (9)$$

where dP/dt is the rate that new proteins are synthesized to replace proteins that have formed soluble oligomers and ν is the number of proteins making up the soluble oligomers. Equations 5, 6, and 9 predict that inbred organisms should have higher rates of protein turnover than outbred organisms, which has been confirmed by Hawkins et al. (1986).

Finally, the removal of soluble oligomers and their replacement with new proteins is an energy consuming process. The overall calorie consumption rate due to protein maintenance, dC_{Maint}/dt , is:

$$\begin{aligned} \frac{dC_{Maint}}{dt} = & \sum_j \sum_i (\beta_{ji} + 2\gamma_{ji}) k_{b_1,ji} r_{U,ji}^2 [U_{ji}]^2 \\ & + \sum_j \sum_i (\beta_{ji} + 3\gamma_{ji}) k_{b_2,ji} r_{D,ji} r_{U,ji} [D_{ji}] [U_{ji}] \\ & + \sum_j \sum_i (\beta_{ji} + 4\gamma_{ji}) k_{b_3,ji} r_{T,ji} r_{U,ji} [T_{ji}] [U_{ji}] \dots \end{aligned} \quad (10)$$

where β is the number of calories consumed during the degradation of each soluble oligomer and γ is the number of calories consumed during the synthesis of each replacement protein. Equation 10 predicts that inbred organisms should be less metabolically efficient than outbred organisms, which has been confirmed by numerous studies (see Mitton 1997 for review). A simplified version of Equation 10 is obtained by assuming that only the dimerization reaction has a non-negligible rate, and by assigning every aggregating protein the same value for k_b and $[U]^2$, which are then combined into a single parameter, k_1 :

$$\frac{dC_{Maint}}{dt} = k_1 [\beta + 2\gamma] [T - 0.5N_{Het}] \quad (11)$$




Figure 2: Results from Equation 11 showing that metabolic efficiency increases with heterozygosity. The dashed curve shows the results for a higher k_1 value, which increases with both the stressfulness of the environment and the abundance of aggregation-prone proteins synthesized by the organism.

where T is the total number of gene loci that synthesize aggregation-prone proteins and N_{Het} is the number of heterozygous gene loci. As shown in Figure 2, Equation 11 is linear with respect to heterozygosity, which means that the overdominant loci exhibit modest epistasis for metabolic heterosis, otherwise the equation should decline exponentially with heterozygosity (Charlesworth and Willis 2009). A linear relationship between metabolic efficiency (as measured by oxygen consumption and growth rate) and heterozygosity has been found for several species of marine bivalves (Koehn and Shumway 1982; Garton et al. 1984; Mitton and Grant 1984; Hawkins et al. 1986).

Equations 1–10 provide a heuristic model for the relationship between protein aggregation and metabolic heterosis. The model shows that heterozygosity can be beneficial even in the absence of deleterious mutations, and is in agreement with previous research that found a correlation between heterozygosity and: (1) metabolic efficiency (Koehn and Shumway 1982; Garton et al. 1984; Mitton 1997), (2) protein turnover (Hawkins et al. 1986; Hawkins et al. 1989), and (3) expression of molecular chaperones (Kristensen et al. 2002; Pedersen et al. 2005). As shown in Figure 2, the model predicts that inbreeding depression should increase with the stressfulness of the environment because the values of k_b and $[U]_{steady}$ for each allele should increase with the stressfulness of the environment (see Subsection 3.3). This is also supported by previous research (see Armbruster and Reed 2005 for review).

2.3. Thermodynamic Considerations

It may be helpful to conceptualize the underlying physics of the model developed in this paper. The model

assumes steady-state conditions in which the rates of soluble oligomer addition, soluble oligomer removal, and synthesis of replacement proteins are equal (Equation 9). This results in steady-state concentrations of unfolded polypeptide chains, soluble oligomers, and native proteins. These steady-state concentrations are not in chemical equilibrium, otherwise the reactions for soluble oligomer formation would not proceed forward. For example, the fact that two unfolded polypeptide chains bind to each other to form dimers suggests that they have greater chemical potentials than the dimers (i.e. $2\mu_{unfolded} > \mu_{dimer}$). We can express this in terms of chemical affinity, A , which is the difference in chemical potential between reactants and products (Kondepudi 2008).

$$A \equiv 2\mu_{unfolded} - \mu_{dimer} > 0 \quad (12)$$

The greater the chemical affinity of a reaction, the further away it is from equilibrium.

However, the aggregation reactions proceed forward more slowly in hybrid organisms than in non-hybrids because the hybrids produce more allozymes. Point mutations can change the probability distributions for unfolded polypeptide chains assuming particular conformations without affecting the stability of the polypeptides' native conformations (King et al. 1996; Sinha and Nussinov 2001; Xu et al. 2013). This lowers the probability of two allozymes co-aggregating because polypeptide chains must have similar conformations in order to form the necessary cross- β interactions that allow them to bind to each other (O'Nuallain et al. 2004). Point mutations in polypeptide chains can also introduce incompatibilities (e.g. steric repulsion between side chains) that prevent two chains from co-aggregating (Apostol et al. 2010). Therefore, hybrid organisms benefit from lower rates of protein aggregation due to their greater degree of "mixedupness."

This can be thought of in terms of chemical potential and chemical affinity. The chemical potentials of the unfolded chains and dimers are related to their concentrations by:

$$\mu_{unfolded} = \mu_{unfolded}^{\circ} + kT \ln[U] \quad (13a)$$

$$\mu_{dimer} = \mu_{dimer}^{\circ} + kT \ln[D] \quad (13b)$$

where μ° is the standard-state chemical potential, kT is the temperature, $[U]$ is the concentration of unfolded polypeptide chain, and $[D]$ is the concentration of dimer. Since an organism's molecular chaperones

work to keep the concentration of dimer low, and the concentration of unfolded polypeptide chain should be about half as much in an organism that is heterozygous for the polypeptide than in an organism that is homozygous for the polypeptide, then the chemical affinity of the dimerization reaction should be lower in the heterozygous organism than in the homozygous organism. In other words, the homozygous organism is further away from chemical equilibrium.

Chemical affinity can be related to the rates of elementary step reactions by (Kondepudi 2008):

$$\frac{A}{kT} = \ln\left(\frac{R_f}{R_r}\right) \quad (14)$$


where R_f is the rate of the forward reaction and R_r is the rate of the reverse reaction. Finally, the relationship between the rate of progression and affinity for a chemical reaction is:

$$\frac{d\xi}{dt} = R_f(1 - e^{-\frac{A}{kT}}) \quad (15)$$


where the rate of chemical progression, $\frac{d\xi}{dt}$, is the difference between the forward and reverse reactions. Comparing Equations 12, 13, 14, and 15, it follows that the rate of progression for a polypeptide chain's dimerization reaction will be slower in an organism that is heterozygous for the polypeptide than in an organism that is homozygous for the polypeptide because the heterozygous organism is closer to chemical equilibrium.

Thus, organisms must expend energy to remove soluble oligomers because they maintain steady-state concentrations of native proteins, unfolded polypeptide chains, and soluble oligomers that are different from their equilibrium values. However, the aggregation reactions proceed toward chemical equilibrium more slowly in hybrid organisms than in non-hybrids, so the hybrids need a lower rate of calorie consumption to maintain steady-state conditions (Equation 10).

An analogy can be drawn with refrigerators, which consume energy to maintain steady-state thermal gradients. The amount of power a refrigerator consumes increases with the difference between its inside and outside temperatures because the rate of inward heat flow increases with the refrigerator's thermal gradient. Likewise, the constant movements toward and away from chemical equilibrium are responsible for organisms' maintenance costs, and the different rates of these movements give rise to the differences in performance (growth rate, size, etc.) between hybrid and non-hybrid organisms.



(a)



(b)

Figure 3: (a) Reproduction of data in Fig. 2 of Rumball et al. (1994). The curve shows the number of offspring produced by *D. melanogaster* after multiple generations of full-sib mating. (b) Equation 16 gives the relative fitness of an organism with increasing concentration of soluble oligomers.

3. Epistasis and Truncation Selection

3.1. Inbreeding and Epistasis

Subsection 2.2 discussed how heterozygosity affects an organism's metabolic performance when the organism maintains steady-state conditions. This subsection will consider the effect heterozygosity has on an organism's viability when the organism's defenses against protein aggregation are overwhelmed and steady-state conditions are no longer maintained. Some studies have found a steep drop in viability at high levels of inbreeding depression, especially in *D. melanogaster* (Kosuda 1972; Rumball et al. 1994; Figure 3). These results suggest that extinction of an inbreeding line is likely to occur when the level of inbreeding exceeds a threshold value (Frankham 1995). This might be evidence for synergistic epistasis between deleterious mutations (Kosuda 1972; Charlesworth 1998). However, it could also be evidence for epistasis between heterozygous gene loci, which Chelo and Teotónio (2012) have observed in their experiments.




Figure 4: Reproduction of plot in Fig. 1 of Tartaglia et al. (2007). The curve shows that the expression level of a protein is negatively correlated with its aggregation rate. This may mean that organisms produce as much protein as they can before it starts to aggregate.

Several authors have shown that molecular chaperones can provide a biochemical basis for epistasis by inhibiting protein aggregation (Fares et al. 2002; Sollars et al. 2003; Maisnier-Patin et al. 2005; de Visser and Elena 2007; de Visser et al. 2011; Lehner 2011). This subsection expands on these ideas and develops a theory that provides a biochemical basis for the steep decline in viability that sometimes occurs with increasing inbreeding. Then it discusses the model's implications for selection for heterozygosity.

One of the reasons that protein aggregation is such a nuisance is that organisms seem to produce as much of a protein as they can just before it starts to aggregate. Tartaglia et al. (2007) provides evidence for this in a plot similar to that shown in Figure 4, which shows that the logarithm of a protein's aggregation rate is negatively correlated with the logarithm of its expression level. From this plot, the authors of Tartaglia et al. (2007) conclude that organisms have “no scope for dealing with any situation in which these [expression] levels rise further or whereby the aggregation rates are increased... In the context of protein solubility, therefore, we are constantly living our lives at the edge of a molecular precipice.” In other words, we should expect steep declines in fitness when protein homeostasis is perturbed and soluble oligomers accumulate within an organism.

Since soluble oligomers are toxic substances (Kayed et al. 2003; Haass and Selkoe 2007; Vieira et al. 2007; Shankar et al. 2008), the equations used by toxicologists may be useful in modeling the fitness declines brought about by the accumulation of soluble oligomers. The responses of organisms to toxic substances typically follow log-normal distributions (Wagner and Løkke 1991; Wheeler et al. 2002; Newman and Unger

2003). Thus, the fitness of individuals with a given concentration of soluble oligomer can be obtained in manner similar to that given in Kimura and Crow (1978):

$$W([O]) = W_{max} \sqrt{\frac{1}{2\pi\sigma^2}} \int_{[O]}^{\infty} \frac{1}{C} e^{-\frac{(\ln C - \ln C_{50})^2}{2\sigma^2}} dC \quad (16)$$

where $W([O])$ is the fitness of individuals with total soluble oligomer concentration $[O]$, W_{max} is the fitness of individuals that produce no soluble oligomers, C is the concentration of soluble oligomer, C_{50} is the concentration of soluble oligomer that kills half of the individuals in a population, and σ is the shape parameter for the log-normal distribution.

I used a log-normal distribution for Equation 16 because log-normal distributions are commonly used in the toxicology literature, but any distribution that yields an S-shaped CCDF (e.g. normal, log-logistic, Weibull) can also be used in a truncation selection model. Use of a log-normal distribution would be justified if the toxicities of soluble oligomers are due to many random variables that result in multiplicative degradation (NIST/SEMATECH 2003), which would be true if, as hypothesized by others, their toxic effects are due to a general disruption of homeostasis caused by the permeabilization of cellular and organelle membranes (Kayed et al. 2003; Bucciantini et al. 2004; Kayed et al. 2004; Glabe 2006). A log-normal distribution would also be justified if the aggregation of regulatory and signaling proteins leads to a disruption in developmental homeostasis as postulated by I. Michael Lerner (see Section 4.2).

Equation 16 assumes that neither allele at a heterozygous gene locus negatively impacts the phenotype of its carrier. This may not be true and may be addressed with:

$$W([O]) = S([O]) \prod_i (1 - h_i s_i) \prod_i (1 - s_i) \quad (17)$$

where $S([O])$ is the right side of Equation 16, s is the fitness cost of the inferior alleles, and h is the dominance of the inferior alleles ($h=1$ is completely dominant and $h=0$ is completely recessive). $\Pi(1-hs)$ is the fitness cost of inferior alleles at heterozygous gene loci, and $\Pi(1-s)$ is the fitness cost of inferior alleles at homozygous gene loci.

The relative fitness of an individual, $(W([O])/W_{max})$, can be calculated once the total concentration of soluble oligomers in the individual is known (Figure 3). There are two approaches to calculate the concentration of

soluble oligomers for Equation 16. The first approach assumes steady-state conditions like in the previous section, but imposes an upper limit to either the amount of molecular chaperone an organism can produce or to the amount of energy available for the chaperone to perform its function². The assumption of an upper limit on available energy is reasonable because there are limits to the amount of oxygen organisms can acquire from their environment (Pörtner 2001; Bickler and Buck 2007; Ramirez et al. 2007). This assumption may apply to organisms in moderately severe environments or to severely inbred animals (e.g. Rumball et al. 1994). Under such conditions, the rate of soluble oligomer removal (Equation 6) may still equal the rate of soluble oligomer addition (Equation 5), but the effective concentration of molecular chaperone (the concentration of molecular chaperone activated by ATP) is constant, which results in a maximum rate of soluble oligomer removal, (R_{max}) . The result is:

$$[O]_{steady} = \frac{(K_m)(\frac{dA}{dt})}{R_{max} - \frac{dA}{dt}} \quad (18)$$

The steady-state concentration of soluble oligomer increases as dA/dt approaches the value of R_{max} . This will occur as the stressfulness of the environment increases or with decreasing heterozygosity.

The second approach assumes that steady-state conditions are disrupted, which would occur if $\frac{dA}{dt} > R_{max}$, or if an organism is unable to remove soluble oligomers once they have formed. There are several reasons why an organism would fail to maintain steady-state conditions:

1. The organism is dormant. Many organisms possess dormant resting stages, such as spores and cysts, that offer protection during adverse environmental conditions. These resting stages are usually associated with high concentrations of compatible solutes (e.g. trehalose) and small heat-shock proteins (sHSPs). Both compatible solutes and sHSPs can prevent unfolded polypeptide chains from binding to each other without consuming ATP. However, they cannot re-fold, disaggregate, or degrade misfolded proteins; they simply inhibit the formation of soluble oligomers (Singer and Lindquist 1998; Garay-Arroyo et al. 2000; Waters et al. 2008; van Leeuwen et al. 2013).

² Some chaperones, such as *Hsp70*, require ATP to bind to unfolded polypeptide substrates (Patterson and Höfled 2006; Lotz et al. 2010)

2. The organism does not have enough readily available energy to maintain protein homeostasis. All organisms have a finite energy supply available to them at any time. The ability of animals to generate ATP, for example, is limited by oxygen availability. Furthermore, several stresses are known to reduce an organism's ability to generate ATP. Thermal stresses can reduce the aerobic scope of ectothermic animals and desiccation can lead to a suspension of metabolism (Pörtner 2001; Alpert and Oliver 2002).
3. The organism is exposed to a sudden environmental shock. This can lead to elevated levels of protein aggregation. The organism should respond to the shock by synthesizing more molecular chaperones, but soluble oligomers can accumulate while the organism is adjusting to the new conditions.
4. The organism has an extracellular space. Multicellular organisms can secrete proteins into their extracellular space, which has orders of magnitude lower ATP than the intracellular space. Little is currently known about the process of preserving protein homeostasis in the extracellular space, but animals appear to produce extracellular chaperones that inhibit formation of soluble oligomers in a manner similar to sHSPs (Poon et al. 2002; Mannini et al. 2012; Wyatt et al. 2013).

In this case, the net rate of soluble oligomer accumulation is given by:

$$\left(\frac{dA}{dt}\right)_{Net} = \frac{dA}{dt} - \frac{dR}{dt} \quad (19)$$

Integrating Equation 19 will give the concentration of soluble oligomer at a given time, but that requires knowledge of how $[U]$ and k_b change with time for each polypeptide chain. The value of dR/dt will also change with time if the organisms respond to the environmental shock by increasing the concentrations of their molecular chaperones. Nevertheless, a $[O]$ value can be used in Equation 16 once it has been obtained. This would give the survivorship of individuals with different heterozygosities after exposure to an environmental shock for a given length of time. Equation 19 does not yield simple solutions like the steady-state equations, but exposures to suddenly elevated stresses may be a more frequent source of viability declines in natural environments.

Since Equation 5 predicts that the rate of soluble oligomer addition increases with decreasing heterozygosity, the above model predicts that viability


should decrease with [decreasing]³ heterozygosity under a given set of environmental conditions. In general, the steep decline in viability should occur at higher heterozygosities with increasing environmental harshness. This may explain why hybrid crops are more drought tolerant than non-hybrids (Duvick 2001), and why a threshold survivorship is seen in some inbreeding studies (Rumball et al. 1994; Frankham 1995). The model also has relevance for explaining truncation selection for heterozygosity.

3.2. Truncation Selection for Heterozygosity


Lewontin and Hubby (1966) provided an influential argument against the hypothesis that natural selection favors heterozygous genotypes. They measured the amount of allozyme diversity in wild *Drosophila pseudoobscura* populations and found polymorphisms segregating at approximately one-third (≈ 2000) of *D. pseudoobscura*'s gene loci. Lewontin and Hubby (1966) argued that such a large number of polymorphisms could not be maintained by balancing selection without enormous fitness costs. For example, if homozygosity at a single gene locus reduces the reproductive potential of an individual by 10%, and only two polymorphisms are segregating at the gene locus at a frequency of 50% each, then the reproductive potential of the whole population will be reduced by 5% for each gene locus. The reproductive fitness of the population would be 0.95^{2000} , or 10^{-46} , its maximum value. This is an unrealistically low number, and they concluded that natural selection could not favor heterozygotes.

Shortly after Lewontin and Hubby (1966) was published, three papers responded with a similar solution to the problem it raised (King 1967; Milkman 1967; Sved et al. 1967; Crow 1992). These papers proposed that truncation selection could maintain a large number of polymorphisms in natural populations without unreasonable fitness costs. In their models, all individuals whose heterozygosities are below a critical value have a fitness of zero, and individuals whose heterozygosities are above the critical value have maximum fitness (Figure 5a). Wills (1978) expanded on these models and showed that the number of polymorphisms that can be maintained in a population by truncation selection depends on the effective population size. However, he found that truncation of the least heterozygous individuals (the bottom


³the published manuscript contains a word omission here that escaped peer review. I have added the word "decreasing" to fix the error.



(a)



(b)



(c)

Figure 5: (a) Idealized truncation selection curve. The relative fitness of an organism is 1 above a critical heterozygosity and 0 below the critical heterozygosity. (b) Truncation selection for heterozygosity generated by combining Equations 16 and 18. The two curves show truncation selection for different levels of environmental stress as quantified by a k_1 parameter (see main text). (c) Equation 17 incorporates the possibility that one of the alleles at a gene locus negatively impacts the phenotype of its carrier.

5%) in a population can maintain polymorphisms at 66,000 gene loci in a population of 100,000 individuals, which is more than enough to support the number of polymorphisms that actually occur in natural populations. The effectiveness of truncation selection comes from its severity and its ability to operate on

many gene loci simultaneously. Several studies have found evidence for truncation selection acting on natural populations, but the results have been mixed (see Mitton 1997 for review and Kaeuffer et al. 2007 as a recent example).

Truncation selection can be considered an extreme form of epistasis in which heterozygosity confers decreasingly small fitness gains with each additional heterozygous gene locus. Both Rumball et al. (1994) and Chelo and Teotónio (2012) have provided experimental evidence that such epistasis exists. The epistasis can be modeled using a S-shaped truncation curve obtained by combining Equations 16 and 5 with either 18 or 19 (Figure 5b). The curve is not strictly a truncation selection curve, but Kimura and Crow (1978) has shown that selection is almost as efficient when it follows a S-shaped cumulative distribution curve (CDF or CCDF). In order to generate the figure, I assigned the same values of k_b and $[U]^2$ to all of the polypeptide chains in Equation 5 (I assumed the formation of trimers and tetramers was negligible). These can be considered the average values of k_b and $[U]^2$ for aggregation-prone proteins in a hypothetical organism. These values were combined into a single parameter $k_1 = k_b [U]^2$, which quantifies the propensity of proteins to aggregate. The value of k_1 should increase with the stressfulness of the environment and the susceptibility of an organism's proteins to aggregation (see below). Figure 5c shows the results when Equation 17 is used to generate a truncation curve. The same value of h_s is assigned to each suboptimal allele, which can be taken as an average of the h_s values for all suboptimal alleles.

The model depicted in Figures 5b and 5c may provide a physical basis for truncation selection for heterozygosity. According to the model, organisms are periodically exposed to stresses that lead to an accumulation of soluble oligomers, which in turn results in a sharp decline in viability. Only highly heterozygous individuals will be found in stressful environments because less heterozygous individuals will be on the wrong side of the truncation curve. This illustrates the severity of truncation selection, which is not typically associated with gradual evolution. If all the individuals in a species are on the wrong side of a truncation curve, then the species will go extinct. Thus, some individuals must be on the right side of the truncation curve prior to the species being subjected to truncation selection. In other words, some of the individuals in the species must be "pre-adapted" to the environment.

Another interesting property of the truncation selection model is that it depends only on the number of heterozygous gene loci, and not on the identity of the

gene loci. This is analogous to a “colligative property” in chemistry. This feature is important because it makes selection for heterozygosity compatible with sexual reproduction. Offspring are not going to be heterozygous at the same gene loci as their parents, but they will have, on average, the same number of heterozygous gene loci as their parents (assuming random mating in the population), so they should have the same overall fitness. Truncation selection allows organisms to substitute heterozygosity at one gene locus for heterozygosity at another gene locus without suffering significant fitness costs because the selection is acting on the overall level of heterozygosity, not on heterozygosity at any particular gene locus. The colligative nature of heterozygous advantage also applies to Equation 11, which relates heterozygosity to the metabolic efficiency of an organism. Again, overall metabolic efficiency depends on the number of heterozygous gene loci, not on the identity of the gene loci. Of course, there is the caveat that only some of the proteins produced by an organism are aggregation-prone, thus the identity of the gene loci is not completely unimportant. Rather, the colligative nature of heterozygous advantage is confined to the subset of gene loci that code for aggregation-prone proteins.

The truncation selection model has one important consequence. Factors that increase the accumulation of soluble oligomers in organisms should lead to truncation selection for higher heterozygosities. These factors can be broken up into two categories: 1) environmental stresses that promote protein aggregation, and 2) inherent characteristics of proteins that affect their propensity to aggregate. Section 4 will discuss how these factors interact to create selection for different ploidy levels, so it will be useful to describe how each promotes protein aggregation.

3.3. Environmental Stress

Temperature, water stress, and hypoxia can promote protein aggregation. High and low temperatures denature proteins, which leads to high concentrations of unfolded polypeptide chains (Becktel and Schellman 1987). The values of the rate law constants for protein aggregation also increase with temperature (Wang and Roberts 2013). Water stress (desiccation, freezing, salinity⁴) is created by the limitation of liquid water inside the organism, which results in high

concentrations of unfolded polypeptide chain (because the volume of solvent is limited) (Goyal et al. 2005). Water stress can also cause proteins to unfold by weakening the hydrophobic effect (Prestrelski et al. 1993; Allison et al. 1999). In addition, water stress results in molecular crowding, which increases the value of the rate law constants for protein aggregation (Ellis 2001; Smallwood and Bowles 2002; Chebotareva et al. 2004; Ellis and Minton 2006; White et al. 2010). Combining the effects of temperature and water stress is particularly harsh. For example, cold temperatures can cause proteins to denature, which greatly enhances the rate of protein aggregation when combined with freezing (Franks et al. 1990; Smallwood and Bowles 2002; Dias et al. 2009; Singh et al. 2009). Likewise, combining high temperatures with arid conditions should also promote aggregation. Finally, hypoxia can hinder the ability of organisms to remove soluble oligomers as they form because many molecular chaperones require ATP to bind to unfolded polypeptide chains (Patterson and Höhfeld 2006; Lotz et al. 2010). The physical stresses that promote protein aggregation should cause the truncation selection curves depicted in Figure 5 to shift to higher heterozygosity values, so only highly heterozygous individuals should be able to live in harsh environments.

3.4. Protein Length and IURs

The intrinsic characteristics of proteins can also affect their propensity to aggregate. Olzscha et al. (2011) determined what structural features increase a protein’s susceptibility to aggregation using molecular templates that seed protein aggregation in human cells. They found that protein size and the presence of intrinsically unstructured regions (IURs) were the two characteristics that most enhanced a protein’s susceptibility to aggregation. Both of these characteristics are associated with proteins that contain multiple domains (Dunker et al. 2005; Fong and Panchenko 2010).

A domain is a sequence of amino acids that, if separated from the rest of the polypeptide chain, would still fold into its proper conformation and function normally. A multi-domain protein is a protein that contains multiple domains. Some multi-domain proteins can be considered a string of proteins that are joined together. In fact, proteins that exist separately in some species may be found as parts of multi-domain proteins in other species, a phenomena called “domain accretion” (Koonin et al. 2002; Basu et al. 2009). Multi-domain proteins are important because their modularity makes them capable of adapting to rapidly

⁴Both high and low salinities can cause protein aggregation (e.g. Chang 2005; Downs et al. 2009; Tine et al. 2010; Monari et al. 2011)

changing environments (Sun and Deem 2007; He et al. 2009; Lorenz et al. 2011; Park et al. 2015). As expected, multi-domain proteins tend to be larger than single-domain proteins (Tan et al. 2005; Tordai et al. 2005).

Multi-domain proteins also tend to contain IURs. The reasons for this are less obvious and still debated (Dunker et al. 2005; Fong and Panchenko 2010). IURs can serve as inter-domain linkers that hold multiple domains together in a protein (Tomba 2002; Tomba 2005). IURs can also occur in or near the binding sites of proteins, and may facilitate protein interactions (Dunker et al. 2005; Uversky and Dunker 2010). This would make IURs particularly relevant to multi-domain proteins because such proteins tend to have multiple interaction partners (Tordai et al. 2005; Ekman et al. 2007). Indeed, the need to interact with multiple partners is one reason why some proteins contain multiple domains (Patthy 2003; Tordai et al. 2005; Tan et al. 2005). Each domain can facilitate interactions with different partners.

As stated previously, Olzscha et al. (2011) found that aggregation-prone proteins tend to be large and tend to contain IURs, which are characteristics of multi-domain proteins. Large proteins are more aggregation-prone than smaller proteins because they take longer to fold (or refold) into their native conformations, which increases the exposure of aggregation-prone amino acid sequences that are typically buried within the interior of the folded protein (Netzer and Hartl 1997; Goldschmidt et al. 2010; Olzscha et al. 2011). This would be true for both recently synthesized proteins and for proteins that denature due to an environmental stress. Large proteins also potentially contain more binding sites than smaller proteins, which would increase the likelihood for an oligomerization reaction to occur when large proteins approach each other (e.g. Andersen et al. 2010). Multi-domain proteins can also aggregate via “domain swapping.” This occurs when two domains in a multi-domain protein molecule are supposed to bind to each other, but instead they bind to the domains on another molecule (Nelsen and Eisenberg 2006; Rousseau et al. 2012).

Proteins that contain IURs tend to be aggregation-prone because the conformational flexibility of IURs allows them to align properly to form the cross- β sheet structure that holds aggregates together (Carrió et al. 2005; Nelson et al. 2005; Ventura 2005; Wang et al. 2008; Olzscha et al. 2011; Ramshini et al. 2011; Stroud et al. 2012). Zhang et al. (2013) have proposed that making a distinction between semi-disordered and completely disordered proteins

is useful, as only the former are aggregation prone. Hence, it is important to realize that aggregation-prone proteins contain intrinsically unstructured regions (IURs), not that the entire protein is intrinsically disordered. Nevertheless, several amyloid-associated diseases are due to polypeptides that contain IURs (Uversky et al. 2008; Uversky et al. 2009; Babu et al. 2011). In addition, proteins containing IURs are metastable and are often in an unfolded state. As a consequence, proteins containing IURs are thought to have relatively short life-spans and rapid turnovers because they are continuously degraded by each cell’s proteolytic machinery (Wright and Dyson 1999).

The abundance of large and IUR-containing proteins varies significantly across the domains of life. On the whole, prokaryotes produce shorter proteins than eukaryotes. About 65% of prokaryote proteins are multi-domain whereas 80% of eukaryote proteins are multi-domain (Apic et al. 2001). Furthermore, the median length of eukaryote proteins is 50% longer than the median length of prokaryote proteins (Brocchieri and Karlin 2005). The relative abundance of proteins containing IURs are 2%, 4%, and 33% for archaea, bacteria, and eukaryotes, respectively (Ward et al. 2004). These differences have caused prokaryotes and eukaryotes to process their proteins differently. For example, prokaryotes typically translate proteins at a rate of 10-20 amino acids per second whereas eukaryotes typically translate proteins at a rate of 3-8 amino acids per second (Siller et al. 2010). As a consequence, most protein folding is post-translational in prokaryotes but co-translational in eukaryotes (Netzer and Hartl 1997). Eukaryotes also have more complex chaperone systems that assist with the proper folding of their *de novo* and stress-denatured proteins (Albanèse et al. 2006). Thus, protein folding is more elaborate in eukaryotes than prokaryotes because their proteins are more aggregation prone. The truncation selection curves depicted in Figure 5 should be shifted to higher heterozygosity values for organisms that synthesize more aggregation-prone proteins, and indeed, heterozygosity is essentially a eukaryotic phenomena. There are no heterozygous bacteria and archaea. This argument will be expanded later to explain the haploid-diploid transition.

A species’ geographic distribution is affected by both environmental stress and its proteins’ susceptibility to aggregation. Both Koonin et al. (2002) and Brocchieri and Karlin (2005) proposed that hyperthermophiles are common in the Domain Archaea because the species in this domain produce short proteins that are not prone to aggregation. However, the idea can

be extended to all prokaryotes that live in extreme environments. Thermophiles can be found among both the Bacteria and the Archaea. Additionally, both domains of life contain species that are able to grow at extremely high salinities (Kunte et al. 2002), withstand desiccation (Potts 1994; Alpert 2006), or grow in freezing conditions (Russell 1998). Not all prokaryotes can grow in extreme environments because additional adaptations, such as those promoting membrane integrity and DNA stability, are required (Konings et al. 2002; Trivedi et al. 2005). However, prokaryotes should be “pre-adapted” to these extreme environments because they are not removed by truncation selection when they migrate into them. In contrast, truncation selection could prevent organisms that synthesize more aggregation-prone proteins, including many eukaryotes, from migrating into these environment (because their relative fitness would be zero), thereby limiting their distribution to less harsh environments.

4. Ploidy Level

4.1. Environmental Stress and Polyploidy

If the expression of two allozymes can inhibit the formation of soluble oligomers, then the expression of three or more allozymes should further inhibit their formation. Equation 4 predicts that polyploid organisms should have lower rates of protein aggregation than diploid organisms. In addition, many polyploids are hybrid species (allopolyploids), and are heterozygous at more gene loci than diploids (Otto and Whitton 2000). However, even autopolyploids can express more than two allozymes at a given gene locus. Therefore, polyploid organisms should be able to tolerate more severe physical stresses than diploids, and in fact, may be “pre-adapted” to harsh environments in which diploid organisms cannot survive due to truncation selection. Indeed, since polyploidy and hybridization may be accompanied by fitness costs (such as high genetic loads, slower growth rates, and outbreeding depression), polyploid species may be restricted to harsh environments where they do not have to compete with diploid species (Otto and Whitton 2000).

The occurrence of polyploid plants at high latitudes and altitudes was first observed in the 1940’s (Stebbins 1950; Stebbins 1984), and recent research has confirmed that polyploid plants and animals frequently occur in frozen environments (Beaton and Hebert 1988; Adamowicz et al. 2002; Brochmann et al. 2004; Lundmark and Saura 2006; Aguilera et al. 2007; Otto et al. 2007; Adolffson et al. 2009). Brochmann et al.

(2004) analyzed data from the *Pan-Arctic Flora (PAF) Checklist* (Elven et al. 2003) and found that 73.7% of arctic plants are polyploid. Plants in the most northerly arctic zone were hexaploid on average. In addition, 39.2% of species within this zone were 7-ploid or higher, and 17.8% of the species were 9-ploid or higher. However, since these plants reproduce primarily through self-fertilization, the heterozygosity of these plants is about half what their ploidy level would indicate.

Polyploid plants are also positively associated with arid zones and deserts (Spellenberg 1981; Rossi et al. 1999; Hunter et al. 2001; Pannell et al. 2004; Joly et al. 2006; Schuettpelz et al. 2008). Senock et al. (1991) and Hao et al. (2013) showed that the ploidy level of *Atriplex canescens* increases in regions of the Chihuahuan Desert with increasing drought stress. Most resurrection plants, which grow in deserts and are capable of withstanding very high levels of desiccation, are polyploid (Bartels and Salamini 2001; Rodriguez et al. 2010). Several studies have shown that a plant’s drought tolerance increases with its ploidy level (Al Hakimi et al. 1998; Xiong et al. 2006). For example, Ramsey (2011) compared the drought tolerances of hexaploid and tetraploid individuals belonging to *Achillea borealis* and found the hexaploids were more tolerant. Furthermore, Ramsey compared the drought tolerances of newly formed hexaploid *A. borealis* individuals to hexaploid individuals collected from the wild, and determined that a third of the the drought tolerance was achieved via genome duplication alone rather than adaptation. Another interesting example of polyploid adaptation to water stress may be the redwood, *Sequoia sempervirens*, which is hexaploid and must cope with water stress due to its extreme height (Ahuja and Neale 2002; Koch et al. 2004; Oldham et al. 2010). Polyploid animals are also positively associated with arid zones. For example, polyploid lizards are found in deserts (Tocidowski et al. 2001; Kearney 2003) and so is the only known polyploid mammal (Gallardo et al. 1999; Gallardo et al. 2004; Svartman et al. 2005; Gallardo et al. 2006; Teta et al. 2014).

Polyploidy should also be associated with salinity stress because many species change their expression of *Hsps* in both hypersaline and hyposaline environments (Chang 2005; Downs et al. 2009; Tine et al. 2010; Monari et al. 2011). However, evidence for a link between polyploidy and salinity is unfortunately tenuous. Polyploid plants appear to have greater tolerance to salt stress than their diploid relatives (Tal and Gardi 1976; Shannon and Grieve 1999; Ashraf et al. 2001; Kumar et al. 2009). Also, several species of

polyploid brine shrimp, *Artemia*, have been identified (Browne and Bowen 1991; Amat et al. 2007). Several papers have suggested that the radiation of polyploid *Artemia* is related to the Messinian salinity crisis (e.g. Agh et al. 2007), but other papers have disputed this claim (Baxevanis et al. 2006). Various papers have found that parthenogenic *Artemia* tolerate both higher and lower salinities than sexuals, but none of these papers distinguish between polyploid and diploid parthenogens (Browne and MacDonald 1982; Zhang and King 1993; El-Bermawi et al. 2004; Agh et al. 2007). The distribution of polyploid *Artemia* may instead be driven by latitude (Zhang and Lefcort 1991).

The geographical distribution of polyploid organisms suggests that they have an advantage in environments that promote protein aggregation. These trends can be understood in terms of a truncation selection model. Truncation selection would prevent individuals with low heterozygosities from migrating into harsh environments, thereby preventing some diploid species from expanding into harsh environments and adapting to them. In contrast, polyploid individuals may be partially pre-adapted to harsh environments because their heterozygosities are already sufficiently high to avoid truncation selection. Upon time, the polyploids will further adapt as favorable alleles increase in frequency. A dramatic illustration of this may be given in Fawcett et al. (2009), which argues that polyploids may have preferentially survived the Cretaceous-Paleogene mass extinction event.

4.2. Organism Complexity, Protein Interaction Networks, and the Haploid/Diploid Transition

Protein aggregation may also potentially explain why some species are haploid and others are diploid. The most suggestive evidence for this is the relative stress tolerances of haploid and diploid organisms. Animals, for instance, have lower thermotolerances than bacteria and archaea, and some fungi species (Pörtner 2001; Salar and Aneja 2007). Even the Pompeii worm, which grows in hydrothermal vents, can not withstand temperatures greater than 55°C for 2 hours (Ravaux et al. 2013). Similarly, animals are less tolerant to desiccation stresses than prokaryotes and fungi, with the notable exceptions of bdelloid rotifers and tardigrades⁵ (Alpert 2006). The general trend appears to be that animals are less stress tolerant than fungi, which in

⁵Polyploidy may explain these exceptions. Bdelloid rotifers are descended from a tetraploid ancestor, and most freshwater and terrestrial tardigrades are polyploid (Bertolani 2001; Hur et al. 2009)

turn are less stress tolerant than bacteria and archaea. Considering that the ability to produce *Hsps* appears to control the upper thermal tolerances of organisms (Pörtner 2001; G. F. Dilly et al. 2012), the relative stress tolerances of animals, fungi, bacteria, and archaea may reflect the relative levels of protein aggregation that they must cope with.

The trend in stress tolerance seems to match a trend in the ploidy levels of these organisms. Bacterial and archaeal species are haploid, animal species are diploid, and fungal species may either be haploid or diploid. Thus, bacterial and archaeal species may never have to cope with truncation selection for heterozygosity while all animal species may experience such truncation selection, even in relatively mild conditions. Fungal species may or may not experience truncation selection for heterozygosity depending on the environmental conditions or the stage in their life-cycle (see below). An underlying mechanism may generate the different levels of protein aggregation in these organisms, which may explain their relative stress tolerances and their ploidy levels.

But why do different species experience different levels of protein aggregation? One explanation is that some species produce more aggregation-prone proteins than others. Olzscha et al. (2011) found that aggregation-prone proteins in human cells tend to be large and tend to contain IURs, which are characteristics of multi-domain proteins (Dunker et al. 2005; Fong and Panchenko 2010). These proteins often have numerous interaction partners and are often involved in signal transduction and regulatory processes (Dunker et al. 2005; Warringer and Blomberg 2006; Uversky and Dunker 2010; Olzscha et al. 2011). Such proteins facilitate organism complexity because they have important roles in coordinating and regulating the biochemical activities within and between cells, which is necessary for the development of complex organisms (Rubin et al. 2000; Patthy 2003; Dunker et al. 2005; Tan et al. 2005; Tordai et al. 2005; Ekman et al. 2007; Uversky and Dunker 2010). For example, Stumpf et al. (2008) found that the human protein interaction network (PIN) contains approximately 650,000 protein interactions while the *S. cerevisiae* PIN contains approximately 25,000-35,000 protein interactions.

PINs are important for communication and coordinating all of the activities that occur in cells (Barabási and Oltvai 2004; Vinayagam et al. 2014). Communication usually occurs through signaling cascades involving large numbers of proteins (Pawson and Nash 2000; Breitkreutz et al. 2010; Vinayagam et al. 2011). Coordination can involve the switching

on or off of proteins via phosphorylation, transcription factors, proteolysis, etc. (LeMosy et al. 2001; Walhout 2006; López-Otín and Hunter 2010; Wang and Chen 2010; Cheng et al. 2011). Thus, the cell needs a sort of managerial class of proteins whose specialty is interacting with other proteins. These proteins tend to be larger than the median protein because they need multiple domains to facilitate all of their interactions (Tan et al. 2005; Tordai et al. 2005; Ekman et al. 2007). The managerial proteins also tend to be metastable (or semi-disordered) which allows them to fold into multiple conformations that can bind to different partners. This flexibility is facilitated by IURs (Dunker et al. 2008). The managerial proteins appear to be prone to aggregation, and the molecular chaperone Hsp90 appears to specialize in preventing such aggregation of this class of proteins (Picard 2002; Sangster et al. 2004; Vabulas et al. 2010).

The proteins that facilitate complex PINs tend to be large and multi-domain because multiple domains are necessary to facilitate different interactions (Rubin et al. 2000; Tan et al. 2005; Tordai et al. 2005; Ekman et al. 2007; Zmasek and Godzik 2011). For example, Xia et al. (2008) found that the average number of interaction domains per protein increases with the number of cell types in an organism. Likewise, Wang et al. (2005) found that proteins shared by *S. cerevisiae*, *D. melanogaster*, and *H. sapiens* are similar in length; but proteins found in *D. melanogaster* and *H. sapiens*, but not in *S. cerevisiae*, are on average 22% longer than proteins shared by all three species. Finally, Warringer and Blomberg (2006) found that *S. cerevisiae* proteins longer than 770 amino acids have more interaction partners, on average, than shorter proteins. Particularly abundant among these large proteins were transport proteins, proteases, kinases, and other signaling proteins. These proteins are responsible for signal transduction and regulating biochemical pathways (Sopory and Munshi 1998; Manning et al. 2002; López-Otín and Hunter 2010). In multicellular organisms, these proteins play vital roles in intercellular communication, regulation of the cell cycle, and cellular differentiation (LeMosy et al. 2001; Schaller 2004; Turk 2006; van der Hoorn 2008; Keshet and Seger 2010). Thus, large proteins play an important role in coordinating and regulating the biochemical activities in organisms, and such proteins appear to be more abundant in complex organisms (Tordai et al. 2005; Ekman et al. 2007).

Proteins containing IURs also facilitate complex PINs. IURs can serve as flexible inter-domain linkers that allow domains to move freely with respect to each

other (Tompa 2002; Tompa 2005). These inter-domain linker regions are not merely structural, but often times serve as binding sites between proteins and their interaction partners (Balázs et al. 2009). IURs allow proteins to assume multiple conformations which in turn allows the proteins to bind to multiple partners (or allows multiple partners to bind to the same protein) (Kriwacki et al. 1996; Tompa et al. 2005; Oldfield et al. 2008; Tyagi et al. 2009; Bustos 2012). IURs may also speed up binding reactions via a “fly-casting” mechanism (Shoemaker et al. 2000). Finally, IURs provide easily accessible sites for post-translational modifications to proteins, which makes them important for biochemical regulation (Dunker et al. 2002; Kurotani et al. 2014). As a consequence of these structural properties, up to 94% of transcription factors and more than 70% of signaling proteins in eukaryotes contain IURs (Iakoucheva et al. 2002; Liu et al. 2006; Uversky and Dunker 2010). In short, IURs are important mediators of protein interactions, and proteins containing IURs are responsible for most signal transduction and biochemical regulation in eukaryotes.

Thus, it appears that the proteins that facilitate organism complexity are also the proteins that make complex organisms more sensitive to environmental stresses. This can explain why higher ploidy levels are associated with complex organisms. Bacteria and archaea never experience truncation selection for heterozygosity, even in harsh environments. In contrast, animals must be diploid, even in mild environments, because they produce an abundance of proteins with numerous interaction partners. These proteins are typically large and typically contain IURs, so they tend to be aggregation-prone. As a consequence, animals must be heterozygous in order to be on the right side of the truncation curve (Figure 5), and that requires them to be at least diploid. Organisms of intermediate complexity, such as plants and fungi, may be either haploid or diploid, depending on the species. Whether such species are haploid or diploid will depend on the abundance of aggregation-prone proteins that they produce and on the stresses to which their proteins are exposed. If this hypothesis is true, then the diploidy of complex organisms is largely necessitated by the physical constraints imposed by PIN complexity. It may be the case that the thermodynamic benefits of heterozygosity (Subsection 2.3) increase with an organism’s complexity.

4.3. Developmental Homeostasis

The hypothesis presented in the previous subsection shares several connections with the pioneering work

from early metabolic heterosis theorists, such as I. Michael Lerner. Lerner (1954) found that inbred plants and animals exhibited more morphological variability than outbreds, which Lerner attributed to a decline in developmental homeostasis (or developmental stability). In other words, less heterozygous individuals display greater degrees of aberrant growth and development, which result in morphological imperfections such as bilateral asymmetry. Over the years, other researchers have corroborated Lerner's findings (Robertson and Reeve 1952; Eanes 1978; Mitton 1978; Soulé 1979; Mitton 1995), which in turn lead to the studies that found heterozygosity correlates with growth rate, metabolic efficiency, and protein turnover (Singh and Zouros 1978; Zouros et al. 1980; Koehn and Shumway 1982; Garton et al. 1984; Mitton and Grant 1984; Hawkins et al. 1986; Hawkins et al. 1989). Thus, the field of metabolic heterosis studies can trace its roots back to Lerner's work on developmental homeostasis.

The hypothesis presented in this subsection can explain how low heterozygosity would disrupt developmental homeostasis. The proteins that regulate proper growth and development are also the proteins that are most susceptible to aggregation. Both environmental stress and inbreeding can lead to elevated rates of aggregation for signaling and regulatory proteins, which could potentially disrupt intercellular communication, regulation of the cell cycle, proper cell differentiation, etc. The cumulative result would be a disruption in developmental homeostasis as described by I. Michael Lerner. Thus, protein aggregation can potentially explain much of the phenomena that concerned early researchers working with allozymes (see Mitton 1997 for review.)

4.4. *Plants and Alternation of Generations*

The trends described in the previous subsections hold for plant species as well. Plants typically alternate between a haploid gametophyte generation and a diploid sporophyte generation. However, the dominant generation varies between divisions. For example, bryophytes have a dominant gametophyte generation and a short-lived sporophyte generation while spermatophytes are primarily diploid (the sporophyte generation is dominant). In addition, ferns have independent gametophyte and sporophyte generations. The theory presented in Subsection 4.2 predicts that fewer aggregation-prone proteins should be produced by the haploid gametophytes than in the diploid sporophytes. Two types of circumstantial evidence support this prediction.

First, there is a relationship between complexity and ploidy level. The haploid-dominant bryophytes are relatively simple plants, typically 2 cm tall and one cell thick. The diploid-dominant spermatophytes are complex and include flowering plants. The ferns alternate between a haploid generation that is simple, resembling bryophytes, and a diploid generation that is significantly larger and more complex. Thus, according to the hypothesis presented in the previous subsection, spermatophytes and fern sporophytes should have more complex PINs and should synthesize more aggregation-prone proteins than bryophytes and fern gametophytes.

The second type of evidence is the relative stress tolerances of the different plant divisions. The relationship between ploidy level and stress tolerance in plants is similar to the trend described in Subsection 4.2 for animals, fungi, and prokaryotes. Bryophytes are much more tolerant of freezing, desiccation, and salinity stresses than spermatophytes (Alpert 2000; Oliver et al. 2005; Wang et al. 2009; Gaff and Oliver 2013). Their ability to tolerate such stresses is comparable to lichens, and they can be found, along with lichens, in extremely cold and arid environments not inhabited by more complex plants (Longton 1988; Alpert 2006; Proctor and Tuba 2002; Kranner et al. 2008). The desiccation tolerance of ferns is more complex. Fern sporophytes are comparable to spermatophytes in their ability to tolerate desiccation, but fern gametophytes are comparable to bryophytes (Watkins, Jr. et al. 2007; Hietz 2010). In fact, some tropical fern species have lost the sporophyte stage of their life cycle and now exist as asexually reproducing gametophytes, which has allowed them to migrate into colder and drier habitats than their sporophyte-producing relatives (Farrar 1978; Farrar 1990). Thus, bryophytes and fern gametophytes probably express fewer aggregation-prone proteins than spermatophytes and fern sporophytes. This would explain the relative order of plant stress tolerances: bryophyte \approx fern gametophyte $>$ spermatophyte \approx fern sporophyte. The trend corresponds to the relative complexity of the plant divisions and to their ploidy levels.

The above observations on relative stress tolerances should not be taken to imply that bryophytes are never subjected to truncation selection for heterozygosity. For example, allodiploid species of bryophytes (hybrids with two sets of chromosomes) have been identified, and they increase in frequency with latitude (Wyatt et al. 1988; Ricca et al. 2008). Bryophyte species probably produce few enough aggregation-prone proteins that they do not experience truncation

selection for heterozygosity in mild environments, but they may produce enough aggregation-prone proteins that they experience truncation selection in harsher environments. Thus, bryophytes, ferns, and spermatophytes all exhibit ploidy level increases in harsh environments. The difference between bryophytes and spermatophytes is that bryophytes have a haploid chromosome set baseline whereas spermatophytes have a diploid chromosome set baseline. Bryophytes may also experience truncation selection for heterozygosity in their sporophyte generations. Like ferns, the sporophyte generations of bryophytes are diploid and cannot survive in as harsh of environments as the gametophyte generations (Stark et al. 2007). This may indicate that the sporophyte generation of bryophytes express aggregation-prone proteins that are not produced by the gametophyte generation.

4.5. *Alternation of Generations and the Evolution of Complexity*

When comparing the life-cycles of green algae, bryophytes, ferns, and spermatophytes, there appears to be a progression from haploid dominant species, to species that alternate between haploid and diploid generations, to diploid dominant species. In other words, diploid dominant species did not evolve directly from haploid dominant species, but instead, from species that alternated between haploid and diploid generations. The truncation selection model developed in this paper can explain this evolutionary sequence.

Recall that a species will go extinct if all individuals are on the wrong side of a truncation curve, and that diploid species produce aggregation-prone proteins that contain multiple domains and IURs. These two statements imply that it would be impossible for strictly haploid species to produce an abundance of aggregation-prone proteins because truncation selection would guarantee their extinction. Hence, the need for diploidy would never arise. Diploid individuals are unlikely to successfully compete against haploid individuals if there is no immediate advantage to diploidy, especially if there are advantages to haploidy as described in the next subsection.

However, a spore-producing generation can circumvent this barrier by providing an immediate advantage to diploidy. Sexual reproduction requires haploid dominant species to possess at least a temporary diploid generation. For such organisms, sexual reproduction is often accompanied by the production of spores or cysts. This may facilitate inbreeding avoidance since spores often serve as a means of dispersal, or a mixed cytoplasm may increase

the ability of a spore or cyst to tolerate environmental stresses (Equation 16). A mixed cytoplasm may also increase the longevity of a spore or cyst since it would slow down the accumulation of soluble oligomers in the dormant organism over time. For fungi, there is some evidence that sexual spores are more stress tolerant than asexual spores⁶ (Grishkan et al. 2003; Dijksterhuis 2007; Trapero-Casas and Kaiser 2007). Regardless, sexual reproduction requires organisms to possess at least a diploid zygote in their life-cycle, and this often accompanies spore production. Some species have evolved a separate diploid (or dikaryotic) spore-producing generation, such as sporophytes in plants or ascocarps and basidiocarps in dikaryotic fungi, because it aids with spore dispersal. This spore-producing generation has the potential to evolve complexity over time because truncation selection would not prevent it from expressing genes that encode the aggregation-prone proteins that facilitate complexity. In contrast, truncation selection would prevent strictly haploid spore-producing species from becoming more complex over time.

Given the constraints that truncation selection would impose on haploid organisms, the evolution of diploid-dominant organisms may have proceeded along the following steps: 1) Bacterial and archaeal species produce proteins that are not susceptible to aggregation, so they do not have a heterozygous diploid stage in their life-cycle, even when producing spores; 2) some eukaryotic species (e.g. some algal and fungal species) produce enough aggregation-prone proteins that their spores benefit from a mixed cytoplasm, which requires at least a temporary diploid stage in their life-cycle⁷; 3) Some species have longer-lived diploid, spore-producing generations in their life-cycles, perhaps because they facilitate spore dispersal (e.g. bryophytes) 4) the diploid stages of some species' life-cycles have become more complex over time because truncation selection has not prevented them from producing aggregation-prone proteins with numerous interaction partners (e.g. ferns, dikaryotic fungi); and 5) the haploid life-cycle stage has become temporary in some multicellular species, which has resulted in diploid-dominant species (e.g. animals and spermatophytes). Thus, the evolution

⁶The spores in these studies were haploid, but they could have inherited a mixed cytoplasm from their diploid parent cells during meiosis.

⁷This could have evolved when cells from closely related species fused together to form a diploid cell, which would be analogous to the formation of polyploid species via hybridization

of diploid-dominant species would be a very gradual process, unlike the evolution of polyploidy, which occurs in a single generation.

The freshwater green algae Charales may provide support for the above evolutionary sequence. Charales are exclusively haploid, lacking a sporophyte stage in their life-cycle (Becker and Marin 2009). They are also relatively complex compared to other green algae, but are still much simpler than angiosperms (Lee 2008). However, Graham and Gray (2001) argues that Charales “are not competitive with freshwater angiosperms and rarely share freshwater habitats with them.” In fact, the fossil record shows that Charales’s species diversity has declined over time since the appearance of angiosperms, and that Charales may be an evolutionary dead end (Graham and Gray 2001). Given that Charales species are closely related to embryophytes, and that complexity seems to have given angiosperms a competitive advantage over Charales species, it is natural to ask why Charales species did not become more complex over time. Competition between Charales individuals should have lead to an increase in complexity over time, just as competition favored the evolution of complex angiosperms.

The absence of a sporophyte stage in the Charales life-cycle may explain their relatively simple morphology since, as argued in the previous four paragraphs, truncation selection would impose an upper limit on the abundance of aggregation-prone proteins that the haploid Charales species can produce. This in turn would impose an upper limit on the complexity of Charales species because proteins with numerous interaction partners, which are required in complex organisms, tend to be aggregation-prone. Thus, Charales species may have hit an evolutionary dead end because they do not possess a sporophyte generation in their life-cycle. In contrast, an upward-growing sporophyte generation is beneficial to embryophytes because it facilitates spore dispersal through the air. The possession of a diploid sporophyte generation may have removed a barrier to the evolution of complexity in some embryophyte lineages, which has allowed the spermatophytes to become large, complex organisms.

4.6. *The Advantages of Haploidy*

The theory developed in Subsection 4.2 attempts to explain why diploidy is advantageous. However, the occurrence of organisms that alternate between haploid and diploid life-cycles stages (e.g. ferns and *Ulva*) suggests that haploidy has advantages. Such organisms could stay permanently diploid (with brief haploid stages for the purpose of sexual reproduction) if

there was no advantage to haploidy. Two such potential advantages are genetic load and growth rate.

Genetic load should favor the evolution of lower ploidy levels (Mable and Otto 1998). The genetic load in a population is directly proportional to the mutation rate (Haldane 1937). Thus, if mutation rates are relatively constant at all ploidy levels, then a population of diploid individuals should have twice the genetic load of a population of haploid individuals (Otto and Whitton 2000, Gerstein and Otto 2009). As a consequence, populations of haploid individuals should have higher mean fitnesses than populations of otherwise identical diploid individuals (Mable and Otto 1998).

Higher ploidy levels are also disadvantageous because they lead to slower growth rates. This has been observed in polyploid plants, which typically grow and mature more slowly than their diploid relatives (Otto and Whitton 2000; Hesse et al. 2009). Also, diploid gametophyte lines of one bryophyte species grow $\approx 70\%$ as fast as haploid lines on full medium (Schween et al. 2005). Thus, haploidy might be beneficial to organisms that face strong selection for high growth rates. Haploidy would also be particularly beneficial to single-celled organisms, in which cell division rates are directly tied to fecundity.

The growth rate hypothesis is particularly useful when trying to understand the life cycles of plants. All plants must compete for limited space (Gurevitch et al. 1990; Gremer et al. 2013), so a faster growth rate would give haploid plants an advantage over diploid plants when attempting to compete for access to land. For instance, bryophytes reproduce asexually via fragmentation and sexually via spore production. In both cases, the gametophyte plants must quickly grow from only a few cells in order to establish themselves in a partition of land. Ferns also disperse themselves via spores made up of only a few cells. Their simple haploid gametophyte generations may also be beneficial for competition over access to land. Once established, the ferns reproduce sexually and produce their complex, diploid sporophyte generations, which don’t have to compete for access to land because they grow out of their gametophyte parents. In contrast, spermatophytes disperse themselves via seeds that carry entire diploid plant embryos. The plant embryos can quickly establish themselves in a partition of land, despite a slower growth rate, because they are already partially developed prior to germination. This might allow spermatophytes to be diploid, complex organisms for the bulk of their life-cycle, which in turn might allow them to utilize complex structures (e.g. flowers) for all of their life processes, including sexual reproduction.

5. Conclusion

This paper attempts to provide a biochemical basis for heterosis and selection for heterozygosity. It then shows how individuals with higher heterozygosities are favored with increasing environmental harshness and organism complexity, which will also favor higher ploidy levels. In addition, the hypothesis was used to explain the different life-cycles of plants and the results of numerous experiments that have found heterozygosity correlates with reduced protein turnovers and higher metabolic efficiencies. Thus, the hypothesis can explain numerous field observations and experimental data in which heterozygosity and ploidy level are variables. Future research should be able to establish whether heterozygous advantage has a thermodynamic basis, and whether organism complexity and environmental harshness are in fact determinants of each species ploidy level.

The hypotheses developed in this paper have important implications for breeding more stress tolerant varieties of crops. Several algorithms have been developed for identifying aggregation-prone amino acid sequences in proteins (Tartaglia et al. 2008; Goldschmidt et al. 2010). These algorithms can be used to identify the proteins that are susceptible to aggregation, which in turn may identify gene loci where heterozygosity is most beneficial. Such identification may prove helpful in developing crop varieties that can withstand the physical stresses caused by global climate change (Lobell et al. 2013). There is already some evidence that global climate change is leading to selection for heterozygosity in animal populations (Forcada and Hoffman 2014). Identifying aggregation-prone proteins may also be helpful in further improving crop yields as has occurred throughout the 20th century.

Acknowledgments

I would like to thank Jeffrey Mitton and Anthony J. Hawkins for their recommendations to improve this paper and encouragement. I would especially like to thank Dr. Mitton for his thorough review of the manuscript.

References

Adamowicz, S. J., Gregory, M. C., Marinone, M. C., and Herbert, P. D. N. (2002). New insights into the distribution of polyploid *Daphnia*: the Holarctic revisited and Argentina explored. *Molecular Ecology*, 11:1209–1217.

- Adolfsson, S., Michalakis, Y., Paczesniak, D., Bode, S. N. S., Butlin, R. K., Lamatsch, D. K., Martins, M. J. F., Schmit, O., Vandekerckhove, J., and Jokela, J. (2009). Evaluation of elevated ploidy and asexual reproduction as alternative explanations for geographic parthenogenesis in *Eucypris virens* ostracods. *Evolution*, 64:986–997.
- Agh, N., Abatzopoulos, T. J., Kappas, I., van Stappen, G., Rouhani, S. M. R., and Sorgeloos, P. (2007). Coexistence of sexual and parthenogenetic *Artemia* populations in Lake Urmia and neighbouring lagoons. *International Review of Hydrobiology*, 92:48–60.
- Aguilera, X., Mergeay, J., Wollebrants, A., Declercq, S., and Meester, L. D. (2007). Asexuality and polyploidy in *Daphnia* from the tropical Andes. *Limnology and Oceanography*, 52:2079–2088.
- Ahuja, M. R. and Neale, D. B. (2002). Origins of polyploidy in coast redwood (*Sequoia sempervirens* (D. DON) ENDL.) and relationship of coast redwood to other genera of Taxodiaceae. *Silvae Genetica*, 51:93–100.
- Al Hakimi, A., Monneveux, P., and Nachit, M. M. (1998). Direct and indirect selection for drought tolerance in alien tetraploid wheat x durum wheat crosses. *Euphytica*, 100:287–294.
- Albanèse, V., Yam, A. Y.-W., Baughman, J., Parnot, C., and Frydman, J. (2006). Systems analyses reveal two chaperone networks with distinct functions in eukaryotic cells. *Cell*, 124:75–88.
- Albertin, W. and Marullo, P. (2012). Polyploidy in fungi: evolution after whole-genome duplication. *Proceedings of the Royal Society of London B*, 279:2497–2509.
- Allison, S. D., Chang, B., Randolph, T. W., and Carpenter, J. F. (1999). Hydrogen bonding between sugar and protein is responsible for inhibition of dehydration-induced protein unfolding. *Archives of Biochemistry and Biophysics*, 365:289–298.
- Alpert, P. (2000). The discovery, scope, and puzzle of desiccation tolerance in plants. *Plant Ecology*, 151:5–17.
- Alpert, P. (2006). Constraints of tolerance: why are desiccation-tolerant organisms so small or rare? *Journal of Experimental Biology*, 209:1575–1584.
- Alpert, P. and Oliver, M. J. (2002). Drying without dying. In Black, M. and Pritchard, H. W., editors, *Desiccation and Survival in Plants: Drying Without Dying*, pages 3–44. CAB International, Wallingford, UK.
- Alström-Rapaport, C., Leskinen, E., and Pamilo, P. (2010). Seasonal variation in the mode of reproduction of *Ulva intestinalis* in a brackish water environment. *Aquatic Botany*, 93:244–249.
- Amat, F., Hontoria, F., Navarro, J. C., Vieira, N., and Mura, G. (2007). Biodiversity loss in the genus *Artemia* in the Western Mediterranean Region. *Limnetica*, 26:387–404.
- Andersen, C. B., Manno, M., Rischel, C., Thórólfsson, M., and Martorana, V. (2010). Aggregation of a multidomain protein: a coagulation mechanism governs aggregation of a model IgG1 antibody under weak thermal stress. *Protein Science*, 19:279–290.
- Apic, G., Gough, J., and Teichmann, S. A. (2001). Domain combinations in Archaeal, Eubacterial and Eukaryotic proteomes. *Journal of Molecular Biology*, 310:311–325.
- Apostol, M. I., Sawaya, M. R., Cascio, D., and Eisenberg, D. (2010). Crystallographic studies of prion protein (PrP) segments suggest how structural changes encoded by polymorphism at residue 129 modulate susceptibility to human prion disease. *Journal of Biological Chemistry*, 285:29671–29675.
- Armbruster, P. and Reed, D. H. (2005). Inbreeding depression in benign and stressful environments. *Heredity*, 95:235–242.
- Ashraf, M., Nazir, N., and McNeilly, T. (2001). Comparative salt tolerance of amphidiploid and diploid *Brassica* species. *Plant Science*, 160:683–689.
- Babu, M. M., van der Lee, R., de Groot, N. S., and Gsponer, J. (2011). Intrinsically disordered proteins: regulation and disease. *Current*

- Opinion in Structural Biology*, 21:1–9.
- Balázs, A., Csizmok, V., Buday, L., Rakács, M., Kiss, R., Bokor, M., Udupa, R., Tompa, K., and Tompa, P. (2009). High levels of structural disorder in scaffold proteins as exemplified by a novel neuronal protein, CASK-interactive protein1. *FEBS Journal*, 276:3744–3756.
- Balloux, F., Amos, W., and Coulson, T. (2004). Does heterozygosity estimate inbreeding in real populations? *Molecular Ecology*, 13:3021–3031.
- Barabási, A.-L. and Oltvai, Z. N. (2004). Network biology: Understanding the cell's functional organization. *Nature Reviews Genetics*, 5:101–113.
- Bartels, D. and Salamini, F. (2001). Desiccation tolerance in the resurrection plant *Craterostigma plantagineum*. a contribution to the study of drought tolerance at the molecular level. *Plant Physiology*, 127:1346–1353.
- Basu, M. K., Poliakov, E., and Rogozin, I. B. (2009). Domain mobility in proteins: functional and evolutionary implications. *Briefings in Bioinformatics*, 10:205–216.
- Baxevasis, A. D., Kappas, I., and Abatzopoulos, T. J. (2006). Molecular phylogenetics and asexuality in the brine shrimp *Artemia*. *Molecular Phylogenetics and Evolution*, 40:724–738.
- Bayne, B. L. (2004). Phenotypic flexibility and physiological tradeoffs in the feeding and growth of marine bivalve molluscs. *Integrative and Comparative Biology*, 44:425–432.
- Bayne, B. L., Hedgecock, D., McGoldrick, D., and Rees, R. (1999). Feeding behaviour and metabolic efficiency contribute to growth heterosis in Pacific oysters [*Crassostrea gigas* (Thunberg)]. *Journal of Experimental Marine Biology and Ecology*, 233:115–130.
- Bazin, E., Glémin, S., and Galtier, N. (2006). Population size does not influence mitochondrial genetic diversity in animals. *Science*, 312:570–572.
- Beaton, M. J. and Hebert, P. D. N. (1988). Geographical parthenogenesis and polyploidy in *Daphnia pulex*. *American Naturalist*, 132:837–845.
- Becker, B. and Marin, B. (2009). Streptophyte algae and the origin of embryophytes. *Annals of Botany*, 103:999–1004.
- Becktel, W. J. and Schellman, J. A. (1987). Protein stability curves. *Biopolymers*, 26:1859–1877.
- Bernales, S., Papa, F. R., and Walter, P. (2006). Intracellular signaling by the Unfolded Protein Response. *Annual Reviews of Cell and Developmental Biology*, 22:487–508.
- Bertolani, R. (2001). Evolution of the reproductive mechanisms in tardigrades – a review. *Zoologischer Anzeiger*, 240:247–252.
- Bickler, P. E. and Buck, L. T. (2007). Hypoxia tolerance in reptiles, amphibians, and fishes: life with variable oxygen availability. *Annual Review of Physiology*, 69:145–170.
- Birchler, J. A., Yao, H., Chudalayandi, S., Vaiman, D., and Veitia, R. (2010). Heterosis. *The Plant Cell*, 22:2105–2112.
- Bitan, G., Lomakin, A., and Teplow, D. B. (2001). Amyloid β -protein oligomerization: prenucleation interactions revealed by photo-induced cross-linking of unmodified proteins. *Journal of Biological Chemistry*, 276:35176–35184.
- Borgia, M. B., Nickson, A. A., Clarke, J., and Hounslow, M. J. (2013). A mechanistic model for amorphous protein aggregation of immunoglobulin-like domains. *Journal of the American Chemical Society*, 135:6456–6464.
- Borrell, Y. J., Pineda, H., McCarthy, I., Vázquez, E., Sánchez, J. A., and Lizana, G. B. (2004). Correlations between fitness and heterozygosity at allozyme and microsatellite loci in the Atlantic salmon, *Salmo salar* L. *Heredity*, 92:585–593.
- Boucher, V., Buitink, J., Lin, X., Boudet, J., Hoekstra, F. A., Hundertmark, M., Renard, D., and Leprince, O. (2010). MtPM25 is an atypical hydrophobic late embryogenesis-abundant protein that dissociates cold and desiccation-aggregated proteins. *Plant, Cell and Environment*, 33:418–430.
- Breitkreutz, A., Choi, H., Sharom, J. R., Boucher, L., Neduva, V., Larsen, B., Lin, Z.-Y., Breitkreutz, B.-J., Stark, C., Liu, G., Ahn, J., Dewar-Darch, D., Reguly, T., Tang, X., Almeida, R., Qin, Z. S., Pawson, T., Gingras, A.-C., Nesvizhskii, A. I., and Tyers, M. (2010). A global protein kinase and phosphatase interaction network in yeast. *Science*, 328:1043–1046.
- Brocchieri, L. and Karlin, S. (2005). Protein length in eukaryotic and prokaryotic proteomes. *Nucleic Acids Research*, 33:3390–3400.
- Brochmann, C., Brysting, A. K., Alsos, I. G., Borgen, L., Grundt, H. H., Scheen, A. C., and Elven, R. (2004). Polyploidy in arctic plants. *Biological Journal of the Linnean Society*, 82:521–536.
- Browne, R. A. and Bowen, S. T. (1991). Taxonomy and population genetics of *Artemia*. In Browne, R. A., Sorgeloos, P., and Trotman, C. N., editors, *Artemia Biology*, pages 221–235. CRC Press, Boca Raton, FL.
- Browne, R. A. and MacDonald, G. H. (1982). Biogeography of the brine shrimp *Artemia*: distribution of parthenogenetic and sexual population. *Journal of Biogeography*, 9:331–338.
- Bruce, A. B. (1910). The Mendelian theory of heredity and the augmentation of vigor. *Science*, 32:627–628.
- Bucciantini, M., Calloni, G., Chiti, F., Formigli, L., Nosi, D., Dobson, C. M., and Stefani, M. (2004). Prefibrillar amyloid protein aggregates share common features of cytotoxicity. *Journal of Biological Chemistry*, 279:31374–31382.
- Bustos, D. M. (2012). The role of protein disorder in the 14-3-3 interaction network. *Molecular BioSystems*, 8:178–184.
- Carrió, M., González-Montalbán, N., Vera, A., Villaverde, A., and Ventura, S. (2005). Amyloid-like properties of bacterial inclusion bodies. *Journal of Molecular Biology*, 347:1025–1037.
- Chakrabortee, S., Boschetti, C., Walton, L. J., Sarkar, S., Rubinsztein, D. C., and Tunnacliffe, A. (2008). Hydrophilic protein associated with desiccation tolerance exhibits broad protein stabilization function. *Proceedings of the National Academy of Sciences of the USA*, 104:18073–18078.
- Chang, E. S. (2005). Stressed-out lobsters: crustacean hyperglycemic hormone and stress proteins. *Integrative and Comparative Biology*, 45:43–50.
- Charlesworth, B. (1991). When to be diploid. *Nature*, 351:273–274.
- Charlesworth, B. (1998). The effect of synergistic epistasis on the inbreeding load. *Genetical Research*, 71:85–89.
- Charlesworth, B. and Willis, J. H. (2009). The genetics of inbreeding depression. *Nature Reviews Genetics*, 10:783–796.
- Chebotareva, N. A., Kurganov, B. I., and Livanova, N. B. (2004). Biochemical effects of molecular crowding. *Biochemistry (Moscow)*, 69:1239–1251.
- Chelo, I. M. and Teotónio, H. (2012). The opportunity for balancing selection in experimental populations of *Caenorhabditis elegans*. *Evolution*, 67:142–156.
- Chen, M. and von Mikecz, A. (2005). Formation of nucleoplasmic protein aggregates impairs nuclear function in response to SiO₂ nanoparticles. *Experimental Cell Research*, 305:51–62.
- Chen, P., Liu, X., Zhang, G., and Deng, Y. (2006). Heat-shock protein70 gene expression in four hatchery Pacific Abalone *Haliotis discus hannai* Ino populations using for marker-assisted selection. *Aquaculture Research*, 37:1290–1296.
- Cheng, H.-C., Qi, R. Z., Paudel, H., and Zhu, H.-J. (2011). Regulation and function of protein kinases and phosphatases. *Enzyme Research*, 2011:794089.
- Choe, K. P. and Strange, K. (2008). Genome-wide RNAi screen and in vivo protein aggregation reporters identify degradation of damaged proteins as an essential hypertonic stress response. *American Journal of Physiology: Cell Physiology*, 295:C1488–C1498.
- Clark, M. S. and Worland, M. R. (2008). How insects survive the

- cold: molecular mechanisms – a review. *Journal of Comparative Physiology B*, 178:917–933.
- Cleary, J. P., Walsh, D. M., Hofmeister, J. J., Shankar, G. M., Kuskowski, M. A., Selkoe, D. J., and Ashe, K. H. (2005). Natural oligomers of the amyloid- β protein specifically disrupt cognitive function. *Nature Neuroscience*, 8:79–84.
- Crow, J. F. (1992). Twenty-five years ago in GENETICS: identical triplets. *Genetics*, 130:395–398.
- Crow, J. F. (1998). 90 years ago: the beginning of hybrid maize. *Genetics*, 148:923–928.
- Crow, J. F. and Kimura, M. (1965). Evolution in sexual and asexual populations. *American Naturalist*, 99:439–450.
- Dahiya, V. and Chaudhuri, T. K. (2014). GroEL/GroES accelerates the refolding of a multi-domain protein through modulating on pathway intermediates. *Journal of Biological Chemistry*, 289:286–298.
- Danzmann, R. G., Ferguson, M., and Allendorf, E. (1987). Heterozygosity and oxygen consumption rate as predictors of growth and developmental rate in rainbow trout. *Physiological Zoology*, 60:211–220.
- Darwin, C. R. (1876). *The effects of cross and self fertilisation in the vegetable kingdom*. John Murray, London.
- Davenport, C. B. (1908). Degeneration, albinism and inbreeding. *Science*, 28:454–455.
- de Visser, J. A. G. M., Cooper, T. F., and Elena, S. F. (2011). The causes of epistasis. *Proceedings of the Royal Society of London B*, 278:3617–3624.
- de Visser, J. A. G. M. and Elena, S. F. (2007). The evolution of sex: empirical insights into the roles of epistasis and drift. *Nature Reviews Genetics*, 8:139–149.
- Demontis, D., Pertoldi, C., Loeschcke, V., Mikkelsen, K., Axelsson, T., and Kristensen, T. N. (2009). Efficiency of selection, as measured by single nucleotide polymorphism variation, is dependent on inbreeding rate in *Drosophila melanogaster*. *Molecular Ecology*, 18:4551–4563.
- Dias, C. L., Ala-Nissila, T., Wong-ekkabut, J., Vattulainen, I., Grant, M., and Karttunen, M. (2009). The hydrophobic effect and its role in cold denaturation. *Cryobiology*, 60:91–99.
- Dijksterhuis, J. (2007). Heat-resistant ascospores. In Dijksterhuis, J. and Samson, R. A., editors, *Food Mycology: A multifaceted Approach to Fungi and Food*. CRC Press, Boca Raton, FL.
- Dobson, C. M. (2003). Protein folding and misfolding. *Nature*, 426:884–890.
- Dougan, D. A., Mogk, A., and Bukau, B. (2006). Protein folding and degradation in bacteria: to degrade or not to degrade? That is the question. *CMLS Cellular and Molecular Life Science*, 59:1607–1616.
- Downs, C. A., Kramarsky-Winter, E., Woodley, C. M., Downs, A., Winters, G., Loya, Y., and Ostrander, G. K. (2009). Cellular pathology and histopathology of hypo-salinity exposure on the coral *Stylophora pistillata*. *Science of the Total Environment*, 407:4838–4851.
- Dunker, A. K., Brown, C. J., Lawson, J. D., Iakoucheva, L. M., and Obradović, Z. (2002). Intrinsic disorder and protein function. *Biochemistry*, 41:6573–6582.
- Dunker, A. K., Cortese, M. S., Romero, P., Iakoucheva, L. M., and Uversky, V. N. (2005). Flexible nets: the roles of intrinsic disorder in protein interaction networks. *FEBS Journal*, 272:5129–5148.
- Dunker, A. K., Oldfield, C. J., Meng, J., Romero, P., Yang, J. Y., Chen, J. W., Vacic, V., Obradovic, Z., and Uversky, V. N. (2008). The unfoldomics decade: an update on intrinsically disordered proteins. *BMC Genomics*, 9:S1.
- Duvick, D. N. (2001). Biotechnology in the 1930s: the development of hybrid maize. *Nature Reviews Genetics*, 2:69–74.
- Eanes, W. F. (1978). Morphological variance and enzyme heterozygosity in the monarch butterfly. *Nature*, 276:263–264.
- East, E. M. (1936). Heterosis. *Genetics*, 21:375–397.
- Ekman, D., Björklund, A. K., and Elofsson, A. (2007). Quantification of the elevated rate of domain rearrangements in metazoa. *Journal of Molecular Biology*, 372:1337–1348.
- El-Bermawi, N., Baxevasis, A. D., Abatzopoulos, T. J., Stappen, G. V., and Sorgeloos, P. (2004). Salinity effects on survival, growth and morphometry of four Egyptian *Artemia* populations (International Study on *Artemia*. LXVII). *Hydrobiologia*, 523:175–188.
- Ellis, R. J. (2001). Macromolecular crowding: an important but neglected aspect of the intracellular environment. *Current Opinion in Structural Biology*, 11:114–119.
- Ellis, R. J. and Minton, A. P. (2006). Protein aggregation in crowded environments. *Biological Chemistry*, 387:485–497.
- Elven, R., Murray, D. F., Razzhivin, V., and Yurtsev, B. A. (2003). *Checklist of the Panarctic Flora (PAF). Vascular plants. Final draft version*. University of Oslo, Oslo, Norway.
- Fares, M. A., Ruiz-González, M. X., Moya, A., Elena, S. F., and Barrio, E. (2002). GroEL buffers against deleterious mutations. *Nature*, 417:398.
- Farrar, D. R. (1978). Problems in the identity and origin of the Appalachian *Vittaria* gametophyte, a sporophyteless fern of the Eastern United States. *American Journal of Botany*, 65:1–12.
- Farrar, D. R. (1990). Species and evolution in asexually reproducing independent fern gametophytes. *Systematic Botany*, 15:98–111.
- Fawcett, J. A., Maere, S., and de Peer, Y. V. (2009). Plants with double genomes might have had a better chance to survive the Cretaceous–Tertiary extinction event. *Proceedings of the National Academy of Sciences of the USA*, 106:5737–5742.
- Ferreira, Á. and Amos, W. (2006). Inbreeding depression and multiple regions showing heterozygote advantage in *Drosophila melanogaster* exposed to stress. *Molecular Ecology*, 15:3885–3893.
- Ferreira, S. T., Vieira, M. N. N., and Felice, F. G. D. (2007). Soluble protein oligomers as emerging toxins in Alzheimers and other amyloid diseases. *IUBMB Life*, 59:332–345.
- Fong, J. H. and Panchenko, A. R. (2010). Intrinsic disorder and protein multibinding in domain, terminal, and linker regions. *Molecular BioSystems*, 6:1821–1828.
- Forcada, J. and Hoffman, J. I. (2014). Climate change selects for heterozygosity in a declining fur seal population. *Nature*, 511:462–465.
- Frankham, R. (1995). Inbreeding and extinction: a threshold effect. *Conservation Biology*, 9:792–799.
- Franks, F., Mathias, S. F., and Hatley, R. H. (1990). Water, temperature and life. *Philosophical Transactions of the Royal Society of London B*, 326:517–531.
- Fu, W., Shuai, L., Yao, J., Zheng, B., Zhong, M., and Duan, D. (2010). Molecular cloning and expression analysis of a cytosolic Hsp70 gene from *Ulva pertusa* (Ulvoophyceae, Chlorophyta). *Journal of Applied Phycology*, 23:681–690.
- G. F. Dilly, C. R. Y., Lane, W. S., Pangilinan, J., and Girguis, P. R. (2012). Exploring the limit of metazoan thermal tolerance via comparative proteomics: thermally induced changes in protein abundance by two hydrothermal vent polychaetes. *Proceedings of the Royal Society of London B*, 279:3347–3356.
- Gabrielsen, T. M., Brochmann, C., and Rueness, J. (2002). The Baltic Sea as a model system for studying postglacial colonization and ecological differentiation, exemplified by the red alga *Ceramium tenuicorne*. *Molecular Ecology*, 11:2083–2095.
- Gaff, D. F. and Oliver, M. (2013). The evolution of desiccation tolerance in angiosperm plants: a rare yet common phenomenon. *Functional Plant Biology*, 40:315–328.
- Gajardo, G., Parragué, M., Beardmore, J. A., and Sorgeloos, P.

- (2001). Reproduction in the brine shrimp *Artemia*: evolutionary relevance of laboratory cross-fertility tests. *Journal of Zoology (London)*, 253:25–32.
- Gajardo, G. M. and Beardmore, J. A. (1989). Ability to switch reproductive mode in *Artemia* is related to maternal heterozygosity. *Marine Ecology Progress Series*, 55:191–195.
- Gallardo, M. H., Bickham, J. W., Honeycutt, R. L., Ojeda, R. A., and Köhler, N. (1999). Discovery of tetraploidy in a mammal. *Nature*, 401:341.
- Gallardo, M. H., Gonz'alez, C. A., and Cebri'an, I. (2006). Molecular cytogenetics and allotetraploidy in the red vizcacha rat, *Tympanoctomys barrerae* (Rodentia, Octodontidae). *Genomics*, 88:214–221.
- Gallardo, M. H., Kausel, G., Jiménez, A., Bacquet, C., González, C., Figueroa, J., Köhler, N., and Ojeda, R. (2004). Whole-genome duplications in South American desert rodents (Octodontidae). *Biological Journal of the Linnean Society*, 82:443–451.
- Garay-Arroyo, A., Colmenero-Flores, J. M., Garciarubio, A., and Covarrubias, A. A. (2000). Highly hydrophilic proteins in prokaryotes and eukaryotes are common during conditions of water deficit. *Journal of Biological Chemistry*, 275:5668–5674.
- García-Mata, R., Gao, Y.-S., and Sztul, E. (2002). Hassles with taking out the garbage: aggravating aggregates. *Traffic*, 3:388–396.
- Garton, D. W., Koehn, R. K., and Scott, T. M. (1984). Multiple-locus heterozygosity and the physiological energetics of growth in the coot clam, *Mulinia lateralis*, from a natural population. *Genetics*, 108:445–455.
- Gerstein, A. Z. and Otto, S. P. (2009). Ploidy and the causes of genomic evolution. *Journal of Heredity*, 100:571–581.
- Ginn, B. R. (2010). The implications of gene heterozygosity for protein folding and protein turnover. *Journal of Theoretical Biology*, 265:554–564.
- Glabe, C. G. (2006). Common mechanisms of amyloid oligomer pathogenesis in degenerative disease. *Neurobiology of Aging*, 27:570–575.
- Goff, S. A. (2011). A unifying theory for general multigenic heterosis: energy efficiency, protein metabolism, and implications for molecular breeding. *New Phytologist*, 189:923–937.
- Goldschmidt, L., Teng, P. K., Riek, R., and Eisenberg, D. (2010). Identifying the amyloids, proteins capable of forming amyloid-like fibrils. *Proceedings of the National Academy of Sciences of the USA*, 107:3487–3492.
- Goyal, K., Walton, L. J., and Tunnacliffe, A. (2005). LEA proteins prevent aggregation due to water stress. *Biochemical Journal*, 388:151–157.
- Graham, L. E. and Gray, J. (2001). The origin, morphology, and ecophysiology of early embryophytes: neontological and paleontological perspectives. In Gensel, P. G. and Edwards, D., editors, *Plants Invade the Land: Evolutionary and Environmental Perspectives*, pages 140–158. Columbia University Press, New York, NY.
- Gremer, J. R., Kimball, S., Keck, K. R., Huxman, T. E., Angert, A. L., and Venable, D. L. (2013). Water-use efficiency and relative growth rate mediate competitive interactions in Sonoran Desert winter annual plants. *American Journal of Botany*, 100:2009–2015.
- Grishkan, I., Korol, A. B., Nevo, E., and Wasser, S. P. (2003). Ecological stress and sex evolution in soil microfungi. *Proceedings of the Royal Society of London B*, 270:13–18.
- Groose, R. W., Talbert, L. E., Kojis, W. P., and Bingham, E. T. (1989). Progressive heterosis in autotetraploid alfalfa: studies using two types of inbreds. *Crop Science*, 29:1173–1177.
- Gurevitch, J., Wilson, P., Stone, J. L., Teese, P., and Stoutenburgh, R. J. (1990). Competition among old-field perennials at different levels of soil fertility and available space. *Journal of Ecology*, 78:727–744.
- Haass, C. and Selkoe, D. J. (2007). Soluble protein oligomers in neurodegeneration: lessons from the Alzheimer's amyloid β -peptide. *Nature Reviews Molecular Cell Biology*, 8:101–112.
- Hahn, M. W. (2008). Toward a selection theory of molecular evolution. *Evolution*, 62:255–265.
- Haldane, J. B. S. (1937). The effect of variation on fitness. *American Naturalist*, 71:337–349.
- Hao, G.-Y., Lucero, M. E., Sanderson, S. C., Zacharias, E. H., and Holbrook, N. M. (2013). Polyploidy enhances the occupation of heterogeneous environments through hydraulic related trade-offs in *Atriplex canescens* (Chenopodiaceae). *New Phytologist*, 197:970–978.
- Harrang, E., Lapègue, S., Morga, B., and Bierne, N. (2013). A high load of non-neutral amino-acid polymorphisms explains high protein diversity despite moderate effective population size in a marine bivalve with sweepstakes reproduction. *G3*, 3:333–341.
- Hasegawa, K., Yamaguchi, I., Omata, S., Gejyo, F., and Naiki, H. (1999). Interaction between $A\beta(1-42)$ and $A\beta(1-40)$ in Alzheimer's β -amyloid fibril formation *in vitro*. *Biochemistry*, 38:15514–15521.
- Hawkins, A. J. S., Bayne, B. L., and Day, A. J. (1986). Protein turnover, physiological energetics and heterozygosity in the blue mussel, *Mytilus edulis*: the basis of variable age-specific growth. *Proceedings of the Royal Society of London B*, 229:161–176.
- Hawkins, A. J. S., Bayne, B. L., Day, A. J., Rusin, J., and Worrall, C. M. (1989). Genotype-dependent interrelations between energy metabolism, protein metabolism and fitness. In Ryland, J. S. and Tyler, P. A., editors, *Reproduction, Genetics and Distributions of Marine Organisms, 23rd European Marine Biology Symposium*, pages 283–292. Olsen & Olsen, Fredensborg, Denmark.
- Hawkins, A. J. S. and Day, A. J. (1999). Metabolic interrelations underlying the physiological and evolutionary advantages of genetic diversity. *American Zoology*, 39:401–411.
- Hawkins, A. J. S., Magoulas, A., Héral, M., Bougrier, S., Naciri-Graven, Y., Day, A. J., and Kotoulas, G. (2000). Separate effects of triploidy, parentage and genomic diversity upon feeding behaviour, metabolic efficiency and net energy balance in the Pacific oyster *Crassostrea gigas*. *Genetic Research*, 76:273–284.
- Hawksworth, D. L. (1988). The variety of fungal-algal symbioses, their evolutionary significance, and the nature of lichens. *Botanical Journal of the Linnean Society*, 96:3–20.
- Hayes, S. A. and Dice, J. F. (1996). Roles of molecular chaperones in protein degradation. *Journal of Cell Biology*, 132.
- He, J., Sun, J., and Deem, M. W. (2009). Spontaneous emergence of modularity in a model of evolving individuals and in real networks. *Physical Review E*, 79:031907.
- Hedgecock, D., McGoldrick, D. J., Manahan, D. T., Vavra, J., Appelmans, N., and Bayne, B. L. (1996). Quantitative and molecular genetic analyses of heterosis in bivalve mollusks. *Journal of Experimental Marine Biology and Ecology*, 203:49–59.
- Henter, H. J. (2003). Inbreeding depression and haplodiploidy: experimental measures in a parasitoid and comparisons across diploid and haplodiploid insect taxa. *Evolution*, 57:1793–1803.
- Hessen, D. O., Jeyasingh, P. D., Neiman, M., and Weider, L. J. (2009). Genome streamlining and the elemental costs of growth. *TRENDS in Ecology and Evolution*, 25:75–80.
- Hietz, P. (2010). Fern adaptations to xerix environments. In Mehlreter, K., Walker, L. R., and Sharpe, J. M., editors, *Fern Ecology*, pages 140–176. Cambridge University Press, New York, NY.
- Hiraoka, M. and Yoshida, G. (2010). Temporal variation in isomorphic phase and sex ratios of a natural population of *Ulva pertusa* (Chlorophyta). *Journal of Phycology*, 46:882–888.
- Hochholdinger, F. and Hoecker, N. (2007). Towards the molecular

- basis of heterosis. *TRENDS in Plant Science*, 12:427–432.
- Hoek, C., Mann, D. G., and Jahns, H. M. (1995). *Algae: An Introduction to Phycology*. Cambridge University Press, New York, NY.
- Hoffman, J. I., Forcada, J., and Amos, W. (2010). Exploring the mechanisms underlying a heterozygosity/fitness correlation for canine size in the Antarctic fur seal *Arctocephalus gazella*. *Journal of Heredity*, 101:539–552.
- Hundertmark, M. and Hinch, D. K. (2008). LEA (Late Embryogenesis Abundant) proteins and their encoding genes in *Arabidopsis thaliana*. *BMC Genomics*, 9:118.
- Hunter, K. L., Betancourt, J. L., Riddle, B. R., Devender, T. R. V., Cole, K. L., and Spaulding, W. G. (2001). Ploidy race distributions since the Last Glacial Maximum in the North American desert shrub, *Larrea tridentata*. *Global Ecology and Biogeography*, 10:521–533.
- Hur, J. H., Doninck, K. V., Mandigo, M. L., and Meselson, M. (2009). Degenerate tetraploidy was established before bdelloid rotifer families diverged. *Molecular Biology and Evolution*, 26:375–383.
- Husband, B. C. and Schemske, D. W. (1996). Evolution of the magnitude and timing of inbreeding depression in plants. *Evolution*, 50:54–70.
- Iakoucheva, L. M., Brown, C. J., Lawson, J. D., Obradović, Z., and Dunker, A. K. (2002). Intrinsic disorder in cell-signaling and cancer-associated proteins. *Journal of Molecular Biology*, 323:573–584.
- Iidicula-Thomas, S. and Balaji, P. V. (2007). Protein aggregation: A perspective from amyloid and inclusion-body formation. *Current Science*, 92:758–767.
- Ivankov, D. N., Garbuzynskiy, S. O., Alm, E., Plaxco, K. W., Baker, D., and Finkelstein, A. V. (2003). Contact order revisited: influence of protein size on the folding rate. *Protein Science*, 12:2057–2062.
- Janská, A., Maršík, P., Zelenková, S., and Ovesna, J. (2010). Cold stress and acclimation – what is important for metabolic adjustment? *Plant Biology (Stuttgart, Germany)*, 12:395–405.
- Joly, S., Starr, J. R., Lewis, W. H., and Bruneau, A. (2006). Polyploid and hybrid evolution in roses east of the Rocky Mountains. *American Journal of Botany*, 93:412–425.
- Jones, D. F. (1917). Dominance of linked factors as a means of accounting for heterosis. *Genetics*, 2:466–479.
- Kacser, H. and Burns, J. A. (1981). The molecular basis of dominance. *Genetics*, 97:639–666.
- Kaeuffer, R., Coltman, D. W., Chapuis, J.-L., Pontier, D., and Réale, D. (2007). Unexpected heterozygosity in an island mouflon population founded by a single pair of individuals. *Proceedings of the Royal Society of London B*, 274:527–533.
- Kaganovich, D., Kopito, R., and Frydman, J. (2008). Misfolded proteins partition between two distinct quality control compartments. *Nature*, 454:1088–1096.
- Kayed, R., Head, E., Thompson, J. L., McIntire, T. M., Milton, S. C., Cotman, C. W., and Glabe, C. G. (2003). Common structure of soluble amyloid oligomers implies common mechanism of pathogenesis. *Science*, 300:486–489.
- Kayed, R., Sokolov, Y., Edmonds, B., McIntire, T. M., Milton, S. C., Hall, J. E., and Glabe, C. G. (2004). Permeabilization of lipid bilayers is a common conformation-dependent activity of soluble amyloid oligomers in protein misfolding diseases. *Journal of Biological Chemistry*, 279:46363–46366.
- Kearney, M. R. (2003). Why is sex so unpopular in the Australian desert? *TRENDS in Ecology and Evolution*, 18:605–607.
- Keshet, Y. and Seger, R. (2010). The MAP kinase signaling cascades: a system of hundreds of components regulates a diverse array of physiological functions. *Methods in Molecular Biology*, 661:3–38.
- Ketola, T. and Kotiaho, J. S. (2009). Inbreeding, energy use and condition. *Journal of Evolutionary Biology*, 22:770–781.
- Kiefhaber, T., Rudolph, R., Kohler, H.-H., and Buchner, J. (1991). Protein aggregation *in vitro* and *in vivo*: a quantitative model of the kinetic competition between folding and aggregation. *Nature Biotechnology*, 9:825–829.
- Kimura, M. (1968). Evolutionary rate at the molecular level. *Nature*, 217:624–626.
- Kimura, M. and Crow, J. F. (1978). Effect of overall phenotypic selection on genetic change at individual loci. *Proceedings of the National Academy of Sciences of the USA*, 75:6168–6171.
- King, J., Haase-Pettingell, C., Robinson, A. S., Speed, M., and Mitraki, A. (1996). Thermolabile folding intermediates: inclusion body precursors and chaperonin substrates. *FASEB Journal*, 10:57–66.
- King, J. L. (1967). Continuously distributed factors affecting fitness. *Genetics*, 55:483–492.
- Koch, G. W., Sillett, S. C., Jennings, G. M., and Davis, S. D. (2004). The limits to tree height. *Nature*, 428:851–854.
- Koehn, R. K. and Shumway, S. E. (1982). A genetic/physiological explanation for differential growth rate among individuals of the American oyster, *Crassostrea virginica* (Gmelin). *Marine Biology Letters*, 3:35–42.
- Kondepudi, D. (2008). *Introduction to Modern Thermodynamics*. John Wiley & Sons Ltd., West Sussex, England.
- Konings, W. N., Albers, S.-V., Konging, S., and Driessen, A. J. M. (2002). The cell membrane plays a crucial role in survival of bacteria and archaea in extreme environments. *Antonie van Leeuwenhoek*, 81:61–72.
- Koonin, E. V., Wolf, Y. I., and Karev, G. P. (2002). The structure of the protein universe and genome evolution. *Nature*, 420:218–223.
- Kopito, R. R. (2000). Aggresomes, inclusion bodies and protein aggregation. *TRENDS in Cell Biology*, 10:524–530.
- Kosuda, K. (1972). Synergistic effect of inbreeding on viability in *Drosophila virilis*. *Genetics*, 72:461–468.
- Kranner, I., Beckett, R., Hochman, A., and Nash III, T. H. (2008). Desiccation-tolerance in lichens: a review. *Bryologist*, 111:576–593.
- Krebs, M. R. H., Morozova-Roche, L. A., Daniel, K., Robinson, C. V., and Dobson, C. M. (2004). Observation of sequence specificity in the seeding of protein amyloid fibrils. *Protein Science*, 13:1933–1938.
- Kristensen, T. N., Dahlggaard, J., and Loeschcke, V. (2002). Inbreeding affects *Hsp70* expression in two species of *Drosophila* even at benign temperatures. *Evolutionary Ecology Research*, 4:1209–1216.
- Kristensen, T. N., Pedersen, K. S., Vermeulen, C. J., and Loeschcke, V. (2009). Research on inbreeding in the 'omic' era. *TRENDS in Ecology and Evolution*, 25:44–52.
- Kriwacki, R. W., Hengst, L., Tennant, L., Reed, S. I., and Wright, P. E. (1996). Structural studies of p21^{Waf1/Cip1/Sdi1} in the free and Cdk2-bound state: conformational disorder mediates binding diversity. *Proceedings of the National Academy of Sciences of the USA*, 93:11504–11509.
- Kruse, K. B., Brodsky, J. L., and McCracken, A. A. (2006). Characterization of an *ERAD* gene as *VPS30/ATG6* reveals two alternative and functionally distinct protein quality control pathways: one for soluble Z variant of human α -1 proteinase inhibitor (A1PiZ) and another for aggregates of A1PiZ. *Molecular Biology of the Cell*, 17:203–212.
- Küess, U. (2000). Life history and developmental processes in the basidiomycete *Coprinus cinereus*. *Microbiology and Molecular Biology Reviews*, 64:316–333.
- Kumar, G., Purty, R. S., Sharma, M. P., Singla-Pareek, S. L., and Pareek, A. (2009). Physiological responses among *Brassica* species under salinity stress show strong correlation with transcript

- abundance for SOS pathway-related genes. *Journal of Plant Physiology*, 166:507–520.
- Kunte, H. J., Trüper, H. G., and Stan-Lotter, H. (2002). Halophilic microorganisms. In Horneck, G. and Baumstark-Khan, C., editors, *Astrobiology: The Quest for the Conditions of Life*, pages 185–200. Springer, Berlin, Germany.
- Kurotani, A., Tokmakov, A. A., Kuroda, Y., Fukami, Y., Shinozaki, K., and Sakurai, T. (2014). Correlations between predicted protein disorder and post-translational modifications in plants. *Bioinformatics*, 30:1095–1103.
- Lashuel, H. A., Hartley, D. M., Petre, B. M., Wall, J. S., Simon, M. N., Walz, T., and P. T. Lansbury, J. (2003). Mixtures of wild-type and a pathogenic (E22G) form of A β 40 *in vitro* accumulate protofibrils, including amyloid pores. *Journal of Molecular Biology*, 332:795–808.
- Lee, R. E. (2008). *Phycology*. Cambridge University Press, New York, NY.
- Leffler, E. M., Bullaughey, K., Matute, D. R., Meyer, W. K., Ségurel, L., Venkat, A., Andolfatto, P., and Przeworski, M. (2012). Revisiting an old riddle: what determines genetic diversity levels within species? *PLoS Biology*, 10:e1001388.
- Lehner, B. (2011). Molecular mechanisms of epistasis within and between genes. *TRENDS in Genetics*, 27:323–331.
- LeMosy, E. K., Tan, Y.-Q., and Hashimoto, C. (2001). Activation of a protease cascade involved in patterning the *Drosophila* embryo. *Proceedings of the National Academy of Sciences of the USA*, 98:5055–5060.
- Lerner, I. M. (1954). *Genetic Homeostasis*. Oliver & Boyd, Edinburgh, UK.
- Lesica, P. and Allendorf, F. W. (1992). Are small populations of plants worth preserving? *Conservation Biology*, 6:135–139.
- Levy, Y., Cho, S. S., Shen, T., Onuchic, J. N., and Wolynes, P. G. (2005). Symmetry and frustration in protein energy landscapes: a near degeneracy resolves the Rop dimer-folding mystery. *Proceedings of the National Academy of Sciences of the USA*, 102:2373–2378.
- Lewis, P. A., Tattum, M. H., Jones, S., Bhelt, D., Batchelor, M., Clarke, A. R., Collinge, J., and Jackson, G. S. (2006). Codon 129 polymorphism of the human prion protein influences the kinetics of amyloid formation. *Journal of General Virology*, 87:2443–2449.
- Lewontin, R. C. and Hubby, J. L. (1966). A molecular approach to the study of genic heterozygosity in natural populations. II. Amount of variation and degree of heterozygosity in natural populations of *Drosophila pseudoobscura*. *Genetics*, 54:595–609.
- Li, Z.-K., Luo, L. J., Mei, H. W., Wang, D. L., Shu, Q. Y., Tabien, R., Zhong, D. B., Ying, C. S., Stansel, J. W., Khush, G. S., and Paterson, A. H. (2001). Overdominant epistatic loci are the primary genetic basis of inbreeding depression and heterosis in rice. I. Biomass and grain yield. *Genetics*, 158:1737–1753.
- Liberek, K., Lewandowski, A., and Ziętkiewicz, S. (2008). Chaperones in control of protein disaggregation. *EMBO*, 27:328–335.
- Lippman, Z. B. and Zamir, D. (2007). Heterosis: revisiting the magic. *TRENDS in Genetics*, 23:60–66.
- Liu, J., Perumal, N. B., Oldfield, C. J., Su, E. W., Uversky, V. N., and Dunker, A. K. (2006). Intrinsic disorder in transcription factors. *Biochemistry*, 45:6873–6888.
- Lobell, D. B., Hammer, G. L., McLean, G., Messina, C., Roberts, M. J., and Schlenker, W. (2013). The critical role of extreme heat for maize production in the United State. *Nature Climate Change*, 3:497–501.
- London, J., Skrzynia, C., and Goldberg, M. (1974). Renaturation of *Escherichia coli* tryptophanase in aqueous urea solutions. *European Journal of Biochemistry*, 47:409–415.
- Longton, R. E. (1988). *Biology of Polar Bryophytes and Lichens*. Cambridge University Press, New York, NY.
- López-Otín, C. and Hunter, T. (2010). The regulatory crosstalk between kinases and proteases in cancer. *Nature Reviews Cancer*, 10:278–292.
- Lorenz, D. M., Jeng, A., and Deem, M. W. (2011). The emergence of modularity in biological systems. *Physics of Life Reviews*, 8:129–160.
- Lotz, G. P., Legleiter, J., Aron, R., Mitchell, E. J., Huang, S.-Y., Ng, C., Glabe, C., Thompson, L. M., and Muchowski, P. J. (2010). Hsp70 and Hsp40 functionally interact with soluble mutant Huntingtin oligomers in a classic ATP-dependent reaction cycle. *Journal of Biological Chemistry*, 285:38183–38193.
- Lundmark, M. and Saura, A. (2006). Asexuality alone does not explain the success of clonal forms in insects with geographical parthenogenesis. *Heredity*, 143:23–32.
- Ma, B. and Nussinov, R. (2012). Selective molecular recognition in amyloid growth and transmission and cross-species barriers. *Journal of Molecular Biology*, 421:172–184.
- Mable, B. K. (2004). ‘why polyploidy is rarer in animals than in plants’: myths and mechanisms. *Biological Journal of the Linnean Society*, 82:453–466.
- Mable, B. K. and Otto, S. P. (1998). The evolution of life cycles with haploid and diploid phases. *BioEssays*, 20:453–462.
- Madlung, A. (2013). Polyploidy and its effect on evolutionary success: old questions revisited with new tools. *Heredity*, 110:99–104.
- Maisnier-Patin, S., Roth, J. R., Fredriksson, Å., Nyström, T., Berg, O. G., and Andersson, D. I. (2005). Genomic buffering mitigates the effects of deleterious mutations in bacteria. *Nature Genetics*, 37:1376–1379.
- Mäkinen, H. S., Cano, J. M., and Merilä, J. (2008). Identifying footprints of directional and balancing selection in marine and freshwater three-spined stickleback (*Gasterosteus aculeatus*) populations. *Molecular Ecology*, 17:3565–3582.
- Manning, G., Whyte, D. B., Martinez, R., Hunter, T., and Sudarsanam, S. (2002). The protein kinase complement of the human genome. *Science*, 298:1912–1934.
- Mannini, B., Cascella, R., Zampagni, M., van Waarde-Verhagen, M., Meehan, S., Roodveldt, C., Campioni, S., Boninsegna, M., Penco, A., Relini, A., Kampinga, H. H., Dobson, C. M., Wilson, M. R., Cecchi, C., and Chiti, F. (2012). Molecular mechanisms used by chaperones to reduce the toxicity of aberrant protein oligomers. *Proceedings of the National Academy of Sciences of the USA*, 109:12479–12484.
- Maurizi, M. R. (2002). Love it or cleave it: tough choices in protein quality control. *Nature Structural Biology*, 9:410–412.
- McClellan, A. J., Tam, S., Kaganovich, D., and Frydman, J. (2005). Protein quality control: chaperones culling corrupt conformations. *Nature Cell Biology*, 7:736–741.
- Mead, S., Stumpf, M. P., Whitfield, J., Beck, J. A., Poulter, M., Campbell, T., Uphill, J. B., Goldstein, D., Alpers, M., Fisher, E. M., and Collinge, J. (2003). Balancing selection at the prion protein gene consistent with prehistoric kurulike epidemics. *Science*, 300:640–643.
- Milkman, R. D. (1967). Heterosis as a major cause of heterozygosity in nature. *Genetics*, 55:493–495.
- Mitton, J. B. (1978). Relationship between heterozygosity for enzyme loci and variation of morphological characters in natural populations. *Nature*, 273:661–662.
- Mitton, J. B. (1985). Shell shape variation in the blue mussel *Mytilus edulis* L., and its association with enzyme heterozygosity. *Journal of Experimental Marine Biology and Ecology*, 90:73–80.
- Mitton, J. B. (1993). Enzyme heterozygosity, metabolism, and developmental stability. *Genetica*, 89:47–65.
- Mitton, J. B. (1995). Enzyme heterozygosity and developmental

- stability. *Acta Theriologica*, 3:33–54.
- Mitton, J. B. (1997). *Selection in Natural Populations*. Oxford University Press, New York, NY.
- Mitton, J. B. (1998). Molecular markers and natural selection. In Carvalho, G. R., editor, *Advances in Molecular Ecology*, pages 225–242. IOS Press, Amsterdam, Netherlands.
- Mitton, J. B. and Grant, M. C. (1984). Associations among protein heterozygosity, growth rate, and developmental homeostasis. *Annual Review of Ecology and Systematics*, 15:479–499.
- Mitton, J. B. and Pierce, B. A. (1980). The distribution of individual heterozygosity in natural populations. *Genetics*, 95:1043–1054.
- Mohsen, A. F., Khaleata, A. F., Hashem, M. A., and Metwalli, A. (1974). Effect of different nitrogen sources on growth, reproduction, amino acid, fat and sugar contents in *Ulva fasciata* Delile (part iii). *Botanica Marina*, 17:218–222.
- Mohsen, A. F., Nasr, A. H., and Metwalli, A. M. (1973). Effect of temperature variations on growth, reproduction, amino acid synthesis, fat and sugar content in *Ulva fasciata* Delile plants. *Hydrobiologia*, 42:451–460.
- Monari, M., Foschi, J., Rosmini, R., Marin, M. G., and Serrazanetti, G. P. (2011). Heat shock protein 70 response to physical and chemical stress in *Chamelea gallina*. *Journal of Experimental Marine Biology and Ecology*, 397:71–78.
- Morell, M., Bravo, R., Espargaró, A., Sisquella, X., Avilés, F. X., Fernández-Busquets, X., and Ventura, S. (2008). Inclusion bodies: specificity in their aggregation process and amyloid-like structure. *Biochimica et Biophysica Acta*, 1783:1815–1825.
- Muller, H. J. (1925). Why polyploidy is rarer in animals than in plants. *American Naturalist*, 59:346–353.
- Nelsen, R. and Eisenberg, D. (2006). Recent atomic models of amyloid fibril structure. *Current Opinion in Structural Biology*, 16:260–265.
- Nelson, R., Sawaya, M. R., Balbirnie, M., Madsen, A. Ø., Riek, C., Grothe, R., and Eisenberg, D. (2005). Structure of the cross- β spine of amyloid-like fibrils. *Nature*, 435:773–778.
- Netzer, W. J. and Hartl, F. U. (1997). Recombination of protein domains facilitated by co-translational folding in eukaryotes. *Nature*, 388:343–349.
- Nevo, R., Brumfeld, V., Kapon, R., Hinterdorfer, P., and Reich, Z. (2005). Direct measurement of protein energy landscape roughness. *EMBO*, 6:482–486.
- Newman, M. C. and Unger, M. A. (2003). *Fundamentals of Ecotoxicology*. CRC Press, Boca Raton, FL.
- NIST/SEMATECH (2003). e-Handbook of Statistical Methods. <http://www.itl.nist.gov/div898/handbook/apr/apr.htm>. Accessed: 2014-06-25.
- Oldfield, C. J., Meng, J., Yang, J. Y., Yang, M. Q., Uversky, V. N., and Dunker, A. K. (2008). Flexible nets: disorder and induced fit in the associations of p53 and 14-3-3 with their partners. *BMC Genomics*, 9:S1.
- Oldham, A. R., Sillett, S. C., Tomescu, A. M. F., and Koch, G. W. (2010). The hydrostatic gradient, not light availability, drive height-related variation in *Sequoia sempervirens* (Cupressaceae) leaf anatomy. *American Journal of Botany*, 97:1087–1097.
- Oliver, M. J., Velten, J., and Mishler, B. D. (2005). Desiccation tolerance in bryophytes: a reflection of the primitive strategy for plant survival in dehydrating habitats? *Integrative and Comparative Biology*, 45:788–799.
- Olzscha, H., Schermann, S. M., Woerner, A. C., Pinkert, S., Hecht, M. H., Tartaglia, G. G., Vendruscolo, M., Hayer-Hartl, M., Hartl, F. U., and Vabulas, R. M. (2011). Amyloid-like aggregates sequester numerous metastable proteins with essential cellular functions. *Cell*, 144:67–78.
- O’Nuallain, B., Williams, A. D., Westermarck, P., and Wetzell, R. (2004). Seeding specificity in amyloid growth induced by heterologous fibrils. *Journal of Biological Chemistry*, 279:17490–17499.
- Onuchic, J. N., Wolynes, P. G., Luthey-Schulten, Z., and Socci, N. D. (1995). Towards an outline of the topography of a realistic protein folding funnel. *Proceedings of the National Academy of Sciences of the USA*, 92:3626–3630.
- Orr, H. A. and Otto, S. P. (1994). Does diploidy increase the rate of adaptation? *Genetics*, 136:1475–1480.
- Otto, C. R., Snodgrass, J. W., Forester, D. C., Mitchell, J. C., and Miller, R. W. (2007). Climatic variation and the distribution of an amphibian polyploid complex. *Journal of Animal Ecology*, 76:1053–1061.
- Otto, S. P. (2007). The evolutionary consequences of polyploidy. *Cell*, 131:452–462.
- Otto, S. P. and Gerstein, A. C. (2008). The evolution of haploidy and diploidy. *Current Biology*, 18:R1121–R1124.
- Otto, S. P. and Goldstein, D. B. (1992). Recombination and the evolution of diploidy. *Genetics*, 131:745–751.
- Otto, S. P. and Whitton, J. (2000). Polyploid incidence and evolution. *Annual Review of Genetics*, 34:401–437.
- Pannell, J. R., Obbard, D. J., and Buggs, R. J. A. (2004). Polyploidy and the sexual system; what can we learn from *Mercurialis annua*? *Biological Journal of the Linnean Society*, 82:547–560.
- Park, J.-M., Chen, M., Wang, D., and Deem, M. W. (2015). Modularity enhances the rate of evolution in a rugged fitness landscape. *Physical Biology*, 12:025001.
- Patterson, C. and Höhfeld, J. (2006). molecular chaperones and the ubiquitin-proteasome system. In Mayer, R. J., Ciechanover, A., and Rechsteiner, M., editors, *Protein Degradation, Vol. 2: The Ubiquitin-Proteasome System*, pages 1–30. Wiley-VCH, Weinheim, Germany.
- Patthy, L. (2003). Modular assembly of genes and the evolution of new functions. *Genetica*, 118:217–231.
- Pawson, T. and Nash, P. (2000). Protein–protein interactions define specificity in signal transduction. *Genes & Development*, 14:1027–1047.
- Pedersen, K. S., Kristensen, T. N., and Loeschcke, V. (2005). Effects of inbreeding and rate of inbreeding in *Drosophila melanogaster* – Hsp70 expression and fitness. *Journal of Evolutionary Biology*, 18:756–762.
- Picard, D. (2002). Heat-shock protein 90, a chaperone for folding and regulation. *CMLS Cellular and Molecular Life Sciences*, 59:1640–1648.
- Pogson, G. H. and Fevolden, S. E. (1998). DNA heterozygosity and growth rate in the Atlantic cod *Gadus morhua* (L). *Evolution*, 52:915–920.
- Poon, S., Treweek, T. M., Wilson, M. R., Easterbrook-Smith, S. B., and Carver, J. A. (2002). Clusterin is an extracellular chaperone that specifically interacts with slowly aggregating proteins on their off-folding pathway. *FEBS Letters*, 513:259–266.
- Pörtner, H. O. (2001). Climate change and temperature-dependent biogeography: oxygen limitation of thermal tolerance in animals. *Naturwissenschaften*, 88:137–146.
- Potts, M. (1994). Desiccation tolerance of prokaryotes. *Microbiology Reviews*, 58:755–805.
- Powers, L. (1944). An expansion of Jones theory for the explanation of heterosis. *American Naturalist*, 78:275–280.
- Prestrelski, S. J., Tedeschi, N., Arakawa, T., and Carpenter, J. F. (1993). Dehydration-induced conformational transitions in proteins and their inhibition by stabilizers. *Biophysical Journal*, 65:661–671.
- Proctor, M. C. F. and Tuba, Z. (2002). Poikilohydry and homoihydric: antithesis or spectrum of possibilities? *New Phytologist*, 156:327–349.
- Pujolar, J. M., Maes, G. E., Vancoillie, C., and Volckaert, F. A. M.

- (2005). Growth rate correlates to individual heterozygosity in the European eel, *Anguilla anguilla* (L). *Evolution*, 59:189–199.
- Querol, A. and Bond, U. (2009). The complex and dynamic genomes of industrial yeasts. *FEMS Microbiology Letters*, 293:1–10.
- Rajan, R. S., Illing, M. E., Bence, N. F., and Kopito, R. R. (2001). Specificity in intracellular protein aggregation and inclusion body formation. *Proceedings of the National Academy of Sciences of the USA*, 98:13060–13065.
- Ramirez, J.-M., Folkow, L. P., and Blix, A. S. (2007). Hypoxia tolerance in mammals and birds: from the wilderness to the clinic. *Annual Review of Physiology*, 69:113–143.
- Ramsey, J. (2011). Polyploidy and ecological adaptation in wild yarrow. *Proceedings of the National Academy of Sciences of the USA*, 108:7096–7101.
- Ramshini, H., Parrini, C., Relini, A., Zampagni, M., Mannini, B., Pesce, A., Saboury, A. A., Nemat-Gorgani, M., and Chiti, F. (2011). Large proteins have a great tendency to aggregate but a low propensity to form amyloid fibrils. *PLoS One*, 6:e16075.
- Ravaux, J., Hamel, G., Zbinden, M., Tasiemski, A. A., Boutet, I., Léger, N., Tanguy, A., Jollivet, D., and Shillito, B. (2013). Thermal limit for metazoan life in question: in vivo heat tolerance of the Pompeii worm. *PLoS One*, 8:e64074.
- Ricca, M., Beecher, F. W., Boles, S. B., Tensch, E., Greilhuber, J., Karlin, E. F., and Shaw, A. J. (2008). Cytotype variation and allopolyploidy in North American species of the *Sphagnum subsecundum* complex (Sphagnaceae). *American Journal of Botany*, 95:1606–1620.
- Riddle, N. C. and Birchler, J. A. (2008). Comparative analysis of inbred and hybrid maize at the diploid and tetraploid levels. *Theoretical and Applied Genetics*, 116:563–576.
- Rinehart, J. P., Haywood, S. A. L., Elnitsky, M. A., Sandro, L. H., Lee, Jr., R. E., and Denlinger, D. L. (2006). Continuous up-regulation of heat shock proteins in larvae, but not adults, of a polar insect. *Proceedings of the National Academy of Sciences of the USA*, 103:14223–14227.
- Robertson, F. and Reeve, E. C. (1952). Heterozygosity, environmental variation and heterosis. *Nature*, 170:286–287.
- Rochet, J.-C., Conway, K. A., and P. T. Lansbury, J. (2000). Inhibition of fibrillization and accumulation of prefibrillar oligomers in mixtures of human and mouse α -synuclein. *Biochemistry*, 39:10619–10626.
- Rodriguez, M. C. S., Edsgård, D., Hussain, S. S., Alquezar, D., Rasmussen, M., Gilbert, T., Nielsen, B. H., Bartels, D., and Mundy, J. (2010). Transcriptomes of the desiccation-tolerant resurrection plant *Craterostigma plantagineum*. *Plant Journal*, 63:212–228.
- Rossi, B. E., Debandi, G. O., Peralta, I. E., and Martinez Palle, E. (1999). Comparative phenology and floral patterns in *Larrea* species (Zygophyllaceae) in the Monte desert (Mendoza, Argentina). *Journal of Arid Environments*, 43:213–226.
- Rousseau, F., Schymkowitz, J., and Itzhaki, L. S. (2012). Implications of 3D domain swapping for protein folding, misfolding and function. *Advances in Experimental Medicine and Biology*, 747:137–152.
- Roy, M., Brodeur, J., and Cloutier, C. (2003). Temperature and sex allocation in a spider mite. *Oecologia*, 135:322–326.
- Rubin, G. M., Yandell, M. D., Wortman, J. R., Miklos, G. L. G., Nelson, C. R., Hariharan, I. K., Fortini, M. E., Li, P. W., Apweiler, R., Fleischmann, W., Cherry, J. M., Henikoff, S., Skupski, M. P., Misra, S., Ashburner, M., Birney, E., Boguski, M. S., Brody, T., Brokstein, P., Celniker, S. E., Chervitz, S. A., Coates, D., Cravchik, A., Gabrielian, A., Galle, R. F., Gelbart, W. M., George, R. A., Goldstein, L. S. B., Gong, F., Guan, P., Harris, N. L., Hay, B. A., Hoskins, R. A., Li, J., Li, Z., Hynes, R. O., Jones, S. J. M., Kuehl, P. M., Lemaitre, B., Littleton, J. T., Morrison, D. K., Mungall, C., O'Farrell, P. H., Pickeral, O. K., Shue, C., Vossell, L. B., Zhang, J., Zhao, Q., Zheng, X. H., Zhong, F., Zhong, W., Gibbs, R., Venter, J. C., Adams, M. D., and Lewis, S. (2000). Comparative genomics of the eukaryotes. *Science*, 287:2204–2215.
- Rubinsztein, D. C. (2006). The roles of intracellular protein-degradation pathways in neurodegeneration. *Nature*, 443:780–786.
- Rumball, W., Franklin, I. R., Frankham, R., and Sheldon, B. L. (1994). Decline in heterozygosity under full-sib and double first-cousin inbreeding in *Drosophila melanogaster*. *Genetics*, 136:1039–1049.
- Russell, N. J. (1998). Molecular adaptations in psychrophilic bacteria: potential for biotechnological applications. *Advances in Biochemical Engineering/Biotechnology*, 61:1–21.
- Salar, R. K. and Aneja, K. R. (2007). Thermophilic fungi: taxonomy and biogeography. *International Journal of Agricultural Technology*, 3:77–107.
- Sangster, T. A., Lindquist, S., and Queitsch, C. (2004). Under cover: causes, effects and implications of Hsp90-mediated genetic capacitance. *BioEssays*, 26:348–362.
- Schadt, C. W., Martin, A. F., Lipson, D. A., and Schmidt, S. K. (2003). Seasonal dynamics of previously unknown fungal lineages in tundra soils. *Science*, 301:1359–1361.
- Schaller, A. (2004). A cut above the rest: the regulatory function of plant proteases. *Planta*, 220:183–197.
- Schröder, M. and Kaufman, R. J. (2005). ER stress and the unfolded protein response. *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis*, 569:29–63.
- Schuettpelz, E., Grusz, A. L., Windham, M. D., and Pryer, K. M. (2008). The utility of nuclear *gapCp* in resolving polyploid fern origins. *Systematic Botany*, 33:621–629.
- Schultz, J., Milpetz, F., Bork, P., and Ponting, C. P. (1998). SMART, a simple modular architecture research tool: identification of signaling domains. *Proceedings of the National Academy of Sciences of the USA*, 95:5857–5864.
- Schween, G., Schulte, J., Reski, R., and Hohe, A. (2005). Effect of ploidy level on growth, differentiation, and morphology in *Physcomitrella patens*. *Bryologist*, 108:27–35.
- Senock, R. S., Barrow, J. R., Gibbens, R. P., and Herbel, C. H. (1991). Ecophysiology of the polyploid shrub *Atriplex canescens* (Chenopodiaceae) growing *in situ* in the northern Chihuahuan Desert. *Journal of Arid Environments*, 21:45–57.
- Shankar, G. M., Li, S., Mehta, T. H., Garcia-Munoz, A., Shepardson, N. E., Smith, I., Brett, F. M., Farrell, M. A., Rowan, M. J., Lemere, C. A., Regan, C. M., Walsh, D. M., Sabatini, B. L., and Selkoe, D. J. (2008). Amyloid β -protein dimers isolated directly from Alzheimer brains impair synaptic plasticity and memory. *Nature Medicine*, 14:837–842.
- Shannon, M. C. and Grieve, C. M. (1999). Tolerance of vegetable crops to salinity. *Scientia Horticulturae*, 78:5–38.
- Sharpe, J. M. and Mehlreter, K. (2010). Ecological insights from fern population dynamics. In Mehlreter, K., Walker, L. R., and Sharpe, J. M., editors, *Fern Ecology*, pages 61–110. Cambridge University Press, New York, NY.
- Shoemaker, B. A., Portman, J. L., and Wolynes, P. G. (2000). Speeding molecular recognition by using the folding funnel: the fly-casting mechanism. *Proceedings of the National Academy of Sciences of the USA*, 97:8868–8873.
- Shull, G. H. (1948). What is “heterosis”? *Genetics*, 33:439–446.
- Siller, E., DeZwaan, D. C., Anderson, J. F., Freeman, B. C., and Barral, J. M. (2010). Slowing bacterial translation speed enhances eukaryotic protein folding efficiency. *Journal of Molecular Biology*, 396:1310–1318.
- Silow, M. and Oliveberg, M. (1997). Transient aggregates in protein folding are easily mistaken for folding intermediates. *Proceedings*

- of the National Academy of Sciences of the USA, 94:6084–6086.
- Silva, A. D., Luikart, G., Yoccoz, N. G., Cohas, A., and Allainé, D. (2006). Genetic diversity–fitness correlation revealed by microsatellite analyses in European alpine marmots (*Marmota marmota*). *Conservation Genetics*, 7:371–382.
- Singer, M. A. and Lindquist, S. (1998). Multiple effects of trehalose on protein folding *in vitro* and *in vivo*. *Molecular Cell*, 1:639–648.
- Singh, S. K., Kohle, P., Wang, W., and Nema, S. (2009). Large-scale freezing of biologics – a practitioner’s review, part one: fundamental aspects. *BioProcess International*, 7:32–44.
- Singh, S. M. and Zouros, E. (1978). Genetic variation associated with growth rate in the American oyster (*Crassostrea virginica*). *Evolution*, 32:342–353.
- Sinha, N. and Nussinov, R. (2001). Point mutations and sequence variability in proteins: redistributions of preexisting populations. *Proceedings of the National Academy of Sciences of the USA*, 98:3139–3144.
- Smallwood, M. and Bowles, D. J. (2002). Plants in a cold climate. *Philosophical Transactions of the Royal Society of London B*, 357:831–847.
- Sollars, V. E., Lu, X., Xiao, L., Wang, X., and Garfinkel, M. D. (2003). Evidence for an epigenetic mechanism by which Hsp90 acts as a capacitor for morphological evolution. *Nature Genetics*, 33:70–74.
- Sopory, S. K. and Munshi, M. (1998). Protein kinases and phosphatases and their role in cellular signaling in plants. *Critical Reviews in Plant Sciences*, 17:245–318.
- Soulé, M. (1979). Heterozygosity and developmental stability: another look. *Evolution*, 33:396–401.
- Speed, M. A., Wang, D. I., and King, J. (1996). Specific aggregation of partially folded polypeptide chains: the molecular basis of inclusion body composition. *Nature Biotechnology*, 14:1283–1287.
- Spellenberg, R. (1981). Polyploidy in *Dalia formosa* (Fabaceae) on the Chihuahuan Desert. *Brittonia*, 33:309–324.
- Stajich, J. A., Berbee, M. L., Blackwell, M., Hibbett, D. S., James, T. Y., Spatafora, J. W., and Taylor, J. W. (2009). Primer – the fungi. *Current Biology*, 19:R840–R845.
- Stark, L. R., Oliver, M. J., Mishler, B. D., and McLetchie, D. N. (2007). Generational differences in response to desiccation stress in the desert moss *Tortula inermis*. *Annals of Botany*, 99:53–60.
- Stebbins, G. L. (1950). *Variation and Evolution in Plants*. Columbia University Press, New York, NY.
- Stebbins, G. L. (1984). Polyploidy and the distribution of the arctic-alpine flora: new evidence and a new approach. *Botanica Helvetica*, 94:1–13.
- Stroud, J. C., Liu, C., Teng, P. K., and Eisenberg, D. (2012). Toxic fibrillar oligomers of amyloid- β have cross- β structure. *Proceedings of the National Academy of Sciences of the USA*, 109:7717–7722.
- Stumpf, M. P. H., Thorne, T., de Silva, E., Stewart, R., An, H. J., Lappe, M., and Wiuf, C. (2008). Estimating the size of the human interactome. *Proceedings of the National Academy of Sciences of the USA*, 105:6959–6964.
- Sun, J. and Deem, M. W. (2007). Spontaneous emergence of modularity in a model of evolving individuals. *Physical Review Letters*, 99:228107.
- Svartman, M., Stone, G., and Stanyon, R. (2005). Molecular cytogenetics discards polyploidy in mammals. *Genomics*, 85:425–430.
- Sved, J. A., Reed, T. A., and Bodmer, W. F. (1967). The number of balanced polymorphisms that can be maintained in a natural population. *Genetics*, 55:469–481.
- Szulkin, M., Bierne, N., and David, P. (2010). Heterozygosity–fitness correlations: a time for reappraisal. *Evolution*, 64:1202–1217.
- Tahiri-Alaoui, A., Sim, V. L., Caughey, B., and James, W. (2006). Molecular heterosis of prion protein β -oligomers: A potential mechanism of human resistance to disease. *Journal of Biological Chemistry*, 281:34171–34178.
- Tajima, F. (1983). Evolutionary relationship of DNA sequences in finite populations. *Genetics*, 105:437–460.
- Tal, M. and Gardi, I. (1976). Physiology of polyploid plants: water balance in autotetraploid and diploid tomato under low and high salinity. *Physiologia Plantarum*, 38:257–261.
- Tan, T., Frenkel, D., Gupta, V., and Deem, M. W. (2005). Length, protein-protein interactions, and complexity. *Physica A: Statistical Mechanics and its Applications*, 350:52–62.
- Tartaglia, G. G., Pawar, A. P., Campioni, S., Dobson, C. M., Chiti, F., and Vendruscolo, M. (2008). Prediction of aggregation-prone regions in structured proteins. *Journal of Molecular Biology*, 380:425–436.
- Tartaglia, G. G., Pechmann, S., Dobson, C. M., and Vendruscolo, M. (2007). Life on the edge: a link between gene expression levels and aggregation rates of human proteins. *TRENDS in Biochemical Sciences*, 32:204–206.
- Teta, P., Pardiñas, U. F. J., Udrizar Sauthier, D. E., and Gallardo, M. H. (2014). A new species of the tetraploid vizcacha rat *Tympanoctomys* (Caviomorpha, Octodontidae) from central Patagonia, Argentina. *Journal of Mammalogy*, 95:60–71.
- Tetzlaff, J. E., Putcha, P., Outeiro, T. F., Ivanov, A., Berezovska, O., Hyman, B. T., and McClean, P. J. (2008). CHIP targets toxic α -synuclein oligomers for degradation. *Journal of Biological Chemistry*, 283:17962–17968.
- Thamatrakoln, K. and Hildebrand, M. (2008). Silicon uptake in diatoms revisited: a model for saturable and nonsaturable uptake kinetics and the role of silicon transporters. *Plant Physiology*, 146:1397–1407.
- Tine, M., Bonhomme, F., McKenzie, D. J., and Durand, J.-D. (2010). Differential expression of the heat shock protein Hsp70 in natural populations of the tilapia, *Sarotherodon melanothron*, acclimatised to a range of environmental salinities. *BMC Ecology*, 10:11.
- Tocidlowski, M. E., Merrill, C. L., Loomis, M. R., and Wright, J. F. (2001). Teratoma in desert grassland whiptail lizards (*Cnemidophorus uniparens*). *Journal of Zoo and Wildlife Medicine*, 32:257–259.
- Tomba, P. (2002). Intrinsically unstructured proteins. *TRENDS in Biochemical Sciences*, 27:527–533.
- Tomba, P. (2005). The interplay between structure and function in intrinsically unstructured proteins. *FEBS Letters*, 579:3346–3354.
- Tomba, P., Szász, C., and Buday, L. (2005). Structural disorder throws new light on moonlighting. *Trends in Biochemical Sciences*, 30:484–489.
- Tordai, H., Nagy, A., Farkas, K., Bányai, L., and Patthy, L. (2005). Modules, multidomain proteins and organismic complexity. *FEBS Journal*, 272:5064–5078.
- Tortajada, A. M., Carmona, M. J., and Serra, M. (2009). Does haplodiploidy purge inbreeding depression in rotifer populations? *PLoS One*, 4:e8195.
- Trapero-Casas, A. and Kaiser, W. J. (2007). Differences between ascospores and conidia of *Didymella rabiei* in spore germination and infection of chickpea. *Phytopathology*, 97:1600–1607.
- Trivedi, S., Rao, S. R., and Gehlot, H. S. (2005). Nucleic acid stability in thermophilic prokaryotes: a review. *Journal of Cell and Molecular Biology*, 4:61–69.
- Turk, B. (2006). Targeting proteases: successes, failures and future prospects. *Nature Reviews Drug Discovery*, 5:785–799.
- Tyagi, M., Shoemaker, B. A., Bryant, S. H., and Panchenko, A. R. (2009). Exploring functional roles of multibinding protein interfaces. *Protein Science*, 18:1674–1683.
- Uversky, V. N. and Dunker, A. K. (2010). Understanding protein

- non-folding. *Biochimica et Biophysica Acta*, 1804:1231–1264.
- Uversky, V. N., Oldfield, C. J., and Dunker, A. K. (2008). Intrinsically disordered proteins in human diseases: introducing the D² concept. *Annual Review of Biophysics*, 37:215–246.
- Uversky, V. N., Oldfield, C. J., Midic, U., Xie, H., Xue, B., Vucetic, S., Iakoucheva, L. M., Obradovic, Z., and Dunker, A. K. (2009). Unfoldomics of human diseases: linking protein intrinsic disorder with diseases. *BMC Genomics*, 10:S7.
- Vabulas, R. M., Raychaudhuri, S., Hayer-Hartl, M., and Hartl, F. U. (2010). Protein folding in the cytoplasm and the heat shock response. *Cold Spring Harbor Perspectives in Biology*, 2:a004390.
- van der Hoorn, R. A. L. (2008). plant proteases: from phenotypes to molecular mechanisms. *Annual Review of Plant Biology*, 59:191–223.
- van Leeuwen, M. R., Krijgheld, P., Bleichrodt, R., Menke, H., Stam, H., Stark, J., Wösten, H. A. B., and Dijksterhuis, J. (2013). Germination of conidia of *Aspergillus niger* is accompanied by major changes in RNA profiles. *Studies in Mycology*, 74:59–70.
- Veerman, A. (1985). Diapause. In Helle, W. and Sabelis, M. W., editors, *Spider Mites: Their Biology, Natural Enemies and Control*, pages 279–316. Elsevier, Amsterdam, Netherlands.
- Ventura, S. (2005). Sequence determinants of protein aggregation: tools to increase protein solubility. *Microbial Cell Factories*, 4:11.
- Vieira, M. N. N., Forný-Germano, L., Saraiva, L. M., Sebollela, A., Martinez, A. M. B., Houzel, J.-C., de Felice, F. G., and Ferreira, S. T. (2007). Soluble oligomers from a non-disease related protein mimic A β -induced tau hyperphosphorylation and neurodegeneration. *Journal of Neurochemistry*, 103:736–748.
- Vinayagam, A., Stelzl, U., Foulle, R., Plassmann, S., Zenkner, M., Timm, J., Assmus, H. E., and Andrade-Navarro, M. A. (2011). A directed protein interaction network for investigating intracellular signal transduction. *Science Signaling*, 4:rs8.
- Vinayagam, A., Zirin, J., Roesel, C., Hu, Y., Yilmazel, B., Samsonova, A. A., Neumüller, R. A., Mohr, S. E., and Perrimon, N. (2014). Integrating protein-protein interaction networks with phenotypes reveals signs of interactions. *Nature Methods*, 11:94–99.
- Wagner, C. and Løkke, H. (1991). Estimation of ecotoxicological protection levels from NOEC toxicity data. *Water Research*, 25:1237–1242.
- Walhout, A. J. M. (2006). Unraveling transcription regulatory networks by protein–dna and protein–protein interaction mapping. *Genome Research*, 16:1445–1454.
- Wang, D., Hsieh, M., and Li, W.-H. (2005). A general tendency for conservation of protein length across eukaryotic kingdoms. *Molecular Biology and Evolution*, 22:142–147.
- Wang, H., Guo, S., Huang, M., Thorsten, L. H., and Wei, J. (2010). Ascomycota has a faster evolutionary rate and higher diversity than Basidiomycota. *Science China*, 53:1163–1169.
- Wang, L., Maji, S. K., Sawaya, M. R., Eisenberg, D., and Riek, R. (2008). Bacterial inclusion bodies contain amyloid-like structure. *PLoS Biology*, 6:e195.
- Wang, W. and Roberts, C. J. (2013). Non-arrhenius protein aggregation. *AAPS Journal*, 15:840–851.
- Wang, X. Q., Yang, P. F., Liu, Z., Liu, W. Z., Hu, Y., Chen, H., Kuang, T. Y., Pei, Z. M., Shen, S. H., and He, Y. K. (2009). Exploring the mechanism of *Physcomitrella patens* desiccation tolerance through a proteomic strategy. *Plant Physiology*, 149:1739–1750.
- Wang, Y.-C. and Chen, B.-S. (2010). Integrated cellular network of transcription regulations and protein-protein interactions. *BMC Systems Biology*, 4:20.
- Ward, J. J., Sodhi, J. S., McGuffin, L. J., Buxton, B. F., and Jones, D. T. (2004). Prediction and functional analysis of native disorder in proteins from the three kingdoms of life. *Journal of Molecular Biology*, 337:635–645.
- Warringer, J. and Blomberg, A. (2006). Evolutionary constraints on yeast protein size. *BMC Evolutionary Biology*, 6:61.
- Waters, E. R., Aebermann, B. D., and Sanders-Reed, Z. (2008). Comparative analysis of the small heat shock proteins in three angiosperm genomes identifies new subfamilies and reveals diverse evolutionary patterns. *Cell Stress Chaperones*, 13:127–142.
- Watkins, Jr., J. E., Mack, M. C., Sinclair, T. R., and Mulkey, S. S. (2007). Ecological and evolutionary consequences of desiccation tolerance in tropical fern gametophytes. *New Phytologist*, 176:708–717.
- Wei, G., Mousseau, N., and Derreumaux, P. (2007). Computational simulations of the early steps of protein aggregation. *Prion*, 1:3–8.
- Wheeler, J. R., Grist, E. P. M., Leung, K. M. Y., Morrill, D., and Crane, M. (2002). Species sensitivity distributions: data and model choice. *Marine Pollution Bulletin*, 45:192–202.
- White, D. A., Buell, A. K., Knowles, T. P. J., Welland, M. E., and Dobson, C. M. (2010). Protein aggregation in crowded environments. *Journal of the American Chemical Society*, 132:5170–5175.
- Williams, W. (1959). Heterosis and the genetics of complex characters. *Nature*, 184:527–530.
- Willis, J. H. (1993). Effects of different levels of inbreeding on fitness components in *Mimulus guttatus*. *Evolution*, 47:864–876.
- Wills, C. (1978). Rank-order selection is capable of maintaining all genetic polymorphisms. *Genetics*, 89:403–417.
- Wright, C. F., Teichmann, S. A., Clarke, J., and Dobson, C. M. (2005). The importance of sequence diversity in the aggregation and evolution of proteins. *Nature*, 438:878–881.
- Wright, P. E. and Dyson, H. J. (1999). Intrinsically unstructured proteins: Re-assessing the protein structure-function paradigm. *Journal of Molecular Biology*, 293:321–331.
- Wyatt, A. R., Yerbury, J. J., Ecroyd, H., and Wilson, M. R. (2013). Extracellular chaperones and proteostasis. *Annual Reviews of Biochemistry*, 82:295–322.
- Wyatt, R., Odrzykoski, I. J., Stoneburner, A., Bass, H. W., and Galau, G. A. (1988). Allopolyploidy in bryophytes: multiple origins of *Plagiomnium medium*. *Proceedings of the National Academy of Sciences of the USA*, 85:5601–5604.
- Xia, K., Fu, Z., Hou, L., and Han, J.-D. J. (2008). Impacts of protein-protein interaction domains on organism and network complexity. *Genome Research*, 18:1500–1508.
- Xiong, Y.-C., Li, F.-M., and Zhang, T. (2006). Performance of wheat crops with different chromosome ploidy: root-sourced signals, drought tolerance, and yield performance. *Planta*, 224:710–718.
- Xu, L., Shan, S., and Wang, X. (2013). Single point mutation alters the microstate dynamics of amyloid β protein A β 42 as revealed by dihedral dynamics analyses. *Journal of Physical Chemistry B*, 117:6206–6216.
- Yagi, H., Kusaka, E., Hongo, K., baobata, T., and Kawata, Y. (2005). Amyloid fibril formation of α -synuclein is accelerated by preformed amyloid seeds of other proteins: implications for the mechanism of transmissible conformational diseases. *Journal of Biological Chemistry*, 280:38609–38616.
- Yorimitsu, T. and Klionsky, D. J. (2007). Eating the endoplasmic reticulum: quality control by autophagy. *TRENDS Cell Biology*, 17:279–285.
- Zhang, L. and King, C. E. (1993). Life history divergence of sympatric diploid and polyploid populations of brine shrimp *Artemia parthenogenetica*. *Oecologia*, 93:177–183.
- Zhang, L. and Lefcort, H. (1991). The effects of ploidy level on the thermal distributions of brine shrimp *Artemia parthenogenetica* and its ecological implications. *Heredity*, 66:445–452.
- Zhang, T., Faraggi, E., Li, Z., and Zhou, Y. (2013). Intrinsically semi-disordered state and its role in induced folding and protein aggregation. *Cell Biochemistry and Biophysics*, 67:11931205.

- Zhdanov, V. P. and Kasemo, B. (2004). Kinetics of protein aggregation with formation of unreactive intermediates. *Langmuir*, 20:2543–2545.
- Zhou, G., Chen, Y., Yao, W., Zhang, C., Xie, W., Hua, J., Xing, Y., Xiao, J., and Zhang, Q. (2012). Genetic composition of yield heterosis in an elite rice hybrid. *Proceedings of the National Academy of Sciences of the USA*, 109:15847–15852.
- Zhu, D., Porter, W. R., Long, M. A., Fraunhofer, W., Gleason, K. M., and Gao, Y. (2010). Using heat conduction microcalorimetry to study thermal aggregation kinetics of proteins. *Thermochimica Acta*, 499:1–7.
- Zmasek, C. M. and Godzik, A. (2011). Strong functional patterns in the evolution of eukaryotic genomes revealed by the reconstruction of ancestral protein domain repertoires. *Genome Biology*, 12:R4.
- Zmasek, C. M. and Godzik, A. (2012). This *déjà vu* feeling—analysis of multidomain protein evolution in eukaryotic genomes. *PLoS Computational Biology*, 8:e1002701.
- Zou, D., Gao, K., Xia, J., Xu, Z., Zhang, X., and Liu, S. (2007). Responses of dark respiration in the light to desiccation and temperature in the intertidal macroalga, *Ulva lactuca* (Chlorophyta) during emersion. *Phycologia*, 46:363–370.
- Zouros, E., Singh, S. M., and Miles, H. E. (1980). Growth rate in oysters: an overdominant phenotype and its possible explanations. *Evolution*, 34:856–867.

Appendix A.

The purpose of this appendix is to highlight some applications of the theory presented in the main text in light of some papers published since the text's publication. The applications involve the evolution of syngamy, polygenic adaptation, and the evolution of sexual reproduction. I think our understanding of each of these phenomena is enhanced by the theory of truncation selection for heterozygosity presented in the main text. I do not think the ideas presented here warrant a new paper at the present moment, but I think the ideas may serve as food for thought, which may benefit other theorists. I will start with the evolution of syngamy, which I regard to be the strongest section of this appendix. I will then move on to polygenic adaptation and the evolution of sexual reproduction, which is much more speculative.

Appendix A.1. The evolution of syngamy

One of the key ideas presented in the main text is that heterozygosity throughout an organism's genome confers resistance to physical stresses that promote protein aggregation. Figure 5b depicts two truncation curves for different levels of stress. Individuals with sufficiently high heterozygosity will survive a given level of stress. In contrast, individuals with too low of a heterozygosity will have a very low, if not negligible, fitness. The critical heterozygosity level depends on the stressfulness of the environment. The more stressful the environment, the higher the requisite heterozygosity for surviving in the environment will be. Hence, individuals with low heterozygosity will only survive in mildly stressful environments.

The relationship between heterozygosity and stress tolerance has important implications for spores. The main text briefly touched on this subject in Section 4.5, but I would like to elaborate more here. Many fungal species produce both asexual spores, usually referred to as conidia, and sexual spores, which have many different names. For the purposes of simplicity, I will focus on ascospores and basidiospores for the sexual spores. There is ample evidence that sexual fungal spores are more stress tolerant than asexual spores (Dijksterhuis 2007; Dijksterhuis et al. 2007; Trapero-Casas and Kaiser 2007). In fact, Grishkan et al. (2003) showed that the abundance of sexual spores increases relative to asexual spores along an aridity gradient in Israel, suggesting that that likelihood of fungal species producing sexual spores increases with aridity.

The theory presented in the main text provides a reason for the relative stress tolerance of fungal spores. Sexual spores are produced after meiosis has occurred and should possess a mixed cytoplasm. Typically, only the nucleus of the proto-spore cell undergoes meiosis. The four daughter nuclei then align, and barriers form between the nuclei to produce four separate spore cells (Mauseth 2009; Neiman 2005). Therefore, sexual spores can inherit a mixed cytoplasm that contains two different allozymes, even though the spore is technically haploid. In contrast, asexual spores are produced via mitosis of haploid parent cells and have no possibility of containing a mixed cytoplasm. Thus, sexual spores should have lower total rates of soluble oligomer formation than asexual spores, and thus greater stress tolerance (Sections 2 and 3 of the main text).

Spore longevity plays a role as well. Hong et al. (1997) shows that the longevity of conidia decreases with temperature and desiccation stress. This would make sense if the toxicity of soluble oligomers played a key role in spore longevity. Suppose that spore viability followed an equation similar to Equation 16 from the main text. There would be some concentration of total soluble oligomers that would kill 50% of the spores (represented as C_{50} in Equation 16). The concentration of soluble oligomers in the spore would be affected by both time and the rate at which the oligomers form. Spores in a stressful environment would reach the C_{50} concentration faster than spores in a milder environment. If possessing a mixed cytoplasm lowers the rate at which soluble oligomers accumulate, then sexual spores should have higher longevities than asexual spores under equivalent environmental conditions.

Everything that I have summarized in this section can now be used to develop a theory for the evolution of syngamy. Since sexual spores possess higher stress tolerances and longevities than asexual spores, they should be able to weather harsher environmental conditions. Dijksterhuis et al. (2007) and Grishkan et al. (2003) provide evidence for the greater hardiness of sexual spores relative to asexual spores. Thus, the ability to produce sexual spores would provide an immediate advantage to possessing a mixed cytoplasm that contains different allozymes for a given protein. For haploid organisms, the easiest way to obtain a mixed cytoplasm is by fusing two cells with different cytoplasm contents together, a process usually called syngamy.

Thus, haploid organisms may fuse their cells with the cells from other individuals in order to obtain a mixed cytoplasm that enables the production of spores that can

withstand harsh conditions. This benefit would apply not only to fungi, but also to other organisms possessing an extended haploid stage in their life-cycle, such as green algae and bryophytes. This would provide an immediate advantage to the evolution of syngamy in haploid organisms.

Since the benefits of syngamy only arise if the fusing cells contain different cytoplasm contents, the organisms undergoing syngamy must guarantee that their cytoplasm contents are different. This may underlie the evolution of mating systems in organisms with extended haploid life-stages. Most such organisms follow a simple +/- mating system, wherein half the population consists of (+)-cells and the other half consists of (-)-cells. Only cells of opposite sign can fuse together, thus decreasing the likelihood that the fusing cells contain similar cytoplasm contents. The green algae *Chlamydomonas reinhardtii* is an example of a species with such a life cycle. However, some fungi have evolved quite elaborate mating systems to reduce the risk of relatives fusing together (Kothe 1996).

Sibling avoidance may also affect the timing of meiosis itself. Some spores, such as zygosporangia and the spores of *C. reinhardtii*, undergo meiosis after spore germination. Other spores, such as ascospores, undergo meiosis prior to spore formation. The former process would seem to guarantee that the spores contain a mixed cytoplasm since the spores are diploid, but the process would also guarantee that siblings occupy the same area since meiosis occurs after germination. On the other hand, the latter process would allow sibling spores to disperse separately, thereby reducing the odds that siblings germinate in the same area. Thus, the latter process would be beneficial for reducing the risk of siblings undergoing syngamy. However, the latter process would only work if meiosis can be carried out in such a way as to generate haploid spores with mixed cytoplasm contents (Mauseth 2009; Neiman 2005).

Finally, I want to discuss why cells started fusing together in the first place. Desiccation and desiccating agents, such as polyethylene glycol (PEG), are known to promote fusion of membranes by weakening hydrophobic interactions (Roy et al. 2016; Erkut et al. 2012; Pedrazzoli et al. 2011; Hoekstra et al. 2001; MacDonald 1985). Thus, it is quite possible that cell fusion first occurred by accident when cells were exposed to desiccating conditions. Ironically, the cells that fused together were more likely to survive the desiccating conditions because their cytoplasm contents were mixed. The necessary allozyme diversity could have been provided by nearly neutral mutations that had accumulated in the population or by hybridization

between closely related species. Later, cells would evolve mechanisms to make the fusion process less haphazard and more systematic, which would give rise to proper syngamy.

The resulting fusion process would be similar to the parasexual cycle that has been observed in the yeast *Candida albicans*, which undergoes cell fusion under stressful conditions and increases its ploidy from diploid to tetraploid. *C. albicans* does not undergo traditional meiosis in order to reduce its ploidy level. Instead, it destroys chromosomal material until the diploid state is restored, hence why the process is called parasexual (Sherwood and Bennett 2009; Forche et al. 2008; Bennett and Johnson 2003). Organisms that alternate between haploid and diploid life-stages may have used a similar parasexual process prior to the evolution of crossing over, chromosome segregation, and proper meiosis.

Thus, the benefits of heterozygosity discussed in the main text may provide an evolutionary advantage to syngamy that is independent of recombination. The reduction of soluble oligomer formation would have provided an immediate evolutionary benefit to fusing cells. Hence, the evolution of syngamy to form hardier spores may have been a prerequisite step to the evolution of meiosis and proper sexual reproduction.

Appendix A.2. Polygenic adaptation and the benefits of sexual reproduction

Several authors have argued that adaptation, especially in humans, may be highly polygenic, resulting from allele frequency shifts across many gene loci (Gneccchi-Ruscione et al. 2018; Bergey et al. 2018; Boyle et al. 2017; Berg and Coop 2014; Hancock et al. 2010; Pritchard and Rienzo 2010; Pritchard et al. 2010). However, there is some debate over the likelihood of such adaptation and proper interpretation of the data (Höllinger et al. 2019; Berg et al. 2019). For the sake of argument, this section will assume that such adaptation occurs.

Höllinger et al. (2019) extensively modeled the conditions under which polygenic adaptation is likely to occur. They concluded that there were only two such conditions. Either the background mutation rate for the trait must be very large (requiring that the trait is controlled by hundreds or thousands of gene loci), or genetic variation must be maintained by balancing selection. The authors rule out the possibility of the former condition since it is unlikely that many traits are controlled by so many gene loci, and the authors are skeptical of the second condition since it would require balancing selection to be operating on many gene loci.

Therefore, Höllinger et al. (2019) argues that adaptation typically occurs from some combination of hard and soft sweeps.

However, one of the goals of my paper was to argue that balancing selection could be operating throughout the genomes of complex organisms. The truncation selection for heterozygosity model developed in the main text could easily maintain polymorphisms at frequencies necessary for polygenic adaptation Wills (1978). Moreover, the model should allow any number of alleles to be maintained at a given gene locus, as long as they do not co-aggregate. The model also does not require selection to occur at any particular gene locus, as long as individuals in the population have a sufficiently high heterozygosity. This generality of the truncation selection model means that any number of alleles could be segregating within a population at elevated frequencies, thereby providing a substantial source of genetic diversity within the population.

An important implication emerges when the ideas from my paper are combined with Höllinger et al. (2019). If polygenic adaptation requires balancing selection, and balancing selection is mostly restricted to complex organisms, then polygenic adaptation should also be restricted to complex organisms. In contrast, simple organisms (especially prokaryotes) should primarily adapt through a combination of hard and soft sweeps (e.g. Barroso-Batista et al. (2014) and Barrick and Lenski (2009)). Thus, selective sweeps at a few gene loci likely play a larger role in prokaryotic adaptation, while polygenic adaptation via allele frequency shifts is likely more common in complex plants and animals (Radwan and Babik 2012). Note that the modes of adaptation are not dichotomous, instead they represent opposite ends of an adaptation spectrum. Hence, simple organisms adapt in a more sweep-like fashion whereas the adaptation of complex organisms is more “polygenicish.”

If this theory is correct, then the level of genetic diversity is not the variable that causes simple and complex organisms to adapt differently. Instead, differences in the composition of standing genetic variation are what cause populations of simple and complex organisms to adapt differently. In the absence of balancing selection, the relative frequencies of alleles that constitute a population’s standing variation are determined by mutation-selection-drift balance (Höllinger et al. 2019). The relative allele frequencies will be highly heterogeneous, with a few alleles occurring at relatively high frequencies and many alleles occurring at very low frequencies. When the environment shifts, some of the alleles in the pool of

standing variation will become beneficial. The few beneficial alleles that achieve high frequencies due to genetic drift are more likely to become established in the population (i.e. achieve a frequency that is less influenced by genetic drift, which allows further frequency changes to be more deterministic). As a result, only a few beneficial alleles will become established in the population at any given time and adaptation will be sweep-like. In contrast, balancing selection can homogenize the distribution of allele frequencies in the pool of standing genetic variation by elevating the frequencies of many alleles above the level maintained by mutation-selection-drift balance. This allows beneficial alleles at more gene loci to become established simultaneously when the environment shifts. Since alleles at many gene loci have reached frequencies that allow for more deterministic frequency changes, adaptation is more polygenic with balancing selection (Höllinger et al. 2019). Therefore, balancing selection allows populations of complex organisms to adapt in a more polygenic manner than populations of simple organisms, even though the populations of complex organisms may contain smaller pools of standing genetic variation (due to smaller population sizes, for example).

My primary interest in this subject is not the mechanism of adaptation, per se, but rather its implications for the evolution of sexual reproduction. Hickey and Golding (2018) argues that sexual reproduction could be maintained in natural populations, even if asexually reproducing individuals have a two-fold reproductive advantage, when adaptation is polygenic. The authors support their hypothesis with computer simulations that generate results similar to those published in previous papers (Roze 2014; Otto and Barton 2001). Hence, the hypothesis presented in Hickey and Golding (2018) is not wholly original, but is an update to a theoretical approach that extends all the way back to August Weismann, R. A. Fisher, and Hermann Muller (Hickey and Golding 2018).

The simulations in Hickey and Golding (2018) consider a population of organisms with a single chromosome containing 100 bi-allelic gene loci. At each gene locus, an allele occurring at 5% frequency within the population is undergoing positive selection. The authors point out that the probability of a single chromosome containing the favored allele at 25 gene loci is less than one in a billion. Hence, no individual organism within the population will possess the favored allele at all 100 gene loci.

The results of the simulation show that each favored

allele increases from 5% to approximately 35% on average. Thus, adaptation was occurring via allele frequency shifts. Furthermore, the sexually reproducing individuals outcompete the asexually reproducing individuals even though the asexual individuals possess a two-fold reproductive advantage over the sexual individuals. Therefore, sexual reproduction may be favored in species undergoing repeated polygenic adaptation.

Here is where my paper comes into the picture. Remember that the main text says protein aggregation becomes worse as organisms increase in complexity. This is because complexity is facilitated by a class of “managerial proteins” that have numerous interaction partners. The managerial proteins serve as hubs in complex protein interaction networks, coordinating various activities and keeping the regulatory and signal transduction machinery running in sync. Among the ranks of these managerial proteins are kinases, proteases, phosphatases, transcription factors, etc. (see main text for elaboration). The managerial proteins are typically large and multi-domain, which facilitates numerous protein interactions. The proteins also typically contain intrinsically unstructured regions (IURs) that allow the proteins to assume multiple conformations (they are metastable), which provides another method of facilitating numerous interactions. Unfortunately, these physical properties tend to make the managerial proteins prone to aggregation. Thus, complex organisms face a greater protein aggregation burden than simple organisms and are more likely to experience truncation selection for heterozygosity. For this reason, complex organisms are typically diploid, whereas simple organisms are typically haploid (see main text for more elaboration).

Combining the hypotheses of Höllinger et al. (2019), Hickey and Golding (2018), and my paper together yields a potential explanation for why sexual reproduction is widespread among complex organisms, but rarer among simple organisms. Sexual reproduction is evolutionarily advantageous when adaptation occurs via allele frequency shifts at many, possibly hundreds of loci (Hickey and Golding 2018). However, this mechanism of adaptation requires balancing selection to be operating on many gene loci throughout the genome (Höllinger et al. 2019). Such balancing selection only operates on organisms of sufficient complexity to experience a heavy protein aggregation burden (my paper). ***Thus, sexual reproduction is widespread among complex organisms because they experience the balancing selection necessary to make polygenic adaptation work. In contrast, simple organisms***

typically reproduced asexually because they do not experience balancing selection and are more likely to adapt via selective sweeps.

It is almost as if simple and complex organisms are living in different worlds. Simple organisms do not experience truncation selection for heterozygosity, are haploid, and adapt via a combination of hard and soft sweeps. Complex organisms do experience truncation selection for heterozygosity, are diploid, and adapt via frequency shifts at many gene loci (polygenic adaptation). It would seem that genetic diversity is much more important for complex organisms than for simple organisms, and this explains why complex organisms are more likely to reproduce sexually.

Is the hypothesis true? I do not know. The hypothesis is highly speculative, but I think it does warrant some consideration because it may assist other theorists thinking about these matters, which is why I have included it in this appendix.

Appendix A.3. Recombination and simple organisms

Many critics would object to the claim that simple organisms do not reproduce sexually (e.g. Vos 2009). There is no doubt that homologous recombination is widespread among bacteria, for example. However, there are important differences. Recombination in bacteria is not coupled to reproduction, and bacteria can go many generations without recombination. In contrast, animals generally undergo sexual reproduction every generation (though cyclical parthenogens are a notable exception). Bacterial recombination only modifies a short and localized sequence of DNA in the bacterial chromosome, whereas crossing over in eukaryotes typically involves recombination at multiple sites along the chromosome, resulting in more shuffling of genetic material. As a consequence, bacterial recombination usually reduces genetic diversity while sexual reproduction usually increases genetic diversity (Iranzo et al. 2019; Ambur et al. 2016; Gorelick and Heng 2010). Therefore, I am partial to alternative explanations for bacterial recombination than those used to explain sexual reproduction in organisms that undergo a complete meiotic cycle.

My preferred explanation for bacterial homologous recombination is that it assists with DNA repair (Bernstein et al. 2011; Bernstein et al. 1989). Bernstein et al. (2011) summarizes many empirical and theoretical justifications for the DNA repair hypothesis. The most convincing are that Rec proteins play a role in both DNA repair and homologous recombination, and that homologous recombination seems to be upregulated

in bacteria exposed to conditions that promote DNA damage.

If DNA repair is the primary function of homologous recombination in bacteria, and combining beneficial alleles is the primary function of crossing over in complex eukaryotes, then there must have been a transition in function during the evolution of eukaryotes. My current thinking on the matter is that the transition occurred over four phases:

1. Homologous recombination: Homologous recombination evolved in the prokaryotes because it plays an important role in DNA repair (Bernstein et al. 2011; Bernstein et al. 1989). Prokaryotes do not alternate between haploid and diploid generations because protein aggregation is not a substantial enough problem for them to benefit from heterozygosity, even under stressful conditions (main text). DNA repair probably reduces genetic diversity in prokaryotic populations (Iranzo et al. 2019; Ambur et al. 2016; Gorelick and Heng 2010), but this may not significantly affect the ability of prokaryotes to adapt because their adaptation is typically driven by hard and soft sweeps rather than via polygenic adaptation, which relies more on standing genetic variation (see previous section).
2. Syngamy: Protein aggregation still does not impose much of a burden for simple eukaryotes, so they can survive with a prolonged haploid generation. However, protein aggregation is more burdensome under stressful conditions, and these eukaryotes survive such conditions by producing spores with a mixture of allozymes. The eukaryotes undergo syngamy to produce the needed mixed cytoplasm and, thus, are temporarily diploid (this appendix). The result is a parasexual cycle wherein extra DNA is lost by degradation analogous to what is seen in *C. albicans* (Sherwood and Bennett 2009; Forche et al. 2008; Bennett and Johnson 2003). DNA repair is still the primary function of homologous recombination.
3. Synapsis: A key step in the evolution of meiosis from mitosis is the synapsis of homologous chromosomes (Wilkins and Holliday 2009). DNA repair proteins seem to have been recruited to assist with the formation of chiasmata (Bernstein et al. 2011). The formation of chiasmata may have initially improved the fidelity of recombination (by preventing ectopic recombination for example), but the chiasmata

also help homologous chromosomes to properly align, so they can segregate correctly during meiosis and ensure that each daughter cell has a proper karyotype (Bernstein et al. 2011; Gorelick and Heng 2010; Wilkins and Holliday 2009; Heng 2007). A true meiotic cycle exists by the time this phase is completed, which allows some eukaryotes to undergo systematic transitions between haploid and diploid life stages. This alternation between haploid and diploid life stages is widespread among fungi, algae, and plants.

4. Polygenic Adaptation: Some eukaryotes (e.g. animals) are primarily diploid. Meiosis would seem to be unnecessary for these species, which could reproduce solely through mitosis. Yet, meiosis has not been lost in these species because sexual reproduction is beneficial when species adapt to their environment via allele frequency shifts (Hickey and Golding 2018). Ironically, balancing selection is needed to maintain polymorphisms at sufficient frequencies for polygenic adaptation to occur (Höllinger et al. 2019), but diploid species are subjected to truncation selection for heterozygosity because they produce many aggregation-prone proteins (main text). That is why the species are diploid to begin with. In this final phase, crossing over is clearly generating genetic diversity that enables rapid environmental adaptation rather than solely facilitating proper chromosome segregation during meiosis, as in the previous phase.

If this four-phase sequence is correct, then meiosis did not originally evolve to support sexual reproduction. Instead, the original function of meiosis was to enable eukaryotes to systematically alternate between haploid and diploid life stages. Meiosis has since been maintained in diploid-dominant species because it promotes polygenic adaptation. Thus, the benefits of meiosis are near-term for haplodiplontic eukaryotes, but long-term for diploid eukaryotes.

Appendix A.4. Asexual reproduction among complex organisms

Two sections ago, I argued that the prokaryotes do not reproduce sexually because they are not complex enough to be subjected to the balancing selection that is required for polygenic adaptation to work. However, some complex organisms that should experience truncation selection for heterozygosity do not reproduce sexually. This section will attempt to explain why.

Haag and Ebert (2004) provides an excellent foundation for explaining asexuality among complex organisms. The authors of the paper argue that asexual species are often found in marginal habitats where species experience many local extinction and recolonization events. The species living in these environments should experience many genetic bottlenecks (founder effect), which would cause substantial genetic drift within the species' populations. As a consequence, these species should experience a decline in genetic diversity over time. The decline in genetic diversity would inflict a large fitness cost on a sexual population because the average individual heterozygosity would decrease with each passing generation. The sexual population would essentially become inbred over time. An asexual population that lacks recombination would not experience this fitness cost because the heterozygosity within each individual organism would be fixed at a constant value, so the asexual population would not experience any inbreeding depression.

The theory presented in Haag and Ebert (2004) works well for asexual hybrids, polyploids, apomictic parthenogens, and any other asexual species with low recombination rates and fixed heterozygosity. However, the theory must be incomplete because some asexual species, such as self-fertilizers and automictic parthenogens, are highly inbred (Barrière and Félix 2007; Stenøien et al. 2005; Birky 1996). Thus, fixed heterozygosity cannot be the sole advantage of asexual reproduction. However, I do think Haag and Ebert (2004) is on the right track.

Any species subjected to metapopulation dynamics should experience substantial genetic bottlenecks and drift as described in Haag and Ebert (2004). Polygenic adaptation would be impossible in such circumstances because genetic drift would govern allele frequency changes, making adaptation via allele frequency shifts unlikely. Since sexual reproduction requires polygenic adaptation (see Appendix A.2), a species following metapopulation dynamics would not benefit from sexual reproduction. However, sexual reproduction would still be accompanied by fitness costs, such as the two-fold cost of males. Hence, asexual reproduction should be more advantageous than sexual reproduction for species following metapopulation dynamics, as described in Haag and Ebert (2004). This revised hypothesis should apply to highly inbred asexual species as well as those with fixed heterozygosity.

The importance of heterozygosity has diminished in the revised theory, but it is not irrelevant. Most, but not all, asexual species can be placed in one of two

broad categories. There are the highly heterozygous asexual species (hybrids and polyploids) and there are the minimally heterozygous species (self-fertilizers and automicts) (Jaron et al. 2018). Both types of asexual species might occupy ephemeral environments that promote metapopulation dynamics, but the harshness of the environments should be considerably different. The highly heterozygous asexual species should be associated with stressful environments that promote high levels of protein aggregation while the minimally heterozygous asexual species should be associated with very mild environments that promote very low levels of protein aggregation. I want to discuss each kind of environment in some detail.

The highly heterozygous asexual species should occupy environments that promote high levels of protein aggregation. As discussed in Section 4.1 of the main text, polyploid and hybrid species are often found in such environments. Asexual species are also found in harsh environments (Lundmark and Saura 2006; Tilquin and Kokko 2016), which is consistent with the metapopulation hypothesis because populations in harsh environments may be restricted local patches that are temporarily hospitable. These species most closely conform to Haag and Ebert (2004)'s original metapopulation hypothesis because metapopulation dynamics should result in a significant loss of heterozygosity in sexual populations. Thus, there are two reasons why asexual species are associated with really harsh environments: (1) asexual species can retain high levels of heterozygosity better than sexual species and (2) genetic drift renders polygenic adaptation ineffectual, thereby removing the main advantage sexual species have over asexual species.

The bdelloid rotifers could provide a good example of highly heterozygous asexual species. Bdelloid rotifers are degenerate tetraploids, and many species survive in environments where the rotifers are routinely exposed to desiccation stress (Nowell et al. 2018; Hur et al. 2009; Welch et al. 2008). However, Nowell et al. (2018) has found that two limnoterrestrial bdelloid species that are frequently exposed to desiccating conditions possess much higher homologous sequence divergence than two aquatic bdelloid species that are not frequently exposed to desiccation. I think this could be evidence that the limnoterrestrial species are exposed to selection for high levels of heterozygosity. Some form of mitotic recombination seems to be responsible for the homogenization of the aquatic species' genomes (Nowell et al. 2018), and these processes should operate on the limnoterrestrial species as well, but the recurring exposure to desiccation stress

that limnoterrestrial species experience has probably prevented the limnoterrestrial species from losing as much heterozygosity as the aquatic species. Thus, limnoterrestrial bdelloids probably benefit from asexual reproduction because it helps maintain high levels of heterozygosity as described in Haag and Ebert (2004). Furthermore, bdelloids constitute 95% of all rotifers in limnoterrestrial environments, but only 20-30% of all rotifers in aquatic environments (Ricci 1987), which is more evidence that asexual species are more competitive in ephemeral environments than in more permanent environments.

Now consider the opposite extreme of asexual species with very low heterozygosity levels, such as self-fertilizers and automictic parthenogens. According to the main text, such species should only be able to survive if they never experience high levels of protein aggregation. Otherwise, they would be weeded out by truncation selection for heterozygosity. These species should be subject to metapopulation dynamics because the very mild conditions in which they can survive should only last temporarily in localized patches. Genetic drift should inhibit polygenic adaptation in these species, thereby removing the advantage of sexual reproduction as described previously. In addition, these species are subject to low levels of balancing selection, a key requirement for polygenic adaptation (Höllinger et al. 2019), which would also render sexual reproduction unbeneficial. Thus, asexual reproduction should be common in species that rarely experience high levels of protein aggregation for two reasons: (1) the absence of balancing selection weakens the efficacy of polygenic adaptation and (2) genetic drift also renders polygenic adaptation ineffectual, thereby removing the main advantage sexual species have over asexual species. *Caenorhabditis elegans* and *Arabidopsis thaliana* could serve as examples of this type of asexual species (Barrière and Félix 2007; Stenøien et al. 2005).

I think it would be useful here to make a distinction between stress tolerating species and stress avoiding species. I see this as a generalization of the freeze tolerating and freeze avoiding distinction made in the insect literature (Sinclair 1999; Bale 1996). Briefly, freeze tolerating species can withstand the formation of ice in their tissues while freeze avoiding species rely on supercooling to prevent the formation of ice in their tissues. Similarly, I think stress tolerating species can withstand the stresses of their environment, but stress avoiding species have evolved mechanisms to limit the damage that physical stresses might inflict upon them. Mechanisms used by stress avoiding species include

supercooling, vitrification, and dormancy within seeds or encysted eggs (Crowe et al. 1998; Siepel 1994).

Thus, self-fertilizers and automictic parthenogens with very low heterozygosity levels may be stress avoiding species, and hybrid parthenogens and asexual polyploids with high heterozygosity levels may be stress tolerating species. *C. elegans*⁸ and the bdelloid rotifers may serve as models for the two types of species. *C. elegans* dauer larva are able to survive desiccation, but only after they have been preconditioned for 48 hours at 98% relative humidity. During the preconditioning period, *C. elegans* synthesis trehalose, heat shock proteins (HSPs), and late embryogenesis abundant proteins (LEAs) that prevent protein aggregation from occurring (Erkut et al. 2013). The necessity of a precondition period, the metapopulation dynamics, and the low genetic diversity of *C. elegans* all suggest that it is a typical stress avoiding species (Teotónio et al. 2017; Erkut et al. 2013; Barrière and Félix 2007). In contrast, many bdelloid rotifer individuals can survive a rapid drop in humidity from 95% to 40% over one hour (Caprioli and Ricci 2001). The high genetic diversity of bdelloid rotifers combined with their ability to survive rapid desiccation would suggest that bdelloid rotifers are typical stress tolerant species.

I have spent much of this section discussing asexual species whose heterozygosity levels are at the high and low end of the spectrum. This is because hybridization, polyploidization, self-fertilization, and automixis are common routes to asexual reproduction. However, asexual species do not have to fall into the two categories I have been discussing. The important issue is whether genetic drift is interfering with polygenic adaptation, which would remove the main advantage of sexual reproduction regardless of the stressfulness of the environment. However, there are extra factors that promote asexual reproduction in stress-tolerating and stress-avoiding species, namely asexual reproduction allows for fixed heterozygosity levels in stress-tolerating species, and the absence of balancing selection in stress-avoiding species provides additional interference inhibiting polygenic adaptation. Therefore, stress-tolerating and stress-avoiding asexual species have extra reinforcements promoting asexual reproduction that other asexual species solely following metapopulation dynamics do not.

I would like to finish this section by briefly mentioning some other asexual species that provide support for the metapopulation hypothesis. The

⁸*A. thaliana* could also serve as a model for stress avoidance (Hannah et al. 2006; Reyes-Díaz et al. 2006)

ant species *Mycocephurus smithii* contains sexual and asexual populations. The sexual populations are concentrated in the central part of the species distribution around the Amazon River Basin whereas the asexual species are spread throughout South America and Central America (Rabeling et al. 2011). The oribatid mites contain numerous asexual species (Maraun et al. 2009; Palmer and Norton 1992). Asexual oribatid species are rare in arboreal environments, but are common in soil environments (Domes et al. 2007). In fact, asexual oribatid species increase in frequency with soil depth, perhaps because the distribution of food resources becomes increasingly patchy (Smelansky 2006). Asexual springtail (Collembola) species also increase in frequency with soil depth (Chernova et al. 2010). The gecko species *Nactus Arnouxii* contains sexual and asexual populations. The sexual populations are found in Australia, Papua New Guinea, and neighboring islands whereas the asexual populations are found in farther away Pacific islands extending out into Micronesia (Moritz 1987). This provides more evidence for the significance of genetic drift in promoting asexual reproduction. Two asexual species of night lizard, *Lepidophyma reticulatum* and *Lepidophyma lavimaculatum*, are found in Costa Rica and Panama, respectively (Sinclair et al. 2010). These species are located at the southernmost end of their genus's range, which may again signify a role for genetic drift in promoting asexual reproduction. Asexual non-marine ostracod species are broadly distributed throughout Europe, but sexual species are concentrated near the Mediterranean Sea. Horne and Martens (1999) argues that asexual species occur in regions frequently disrupted by climate changes and that sexual species occur in regions with a more stable climate.

Appendix A.5. The importance of spores

Spores may not seem important, but if the ideas presented in the main text and this appendix are true, then spores were critical to the evolution of complexity, diploidy, and sexual reproduction. That is why I think complexity, diploidy, and sexual reproduction are associated with the sporophyte generation in plants, and with ascocarps and basidiocarps in fungi. I want to briefly discuss the importance of spores in the evolutionary history of life.

The importance of spores is immediately obvious in the evolution of syngamy, a critical step needed for sexual reproduction. As discussed in the beginning of this appendix, many organisms undergo syngamy prior to spore production. This allows the spores to contain a mixed cytoplasm, which could lower protein

aggregation rates within the spore. Lowering the rate at which soluble oligomers form is beneficial because it would increase the longevity of the spore and increase the spore's resistance to physical stresses. Thus, syngamy may have evolved because it allowed for the creation of hardier spores.

Spores are great for dispersing offspring over a wide area. In the case of land plants and fungi, spore dispersal is aided by structures that rise above the ground, such as sporophytes, ascocarps, and basidiocarps. These structures must be diploid otherwise the spores they produce would not contain mixed cytoplasm contents. Hence, diploidy is associated with spore dispersing stages in the plant and fungal life-cycles.

It seems inevitable that some spore dispersing life-stages would become larger and more complex over time in order to more effectively scatter their spores. However, I think there is an additional reason why complexity is associated with spore dispersing structures. As discussed in Section 4.5 of the main text, there is a problem with the transition from haploid-dominant life-cycles to diploid-dominant life-cycles. I think spore dispersing life-stages helped to overcome this problem.

To recap, truncation selection would prevent a strictly haploid species from producing an abundance of aggregation-prone proteins. Otherwise, such a species would sit at the precipice of a rapid fall in fitness, and probable extinction, should the environment deteriorate. Diploid individuals that might arise in the species would not have any competitive advantage because protein aggregation would not be a significant problem, and there could be advantages to being haploid (Section 4.6 of main text). Thus, the evolution of complexity contains a roadblock. Strictly haploid species will not become more complex over time because they cannot produce the aggregation-prone proteins that facilitate complexity, and no diploid species exist because diploidy does not confer any immediate competitive advantage.

However, the previously discussed spore dispersing structures provide a path around the roadblock. The spore dispersing structures are diploid because their offspring benefit from possessing mixed cytoplasm contents, not because the structures themselves benefit from diploidy. This provides enough scope for the spore dispersing structures to evolve the aggregation-prone proteins that facilitate complexity. Hence, some spore dispersing structures have become more complex over time.

I think the roadblock in the evolution of complexity can potentially explain why complexity is often

associated with spore dispersing structures. I also think the roadblock can explain why haploid dominant species do not seem to give rise to diploid dominant species directly. Instead, diploid-dominant species seem to typically evolve from species that alternate between haploid and diploid generations, which in turn evolved from haploid-dominant species. This contrasts with the evolution of polyploidy, which occurs in a single generation when diploid parents produce polyploid offspring. The key difference is that the evolution of diploidy is tied to the evolution of complexity, whereas polyploids are typically descended from species that are already complex.

Thus, spores played an important role in the evolutionary history of life. They helped life find a path around the roadblock to the evolution of complexity, and they were the first structures to benefit from diploidy, which favored the evolution of syngamy, a necessary step in the evolution of sexual reproduction. I think this illustrates how complexity, diploidy, sexual reproduction, and spore dispersal are all interconnected.

Appendix A.6. Shifts in truncation selection

Selection for heterozygosity would definitely fall within the parameter space for polygenic adaptation given in Höllinger et al. (2019). An individual organism's heterozygosity is controlled by thousands of gene loci, and balancing selection maintains multiple alleles at many of the gene loci, at least if the ideas presented in the main text are true. Höllinger et al. (2019) even assumed diminishing returns epistasis for fitness in their models. The truncation selection model presented in the main text may be taken as an extreme form of diminishing returns epistasis. Thus, an individual organism's heterozygosity is a prime candidate for polygenic adaptation via allele frequency shifts.

Some numbers may provide a useful example. Suppose that a population of organisms contains two kinds of alleles at each gene locus subjected to balancing selection. One allele, the major allele, has an average frequency of $p = 0.9$. The average frequency of a collection of minor alleles is $q = 0.1$. Assuming Hardy-Weinberg equilibrium, then the average individual heterozygosity will be $H = 0.18$. Now suppose that the average individual is heterozygous at 1000 gene loci. This must mean that $1000/0.18 = 5555$ gene loci are subjected to balancing selection. Individual heterozygosity follows a binomial probability distribution because an organism is either heterozygous at a gene locus or it is not. Hence,

the standard deviation would be $\sqrt{((5555)(0.18)(1 - 0.18))} = 29$. If the average allele frequencies were to shift to $p = 0.88$ for the major alleles and $q = 0.12$ for the minor alleles, then the average heterozygosity would shift to $H = 0.21$, and the average individual would be heterozygous at 1167 gene loci. Thus, very minor allele frequency shifts at many gene loci would cause the average number of heterozygous gene loci, per individual, to move over five standard deviations.

Changes in the average number of heterozygous gene loci per individual could represent an important mode of adaptation for complex organisms. According to the main text, selection for heterozygosity follows a truncation curve in which an individual's fitness dramatically declines below a critical heterozygosity. However, the truncation selection curve could move to either the left or the right, depending on the stressfulness of the environment, as shown in Figure 5b. When the environment has a high level of stress that promotes protein aggregation (the value of k_1 in Figure 5b is high), then only individuals that are heterozygous at many gene loci will be able to survive in the environment. On the other hand, when the environment is only mildly stressful (the value of k_1 in Figure 5b is low), then individuals with very low heterozygosity levels will be able to survive. Individuals with not heterozygosity at all may be able to survive in very mild environments. Therefore, the individual heterozygosity levels may need to shift when the truncation curve moves to the left and the right.

It is easy to conceive how the shifts in average heterozygosity could occur. Assume that the number of heterozygous gene loci per individual follows a binomial distribution, and that the truncation curve generally stays on the left side of the binomial distribution (otherwise more than half the population would fail to survive each generation). When the truncation curve moves to the right along the binomial distribution, a larger fraction of the population will fail to survive and the average individual heterozygosity for the population will increase. The allele frequencies should shift to accommodate the increase in average individual heterozygosity (see Wills (1978) for an example). On the other hand, only a small fraction of the population will fail to survive when the truncation curve moves to the left, out to the tail-end of the binomial distribution. In this case, a combination of genetic drift and natural selection (assuming that some alleles confer higher fitness than other alleles) would cause allele frequencies to shift in favor of the most common allele, and average individual heterozygosity

will decrease in the population (see Equation 17 in the main text, and Wills (1978) again). Hence, individual heterozygosity may be constantly changing in natural populations due to a moving truncation curve.

References

- Ambur, O. H., Engelstädter, J., Johnsen, P. J., Miller, E. L., and Rozen, D. E. (2016). Steady at the wheel: conservative sex and the benefits of bacterial transformation. *Philosophical Transactions of the Royal Society of London B*, 371:20150528.
- Bale, J. S. (1996). Insect cold hardiness: A matter of life and death. *European Journal of Entomology*, 93:369–382.
- Barrick, J. E. and Lenski, R. E. (2009). Genome-wide mutational diversity in an evolving population of *Escherichia coli*. *Cold Spring Harbor Symposia on Quantitative Biology*, 74:119–129.
- Barrière, A. and Félix, M.-A. (2007). Temporal dynamics and linkage disequilibrium in natural *Caenorhabditis elegans* populations. *Genetics*, 176:999–1011.
- Barroso-Batista, J., Sousa, A., Lourenço, M., Bergman, M.-L., Sobral, D., Demengeot, J., Xavier, K. B., and Gordo, I. (2014). The first steps of adaptation of *Escherichia coli* to the gut are dominated by soft sweeps. *PLoS Genetics*, 10:e1004182.
- Bennett, R. J. and Johnson, A. D. (2003). Completion of a parasexual cycle in *Candida albicans* by induced chromosome loss in tetraploid strains. *EMBO*, 22:2505–2515.
- Berg, J. J. and Coop, G. (2014). A population genetic signal of polygenic adaptation. *PLoS Genetics*, 10:e1004412.
- Berg, J. J., Harpak, A., Sinnott-Armstrong, N., Jørgensen, A. M., Mostafavi, H., Field, Y., Boyle, E. A., Zhang, X., Racimo, F., Pritchard, J. K., and Coop, G. (2019). Reduced signal for polygenic adaptation of height in UK Biobank. *eLife*, 8:e39725.
- Bergey, C. M., Lopez, M., Harrison, G. F., Patin, E., Cohen, J., Quintana-Murci, L., Barreiro, L. B., and Perry, G. H. (2018). Polygenic adaptation and convergent evolution on growth and cardiac genetic pathways in African and Asian rainforest hunter-gatherers. *Proceedings of the National Academy of Sciences of the USA*, 115:E11256–E11263.
- Bernstein, H., Bernstein, C., and Michod, R. E. (2011). Meiosis as an evolutionary adaptation for DNA repair. In Kruman, I., editor, *DNA Repair*, pages 357–382. IntechOpen Limited, London, UK.
- Bernstein, H., Hopf, F. A., and Michod, R. E. (1989). The evolution of sex: DNA repair hypothesis. In Rasa, C. V. and Voland, E., editors, *Sociobiology of Reproductive Strategies in Animals and Man*, pages 3–18. Chapman and Hall, London, UK.
- Birky, Jr, C. W. (1996). Heterozygosity, heteromorphy, and phylogenetic trees in asexual eukaryotes. *Genetics*, 144:427–437.
- Boyle, E. A., Li, Y. I., and Pritchard, J. K. (2017). An expanded view of complex traits: From polygenic to omnigenic. *Cell*, 169:1177–1186.
- Caprioli, M. and Ricci, C. (2001). Recipes for successful anhydrobiosis in bdelloid rotifers. *Hydrobiologia*, 446/447:13–17.
- Chernova, N. M., Potapov, M. B., Savenkova, Y. Y., and Bokova, A. I. (2010). Ecological significance of parthenogenesis in Collembola. *Entomological Review*, 90:23–38.
- Crowe, J. H., Carpenter, J. F., and Crowe, L. M. (1998). The role of vitrification in anhydrobiosis. *Annual Reviews in Physiology*, 60:73–103.
- Dijksterhuis, J. (2007). Heat-resistant ascospores. In Dijksterhuis, J. and Samson, R. A., editors, *Food Mycology: A multifaceted Approach to Fungi and Food*, pages 101–117. CRC Press, Boca Raton, FL.
- Dijksterhuis, J., Nijse, J., Hoekstra, F. A., and Golovina, E. A. (2007). High viscosity and anisotropy characterize the cytoplasm of fungal dormant stress-resistant spores. *Eukaryotic Cell*, 6:157–170.
- Domes, K., Norton, R. A., Maraun, M., and Scheu, S. (2007). Reeolution of sexuality breaks Dollos law. *Proceedings of the National Academy of Sciences of the USA*, 104:7139–7144.
- Erkut, C., Penkov, S., Fahmy, K., and Kurzchalia, T. V. (2012). How worms survive desiccation. *Worm*, 1:61–65.
- Erkut, C., Vasilij, A., Boland, S., Habermann, B., Shevchenko, A., and Kurzchalia, T. V. (2013). Molecular strategies of the *Caenorhabditis elegans* dauer larva to survive extreme desiccation. *PLoS One*, 8:e82473.
- Fledel-Alon, A., Wilson, D. J., Broman, K., Wen, X., Ober, C., Coop, G., and Przeworski, M. (2009). Broad-scale recombination patterns underlying proper disjunction in humans. *PLoS Genetics*, 5:e1000658.
- Forche, A., Alby, K., Schaefer, D., Johnson, A. D., Berman, J., and Bennett, R. J. (2008). The parasexual cycle in *Candida albicans* provides an alternative pathway to meiosis for the formation of recombinant strains. *PLoS Biology*, 6:e110.
- Gnecchi-Ruscione, G. A., Abondio, P., Fantì, S. D., Sarno, S., Sherpa, M. G., Sherpa, P. T., Marinelli, G., Natali, L., Marcello, M. D., Peluzzi, D., Luiselli, D., Pettener, D., and Sazzini, M. (2018). Evidence of polygenic adaptation to high altitude from Tibetan and Sherpa genomes. *Genome Biology and Evolution*, 10:2919–2930.
- Gorelick, R. and Heng, H. H. Q. (2010). Sex reduces genetic variation: A multidisciplinary review. *Evolution*, 65:1088–1098.
- Grishkan, I., Korol, A. B., Nevo, E., and Wasser, S. P. (2003). Ecological stress and sex evolution in soil microfungi. *Proceedings of the Royal Society of London B*, 270:13–18.
- Haag, C. R. and Ebert, D. (2004). A new hypothesis to explain geographic parthenogenesis. *Annales Zoologici Fennici*, 41:539–544.
- Hancock, A. M., Witonsky, D. B., Ehler, E., Alkorta-Aranburu, G., Beall, C., Gebremedhin, A., Sukernik, R., Utermann, G., Pritchard, J., Coop, G., and Rienzo, A. D. (2010). Human adaptations to diet, subsistence, and ecoregion are due to subtle shifts in allele frequency. *Proceedings of the National Academy of Sciences of the USA*, 107:8924–8930.
- Hannah, M. A., Wiese, D., Freund, S., Fiehn, O., Heyer, A. G., and Hinch, D. K. (2006). Natural genetic variation of freezing tolerance in *Arabidopsis*. *Plant Physiology*, 142:98–112.
- Heng, H. H. Q. (2007). Elimination of altered karyotypes by sexual reproduction preserves species identity. *Genome*, 50:517–524.
- Hickey, D. A. and Golding, G. B. (2018). The advantage of recombination when selection is acting at many genetic loci. *Journal of Theoretical Biology*, 442:123–128.
- Hoekstra, F. A., Golovina, E. A., and Buitink, J. (2001). Mechanisms of plant desiccation tolerance. *Trends in Plant Science*, 6:431–438.
- Höllinger, I., Pennings, P., and Hermisson, J. (2019). Polygenic adaptation: From sweeps to subtle frequency shifts. *PLoS Genetics*, 15:e1008035.
- Hong, T. D., Ellis, R. H., and Moore, D. (1997). Development of a model to predict the effect of temperature and moisture on fungal spore longevity. *Annals of Botany*, 79:121–128.
- Horne, D. J. and Martens, K. (1999). Geographical parthenogenesis in european non-marine ostracods: post-glacial invasion or holocene stability? *Hydrobiologia*, 391:1–7.
- Hur, J. H., Doninck, K. V., Mandigo, M. L., and Meselson, M. (2009). Degenerate tetraploidy was established before bdelloid rotifer families diverged. *Molecular Biology and Evolution*, 26:375–383.
- Iranzo, J., Wolf, Y. I., Koonin, E. V., and Sela, I. (2019). Homologous recombination substantially delays sequence but not gene content divergence of prokaryotic populations. *BioRxiv*, page 518852.
- Jaron, K. S., Bast, J., Ranallo-Benavidez, T. R., Robinson-Rechavi, M., and Schwander, T. (2018). Genomic features of asexual

- animals. *BioRxiv*, page 497495.
- Kothe, E. (1996). Tetrapolar fungal mating types: Sexes by the thousands. *FEMS Microbiology Reviews*, 18:65–87.
- Lewontin, R. C. and Hubby, J. L. (1966). A molecular approach to the study of genic heterozygosity in natural populations. II. Amount of variation and degree of heterozygosity in natural populations of *Drosophila pseudoobscura*. *Genetics*, 54:595–609.
- Lundmark, M. and Saura, A. (2006). Asexuality alone does not explain the success of clonal forms in insects with geographical parthenogenesis. *Hereditas*, 143:23–32.
- MacDonald, R. I. (1985). Membrane fusion due to dehydration by polyethylene glycol, dextran, or sucrose. *Biochemistry*, 24:4058–4066.
- Maraun, M., Erdmann, G., Schulz, G., Norton, R. A., Scheu, S., and Domes, K. (2009). Multiple convergent evolution of arboreal life in oribatid mites indicates the primacy of ecology. *Proceedings of the Royal Society of London B*, 276:3219–3227.
- Mauseth, J. D. (2009). *Botany: An Introduction to Plant Biology*. Jones and Bartlett Publishers, Sudbury, MA.
- Moritz, C. (1987). parthenogenesis in the tropical gekkonid lizard, *Nactus arnouxi* (Sauria: Gekkonidae). *Evolution*, 41:1252–1266.
- Neiman, A. M. (2005). Ascospore formation in the yeast *Saccharomyces cerevisiae*. *Microbiology and Molecular Biology Reviews*, 69:565–584.
- Nowell, R. W., Almeida, P., Wilson, C. G., Smith, T. P., Fontaneto, D., Crisp, A., Micklem, G., Tunnacliffe, A., Boschetti, C., and Barraclough, T. G. (2018). Comparative genomics of bdelloid rotifers: Insights from desiccating and nondesiccating species. *PLoS Biology*, 16:e2004830.
- Otto, S. P. and Barton, N. H. (2001). Selection for recombination in small populations. *Evolution*, 55:1921–1931.
- Palmer, S. C. and Norton, R. A. (1992). Genetic diversity in thelytokous oribatid mites (Acari; Acariformes; Desmonomata). *Biochemical Systematics and Ecology*, 20:219–231.
- Pedrazzoli, F., Chrysantzas, I., Dezzani, L., Rosti, V., Vincitorio, M., and Sitar, G. (2011). Cell fusion in tumor progression: the isolation of cell fusion products by physical methods. *Cancer Cell International*, 11:32.
- Pritchard, J. K., Pickrell, J. K., and Coop, G. (2010). The genetics of human adaptation: Hard sweeps, soft sweeps, and polygenic adaptation. *Current Biology*, 20:R208R215.
- Pritchard, J. K. and Rienzo, A. D. (2010). Adaptation – not by sweeps alone. *Nature Reviews Genetics*, 11:665–667.
- Rabeling, C., Gonzales, O., Schultz, T. R., Bacci, Jr, M., Garcia, M. V. B., Verhaagh, M., Ishak, H. D., and Mueller, U. G. (2011). Cryptic sexual populations account for genetic diversity and ecological success in a widely distributed, asexual fungus-growing ant. *Proceedings of the National Academy of Sciences of the USA*, 108:1236612371.
- Radwan, J. and Babik, W. (2012). The genomics of adaptation. *Proceedings of the Royal Society of London B*, 279:5024–5028.
- Reyes-Díaz, M., Ulloa, N., Zúñiga-Feest, A., Guti'érrez, A., Gidekel, M., Alberdi, M., Corcuera, L. J., and Bravo, L. A. (2006). *Arabidopsis thaliana* avoids freezing by supercooling. *Journal of Experimental Botany*, 57:36873696.
- Ricci, C. N. (1987). Ecology of bdelloids: how to be successful. *Hydrobiologia*, 147:117–127.
- Roy, A., Dutta, R., Kundu, N., Banik, D., and Sarkar, N. (2016). A comparative study of the influence of sugars sucrose, trehalose, and maltose on the hydration and diffusion of DMPC lipid bilayer at complete hydration: Investigation of structural and spectroscopic aspect of lipid–sugar interaction. *Langmuir*, 32:5124–5134.
- Roze, D. (2014). Selection for sex in finite populations. *Journal of Evolutionary Biology*, 27:1304–1322.
- Sherwood, R. K. and Bennett, R. J. (2009). Fungal meiosis and parasexual reproduction – lessons from pathogenic yeast. *Current Opinion in Microbiology*, 12:599–607.
- Siepel, H. (1994). Life–history tactics of soil microarthropods. *Biology and Fertility of Soils*, 18:263–278.
- Sinclair, B. J. (1999). Insect cold tolerance: How many kinds of frozen? *European Journal of Entomology*, 96:157–164.
- Sinclair, E. A., Pramuk, J. B., Bezy, R. L., Crandall, K. A., and Sites, Jr, J. W. (2010). DNA evidence for nonhybrid origins of parthenogenesis in natural populations of vertebrates. *Evolution*, 64:1346–1357.
- Smelansky, I. E. (2006). Some population characteristics of oribatid mites in steppe habitats. *Acarina*, 14:123–130.
- Stenøien, H. K., Fenster, C. B., Tonteri, A., and Savolainen, O. (2005). Genetic variability in natural populations of *Arabidopsis thaliana* in northern Europe. *Molecular Ecology*, 14:137–148.
- Teotónio, H., Estes, S., Phillips, P. C., and Baer, C. F. (2017). Experimental evolution with *Caenorhabditis nematodes*. *Genetics*, 206:691–716.
- Tilquin, A. and Kokko, H. (2016). What does the geography of parthenogenesis teach us about sex? *Philosophical Transactions of the Royal Society of London B*, 371:20150538.
- Trapero-Casas, A. and Kaiser, W. J. (2007). Differences between ascospores and conidia of *Didymella rabiei* in spore germination and infection of chickpea. *Phytopathology*, 97:1600–1607.
- Vos, M. (2009). Why do bacteria engage in homologous recombination? *Trends in Microbiology*, 17:226–232.
- Welch, D. B. M., Welch, J. L. M., and Meselson, M. (2008). Evidence for degenerate tetraploidy in bdelloid rotifers. *Proceedings of the National Academy of Sciences of the USA*, 105:5145–5149.
- Wilkins, A. S. and Holliday, R. (2009). The evolution of meiosis from mitosis. *Genetics*, 181:3–12.