# CHERN NUMBERS OF MANIFOLDS WITH TORUS ACTION.

#### ANDREY KUSTAREV

ABSTRACT. We show that every set of numbers that occurs as the set of Chern numbers of an almost complex manifold  $M^n$ ,  $n \geq 6$ , may be realized as the set of Chern numbers of a connected almost complex manifold  $M^n$  with an almost complex action of two-dimensional compact torus.

## 1. INTRODUCTION

There is a well-known problem (posed by Hirzebruch) asking which combinations of Chern numbers of almost complex manifolds (or, equivalently, which complex cobordism classes) may be realized by a smooth connected complex projective variety. As shown by Milnor, any complex cobordism class may be realized by a disjoint union of smooth projective varieties, so the connectivity condition in the formulation of Hirzebruch problem is crucial. As shown in ([\[4\]](#page-4-0)), any complex cobordism class may be realized by a connected smooth almost complex manifold.

Chern numbers of compact almost complex and symplectic manifolds with compact Lie group action have been an object of intensive research in recent years  $([6], [7], [9], [12], [13])$  $([6], [7], [9], [12], [13])$  $([6], [7], [9], [12], [13])$  $([6], [7], [9], [12], [13])$  $([6], [7], [9], [12], [13])$  $([6], [7], [9], [12], [13])$  $([6], [7], [9], [12], [13])$  $([6], [7], [9], [12], [13])$  $([6], [7], [9], [12], [13])$  $([6], [7], [9], [12], [13])$  $([6], [7], [9], [12], [13])$ . Any smooth manifold M with an almost complex action having only isolated fixed points necessarily has  $c_n(M) \geq 0$ ; many other restrictions are shown to exist in this case. In this paper we show that if fixed points are allowed to be non-isolated, then all restrictions on Chern numbers vanish.

<span id="page-0-0"></span>**Theorem 1.1.** Every complex cobordism class of dimension  $n \geq 6$  may be realized by a connected almost complex manifold  $M^n$  with an almost complex action of two-dimensional compact torus.

Moreover, the set of fixed points of the action consists only of isolated fixed points and two-dimensional oriented closed surfaces.

**Definition 1.2.** A non-singular complex projective variety  $M = M^n$ will be called *good* if it is a product of one or several varieties belonging to following classes:

- compact complex curves of genus  $q > 1$ ,
- non-singular compact projective toric varieties  $X^n$ ,
- Milnor hypersurfaces  $H_{i,j}$  with  $4 \leq i \leq j$ ,

such that there is at most one complex curve in the product.

#### 2 ANDREY KUSTAREV

<span id="page-1-0"></span>Lemma 1.3. Any complex cobordism class may be represented by a disjoint union of good varieties.

We will obtain the proof of theorem [1.1](#page-0-0) by applying the construction of fiber connected sum (see lemma [3.2,](#page-3-0) cf. [\[5\]](#page-4-4) for contact structures) after establishing lemma [1.3.](#page-1-0)

### 2. Proof of lemma [1.3](#page-1-0)

Recall that the complex cobordism ring  $\Omega_t^*$  $_{U}^{*}$  is generated by smooth stably complex manifolds modulo the equivalence relation:  $M_1 \sim M_2$ if and only if there exists stably complex manifold  $N$  such that  $\partial N = M_1 \cup (-M_2)$  where  $(-M_2)$  is  $M_2$  with reversed orientation and restriction of stably complex structure of N to  $M_1$  and  $M_2$  is equivalent to stably complex structures on  $M_1$  and  $M_2$  respectively. The operations of taking disjoint union and product of manifolds turn  $\Omega^*_L$  $_{U}^{*}$  into a graded commutative ring; stably complex manifold  $M^{n}$ determines a class  $[M^n] \in \Omega_{U}^{-n}$  $\frac{-n}{U}$ .

As shown by Milnor and Novikov ([\[10\]](#page-5-2), [\[11\]](#page-5-3)), the ring  $\Omega_t^*$  $\mathrm{d}^*$  is isomorphic to the polynomial ring  $\mathbb{Z}[a_1, a_2, \ldots]$ , where deg  $a_i = -2i$ . In particular, the manifold is zero in  $\Omega_t^*$  $_U^*$  if and only if all of its Chern numbers are zero. Moreover, the set of stably complex manifolds  $\{M^{2n}\}, n \geq 1$ , may solve as the set of polynomial generators for  $\Omega_L^*$  $_U^*$  if and only if for any *n* we have  $s_n([M^{2n}]) = \eta(n)$ , where

$$
\eta(n) = p
$$
, if  $n + 1 = p^k$  for some prime  $p$ ,

 $\eta(n) = 1$  otherwise,

and  $s_n$  is a *Milnor number* which is a Chern number corresponding to Newton symmetric polynomial of dimension  $n$  (e.g.  $s_1 = c_1$ ,  $s_2 = c_1^2 - 2c_2$ ,  $s_3 = c_1^3 - 3c_1c_2 + 3c_3$ , etc.).

We proceed with the proof of lemma [1.3](#page-1-0) by induction. Any complex cobordism class of dimension two is represented by a disjoint union of complex curves, since any two-dimensional almost complex manifold is automatically a complex curve.

Suppose that the statement of theorem [1.3](#page-1-0) is true for all complex cobordism classes of dimension  $\leq 2n$ .

<span id="page-1-1"></span>**Lemma 2.1.** Any complex cobordism class  $[M] \in \Omega_U^{-2n}$  satisfying  $s_n([M]) = 0$  may be represented by a good variety.

*Proof.* Since  $s_n([M]) = 0$ , the class [M] is decomposable and we have  $[M] = \sum$ K  $k=1$  $[N_k \times N'_k]$  where  $N_k$  and  $N'_k$  are good varieties. If for some k both  $N_k$  and  $N'_k$  contain complex curves of genus g and g' respectively, we replace these curves with disjoint unions of  $(g - 1)$  and  $(g' - 1)$ copies of  $\mathbb{C}P^1$  correspondingly.  $\square$ 

**Lemma 2.2.** If the class  $[M] \in \Omega_U^{-2n}$  may be represented by a good variety, the same is true for  $(-[M])$ .

Proof. Recall that blowing up a point on a compact complex variety X lowers its Milnor number  $s_n([X])$  by the value  $(n+(-1)^n)$ . This implies that in every dimension  $n > 1$  there exist toric varieties with positive and negative Milnor numbers – namely, the complex projective space  $\mathbb{C}P^n$  and the variety obtained from  $\mathbb{C}P^n$  by blowing up three different fixed points of torus action.

This argument shows that there exists a toric variety  $N$  such that  $s_n(a[M] + b[N]) = 0$ , where  $a, b \in \mathbb{Z}_{>0}$ . By lemma [2.1,](#page-1-1) the class  $(-a[M] - b[N])$  may be represented by a disjoint union of good varieties. Then the same is true for the class  $-[M] = (-a[M] - b[N]) + b[N] + (a-1)[M]$ .

**Lemma 2.3.** For every  $n \ge 1$  there exists a good variety  $G_n$  (of complex dimension n) with  $s_n([G_n]) = \pm \eta(n)$ .

Proof. We will use the following result on toric generators in complex cobordism.

<span id="page-2-0"></span>**Theorem 2.4.** [\[14\]](#page-5-4) For every odd n and every even  $n \le 100001$  there exists a compact non-singular toric variety  $G_n$  (of complex dimension n) satisfying  $s_n([G_n]) = \pm \eta(n)$ .

Therefore, it is enough to show that for every even  $n > 100001$  the complex cobordism generator  $a_n \in \Omega_U^{-2n}$  may be realized by a disjoint union of good varieties.

Recall that we have  $s_n(\mathbb{C}P^n) = n+1$  and  $s_n(H_{i,j}) = -\binom{i+j}{i}$  $\binom{+j}{i}$ for  $2 \leq i \leq j$ . By Kummer theorem, the maximum degree k for which  $p^k$  divides  $\binom{i+j}{i}$  $\binom{+j}{i}$  is equal to the number of carries when  $i$  is added to  $j$  in base  $p$ . We need to show that  $\gcd\left(\frac{(n+1)}{i},\binom{i+j}{i}\right)$  $\{i+j\}$  |  $i+j = n+1, 2 \leqslant i \leqslant j$   $= \eta(n)$  for every even  $n$  large enough.

Suppose first that  $n + 1 = p^k$  and  $\eta(n) = p$  for some prime p. Using [2.4,](#page-2-0) we may only consider the case  $p > 2$ . If  $k = 1$ , then  $s_n([\mathbb{C}P^n]) = n+1 = p = \eta(n)$ , so we may also assume that  $k > 1$ . Adding up numbers  $p^{k-1}$  and  $p^k - p^{k-1}$  in base p has only one carry and by Kummer theorem we see that  $\binom{p^k}{p^k}$  $p_{p^{k-1}}^{p^{k}}$  is divisible by p and not by  $p^2$ . So the statement of the lemma is true unless  $p^{k-1} \leq 3$ . But in this case  $p = 3$  and  $k = 2$ , so  $n = 8$  and toric generator  $G_8$  exists by [2.4.](#page-2-0)

Next, let us assume that  $(n+1)$  is not a degree of a prime p. If p does not divide  $(n+1)$ , then there is nothing to prove since  $s_n([\mathbb{C}P^n]) = n+1$ . The representation of  $(n+1)$  in base p has at least two non-zero digits. Let  $l > 0$  be the maximum number such that  $p^l < (n+1)$ . Then  $\binom{n+1}{n^l}$  $p^{l-1 \choose p^l}$ does not divide by p. Recalling that p divides  $(n + 1)$ , we see that if

#### 4 ANDREY KUSTAREV

 $(n+1) - p<sup>l</sup> \leq 3$ , then either  $n+1 = 2<sup>l</sup> + 2$  or  $n+1 = 3<sup>l</sup> + 3$ , so n is odd and we may again apply [2.4.](#page-2-0)  $\Box$ 

### 3. Fiber connected sums of almost complex manifolds

In this section we apply lemma [1.3](#page-1-0) to prove the main result, theorem [1.1.](#page-0-0)

<span id="page-3-1"></span>Lemma 3.1. Any disjoint union of good varieties admits a faithful action of compact torus  $T^{\min(4,n-1)}$  with the set of fixed points being a disjoint union of nonsingular curves and isolated fixed points.

*Proof.* Recall that Milnor hypersurface  $H_{i,j} \subset \mathbb{C}P^i \times \mathbb{C}P^j$ ,  $i \leq j$ , is given by the equation  $\{z_0w_0 + \ldots + z_iw_i = 0\}$  and has the natural faithful action of compact torus  $T<sup>i</sup>$  given by the formula

$$
(t_1,\ldots,t_i)\cdot (z_0,w_0,\ldots,z_i,w_i)=(z_0,w_0,t_1z_1,t_1^{-1}w_1,\ldots,t_iz_i,t_i^{-1}w_i).
$$

The action is faithful and has only isolated fixed points. Furthermore, every complex projective toric variety  $X<sup>n</sup>$  has a natural faithful action of compact torus  $T^n$  and any curve of genus  $g > 1$  has a trivial torus action.

Taking products of torus actions, we see that every good variety  $X^n$ possesses a faithful holomorphic action of a torus of dimension equal to at least min $(4, n-1)$ .  $\Box$ 

Let  $X_1 = X_1^{2n}$  and  $X_2 = X_2^{2n}$  be two smooth manifolds with a smooth faithful action of torus  $T^k$ . Any principal orbit of the action has an equivariant tubilar neighborhood diffeomorphic to  $T^k \times D^{2n-k}$ , where  $D^{2n-k}$  is an open unit ball in  $\mathbb{R}^{2n-k}$  centered at the origin. Let us choose two principal orbits  $O_1 \subset X_1$  and  $O_2 \subset X_2$  equipped with equivariant tubilar neigborhoods  $U_1$  and  $U_2$  respectively. One can now construct a *fiber connected sum*  $Y = X_1 \#_{T^k} X_2$  by gluing together diffeomorphic parts  $U_1 - O_1$  and  $U_2 - O_2$  of open manifolds  $X_1 - O_1$ and  $X_2 - O_2$  by the diffeomorphism  $(t, r, \varphi) \sim (t, \frac{1}{r}, \varphi)$ , where  $t \in T^k$ ,  $(r, \varphi) \in D^{2n-k}, r > 0, \varphi \in S^{2n-k-1}.$ 

There are smooth embeddings  $(X_1 - U_1) \hookrightarrow Y$  and  $(X_2 - U_2) \hookrightarrow Y$ . If  $X_1$  and  $X_2$  are almost complex manifolds, there is a natural question whether we can extend a  $T^k$ -invariant almost complex structure from a disjoint union of manifolds  $(X_1 - U_1) \cup (X_2 - U_2)$  to the entire manifold Y.

<span id="page-3-0"></span>**Lemma 3.2.** Suppose that  $X_1 = X_1^{2n}$  and  $X_2 = X_2^{2n}$  are compact manifolds with a smooth faithful almost complex action of compact torus  $T^k$ . If the stable homotopy group  $\pi_{2n-k-1}(SO/U)$  is trivial, then the invariant almost complex structure can be extended from a disjoint union  $(X_1 - U_1) \cup (X_2 - U_2)$  to an invariant almost complex structure on Y. Moreover,  $[Y] = [X_1] + [X_2]$  in  $\Omega_t^*$  $_U^*$ .

*Proof.* The space  $U_1 - O_1 \simeq U_2 - O_2$  is a trivial fibration over the torus  $T^k$ . We denote by  $A_t$  its fiber (an open annulus) over the point  $t \in T^k$ . Let us fix some point  $t_0 \in T^k$ . Suppose that we managed to extend somehow the almost complex structure operator  $J$ from  $(X_1 - U_1) \cup (X_2 - U_2)$  to the restriction of the tangent bundle  $\tau_{\mathbb{R}}(Y)|_{A_{t_0}}$ . Then this new structure may be automatically extended by the action of torus  $T^k$  to an invariant structure on the space  $U_1 - O_1 \simeq U_2 - O_2$ , and therefore, on the entire space Y.

The obstruction to extending  $J$  to  $\tau_{\mathbb{R}}(Y)|_{A_{t_0}}$  lies in a homotopy group  $\pi_{2n-k-1}(SO(2n)/U(n))$ . Since  $k > 0$ , this group is equal to the stable homotopy group  $\pi_{2n-k-1}(SO/U)$ .

The equality  $[Y] = [X_1] + [X_2]$  now follows from the localization theorem ([\[1\]](#page-4-5), [\[2\]](#page-4-6), [\[8\]](#page-4-7)), which implies that Chern numbers of a manifold with an almost complex torus action are determined by the behaviour near fixed points.  $\square$ 

Proof of the theorem [1.1.](#page-0-0) By Bott periodicity, we have  $\pi_{i-1}(SO/U) = \pi_i(O)$  and the stable homotopy group  $\pi_i(O)$  is trivial if and only if  $j \equiv 2, 4, 5, 6 \mod 8$ .

Consider the class  $[M^n] \in \Omega_U^{-2n}, n \geq 3$ . By lemma [1.3](#page-1-0) it can be realized by a disjoint union of good varieties and by lemma [3.1](#page-3-1) every of these varieties admits an action of torus  $T^l$ , where  $l \geq \min(4, n-1)$ . Let  $k = 4$ , if  $n \equiv 1 \pmod{4}$ , and  $k = 2$  otherwise. We see that

- $k \leq l$  and therefore every good variety admits a faithful almost complex action of torus  $T^k$ ;
- the stable homotopy group  $\pi_{2n-k-1}(SO/U) = \pi_{2n-k}(O)$  is trivial.

Applying lemma [3.2](#page-3-0) finishes the proof of [1.1.](#page-0-0)  $\Box$ 

### **REFERENCES**

- <span id="page-4-5"></span>[1] M. Atiyah and R. Bott, The moment map and equivariant cohomology, Topology, 23, (1984), 1–28.
- <span id="page-4-6"></span>[2] V. M. Buchstaber, T. E. Panov and N. Ray, Toric genera, Internat. Math. Research Notices 16, (2010), 3207–3262.
- [3] K.E.Feldman. Hirzebruch genera of manifolds equipped with a Hamiltonian circle action. [arXiv:math/0110028v](http://arxiv.org/abs/math/0110028)2
- <span id="page-4-0"></span>[4] H. Geiges, Chern numbers of almost complex manifolds, Proc. AMS, Volume 129, Number 12, Pages 3749-3752.
- <span id="page-4-4"></span>[5] H. Geiges, An introduction to contact topology, Cambridge studies in advanced mathematics 109, Cambridge University Press (2008).
- <span id="page-4-1"></span>[6] L. Godinho, A. Pelayo, S. Sabatini, Fermat and the number of fixed points of periodic flows, [arXiv:1404.4541](http://arxiv.org/abs/1404.4541) [math.AT]
- <span id="page-4-2"></span>[7] L. Godinho, S. Sabatini, New tools for classifying Hamiltonian circle actions with isolated fixed points, Foundations of Comp. Math., 14(4), 791-860 (2014).
- <span id="page-4-7"></span>[8] I. M. Krichever, Formal groups and the Atiyah-Hirzebruch formula, Mathematics of the USSR-Izvestiya, no. 6, (1974), 1271–1285.
- <span id="page-4-3"></span>[9] P. Li, K. Liu, Some remarks on circle action on manifolds, Math. Res. Letters, 18, (2011), 3, pp. 437–446.

### 6 ANDREY KUSTAREV

- <span id="page-5-2"></span>[10] J. Milnor. On the cobordism ring  $\Omega^*$  and a complex analogue I, Amer. J. Math. 82 (1960).
- <span id="page-5-3"></span>[11] S. P. Novikov, Methods of algebraic topology from the point of view of cobordism theory, Izv. Akad. Nauk SSSR, Ser. Mat. 31 (1967), no. 4, 855–951 (Russian); Math. USSR – Izv. 1 (1967), no. 4, 827–913 (English translation).
- <span id="page-5-0"></span>[12] A. Pelayo, S. Tolman, *Fixed points of symplectic periodic flows*, *Erg.* Theory and Dyn. Syst. 31, (2011), 1237–1247.
- <span id="page-5-1"></span>[13] S. Tolman, J. Weitsman, On semi-free symplectic circle actions with isolated fixed points, Topology, 39, (2000), no. 2, 299–309.
- <span id="page-5-4"></span>[14] A. Wilfong, Toric polynomial generators of complex cobordism, [arXiv:1308.2010.](http://arxiv.org/abs/1308.2010)