

Exponentiating Higgs

Marco Matone

Dipartimento di Fisica e Astronomia “G. Galilei”
Istituto Nazionale di Fisica Nucleare
Università di Padova, Via Marzolo, 8-35131 Padova, Italy

Abstract

The scalar models with exponential interaction, introduced in arXiv:1506.00987, include scalar theories with nontrivial $\langle\phi(x)\rangle \neq 0$. Here we consider the theory obtained by normal ordering the exponential of the potential $V(\phi) = \mu^D \exp(-\alpha\phi)$, rather than of $V(\phi)$ itself. This leads to the partition function

$$W_R[J] = {}_J\langle 0 | : e^{-\mu^D \int d^D x \exp(-\alpha\phi(x))} : | 0 \rangle_J$$

that corresponds to fill-in the vacuum of the free scalar theory coupled to the external current with the scalar modes. It turns out that

$$\langle\phi(x)\rangle = \alpha\mu^D/m^2$$

The N point functions follow straightforwardly from the formulation. The fact that the purely scalar sector has an exact nontrivial vev, suggests a possible natural way to get the Lagrangian of the Standard Model, without the cubic and quartic terms of the Higgs field, that may be tested in future experiments at LHC.

1 Introduction

The Higgs mechanism [1]-[6] is a basic step in the formulation of the Standard Model [7, 8]. This has been confirmed by the spectacular experimental results at LHC [9, 10]. Despite this, there are still some open questions. The most important one is that the vev of the Higgs field is evaluated at the classical level. On the other other hand, there are models with non-trivial minima for the classical potential, having a trivial $\langle\phi(x)\rangle$. The point is that the SSB is a strictly nonperturbative phenomenon, concerning infinitely many degrees of freedom. As such, even radiative corrections to $\langle\phi(x)\rangle$ should be considered with particular attention. In this respect, one should also recall that, against the evidence coming from the perturbative expansion, $\lambda\phi^4$ is believed to be a free theory. A related aspect in the standard formulation of the Higgs mechanism, is that the initial lagrangian has the opposite sign for the mass term.

The obvious way to escape the above questions, is to find an alternative way to get a nontrivial vev for the scalar field. Recently, in [11], it has been shown that there are scalar theories with exponential interaction leading to a nontrivial $\langle\phi(x)\rangle$. An outcome of [11], is that there appear terms coming from the normal ordering of the exponentiated scalar. Here we show that this indicates a natural way to regularize the theory, by considering the normal ordering prescription of the interaction term, seen as a field insertion in the free path integral. More precisely, the D -dimensional euclidean partition function of the purely scalar sector is

$$W_R[J] = e^{-Z_R[J]} = {}_J\langle 0 | : e^{-\mu^D \int d^D x \exp(-\alpha\phi(x))} : | 0 \rangle_J . \quad (1.1)$$

This can be seen as filling-in by scalar modes the free vacuum in an external field. The results in [11] straightforwardly imply that

$$Z_R[J] = Z_0[J] + \mu^D \int d^D x e^{-\alpha \int d^D y J(y) \Delta_F(y-x)} , \quad (1.2)$$

which, in turn, implies the exact result

$$\langle 0 | \phi(x) | 0 \rangle = \frac{\alpha \mu^D}{m^2} . \quad (1.3)$$

Furthermore, we will show that the N -point function, $N > 2$, is

$$\begin{aligned} & \langle 0 | T \eta(x_1) \dots \eta(x_N) | 0 \rangle \\ &= (-1)^N \frac{\delta^N Z_0[J]}{\delta J(x_1) \dots \delta J(x_N)} \Big|_{J=0} + \alpha^N \int d^D y \Delta_F(y-x_1) \dots \Delta_F(y-x_N) , \end{aligned} \quad (1.4)$$

where

$$\eta(x) = \phi(x) - \frac{\alpha \mu^D}{m^2} . \quad (1.5)$$

In the present investigation we consider the usual lagrangian density of the Standard Model, except for the scalar lagrangian density, which is

$$\mathcal{L}_\Phi = (D_\mu \Phi)^\dagger (D_\mu \Phi) - m^2 \Phi^\dagger \Phi - \mu^4 e^{-2\alpha(\Phi^\dagger \Phi)^{\frac{1}{2}}} . \quad (1.6)$$

Parameterizing the scalar doublet in polar coordinates and absorbing the phase by a gauge transformation, we are left with

$$\Phi(x) = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ \phi(x) \end{pmatrix} . \quad (1.7)$$

We then focus on the purely scalar part of the theory, described by the lagrangian density

$$\mathcal{L}_\phi = \partial_\mu \phi \partial^\mu \phi - \frac{1}{2} m^2 \phi^2 - \mu^4 e^{-\alpha \phi} . \quad (1.8)$$

The key points of the present investigation are the use of the formalism introduced in [11], together with the prescription of considering the exponential interaction in (1.8) as the insertion in the free theory of the normal ordered term in (1.1).

2 The Model

Let us start by introducing the notation, which follows the one in Ramond's book [12], and shortly reviewing the investigation in [11]. We consider the partition function in the D -dimensional euclidean space

$$W[J] = e^{-Z[J]} = N \int D\phi \exp \left[- \int d^D x \left(\frac{1}{2} \partial_\mu \phi \partial_\mu \phi + \frac{1}{2} m^2 \phi^2 + V(\phi) - J\phi \right) \right] . \quad (2.1)$$

Define

$$\langle f(x_1, \dots, x_n) \rangle_{x_j \dots x_k} \equiv \int d^D x_j \dots d^D x_k f(x_1, \dots, x_n) , \quad (2.2)$$

and denote by $\langle f(x_1, \dots, x_n) \rangle$ integration of f over x_1, \dots, x_n . Let

$$\Delta_F(x-y) = \int \frac{d^D p}{(2\pi)^D} \frac{e^{ip(x-y)}}{p^2 + m^2} , \quad (2.3)$$

be the Feynman propagator and set

$$Z_0[J] = -\frac{1}{2} \langle J(x) \Delta_F(x-y) J(y) \rangle . \quad (2.4)$$

We use the Schwinger trick to compute $W[J]$, namely

$$W[J] = N e^{-\langle V(\frac{\delta}{\delta J}) \rangle} e^{-Z_0[J]} . \quad (2.5)$$

The starting point in [11] has been the observation that Schwinger's trick can be extended to get exact results. In particular, it has been observed that exponential interactions can

be obtained by acting on $\exp(-Z_0[J])$ with the translation operator. We will follow [11] except for the sign of α . Namely, we will consider the potential

$$V(\phi) = \mu^D e^{-\alpha\phi} . \quad (2.6)$$

Dropping the constant N in the expression of $W[J]$

$$\begin{aligned} W[J] &= \exp \left[-\mu^D \langle \exp(-\alpha \frac{\delta}{\delta J}) \rangle \right] \exp(-Z_0[J]) \\ &= \sum_{k=0}^{\infty} \frac{(-1)^k}{k!} \mu^{kD} \langle \exp(-\alpha \frac{\delta}{\delta J}) \rangle^k \exp(-Z_0[J]) . \end{aligned} \quad (2.7)$$

We have

$$\exp(-\alpha \frac{\delta}{\delta J(x)}) \exp(-Z_0[J]) = \exp(-Z_0[J - \alpha_x]) \exp(-\alpha \frac{\delta}{\delta J(x)}) = \exp(-Z_0[-\alpha_x]) , \quad (2.8)$$

where

$$\begin{aligned} Z_0[J - \alpha_x] &:= -\frac{1}{2} \int d^D y d^D z (J(y) - \alpha \delta(x-y)) \Delta_F(y-z) (J(z) - \alpha \delta(x-z)) \\ &= Z_0[J] - \frac{\alpha^2}{2} \Delta_F(0) + \alpha \int d^D y J(y) \Delta_F(y-x) . \end{aligned} \quad (2.9)$$

Therefore,

$$\begin{aligned} W[J] &= \exp(-Z_0[J]) \sum_{k=0}^{\infty} \left[\frac{(-\mu^D)^k}{k!} \exp \left(\frac{k\alpha^2}{2} \Delta_F(0) \right) \right. \\ &\quad \left. \int d^D z_1 \dots \int d^D z_k \exp \left(-\alpha \int d^D z J(z) \sum_{j=1}^k \Delta_F(z-z_j) + \alpha^2 \sum_{j>l}^k \Delta_F(z_j-z_l) \right) \right] \end{aligned} \quad (2.10)$$

The terms $\exp(\frac{k\alpha^2}{2} \Delta_F(0))$ and $\exp(\alpha^2 \sum_{j>l}^k \Delta_F(z_j-z_l))$ are related to normal ordering. To see this, note that the expansion (2.7) corresponds to the expansion of the potential, namely

$$\exp \left(\int d^D x V(x) \right) = \sum_{k=0}^{\infty} \frac{1}{k!} \left(\int d^D x V(x) \right)^k . \quad (2.11)$$

Let us consider again the expansion (2.7). This corresponds to

$$\begin{aligned} W[J] &= {}_J \langle 0 | T e^{-\mu^D \int d^D x \exp(-\alpha\phi(x))} | 0 \rangle_J \\ &= \sum_{k=0}^{\infty} \frac{(-1)^k}{k!} \mu^{kD} \int d^D x_1 \dots d^D x_k J \langle 0 | T e^{-\alpha\phi(x_1)} \dots e^{-\alpha\phi(x_k)} | 0 \rangle_J , \end{aligned} \quad (2.12)$$

where the vacua are the ones of the free scalar theory coupled to the external source J . The fact that the normal ordering problem is the cause of some of the infinities arising in

perturbation theory, suggests that the natural way to absorb them is to replace (2.12) by

$$\begin{aligned} W_R[J] &= {}_J\langle 0 | : e^{-\mu^D \int d^D x \exp(-\alpha\phi(x))} : |0\rangle_J \\ &= \sum_{k=0}^{\infty} \frac{(-1)^k}{k!} \mu^{kD} \int d^D x_1 \dots d^D x_k {}_J\langle 0 | : e^{-\alpha\phi(x_1)} \dots e^{-\alpha\phi(x_k)} : |0\rangle_J . \end{aligned} \quad (2.13)$$

To see this, note that

$$: e^{-\alpha\phi(x)} : := e^{-\frac{\alpha^2}{2} \Delta_F(0)} e^{-\alpha\phi(x)} , \quad (2.14)$$

and

$$T : e^{-\alpha\phi(x_1)} : \dots : e^{-\alpha\phi(x_k)} : := e^{\alpha^2 \sum_{j>l}^k \Delta_F(x_j - x_k)} : e^{-\alpha\phi(x_1)} \dots e^{-\alpha\phi(x_k)} : . \quad (2.15)$$

It follows that,

$$: e^{-\alpha\phi(x_1)} \dots e^{-\alpha\phi(x_k)} : := e^{-\alpha^2 \left(\frac{1}{2} \Delta_F(0) + \sum_{j>l}^k \Delta_F(x_j - x_l) \right)} T e^{-\alpha\phi(x_1)} \dots e^{-\alpha\phi(x_k)} . \quad (2.16)$$

Considering the normal ordering of $e^{-\mu^D \int d^D x \exp(-\alpha\phi(x))}$ has the nice consequence that now the expansion on right hand side in (2.13) exponentiates. Actually, (2.10), (2.12), (2.13) and (2.15) yield

$$W_R[J] = \exp(-Z_R[J]) , \quad (2.17)$$

where

$$Z_R[J] = Z_0[J] + \mu^D \int d^D x e^{-\alpha \int d^D y J(y) \Delta_F(y-x)} . \quad (2.18)$$

Note that

$$W_R[0] = 1 . \quad (2.19)$$

Interestingly, the removal of the term $\exp(\alpha^2 \sum_{j>l}^k \Delta_F(x_j - x_l))$, coming from the normal ordering, is equivalent to remove a term $\langle \exp(-\alpha \frac{\delta}{\delta J}) \rangle$ in (2.7). To show this, recall that for any suitable function F , if A and B are operators, then $A^{-1}F(B)A = F(A^{-1}BA)$. Therefore,

$$\begin{aligned} W[J] &= \exp \left[-\mu^D \langle \exp(-\alpha \frac{\delta}{\delta J}) \rangle \right] \exp(-Z_0[J]) \\ &= \exp(-Z_0[J]) \exp \left[-\mu^D \exp(Z_0[J]) \langle \exp(-\alpha \frac{\delta}{\delta J}) \rangle \exp(-Z_0[J]) \right] \\ &= \exp(-Z_0[J]) \exp \left[-\mu_0^D \langle \exp(-\alpha \langle J(y) \Delta_F(x-y) \rangle_y) \rangle_x \langle \exp(-\alpha \frac{\delta}{\delta J}) \rangle \right] \end{aligned} \quad (2.20)$$

where

$$\mu^D = \mu_0^D \exp \left(-\frac{\alpha^2}{2} \Delta_F(0) \right) . \quad (2.21)$$

Eq.(2.20) differs from $W_R[J]$ by the term $\langle \exp(-\alpha \frac{\delta}{\delta J}) \rangle$ in the last member, and by the relabeling of μ_0 .

A similar analysis can be extended to the case of more general potentials, such as

$$V(\phi) = \sum_{k=1}^N \mu_k^D \exp(\alpha_k \phi) . \quad (2.22)$$

3 Nonperturbative evaluation of $\langle \phi(x) \rangle$

Consider the field

$$\phi_{\text{cl}}(x) := {}_J \langle 0 | \phi(x) | 0 \rangle_J = -\frac{\delta Z_R[J]}{\delta J(x)} , \quad (3.1)$$

and note that by (2.18)

$$\phi_{\text{cl}}(x) = \langle J(y) \Delta_F(x-y) \rangle_y + \alpha \mu^D \langle \Delta_F(y-x) \exp(-\alpha \langle J(z) \Delta_F(y-z) \rangle_z) \rangle_y , \quad (3.2)$$

that satisfies the equation of motion

$$(-\partial_\mu \partial_\mu + m^2) \phi_{\text{cl}}(x) = J(x) + \alpha \mu^D \exp(-\alpha \langle J(y) \Delta_F(x-y) \rangle_y) . \quad (3.3)$$

By (3.2) the effective action

$$\Gamma_R[\phi_{\text{cl}}] = Z_R[J] - \langle J(x) \phi_{\text{cl}}(x) \rangle_x , \quad (3.4)$$

reads

$$\Gamma_R[\phi_{\text{cl}}] = Z_R[J] - \langle J(x) \Delta_F(x-y) J(y) \rangle_{xy} - \alpha \mu^D \langle J(x) \Delta_F(x-y) \exp(-\alpha \langle J(z) \Delta_F(z-y) \rangle_z) \rangle_{xy} \quad (3.5)$$

Note that the theory provides an exact simple expression for $\langle \phi(x) \rangle$. Actually, by (3.2) it follows

$$\langle 0 | \phi(x) | 0 \rangle = \frac{\alpha \mu^D}{m^2} , \quad (3.6)$$

where we used

$$\langle \Delta_F(x-y) \rangle_y = \frac{1}{m^2} . \quad (3.7)$$

Such a relation follows, for example, by

$$\int d^D x (-\partial_\mu \partial_\mu + m^2) \Delta_F(x-y) = 1 , \quad (3.8)$$

and then using the translation invariance of the Feynman propagator to check that there are no additional terms in (3.7), solutions of the homogeneous Klein-Gordon equation. It follows that the higher derivatives of $Z_R[J]$, evaluated at $J=0$, are the connected Green functions associated to

$$\eta(x) = \phi(x) - \frac{\alpha \mu_0^D}{m^2} , \quad (3.9)$$

that is

$$(-1)^N \frac{\delta^N Z_R[J]}{\delta J(x_1) \dots \delta J(x_N)} \Big|_{J=0} = \langle 0 | T \eta(x_1) \dots \eta(x_N) | 0 \rangle . \quad (3.10)$$

By (2.18) the evaluation of the N -point function (3.10) is immediate

$$\begin{aligned} & \langle 0 | T \eta(x_1) \dots \eta(x_N) | 0 \rangle \\ &= (-1)^N \frac{\delta^N Z_0[J]}{\delta J(x_1) \dots \delta J(x_N)} \Big|_{J=0} + \alpha^N \int d^D y \Delta_F(y-x_1) \dots \Delta_F(y-x_N) . \end{aligned} \quad (3.11)$$

Acknowledgements

It is a pleasure to thank Antonio Bassetto, Giuseppe De Grassi, Pieralberto Marchetti, Paride Paradisi, Paolo Pasti, Javi Serra, Dima Sorokin, Mario Tonin, Roberto Volpato and Andrea Wulzer for interesting discussions.

References

- [1] F. Englert and R. Brout, Phys. Rev. Lett. **13** (1964) 321.
- [2] P. W. Higgs, Phys. Lett. **12** (1964) 132.
- [3] P. W. Higgs, Phys. Rev. Lett. **13** (1964) 508.
- [4] G. S. Guralnik, C. R. Hagen and T. W. B. Kibble, Phys. Rev. Lett. **13** (1964) 585.
- [5] P. W. Higgs, Phys. Rev. **145** (1966) 1156.
- [6] T. W. B. Kibble, Phys. Rev. **155** (1967) 1554.
- [7] S. Weinberg, Phys. Rev. Lett. **19** (1967) 1264.
- [8] A. Salam, Conf. Proc. C **680519** (1968) 367.
- [9] G. Aad *et al.* [ATLAS Collaboration], Phys. Lett. B **716** (2012) 1 [arXiv:1207.7214 [hep-ex]].
- [10] S. Chatrchyan *et al.* [CMS Collaboration], Phys. Lett. B **716** (2012) 30 [arXiv:1207.7235 [hep-ex]].
- [11] M. Matone, “Quantum Field Perturbation Theory Revised,” arXiv:1506.00987.
- [12] P. Ramond, “Field Theory. A Modern Primer,” Front. Phys. **51** (1981) 1.