

The role of correlation and solvation in ion interactions with B-DNA

Maria L. Sushko*, Dennis G. Thomas†, Suzette A. Pabit‡, Lois Pollack§, Alexey V. Onufriev¶, Nathan A. Baker||

Abstract

The ionic atmospheres around nucleic acids play important roles in biological function. Large-scale explicit solvent simulations coupled to experimental assays such as anomalous small-angle X-ray scattering (ASAXS) can provide important insights into the structure and energetics of such atmospheres but are time- and resource-intensive. In this paper, we use classical density functional theory (cDFT) to explore the balance between ion-DNA, ion-water, and ion-ion interactions in ionic atmospheres of RbCl, SrCl₂, and CoHexCl₃ (cobalt hexammine chloride) around a B-form DNA molecule. The accuracy of the cDFT calculations was assessed by comparison between simulated and experimental ASAXS curves, demonstrating that an accurate model should take into account ion-ion correlation and ion hydration forces, DNA topology, and the discrete distribution of charges on DNA strands. As expected, these calculations revealed significant differences between monovalent, divalent, and trivalent cation distributions around DNA. About half of the DNA-bound Rb⁺ ions penetrate into the minor groove of the DNA and half adsorb on the DNA strands. The fraction of cations in the minor groove decreases for the larger Sr²⁺ ions and becomes zero for CoHex³⁺ ions, which all adsorb on the DNA strands. The distribution of CoHex³⁺ ions is mainly determined by Coulomb and steric interactions, while ion-correlation forces play a central role in the monovalent Rb⁺ distribution and a combination of ion-correlation and hydration forces affect the Sr²⁺ distribution around DNA.

Please address correspondence to Nathan Baker (nathan.baker@pnnl.gov).

Introduction

Interactions with ions stabilize nucleic acid secondary and tertiary structure, have a major impact on DNA packing in cells, and strongly influence protein and drug binding (1–9). A fraction of counterions bind to specific sites on nucleic acids and can be detected in crystallographic structures (10), while other counterions form a dynamic ion atmosphere around DNA, diffusing along the molecule and exchanging with ions in bulk solution (11). Mean-field approaches such as Manning counterion condensation (12) and Poisson-Boltzmann (PB) (13–16) theory have been used to obtain insight into ion distributions around biomolecules and ion-mediated interactions between macro-ions. While successful in describing some properties of nucleic acids in electrolyte solutions (e.g., RNA pK_a shifts (17), monovalent ion concentration linkages to ligand-DNA binding (8, 9, 18, 19), and low valency ion distributions around DNA), these mean-field methods often fail when the ion charge concentration increases. For example, PB models cannot capture the displacement of Na⁺ by Mg²⁺ around DNA in mixed solutions (20) or ion-mediated DNA-DNA attractive interactions (21). By imposing the constraint that a fraction of the counterions are bound (condensed) to polyelectrolyte and part form ionic atmosphere in the mean-field counterion condensation theory, it has been possible to reproduce attraction between like-charged polyelectrolytes in the presence of monovalent counterions in the intermediate range of separations (21–25). Manning suggests that the origin of this effect lies in the increase in entropy due to the increase in the effective volume available for condensed counterions as two DNA molecules approach (23). Such condensation implies penetration of ions through the DNA hydration layer and their partial desolvation to form direct bonds with DNA (26–28). Describing this process requires atomistic or coarse-grained representation of the macro-ion,

*Physical Sciences Division, Pacific Northwest National Laboratory, Richland, WA 99352.

†Biological Sciences Division, Pacific Northwest National Laboratory, Richland, WA 99352

‡School of Applied and Engineering Physics, Cornell University, Ithaca, NY 14853-3501.

§School of Applied and Engineering Physics, Cornell University, Ithaca, NY 14853-3501.

¶Department of Computer Science and Department of Physics, Virginia Tech, Blacksburg, VA 24061.

||Computational and Statistical Analytics Division, Pacific Northwest National Laboratory, Richland, WA 99352




Figure 1: Macroion models used in classical DFT simulations: (a) Model of a cylinder with uniform axial charge density; (b) discrete charge model.

which captures both the discreteness of charge distribution on the DNA strands and DNA topology, as well as a model for ion desolvation. Such characteristics are not currently present in the PB equation or other popular models of biomolecular electrostatics.

These failures suggest that, to reliably describe ion distribution around nucleic acids, the theoretical model must be refined to include more detailed interactions and incorporate higher-order non-mean-field interactions such as fluctuations. Such extensions of PB approach have been developed for simple geometries (e.g., plates, rods, spheres, etc.) to include second-order terms representing the interactions between fluctuations in ionic densities (29–35). These extended models and molecular simulations (36–44) as well as experimental data (45–56) predict attraction between like-charged objects in the presence of multivalent electrolytes.

In this study, we establish a minimal model based on classical density functional theory (cDFT) to systematically study the influence of the discrete DNA molecular charge representation, ion-ion correlations, and ion-solvent interactions on the distribution of monovalent and multivalent ions around highly charged macromolecules. We show that this model is able to accurately reproduce the results of anomalous small-angle X-ray scattering (ASAXS) experiments (57–60) for B-DNA in RbCl, SrCl₂ and CoHexCl₃ solutions. As expected, ion-ion correlations play a significant role in the accurate prediction of ASAXS curves. However, our results also demonstrate the importance of ion solvation in cation-DNA interactions and show that for doubly-charged cations these interactions can be as important as ion-ion correlations in modeling ion distributions around DNA.

Methods

DNA models

We used two coarse-grained models for the DNA macro-ion in the cDFT simulations: an infinitely long cylinder with a uniform line charge density along its z -axis (charge distribution -1 e per 0.17 nm and the 2 nm cylinder diameter) and a model with a discrete charge distribution (Fig. 1). The discrete charge distribution of the second model is described by three particle types: two helical arrays of charged spheres that represent the phosphate groups (charge -1 e, diameter 0.42 nm), two helical arrays of neutral spheres (diameter 0.42 nm) that represent the sugar/base groups, and an array of overlapping neutral spheres (diameter 0.78 nm) defining the DNA axis (61). The positions of these spheres were chosen to mimic B-form DNA using a cylindrical coordinate system (r_j^s, ϕ_j^s, z_j^s) for DNA strand s and residue j . The phosphate spheres have coordinates $r_j^s = 0.89$ nm, $\phi_j^s = \phi_0^s + 36j$ degrees, and $z_j^s = z_0^s + 0.34j$ nm; the sugar/base spheres have coordinates $r_j^s = 0.59$ nm, $\phi_j^s = \phi_0^s + 36j$ degrees, and $z_j^s = z_0^s + 0.34j$ nm; and the axis spheres have coordinates $r = 0$ nm, $\phi = 0$ degrees, and $z_j = 0.5 + 0.34j$ nm. There are 9 residues ($j = 0, \dots, 9$) per turn of B-DNA; the angular cylindrical coordinates for the first and second strands start at $\phi_0^{(1)} = 0$ and $\phi_0^{(2)} = 154$ degrees, respectively.

Table 1: A summary of the different computational models used in this paper to assess the influence of different energetic contributions (ion-ion electrostatic correlations, ion-ion steric correlations, ion-solvent interactions, and water structural changes) on DNA-ion distributions and compare the resulting distribution functions with experimental ASAXS data. The rows provide model descriptions while the columns indicate which physical phenomena are included by the models.

Model	Ion-ion elec. correl.	Ion-ion steric correl.	Ion-solvent interact.	Water struct. change	Rb ⁺ exp. agreement	Sr ²⁺ exp. agreement	CoHex ³⁺ exp. agreement
NLPB	no	no	no	no	no*	no*	yes
cDFT, no correlation	no	yes	no	yes	no*	no*	yes
cDFT, no ion solvation	yes	yes	no	yes	yes	no*	yes
cDFT, full model	yes	yes	yes	yes	yes	yes	yes

*Agreement with experiment can be obtained by fitting ion radii.

Computational models

A variety of computational models were used with the DNA models described above to assess the influence of different energetic contributions on DNA-ion interactions. These models are summarized in Table 1 and described in detail in the following sections.

Classical density functional theory (cDFT)

Classical DFT (cDFT) was used to determine the equilibrium distributions of multicomponent salt solutions surrounding DNA (62, 63). In our cDFT models, the aqueous salt solution was modeled as a dielectric medium with $\epsilon = 78.5$, charged spherical particles representing ions, and neutral spherical particles representing water molecules. The concentration of spherical “water molecules” was 55.5 M, chosen to model experimental water density. The solutions considered in this work were aqueous NaCl, RbCl, SrCl₂ and CoHexCl₃ electrolytes in Na⁺ buffer. These electrolytes were chosen based on the availability of the experimental data for these systems (57, 58). We used experimental crystalline ionic diameters for mobile ions: $\sigma_{\text{Na}} = 0.204$ nm, $\sigma_{\text{CoHex}} = 1.166$ nm, $\sigma_{\text{Sr}} = 0.252$ nm, $\sigma_{\text{Rb}} = 0.322$ nm, $\sigma_{\text{Cl}} = 0.362$ nm, and $\sigma_{\text{water}} = 0.275$ nm (64). The ion charges were $q_{\text{Na}} = +1$, $q_{\text{CoHex}} = +3$, $q_{\text{Sr}} = +2$, $q_{\text{Rb}} = +1$, $q_{\text{Cl}} = -1$, and $q_{\text{water}} = 0$. Parameterization of the cDFT model was performed against experimental data for the concentration dependence of mean activity coefficients in bulk electrolyte solutions (see Supporting Information). All calculations were performed at 298 K temperature.

To determine the equilibrium water and ion distributions via cDFT, the total Helmholtz free energy functional is minimized with respect to the densities of all the species in the presence of rigid DNA models. For this optimization, it is convenient to partition the total free energy of the system into so-called ideal (\mathcal{F}^{id}) and excess components (\mathcal{F}^{ex}) (62). The ideal free energy corresponds to the non-interacting system and is determined by the configurational entropy contributions from water and small ions,

$$\mathcal{F}^{\text{id}} = kT \sum_i^N \int_{\Omega} (\rho_i(\mathbf{r}) \log \rho_i(\mathbf{r}) - \rho_i(\mathbf{r})) d\mathbf{r} \quad (1)$$

where k is Boltzmann’s constant, T is the temperature, $\rho_i : \Omega \mapsto [0, 1]$ is the density profile of ion species i , N is the number of ion species, $\mathbf{r} \in \Omega$ is the ion coordinate, and $\Omega \in \mathbb{R}^3$ is the calculation domain. The excess free energy is generally not known exactly but can be approximated by

$$\mathcal{F}^{\text{ex}} \approx \mathcal{F}_{\text{hs}}^{\text{ex}} + \mathcal{F}_{\text{corr}}^{\text{ex}} + \mathcal{F}_{\text{C}}^{\text{ex}} + \mathcal{F}_{\text{solv}}^{\text{ex}} \quad (2)$$

where $\mathcal{F}_{\text{hs}}^{\text{ex}}$ is the hard-sphere repulsion term, $\mathcal{F}_{\text{corr}}^{\text{ex}}$ is the electrostatic correlation term, $\mathcal{F}_{\text{C}}^{\text{ex}}$ is the direct Coulomb term, and $\mathcal{F}_{\text{solv}}^{\text{ex}}$ is the ion solvation term included in some cDFT calculations (as described below).

The hard sphere term excess free energy describes ion and water many-body interactions in condensed phase due to density fluctuations and can be approximated by Fundamental Measure Theory (65) as

$$\mathcal{F}_{\text{hs}}^{\text{ex}} \approx \int_{\Omega} \Phi^{\text{hs}} [n_w(\mathbf{r})] d\mathbf{r} \quad (3)$$

where the functional Φ^{hs} has the form (66)

$$\Phi_{\text{hs}}(\mathbf{r}) = -n_0 \ln(1 - n_3) + \frac{n_1 n_2}{1 - n_3} + \left[\frac{1}{36\pi n_3^2} \ln(1 - n_3) + \frac{1}{36\pi n_3 (1 - n_3)^2} \right] n_2^3 - \frac{\mathbf{n}_1 \cdot \mathbf{n}_2}{1 - n_3} - \left[\frac{1}{12\pi n_3^2} \ln(1 - n_3) + \frac{1}{12\pi n_3 (1 - n_3)^2} \right] n_2 (\mathbf{n}_2 \cdot \mathbf{n}_2). \quad (4)$$

where n_α and \mathbf{n}_β are the scalar and vector weighted averages of the density distribution functions $\rho_i(\mathbf{r})$ and are defined by:

$$n_\alpha(\mathbf{r}) = \sum_i \int_\Omega \rho_i(\mathbf{r}') \omega_i^{(\alpha)}(\mathbf{r}' - \mathbf{r}) d\mathbf{r}', \quad \text{for } \alpha = 0, 1, 2, 3$$

$$\mathbf{n}_\beta(\mathbf{r}) = \sum_i \int_\Omega \rho_i(\mathbf{r}') \omega_i^{(\beta)}(\mathbf{r}' - \mathbf{r}) d\mathbf{r}', \quad \text{for } \beta = 1, 2.$$

In the limit of a bulk hard-sphere fluid in the absence of external fields, vector densities \mathbf{n}_1 and \mathbf{n}_2 vanish. In the same limit, the four scalar weighted densities reduce to the sum of bulk densities for all species (n_0) and the 1D (n_1), 2D (n_2), and 3D (n_3) packing fractions. The ‘‘weight functions’’ $\omega_i^{(\alpha)}$ and $\omega_i^{(\beta)}$, characterizing the geometry of particles (hard sphere with radius R_i for ion species i), are given by (66)

$$\omega_i^{(3)}(\mathbf{r}) = \theta(|\mathbf{r}| - R_i) \quad (5)$$

$$\omega_i^{(2)}(\mathbf{r}) = |\nabla \theta(|\mathbf{r}| - R_i)| = \delta(|\mathbf{r}| - R_i) \quad (6)$$

$$\omega_i^{(2)}(\mathbf{r}) = \nabla \theta(|\mathbf{r}| - R_i) = \frac{\mathbf{r}}{r} \delta(|\mathbf{r}| - R_i) \quad (7)$$

$$\omega_i^{(0)}(\mathbf{r}) = \omega_i^{(2)}(\mathbf{r}) / (4\pi R_i^2) \quad (8)$$

$$\omega_i^{(1)}(\mathbf{r}) = \omega_i^{(2)}(\mathbf{r}) / (4\pi R_i) \quad (9)$$

$$\omega_i^{(1)}(\mathbf{r}) = \omega_i^{(2)}(\mathbf{r}) / (4\pi R_i). \quad (10)$$

In the preceding formula, θ is the Heaviside step function, with $\theta(x) = 0$ for $x > 0$ and $\theta(x) = 1$ for $x \leq 0$, and δ denotes the Dirac delta function.

The electrostatic correlation term ($\mathcal{F}_{\text{corr}}^{\text{ex}}$) can be derived using the Mean Spherical Approximation (31, 33)

$$\mathcal{F}_{\text{corr}}^{\text{ex}} = \mathcal{F}_{\text{corr}}^{\text{ex}} [\{\rho_i^{\text{bulk}}\}] - kT \int_\Omega \sum_i^N c_i^{(1)} (\rho_i(\mathbf{r}) - \rho_i^{\text{bulk}}) d\mathbf{r} - \frac{kT}{2} \int_\Omega \int_\Omega \sum_{i,j}^N c_{ij}^{(2)} (\rho_i(\mathbf{r}) - \rho_i^{\text{bulk}}) (\rho_j(\mathbf{r}') - \rho_j^{\text{bulk}}) d\mathbf{r} d\mathbf{r}' \quad (11)$$

where ρ_i^{bulk} is the bulk concentration of ion species i and the first term describes ion correlation free energy in bulk electrolyte solution in the absence of DNA. The first-order direct correlation functions are defined as

$$c_i^{(1)} = -\frac{\mu_i}{kT}, \quad (12)$$

where μ_i is the chemical potential of ion species i . The second-order direct correlation functions are defined as

$$c_{ij}^{(2)}(\mathbf{r} - \mathbf{r}') = \begin{cases} -\frac{q_i q_j}{kT\epsilon} \left(\frac{2B}{\sigma_{ij}} - \left(\frac{B}{\sigma_{ij}} \right)^2 |\mathbf{r} - \mathbf{r}'| - \frac{1}{|\mathbf{r} - \mathbf{r}'|} \right) & |\mathbf{r} - \mathbf{r}'| \leq \sigma_{ij} \\ 0 & |\mathbf{r} - \mathbf{r}'| > \sigma_{ij}, \end{cases} \quad (13)$$

where q_i is the charge of ion species i , ϵ is the dielectric constant of the solvent, $\sigma_{ij} = (\sigma_i + \sigma_j) / 2$ is the hard-sphere contact distance between ions of diameters σ_i and σ_j , B is given by

$$B = \frac{1}{\xi} \left(\xi + 1 - \sqrt{1 + 2\xi} \right), \quad (14)$$

$\xi = \kappa\sigma_{ij}$, κ is the inverse Debye length $\kappa^2 = l_B \sum_i q_i^2 \rho_i^{\text{bulk}}$, $l_B = \frac{e^2}{kT\epsilon}$ is the Bjerrum length, and e is the unit charge. The direct Coulomb free energy term can be calculated exactly

$$\mathcal{F}_C^{\text{ex}} = \frac{kTl_B}{2} \int_{\Omega} \int_{\Omega} \sum_{i,j} \frac{q_i q_j}{|\mathbf{r} - \mathbf{r}'|} \rho_i(\mathbf{r}) \rho_j(\mathbf{r}') d\mathbf{r} d\mathbf{r}'. \quad (15)$$

Finally, the solvation term $\mathcal{F}_{\text{solv}}^{\text{ex}}$ models ion-water interactions with a square well potential

$$V(\mathbf{r} - \mathbf{r}') = \begin{cases} \infty & |\mathbf{r} - \mathbf{r}'| < \sigma_{ij} \\ -\varepsilon & \sigma_{ij} \leq |\mathbf{r} - \mathbf{r}'| \leq \sigma_{ij} + h \\ 0 & \sigma_{ij} + h < |\mathbf{r} - \mathbf{r}'|, \end{cases} \quad (16)$$

where ε is the well depth, and h is the well width. For the current study, $h = 0.2\zeta$ is the well width for interactions between ions and water and ζ is the sum of radii of interacting particles (63). The following well depths were calculated using SPC/E water using the parameters from Horinek et al: $\varepsilon_{\text{Sr}} = 0.01038$ eV, $\varepsilon_{\text{Cl}} = 0.0053894$ eV, $\varepsilon_{\text{Rb}} = \varepsilon_{\text{CoHex}} = 0.0021$ eV (67). Simulations of concentration dependence of ion activity coefficients in RbCl and CoHexCl₃ solutions demonstrated that adding attractive ion-water interactions does not affect the ion chemical potential (see Supporting Information).

Minimization of the excess free energy functional \mathcal{F}^{ex} with respect to the water and ion densities gives

$$\rho_i(\mathbf{r}) = \exp\left(\frac{\mu_i}{kT} - \frac{1}{kT} \frac{\delta \mathcal{F}^{\text{ex}}}{\delta \rho_i(\mathbf{r})}\right). \quad (17)$$

We solve Poisson's equation

$$-\nabla \cdot \epsilon(\mathbf{r}) \nabla \varphi(\mathbf{r}) = \sum_i q_i \rho_i(\mathbf{r}) \quad (18)$$

for the electrostatic potential ($\varphi(\mathbf{r})$) where $\epsilon(\mathbf{r})$ is the dielectric coefficient. For an infinitely long uniformly charged cylinder in electroneutral conditions, the potential

$$\varphi(r) = \frac{4\pi}{\epsilon} \int_r^\infty t \log\left(\frac{r}{t}\right) \sum_i q_i \rho_i(t) dt. \quad (19)$$

Using this potential for the cylinder model and a numerical solution to Poisson's equation (Eq. 18) for the 3D DNA model, the expression for the densities is

$$\rho_i(\mathbf{r}) = \exp\left(\frac{\mu_i}{kT} - \frac{q_i \varphi(\mathbf{r})}{kT} - \frac{1}{kT} \frac{\delta (\mathcal{F}_{\text{hs}}^{\text{ex}} + \mathcal{F}_{\text{corr}}^{\text{ex}} + \mathcal{F}_{\text{solv}}^{\text{ex}})}{\delta \rho_i(\mathbf{r})}\right). \quad (20)$$

The resulting system of Eqs. 18 and 20 was solved iteratively to self-consistency using the numerical procedure described by Meng (66). In particular, equilibrium ion density distributions were obtained using a relaxed Gummel iterative procedure for 3D systems and Picard iterations in 1D. Convergence was considered to be achieved when the maximum difference between the input and the output density profiles between iterations was smaller than 10^{-6} .

Three main features distinguish our approach from previous cDFT models (68–70). First, our model includes a full representation of the coarse-grained DNA topology and a discrete distribution of charges. Second, we use Pauling diameters for ions and van der Waals diameters for water molecules as opposed to previous restricted models where all species have the same diameter. Finally, our model includes water-ion attractive interactions.

Anomalous small-angle X-ray scattering curve calculations

ASAXS profiles were calculated using the ion density distributions $\rho_i(r)$ around DNA. In the 3D model, ion densities were averaged in cylindrical coordinates over the cylinder azimuthal angle ϕ and length z for each radial distance r from the DNA axis. The excess form factor for ion species α was calculated as

$$F_{\text{ion},\alpha}(Q) = a_\alpha \int \rho_\alpha(r) e^{-iQr} dr, \quad (21)$$




Figure 2: Ionic distributions around a uniformly charged cylinder. (a) Solutions of 100 mM NaCl, 100 mM RbCl, and 10 mM SrCl_2 in a 1 mM NaCl buffer; concentration profiles are shown for Na^+ (solid line), Rb^+ (dashed line), and Sr^{2+} (dot-dashed line). (b) Solutions of 5 mM CoHexCl_3 in a 20 mM NaCl buffer; concentration profiles are shown for Na^+ (solid line), CoHex^{3+} (dashed line), and Cl^- (dot-dashed line).

where a_α is a constant related to the average electron density of ion species α and Q is the scattering vector. In the current study, we only consider the excess form factor due to cation species; the chloride anion has no ASAXS response. Furthermore, we only consider a single cation species at a time so that $F_{\text{ion}}(Q) = F_{\text{ion},\alpha}(Q)$.

The excess form factor of DNA ($F_{\text{DNA}}(Q)$) was calculated using AquaSAXS (71). from the form factor of DNA in vacuo ($F_{\text{DNA}}^{\text{vac}}(Q)$), the form factor of the volume of water excluded by DNA ($F_{\text{DNA}}^{\text{excl}}(Q)$), and the form factor of hydration shell of the DNA ($F_{\text{hsh}}(Q)$):

$$F_{\text{DNA}}(Q) = F_{\text{DNA}}^{\text{vac}}(Q) - \rho_w F_{\text{DNA}}^{\text{excl}}(Q) + \rho_w F_{\text{hsh}}(Q), \quad (22)$$

where ρ_w is the bulk density of water. The form factor of the hydration shell is calculated using water density maps, $\rho_{\text{hsh}}(r)$, obtained via AquaSol (72), which employs the Poisson-Boltzmann formalism with water treated as an assembly of self-oriented dipoles:

$$F_{\text{hsh}}(Q) = b \int \left(\frac{\rho_{\text{hsh}}(r)}{\rho_w} - 1 \right) e^{-iQr} dr \quad (23)$$

where b is a scale factor to adjust the hydration shell contribution (usually $b = 1.0$) and integration is performed over the region where solvent density deviates from the bulk by a factor larger (in magnitude) than $\pm 10^{-4}$.

The ASAXS intensity is then calculated from these quantities as

$$I(Q) = 2(f'_{\text{ion}}(E_1) - f'_{\text{ion}}(E_2)) (f_{\text{DNA}} N_{\text{ion}} F_{\text{DNA}}(Q) F_{\text{ion}}(Q) + f_{\text{ion}0} N_{\text{ion}}^2 F_{\text{ion}}(Q)^2) + (f''_{\text{ion}}(E_1) - f''_{\text{ion}}(E_2)) N_{\text{ion}}^2 F_{\text{ion}}^2(Q) \quad (24)$$

where $f'_{\text{ion}}(E_i)$ is the energy-dependent real part of ion anomalous scattering factor, E_1 is the energy far from the X-ray absorption edge of the ion, E_2 is the energy near the edge where ion scattering is suppressed by absorption, $f_{\text{ion}0}$ is the energy independent solvent-corrected scattering factor, f_{DNA} is the effective number of electrons from DNA and N_{ion} is the number of excess ions around DNA (57) (see Supporting Information for more details). Since experimental data are available in arbitrary units, theoretical intensities were uniformly scaled with a common scaling factor, chosen to match the experimental and calculated intensities, obtained using 3D cDFT-full model, at low Q .

Results

Comparison between DNA Model systems

The uniformly charged cylinder model (Fig. 1 left) represents a one-dimensional case for which ionic distribution is only a function of the radial distance from the cylinder axis. The results for this 1D system are shown in Fig. 2. For monovalent ions, the 1D cDFT calculations predict 91.5% and 77.5% DNA charge neutralization by Na^+ and Rb^+ , respectively. Such differences in monovalent cation condensation on DNA were not observed experimentally (73), demonstrating a fundamental




Figure 3: Panoramic view of cation distributions around DNA in 100 mM RbCl calculated using (a) cDFT, and (b) and NLPB (solid line) and the cDFT model with no ion-correlation interactions (cDFT-nc, dotted line). Rb⁺ distributions on the DNA strand are shown as black lines and in the minor groove as red lines with (c) a zoom-in into a low-density region for Rb⁺ in plot (b). Na⁺ distributions are shown in blue. (d) Radial distribution function of Rb⁺ ions around DNA molecule.

deficiency of a uniformly charged cylinder model for simulating ionic atmosphere around DNA. Furthermore, for divalent ions, the 1D cDFT calculations predict charge inversion at the DNA surface in SrCl₂ solution. Note that charge inversion in the presence of multivalent salts has also been observed in cDFT and MC simulations for a cylinder DNA model (68, 69). However, we do not see this effect in our more detailed 3D DNA geometry simulations (see below). Finally, for trivalent ions, 90% DNA charge neutralization is found within 5 CoHex radii from the cylinder surface for the 1D cDFT calculations. Competitive cation condensation in mixed 5 mM CoHexCl₃ + 20 mM NaCl solutions results in preferential CoHex³⁺ condensation on the cylinder surface: sodium ions are not found in the immediate vicinity of the DNA. This competition is in qualitative agreement with experimental observations of a negligible effect of Na⁺ on CoHex³⁺ binding when NaCl concentration is below 40 mM (74).

We also performed 3D cDFT calculations of the same electrolyte solutions surrounding the helical discrete charge model (Fig. 1 right). Fig. 3 shows cDFT results for the monovalent ion Rb⁺. As shown in the panoramic density profiles, cDFT




Figure 4: Panoramic density distributions of Sr^{2+} ions on DNA (a) strands and (b) minor grooves obtained using 3D cDFT (solid red line), NLPB (solid black line) and cDFT-nc (dotted line). SrCl_2 concentration is 10 mM. (c) Radial distribution function of Sr^{2+} ions around DNA molecule obtained using a complete cDFT model (solid line), cDFT model without ion-water attractive interactions (dotted line).

predicts a two-peak radial density distribution of Rb^+ : first peak at around 0.6 nm is due to cation penetration into DNA minor grooves and the second peak at 1.2 nm to Rb^+ condensation on DNA strands (Fig. 3a,c). These data are in good quantitative agreement with molecular dynamics results obtained using TIP3P water model (28). Fig. 4 shows cDFT results for the divalent ion Sr^{2+} . In the case of Sr^{2+} , the effect of ion solvation can be clearly seen in the density distribution of Sr^{2+} with respect to the DNA axis. While both cDFT models - with and without ion-solvation - produce two-peak Sr^{2+} density distributions at the same positions with respect to the DNA axis, the density distributions are qualitatively different. In particular, the model without ion solvation predicts much higher Sr^{2+} concentration in the DNA grooves than on strands, while the trend is reversed in the model with ion-solvation. The 3D cDFT results for trivalent CoHexCl_3 solutions are shown in Fig. 5b and are very similar to those obtained from the 1D cDFT model.

Finally, we used the results of our cDFT and NLPB calculations to determine ASAXS profiles as described in the *Methods* section. The results of these calculations for Rb^+ and Sr^{2+} , together with experimental data, are shown in Fig. 6. Similar




Figure 5: (a) Panoramic density distributions of CoHex^{3+} ions on DNA strand obtained using 3D cDFT (solid red line), NLPB (solid black line; the curve is shifted up by 0.5 mM for clarity) and cDFT-nc (dotted line). CoHexCl_3 concentration is 0.5 mM. (b) Radial distribution function of CoHex^{3+} ions around DNA molecule. Note: as seen from the RDF, there is zero CoHex density in the minor groove, so the corresponding panoramic density is not shown as a separate plot.




Figure 6: Simulated and experimental ASAXS profiles for 25 bp DNA in (a) 100 mM RbCl and (b) 10 mM SrCl_2 solutions. Experimental data (57, 58) are shown as black dots, simulations results obtained using 3D cDFT as blue lines, NLPB as green lines. Simulation results in (a) obtained using 1D cDFT and NLPB results, are shown as dotted blue and green lines, respectively. The red line in (b) corresponds to cDFT results obtained in the model with no ion-water attractive interactions.

results for CoHexCl_3 are shown in Fig. 7.




Figure 7: ASAXS profiles for 0.5 mM CoHexCl₃ DNA solutions. Experimental data are shown with a thick black line (unpublished) and those from Andresen et al (58) with a blue line. 1D cDFT results are shown as thin black line, 1D NLPB results as red line, 3D cDFT and NLPB data coincide and are shown as green line.

Discussion

Comparison to Manning condensation

The cDFT calculations of ionic distributions (Fig. 2) for the uniformly charged cylinder model reproduce the Manning condensation limits (12) with approximately 1M concentrations of singly-charged cations at the cylinder surface. A complete 3D cDFT model also reproduces the Manning condensation limit for monovalent cations: the total concentration of Rb⁺ on strands and in minor groove is about 1 M (Fig. 3). Additionally, the 3D cDFT model predicts that the multivalent ions form much denser layers at the DNA surface than the monovalent cations (Figures 3, 4, and 5), consistent with Manning theory. The good correlation between our non-mean-field cDFT model (with full ion-ion correlations included) and the mean-field Manning theory is somewhat surprising, particularly given the significant differences observed in the total condensed ion densities between cDFT and the mean-field NLPB approaches. However, Manning theory indirectly accounts for interactions beyond first-order electrostatics through partitioning the total ion density into condensed ions and the surrounding ionic atmosphere. This accounts for the success of Manning theory in predicting the total concentration of 1:1 electrolyte counterions condensed on DNA as observed in experiments (12) and recent MD simulations (28).

Ion interaction with DNA grooves

By definition, the cylinder model does not allow ion penetration inside DNA and therefore yields well-known monotonically decreasing counterion distributions shown in Fig. 2. Thus, the model is not adequate for describing the interaction between DNA and small weakly solvated Na⁺ and Rb⁺ ions, which are known to penetrate into the minor grooves of DNA (28, 75). However, our more detailed helical charge model allows ion penetration. Simulations of RbCl solutions using this model showed that about half of the condensed Rb⁺ ions are bound to the minor groove of the DNA molecule (Fig. 3). The distributions of cation densities on the DNA strands and in the minor groove are highly structured: they exhibit a periodicity correlated with the periodic spacing of phosphate groups on DNA strands. In contrast, cation distributions in the major groove are mostly featureless (see Fig. S1 in Supporting Information), in agreement with previous simulations and experimental data (27). Penetration of some cations into the grooves lowers the effective charge density on the DNA, limiting cation condensation on strands.

Increasing cation valency correlates with a stronger preference of cation binding to phosphate groups on the DNA strands (Figures 3, 4, and 5). A similar preference for CoHex³⁺ binding to phosphates of B-DNA was also observed in MD simulations (76) and is determined by the strong electrostatic attraction of the trivalent cations to phosphate groups, CoHex³⁺-CoHex³⁺ repulsion, and steric inaccessibility of B-DNA minor groove to the large CoHex³⁺ ions. As shown in Figures 4 and 5, both

Sr^{2+} and CoHex^{3+} ions preferentially bind to every fourth phosphate on the strand. Further away from the DNA axis, the Sr^{2+} density variations along the angular cylindrical coordinate have the same period as the period of the angular phosphate distribution. The period of the density variations for CoHex^{3+} is two times larger than for Sr^{2+} ; i.e., some Sr^{2+} ions can penetrate into the minor groove, while CoHex^{3+} ions bind exclusively to phosphate groups on the strands.

Influence of correlation on ion distributions

To investigate the influence of ion correlation forces on the distribution of ions around DNA, we used a cDFT model without ion-correlation interactions (cDFT-nc) as well as the nonlinear Poisson-Boltzmann (NLPB) model, which also lacks correlation (Figures 3, 4, and 5). Both models without correlations yield qualitatively different ion distributions than the 3D cDFT calculations which include correlations. In the presence of correlations, sterically allowed ions accumulate in the minor groove; in the absence of correlations, ions accumulate near phosphate groups on the exterior of the DNA strand. The largest qualitative difference between NLPB and cDFT ion distributions was observed for the Rb^+ density distribution. In NLPB, Rb^+ ions decorate the phosphate groups, driven by Coulombic interactions; the panoramic distribution of Rb^+ ions condensed on DNA strands in NLPB model has a larger peak at 45° and a smaller one at 135° (Figures 3, 4, and 5). In contrast, ion-ion correlations reduce the effective electrostatic repulsion between cations promoting their penetration into the grooves. Due to stronger Coulomb interactions between multiply-charged cations the effect of correlations is weaker for Sr^{2+} and CoHex^{3+} resulting in the decrease in the fraction of counterions in the grooves with ion radius and charge (Fig. 4 and Fig. 5). For CoHex^{3+} the concentration of counterions in the grooves becomes insignificant. As a result, NLPB and cDFT predict qualitatively similar panoramic density distributions on DNA strands for Sr^{2+} and CoHex^{3+} (Figures 4 and 5).

The models without correlations (NLPB and cDFT-nc) are very similar to each other, indicating the major influence of correlation on even low charge-density (monovalent) ion behavior. This result contrasts the conclusion that correlations are insignificant in monovalent electrolytes from early theories of ion correlations (77). However, these theories considered electrolytes at uniformly charged surfaces ignoring the influence of the discreteness of charge distribution on fluctuations in ionic atmosphere. Not surprisingly, these models do not capture the experimentally observed attraction between like-charged polyelectrolytes in low concentration monovalent electrolytes (23, 52). Recent molecular dynamic simulations also point to the importance of nonmean-field interactions between biomolecules and monovalent electrolytes manifested in a more structured ionic atmosphere than that predicted by NLPB (28, 78). The small difference between the cDFT-nc and NLPB models (in the height of the double peak around 45°) for Rb^+ is due to the solvent excluded-volume effects included in the cDFT-nc model and absent from NLPB theory (Fig. 3).

Ion correlations also influence ion-specific details in density distributions for counterions of the same valency. 3D cDFT results demonstrate that smaller Na^+ ions tend to accumulate on DNA strands and minor groove while Rb^+ ions are more evenly distributed along the DNA helix (see Fig. 3a). In the cDFT-nc and NLPB models with no correlation, the differences between Na^+ and Rb^+ distributions are significantly smaller (see Fig. 3b), suggesting that ion-correlation interactions are responsible for this effect.

Comparison with ASAXS experiments

ASAXS profiles calculated using the 3D cDFT model show very good agreement with experimental data (58) for RbCl solutions (Fig. 6). The shapes of the scattering curves are very similar in the 1D cDFT and 3D cDFT models, with similar average numbers of condensed counterions: the 1D and 3D cDFT calculations predict 34.9 and 34.6 condensed Rb^+ ions, respectively. Both predictions are within error of the experimental measurement of 34 ± 3 ions (57). However, the distribution of Rb^+ ions is different in these models: all condensed cations decorate the cylinder surface (by definition) in the 1D model while half of the condensed cations are on DNA strands and the other half are in minor grooves in the 3D models. In contrast, the 3D NLPB model shows a significant deviation of the simulated scattering curve from the experimental one (Fig. 6). As discussed in the previous section, penetration of some cations into DNA grooves reduces the negative electrostatic potential acting on cations in solution. In the absence of any interactions beyond Coulomb forces, this penetration leads to lower concentrations of cations on the DNA surface and lower total concentrations of condensed counterions. Previous NLPB simulations demonstrated that adjusting the ionic radius of Rb^+ to its hydrated radius and prohibiting ion penetration into the DNA hydration shell can lead to closer agreement between 3D NLPB results and experiment (73) – but at the price of an incorrect ion distribution around DNA.

Comparison of calculated and experimental SrCl_2 data highlight the importance of solvation on ion distributions around DNA. Sr^{2+} ions have a significant hydration energy; approximately 3 times higher than that of monovalent alkali metal ions. These strong cation-water interactions lower the entropy of water molecules around cations, but introduce a higher enthalpy

cost for partial ion desolvation (79). Simulations with the solvent approximated as dielectric continuum do not account for such desolvation, limiting interactions in the system to first- and second-order electrostatic interactions: Coulomb and ion-correlation forces. To understand the importance of these desolvation contributions, we used two variants of the cDFT model: one with attractive cation-water interactions and another without. As shown in Fig. 6, ASAXS curves calculated using cDFT without desolvation contributions deviate significantly from the experimental data and the ASAXS curves calculated using the complete cDFT model. On the other hand, the experimental ASAXS curves agree with those calculated from the complete cDFT model. The importance of desolvation is also emphasized by the fact that inclusion of such interactions is essential for reproducing the chemical potentials of divalent cations but is not required for weakly hydrated alkali metal ions or CoHex^{3+} (see Supporting Information). In summary, ion solvation behavior is important for accurately modeling ion-DNA interactions: desolvation reduces the excess chemical potential of cations and anions, lowering the effective concentration of electrolyte and weakening ion-ligand interactions. Surprisingly, an NLPB model includes neither solvation nor ion-correlation interactions reproduces the experimental ASAXS curves for Sr^{2+} (Fig. 6), although some differences are obvious in the more detailed radial distribution functions (Fig. 4). This agreement is serendipitous and is due to cancellation of errors from the lack of ion-correlation, that favors ion accumulation in the grooves, and solvation, that limits ion concentration in the grooves.

Unexpectedly, the trivalent CoHexCl_3 solution is the simplest ion to model around DNA; CoHex^{3+} can be reliably described by first-order electrostatics (i.e., direct Coulomb interactions). CoHex^{3+} ions decorate DNA phosphate strands and do not penetrate inside B-DNA grooves. All models explored in this paper show reasonable agreement between the calculated ASAXS profiles and the experimental data (Fig. 7), and is consistent with all-atom MD simulations in explicit solvent (76). For these triply-charged ions, the good agreement between cDFT and NLPB is caused by the dominance of first-order electrostatics in ion-DNA interactions over higher-order ion-ion correlations. Moreover, the large CoHex^{3+} ionic diameter creates a steric barrier for ion penetration inside the grooves, rendering the 1D cylinder models adequate for calculating the average number of condensed CoHex^{3+} ions. Finally, because the diameter of CoHex^{3+} is large, the field at its surface is comparable to Na^+ and the effects of solvation are lower than for the smaller divalent Sr^{2+} ions. It follows from the current study that the models required to describe CoHex^{3+} around a single B-DNA strands are relatively simple. However, we expect that ion correlation forces will dominate DNA-DNA interactions between multiple strands due to the higher local phosphate charge density.

Conclusions

We have studied the details of ionic atmospheres around DNA molecule for 1:1, 2:1 and 3:1 electrolytes using a combination of cDFT and NLPB methods. Our calculations demonstrated that ion-ion correlation interactions induce counterion penetration into the DNA grooves, unless sterically prohibited by large ion radii. In particular, ion binding in the grooves – compared to binding on strands – has a profound effect on ion-induced nucleic acid condensation as demonstrated in our previous work (76). Solvation interactions have an opposite effect: when the enthalpy cost of desolvation is high (e.g., for Sr^{2+} ions), ion-water interactions limit ion penetration into the DNA grooves. Partial compensation of these two opposing effects explains the success of NLPB in reproducing the average number of condensed cations and the shape of the ASAXS curves of the ion-counting experiments. In contrast, cDFT model without ion-desolvation interactions was found to systematically overestimate ion concentration in DNA grooves. Through the comparison of several cDFT models and experimental data, we demonstrated that a minimum model to describe ion-polyelectrolyte interactions should include long-range correlations arising from density and charge density fluctuations in electrolyte solution as well as short-range ion (de)solvation forces. The latter interactions are often ignored in reduced models of electrolyte solutions limiting their applicability to the classes of weakly solvated ions. Ion hydration forces are particularly pronounced in solutions of multiply-charged ions and give significant contribution to ion activity and, therefore, to ion-polyelectrolyte interactions. Our results highlight important aspects of the properties of electrolyte solutions influencing ionic atmosphere around biomolecules that may significantly impact DNA condensation and biomolecules-ligand interactions.

Author contributions

MLS, DGT, and SAP performed the research, analyzed the data, and helped write the manuscript. LP, AVO, and NAB designed the research, analyzed the data, and assisted in writing the manuscript.

Acknowledgments

This work was supported by NIH Grant R01 GM099450. NAB and MLS acknowledge fruitful discussions of hydration effects with Chris Mundy and Greg Schenter. The cDFT calculations were performed using PNNL Institutional Computing resources at Pacific Northwest National Laboratory. PNNL is a multiprogram national laboratory operated for DOE by Battelle under Contract DE-AC05-76RL01830.

SUPPLEMENTARY MATERIAL

An on-line supplement to this article can be found by visiting BJ Online at <http://www.biophysj.org>.

References

1. Rau, D. C., and V. A. Parsegian, 1984. Attractive Long-Range Hydration Forces Can Cause Macromolecular Assembly - the Condensation of DNA by Cross-Linking Agents. *Biophysical Journal* 45:A114–A114.
2. Rau, D. C., B. Lee, and V. A. Parsegian, 1984. Measurement of the Repulsive Force between Poly-Electrolyte Molecules in Ionic Solution - Hydration Forces between Parallel DNA Double Helices. *Proceedings of the National Academy of Sciences of the United States of America* 81:2621–2625.
3. Knobler, C. M., and W. M. Gelbart, 2009. Physical Chemistry of DNA Viruses. *Annual Review of Physical Chemistry* 60:367–383.
4. Xiang, Y., A. J. Tong, and Y. Lu, 2009. Abasic Site-Containing DNAzyme and Aptamer for Label-Free Fluorescent Detection of Pb²⁺ and Adenosine with High Sensitivity, Selectivity, and Tunable Dynamic Range. *Journal of the American Chemical Society* 131:15352–15357.
5. Draper, D. E., 2008. RNA Folding: Thermodynamic and Molecular Descriptions of the Roles of Ions. *Biophysical Journal* 95:5489–5495.
6. Draper, D. E., D. Grilley, and A. M. Soto, 2005. Ions and Rna Folding. *Annual Review of Biophysics and Biomolecular Structure* 34:221–243.
7. Woodson, S. A., 2005. Metal ions and RNA folding: a highly charged topic with a dynamic future. *Current Opinion in Chemical Biology* 9:104–109.
8. Misra, V. K., J. L. Hecht, K. A. Sharp, R. A. Friedman, and B. Honig, 1994. Salt Effects on Protein-DNA Interactions - the λ CI Repressor and EcoRI Endonuclease. *Journal of Molecular Biology* 238:264–280.
9. Misra, V. K., K. A. Sharp, R. A. Friedman, and B. Honig, 1994. Salt Effects on Ligand-DNA Binding - Minor-Groove Binding Antibiotics. *Journal of Molecular Biology* 238:245–263.
10. Frederiksen, J. K., and J. A. Piccirilli, 2009. Identification of catalytic metal ion ligands in ribozymes. *Methods* 49:148–166.
11. Freisinger, E., and R. K. O. Sigel, 2007. From nucleotides to ribozymes - A comparison of their metal ion binding properties. *Coordination Chemistry Reviews* 251:1834–1851.
12. Manning, G. S., 1978. Molecular Theory of Polyelectrolyte Solutions with Applications to Electrostatic Properties of Polynucleotides. *Quarterly Reviews of Biophysics* 11:179–246.
13. Lamm, G., 2003. The Poisson-Boltzmann Equation. In K. B. Lipkowitz, R. Larter, and T. R. Cundari, editors, *Reviews in Computational Chemistry*, John Wiley & Sons, Inc., Hoboken, NJ, USA, volume 19, chapter 4, 147–365.
14. Baker, N. A., 2004. Poisson-Boltzmann Methods for Biomolecular Electrostatics. In *Methods in Enzymology*, Elsevier, volume 383, 94–118.
15. Fixman, M., 1979. The Poisson-Boltzmann equation and its application to polyelectrolytes. *The Journal of Chemical Physics* 70:4995–5005.
16. Anderson, C., and M. Record, 1990. Ion Distributions Around DNA and other Cylindrical Polyions: Theoretical Descriptions and Physical Implications. *Annual Review of Biophysics and Biophysical Chemistry* 19:423–463. <http://dx.doi.org/10.1146/annurev.bb.19.060190.002231>.
17. Tang, C. L., E. Alexov, A. M. Pyle, and B. Honig, 2007. Calculation of pK(a)s in RNA: On the structural origins and functional roles of protonated nucleotides. *Journal of Molecular Biology* 366:1475–1496.
18. Shkel, I., and T. Record, 2012. Coulombic free energy and salt ion association per phosphate of all-atom models of DNA oligomer: dependence on oligomer size. *Soft Matter* 8:9345–9355. <http://dx.doi.org/10.1039/c2sm25607j>.
19. Misra, V. K., J. L. Hecht, A. S. Yang, and B. Honig, 1998. Electrostatic contributions to the binding free energy of the λ CI repressor to DNA. *Biophysical Journal* 75:2262–2273.
20. Chu, V. B., Y. Bai, J. Lipfert, D. Herschlag, and S. Doniach, 2007. Evaluation of ion binding to DNA duplexes using a size-modified Poisson-Boltzmann theory. *Biophysical Journal* 93:3202–3209.
21. Kornyshev, A. A., and S. Leikin, 1999. Electrostatic zipper motif for DNA aggregation. *Physical Review Letters* 82:4138–4141.
22. Ray, J., and G. S. Manning, 1994. An Attractive Force between 2 Rodlike Polyions Mediated by the Sharing of Condensed Counterions. *Langmuir* 10:2450–2461.
23. Manning, G. S., 2011. Counterion condensation theory of attraction between like charges in the absence of multivalent counterions.

- European Physical Journal E 34:132.
24. Perico, A., and A. Rapallo, 2011. Clusters in strong polyelectrolyte solutions in the condensation theory approach. Journal of Chemical Physics 134:055108.
 25. Pietronave, S., L. Arcesi, C. D'Arrigo, and A. Perico, 2008. Attraction between Like-Charged Polyelectrolytes in the Extended Condensation Theory. Journal of Physical Chemistry B 112:15991–15998.
 26. Long, H., A. Kudlay, and G. C. Schatz, 2006. Molecular dynamics studies of ion distributions for DNA duplexes and DNA clusters: Salt effects and connection to DNA melting. Journal of Physical Chemistry B 110:2918–2926.
 27. Allahyarov, E., H. Lowen, and G. Gompper, 2003. Adsorption of monovalent and multivalent cations and anions on DNA molecules. Physical Review E 68:061903.
 28. Giambasu, G. M., T. Luchko, D. Herschlag, D. M. York, and D. A. Case, 2014. Ion Counting from Explicit-Solvent Simulations and 3D-RISM. Biophysical Journal 106:883–894.
 29. Gonzalez-Amezcuca, O., M. Hernandez-Contreras, and P. Pincus, 2001. Electrostatic correlation force of discretely charged membranes. Physical Review E 64:041603.
 30. Kjellander, R., S. Marcelja, R. M. Pashley, and J. P. Quirk, 1988. Double-Layer Ion Correlation Forces Restrict Calcium Clay Swelling. Journal of Physical Chemistry 92:6489–6492.
 31. Blum, L., 1975. Mean Spherical Model for Asymmetric Electrolytes .1. Method of Solution. Molecular Physics 30:1529–1535.
 32. Henderson, D., E. Waisman, J. L. Lebowitz, and L. Blum, 1978. Equation of State of a Hard-Core Fluid with a Yukawa Tail. Molecular Physics 35:241–255.
 33. Hoye, J. S., and L. Blum, 1978. Mean Spherical Model for Asymmetric Electrolytes - Thermodynamics and Pair Correlation-Function. Molecular Physics 35:299–300.
 34. Jiang, J. W., L. Blum, and O. Bernard, 2001. Charged hard dumbbell in the binding mean-spherical approximation. Molecular Physics 99:1765–1767.
 35. Jiang, J. W., L. Blum, O. Bernard, and J. M. Prausnitz, 2001. Thermodynamic properties and phase equilibria of charged hard sphere chain model for polyelectrolyte solutions. Molecular Physics 99:1121–1128.
 36. Guldbbrand, L., L. G. Nilsson, and L. Nordenskiold, 1986. A Monte-Carlo Simulation Study of Electrostatic Forces between Hexagonally Packed DNA Double Helices. Journal of Chemical Physics 85:6686–6698.
 37. Rouzina, I. F., and V. A. Bloomfield, 1996. Macroion attraction by correlated counterion fluctuations: Application to condensation of DNA. Biophysical Journal 70:Mp418–Mp418.
 38. Rouzina, I., and V. A. Bloomfield, 1996. Macroion attraction due to electrostatic correlation between screening counterions .1. Mobile surface-adsorbed ions and diffuse ion cloud. Journal of Physical Chemistry 100:9977–9989.
 39. Arenzon, J. J., J. F. Stilck, and Y. Levin, 1999. Simple model for attraction between like-charged polyions. European Physical Journal B 12:79–82.
 40. Levin, Y., J. J. Arenzon, and J. F. Stilck, 1999. The nature of attraction between like-charged rods. Physical Review Letters 83:2680–2680.
 41. Shklovskii, B. I., 1999. Screening of a macroion by multivalent ions: Correlation-induced inversion of charge. Physical Review E 60:5802–5811.
 42. Shklovskii, B. I., 1999. Wigner crystal model of counterion induced bundle formation of rodlike polyelectrolytes. Physical Review Letters 82:3268–3271.
 43. Netz, R. R., 2001. Electrostatistics of counter-ions at and between planar charged walls: From Poisson-Boltzmann to the strong-coupling theory. European Physical Journal E 5:557–574.
 44. Naji, A., A. Arnold, C. Holm, and R. R. Netz, 2004. Attraction and unbinding of like-charged rods. Europhysics Letters 67:130–136.
 45. Danilowicz, C., C. H. Lee, K. Kim, K. Hatch, V. W. Coljee, N. Kleckner, and M. Prentiss, 2009. Single molecule detection of direct, homologous, DNA/DNA pairing. Proceedings of the National Academy of Sciences of the United States of America 106:19824–19829.
 46. Lin, S. C., W. I. Lee, and J. M. Schurr, 1978. Brownian-Motion of Highly Charged Poly(L-Lysine) - Effects of Salt and Polyion Concentration. Biopolymers 17:1041–1064.
 47. Drifford, M., and J. P. Dalbiez, 1985. Effect of Salt on Sodium Polystyrene Sulfonate Measured by Light-Scattering. Biopolymers 24:1501–1514.
 48. Zero, K., and B. R. Ware, 1984. Mobilities of Poly-L-Lysine Molecules in Low-Salt Solutions. Journal of Chemical Physics 80:1610–1616.
 49. Matsuoka, H., D. Schwahn, and N. Ise, 1991. Observation of Cluster Formation in Polyelectrolyte Solutions by Small-Angle Neutron-Scattering .1. A Steep Upturn of the Scattering Curves from Solutions of Sodium Poly(Styrenesulfonate) at Scattering Vectors Below 0.01(a)⁻¹. Macromolecules 24:4227–4228.
 50. Sedlak, M., 1993. Domain-Structure of Polyelectrolyte Solutions - Is It Real. Macromolecules 26:1158–1162.
 51. Sedlak, M., 1994. On the Possible Role of Nonelectrostatic Interactions in the Mechanism of the Slow Polyelectrolyte Mode Observed by Dynamic Light-Scattering. Journal of Chemical Physics 101:10140–10144.
 52. Sedlak, M., 1996. The ionic strength dependence of the structure and dynamics of polyelectrolyte solutions as seen by light scattering: The slow mode dilemma. Journal of Chemical Physics 105:10123–10133.
 53. Sedlak, M., 1997. Dynamic light scattering from binary mixtures of polyelectrolytes. II. Appearance of the medium polyelectrolyte mode upon mixing and comparison with experiments on binary mixtures of neutral polymers. Journal of Chemical Physics

- 107:10805–10815.
54. Sedlak, M., and E. J. Amis, 1992. Concentration and Molecular-Weight Regime Diagram of Salt-Free Polyelectrolyte Solutions as Studied by Light-Scattering. *Journal of Chemical Physics* 96:826–834.
 55. Sedlak, M., and E. J. Amis, 1992. Dynamics of Moderately Concentrated Salt-Free Polyelectrolyte Solutions - Molecular-Weight Dependence. *Journal of Chemical Physics* 96:817–825.
 56. Bockstaller, M., W. Kohler, G. Wegner, D. Vlassopoulos, and G. Fytas, 2001. Levels of structure formation in aqueous solutions of anisotropic association colloids consisting of rodlike polyelectrolytes. *Macromolecules* 34:6359–6366.
 57. Pabit, S. A., S. P. Meisburger, L. Li, J. M. Blose, C. D. Jones, and L. Pollack, 2010. Counting Ions around DNA with Anomalous Small-Angle X-ray Scattering. *Journal of the American Chemical Society* 132:16334–16336.
 58. Andresen, K., X. Y. Qiu, S. A. Pabit, J. S. Lamb, H. Y. Park, L. W. Kwok, and L. Pollack, 2008. Mono- and trivalent ions around DNA: A small-angle scattering study of competition and interactions. *Biophysical Journal* 95:287–295.
 59. Pabit, S. A., K. D. Finkelstein, and L. Pollack, 2009. Using Anomalous Small Angle X-Ray Scattering to Probe the Ion Atmosphere around Nucleic Acids. *Methods in Enzymology, Vol 469: Biophysical, Chemical, and Functional Probes of Rna Structure, Interactions and Folding, Pt B* 469:391–410.
 60. Pabit, S. A., X. Y. Qiu, J. S. Lamb, L. Li, S. P. Meisburger, and L. Pollack, 2009. Both helix topology and counterion distribution contribute to the more effective charge screening in dsRNA compared with dsDNA. *Nucleic Acids Research* 37:3887–3896.
 61. Allahyarov, E., G. Gompper, and H. Löwen, 2005. DNA condensation and redissolution: interaction between overcharged DNA molecules. *Journal of Physics: Condensed Matter* 17:S1827. <http://dx.doi.org/10.1088/0953-8984/17/20/012>.
 62. Wu, J. Z., and Z. D. Li, 2007. Density-functional theory for complex fluids. *Annual Review of Physical Chemistry* 58:85–112.
 63. Cao, D. P., and J. Z. Wu, 2005. Microstructure of block copolymers near selective surfaces: Theoretical predictions and configurational-bias Monte Carlo simulation. *Macromolecules* 38:971–978.
 64. Marcus, Y., 1988. Ionic-Radii in Aqueous-Solutions. *Chemical Reviews* 88:1475–1498.
 65. Yu, Y. X., and J. Z. Wu, 2002. Structures of hard-sphere fluids from a modified fundamental-measure theory. *Journal of Chemical Physics* 117:10156–10164.
 66. Meng, D., B. Zheng, G. Lin, and M. L. Sushko, 2014. Numerical Solution of 3D Poisson-Nernst-Planck Equations Coupled with Classical Density Functional Theory for Modeling Ion and Electron Transport in a Confined Environment. *Communications in Computational Physics* 16:1298–1322.
 67. Horinek, D., A. Herz, L. Vrbka, F. Sedlmeier, S. I. Mamatkulov, and R. R. Netz, 2009. Specific ion adsorption at the air/water interface: The role of hydrophobic solvation. *Chemical Physics Letters* 479:173–183.
 68. Goel, T., C. N. Patra, S. K. Ghosh, and T. Mukherjee, 2008. Molecular solvent model of cylindrical electric double layers: A systematic study by Monte Carlo simulations and density functional theory. *Journal of Chemical Physics* 129.
 69. Goel, T., C. N. Patra, S. K. Ghosh, and T. Mukherjee, 2011. Effect of Ionic Size on the Structure of Cylindrical Electric Double Layers: A Systematic Study by Monte Carlo Simulations and Density Functional Theory. *Journal of Physical Chemistry B* 115:10903–10910.
 70. Patra, C. N., and A. Yethiraj, 1999. Density functional theory for the distribution of small ions around polyions. *Journal of Physical Chemistry B* 103:6080–6087.
 71. Poitevin, F., H. Orland, S. Doniach, P. Koehl, and M. Delarue, 2011. AquaSAXS: a web server for computation and fitting of SAXS profiles with non-uniformly hydrated atomic models. *Nucleic Acids Research* 39:W184–W189.
 72. Koehl, P., and M. Delarue, 2010. AQUASOL: An efficient solver for the dipolar Poisson-Boltzmann-Langevin equation. *Journal of Chemical Physics* 132:064101.
 73. Andresen, K., R. Das, H. Y. Park, H. Smith, L. W. Kwok, J. S. Lamb, E. J. Kirkland, D. Herschlag, K. D. Finkelstein, and L. Pollack, 2004. Spatial distribution of competing ions around DNA in solution. *Physical Review Letters* 93:248103.
 74. Braunlin, W. H., C. F. Anderson, and M. T. Record, 1987. Competitive interactions of Co(NH₃)₆(3+) and Na⁺ with helical B-DNA probed by ⁵⁹Co and ²³Na NMR. *Biochemistry* 26:7724–31. <http://www.ncbi.nlm.nih.gov/pubmed/3427101>.
 75. Robbins, T. J., J. D. Ziebarth, and Y. Wang, 2014. Comparison of Monovalent and Divalent Ion Distributions Around a DNA Duplex with Molecular Dynamics Simulation and A Poisson-Boltzmann Approach. *Biopolymers* 101:834–848.
 76. Tolokh, I. S., S. A. Pabit, A. M. Katz, Y. Chen, A. Drozdetski, N. Baker, L. Pollack, and A. V. Onufriev, 2014. Why double-stranded RNA resists condensation. *Nucleic Acids Research* 42:10823–10831.
 77. Grochowski, P., and J. Trylska, 2008. Continuum molecular electrostatics, salt effects, and counterion binding—a review of the Poisson-Boltzmann theory and its modifications. *Biopolymers* 89:93–113.
 78. Robbins, T. J., J. D. Ziebarth, and Y. Wang, 2014. Comparison of monovalent and divalent ion distributions around a DNA duplex with molecular dynamics simulation and a Poisson-Boltzmann approach. *Biopolymers* 101:834–848.
 79. Lightstone, F. C., E. Schwegler, R. Q. Hood, F. Gygi, and G. Galli, 2001. A first principles molecular dynamics simulation of the hydrated magnesium ion. *Chemical Physics Letters* 343:549–555.

List of Figures

- | | | |
|---|--|----|
| 1 | Macroion models used in classical DFT simulations: (a) Model of a cylinder with uniform axial charge density; (b) discrete charge model. | 2 |
| 2 | Ionic distributions around a uniformly charged cylinder. (a) Solutions of 100 mM NaCl, 100 mM RbCl, and 10 mM SrCl ₂ in a 1 mM NaCl buffer; concentration profiles are shown for Na ⁺ (solid line), Rb ⁺ (dashed line), and Sr ²⁺ (dot-dashed line). (b) Solutions of 5 mM CoHexCl ₃ in a 20 mM NaCl buffer; concentration profiles are shown for Na ⁺ (solid line), CoHex ³⁺ (dashed line), and Cl ⁻ (dot-dashed line). | 6 |
| 3 | Panoramic view of cation distributions around DNA in 100 mM RbCl calculated using (a) cDFT, and (b) and NLPB (solid line) and the cDFT model with no ion-correlation interactions (cDFT-nc, dotted line). Rb ⁺ distributions on the DNA strand are shown as black lines and in the minor groove as red lines with (c) a zoom-in into a low-density region for Rb ⁺ in plot (b). Na ⁺ distributions are shown in blue. (d) Radial distribution function of Rb ⁺ ions around DNA molecule. | 7 |
| 4 | Panoramic density distributions of Sr ²⁺ ions on DNA (a) strands and (b) minor grooves obtained using 3D cDFT (solid red line), NLPB (solid black line) and cDFT-nc (dotted line). SrCl ₂ concentration is 10 mM. (c) Radial distribution function of Sr ²⁺ ions around DNA molecule obtained using a complete cDFT model (solid line), cDFT model without ion-water attractive interactions (dotted line). | 8 |
| 5 | (a) Panoramic density distributions of CoHex ³⁺ ions on DNA strand obtained using 3D cDFT (solid red line), NLPB (solid black line; the curve is shifted up by 0.5 mM for clarity) and cDFT-nc (dotted line). CoHexCl ₃ concentration is 0.5 mM. (b) Radial distribution function of CoHex ³⁺ ions around DNA molecule. Note: as seen from the RDF, there is zero CoHex density in the minor groove, so the corresponding panoramic density is not shown as a separate plot. | 9 |
| 6 | Simulated and experimental ASAXS profiles for 25 bp DNA in (a) 100 mM RbCl and (b) 10 mM SrCl ₂ solutions. Experimental data (57, 58) are shown as black dots, simulations results obtained using 3D cDFT as blue lines, NLPB as green lines. Simulation results in (a) obtained using 1D cDFT and NLPB results, are shown as dotted blue and green lines, respectively. The red line in (b) corresponds to cDFT results obtained in the model with no ion-water attractive interactions. | 9 |
| 7 | ASAXS profiles for 0.5 mM CoHexCl ₃ DNA solutions. Experimental data are shown with a thick black line (unpublished) and those from Andresen et al (58) with a blue line. 1D cDFT results are shown as thin black line, 1D NLPB results as red line, 3D cDFT and NLPB data coincide and are shown as green line. | 10 |

List of Tables

- 1 A summary of the different computational models used in this paper to assess the influence of different energetic contributions (ion-ion electrostatic correlations, ion-ion steric correlations, ion-solvent interactions, and water structural changes) on DNA-ion distributions and compare the resulting distribution functions with experimental ASAXS data. The rows provide model descriptions while the columns indicate which physical phenomena are included by the models. 3