

Memory constraints for power-law series

Fangjian Guo,^{1,2} Zimo Yang,¹ Zhi-Dan Zhao,¹ and Tao Zhou^{1,3,*}

¹*CompleX Lab, Web Sciences Center, University of Electronic Science and Technology of China, Chengdu 611731, People's Republic of China*

²*Department of Computer Science, Duke University, Durham, NC 27708, USA*

³*Big Data Research Center, University of Electronic Science and Technology of China, Chengdu 611731, People's Republic of China*

Many time series produced by complex systems are empirically found to follow power-law distributions with different exponents α . By permuting the independently drawn samples from a power-law distribution, we present non-trivial bounds on the memory (1st-order autocorrelation) as a function of α , which are markedly different from the ordinary ± 1 bounds for Gaussian or uniform distributions. When $1 < \alpha \leq 3$, as α grows bigger, the upper bound increases from 0 to +1 while the lower bound remains 0; when $\alpha > 3$, the upper bound remains +1 while the lower bound descends below 0. Theoretical bounds agree well with numerical simulations. Based on the ratings of *MovieLens* and posts on *Twitter*, we also find that empirical power-law distributed data produced by human activities conform to such constraints. The present findings explain some observed constraints in bursty time series and scale-free networks, clear up possible misunderstanding from the ostensible values of some statistical measures like autocorrelation and assortativity coefficient, and challenge the validity of those measures in heterogeneous systems.

PACS numbers: 89.75.Da, 05.45.Tp, 89.75.-k, 02.50.-r

I. INTRODUCTION

Heterogeneity and memory are two remarkable features in characterizing time series produced by complex systems. The former quantifies the extent that the distribution of time series differs from a normal distribution. In particular, many real time series, denoted by $\{t_1, t_2, \dots\}$, are empirically found to follow power-law distributions $p(t) \sim t^{-\alpha}$ [1]. Examples include the magnitudes of earthquake [2], the intensity of wars [3], the severity of terrorist attacks [4], the strength of fluctuations in finance market [5], the time intervals between human-activated events [6], and so on. The latter characterizes the correlation between neighboring or nearby elements: a positive memory indicates that a high-value (low-value) element is likely to be followed by some high-value (low-value) elements. Memory effects are observed in a large variety of real-world time series [7], such as natural phenomena (e.g., earthquake and daily precipitation) [8], online human activities (e.g., visits on movies, books and music) [9], physiological signals (e.g., respiratory and liver cirrhosis time series) [10], and so on. Such memory effects also play a significant role in modeling spatial mobility [11–13] and temporal activities [14, 15] of human beings.

Here we adopt the simplest measure, first-order autocorrelation [8], to characterize the memory effect of a finite time series $\{t_1, t_2, \dots, t_n\}$, namely the Pearson's correlation coefficient between $\{t_1, t_2, \dots, t_{n-1}\}$

and $\{t_2, t_3, \dots, t_n\}$, defined as

$$M = \frac{1}{n-1} \sum_{i=1}^{n-1} \frac{(t_i - m_1)(t_{i+1} - m_2)}{\sigma_1 \sigma_2}, \quad (1)$$

where m_1, m_2 and σ_1, σ_2 refer to the means and standard deviations of the two series. In the recent studies [8, 13, 16, 17], the memory coefficient M was usually treated as an independent measure of the heterogeneity, with natural bounds $-1 \leq M \leq 1$. Such seemingly correct statement is challenged by the sensitivity of autocorrelation function to the fat tails in $p(t)$ [18], as well as related evidence on the nontrivial bounds of assortativity coefficients in networks with heterogeneous degree distributions [19].

In this paper, given the elements t_1, t_2, \dots, t_n , we calculate the maximum and minimum values of M under any permutation, suggesting the dependence between memory coefficient and series heterogeneity. In particular, given a power-law distribution $p(t) \sim t^{-\alpha}$, we show unreported nontrivial bounds of M in the thermodynamic limit — when $1 < \alpha \leq 3$, as α grows bigger, the upper bound increases from 0 to +1 while the lower bound remains 0; when $\alpha > 3$, the upper bound remains +1 while the lower bound descends below 0 but strictly above -1. The theoretical bounds agree very well with numerical simulations. In addition, according to the empirical analysis on *MovieLens* and *Twitter*, power-law distributed inter-event time series produced by human activities are found to conform to such bounds. Our findings add novel insights in characterizing not only heterogeneous time series, but also heterogeneous networks and other complex systems with heavy-tailed distributions.

The rest of this paper is organized as follows. In Section II, we will derive the theoretical bounds of M given

* zhutou@ustc.edu

$p(t) \sim t^{-\alpha}$. Such bounds will get validated based on extensive numerical tests in Section III and empirical results in Section IV. We will draw the main conclusion, discuss the relevance and implication of our findings in Section V.

II. THEORETICAL BOUNDS

Given a finite set of real numbers $\{t_1, t_2, \dots, t_n\}$ independently sampled from a certain distribution, without loss of generality, we order them as $t_1 \leq t_2 \leq \dots \leq t_n$, which are referred to as *order statistics*. Then, by applying a permutation θ (a one-to-one mapping from the set $\{1, 2, \dots, n\}$ to itself) to $\{t_i\}$, we have a new sequence $\{t_{\theta_1}, t_{\theta_2}, \dots, t_{\theta_n}\}$ with a different interdependence structure among elements. By changing θ , we can expect series with different values of M . For example, if series are permuted such that big elements tend to be followed by big ones and small followed by small ones, M would be positive; on the contrary, if big elements followed by small ones and small elements followed by big ones, M would be negative. Interestingly, there exists explicit θ_{\max} and θ_{\min} (though not unique) that respectively maximizes and minimizes M among all possible permutations. And we can use these two extremes to derive the bounds for memory in the sense of all permuted independent samples.

To see this, we need to find out how a permutation affects M . As the values of the two aforementioned series are different in only one element (t_1 in head and t_n in tail), we assume $m_1 = m_2 = m$ and $\sigma_1 = \sigma_2 = \sigma$ when n is large, where m and σ are the mean and standard deviation of the whole series. The memory of $\{t_{\theta_i}\}$ can hence be rewritten as

$$M(t_\theta) = \frac{1}{\sigma^2} \left(\frac{1}{n-1} \sum_{i=1}^{n-1} t_{\theta_i} t_{\theta_{i+1}} - m^2 \right), \quad (2)$$

where the reordering of the series only affects the summed products of adjacent terms

$$S_\theta = \sum_{i=1}^{n-1} t_{\theta_i} t_{\theta_{i+1}}, \quad (3)$$

while m and σ are invariant to permutations. The desired extreme permutations for M are just those maximize/minimize S_θ , denoted by θ_{\max} and θ_{\min} respectively. It has been shown [20] that, for any n , there are explicit solutions to θ_{\max} and θ_{\min} for any real numbers $t_1 \leq t_2 \leq \dots \leq t_n$. Notice that, whereas Hallin *et al.* [20] use an objective function that sums the products in a cycle (and therefore they call the problem *optimal Hamiltonian cycles*), i.e. $S'_\theta = \sum_{i=1}^{n-1} t_{\theta_i} t_{\theta_{i+1}} + t_{\theta_1} t_{\theta_n}$, the results can be reduced to our case by introducing an additional element $t_0 = 0$ to the series, which makes zero contribution to the sum.

θ_{\max} achieves the maximum memory by first arranging the odd elements of order statistics in the increasing order, followed by even elements in the decreasing order, which is

$$\begin{aligned} t_1, t_3, \dots, t_{2l-1}, t_{2l}, t_{2l-2}, \dots, t_4, t_2 \quad (n = 2l), \\ t_1, t_3, \dots, t_{2l-1}, t_{2l+1}, t_{2l}, \dots, t_4, t_2 \quad (n = 2l + 1). \end{aligned} \quad (4)$$

For simplicity, we only address the case when $n = 2l$, the sum with order statistics is expressed as

$$S_{\theta_{\max}} = \sum_{i=1}^{2l-2} t_i t_{i+2} + t_{2l} t_{2l-1} \quad (n = 2l), \quad (5)$$

while the case of $n = 2l + 1$ can be handled analogously.

On the contrary, θ_{\min} arranges the order statistics by alternating small and big terms, namely

$$\begin{aligned} t_{2l}, t_1, t_{2l-2}, \dots, t_{2l-3}, t_2, t_{2l-1} \quad (n = 2l), \\ t_{2l}, t_2, \dots, t_l, \dots, t_1, t_{2l+1} \quad (n = 2l + 1), \end{aligned} \quad (6)$$

where when n is even, even and odd terms are interlaced; when n is odd, half of the sequence is made up of even terms while the other half odd. Similarly, for even n , we have

$$S_{\theta_{\min}} = \sum_{i=1}^{l-1} (t_i t_{2l+1-i} + t_i t_{2l-1-i}) + t_l t_{l+1} \quad (n = 2l), \quad (7)$$

and the case of $n = 2l + 1$ is analogous.

We then define the upper bound and lower bound for memory as the expectation of M under θ_{\max} and θ_{\min} in the limit of infinitely long series. The bounds, defined as

$$M_{\max} = \mathbb{E} \lim_{n \rightarrow \infty} M(t_{\theta_{\max}}) \quad (8)$$

and

$$M_{\min} = \mathbb{E} \lim_{n \rightarrow \infty} M(t_{\theta_{\min}}), \quad (9)$$

measure the memory constraints imposed by a marginal distribution. As we will see, these bounds can be derived in a closed form or effectively approximated for several distributions.

It can be shown that $M_{\max} = 1$ and $M_{\min} = -1$ hold for uniform and Gaussian distributions, corresponding to the natural range of M (see details in *Appendix A* and *Appendix B*). However, a much narrower range is found for power-law distributions, where the bounds rely on the exponent α . While the rest of the paper focuses on power-law, we also notice a few other distributions with non-trivial memory constraints, which are discussed in *Appendix C*.

Supposing that the series $\{t_1, t_2, \dots, t_n\}$ are independently sampled from a power-law distribution with density

$$p(t) = (\alpha - 1)t^{-\alpha}. \quad (10)$$

To derive how the bounds rely on α , we first consider the case where $\alpha > 3$, which is necessary for the population variance $\sigma(\alpha)^2$ (appearing in the denominator of M) to converge. When $\alpha > 3$, by the strong law of large numbers and continuous mapping theorem, we have

$$M(t_\theta) \rightarrow \frac{1}{\sigma(\alpha)^2} \left(\lim_{n \rightarrow \infty} \frac{1}{n-1} \sum_{i=1}^{n-1} t_{\theta_i} t_{\theta_{i+1}} - m(\alpha)^2 \right)$$

almost surely as $n \rightarrow \infty$, where the sample moments are replaced by the corresponding population moments

$$m(\alpha) = \int_1^{+\infty} t p(t) dt = \frac{\alpha-1}{\alpha-2}, \quad (11)$$

and

$$\sigma(\alpha)^2 = \int_1^{+\infty} (t - m(\alpha))^2 p(t) dt = \frac{\alpha-1}{\alpha-3} - \left(\frac{\alpha-1}{\alpha-2} \right)^2. \quad (12)$$

Here we assume $x_{\min} = 1$ because M would remain the same if every t_i is divided by the same constant. Since $|M| \leq 1$, by Lebesgue's dominated convergence theorem, we can switch the order between limit and expectation and have

$$\mathbb{E} \lim_{n \rightarrow \infty} M(t_\theta) = \frac{1}{\sigma(\alpha)^2} \left(\lim_{n \rightarrow \infty} \frac{1}{n-1} \mathbb{E} S_\theta - m(\alpha)^2 \right). \quad (13)$$

Therefore, in the $\alpha > 3$ case, M_{\max} and M_{\min} can be respectively determined by $\frac{1}{n-1} \mathbb{E} S_{\theta_{\max}}$ and $\frac{1}{n-1} \mathbb{E} S_{\theta_{\min}}$ in the limit of large n .

According to Eq. (5), for the case of $n = 2l$, we have

$$\frac{1}{n-1} \mathbb{E} S_{\theta_{\max}} = \frac{1}{2l-1} \sum_{i=1}^{2l-2} \mathbb{E} t_i t_{i+2} + \frac{\mathbb{E} t_{2l} t_{2l-1}}{2l-1}, \quad (14)$$

whereas the case of $n = 2l + 1$ can be worked out in a similar fashion to arrive at the same result.

The expected value of each term can be obtained by using the joint distribution of order statistics. The probability density function for the joint distribution of two order statistics t_j, t_k ($j < k$) is given by [21]

$$\begin{aligned} p(t_j = x, t_k = y) &= n! \frac{[P(x)]^{j-1} [P(y) - P(x)]^{k-1-j}}{(j-1)! (k-1-j)!} \\ &\times \frac{[1 - P(y)]^{n-k}}{(n-k)!} p(x) p(y) \quad (x \leq y), \end{aligned} \quad (15)$$

where $P(x)$ is the corresponding cumulative distribution function for power-law, i.e.

$$P(x) = \int_1^x p(t) dt = 1 - x^{1-\alpha}.$$

Therefore, we have

$$\begin{aligned} \mathbb{E} t_i t_{i+2} &= \iint_{1 \leq x \leq y < \infty} x y p(t_i = x, t_{i+2} = y) dx dy \\ &= \frac{\Gamma(2l+1)}{\Gamma(2l+1-2c)} \frac{\Gamma(2l-i+1-2c)}{\Gamma(2l-i-1)} \\ &\times \frac{1}{(2l-i-c)(2l-i-\alpha c)} \end{aligned}$$

for $1 \leq i \leq 2l-2$ and the boundary term

$$\begin{aligned} \mathbb{E} t_{2l} t_{2l-1} &= \iint_{1 \leq x \leq y < \infty} x y f_{t_{(2l-1)}, t_{(2l)}}(x, y) dx dy \\ &= \frac{\alpha-1}{\alpha-2} \frac{\Gamma(2-2c)\Gamma(2l+1)}{\Gamma(2l+1-2c)}, \end{aligned}$$

where a shorthand $c = \frac{1}{\alpha-1}$ ($0 < c < \frac{1}{2}$) is adopted.

In the limit of $n \rightarrow \infty$, the first term in (14) would be

$$\begin{aligned} &\lim_{l \rightarrow \infty} \frac{1}{2l-1} \sum_{i=1}^{2l-2} \mathbb{E} t_i t_{i+2} \\ &= \lim_{l \rightarrow \infty} (2l+1)^{-(1-2c)} \sum_{k=2}^{2l-1} \frac{1}{(k-c)(k-c-1)} \frac{\Gamma(k+1-2c)}{\Gamma(k-1)} \\ &= \lim_{l \rightarrow \infty} \sum_{k=2}^{2l-1} \left(\frac{k}{2l+1} \right)^{-2c} \frac{1}{2l+1} = \int_0^1 t^{-2c} dt = \frac{\alpha-1}{\alpha-3}, \end{aligned} \quad (16)$$

where we have applied the property $\lim_{x \rightarrow \infty} \frac{\Gamma(x)}{\Gamma(x+\gamma)} = x^{-\gamma}$ for real γ , and rewrote the summation with $k = 2l - i$.

Meanwhile, the boundary term in the right hand side of Eq. (14) vanishes in the limit of large n , i.e.

$$\lim_{l \rightarrow \infty} \frac{1}{2l-1} \mathbb{E} t_{2l} t_{2l-1} = 0. \quad (17)$$

Substituting Eqs. (12), (13), (16) and (17) into (13), we arrive at

$$M_{\max} = \mathbb{E} \lim_{n \rightarrow \infty} M_{\theta_{\max}} = 1 \quad (\alpha > 3). \quad (18)$$

Similarly, to obtain the lower bound when $\alpha > 3$, we rewrite the summed products as

$$\begin{aligned} \frac{1}{n-1} \mathbb{E} S_{\theta_{\min}} &= \frac{1}{2l-1} \sum_{i=1}^{l-1} (\mathbb{E} t_i t_{2l+1-i} + \mathbb{E} t_i t_{2l-1-i}) + \\ &\frac{1}{2l-1} \mathbb{E} t_l t_{l+1}. \end{aligned}$$

Again, we assume $n = 2l$ for convenience, while the case of $n = 2l + 1$ gives the same result. Taking the large n

limit, we have

$$\begin{aligned}
& \lim_{l \rightarrow \infty} \frac{1}{2l-1} \sum_{i=1}^{l-1} \mathbb{E} t_i t_{2l+1-i} \\
&= \lim_{l \rightarrow \infty} (2l+1)^{-(1-2c)} \sum_{i=1}^{l-1} (i+2-c)^{-c} (2l-i+1-2c)^{-c} \\
&= \lim_{l \rightarrow \infty} \frac{1}{2l-1} \sum_{i=1}^{l-1} \left(\frac{i}{2l+1} + \frac{2-c}{2l+1} \right)^{-c} \left(1 - \frac{i}{2l+1} - \frac{2c}{2l+1} \right)^{-c} \\
&= \lim_{l \rightarrow \infty} \frac{1}{2l-1} \sum_{i=1}^{l-1} \left(\frac{i}{2l+1} \right)^{-c} \left(1 - \frac{i}{2l+1} \right)^{-c} \\
&= \int_0^{\frac{1}{2}} u^{-c} (1-u)^{-c} du = B\left(\frac{1}{2}; \frac{\alpha-2}{\alpha-1}, \frac{\alpha-2}{\alpha-1}\right),
\end{aligned}$$

where $u = i/(2l+1)$ and $B(\cdot)$ is the *incomplete beta function*

$$B(x; a, b) = \int_0^x t^{a-1} (1-t)^{b-1} dt.$$

The other term has the same limit, namely

$$\lim_{l \rightarrow \infty} \frac{1}{2l-1} \sum_{i=1}^{l-1} \mathbb{E} t_i t_{2l-1-i} = B\left(\frac{1}{2}; \frac{\alpha-2}{\alpha-1}, \frac{\alpha-2}{\alpha-1}\right),$$

and again the remaining term vanishes as

$$\lim_{l \rightarrow \infty} \frac{1}{2l-1} \mathbb{E} t_l t_{l+1} = 0.$$

Therefore, the minimum memory for $\alpha > 3$ is

$$M_{\min} = \frac{1}{\sigma(\alpha)^2} \left[2B\left(\frac{1}{2}; \frac{1}{m(\alpha)}, \frac{1}{m(\alpha)}\right) - m(\alpha)^2 \right], \quad (19)$$

where the population moments are given by Eqs. (11) and (12). Noticeably, M_{\min} is a decreasing function of α (see Figure 1) that approaches -0.65 as α gets large.

Having obtained the upper bound (18) and the lower bound (19) for $\alpha > 3$, we now turn to the case for $1 < \alpha \leq 3$ ($\alpha > 1$ is necessary for power-law to be normalized). In this case, the corresponding population moment for σ^2 would diverge ($m(\alpha)$ also diverges when $\alpha < 2$), rendering the moment-substitution technique infeasible. Meanwhile, it seems formidable to deal with the probability distributions of $M(t_{\theta_{\max}})$ and $M(t_{\theta_{\min}})$ directly. Therefore, we present an approximation method that recovers the asymptotic behavior of statistics when $n \rightarrow \infty$ by substituting random variables with deterministic surrogates.

To do so, we pick the points $\{\hat{t}_1, \hat{t}_2, \dots, \hat{t}_n\}$ that cut the area under the probability density function $p(t)$ into slices of equal area $1/n$, with $\hat{t}_1 = x_{\min} = 1$ and the area from \hat{t}_n extending to infinity also being $1/n$. Then we approximate the random samples $\{t_{(1)}, t_{(2)}, \dots, t_{(n)}\}$ with these deterministic points $\{\hat{t}_1, \hat{t}_2, \dots, \hat{t}_n\}$. It should




FIG. 1. (Color online) Theoretical bounds (black solid) compared to simulated sample mean (dots) of $M(t_{\theta_{\max}})$ and $M(t_{\theta_{\min}})$ under different sequence lengths $n = 10^3, 10^4, 10^6$ and 10^8 . Each dot is averaged over 1,000 independent draws of the sequence $\{t_i\}$ and the error bar denotes one standard deviation.

be noted that such approximation imposes a cut-off on the maximum value of $\{t_i\}$ and the probability of drawing a sample exceeding the cut-off is $1/n$, which diminishes to zero as $n \rightarrow \infty$. As $\int_1^{\hat{t}_i} p(t) dt = (i-1)/n$, we have

$$\hat{t}_i = \left(1 - \frac{i-1}{n}\right)^{-c} \quad (i = 1, 2, \dots, n), \quad (20)$$

where $c = 1/(\alpha-1) > \frac{1}{2}$ in this case.

Rewriting M in terms of samples as

$$M = \frac{s - m^2}{\bar{t}^2 - m^2}, \quad (21)$$

where $s = \frac{1}{n-1} S = \frac{1}{n-1} \sum_{i=1}^{n-1} t_i t_{i+1}$, $\bar{t}^2 = \frac{1}{n} \sum_{i=1}^n t_i^2$ and $m^2 = \left(\frac{1}{n} \sum_{i=1}^n t_i\right)^2$ are the statistics in concern. By substituting t_i with \hat{t}_i , we seek approximations for these statistics (denoted by \hat{s} , $\bar{\hat{t}}^2$ and \hat{m}^2) in the form of $n^{\mu(\alpha)} g(n, \alpha)$, where when $n \rightarrow \infty$, $g(n, \alpha)$ converges to a non-zero function of α while the divergence is characterized by the polynomial term $n^{\mu(\alpha)}$. This family of functions are denoted by $O(n^{\mu(\alpha)})$ generically. With some algebra, we have

$$\hat{m}^2 = \begin{cases} O(1) & (2 < \alpha < 3) \\ n^{2c-2} (\sum_{k=1}^n k^{-c})^2 = O(n^{2c-2}) & (1 < \alpha \leq 2) \end{cases}$$

and

$$\bar{\hat{t}}^2 = n^{2c-1} \sum_{k=1}^n k^{-2c} = O(n^{2c-1}),$$

which two hold for both θ_{\max} and θ_{\min} .

Then, for the upper bound, we have

$$\hat{s}_{\theta_{\max}} = \frac{n^{2c}}{n-1} \left[\sum_{k=2}^{n-1} (k^2 - 1)^{-c} + 2^{-c} \right], \quad (22)$$

which diverges with the order of $O(n^{2c-1})$. By comparing it with the order of \hat{m}^2 and \hat{t}^2 , in the limit of large n , we know $M_{\theta_{\max}}$ can be approximated by neglecting \hat{m}^2 . Therefore, for $1 < \alpha \leq 3$ we have

$$M_{\max} \approx \lim_{n \rightarrow \infty} \hat{M}_{\theta_{\max}} = \lim_{n \rightarrow \infty} \frac{\sum_{k=2}^{n-1} (k^2 - 1)^{-c} + 2^{-c}}{\sum_{k=1}^n k^{-2c}}, \quad (23)$$

where both the numerator and the denominator are convergent and can thus be approximately computed by taking a large n .

Meanwhile, supposing $n = 2l$ for convenience, we have

$$\hat{s}_{\theta_{\min}} = \frac{n^{2c}}{n-1} \left[\frac{1}{2} \sum_{i=1}^n [i(n+1-i)]^{-c} + \frac{1}{2} \sum_{i=1}^n [(i+2)(n+1-i)]^{-c} - (n+2)^{-c}/2 - (2n+2)^{-c}/2 \right].$$

By observing when $c > 1/2$

$$\sum_{i=1}^n [i(n+1-i)]^{-c} \leq \sum_{i=1}^n (n+1)^{-2c} = n(n+1)^{-2c} \rightarrow 0$$

and similarly for the other sum, we know $\hat{s}_{\theta_{\min}} = o(n^{2c-1})$, which diverges more slowly than \hat{t}^2 . Here $f(n) = o(g(n))$ means that $f(n)/g(n) \rightarrow 0$ in the limit of large n . Because the term with the biggest order only appears in the denominator, we have

$$M_{\min} \approx \lim_{n \rightarrow \infty} \hat{M}_{\theta_{\min}} = 0 \quad (24)$$

for $1 < \alpha \leq 3$. This non-negative constraint is particularly interesting because many power-law series are empirically found to have α in this region [22–41].




FIG. 2. (Color online) Memory and fitted α for power-law distributed inter-event time series produced by (a) rating on *MovieLens*, and (b) posting to *Twitter*. Each circle corresponds to a user. Theoretical bounds are drawn by blue solid line.

III. SIMULATIONS

In Figure 1, we compare theoretical bounds with simulated $M(t_{\theta_{\max}})$ and $M(t_{\theta_{\min}})$ under difference lengths n

of the sequence $\{t_1, t_2, \dots, t_n\}$. First, we observe that with a larger n , the theoretical bounds, defined as the expectation under infinite n , match the sample mean of $M(t_{\theta_{\max}})$ and $M(t_{\theta_{\min}})$ more closely. Second, in terms of accuracy, the effect of approximation is noticeable but satisfactory in the upper bound when $1 < \alpha \leq 3$; meanwhile, the lower bound and the upper bound for $\alpha > 3$ are both accurately predicted. Third, because $M(t_{\theta_{\max}})$ and $M(t_{\theta_{\min}})$ are random variables for each α , we quantify their variance around the mean in terms of standard deviation in the figure. Although our theoretical bounds do not predict their variance, we speculate that as n tends to infinity, $M(t_{\theta_{\min}})$ converges almost surely to the deterministic theoretical lower bounds; similarly, $M(t_{\theta_{\max}})$ converges to constant 1 for $\alpha > 3$ because $M \leq 1$. However, when $1 < \alpha \leq 3$, $M(t_{\theta_{\max}})$ converges to a non-degenerate random variable whose distribution depends on α , as evidenced by the non-vanishing error bars in that region.

IV. EMPIRICAL RESULTS

In this section, we examine the empirical distribution of (M, α) of power-law distributed sequences to see if real physical processes conform to the memory constraints predicted by our theory. To this end, we use inter-event time series collected from online human activities [38]. Inter-event time series refer to the series made up of time intervals between every two consecutive events and have been widely found to follow power-law distributions [6].

Figure 2 reports findings from *MovieLens* and *Twitter* datasets (each circle represents an individual user) compared to theoretical bounds (blue solid). Similar graphs have been proposed as a “phase diagram” in which different systems are grouped into different regions [8]. *MovieLens* is a website where users rate movies and get recommendations based on their ratings. The *MovieLens* 10M dataset collects 10^7 time stamps from 71,567 users when they rate a movie online (the collection of *MovieLens* data started years ago [45] and can be freely downloaded from <http://grouplens.org/>). Similarly, the *Twitter* dataset (a year-long subset of tweets crawled by Choudhury *et al.* [46], starting from Nov 2008) collects the time stamps from 9,832,781 tweets posted by 117,436 users. Therefore, the series correspond to the time intervals between two consecutive rating and tweeting respectively.

For comparison, we rule out the series in the datasets that are either too short ($n < 200$ for *MovieLens* and $n < 180$ for *Twitter*) or are unlikely to follow a power-law (p -value < 0.1). The p -value is computed from a goodness-of-fit test based on the Kolmogorov-Smirnov statistic as suggested by [1]. As a result, we have 2,261 series from *MovieLens* and 5,517 series from *Twitter*, of which α is estimated with maximum likelihood [1]. As can be seen from the plot, except for a few outliers, most of the sequences fall into the predicted region. It

is worth recapitulating that the predicted upper bound for $1 < \alpha < 3$ is an approximate to the mean, and the non-zero variance of $M(t_{\theta_{\max}})$ (see Figure 1) allows some points to exceed the predicted upper bound.

V. CONCLUSION AND DISCUSSION

In this article, we explore non-trivial bounds on the memory of power-law distributed series, which challenge the common treatment of memory as a measure independent of heterogeneity. We seek the bounds inside an ensemble formed by permuting independently and identically distributed power-law series, which covers a wide range of dependent structures while preserving the marginal distribution. Based on results in permutational extreme values, we present the bounds in either closed form or with an effective approximation.

Our bounding technique relies on a permutation ensemble of which the extremes are solvable. Although it does not subsume all possible power-law distributed series, we speculate that it is reasonably flexible that it covers most “natural” series. For example, series with $t_1 \sim p(x)$ and $t_1 = t_2 = \dots = t_n$ is an “unnatural” construction that is not contained by the ensemble. It is worth further investigation on the applicability of the permutation ensemble and the possibility of constructing other tractable ensembles with larger flexibility and fewer assumptions. Nevertheless, the usefulness of permutation ensemble is evidenced by empirical data (human activities in this case) found to closely conform to such bounds.

Without the knowledge of the present non-trivial bounds of memory, one may straightforwardly think that the values of M in the permutation ensemble are uniformly distributed in the range $[-1, 1]$ with average value being equal to 0. If so, when one observes a positive value of M as in many real human-activated systems [8], he will feel that the memory effect of this time series is stronger than the average of the permutation ensemble. Analogously, when one observes a positive assortativity coefficient [47] in a network, he will feel that the large-degree nodes tend to connect with other large-degree nodes, compared with the randomized networks with the same degree sequence. However, as indicated by the present results, the above intuition is incorrect. For example, for a power-law time series with exponent α in the range $(1, 3]$, the memory M is non-negative so that a small positive value of M is possibly still *smaller* than the average of the permutation ensemble. That is to say, given the values of elements in the time series, a high-value element is indeed more likely to be followed by some lower-value elements than the randomized counterpart even for some $M > 0$. Analogously, in a scale-free network [48] with power-law exponent γ in the range $(2, 4]$, the lower bound of assortativity coefficient is zero (details will be published elsewhere), hence in a real network with assortativity coefficient larger than zero, a large-degree

node may be more likely to connect with some smaller-degree nodes than the average in the corresponding null networks with the same degree sequence [49]. Therefore, bringing to light the existence of non-trivial bounds of M (and possibly other statistical measures to be revealed by future studies) in heterogeneous systems, this work could clear up some misunderstanding from the ostensible values of autocorrelation, assortativity coefficient, etc. Since many power-law series have α in the range $(1, 3]$ and many scale-free networks have γ in the range $(2, 4]$, our findings are relevant and significant to the academic society with interests in real complex systems.

Lastly, this work gives rise to a natural question, namely how to do proper statistics on heterogeneous systems, whose characteristic distributions may be mathematically ill-posed and/or have divergent finite moments, challenging the validity and usefulness of many commonly-used statistical measures. Given a certain statistical measure computed from a system, it would be important to know the relative position of this value compared to the distribution of the same measure computed from a “reference ensemble” of systems, such as the permutation ensemble in this case. However, such a task is usually difficult due to the computational complexity associated with the ensemble analysis. Therefore, unfolding the mathematical structure of reference ensembles could be very helpful, which is left as our future aim.

ACKNOWLEDGMENTS

The authors acknowledge valuable discussion with Chenmin Sun and Dan Yang. This work is partially supported by National Natural Science Foundation of China under Grants Nos. 61433014 and 11222543. T.Z. acknowledges the Program for New Century Excellent Talents in University under Grant No. NCET-11-0070, and Special Project of Sichuan Youth Science and Technology Innovation Research Team under Grant No. 2013TD0006.

Appendix A: Bounds for Uniform Distribution

Since $M(\cdot)$ is invariant to translation and scaling, without loss of generality, we assume that t_1, t_2, \dots, t_n are sampled from a uniform distribution in the range $(0, 1)$, namely $\{t_i\} \sim \text{Unif}(0, 1)$. We use the same technique as in the paper to show that the theoretical bounds for uniform distributed sequence are the ordinary ± 1 bounds.

Obviously $m = 1/2$ and $\sigma^2 = 1/12$, and the joint distribution for two order statistics is given by

$$p(t_j = x, t_k = y) = \frac{n!}{(j-1)!(k-j-1)!(n-k)!} \times x^{j-1}(y-x)^{k-j-1}(1-y)^{n-k},$$

for $1 \leq j < k \leq n$ and $x \leq y$.




FIG. 3. $M(t_{\theta_{\min}})$ simulated for different n , where $\{t_i\}$ are sampled from a Gaussian distribution. The dot represents sample mean from 1,000 independent draws and error bar represents one standard deviation.

M_{\max} is determined by

$$\lim_{n \rightarrow \infty} \frac{1}{n-1} S_{\theta_{\max}} = \lim_{l \rightarrow \infty} \frac{1}{2l-1} \sum_{i=1}^{2l-2} \mathbb{E} t_i t_{i+2}.$$

By plugging in

$$\begin{aligned} \mathbb{E} t_i t_{i+2} &= \iint_{0 \leq x \leq y \leq 1} xy f(t_i = x, t_{i+2} = y) dx dy \\ &= \frac{i(i+3)}{(2l+1)(2l+2)}, \end{aligned}$$

we have $\lim_{n \rightarrow \infty} \frac{1}{n-1} \mathbb{E} S_{\theta_{\max}} = 1/3$ and therefore

$$M_{\max} = \frac{1}{\sigma^2} \left(\lim_{n \rightarrow \infty} \frac{1}{n-1} \mathbb{E} S_{\theta_{\max}} - m^2 \right) = 1. \quad (\text{A1})$$

M_{\min} is determined by

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{1}{n-1} \mathbb{E} S_{\theta_{\min}} &= \lim_{l \rightarrow \infty} \frac{1}{2l-1} \sum_{i=1}^{l-1} \mathbb{E} t_i t_{2l+1-i} \\ &\quad + \lim_{l \rightarrow \infty} \frac{1}{2l-1} \sum_{i=1}^{l-1} \mathbb{E} t_i t_{2l-1-i}. \end{aligned}$$

Since

$$\begin{aligned} \mathbb{E} t_i t_{2l+1-i} &= \iint_{0 \leq x \leq y \leq 1} xy f(t_i = x, t_{2l+1-i} = y) dx dy \\ &= \frac{i(2l-i+2)}{(2l+1)(2l+2)}, \end{aligned}$$

we have $\lim_{l \rightarrow \infty} \frac{1}{2l-1} \sum_{i=1}^{l-1} \mathbb{E} t_i t_{2l+1-i} = \frac{1}{12}$ and similarly $\lim_{l \rightarrow \infty} \frac{1}{2l-1} \sum_{i=1}^{l-1} \mathbb{E} t_i t_{2l-1-i} = \frac{1}{12}$. Substituting these two results, we have $\lim_{n \rightarrow \infty} \frac{1}{n-1} \mathbb{E} S_{\theta_{\min}} = \frac{1}{6}$. Therefore, we arrive at

$$M_{\min} = \frac{1}{\sigma^2} \left(\lim_{n \rightarrow \infty} \frac{1}{n-1} \mathbb{E} S_{\theta_{\min}} - m^2 \right) = -1. \quad (\text{A2})$$

Appendix B: Bounds for Gaussian Distribution

In the case of Gaussian distributions, since the joint distribution in Eq. (15) does not have an analytical form, our previous technique that evaluates the expectation of $S_{\theta_{\max}}$ and $S_{\theta_{\min}}$ becomes infeasible. Therefore, we will instead derive the upper bound for Gaussian sequences with a bounding technique, and resort to simulations to observe the lower bound.

As $M(\cdot)$ is invariant to translation and scaling, we can consider $\{t_i\} \sim N(\mu, \sigma^2)$ for any μ and σ^2 . In particular, to get rid of the signs, we let μ depend on n such that μ is large enough to ensure all samples to be non-negative (μ grows like $O(\sqrt{\log n})$ [42]). Following Eq. (2), $M(t_{\theta_{\max}})$ is rewritten as

$$M(t_{\theta_{\max}}) = \frac{n}{n-1} \frac{S_{\theta_{\max}} - (n-1)m^2}{\sum_{i=1}^n t_i^2 - nm^2}, \quad (\text{B1})$$

where $m = \frac{1}{n} \sum_{i=1}^n t_i$. Assuming $n = 2l$, since all samples are non-negative, by the ordering $t_1 \leq t_2 \leq \dots \leq t_n$ we have

$$\begin{aligned} S_{\theta_{\max}} &= \sum_{i=1}^{2l-2} t_i t_{i+2} + t_{2l} t_{2l-1} \\ &\geq \sum_{i=1}^{2l-2} t_i^2 + t_{2l-1}^2 = \sum_{i=1}^n t_i^2 - t_n^2. \end{aligned}$$

Now dividing both numerator and denominator by $\sum_{i=1}^n t_i^2$ in Eq. (B1), we have

$$M(t_{\theta_{\max}}) \geq \frac{n}{n-1} \frac{1 - \frac{t_n^2}{\sum_{i=1}^n t_i^2} - \frac{n-1}{n} \frac{m^2}{\frac{1}{n} \sum_{i=1}^n t_i^2}}{1 - \frac{m^2}{\frac{1}{n} \sum_{i=1}^n t_i^2}}, \quad (\text{B2})$$

where in the right hand side, $\frac{m^2}{\frac{1}{n} \sum_{i=1}^n t_i^2} \rightarrow \frac{\mu^2}{\sigma^2 + \mu^2}$ almost surely. Also, it has been shown that the ratio $t_n^2 / \sum_{i=1}^n t_i^2$ between the largest element and the sum converges to 0 almost surely for independent sequence $\{t_i^2\}$ drawn from a common distribution with finite expectation [43, 44]. Hence, the right hand side of Eq. (B1) converges to 1 almost surely. Meanwhile, we know $M(t_{\theta_{\max}}) \leq 1$. Therefore, we know the bounded sequence $M(t_{\theta_{\max}}) \rightarrow 1$ almost surely and thus $M_{\max} = 1$.

We conjecture that $M(t_{\theta_{\min}}) \rightarrow -1$ almost surely also holds for the lower bound, although a proof is not obvious. In support of this result, Figure 3 presents the simulation for different lengths n . As n gets bigger, $M(t_{\theta_{\min}})$ converges to -1 with vanishing variance. Hence, we observe a tight lower bound $M_{\min} = -1$ for Gaussian distributions.

Appendix C: Non-trivial Bounds for Other Distributions

In this section, we present non-trivial memory constraints that we have found for several other distribu-




FIG. 4. Simulated $M(t_{\theta_{\max}})$ and $M(t_{\theta_{\min}})$ for various distributions also with non-trivial memory constraints: (a) Gamma distribution $p(x) \propto (x/\lambda)^{k-1} e^{-x/\lambda}$ with shape parameter k , (b) Student's t-distribution $p(x) \propto (1+x^2/\nu)^{-(\nu+1)/2}$ with degree of freedom ν , (c) Weibull distribution $p(x) \propto (x/\lambda)^{k-1} e^{-(x/\lambda)^k}$ with shape parameter k , and (d) Log-normal distribution $p(x) \propto x^{-1} e^{-\frac{(\ln x - \mu)^2}{2\sigma^2}}$ with scale parameter σ . Each dot is averaged over 1,000 independent draws with series length $n = 1,000$. Error bars represent one standard deviation.

tions, including (a) Gamma distribution, (b) Student's t-distribution, (c) Weibull distribution, and (d) log-normal distribution. In Figure 4, for each distribution, we plot simulated $M(t_{\theta_{\max}})$ and $M(t_{\theta_{\min}})$ against the parameter that controls the shape of the distribution. Gamma,

Weibull and log-normal distribution are also parameterized by a parameter that controls the scale (λ for Gamma and Weibull, μ for log-normal), which does not affect the memory.

-
- [1] A. Clauset, C. Shalizi, and M. E. J. Newman, *SIAM Rev.* **51**, 661 (2009).
- [2] V. F. Pisarenko and D. Sornette, *Pure Appl. Geophys.* **160**, 2343 (2003).
- [3] D. C. Roberts and D. L. Turcotte, *Fractals* **6**, 351 (1998).
- [4] A. Clauset, M. Young, and K. S. Gleditsch, *J. Conflict Resolution* **51**, 58 (2007).
- [5] B.-H. Wang and P. M. Hui, *Eur. Phys. J. B* **20**, 573 (2001).
- [6] A.-L. Barabási, *Nature (London)* **435**, 207 (2005).
- [7] W. Palma, *Long-Memory Time Series: Theory and Methods* (John Wiley & Sons, New Jersey, 2007).
- [8] K.-I. Goh and A.-L. Barabási, *EPL* **81**, 48002 (2008).
- [9] Z.-D. Zhao, S.-M. Cai, J. Huang, Y. Fu, and T. Zhou, *EPL* **100**, 48004 (2012).
- [10] A. H. Shirazi, *et al.*, *PLoS ONE* **8**, e72854 (2013).
- [11] M. Szell, R. Sinatra, G. Petri, S. Thurner, and V. Latora, *Sci. Rep.* **2**, 457 (2012).
- [12] J. Choi, J.-I. Sohn, K.-I. Goh, and I.-M. Kim, *EPL* **98**, 50001 (2012).
- [13] Z.-D. Zhao, Z. Yang, Z.-K. Zhang, T. Zhou, Z.-G. Huang, and Y.-C. Lai, *Sci. Rep.* **3**, 3472 (2013).
- [14] A. Vázquez, *Physica A* **373**, 747 (2007).
- [15] X.-P. Han, T. Zhou, and B.-H. Wang, *New J. Phys.* **10**, 073010 (2008).
- [16] Z.-D. Zhao, H. Xia, M.-S. Shang, and T. Zhou, *Chin. Phys. Lett.* **28**, 068901 (2011).
- [17] P. Wang, T. Zhou, X.-P. Han, and B.-H. Wang, *Physica A* **398**, 145 (2014).
- [18] M. Karsai, K. Kaski, A.-L. Barabási, and J. Kertész, *Sci. Rep.* **2**, 397 (2012).
- [19] S. Zhou and R. J. Mondragón, *New J. Phys.* **9**, 173 (2007).
- [20] M. Hallin, G. Melard, and X. Milhaud, *Annals of Statistics* **20**, 523 (1992).
- [21] H. A. David and H. N. Nagaraja, *Order Statistics* (Wiley, New Jersey, 2003, 3rd ed).
- [22] J.-G. Oliveira and A.-L. Barabási, *Nature (London)* **437**, 1251 (2005).
- [23] N. Eagle and A. Pentland, *Personal and Ubiquitous Computing* **10**, 255 (2005).
- [24] A. Vázquez, J.-G. Oliveira, Z. Dezső, K.-I. Goh, I. Kondor, and A.-L. Barabási, *Phys. Rev. E* **73**, 036127 (2006).
- [25] Z. Dezső, E. Almaas, A. Lukács, B. Rácz, I. Szakadát, and A.-L. Barabási, *Phys. Rev. E* **73**, 066132 (2006).
- [26] R. Lambiotte, M. Ausloos, and M. Thelwall, *J. Informetrics* **1**, 277 (2007).
- [27] T. Zhou, H. A. T. Kiet, B. J. Kim, B.-H. Wang, and P. Holme, *EPL* **82**, 28002 (2008).
- [28] S. C. Wang, J. J. Tseng, C. C. Tai, K. H. Lai, W. S. Wu, S. H. Chen, and S. P. Li, *Eur. Phys. J. B* **62**, 105 (2008).
- [29] N.-N. Li, N. Zhang, and T. Zhou, *Physica A* **387**, 6391 (2008).
- [30] B. Gonçalves and J. J. Ramasco, *Phys. Rev. E* **78**, 026123 (2008).
- [31] S. K. Baek, T. Y. Kim, and B. J. Kim, *Physica A* **387**, 3660 (2008).
- [32] W. Hong, X.-P. Han, T. Zhou, and B.-H. Wang, *Chin. Phys. Lett.* **26**, 028902 (2009).
- [33] F. Radicchi, *Phys. Rev. E* **80**, 026118 (2009).
- [34] Y. Wu, C. Zhou, J. Xiao, J. Kurths, and H. J. Schellnhuber, *Proc. Natl. Sci. Acad. U.S.A.* **107**, 18803 (2010).
- [35] P. Wang, X. Y. Xie, C. H. Yeung, and B.-H. Wang, *Physica A* **390**, 2395 (2011).
- [36] T. Takaguchi, M. Nakamura, N. Sato, K. Yano, and N. Masuda, *Phys. Rev. X* **1**, 011008 (2011).
- [37] T. Zhou, Z.-D. Zhao, Z. Yang, and C. Zhou, *EPL* **97**, 18006 (2012).
- [38] Z.-D. Zhao and T. Zhou, *Physica A* **391**, 3308 (2012).

- [39] D. Kondor, M. Pósfai, I. Csabai, and G. Vattay, PLoS ONE **9**, e86197 (2014).
- [40] S. Picoli, M. del Castillo-Mussot, H. V. Ribeiro, E. K. Lenzi, and R. S. Mendes, Sci. Rep. **4**, 4773 (2014).
- [41] L. Hou, X. Pan, Q. Guo, and J.-G. Liu, Sci. Rep. **4**, 6560 (2014).
- [42] S. Coles, J. Bawa, L. Trenner, and P. Dorazio, An introduction to statistical modeling of extreme values (Vol. 208) (Springer, London, 2001).
- [43] G. L. O'Brien, J. Appl. Prob. **17**, 539 (1980).
- [44] P. J. Downey, and P. E. Wright, Extremes **10**, 249 (2007).
- [45] P. Resnick, N. Iacovou, M. Suchak, P. Bergstrom, and J. Riedl, In *Proceedings of the 1994 ACM Conference on Computer Supported Cooperative Work* (ACM Press, New York, 1994), pp. 175-186.
- [46] M. De Choudhary, Y.-R. Lin, H. Sundaram, K.S. Candan, L. Xie, and A. Kelliher, In *Proceedings of the Fourth International AAAI Conference on Weblogs and Social Media* (ACM Press, New York, 2010), pp. 34-41.
- [47] M. E. J. Newman, Phys. Rev. Lett. **89**, 208701 (2002).
- [48] A.-L. Barabási and R. Albert, Science **286**, 509 (1999).
- [49] S. Maslov and K. Sneppen, Science **296**, 910 (2002).