

# $t$ -perfection in $P_5$ -free graphs

Henning Bruhn and Elke Fuchs

## Abstract

A graph is called  $t$ -perfect if its stable set polytope is fully described by non-negativity, edge and odd-cycle constraints. We characterise  $P_5$ -free  $t$ -perfect graphs in terms of forbidden  $t$ -minors. Moreover, we show that  $P_5$ -free  $t$ -perfect graphs can always be coloured with three colours, and that they can be recognised in polynomial time.

## 1 Introduction

There are three quite different views on *perfect graphs*, a view in terms of colouring, a polyhedral and a structural view. Perfect graphs can be seen as:

- the graphs for which the chromatic number  $\chi(H)$  always equals the clique number  $\omega(H)$ , and that in any induced subgraph  $H$ ;
- the graphs for which the *stable set polytope*, the convex hull of stable sets, is fully described by non-negativity and clique constraints; and
- the graphs that do not contain any *odd hole* (an induced cycle of odd length at least 5) or their complements, *odd antiholes*.

(The polyhedral characterisation is due to Fulkerson [15] and Chvátal [9], while the third item, the strong perfect graph theorem, was proved by Chudnovsky, Robertson, Seymour and Thomas [6].)

In this article, we work towards a similar threefold view on  $t$ -perfect graphs. These are graphs that, similar to perfect graphs, have a particularly simple stable set polytope. For a graph to be  $t$ -perfect its stable set polytope needs to be given by non-negativity, edge and odd-cycle constraints; for precise definitions we defer to the next section. The concept of  $t$ -perfection, due to Chvátal [9], thus takes its motivation from the polyhedral aspect of perfect graphs. The corresponding colouring and structural view, however, is still missing. For some graph classes, though, claw-free graphs for instance [5], the list of minimal obstructions for  $t$ -perfection is known. We extend this list to  $P_5$ -free graphs. (A graph is  $P_5$ -free if it does not contain the path on five vertices as an induced subgraph.)

Perfection is preserved under vertex deletion, and the same is true for  $t$ -perfection. There is a second simple operation that maintains  $t$ -perfection: a  $t$ -contraction, which is only allowed at a vertex with stable neighbourhood, contracts all the incident edges. Any graph obtained by a sequence of vertex deletions and  $t$ -contractions is a  $t$ -minor. The concept of  $t$ -minors makes it more convenient to characterise  $t$ -perfection in certain graph classes as it allows for more succinct lists of obstructions.

For that characterisation denote by  $C_n^k$  the  $k$ th power of the  $n$ -cycle  $C_n$ , that is, the graph obtained from  $C_n$  by adding an edge between any two vertices of distance at most  $k$  in  $C_n$ . We, moreover, write  $\overline{G}$  for the complement of a graph  $G$ , and  $K_n$  for the complete graph on  $n$  vertices and  $W_n$  for the wheel with  $n + 1$  vertices.

**Theorem 1.** *Let  $G$  be a  $P_5$ -free graph. Then  $G$  is  $t$ -perfect if and only if it does not contain any of  $K_4$ ,  $W_5$ ,  $C_7^2$ ,  $\overline{C_{10}^2}$  or  $\overline{C_{13}^3}$  as a  $t$ -minor.*

This answers a question of Benchetrit [2, p. 76].




Figure 1: Forbidden  $t$ -minors in  $P_5$ -free graphs

The forbidden graphs of the theorem are *minimally  $t$ -imperfect*, in the sense that they are  $t$ -imperfect but any of their proper  $t$ -minors are  $t$ -perfect. Odd wheels, even Möbius ladders (see Section 3), the cycle power  $C_7^2$  and the graph  $\overline{C_{10}^2}$  are known to be minimally  $t$ -imperfect. The graph  $\overline{C_{13}^3}$  appears here for the first time as a minimally  $t$ -imperfect graph. We prove this in Section 4, where we also present two more minimally  $t$ -imperfect graphs.

A starting point for Theorem 1 was the observation of Benchetrit [2, p. 75] that  $t$ -minors of  $P_5$ -free graphs are again  $P_5$ -free. Thus, any occurring minimally  $t$ -imperfect graph will be  $P_5$ -free, too. This helped to whittle down the list of prospective forbidden  $t$ -minors. We prove Theorem 1 in Sections 5 and 6.

A graph class in which  $t$ -perfection is quite well understood is the class of *near-bipartite* graphs; these are the graphs that become bipartite whenever the neighbourhood of any vertex is deleted. In the course of the proof of Theorem 1 we make use of results of Shepherd [26] and of Holm, Torres and Wagler [20]: together they yield a description of  $t$ -perfect near-bipartite graphs in terms of forbidden induced subgraphs. We discuss this in Section 3.

As a by-product of the proof of Theorem 1 we also obtain a polynomial-time algorithm to check for  $t$ -perfection in  $P_5$ -free graphs (Theorem 20).

Finally, in Section 7, we turn to the third defining aspect of perfect graphs: colouring. Shepherd and Sebő conjectured that every  $t$ -perfect graph can be coloured with four colours, which would be tight. For  $t$ -perfect  $P_5$ -graphs we show (Theorem 23) that already three colours suffice. We, furthermore, offer a conjecture that would, if true, characterise  $t$ -perfect graphs in terms of (fractional) colouring, in a way that is quite similar as for perfect graphs.

We end the introduction with a brief discussion of the literature on  $t$ -perfect graphs. A general treatment may be found in Grötschel, Lovász and Schrijver [19, Ch. 9.1] as well as in Schrijver [25, Ch. 68]. The most comprehensive source of literature references is surely the PhD thesis of Benchetrit [2]. A part of the literature is devoted to proving  $t$ -perfection for certain graph classes. For instance, Boulala and Uhry [3] established the  $t$ -perfection of series-parallel graphs. Gerards [16] extended this to graphs that do not contain an *odd- $K_4$*  as a subgraph (an *odd- $K_4$*  is a subdivision of  $K_4$  in which every triangle becomes an odd circuit). Gerards and Shepherd [17] characterised the graphs with all subgraphs  $t$ -perfect, while Barahona and Mahjoub [1] described the  $t$ -imperfect subdivisions of  $K_4$ . Wagler [29] gave a complete description of the stable set polytope of antiwebs, the complements of cycle powers. These are near-bipartite graphs that also play a prominent role in the proof of Theorem 1. See also Wagler [30] for an extension to a more general class of near-bipartite graphs. The complements of near-bipartite graphs are the quasi-line graphs. Chudnovsky and Seymour [8], and Eisenbrand, Oriolo, Stauffer and Ventura [12] determined the precise structure of the stable set polytope of quasi-line graphs. Previously, this was a conjecture of Ben Rebea [24].

Algorithmic aspects of  $t$ -perfection were also studied: Grötschel, Lovász and Schrijver [18] showed that the max-weight stable set problem can be solved in polynomial-time in  $t$ -perfect graphs. Eisenbrand et al. [11] found a combinatorial algorithm for the unweighted case.

## 2 Definitions

All the graphs in this article are finite, simple and do not have parallel edges or loops. In general, we follow the notation of Diestel [10], where also any missing elementary facts about graphs may be found.

Let  $G = (V, E)$  be a graph. The *stable set polytope*  $\text{SSP}(G) \subseteq \mathbb{R}^V$  of  $G$  is defined as the convex hull of the characteristic vectors of stable, i.e. independent, subsets of  $V$ . The characteristic vector of a subset  $S$  of the set  $V$  is the vector  $\chi_S \in \{0, 1\}^V$  with  $\chi_S(v) = 1$  if  $v \in S$  and 0 otherwise. We define a second polytope  $\text{TSTAB}(G) \subseteq \mathbb{R}^V$  for  $G$ , given by

$$\begin{aligned} x &\geq 0, \\ x_u + x_v &\leq 1 \text{ for every edge } uv \in E, \\ \sum_{v \in V(C)} x_v &\leq \left\lfloor \frac{|C|}{2} \right\rfloor \text{ for every induced odd cycle } C \text{ in } G. \end{aligned}$$

These inequalities are respectively known as non-negativity, edge and odd-cycle inequalities. Clearly,  $\text{SSP}(G) \subseteq \text{TSTAB}(G)$ .

Then, the graph  $G$  is called  *$t$ -perfect* if  $\text{SSP}(G)$  and  $\text{TSTAB}(G)$  coincide. Equivalently,  $G$  is  $t$ -perfect if and only if  $\text{TSTAB}(G)$  is an integral polytope, i.e. if all its vertices are integral vectors. It is easy to see that bipartite graphs are  $t$ -perfect. The smallest  *$t$ -imperfect* graph is  $K_4$ . Indeed, the vector  $\frac{1}{3}\mathbf{1}$  lies in  $\text{TSTAB}(K_4)$  but not in  $\text{SSP}(K_4)$ .

It is easy to verify that vertex deletion preserves  $t$ -perfection. Another operation that keeps  $t$ -perfection was found by Gerards and Shepherd [17]: whenever there is a vertex  $v$ , so that its neighbourhood is stable, we may contract all

edges incident with  $v$  simultaneously. We will call this operation a  $t$ -contraction at  $v$ . Any graph that is obtained from  $G$  by a sequence of vertex deletions and  $t$ -contractions is a  $t$ -minor of  $G$ . Let us point out that any  $t$ -minor of a  $t$ -perfect graph is again  $t$ -perfect.

### 3 $t$ -perfection in near-bipartite graphs

Part of the proof of Theorem 1 consists in a reduction to *near-bipartite* graphs. A graph is near-bipartite if it becomes bipartite whenever the neighbourhood of any of its vertices is deleted. We will need a characterisation of  $t$ -perfect near-bipartite graphs in terms of forbidden induced subgraphs. Fortunately, such a characterisation follows immediately from results of Shepherd [26] and of Holm, Torres and Wagler [20].

We need a bit of notation. Examples of near-bipartite graphs are antiwebs: an *antiweb*  $\overline{C}_n^k$  is the complement of the  $k$ th power of the  $n$ -cycle  $C_n$ . The antiweb is *prime* if  $n \geq 2k + 2$  and  $k + 1, n$  are relatively prime. We simplify the notation for antiwebs  $\overline{C}_n^k$  slightly by writing  $A_n^k$  instead. *Even Möbius ladders*, the graphs  $A_{4t+4}^{2t}$ , are prime antiwebs; see Figure 2 for the Möbius ladder  $\overline{C}_8^2$ . We view  $K_4$  alternatively as the smallest odd wheel  $W_3$  or as the smallest even Möbius ladder  $\overline{C}_4^0$ . Trotter [27] found that prime antiwebs give rise to facets in the stable set polytope—we only need that prime antiwebs other than odd cycles are  $t$ -imperfect, a fact that is easier to check.




Figure 2: Two views of the Möbius ladder on 8 vertices

Shepherd proved:

**Theorem 2** (Shepherd [26]). *Let  $G$  be a near-bipartite graph. Then  $G$  is  $t$ -perfect if and only if*

- (i)  $G$  contains no induced odd wheel; and
- (ii)  $G$  contains no induced prime antiweb other than possibly an odd hole.

Holm, Torres and Wagler [20] gave a neat characterisation of  $t$ -perfect antiwebs. For us, however, a direct implication of the proof of that characterisation is more interesting: an antiweb is  $t$ -perfect if and only if it does not contain any even Möbius ladder, or any of  $A_7^1, A_{10}^2, A_{13}^3, A_{13}^4, A_{17}^4$  and  $A_{19}^7$  as an induced subgraph. We may omit  $A_{17}^4$  from that list as it contains an induced  $A_{13}^3$ . Combining the theorem of Holm et al. with Theorem 2 one obtains:

**Proposition 3.** *A near-bipartite graph is  $t$ -perfect if and only if it does not contain any odd wheel, any even Möbius ladder, or any of  $A_7^1$ ,  $A_{10}^2$ ,  $A_{13}^3$ ,  $A_{13}^4$  and  $A_{19}^7$  as an induced subgraph.*

## 4 Minimally $t$ -imperfect antiwebs

For any characterisation of  $t$ -perfection in *minimally  $t$ -imperfect*, that is, all graphs that are  $t$ -imperfect but whose proper  $t$ -minors are  $t$ -perfect. Even Möbius ladders and odd wheels, for instance, are known to be minimally  $t$ -imperfect. This follows from the result of Fonlupt and Uhry [14] that *almost bipartite* graphs are  $t$ -perfect; a graph is almost bipartite if it contains a vertex whose deletion renders it bipartite. It is easy to check that any proper  $t$ -minor of an even Möbius ladder or an odd wheel is almost bipartite.

All the other forbidden  $t$ -minors in Theorem 1 or Proposition 3 are minimally  $t$ -imperfect, too. That  $C_7^2$  is minimally  $t$ -imperfect is proved in [5]. There, also minimality for  $C_{10}^2$  is shown, which allows us to verify that  $A_{10}^2$  is minimally  $t$ -imperfect as well. Indeed, for this we first observe that  $A_{10}^2$  can be obtained from  $C_{10}^2$  by adding diagonals of the underlying 10-cycle. The second necessary observation is that any two vertices directly opposite in the 10-cycle form a so called *odd pair*: any induced path between them has odd length. Minimality now follows from the result of Fonlupt and Hadjar [13] that adding an edge between the vertices of an odd pair preserves  $t$ -perfection.

In this section, we prove that  $A_{13}^3$ ,  $A_{13}^4$  and  $A_{19}^7$  are minimally  $t$ -imperfect, which was not observed before. As prime antiwebs these are  $t$ -imperfect. This follows from Theorem 2 but can also be seen directly by observing that the vector  $x \equiv \frac{1}{3}$  lies in TSTAB but not in SSP for any of the three graphs.

To show that the graphs are *minimally  $t$ -imperfect*, it suffices to consider the  $t$ -minors obtained from a single vertex deletion or from a single  $t$ -contraction. If these are  $t$ -perfect then the antiweb is minimally  $t$ -imperfect.

Trotter gave necessary and sufficient conditions when an antiweb contains another antiweb:

**Theorem 4** (Trotter [27]).  *$A_{n'}^{k'}$  is an induced subgraph of  $A_n^k$  if and only if*

$$n(k' + 1) \geq n'(k + 1) \text{ and } nk' \leq n'k.$$

We fix the vertex set of any antiweb  $A_n^k$  to be  $\{0, 1, \dots, n - 1\}$ , so that  $ij$  is an edge of  $A_n^k$  if and only if  $|i - j| \bmod n > k$ .

**Proposition 5.** *The antiweb  $A_{13}^3$  is minimally  $t$ -imperfect.*

*Proof.* For  $A_{13}^3$  to be minimally  $t$ -imperfect, every proper  $t$ -minor  $A_{13}^3$  needs to be  $t$ -perfect. As no vertex of  $A_{13}^3$  has a stable neighbourhood, any proper  $t$ -minor is a  $t$ -minor of a proper induced subgraph  $H$  of  $A_{13}^3$ . Thus, it suffices to show that any such  $H$  is  $t$ -perfect.

By Proposition 3,  $H$  is  $t$ -perfect unless it contains an odd wheel or one of  $A_7^1$ ,  $A_8^2$  or  $A_{10}^2$  as an induced subgraph. Since the neighbourhood of every vertex is stable,  $H$  cannot contain any wheel. For the other graphs, we check the inequalities of Theorem 4 and see that none can be contained in  $H$ . Thus,  $H$  is  $t$ -perfect and  $A_{13}^3$  therefore minimally  $t$ -imperfect.  $\square$




Figure 3: Antiweb  $A_{13}^4$ , and its  $t$ -minor obtained by a  $t$ -contraction at 0

**Proposition 6.** *The antiweb  $A_{13}^4$  is minimally  $t$ -imperfect.*

*Proof.* By Proposition 3, any proper induced subgraph of  $A_{13}^4$  that is  $t$ -imperfect contains one of  $A_7^1$ ,  $A_8^2$ , or  $A_{10}^2$  as an induced subgraph; note that  $A_{13}^4$  does not contain odd wheels. However, routine calculation and Theorem 4 show that  $A_{13}^4$  contains neither of these. Therefore, deleting any vertex in  $A_{13}^4$  always results in a  $t$ -perfect graph.

It remains to consider the graphs obtained from  $A_{13}^4$  by a single  $t$ -contraction. By symmetry, it suffices check whether the graph  $H$  obtained by  $t$ -contraction at 0 is  $t$ -perfect; see Figure 3. Denote by  $\tilde{0}$  the new vertex that resulted from the contraction.

The graph  $H$  is still near-bipartite and still devoid of odd wheels. Thus, by Proposition 3, it is  $t$ -perfect unless it contains  $A_7^1$  and  $A_8^2$  as an induced subgraph—all the  $t$ -imperfect antiwebs of Proposition 3 are too large for the nine-vertex graph  $H$ .

Now,  $A_7^1$  is 4-regular but  $H$  only contains five vertices of degree at least 4. Similarly,  $A_8^2$  is 3-regular but two of the nine vertices of  $H$ , namely 1 and 12, have degree 2. We see that neither of the two antiwebs can be contained in  $H$ , so that  $H$  is  $t$ -perfect and, thus,  $A_{13}^4$  minimally  $t$ -imperfect.  $\square$

**Proposition 7.** *The antiweb  $A_{19}^7$  is minimally  $t$ -imperfect.*

*Proof.* We claim that any proper induced subgraph of  $A_{19}^7$  is  $t$ -perfect. Indeed, as  $A_{19}^7$  does not contain any induced odd wheel, this follows from Proposition 3, unless  $A_{19}^7$  contains one of  $A_7^1$ ,  $A_8^2$ ,  $A_{10}^2$ ,  $A_{12}^4$ ,  $A_{13}^3$ ,  $A_{13}^4$ , or  $A_{16}^6$  as an induced subgraph. We can easily verify with Theorem 4 that this is not the case.

It remains to check that any  $t$ -contraction in  $A_{19}^7$  yields a  $t$ -perfect graph, too. By symmetry, we may restrict ourselves to a  $t$ -contraction at the vertex 0. Let  $H$  be the resulting graph, and let  $\tilde{0}$  be the new vertex; see Figure 4.

The graph  $H$  is a near-bipartite graph on 15 vertices. It does not contain any odd wheel as an induced subgraph. Thus, by Proposition 3,  $H$  is  $t$ -perfect unless it has an induced subgraph  $A$  that is isomorphic to a graph in

$$\mathcal{A} := \{A_7^1, A_8^2, A_{10}^2, A_{12}^4, A_{13}^3, A_{13}^4\}.$$




Figure 4: Antiweb  $A_{19}^7$ , and its  $t$ -minor obtained by  $t$ -contraction at 0

Since this is not the case for  $A_{19}^7$ , we may assume that  $\tilde{0} \in V(A)$ .

Note that the graphs  $A_7^1$ ,  $A_{10}^2$ ,  $A_{13}^3$  and  $A_{13}^4$  have minimum degree at least 4. Yet,  $\tilde{0}$  has only two neighbours of degree 4 or more (namely, 3 and 16). Thus, neither of these four antiwebs can occur as an induced subgraph in  $H$ .

It remains to consider the case when  $H$  contains an induced subgraph  $A$  that is isomorphic to  $A_8^2$  or to  $A_{12}^4$ , both of which are 3-regular graphs. In particular,  $A$  is then contained in  $H' = H - \{1, 18\}$  as the vertices 1 and 18 have degree 2.

As  $H'$  has only 13 vertices,  $A$  cannot be isomorphic to  $A_{12}^4$  since deleting any single vertex of  $H'$  never yields a 3-regular graph. That leaves only  $A = A_8^2$ .

Since  $A_8^2$  is 3-regular, we need to delete exactly one of the four neighbours of  $\tilde{0}$  in  $H'$ . Suppose this is the vertex 3. Then, 12 has degree 2 and thus cannot be part of  $A$ . Deleting 12 as well leads to vertex 2 having degree 2, which thereby is also excluded from  $A$ . This, however, is impossible as 2 is one of the three remaining neighbours of  $\tilde{0}$ .

By symmetry, we may therefore assume that the neighbours of  $\tilde{0}$  in  $A$  are precisely 2, 3, 16. That 17 is not part of  $A$  entails that the vertex 7 has degree 2 and thus cannot lie in  $A$  either. Then, however,  $16 \in V(A)$  has degree 2 as well, which is impossible.  $\square$

## 5 Harmonious cutsets

We investigate the structure of minimally  $t$ -imperfect graphs, whether they are  $P_5$ -free or not. We hope this more general setting might prove useful in subsequent research.

A structural feature that may never appear in a minimally  $t$ -imperfect graph  $G$  is a *clique separator*: any clique  $K$  of  $G$  so that  $G - K$  is not connected.

**Lemma 8** (Chvátal [9]; Gerards [16]). *No minimally  $t$ -imperfect graph contains a clique separator.*

A generalisation of clique separators was introduced by Chudnovsky et al. [7] in the context of colouring  $K_4$ -free graphs without odd holes. A tuple  $(X_1, \dots, X_s)$  of disjoint subsets of the vertex set of a graph  $G$  is  *$G$ -harmonious* if

- any induced path with one endvertex in  $X_i$  and the other in  $X_j$  has even length if and only if  $i = j$ ; and
- if  $s \geq 3$  then  $X_1, \dots, X_s$  are pairwise complete to each other.

A pair of subgraphs  $\{G_1, G_2\}$  of  $G = (V, E)$  is a *separation* of  $G$  if  $V(G_1) \cup V(G_2) = V$  and  $G$  has no edge between  $V(G_1) \setminus V(G_2)$  and  $V(G_2) \setminus V(G_1)$ . If both  $V(G_1) \setminus V(G_2)$  and  $V(G_2) \setminus V(G_1)$  are non-empty, the separation is *proper*.

A vertex set  $X$  is called a *harmonious cutset* if there is a proper separation  $(G_1, G_2)$  of  $G$  so that  $X = V(G_1) \cap V(G_2)$  and if there exists a partition  $X = (X_1, \dots, X_s)$  so that  $(X_1, \dots, X_s)$  is  $G$ -harmonious.

We prove:

**Lemma 9.** *If a  $t$ -imperfect graph contains a harmonious cutset then it also contains a proper induced subgraph that is  $t$ -imperfect. In particular, no minimally  $t$ -imperfect graph admits a harmonious cutset.*

For the proof we need a bit of preparation.

**Lemma 10.** *Let  $S_1 \subsetneq \dots \subsetneq S_k$  and  $T_1 \subsetneq \dots \subsetneq T_\ell$  be nested subsets of a finite set  $V$ . Let  $\sigma := \sum_{i=1}^k \lambda_i \chi_{S_i}$  and  $\tau := \sum_{j=1}^\ell \mu_j \chi_{T_j}$  be two convex combinations in  $\mathbb{R}^V$  with non-zero coefficients. If  $\sigma = \tau$  then  $k = \ell$ ,  $\lambda_i = \mu_i$  and  $S_i = T_i$  for all  $i = 1, \dots, k$ .*

The lemma is not new. It appears in the context of submodular functions, where it may be seen to assert that the *Lovász extension* of a set-function is well-defined; see Lovász [21]. For the sake of completeness, we give a proof here.

*Proof.* By allowing  $\lambda_1$  and  $\mu_1$  to be 0, we may clearly assume that  $S_1 = \emptyset = T_1$ . Moreover, if two elements  $u, v \in V$  always appear together in the sets  $S_i, T_j$  then we may omit one of  $u, v$  from all the sets. So, in particular, we may assume  $S_2$  and  $T_2$  to be singleton-sets.

Let  $s$  be the unique element of  $S_2$ . Then  $\sum_{i=2}^k \lambda_i = \sigma_s = \tau_s \leq \sum_{j=2}^\ell \mu_j$ . By symmetry, we also get  $\sum_{i=2}^k \lambda_i \geq \sum_{j=2}^\ell \mu_j$ , and thus we have equality. We deduce that  $T_2 = \{s\}$ , and that  $\lambda_1 = \mu_1$  as  $\lambda_1 = 1 - \sum_{i=2}^k \lambda_i = 1 - \sum_{j=2}^\ell \mu_j = \mu_1$ . Then

$$(\lambda_1 + \lambda_2) \chi_{S_1} + \sum_{i=3}^k \lambda_i \chi_{S_i \setminus \{s\}} = (\mu_1 + \mu_2) \chi_{T_1} + \sum_{j=3}^\ell \mu_j \chi_{T_j \setminus \{s\}}$$

are two convex combinations. Induction on  $|S_k|$  now finishes the proof, where we also use that  $\lambda_1 = \mu_1$ .  $\square$

**Lemma 11.** *Let  $G$  be a graph, and let  $(X, Y)$  be a  $G$ -harmonious tuple (with possibly  $X = \emptyset$  or  $Y = \emptyset$ ). If  $S_1, \dots, S_k$  are stable sets then there are stable sets  $S'_1, \dots, S'_k$  so that*

- (i)  $S'_1 \cap X \subseteq \dots \subseteq S'_k \cap X$ ;
- (ii)  $S'_1 \cap Y \supseteq \dots \supseteq S'_k \cap Y$ ; and
- (iii)  $\sum_{i=1}^k \chi_{S'_i} = \sum_{i=1}^k \chi_{S_i}$ .



*Proof.* We start with two easy claims. First:

*For any two stable sets  $S, T$  there are stable sets  $S'$  and  $T'$  such that  $\chi_S + \chi_T = \chi_{S'} + \chi_{T'}$  and  $S' \cap X \subseteq T' \cap X$ .* (1)

Indeed, assume there is an  $x \in (S \cap X) \setminus T$ . Denote by  $K$  the component of the induced graph  $G[S \cup T]$  that contains  $x$ , and consider the symmetric differences  $\tilde{S} = S \Delta K$  and  $\tilde{T} = T \Delta K$ . Clearly,  $\chi_S + \chi_T = \chi_{\tilde{S}} + \chi_{\tilde{T}}$ . Moreover,  $K$  meets  $X$  only in  $S$  as otherwise  $K$  would contain an induced  $x$ – $(T \cap X)$  path, which then has necessarily odd length. This, however, is impossible as  $(X, Y)$  is  $G$ -harmonious. Therefore,  $x \notin \tilde{S} \cap X \subset S \cap X$ . By repeating this exchange argument for any remaining  $x' \in (\tilde{S} \cap X) \setminus \tilde{T}$ , we arrive at the desired stable sets  $S'$  and  $T'$ . This proves (1).

We need a second, similar assertion:

*For any two stable sets  $S, T$  with  $S \cap X \subseteq T \cap X$  there are stable sets  $S'$  and  $T'$  such that  $\chi_S + \chi_T = \chi_{S'} + \chi_{T'}$ ,  $S' \cap X = S \cap X$  and  $S' \cap Y \supseteq T' \cap Y$ .* (2)

To see this, assume there is a  $y \in (T \cap Y) \setminus S$ , and let  $K$  be the component of  $G[S \cup T]$  containing  $y$ , and set  $\tilde{S} = S \Delta K$  and  $\tilde{T} = T \Delta K$ . The component  $K$  may not meet  $T \cap X$ , as then it would contain an induced  $y$ – $(T \cap X)$  path. This path would have even length, contradicting the definition of a  $G$ -harmonious tuple. As above, we see, moreover, that  $K$  meets  $Y$  only in  $T$ ; otherwise there would be an induced odd  $y$ – $(S \cap Y)$  path, which is impossible. Thus,  $\tilde{S}, \tilde{T}$  satisfy the first two conditions we want to have for  $S', T'$ , while  $(\tilde{T} \cap Y) \setminus \tilde{S}$  is smaller than  $(T \cap Y) \setminus S$ . Again repeating the argument yields  $S', T'$  as desired. This proves (2).

We now apply (1) iteratively to  $S_1$  (as  $S$ ) and each of  $S_2, \dots, S_k$  (as  $T$ ) in order to obtain stable sets  $R_1, \dots, R_k$  with  $R_1 \cap X \subseteq R_i \cap X$  for every  $i = 2, \dots, k$  and  $\sum_{i=1}^k \chi_{S_i} = \sum_{i=1}^k \chi_{R_i}$ . We continue applying (1), first to  $R_2$  and each of  $R_3, \dots, R_k$ , then to the resulting  $R'_3$  and each of  $R'_4, \dots, R'_k$  and so on, until we arrive at stable sets  $T_1, \dots, T_k$  with  $\sum_{i=1}^k \chi_{S_i} = \sum_{i=1}^k \chi_{T_i}$  that are nested on  $X$ :  $T_1 \cap X \subseteq \dots \subseteq T_k \cap X$ .

In a similar way, we use (2) to force the stable sets to become nested on  $Y$  as well. First, we apply (2) to  $T_1$  (as  $S$ ) and to each of  $T_2, \dots, T_k$  (as  $T$ ), then to the resulting  $T'_3$  and each of  $T'_4, \dots, T'_k$ , and so on. Proceeding in this manner, we obtain the desired stable sets  $S'_1, \dots, S'_k$ .  $\square$

**Lemma 12.** *Let  $(G_1, G_2)$  be a proper separation of a graph  $G$  so that  $X = V(G_1) \cap V(G_2)$  is a harmonious cutset. Let  $z \in \mathbb{Q}^{V(G)}$  be so that  $z|_{G_1} \in \text{SSP}(G_1)$  and  $z|_{G_2} \in \text{SSP}(G_2)$ . Then  $z \in \text{SSP}(G)$ .*

The lemma generalises the result by Chudnovsky et al. [7] that  $G = G_1 \cup G_2$  is 4-colourable if  $G_1$  and  $G_2$  are 4-colourable.

*Proof of Lemma 12.* Let  $(X_1, \dots, X_s)$  be a  $G$ -harmonious partition of  $X$ . As  $z|_{G_j} \in \text{SSP}(G_j)$ , for  $j = 1, 2$ , we can express  $z|_{G_1}$  as a convex combination of stable sets  $S_1, \dots, S_m$  of  $G_1$ , and  $z|_{G_2}$  as a convex combination of stable sets  $T_1, \dots, T_{m'}$  of  $G_2$ . Since  $z$  is a rational vector, we may even assume that

$$z|_{G_1} = \frac{1}{m} \sum_{i=1}^m \chi_{S_i} \quad \text{and} \quad z|_{G_2} = \frac{1}{m} \sum_{i=1}^m \chi_{T_i}.$$

Indeed, this can be achieved by repeating stable sets.

We first treat the case when  $s \leq 2$ . If  $s = 1$ , then set  $X_2 = \emptyset$ , so that whenever  $s \leq 2$ , we have  $X = X_1 \cup X_2$ .

Using Lemma 11, we find stable sets  $S'_1, \dots, S'_m$  of  $G_1$  so that  $z|_{G_1} = \frac{1}{m} \sum_{i=1}^m \chi_{S'_i}$  and

$$S'_1 \cap X_1 \subseteq \dots \subseteq S'_m \cap X_1, \text{ and } S'_1 \cap X_2 \supseteq \dots \supseteq S'_m \cap X_2$$

holds. Analogously, we obtain a convex combination  $z|_{G_2} = \frac{1}{m} \sum_{i=1}^m \chi_{T'_i}$  of stable sets  $T'_1, \dots, T'_m$  of  $G_2$  that are increasingly nested on  $X_1$  and decreasingly nested on  $X_2$ .

Define  $\bar{S}_1 \subsetneq \dots \subsetneq \bar{S}_k$  to be the distinct restrictions of the sets  $S'_i$  to  $X_1$ . More formally, let  $1 = i_1 < \dots < i_k < i_{k+1} = m + 1$  be so that

$$\bar{S}_t = S'_{i_t} \cap X_1 \text{ for all } i_t \leq i < i_{t+1}$$

We set, moreover,  $\lambda_t = \frac{1}{m}(i_{t+1} - i_t)$ . Equivalently,  $m\lambda_t$  is the number of  $S'_i$  with  $\bar{S}_t = S'_i \cap X_1$ . Then  $z|_{X_1} = \sum_{t=1}^k \lambda_t \chi_{\bar{S}_t}$  is a convex combination.

We do exactly the same in  $G_2$  in order to obtain  $z|_{X_1} = \sum_{t=1}^k \mu_t \chi_{\bar{T}_t}$ , where the sets  $\bar{T}_t$  are the distinct restrictions of the  $T'_i$  to  $X_1$ . With Lemma 10, we deduce first that  $\bar{S}_t = \bar{T}_t$  and  $\lambda_t = \mu_t$  for all  $t$ , from which we get that

$$S'_i \cap X_1 = T'_i \cap X_1 \text{ for all } i = 1, \dots, m.$$

The same argument, only applied to the restrictions of  $S'_i$  and of  $T'_i$  to  $X_2$ , yields that also

$$S'_i \cap X_2 = T'_i \cap X_2 \text{ for all } i = 1, \dots, m.$$

Thus,  $R_i = S'_i \cup T'_i$  is, for every  $i = 1, \dots, m$ , a stable set of  $G$ . Consequently,  $z = \frac{1}{m} \sum_{i=1}^m \chi_{R_i}$  is a convex combination of stable sets and thus a point of  $\text{SSP}(G)$ .

It remains to treat the case when the harmonious cutset has at least three parts, that is, when  $s \geq 3$ . We claim that there are sets  $\mathcal{S}_0, \mathcal{S}_1, \dots, \mathcal{S}_s$  of stable sets of  $G_1$  so that

- (a)  $z|_{G_1} = \frac{1}{m} \sum_{j=0}^s \sum_{S \in \mathcal{S}_j} \chi_S$  and  $\sum_{j=0}^s |\mathcal{S}_j| = m$ ;
- (b) for  $j = 1, \dots, s$  if  $S \in \mathcal{S}_j$  then  $X_j \cap S$  is non-empty; and
- (c) for  $j = 0, \dots, s$  if  $S, T \in \mathcal{S}_j$  then  $X_j \cap S \subseteq X_j \cap T$  or  $X_j \cap S \supseteq X_j \cap T$ .

Moreover, there are analogous sets  $\mathcal{T}_0, \mathcal{T}_1, \dots, \mathcal{T}_s$  for  $G_2$ .

To prove the claim note first that each  $S_i$  meets at most one of the sets  $X_j$  as each two induce a complete bipartite graph. Therefore, we can partition  $\{S_1, \dots, S_m\}$  into sets  $\mathcal{S}'_0, \dots, \mathcal{S}'_s$  so that (a) and (b) are satisfied. Next, we apply Lemma 11 to each  $\mathcal{S}'_j$  and  $(X_j, \emptyset)$  in order to obtain sets  $\mathcal{S}''_j$  that satisfy (a) and (c) but not necessarily (b); property (a) still holds as Lemma 11 guarantees  $\sum_{S \in \mathcal{S}'_j} \chi_S = \sum_{S \in \mathcal{S}''_j} \chi_S$  for each  $j$ . If (b) is violated, then only because for some  $j \neq 0$  there is  $S \in \mathcal{S}''_j$  that is not only disjoint from  $X_j$  but also from all other  $X_{j'}$ . In order to repair (b) we remove all stable sets  $S$  in  $\bigcup_{j=1}^s \mathcal{S}''_j$  that

are disjoint from  $\bigcup_{j=1}^s X_j$  from their respective sets and add them to  $\mathcal{S}''$ . The resulting sets  $\mathcal{S}_0, \mathcal{S}_1, \dots, \mathcal{S}_s$  then satisfy (a)–(c). The proof for the  $\mathcal{T}_j$  is the same.

As a consequence of (a) and (b) it follows for  $j = 0, 1, \dots, s$  that

$$\sum_{S \in \mathcal{S}_j} \chi_{S \cap X_j} = m \cdot z|_{X_j} = \sum_{T \in \mathcal{T}_j} \chi_{T \cap X_j} \quad (3)$$

Now, consider  $j \neq 0$ . Then, by (b) and (c), there is a vertex  $v \in X_j$  that lies in every  $S \in \mathcal{S}_j$ . Thus, we have  $\sum_{S \in \mathcal{S}_j} \chi_S(v) = |\mathcal{S}_j|$ .

Evaluating (3) at  $v \in X_j$ , we obtain

$$|\mathcal{S}_j| = m \cdot z(v) = \sum_{T \in \mathcal{T}_j} \chi_T(v) \leq |\mathcal{T}_j|.$$

Reversing the roles of  $\mathcal{S}_j$  and  $\mathcal{T}_j$ , we also get  $|\mathcal{T}_j| \leq |\mathcal{S}_j|$ , and thus that  $|\mathcal{T}_j| = |\mathcal{S}_j|$ , as long as  $j \neq 0$ . That this also holds for  $j = 0$  follows from  $m = \sum_{j=0}^s |\mathcal{S}_j| = \sum_{j=0}^s |\mathcal{T}_j|$ , so that we get  $m_j := |\mathcal{S}_j| = |\mathcal{T}_j|$  for every  $j = 0, 1, \dots, s$ .

Together with (3) this implies, in particular, that

$$\frac{1}{m_j} \sum_{S \in \mathcal{S}_j} \chi_{S \cap X_j} = \frac{1}{m_j} \sum_{T \in \mathcal{T}_j} \chi_{T \cap X_j}$$

We may, therefore, define a vector  $y^j$  on  $V(G)$  by setting

$$y^j|_{G_1} := \frac{1}{m_j} \sum_{S \in \mathcal{S}_j} \chi_S \text{ and } y^j|_{G_2} := \frac{1}{m_j} \sum_{T \in \mathcal{T}_j} \chi_T \quad (4)$$

For any  $j = 0, \dots, s$ , define  $G^j = G - \bigcup_{r \neq j} X_r$ , and observe that  $X_j$  is a harmonious cutset of  $G^j$  consisting of only one part. (That is,  $X_j$  is  $G^j$ -harmonious.) Moreover, as (4) shows, the restriction of  $y^j$  to  $G_1 \cap G^j$  lies in  $\text{SSP}(G_1 \cap G^j)$ , while the restriction to  $G_2 \cap G^j$  lies in  $\text{SSP}(G_2 \cap G^j)$ . Thus, we can apply the first part of this proof, when  $s \leq 2$ , in order to deduce that  $y^j \in \text{SSP}(G^j) \subseteq \text{SSP}(G)$ .

To finish the proof we observe, with (a) and (4), that

$$z = \sum_{j=0}^s \frac{m_j}{m} y^j$$

As, by (a),  $\sum_{j=0}^s m_j = m$ , this means that  $z$  is a convex combination of points in  $\text{SSP}(G)$ , and thus itself an element of  $\text{SSP}(G)$ .  $\square$

**Corollary 13.** *Let  $(G_1, G_2)$  be a proper separation of  $G$  so that  $X = V(G_1) \cap V(G_2)$  is a harmonious cutset. Then  $G$  is  $t$ -perfect if and only if  $G_1$  and  $G_2$  are  $t$ -perfect.*

*Proof.* Assume that  $G_1$  and  $G_2$  are  $t$ -perfect, and consider a rational point  $z \in \text{TSTAB}(G)$ . Then  $z|_{G_1} \in \text{SSP}(G_1)$  and  $z|_{G_2} \in \text{SSP}(G_2)$ , which means that Lemma 12 yields  $z \in \text{SSP}(G)$ . Since this is true for all rational  $z$  it extends to real  $z$  as well.  $\square$

The corollary directly implies Lemma 9.

## 6 $P_5$ -free graphs

Let  $\mathcal{F}$  be the set of graphs consisting of  $P_5$ ,  $K_4$ ,  $W_5$ ,  $C_7^2$ ,  $A_{10}^2$  and  $A_{13}^3$  together with the three graphs in Figure 5. Note that the latter three graphs all contain  $K_4$  as a  $t$ -minor: for (a) and (b)  $K_4$  is obtained by a  $t$ -contraction at any vertex of degree 2, while for (c) both vertices of degree 2 need to be  $t$ -contracted. In particular, every graph in  $\mathcal{F}$  besides  $P_5$  is  $t$ -imperfect. We say that a graph is  $\mathcal{F}$ -free if it contains none of the graphs in  $\mathcal{F}$  as an induced subgraph.




Figure 5: Three graphs that  $t$ -contract to  $K_4$

We prove a lemma that implies directly Theorem 1:

**Lemma 14.** *Any  $\mathcal{F}$ -free graph is  $t$ -perfect.*

We first examine how a vertex may position itself relative to a 5-cycle in an  $\mathcal{F}$ -free graph.

**Lemma 15.** *Let  $G$  be an  $\mathcal{F}$ -free graph. If  $v$  is a neighbour of a 5-hole  $C$  in  $G$  then  $v$  has either exactly two neighbours in  $C$ , and these are non-consecutive in  $C$ ; or  $v$  has exactly three neighbours in  $C$ , and these are not all consecutive.*




Figure 6: The types of neighbours of a 5-hole

*Proof.* See Figure 6 for the possible types of neighbours (up to isomorphism). Of these, (b) and (c) contain an induced  $P_5$ ; (e) and (g) are the same as (a) and (b) in Figure 5 and thus in  $\mathcal{F}$ ; (h) is  $W_5$ . Only (d) and (f) remain.  $\square$

**Lemma 16.** *Let  $G$  be an  $\mathcal{F}$ -free graph, and let  $u$  and  $v$  be two non-adjacent vertices such that both of them have precisely three neighbours in a 5-hole  $C$ .*

Then  $u$  and  $v$  have either all three or exactly two non-consecutive neighbours in  $C$  in common.




Figure 7: The possible configurations of Lemma 16

*Proof.* By Lemma 15, both of  $u$  and  $v$  have to be as in (f) of Figure 6. Figure 7 shows the possible configurations of  $u$  and  $v$  (up to isomorphism). Of these, (b) is impossible as there is an induced  $P_5$ —the other two configurations (a) and (c) may occur.  $\square$

A subgraph  $H$  of a graph  $G$  is *dominating* if every vertex in  $G - H$  has a neighbour in  $H$ .

**Lemma 17.** *Let  $G$  be an  $\mathcal{F}$ -free graph. Then, either any 5-hole of  $G$  is dominating or  $G$  contains a harmonious cutset.*

*Proof.* Assume that there is a 5-hole  $C = c_1 \dots c_5 c_1$  that fails to dominate  $G$ . Our task consists in finding a harmonious cutset. We first observe:

*Let  $u \in N(C)$  be a neighbour of some  $x \notin N(C)$ . Then  $u$  has exactly three neighbours in  $C$ , not all of which are consecutive.* (5)

So, such a  $u$  is as in (f) of Figure 6. Indeed, by Lemma 15, only (d) or (f) in Figure 6 are possible. In the former case, we may assume that the neighbours of  $u$  in  $C$  are  $c_1$  and  $c_3$ . Then, however,  $xuc_1c_4c_5$  is an induced  $P_5$ . This proves (5).




Figure 8:  $x$  in solid black.

Consider two adjacent vertices  $y, z \notin N(C)$ , and assume that there is a  $u \in N(y) \cap N(C)$  that is not adjacent to  $z$ . We may assume that  $N(u) \cap C = \{c_1, c_2, c_4\}$  by (5). Then,  $zyuc_2c_3$  is an induced  $P_5$ , which is impossible. Thus:

$$N(y) \cap N(C) = N(z) \cap N(C) \text{ for any adjacent } y, z \notin N(C). \quad (6)$$

Next, fix some vertex  $x$  that is not dominated by  $C$  (and, by assumption, there is such a vertex). As a consequence of (6),  $N(x) \cap N(C)$  separates  $x$  from  $C$ . In particular,

$$X := N(x) \cap N(C) \text{ is a separator.} \quad (7)$$

Consider two vertices  $u, v \in X$ . Then, by (5), each of  $u$  and  $v$  have exactly three neighbours in  $C$ , not all of which are consecutive. We may assume that  $N(u) \cap V(C) = \{c_1, c_2, c_4\}$ .

First, assume that  $uv \in E(G)$ , and suppose that the neighbourhoods of  $u$  and  $v$  in  $C$  are the same. This, however, is impossible as then  $u, v, c_1, c_2$  form a  $K_4$ . Therefore,  $uv \in E(G)$  implies  $N(u) \cap V(C) \neq N(v) \cap V(C)$ .

Now assume  $uv \notin E(G)$ . By Lemma 16, there are only two possible configurations (up to isomorphism) for the neighbours of  $v$  in  $C$ ; these are (a) and (c) in Figure 7. The first of these, (a) in Figure 8, is impossible, as this is a graph of  $\mathcal{F}$ ; see Figure 5 (c). Thus, we see that  $u, v$  are as in (b) of Figure 8, that is, that  $u$  and  $v$  have the same neighbours in  $C$ .

To sum up, we have proved that:

$$uv \in E(G) \Leftrightarrow N(u) \cap V(C) \neq N(v) \cap V(C) \text{ for any two } u, v \in X \quad (8)$$

An immediate consequence is that the neighbourhoods in  $C$  partition  $X$  into stable sets  $X_1, \dots, X_k$  such that  $X_i$  is complete to  $X_j$  whenever  $i \neq j$ . As  $X$  cannot contain any triangle—together with  $x$  this would result in a  $K_4$ —it follows that  $k \leq 2$ . If  $k = 1$ , we put  $X_2 = \emptyset$  so that always  $X = X_1 \cup X_2$ .

We claim that  $X$  is a harmonious cutset. As  $X$  is a separator, by (7), we only need to prove that  $(X_1, X_2)$  is  $G$ -harmonious. For this, we have to check the parities of induced  $X_1$ -paths and of  $X_2$ -paths; since  $X_1$  is complete to  $X_2$  any induced  $X_1$ - $X_2$  path is a single edge and has therefore odd length.

Suppose there is an odd induced  $X_1$ -path or  $X_2$ -path. Clearly, we may assume there is such a path  $P$  that starts in  $u \in X_1$  and ends in  $v \in X_1$ . As  $X_1$  is stable, and as  $G$  is  $P_5$ -free, it follows that  $P$  has length 3. So, let  $P = upqv$ .

Let us consider the position of  $p$  and  $q$  relative to  $C$ . We observe that neither  $p$  nor  $q$  can be in  $C$ . Indeed, if, for instance,  $p$  was in  $C$  then  $p$  would also be a neighbour of  $v$  since  $N(u) \cap V(C) = N(v) \cap V(C)$ , by (8). This, however, is impossible as  $P$  is induced.

Next, assume that  $p, q \notin N(C)$  holds. Since  $p$  and  $q$  are adjacent, we can apply (6) to  $p$  and  $q$ , which results in  $N(p) \cap N(C) = N(q) \cap N(C)$ . However, as  $u$  lies in  $N(p) \cap N(C)$  it then also is a neighbour of  $q$ , which contradicts that  $upqv$  is induced.

It remains to consider the case when one of  $p$  and  $q$ ,  $p$  say, lies in  $N(C)$ . As  $p$  is adjacent to  $u$  but not to  $v$ , both of which lie in  $X_1$  and are therefore non-neighbours, it follows from (8) that  $p \notin X$ . In particular,  $p$  is not a neighbour of  $x$ , which means that  $pu xv$  is an induced path.

Suppose there is a neighbour  $c \in V(C)$  of  $p$  that is not adjacent to  $u$ . By (8),  $c$  is not adjacent to  $v$  either, so that  $cpu xv$  forms an induced  $P_5$ , a contradiction. Thus,  $N(p) \cap V(C) \subseteq N(u)$  has to hold. By (5), we may assume that the neighbours of  $u$  in  $C$  are precisely  $c_1, c_2, c_4$ . As  $u$  and  $p$  are adjacent,  $p$  cannot be neighbours with both of  $c_1$  and  $c_2$ , as this would result in a  $K_4$ . Thus, we may assume that  $N(p) \cap V(C) = \{c_2, c_4\}$ . (Note, that  $p$  has at least two neighbours in  $C$ , by Lemma 15.)

To conclude, we observe that  $pc_4c_5c_1c_2p$  forms a 5-hole, in which  $u$  has four neighbours, namely  $c_1, c_2, c_4, p$ . This, however, is in direct contradiction to Lemma 15, which means that our assumption is false, and there is no odd induced  $X_1$ -path, and no such  $X_2$ -path either. Consequently,  $(X_1, X_2)$  is  $G$ -harmonious, and  $X = X_1 \cup X_2$  therefore a harmonious cutset.  $\square$

**Proposition 18.** *Let  $G$  be a  $t$ -imperfect graph. Then either  $G$  contains an odd hole or it contains  $K_4$  or  $C_7^2$  as an induced subgraph.*

*Proof.* Assume that  $G$  does not contain any odd hole and neither  $K_4$  nor  $C_7^2$  as an induced subgraph. Observe that any odd antihole of length  $\geq 9$  contains  $K_4$ . Since the complement of a 5-hole is a 5-hole, and since  $C_7^2$  is the odd antihole of length 7, it follows that  $G$  cannot contain any odd antihole at all.

Now, by the strong perfect graph theorem it follows that  $G$  is perfect. (Note that we do not need the full theorem but only the far easier version for  $K_4$ -free graphs; see Tucker [28].) Since  $G$  does not contain any  $K_4$  it is therefore  $t$ -perfect as well.  $\square$

**Lemma 19.** *Let  $G$  be an  $\mathcal{F}$ -free graph. If  $G$  contains a 5-hole, and if every 5-hole is dominating then  $G$  is near-bipartite.*

*Proof.* Let  $G$  contain a 5-hole, and assume every 5-hole to be dominating. Suppose that the lemma is false, i.e. that  $G$  fails to be near-bipartite. In particular, there is a vertex  $v$  such that  $G - N(v)$  is not bipartite, and therefore contains an induced odd cycle  $T$ . As any 5-hole is dominating and any  $k$ -hole with  $k > 5$  contains an induced  $P_5$ ,  $T$  has to be a triangle. Let  $T = xyz$ . We distinguish two cases, both of which will lead to a contradiction.

**Case:**  $v$  lies in a 5-hole  $C$ .

Let  $C = c_1 \dots c_5c_1$ , and  $v = c_1$ . Then  $T$  could meet  $C$  in 0, 1 or 2 vertices. If  $T$  has two vertices with  $C$  in common, these have to be  $c_3$  and  $c_4$  as the others are neighbours of  $v$ . Then, the third vertex of  $T$  has two consecutive neighbours in  $C$ , which means that by Lemma 15 its third neighbour in  $C$  has to be  $c_1 = v$ , which is impossible.

Next, suppose that  $T$  meets  $C$  in one vertex,  $c_3 = z$ , say. By Lemma 15, each of  $x, y$  has to have a neighbour opposite of  $c_3$  in  $C$ , that is, either  $c_1$  or  $c_5$ . As  $c_1 = v$ , both of  $x, y$  are adjacent with  $c_5$ . The vertices  $x, y$  could have a third neighbour in  $C$ ; this would necessarily be  $c_2$ . However, not both can be adjacent to  $c_2$  as then  $x, y, c_2, c_3$  would induce a  $K_4$ . Thus, assume  $x$  to have exactly  $c_3$  and  $c_5$  as neighbours in  $C$ . This means that  $C' = c_3xc_5c_1c_2c_3$  is a 5-hole in which  $y$  has at least three consecutive neighbours,  $c_3, x, c_5$ , which is impossible (again, by Lemma 15).

Finally, suppose that  $T$  is disjoint from  $C$ . Each of  $x, y, z$  has at least two neighbours among  $c_2, \dots, c_5$ , and no two have  $c_3$  or  $c_4$  as neighbour; otherwise we would have found a triangle in  $G - N(v)$  meeting  $C$  in exactly one vertex, and could reduce to the previous subcase. Thus, we may assume that  $x$  is adjacent to  $c_2$  and  $c_5$ . Moreover, since no vertex of  $x, y, z$  can be adjacent to both  $c_3$  and  $c_4$  (as then it would also be adjacent to  $c_1$ , by Lemma 15) and no  $c_i \in C$  can be adjacent to all vertices of  $T$  (because otherwise  $c_i, x, y, z$  would form a  $K_4$ ), it follows that we may assume that  $y$  is adjacent to  $c_2$  but not to  $c_5$ , while  $z$  is adjacent to  $c_5$  but not to  $c_2$ . Then,  $c_1c_2yzc_5c_1$  is a 5-hole in which  $x$  has

four neighbours, in obvious contradiction to Lemma 15. Therefore, this case is impossible.

**Case:**  $v$  does not lie in any 5-hole.

Let  $C = c_1 \dots c_5 c_1$  be a 5-hole. Since every 5-hole is dominating,  $v$  has a neighbour in  $C$ , and thus is, by Lemma 15, either as in (f) of Figure 6 or as in (d). The latter, however, is impossible since then  $v$  would be contained in a 5-hole. Therefore, we may assume that the neighbours of  $v$  in  $C$  are precisely  $\{c_1, c_2, c_4\}$ . As a consequence,  $T$  can meet  $C$  in at most  $c_3$  and  $c_5$ , both not in both as  $C$  is induced.

Suppose  $T = xyz$  meets  $C$  in  $x = c_3$ . If  $y$  is not adjacent to either of  $c_1$  and  $c_4$ , then  $c_1 v c_4 x y$  forms an induced  $P_5$ . If, on the other hand,  $y$  is adjacent to  $c_4$  then, by Lemma 15, also to  $c_1$ . Thus,  $y$  is either adjacent to  $c_1$  or to both  $c_1$  and  $c_4$ . The same holds for  $z$ . Since  $y$  and  $z$  are adjacent, they cannot both have three neighbours in  $C$  (otherwise  $G$  would contain a  $K_4$ ). Suppose  $N(y) \cap C = \{x, c_1\}$ . But then  $x c_4 c_5 c_1 y x$  forms an induced 5-cycle in which  $z$  has at least three consecutive neighbours; a contradiction to Lemma 15.

Consequently,  $T$  is disjoint from every 5-hole. By Lemma 15, each of  $x, y, z$  has neighbours in  $C$  as in (d) or (f) of Figure 6. However, if any of  $x, y, z$  has only two neighbours in  $C$  as in (d) then that vertex together with four vertices of  $C$  forms a 5-hole that meets  $T$ —this is precisely the situation of the previous subcase. Thus, we may assume that all vertices of  $T$  have three neighbours in  $C$  as in (f) of Figure 6. If we consider the possible configurations of two non-adjacent vertices which have three neighbours in  $C$  (namely  $v$  and a vertex of  $T$ ) as we have done in Lemma 7, we see that only (a) and (c) in Figure 7 are possible. But then each vertex of  $T$  has to be adjacent to  $c_4$ , which means that  $T$  together with  $c_4$  induces a  $K_4$ , which is impossible.  $\square$

*Proof of Lemma 14.* Suppose that  $G$  is a  $t$ -imperfect and but  $\mathcal{F}$ -free. By deleting suitable vertices we may assume that every proper induced subgraph of  $G$  is  $t$ -perfect. In particular, by Lemma 9,  $G$  does not admit a harmonious cutset. Since  $G$  is  $t$ -imperfect it contains an odd hole, by Proposition 18, and since  $G$  is  $P_5$ -free, the odd hole is of length 5. From Lemma 17 we deduce that any 5-hole is dominating. Lemma 19 implies that  $G$  is near-bipartite.

Noting that both  $A_{13}^4$  and  $A_{19}^7$ , as well as any Möbius ladder or any odd wheel larger than  $W_5$ , contain an induced  $P_5$ , we see with Proposition 3 that  $G$  is  $t$ -perfect after all.  $\square$

By Lemma 14, a  $P_5$ -free graph is either  $t$ -perfect or contains one of eight  $t$ -imperfect graphs as an induced subgraph. Obviously, checking for these forbidden induced subgraphs can be done in polynomial time, so that we get as immediate algorithmic consequence:

**Theorem 20.**  *$P_5$ -free  $t$ -perfect graphs can be recognised in polynomial time.*

We suspect, but cannot currently prove, that  $t$ -perfection can be recognised as well in polynomial time in near-bipartite graphs.

## 7 Colouring

Can  $t$ -perfect graphs always be coloured with few colours? This is one of the main open questions about  $t$ -perfect graphs. A conjecture by Shepherd and



Sebó asserts that four colours are always enough:

**Conjecture 21** (Shepherd; Sebó [23]). *Every  $t$ -perfect graph is 4-colourable.*

The conjecture is known to hold in a number of graph classes, for instance in claw-free graphs, where even three colours are already sufficient; see [5]. It is straightforward to verify the conjecture for near-bipartite graphs:

**Proposition 22.** *Every near-bipartite  $t$ -perfect graph is 4-colourable.*

*Proof.* Pick any vertex  $v$  of a near-bipartite and  $t$ -perfect graph  $G$ . Then  $G - N(v)$  is bipartite and may be coloured with colours 1, 2. On the other hand, as  $G$  is  $t$ -perfect the neighbourhood  $N(v)$  necessarily induces a bipartite graph as well; otherwise  $v$  together with a shortest odd cycle in  $N(v)$  would form an odd wheel. Thus we can colour the vertices in  $N(v)$  with the colours 3, 4.  $\square$

Near-bipartite  $t$ -perfect graphs can, in general, not be coloured with fewer colours. Indeed, this is even true if we restrict ourselves further to complements of line graphs, which is a subclass of near-bipartite graphs. Two  $t$ -perfect graphs in this class that need four colours are:  $\overline{L(\Pi)}$ , the complement of the line graph of the prism, and  $\overline{L(W_5)}$ . The former was found by Laurent and Seymour (see [25, p. 1207]), while the latter was discovered by Benchetrit [2]. Moreover, Benchetrit showed that any 4-chromatic  $t$ -perfect complement of a line graph contains one of  $\overline{L(\Pi)}$  and  $\overline{L(W_5)}$  as an induced subgraph.

How about  $P_5$ -free  $t$ -perfect graphs? Applying insights of Sebó and of Sumner, Benchetrit [2] proved that  $P_5$ -free  $t$ -perfect graphs are 4-colourable. This is not tight:

**Theorem 23.** *Every  $P_5$ -free  $t$ -perfect graph  $G$  is 3-colourable.*

For the proof we use that there is a finite number of obstructions for 3-colourability in  $P_5$ -free graphs:

**Theorem 24** (Maffray and Morel [22]). *A  $P_5$ -free graph is 3-colourable if and only if it does not contain  $K_4$ ,  $W_5$ ,  $C_7^2$ ,  $A_{10}^2$ ,  $A_{13}^3$  or any of the seven graphs in Figure 9 as an induced subgraph.*

(Maffray and Morel call these graphs  $F_1$ – $F_{12}$ . The graphs  $K_4$ ,  $W_5$ ,  $C_7^2$ ,  $A_{10}^2$ ,  $A_{13}^3$  are respectively  $F_1$ ,  $F_2$ ,  $F_9$ ,  $F_{11}$  and  $F_{12}$ .) A similar result was obtained by Bruce, Hoàng and Sawada [4], who gave a list of five forbidden (not necessarily induced) subgraphs.

*Proof of Theorem 23.* Any  $P_5$ -free graph  $G$  that cannot be coloured with three colours contains one of the twelve induced subgraphs of Theorem 24. Of these twelve graphs, we already know that  $K_4$ ,  $W_5$ ,  $C_7^2$ ,  $A_{10}^2$ ,  $A_{13}^3$  are  $t$ -imperfect, and thus cannot be induced subgraphs of a  $t$ -perfect graph. It remains to consider the seven graphs in Figure 9. These graphs are  $t$ -imperfect, too: each can be turned into  $K_4$  by first deleting the grey vertices and then performing a  $t$ -contraction at the respective black vertex.  $\square$

We mention that Benchetrit [2] also showed that  $P_6$ -free  $t$ -perfect graphs are 4-colourable. This is tight: both  $\overline{L(\Pi)}$  and  $\overline{L(W_5)}$  (and indeed all complements of line graphs) are  $P_6$ -free. We do not know whether  $P_7$ -free  $t$ -perfect graphs are 4-colourable.




Figure 9: The remaining 4-critical  $P_5$ -free graphs of Theorem 24; in Maffray and Morel [22] these are called  $F_3$ – $F_8$  and  $F_{10}$ . In each graph, deleting the grey vertices and then  $t$ -contracting at the black vertex results in  $K_4$ .

We turn now to fractional colourings. A motivation for Conjecture 21 was certainly the fact that the *fractional chromatic number*  $\chi_f(G)$  of a  $t$ -perfect graph  $G$  is always bounded by 3. More precisely, if  $og(G)$  denotes the *odd girth* of  $G$ , that is, the length of the shortest odd cycle, then  $\chi_f(G) = 2 \frac{og(G)}{og(G)-1}$  as long as  $G$  is  $t$ -perfect (and non-bipartite). This follows from linear programming duality; see for instance Schrijver [25, p. 1206].

Recall that a graph  $G$  is perfect if and only if  $\chi(H) = \omega(H)$  for every induced subgraph  $H$  of  $G$ . As odd cycles seem to play a somewhat similar role for  $t$ -perfection as cliques play for perfection, one might conjecture that  $t$ -perfection is characterised in an analogous way:

**Conjecture 25.** *A graph  $G$  is  $t$ -perfect if and only if  $\chi_f(H) = 2 \frac{og(H)}{og(H)-1}$  for every non-bipartite  $t$ -minor  $H$  of  $G$ .*

Note that the conjecture becomes false if, instead of  $t$ -minors, only induced subgraphs  $H$  are considered. Indeed, in the  $t$ -imperfect graph obtained from  $K_4$  by subdividing some edge twice, all induced subgraphs satisfy the condition (but not the  $t$ -minor  $K_4$ ).

An alternative but equivalent formulation of the conjecture is:  $\chi_f(G) > 2 \frac{og(G)}{og(G)-1}$  holds for every minimally  $t$ -imperfect graph  $G$ . It is straightforward to check that all minimally  $t$ -imperfect graphs that are known to date satisfy this. In particular, it follows that the conjecture is true for  $P_5$ -free graphs, for near-bipartite graphs, as well as for claw-free graphs; see [5] for the minimally  $t$ -imperfect graphs that are claw-free.

## Acknowledgment

We thank Oliver Schaudt for pointing out [22]. We also thank one of the referees for bringing the work of Holm et al. [20] to our attention, as well as identifying inaccuracies that led to a (hopefully!) clearer presentation of Theorems 20 and 23.

## References

- [1] F. Barahona and A.R. Mahjoub, *Decompositions of graphs and polyhedra III: Graphs with no  $W_4$  minor*, SIAM J. Discrete Math. **7** (1994), 372–389.
- [2] Y. Benchetrit, *Geometric properties of the chromatic number: polyhedra, structure and algorithms*, Ph.D. thesis, Université de Grenoble, 2015.
- [3] M. Boulala and J.P. Uhry, *Polytope des indépendants d'un graphe série-parallèle*, Disc. Math. **27** (1979), 225–243.
- [4] D. Bruce, C.T. Hoàng, and J. Sawada, *A certifying algorithm for 3-colorability of  $P_5$ -free graphs*, Algorithms and Computation, Lecture Notes in Computer Science, vol. 5878, Springer Berlin Heidelberg, 2009, pp. 594–604.
- [5] H. Bruhn and M. Stein, *On claw-free  $t$ -perfect graphs*, Math. Prog. (2012), 461–480.
- [6] M. Chudnovsky, N. Robertson, P. Seymour, and R. Thomas, *The strong perfect graph theorem*, Ann. Math. **164** (2006), 51–229.
- [7] M. Chudnovsky, N. Robertson, P.D. Seymour, and R. Thomas,  *$K_4$ -free graphs with no odd holes*, J. Combin. Theory (Series B) **100** (2010), 313–331.
- [8] M. Chudnovsky and P. Seymour, *The structure of claw-free graphs*, Surveys in combinatorics **327** (2005), 153–171.
- [9] V. Chvátal, *On certain polytopes associated with graphs*, J. Combin. Theory (Series B) **18** (1975), 138–154.
- [10] R. Diestel, *Graph theory*, 4 ed., Springer-Verlag, 2010.
- [11] F. Eisenbrand, S. Funke, N. Garg, and J. Könemann, *A combinatorial algorithm for computing a maximum independent set in a  $t$ -perfect graph*, In Proc. of the ACM-SIAM Symp. on Discrete Algorithms, ACM, 2003, pp. 517–522.
- [12] F. Eisenbrand, G. Oriolo, G. Stauffer, and P. Ventura, *Circular ones matrices and the stable set polytope of quasi-line graphs*, International Conference on Integer Programming and Combinatorial Optimization (2005), 291–305.
- [13] J. Fonlupt and A. Hadjar, *The stable set polytope and some operations on graphs*, Disc. Math. **252** (2002), 123–140.
- [14] J. Fonlupt and J.P. Uhry, *Transformations which preserve perfectness and  $h$ -perfectness of graphs*, Ann. Disc. Math. **16** (1982), 83–95.
- [15] D.R. Fulkerson, *Anti-blocking polyhedra*, J. Combin. Theory (Series B) **12** (1972), 50–71.
- [16] A.M.H. Gerards, *A min-max relation for stable sets in graphs with no odd- $K_4$* , J. Combin. Theory (Series B) **47** (1989), 330–348.

- [17] A.M.H. Gerards and F.B. Shepherd, *The graphs with all subgraphs  $t$ -perfect*, SIAM J. Discrete Math. **11** (1998), 524–545.
- [18] M. Grötschel, L. Lovász, and A. Schrijver, *Relaxations of vertex packing*, J. Combin. Theory (Series B) **40** (1986), 330–343.
- [19] ———, *Geometric algorithms and combinatorial optimization*, Springer-Verlag, 1988.
- [20] E. Holm, L. M. Torres, and A. K. Wagler, *On the Chvátal rank of linear relaxations of the stable set polytope*, Intl. Trans. in Op. Res. **17** (2010), 827–849.
- [21] L. Lovász, *Submodular functions and convexity*, Mathematical Programming The State of the Art (A. Bachem, B. Korte, and M. Grötschel, eds.), Springer Berlin Heidelberg, 1983, pp. 235–257.
- [22] F. Maffray and G. Morel, *On 3-colorable  $P_5$ -free graphs*, SIAM J. Discrete Math. **26** (2012), 1682–1708.
- [23] A. Sebő, personal communication.
- [24] A. Ben Rebea, *Étude des stables dans les graphes quasi-adjoints*, Ph.D. thesis, Université de Grenoble, 1981.
- [25] A. Schrijver, *Combinatorial optimization. Polyhedra and efficiency*, Springer-Verlag, 2003.
- [26] F.B. Shepherd, *Applying Lehman’s theorems to packing problems*, Math. Prog. **71** (1995), 353–367.
- [27] L.E. Trotter, *A class of facet producing graphs for vertex packing polyhedra*, Disc. Math. **12** (1975), 373–388.
- [28] A. Tucker, *Critical perfect graphs and perfect 3-chromatic graphs*, J. Combin. Theory (Series B) **23** (1977), 143–149.
- [29] A. Wagler, *Antiwebs are rank-perfect*, 4OR **2** (2004), 149–152.
- [30] ———, *On rank-perfect subclasses of near-bipartite graphs*, 4OR **3** (2005), 329–336.

Version October 24, 2016

Henning Bruhn <henning.bruhn@uni-ulm.de>  
 Elke Fuchs <elke.fuchs@uni-ulm.de>  
 Institut für Optimierung und Operations Research  
 Universität Ulm, Ulm  
 Germany