

# On chaotic minimal center of attraction of a Lagrange stable motion for topological semi flows

Xiongping Dai

Department of Mathematics, Nanjing University, Nanjing 210093, People's Republic of China

---

## Abstract

Let  $f: \mathbb{R}_+ \times X \rightarrow X$  be a topological semi flow on a Polish space  $X$ . In 1977, Karl Sigmund conjectured that if there is a point  $x$  in  $X$  such that the motion  $f(t, x)$  has just  $X$  as its minimal center of attraction, then the set of all such  $x$  is *residual* in  $X$ . In this paper, we present a positive solution to this conjecture and apply it to the study of chaotic dynamics of minimal center of attraction of a motion.

*Keywords:* Minimal center of attraction · Chaotic motion · Semi flow

*2010 MSC:* 37B05 · 37A60 · 54H20

---

## 1. Introduction

By a  $C^0$ -semi flow over a metric space  $X$ , we here mean a transformation  $f: \mathbb{R}_+ \times X \rightarrow X$  where  $\mathbb{R}_+ = [0, \infty)$ , which satisfies the following three conditions:

- (1) The initial condition:  $f(0, x) = x$  for all  $x \in X$ .
- (2) The condition of continuity: if there be given two convergent sequences  $t_n \rightarrow t_0$  in  $\mathbb{R}_+$  and  $x_n \rightarrow x_0$  in  $X$ , then  $f(t_n, x_n) \rightarrow f(t_0, x_0)$  as  $n \rightarrow \infty$ .
- (3) The semigroup condition:  $f(t_2, f(t_1, x)) = f(t_1 + t_2, x)$  for any  $x$  in  $X$  and any times  $t_1, t_2$  in  $\mathbb{R}_+$ .

Sometimes we write  $f(t, x) = f^t(x)$  for any  $t \geq 0$  and  $x \in X$ ; and for any given point  $x \in X$  we call  $f(t, x)$  a motion in  $X$  and  $O_f(x) = f(\mathbb{R}_+, x)$  the orbit starting from the point  $x$ . If  $O_f(x)$  is precompact (i.e.  $\overline{O_f(x)}$  is compact) in  $X$ , then we say that  $f(t, x)$  is *Lagrange stable*.

We refer to any subset  $\Lambda$  of  $X$  as an *invariant* set if  $f(t, x) \in \Lambda$  for each point  $x \in \Lambda$  and any time  $t \geq 0$ . In dynamical systems, statistical mechanics and ergodic theory, we shall have to do with “probability of sojourn of a motion  $f(t, x)$  in a given region  $E$  of  $X$ ” as  $t \rightarrow +\infty$ :

$$P(f(t, x) \in E) = \lim_{T \rightarrow +\infty} \frac{1}{T} \int_0^T \mathbb{1}_E(f(t, x)) dt,$$

where  $\mathbb{1}_E(x)$  is the characteristic function of the set  $E$  on  $X$ .

This motivates H.F. Hilmy to introduce following important concept, which was discussed in [13, 15, 16, 12], for example.

---

*Email address:* xpdai@nju.edu.cn (Xiongping Dai)

*Preprint submitted to arXiv, July 2, 2015*

*October 11, 2018*

**Definition 1.1** (Hilmy 1936 [10]). Given any  $x \in X$ , a closed subset  $C_x$  of  $X$  is called the *center of attraction of the motion*  $f(t, x)$  as  $t \rightarrow +\infty$  if  $P(f(t, x) \in B_\varepsilon(C_x)) = 1$  for all  $\varepsilon > 0$ , where  $B_\varepsilon(C_x)$  denotes the  $\varepsilon$ -neighborhood around  $C_x$  in  $X$ . If the set  $C_x$  does not admit a proper subset which is likewise a center of attraction of the motion  $f(t, x)$  as  $t \rightarrow +\infty$ , then  $C_x$  is called the *minimal center of attraction of the motion*  $f(t, x)$  as  $t \rightarrow +\infty$ .

First of all, by the classical Cantor-Baire theorem and Zorn's lemma we can obtain the following basic existence lemma.

**Lemma 1.2.** *Let  $f: \mathbb{R} \times X \rightarrow X$  be a  $C^0$ -semi flow on a metric space  $X$ . Then each Lagrange stable motion  $f(t, x)$  always possesses the minimal center of attraction.*

From now on, by  $C_x$  we will understand the minimal center of attraction of a Lagrange stable motion  $f(t, x)$  as  $t \rightarrow +\infty$ . In [16], Karl Sigmund gave an intrinsic characterization for  $C_x$ . In this paper, we shall study the generic property and chaotic dynamics occurring in  $C_x$  for a Lagrange stable motion  $f(t, x)$ .

Just as the existence of one point which is topologically transitive implies that a residual set is topologically transitive, in 1977 Karl Sigmund raised the following open problem:

**Conjecture 1.3** (K. Sigmund 1977 [16, Remark 4]). For any homeomorphism  $f$  of a compact metric space  $X$ , the set of points  $x \in X$  with  $C_x = X$ , if nonempty, is residual in  $X$ .

Since a residual set contains a dense and  $G_\delta$  subset of  $X$ , it is very large from the viewpoint of topology. Although Sigmund's conjecture is of interest, there has not been any progresses on it since 1977 except  $f$  satisfies the "specification" property ([16, Proposition 6]). In Section 2, we will present a positive solution to Sigmund's conjecture without any imposed shadowing assumption like specification, which is stated as follows:

**Theorem 1.4.** *Let  $f: \mathbb{R}_+ \times X \rightarrow X$  be a  $C^0$ -semi flow on a Polish space  $X$ . If some motion  $f(t, x)$  is such that  $C_x = X$ , then the set  $\{x \in X: C_x = X\}$  is residual in  $X$ .*

Our argument of Theorem 1.4 below also works for discrete-time case. Thus we can obtain the following:

**Corollary.** *For any continuous transformation  $f$  of a Polish space  $X$ , the set of points  $x \in X$  with  $C_x = X$ , if nonempty, is residual in  $X$ .*

We now turn to some applications of Theorem 1.4. For our convenience, we first introduce two notions for a  $C^0$ -semi flow  $f: \mathbb{R}_+ \times X \rightarrow X$  on a Polish space  $X$ .

**Definition 1.5.** An  $f$ -invariant subset  $\Lambda$  of  $X$  is referred to as *generic* if there exists some point  $x \in \Lambda$  with  $\Lambda = C_x$ .

According to Conjecture 1.3 (or precisely speaking Theorem 1.4) for any generic minimal center of attraction  $C_x$  of a motion  $f(t, x)$ , the set of points  $y \in C_x$  with  $C_x = C_y$  is residual in  $C_x$ .

**Definition 1.6.** We say that a motion  $f(t, x)$  is *chaotic* for  $f$  if there can be found some point  $y \in X$  such that

$$\liminf_{t \rightarrow +\infty} d(f(t, x), y) = 0, \quad \limsup_{t \rightarrow +\infty} d(f(t, x), y) > 0$$

and

$$\liminf_{t \rightarrow +\infty} d(f(t, x), f(t, y)) = 0, \quad \limsup_{t \rightarrow +\infty} d(f(t, x), f(t, y)) > 0.$$

By using Theorem 1.4, we will show that if  $C_x$  is not generic, then the chaotic behavior occurs near  $C_x$ ; see Theorems 3.1 and 3.2 stated and proved in Section 3. On the other hand whenever  $C_x$  is generic and it is not “very simple”, then chaotic motions are generic in  $C_x$ ; that is the following

**Theorem 1.7.** *Let  $f(t, p)$  be a Lagrange stable motion in a Polish space  $X$ . If  $C_p$  is generic and itself is not a minimal subset of  $(X, f)$ , then there exists a residual subset  $S$  of  $C_p$  such that  $f(t, x)$  is chaotic for each  $x \in S$ .*

In Section 4 we will consider a relationship of the minimal center of attraction of a motion with the pointwise recurrence.

Finally we shall consider the multiply attracting of the minimal center of attraction of a motion in Section 5.

## 2. Proof of Sigmund’s conjecture

This section will be devoted to proving Karl Sigmund’s Conjecture 1.3 in the continuous-time case; that is, Theorem 1.4.

Recall that if a continuous surjective map  $T: X \rightarrow X$  is topologically transitive, then for a countable basis  $U_1, U_2, \dots, U_n, \dots$  of the underlying space  $X$ , the set

$$\{x \in X \mid \overline{O_T(x)} = X\} = \bigcap_{n=1}^{+\infty} \bigcup_{m=0}^{+\infty} T^{-m}(U_n), \quad \text{where } O_T(x) = \{T^n x : n \in \mathbb{Z}_+\},$$

is a dense  $G_\delta$  set in  $X$  because  $\bigcup_{m=0}^{+\infty} T^{-m}(U_n)$  is open and dense in  $X$ . It is easy to see that this standard argument for topological transitivity does not work here for Sigmund’s conjecture (or Theorem 1.4). So we need a new idea that will explore the times of sojourn of a motion in a domain.

Given any  $x \in X$ , let  $\mathcal{U}_x$  be the neighborhood system of the point  $x$  in  $X$ . To prove Conjecture 1.3, we will need a classical result, which shows that  $C_x$  consists of the points  $y \in X$  which are interesting for  $x$ .

**Lemma 2.1** ([10, 15] for  $C^0$ -flow). *Let  $f: \mathbb{R}_+ \times X \rightarrow X$  be a  $C^0$ -semi flow on a metric space  $X$ . Then for any  $x \in X$ , there holds*

$$C_x = \left\{ y \in X \mid \limsup_{T \rightarrow +\infty} \frac{1}{T} \int_0^T \mathbb{1}_U(f(t, x)) dt > 0 \forall U \in \mathcal{U}_y \right\}.$$

*Proof.* For self-closeness, we present an independent proof for this lemma which is shorter than that of [15]. Let  $x \in X$  and write

$$I(x) = \left\{ y \in X \mid \limsup_{T \rightarrow +\infty} \frac{1}{T} \int_0^T \mathbb{1}_U(f(t, x)) dt > 0 \forall U \in \mathcal{U}_y \right\}.$$

We first claim that  $C_x \subseteq I(x)$ . Indeed, for any  $q \in C_x$ , let  $U \in \mathcal{U}_q$  be a neighborhood of  $q$  in  $X$ ; then

$$\limsup_{T \rightarrow +\infty} \frac{1}{T} \int_0^T \mathbb{1}_U(f(t, x)) dt > 0.$$

Otherwise, one would find some  $\varepsilon > 0$  so that

$$\lim_{T \rightarrow +\infty} \frac{1}{T} \int_0^T \mathbb{1}_{B_{3\varepsilon}(q)}(f(t, x)) dt = 0.$$

Further  $C_x - B_{2\varepsilon}(q)$  is a center of attraction of the motion  $f(t, x)$  and we thus arrive at a contradiction to the minimality of  $C_x$ .

Finally we assert that  $C_x \supseteq I(x)$ . By contradiction, let  $q \in I(x) - C_x$  and then we can take some  $\varepsilon > 0$  such that  $d(q, C_x) \geq 3\varepsilon$ . Set

$$N(B_\varepsilon(q)) = \{t \geq 0 \mid f(t, x) \in B_\varepsilon(q) \text{ for } 1 \leq i \leq l\}$$

and

$$N(B_\varepsilon(C_x)) = \{t \geq 0 \mid f(t, x) \in B_\varepsilon(C_x) \text{ for } 1 \leq i \leq l\}.$$

Clearly  $N(B_\varepsilon(q)) \cap N(B_\varepsilon(C_x)) = \emptyset$ . However, by definitions,  $N(B_\varepsilon(q))$  has positive upper density and  $N(B_\varepsilon(C_x))$  has density 1 in  $[0, \infty)$ . This is a contradiction.

The proof of Lemma 2.1 is thus completed.  $\square$

In Definition 1.1 we do not require the  $f$ -invariance of  $C_x$ . However, it is actually  $f$ -invariant by Lemma 2.1.

**Corollary 2.2.** *Let  $f: \mathbb{R}_+ \times X \rightarrow X$  be a  $C^0$ -semi flow on a metric space  $X$ . For any  $x \in X$ ,  $C_x$  is  $f$ -invariant.*

Since for any real number  $\theta > 0$  and any integer  $N \geq 0$  there holds

$$0 \leq \int_0^\theta \mathbb{1}_U(f(N\theta + t, x)) dt \leq \theta,$$

then Lemma 2.1 implies immediately the following.

**Corollary 2.3.** *Let  $f: \mathbb{R}_+ \times X \rightarrow X$  be a  $C^0$ -semi flow on a Polish space  $X$ . For any  $x \in X$  and any  $\theta > 0$ , there holds*

$$C_x = \left\{ y \in X \mid \limsup_{N \ni N \rightarrow +\infty} \frac{1}{N} \int_0^{N\theta} \mathbb{1}_U(f(t, x)) dt > 0 \forall U \in \mathcal{U}_y \right\}.$$

Here  $\mathbb{N} = \{1, 2, \dots\}$ .

*Proof.* The statement follows from that  $T = N_T\theta + r_T$  where  $N_T \in \mathbb{N}$ ,  $0 \leq r_T < \theta$  for any  $T \geq 1$  and that

$$\limsup_{T \rightarrow +\infty} \frac{1}{T} \int_0^T \mathbb{1}_U(f(t, x)) dt = \limsup_{N \ni N \rightarrow +\infty} \frac{1}{N\theta} \int_0^{N\theta} \mathbb{1}_U(f(t, x)) dt$$

for any  $U \in \mathcal{U}_y$  and any  $y \in X$ .  $\square$

We are now ready to prove one of our main statements—Theorem 1.4—by applying Lemma 2.1 and Corollaries 2.2 and 2.3.

*Proof of Theorem 1.4.* Write  $\Theta = \{x \in X : C_x = X\}$  and let  $x \in X$  be such that  $C_x = X$ . Then from Lemma 2.1, it follows that the orbit  $O_f(x) = f(\mathbb{R}_+, x)$  is dense in  $X$  and that  $O_f(x) \subseteq \Theta$  such that for any  $y \in O_f(x)$ ,

$$\limsup_{\mathbb{N} \ni N \rightarrow \infty} \frac{1}{N} \int_0^N \mathbb{1}_U(f(t, x)) dt = \limsup_{\mathbb{N} \ni N \rightarrow \infty} \frac{1}{N} \int_0^N \mathbb{1}_U(f(t, y)) dt,$$

for every nonempty  $U \in \mathcal{T}_X$ , where  $\mathcal{T}_X$  is the topology of the space  $X$ .

Let  $\mathcal{U} = \{U_i\}_{i=1}^\infty$  be an any given countable base of the topology  $\mathcal{T}_X$  of the state space  $X$ . Then by Corollary 2.3, we can choose a sequence of positive integers  $L_i, i = 1, 2, \dots$ , such that

$$\limsup_{\mathbb{N} \ni N \rightarrow \infty} \frac{1}{N} \int_0^N \mathbb{1}_{U_i}(f(t, x)) dt > \frac{1}{L_i},$$

for all  $i = 1, 2, \dots$ . For each  $i$ , write

$$\Theta_i = \left\{ y \in X \mid \forall n_0 \in \mathbb{N}, \exists n > n_0 \text{ with } \int_0^{nL_i} \mathbb{1}_{U_i}(f(t, y)) dt > n \right\}.$$

From the continuity of  $f(t, y)$  with respect to  $(t, y)$  and

$$\Theta_i = \bigcap_{n_0=1}^\infty \bigcup_{n > n_0} \left\{ y \in X \mid \int_0^{nL_i} \mathbb{1}_{U_i}(f(t, y)) dt > n \right\},$$

it follows that  $\Theta_i$  is a  $G_\delta$  subset of  $X$ . Because  $x$  belongs to  $\Theta_i$  by noting  $N = n_N L_i + r_N$  and  $\lim_{N \rightarrow \infty} \frac{r_N}{N} = 0$  where  $0 \leq r_N < L_i$  and then similarly  $O_f(x) \subseteq \Theta_i$ , there follows  $\bigcap_{i=1}^\infty \Theta_i$  is a dense and  $G_\delta$  set in  $X$ .

Since for any  $y \in \bigcap_{i=1}^\infty \Theta_i$  we have

$$\limsup_{\mathbb{N} \ni N \rightarrow \infty} \frac{1}{N} \int_0^N \mathbb{1}_{U_i}(f(t, y)) dt \geq \frac{1}{L_i} > 0$$

for any  $U_i \in \mathcal{U}$  and  $\mathcal{U}$  is a base of the topology  $\mathcal{T}_X$  of the space  $X$ , we see that  $y \in \Theta$  from Corollary 2.3. Therefore,  $\Theta$  is a residual set in  $X$ .

This completes the proof of Theorem 1.4.  $\square$

Finally we mention that the statement of Theorem 1.4 is also valid for a continuous transformation  $f: X \rightarrow X$  of a Polish space  $X$ .

### 3. Li-Yorke chaotic pairs and sensitive dependence on initial data

In this section, we shall apply Theorem 1.4 to the study of chaos of a topological dynamical system on a Polish space  $X$ .

Let  $f: \mathbb{R}_+ \times X \rightarrow X$  be a  $C^0$ -semi flow on the Polish space  $(X, d)$ . Recall that two points  $x, y \in X$  is called a *Li-Yorke chaotic pair* for  $f$  if

$$\limsup_{t \rightarrow +\infty} d(f(t, x), f(t, y)) > 0 \quad \text{and} \quad \liminf_{t \rightarrow +\infty} d(f(t, x), f(t, y)) = 0.$$

That is to say,  $x$  is proximal to  $y$  but not asymptotical. If there can be found an uncountable set  $S \subset X$  such that every pair of points  $x, y \in S, x \neq y$ , is a Li-Yorke chaotic pair for  $f$ , then we say  $f$  is *Li-Yorke chaotic*; see, e.g., Li and Yorke 1975 [14].

### 3.1. Nongeneric case

Let  $f: \mathbb{R}_+ \times X \rightarrow X$  be a  $C^0$ -semi flow on a compact metric space. The following theorem shows that if  $C_x$  has no the generic dynamics in the sense of Definition 1.5, then  $f$  has the chaotic dynamics.

**Theorem 3.1.** *Given any  $x \in X$ , if  $C_x$  is not generic, then one can find some point  $q \in \Delta$ , for any closed  $f$ -invariant subset  $\Delta \subseteq C_x$ , such that  $(x, q)$  is a Li-Yorke chaotic pair for  $f$ .*

*Proof.* Given any  $x \in X$ , let  $C_x$  be not generic in the sense of Definition 1.5. Let  $\Delta$  be an  $f$ -invariant nonempty closed subset of  $C_x$ . Then by Theorem 1.4 it follows that  $x \notin C_x$ . Moreover from Definition 1.1, we can obtain that  $x$  is proximal to  $\Delta$ ; that is,

$$\liminf_{t \rightarrow +\infty} d(f(t, x), f(t, \Delta)) = 0.$$

Then from [1, 6] also [7, Proposition 8.6], it follows that there exists some point  $q \in \Delta$  such that

$$\liminf_{t \rightarrow +\infty} d(f(t, x), f(t, q)) = 0.$$

We claim that  $\limsup_{t \rightarrow +\infty} d(f(t, x), f(t, q)) > 0$ . Otherwise,  $\lim_{t \rightarrow +\infty} d(f(t, x), f(t, q)) = 0$  and thus  $C_x = C_q$ ; and then  $C_x$  is generic by Theorem 1.4.

The proof of Theorem 3.1 is therefore complete.  $\square$

**Corollary 1.** *Given any  $x \in X$ , if  $C_x$  is not generic, then one can find some point  $q \in C_x$  such that  $(x, q)$  is a Li-Yorke chaotic pair for  $f$  and the set*

$$N_f(x, B_\varepsilon(q)) = \{t \geq 0: f(t, x) \in B_\varepsilon(q)\}$$

*is a central set in  $\mathbb{R}_+$ , which has positive upper density.*

*Proof.* Let  $\Delta \subset C_x$  be a minimal set. Then there can be found by Theorem 3.1 a point  $q \in \Delta$  such that  $(x, q)$  is a Li-Yorke chaotic pair for  $f$ . By definition (cf. [7, Definition 8.3] and [5, Definition 7.2])  $N_f(x, B_\varepsilon(q))$  is a central set of  $\mathbb{R}_+$  for each  $\varepsilon > 0$ . In addition, by Lemma 2.1 it follows that  $N_f(x, B_\varepsilon(q))$  has positive upper density. This proves the corollary.  $\square$

**Corollary 2.** *Let there exist a fixed point or a periodic orbit in the minimal center  $C_x$  of attraction of a motion  $f(t, x)$ . Then we can find a Li-Yorke chaotic pair near  $C_x$ .*

*Proof.* First if  $C_x$  is generic in the sense of Definition 1.5, then  $f$  restricted to it is topologically transitive and has a fixed point or a periodic orbit. Then by Huang and Ye 2002 [11, Theorem 4.1], it follows that  $f$  restricted to  $C_x$  is chaotic in the sense of Li and Yorke.

On the other hand, if  $C_x$  is not generic in the sense of Definition 1.5, then from Theorem 3.1 it follows that there always exists a Li-Yorke chaotic pair near  $C_x$ .

The proof of the corollary is thus complete.  $\square$

Recall that a motion  $f(t, x)$  is referred to as a *Birkhoff recurrent motion* of  $f$  if  $\overline{O_f(x)}$  is minimal (cf. [15, 3]). It is also called “uniformly recurrent” in [7] and “almost periodic” of von Neumann in [9] in the discrete-time case.

Motivated by [2, 8, 4] we can obtain the following theorem on sensitivity on initial data near the minimal center of attraction of a motion.

**Theorem 3.2.** *Let  $C_x$ , for a motion  $f(t, x)$ , be not generic. If the Birkhoff recurrent points of  $f$  are dense in  $C_x$ , then  $f$  has the sensitive dependence on initial data near  $C_x$  in the sense that one can find a sensitive constant  $\epsilon > 0$  such that for any  $a \in X$ ,  $\hat{x} \in C_x$  and any  $U \in \mathcal{W}_{\hat{x}}$ , there exists some point  $c \in U$  with  $\limsup_{t \rightarrow +\infty} d(f(t, a), f(t, c)) \geq \epsilon$ .*

*Proof.* Since  $C_x$  is not generic, by Theorem 1.4 it follows that it is not minimal and so it contains at least two different motions of  $f$  far away each other. Thus one can find a number  $\delta > 0$  such that for all  $\hat{x} \in C_x$  there exists a corresponding motion  $f(t, q)$  in  $C_x$ , not necessarily recurrent, such that

$$d(\hat{x}, \overline{O_f(q)}) \geq \delta,$$

where  $d(\hat{x}, A) = \inf_{a \in A} d(\hat{x}, a)$  for any subset  $A$  of  $X$ . We will show that  $f$  has sensitive dependence on initial data with sensitivity constant  $\epsilon = \delta/4$  following the idea of [4, Theorem 4].

For this, we let  $\hat{x}$  be an arbitrary point in  $C_x$  and let  $U$  be an arbitrary neighborhood of  $\hat{x}$  in  $X$ . Since the Birkhoff recurrent motions of  $(X, f)$  are dense in  $C_x$  from assumption of the theorem, there exists a Birkhoff recurrent point  $p \in U \cap B_{\epsilon/2}(\hat{x}) \cap C_x$ , where  $B_r(\hat{x})$  is the open ball of radius  $r$  centered at  $\hat{x}$  in  $X$ . As we noted above, there must exist another point  $q \in C_x$  whose orbit  $O_f(q)$  is of distance at least  $4\epsilon$  from the given point  $\hat{x}$ .

Let  $\eta > 0$  be such that  $\eta < \epsilon/2$ . Then from the Birkhoff recurrence of the motion  $f(t, p)$ , it follows that one can find a constant  $T = T(\eta, p) > 0$  such that for any  $\gamma \geq 0$ , there is some moment  $t_\gamma \in [\gamma, \gamma + T)$  verifying that

$$d(p, f^{t_\gamma}(p)) < \eta.$$

For the given  $q$ , we simply write

$$V = \bigcap_{t \in [0, 2T)} f^{-t}(B_\epsilon(f^t(q))), \quad \text{where } f^{-t}(\cdot) = f(t, \cdot)^{-1}.$$

Clearly from the continuity of topological flow, it follows that  $V$  is a neighborhood of  $q$  in  $X$  but not necessarily open, and it is nonempty since  $q \in V$ .

Since  $C_x$  is the minimal center of attraction of the motion  $f(t, x)$ , from Lemma 2.1 it follows that there is at least one point  $z \in U \cap B_\epsilon(\hat{x})$  such that  $f^N(z) \in V$  for some sufficiently large number  $N \gg T$ . Let

$$N = jT - r \quad \text{where } 0 \leq r < T, \quad j \in \mathbb{N},$$

and

$$t_{jT} \in [jT, (j+1)T) \quad \text{such that } d(p, f^{t_{jT}}(p)) < \eta.$$

Then  $0 \leq t_{jT} - N < 2T$ .

By construction, one has

$$f^{t_{jT}}(z) = f^{t_{jT}-N}(f^N(z)) \in f^{t_{jT}-N}(V) \subseteq B_\epsilon(f^{t_{jT}-N}(q)).$$

From the triangle inequality of metric, it follows that

$$\begin{aligned} d(f^{t_{jT}}(p), f^{t_{jT}}(z)) &\geq d(p, f^{t_{jT}}(z)) - \eta \\ &\geq d(\hat{x}, f^{t_{jT}}(z)) - d(p, \hat{x}) - \eta \\ &\geq d(\hat{x}, f^{t_{jT}-N}(q)) - d(f^{t_{jT}-N}(q), f^{t_{jT}}(z)) - d(p, \hat{x}) - \eta. \end{aligned}$$

Consequently, since  $\eta < \epsilon/2$ ,  $p \in B_{\epsilon/2}(\hat{x})$  and  $f^{t_{jT}}(z) \in B_{\epsilon}(f^{t_{jT}-N}(q))$ , it holds that

$$d(f^{t_{jT}}(p), f^{t_{jT}}(z)) > 2\epsilon.$$

Therefore from the triangle inequality again, one can obtain either

$$d(f^{t_{jT}}(\hat{x}), f^{t_{jT}}(z)) > \epsilon$$

or

$$d(f^{t_{jT}}(\hat{x}), f^{t_{jT}}(p)) > \epsilon.$$

Repeating this argument for another likewise  $N$  bigger than  $(j+2)T$ , one can find a sequence  $t_n = j_n T \uparrow +\infty$  as  $n \rightarrow +\infty$  such that either

$$d(f^{t_n}(\hat{x}), f^{t_n}(z)) > \epsilon$$

or

$$d(f^{t_n}(\hat{x}), f^{t_n}(p)) > \epsilon,$$

for all  $n \geq 1$ . Thus in either case, we have found a point  $\hat{y} \in U$  such that

$$\limsup_{t \rightarrow +\infty} d(f^t(\hat{x}), f^t(\hat{y})) \geq \epsilon.$$

Now for any  $a \in X$ , using the triangle inequality once more, we see either

$$\limsup_{t \rightarrow +\infty} d(f^t(\hat{x}), f^t(a)) \geq \frac{\epsilon}{3}$$

or

$$\limsup_{t \rightarrow +\infty} d(f^t(\hat{y}), f^t(a)) \geq \frac{\epsilon}{3}.$$

Since  $\hat{x}, U$  both are arbitrary and  $\hat{y} \in U$ , hence the proof of Theorem 3.2 is complete.  $\square$

We note that if  $(C_x, f)$  is distal (cf. [7, Definition 8.2]) and not minimal and even not topologically transitive, then the conditions of Theorem 3.2 hold; i.e., the Birkhoff recurrent points are dense in  $C_x$ . In fact,  $C_x$  consists of Birkhoff recurrent points ([7, Corollary of Theorem 8.7]) and it is not topologically transitive.

### 3.2. Generic case

Let  $f: \mathbb{R}_+ \times X \rightarrow X$  be a  $C^0$ -semi flow on a Polish space and  $f(t, p)$  a Lagrange stable motion. Then the minimal center  $C_p$  of attraction of the motion  $f(t, p)$  is always existent. Theorem 1.7 is just a corollary of the following

**Theorem 3.3.** *Let  $C_p$  be generic and not a minimal subset of  $(X, f)$ . Then there exists a residual subset  $S$  of  $C_p$  such that for any  $x \in S$  and any minimal subset  $\Lambda \subset C_p$ , there corresponds some point  $y \in \Lambda$  with the properties:  $x, y$  form a Li-Yorke chaotic pair for  $f$  and*

$$\liminf_{t \rightarrow +\infty} d(f(t, x), y) = 0 \quad \text{and} \quad \limsup_{t \rightarrow +\infty} d(f(t, x), y) \geq \frac{1}{2} \text{diam}(C_p).$$

*Proof.* Since  $C_p$  is generic in the sense of Definition 1.5, there exists some point  $q \in C_p$  with  $C_q = C_p$ . Therefore by Theorem 1.4, there is a residual subset  $S$  of  $C_p$  such that  $C_x = C_p$  for all point  $x$  in  $S$ . Because  $C_p$  is not a minimal subset of  $X$  by hypothesis of Theorem 1.7,  $\overline{O_f(x)}$  is not minimal for each  $x \in S$ .

Let  $\Lambda$  be a minimal subset of  $C_p$ . Then each  $x \in S$  is proximal to  $\Lambda$ . Moreover by [7, Theorem 8.7], it follows that for every  $x \in S$ , there corresponds some point  $y \in \Lambda$  such that  $x$  is proximal to  $y$  and  $f(t, x)$  is Birkhoff recurrent (or uniformly recurrent). Clearly,  $x$  and  $y$  is a Li-Yorke chaotic pair for  $f$ . In addition, from Lemma 2.1 follows that

$$\limsup_{t \rightarrow +\infty} d(f(t, x), y) \geq \frac{1}{2} \text{diam}(C_p)$$

and

$$\liminf_{t \rightarrow +\infty} d(f(t, x), y) = 0.$$

This completes the proof of Theorem 3.3.  $\square$

Therefore by Theorems 3.1 and 3.3, it follows that every Lagrange stable motion  $f(t, x)$  is chaotic in the sense of Definition 1.6 if its minimal center  $C_x$  of attraction is not a minimal subset of  $(X, f)$ .

#### 4. Quasi-weakly almost periodic motion

In this section, we let  $f: \mathbb{R}_+ \times X \rightarrow X$  be a  $C^0$ -semi flow on the compact metric space  $X$ .

**Definition 4.1** (Huang-Zhou 2012 [12]). A point  $x$  in  $X$  is called a *quasi-weakly almost periodic point* of  $f$  if for any  $\varepsilon > 0$  there exists an integer  $N = N(\varepsilon) \geq 1$  and an increasing positive integer sequence  $\{n_j\}$  with the property that for each  $j$  there are  $0 = t_0 < t_1 < \dots < t_{n_j} < n_j N$  with  $t_{i+1} - t_i \geq 1$  such that  $f(t_i, x) \in B_\varepsilon(x)$  for all  $i = 1, \dots, n_j$ .

As results of the statements of Lemma 2.1 and Theorem 1.4, we can obtain the following two results.

**Proposition 4.2.** The following statements are equivalent to each other.

- (1)  $x \in X$  is a quasi-weakly almost periodic point of  $f$ .
- (2)  $x \in C_x$ .

*Proof.* (1) $\Rightarrow$ (2) follows from Lemma 2.1. (2) $\Rightarrow$ (1) follows from Lemma 2.1 and the local section theorem of Bebutov (cf. [15, Theorem V.2.14]).  $\square$

Proposition 4.2 has been proved in [12] in the case where  $x$  is a Poisson stable point of  $f$ , i.e., there is a sequence  $t_n \rightarrow \infty$  such that  $f(t_n, x) \rightarrow x$  as  $n \rightarrow \infty$ .

**Proposition 4.3.** If  $x \in C_x$ , then the set  $\{y \in C_x \mid y \in C_y = C_x\}$  is dense and  $G_\delta$  relative to the subspace  $C_x$ .

*Proof.* The statement follows from Theorem 1.4 with  $C_x$  replacing of  $X$ .  $\square$

## 5. Minimal center of multi-attraction of a motion

Let  $f: \mathbb{R}_+ \times X \rightarrow X$  be a  $C^0$ -semi flow on a Polish space  $X$ . From now on, by  $\lambda(dt)$  we denote the standard Haar (Lebesgue) measure on  $\mathbb{R}$ . We will need the following simple but useful fact.

**Lemma 5.1.** *Let  $S$  be a measurable subset of  $\mathbb{R}_+$  and  $\tau > 0$ . If  $S$  has the density  $\alpha$ , i.e.,*

$$D(S) := \lim_{T \rightarrow +\infty} \frac{\lambda(S \cap [0, T])}{T} = \alpha,$$

then  $\tau S = \{\tau t: t \in S\}$  also has the density  $\alpha$  in  $\mathbb{R}_+$ .

*Proof.* Let  $\tau > 0$  be any given. Since  $\lambda(\tau A) = \tau \lambda(A)$  and  $\lambda(\tau S \cap [0, T]) = \tau \lambda(S \cap [0, T\tau^{-1}])$ , hence it follows that  $D(\tau S) = 1$ . This proves the lemma.  $\square$

It should be noted here that there is no an analogous result for the discrete-time  $\mathbb{Z}_+$ ; for example,  $S = \{0, 2, 4, 6, \dots\}$  has the density  $\frac{1}{2}$  but  $\frac{1}{2}S$  has the density 1 in  $\mathbb{Z}_+$ .

The following lemma shows that every minimal center of attraction of a motion  $f(t, x)$  is multiply attracting as  $t \rightarrow +\infty$ .

**Lemma 5.2.** *Let  $C_x$  be existent for a motion  $f(t, x)$  as  $t \rightarrow +\infty$ . Then for any  $t_1 > 0, \dots, t_l > 0$  where  $l \in \mathbb{N}$  and any  $\varepsilon > 0$ ,*

$$\lim_{T \rightarrow +\infty} \frac{1}{T} \int_0^T \mathbb{1}_{B_\varepsilon(C_x)}(f(t_1 t, x)) \cdots \mathbb{1}_{B_\varepsilon(C_x)}(f(t_l t, x)) dt = 1.$$

*Proof.* For any  $\tau > 0$  and any open set  $U$ , define an open set

$$N_\tau(x, U) = \{t: 0 \leq t < +\infty, f(\tau t, x) \in U\}.$$

It is easy to check that  $\tau^{-1}N_1(x, U) = N_\tau(x, U)$ . Then by Lemma 5.1 and Definition 1.1, it follows that  $N_{t_1}(x, B_\varepsilon(C_x)), \dots, N_{t_l}(x, B_\varepsilon(C_x))$  all have the density 1. Thus

$$N_{t_1, \dots, t_l}(x, B_\varepsilon(C_x)) := N_{t_1}(x, B_\varepsilon(C_x)) \cap \cdots \cap N_{t_l}(x, B_\varepsilon(C_x))$$

also has the density 1 in  $\mathbb{R}_+$ . This completes the proof of Lemma 5.2.  $\square$

Recall that a motion  $f(t, x)$  is said to be *Lagrange stable* as  $t \rightarrow +\infty$  if the orbit-closure  $\overline{O_f(x)}$  is compact in  $X$  (cf. [15]). As a direct result of Lemma 5.2, we can obtain the following.

**Corollary 5.3.** *For any Lagrange stable motion  $f(t, x)$  as  $t \rightarrow +\infty$ ,  $C_x$  is the minimal closed subset of  $X$  such that for any  $t_1 > 0, \dots, t_l > 0$  and any  $\varepsilon > 0$ ,*

$$\lim_{T \rightarrow +\infty} \frac{1}{T} \int_0^T \mathbb{1}_{B_\varepsilon(C_x)}(f(t_1 t, x)) \cdots \mathbb{1}_{B_\varepsilon(C_x)}(f(t_l t, x)) dt = 1.$$

This result shows that  $C_x$  is the “minimal center of multi-attraction” of a Lagrange stable motion  $f(t, x)$  as  $t \rightarrow +\infty$ .

## Acknowledgment

This work was supported partly by National Natural Science Foundation of China grant #11431012, 11271183 and PAPD of Jiangsu Higher Education Institutions.

## References

- [1] J. Auslander, *On the proximal relation in topological dynamics*, Proc. Amer. Math. Soc., **11** (1960), 890–895.
- [2] J. Banks, J. Brooks, G. Cairns, G. Davis, and P. Stacey, *On Devaney's definition of chaos*, Amer. Math. Monthly, **99** (1992), 332–334.
- [3] B. Chen and X. Dai, *On uniformly recurrent motions of topological semigroup actions*, Discret. Contin. Dyn. Syst., in press, 2015.
- [4] X. Dai, *Chaotic dynamics of continuous-time topological semiflow on Polish spaces*, J. Differential Equations, **258** (2015), 2794–2805.
- [5] X. Dai, *On the Furstenberg-Zimmer structure theorems for Noetherian modules over syndetic rings*, Preprint, 2015.
- [6] R. Ellis, *A semigroup associated with a transformation group*, Trans. Amer. Math. Soc., **94** (1960), 272–281.
- [7] H. Furstenberg, *Recurrence in Ergodic Theory and Combinatorial Number Theory*, Princeton University Press, Princeton, New Jersey, 1981.
- [8] E. Glasner and B. Weiss, *Sensitive dependence on initial conditions*, Nonlinearity, **6** (1993), 1067–1075.
- [9] W.H. Gottschalk and G.A. Hedlund, *Topological Dynamics*, Amer. Math. Soc. Colloq. Publ., vol. 36, Amer. Math. Soc., Providence, RI, 1955.
- [10] H.F. Hilmy, *Sur les centres d'attraction minimaux des systèmes dynamiques*, Compositio Math., **3** (1936), 227–238.
- [11] W. Huang and X.-D. Ye, *Devaney's chaos or 2-scattering implies Li-Yorke's chaos*, Topology and its Appl., **117** (2012), 259–272.
- [12] Y. Huang and Z.-L. Zhou, *Two new recurrent levels for  $C^0$ -flows*, Acta Appl. Math., **118** (2012), 125–145.
- [13] K. Jacobs, *Einige Grundbegriffe der topologischen Dynamik*, Math. Phys. Semesterberichte, **14** (1967), 129–150.
- [14] T. Li and J. Yorke, *Period 3 implies chaos*, Amer. Math. Monthly, **82** (1975), 985–992.
- [15] V.V. Nemytskii and V.V. Stepanov, *Qualitative Theory of Differential Equations*, Princeton University Press, Princeton, New Jersey 1960.
- [16] K. Sigmund, *On minimal centers of attraction and generic points*, J. Reine Angew. Math., **295** (1977), 72–79.