Notes on a theorem of Naji

Lorenzo Traldi Lafayette College Easton, Pennsylvania USA

Abstract

We present a new proof of an algebraic characterization of circle graphs due to W. Naji. For bipartite graphs, Naji's theorem is equivalent to an algebraic characterization of planar matroids due to J. Geelen and B. Gerards. Naji's theorem also yields an algebraic characterization of permutation graphs.

Keywords. circle graph, graphic matroid, permutation graph, split decomposition

Mathematics Subject Classification. 05C50

1 Introduction

This paper is concerned with the following notion.

Definition 1 Let $W = w_1...w_{2n}$ be a double occurrence word in the letters $v_1, ..., v_n$. The interlacement graph $\mathcal{I}(W)$ is the simple graph with vertex-set $V = \{v_1, ..., v_n\}$, in which v_i and v_j are adjacent if and only if they are interlaced in W, i.e., they appear in W in the order $v_i v_j v_i v_j$ or $v_j v_i v_j v_i$. A circle graph is a simple graph that can be realized as the interlacement graph of some double occurrence word.

As far as we know, the idea of interlacement first appeared in the form of a symmetric matrix used in Brahana's 1921 study of curves on surfaces [6]. Interlacement graphs were studied by Zelinka [20], who credited the idea to Kotzig. During the subsequent decades several researchers discussed graphs and matrices defined using interlacement. Cohn and Lempel [7] and Even and Itai [9] used them to analyze permutations, and Bouchet [1] and Read and Rosenstiehl [18] used them to study Gauss' problem of characterizing generic self-intersecting curves in the plane. Recognition algorithms for circle graphs have been introduced by Bouchet [2], Gioan, Paul, Tedder and Corneil [14], Naji [16, 17] and Spinrad [19].

Although Naji's is not the best of the circle graph recognition algorithms in terms of computational complexity, it is particularly interesting for two reasons.

The first reason is that Naji's characterization is only indirectly algorithmic; it involves a system of equations that may be defined for any graph, which is only solvable for circle graphs. The second reason is that the two known proofs of the theorem are quite long. The original argument ends on p. 173 of Naji's thesis [16]. A much shorter argument was given by Gasse [11], but Gasse's argument requires Bouchet's circle graphs obstructions theorem [4], which itself has a long and difficult proof.

A couple of years ago, Geelen and Gerards [13] characterized graphic matroids by a system of equations that resembles Naji's system of equations. (Indeed, they mention that Naji's theorem motivated their result.) The resemblance is limited to the equations; there is a striking contrast between their concise, well-motivated proof and Naji's long, detailed argument. This contrast encouraged us to look for an alternative proof of Naji's theorem; we eventually developed the one presented below. Although our argument is certainly not as elegant as the proof of Geelen and Gerards, it is shorter than either Naji's original proof or the combination of a proof of Bouchet's obstructions theorem and Gasse's derivation of Naji's theorem.

After proving Naji's theorem for circle graphs in general, we briefly discuss two special cases. First, the restriction of Naji's theorem to bipartite graphs is equivalent to the restriction of the Geelen-Gerards characterization to planar matroids. Second, Naji's theorem also characterizes permutation graphs.

Before proceeding we should thank Jim Geelen for his comments on Naji's theorem. In particular, he pointed out that although all circle graphs have solutions of Naji's equations that arise naturally from double occurrence words, some circle graphs also have other Naji solutions that do not seem so natural. He conjectured that these other solutions might correspond in some way to splits. (See Sections 2 and 3 for definitions.) Although we do not address Geelen's conjecture directly we do provide some indirect evidence for it, as the first step of our proof is to show that none of these other solutions occur in circle graphs that have no splits.

2 Naji's equations and their solutions

We begin with some definitions.

Definition 2 [16, 17] Let G be a simple graph. For each pair of distinct vertices v and w of G, let $\beta(v, w)$ and $\beta(w, v)$ be distinct variables. Then the Naji equations for G are the following.

(a) For each edge vw of G, $\beta(v, w) + \beta(w, v) = 1$.

(b) For each edge vw of G and each vertex x not adjacent to either v or w, $\beta(x, v) + \beta(x, w) = 0.$

(c) For each pair of edges vw, vx of G such that wx is not an edge, $\beta(v, w) + \beta(v, x) + \beta(w, x) + \beta(x, w) = 1$.

If the Naji equations of G have a solution over GF(2), the field with two elements, then any such solution is a Naji solution and G is a Naji graph.

Definition 3 For each vertex $v \in V(G)$, let $\delta(v)$ be given by $\delta(v)(v, w) = 1$ $\forall w \neq v, \ \delta(v)(w, v) = 1$ if $vw \in E(G)$, and $\delta(v)(x, y) = 0$ otherwise.

Definition 4 Let ρ be given by $\rho(v, w) = 1 \quad \forall v \neq w \in V(G)$.

The following proposition is easily verified.

Proposition 5 Let β be a Naji solution for G. Then $\beta + \rho$ is a Naji solution and for each $v \in V(G)$, $\beta + \delta(v)$ is a Naji solution.

Corollary 6 If G is a Naji graph and $v \in V(G)$, then for every subset $X \subseteq V(G) - v$, G has a Naji solution β with $X = \{x \in V(G) \mid \beta(x, v) = 1\}$.

Proof. Begin with an arbitrary Naji solution β and consider the Naji solution

$$\beta + \sum_{\substack{x \in X \\ \beta(x,v) = 0}} \delta(x) + \sum_{\substack{y \notin X \cup \{v\} \\ \beta(y,v) = 1}} \delta(y).$$

The next proposition is also easy to verify.

Proposition 7 The set $\{\rho\} \cup \{\delta(v) \mid v \in V(G)\}$ is linearly independent over GF(2) unless G is a complete graph, a star or a trivial (edgeless) graph. In each of these exceptional cases the rank of $\{\rho\} \cup \{\delta(v) \mid v \in V(G)\}$ is |V(G)|.

Every circle graph is a Naji graph. To see why, consider a double occurrence word W. An orientation of W is given by arbitrarily designating one appearance of each letter as "initial"; the other appearance is "terminal." We use the notation v^{in} and v^{out} for the initial and terminal appearances of v, respectively. For each orientation of W, there is an associated Naji solution defined by: $\beta(v, w) = 0$ if and only if when we cyclically permute W to begin with v^{in} , w^{out} precedes v^{out} . We leave to the reader the job of verifying that this definition yields a solution of the Naji equations of $\mathcal{I}(W)$. The reader can also easily verify the following proposition.

Proposition 8 Let β be the Naji solution of $\mathcal{I}(W)$ corresponding to an orientation of W. Then cyclically permuting W results in the same Naji solution β , and reversing W results in the Naji solution $\beta + \rho$. Also, if $v \in V(G)$ then $\beta + \delta(v)$ is the Naji solution corresponding to the orientation obtained by interchanging the appearances of v^{in} and v^{out} in W.

Proposition 9 Suppose v and w are two vertices of a connected circle graph $G = \mathcal{I}(W)$. Then v and w appear consecutively in W if and only if there is an orientation of W for which the corresponding Naji solution has $\beta(x, v) = \beta(x, w)$ $\forall x \notin \{v, w\}$.

Proof. If W has an orientation in which v^{out} and w^{out} appear consecutively, then the corresponding Naji solution has $\beta(x, v) = \beta(x, w) \ \forall x \notin \{v, w\}.$

For the converse, suppose W can be oriented in such a way that the corresponding Naji solution has $\beta(x, v) = \beta(x, w) \ \forall x \notin \{v, w\}$. Permute W cyclically so that it is in the form $Av^{out}Bw^{out}$; this permutation does not affect the associated Naji solution. If A or B is empty, then v^{out} and w^{out} are consecutive. Suppose instead that A and B are both nonempty; we derive contradictions in all cases. Consider an arbitrary $x \notin \{v, w\}$. If x^{in} appears in A and x^{out} appears in B then $\beta(x, v) = 0 \neq \beta(x, w)$, a contradiction. Also, if x^{out} appears in A and x^{in} appears in B then $\beta(x, v) = 1 \neq \beta(x, w)$, another contradiction. Consequently x must appear twice in one of A, B. If neither v^{in} nor w^{in} appears in A, it follows that no vertex that appears in A is interlaced with a vertex that does not appear in A; but then G is not connected, an impossibility. Similarly, if neither v^{in} nor w^{in} appears in B then G is not connected. Consequently one of v^{in} , w^{in} appears in A and the other appears in B. If v^{in} appears in A then the subwords Av^{out} and Bw^{out} are separate double occurrence words; and if w^{in} appears in A then the subwords $v^{out}B$ and $w^{out}A$ are separate double occurrence words. Either way, the connectedness of G is contradicted.

3 Prime graphs and splits

Cunningham's split decomposition [8] is of fundamental importance in analyzing circle graphs.

Definition 10 A split (X, Y) of a simple graph G consists of disjoint subsets $X, Y \subset V(G)$ such that $|X|, |Y| \ge 2, X \cap Y = \emptyset, X \cup Y = V(G)$ and the edges of G between X and Y define a complete bipartite graph. A connected simple graph with no split is prime.

Connected graphs of order 1, 2 or 3 are prime for the trivial reason that 2+2>3, and it is easy to see that no graph of order 4 is prime. Consequently the interesting part of the theory of prime graphs begins with order 5.

If G has a split (X, Y) then Cunningham called G the *composition* of two smaller graphs, G_X and G_Y . G_X is obtained from the induced subgraph G[X]by attaching a new vertex y, whose neighborhood consists of those vertices from X that have neighbors in Y. G_Y is obtained in the same way from G[Y], except the new vertex is denoted x. The new vertices x and y are called *markers*. Cunningham actually used only one marker but it is convenient to use two for the simple reason that the orders of G_X and G_Y are strictly less than the order of G, so inductive arguments can be set up in a natural way.

If a graph has splits then Cunningham showed that the graph can be decomposed in an essentially unique way as a composition of smaller graphs. This unique decomposition is both elegant and useful, but we do not discuss it in detail because uniqueness of the decomposition is not crucial here.

The following simple proposition of Bouchet [2] allows us to focus our attention on prime circle graphs. **Proposition 11** [2] If G has a split (X, Y), then G is a circle graph if and only if G_X and G_Y are both circle graphs.

Proof. If $G_X = \mathcal{I}(W_1 x W_3 x)$ and $G_Y = \mathcal{I}(W_2 y W_4 y)$ then $G = \mathcal{I}(W_1 W_2 W_3 W_4)$.

Suppose conversely that G is a circle graph and $G = \mathcal{I}(W_1W_2...W_{2m})$ where every W_i is nonempty and contains letters from X (resp. Y) if i is odd (resp. even). If no edge of G connects X to Y then $G_X = \mathcal{I}(xxW_1W_3...W_{2m-1})$ and $G_Y = \mathcal{I}(yyW_2W_4...W_{2m})$. Otherwise, let x_0y_0 be an edge of G. After cyclic permutation we may presume that x_0 appears in W_1 and W_{2i-1} , and y_0 appears in W_{2j} and W_{2k} , with i > 1 and j < k. The fact that x_0y_0 is an edge implies that $1 \le j < i \le k$. Then $G_X = \mathcal{I}(W_1...W_{2j-1}yW_{2j+1}...W_{2k-1}yW_{2k+1}...W_{2m-1})$ and $G_Y = \mathcal{I}(xW_2...W_{2i-2}xW_{2i}...W_{2m})$.

4 Local complementation

Definition 12 If v is a vertex of a simple graph G then the local complement G^v is the graph obtained from G by reversing the adjacency status of every pair of neighbors of v. A graph that can be obtained from G through some sequence of local complementations is locally equivalent to G.

That is, G^v includes the same edges vw and wx as G, so long as $x \notin N(v)$; but if $y \neq z \in N(v)$ then $yz \in E(G^v)$ if and only if $yz \notin E(G)$. (Here N(v) denotes the open neighborhood of v, $N(v) = \{w \in V(G) \mid vw \in E(G)\}$.)

Local complementation is important in the theory of circle graphs because the following propositions indicate that inductive proofs involving prime circle graphs can be set up using local complementation. The first two appeared in Bouchet's discussion of his circle graph recognition algorithm [2].

Proposition 13 [2] If $v \in V(G)$ and (X, Y) is a partition of V(G) then (X, Y) is a split of G if and only if (X, Y) is a split of G^v .

Proof. We leave the easy verification to the reader.

Proposition 14 [2] If G and H are locally equivalent then G is a circle graph if and only if H is a circle graph.

Proof. If G is $\mathcal{I}(vW_1vW_2)$ then G^v is $\mathcal{I}(vW_1vW_2)$, where the arrow indicates reversal of the subword.

The next proposition appeared in Gasse's derivation of Naji's theorem [11].

Proposition 15 [11] If G and H are locally equivalent then G is a Naji graph if and only if H is a Naji graph.

Proof. Suppose G is a Naji graph. Then according to Corollary 6, G has a Naji solution β such that $\beta(x, v) = 0$ if and only if $xv \in E(G)$. The reader can easily check that a Naji solution for G^v can then be defined by

$$\beta^{v}(x,y) = \begin{cases} \beta(x,y) + \beta(v,y) \text{ if } x \in N(v) \\ \beta(x,y) + \beta(v,x) \text{ if } x \notin N(v), \end{cases}$$

with the understanding that $v \notin N(v)$ and $\beta(v, v) = 0$.

We can say a little more.

Proposition 16 If $v \in V(G)$ then G has the same number of Naji solutions as G^{v} .

Proof. If G is not a Naji graph then Proposition 15 tells us that G^v is not a Naji graph either.

Let $\mathcal{B}_0(G)$ be the set that includes all the Naji solutions β of G with the property that $\beta(x, v) = 0$ if and only if $x \in N(v)$. Suppose β is an arbitrary Naji solution of G. Let $X = \{x \in N(v) \mid \beta(x, v) = 1\}$ and $Y = \{y \notin N(v) \cup \{v\} \mid \beta(y, v) = 0\}$. Then

$$\beta + \sum_{x \in X} \delta_G(x) + \sum_{y \in Y} \delta_G(y) \in \mathcal{B}_0(G).$$

Moreover, if $W \subseteq V(G - v)$ is any subset other than $X \cup Y$ then

$$\beta + \sum_{w \in W} \delta_G(w) \notin \mathcal{B}_0(G).$$

As no two subsets $W \subseteq V(G-v)$ yield the same sum $\sum_{w \in W} \delta_G(w)$, we conclude that $|\mathcal{B}_0(G)| \cdot 2^{|V(G)|-1}$ is the number of Naji solutions of G. The same argument applies to G^v , so it suffices to prove that $|\mathcal{B}_0(G^v)| = |\mathcal{B}_0(G)|$.

Suppose $\beta \in \mathcal{B}_0(G)$, and let β^v be the Naji solution of G^v defined in Proposition 15. Notice that if $x \in N(v)$, then $\beta^v(x,v) = \beta(x,v) = 0$. Also, if $y \notin N(v) \cup \{v\}$ then $\beta^v(y,v) = \beta(y,v) + \beta(v,y) = 1 + \beta(v,y)$. Consequently if we let $Y_\beta = \{y \notin N(v) \cup \{v\} \mid \beta(v,y) = 1\}$ then

$$f(\beta) \equiv \beta^v + \sum_{y \in Y_\beta} \delta_{G^v}(y) \in \mathcal{B}_0(G^v).$$

We claim that $f : \mathcal{B}_0(G) \to \mathcal{B}_0(G^v)$ is injective, and consequently $|\mathcal{B}_0(G^v)| \ge |\mathcal{B}_0(G)|$. As $G = (G^v)^v$, the claim suffices to complete the proof.

Suppose $\beta, \beta' \in \mathcal{B}_0(G)$. If $Y_{\beta} \neq Y_{\beta'}$, there is a $z \notin N(v) \cup \{v\}$ with $\beta(v, z) \neq \beta'(v, z)$. Then $\beta^v(v, z) = \beta(v, z) \neq (\beta')^v(v, z) = \beta'(v, z)$. Moreover there is no $y \in Y_{\beta} \cup Y_{\beta'}$ such that $\delta_{G^v}(y)$ has a nonzero (v, z) coordinate, because $v \notin N(z)$ and $v \notin Y_{\beta} \cup Y_{\beta'}$. Consequently $f(\beta)(v, z) \neq f(\beta')(v, z)$.

Now, suppose β , $\beta' \in \mathcal{B}_0(G)$ and $f(\beta) = f(\beta')$. As we just saw, $f(\beta) = f(\beta')$ requires $Y_\beta = Y_{\beta'}$, i.e., $\beta(v, y) = \beta'(v, y) \ \forall y \notin N(v) \cup \{v\}$. As $\beta(y, v) = 0 = \beta'(y, v) \ \forall y \in N(v)$, the Naji equations imply $\beta(v, y) = 1 = \beta'(v, y) \ \forall y \in N(v)$. Consequently $\beta(v, y) = \beta'(v, y) \ \forall y \neq v$. The equalities $f(\beta) = f(\beta')$ and $Y_\beta = Y_{\beta'}$ imply $\beta^v = (\beta')^v$, and the definition of β^v in Proposition 15 makes it clear that the equalities $\beta^v = (\beta')^v$ and $\beta(v, y) = \beta'(v, y) \ \forall y \neq v$ imply $\beta = \beta'$.

The next proposition is more difficult; it was first proved by Bouchet using isotropic systems [2, 3].

Proposition 17 [2] If G is prime and |V(G)| > 5 then there is a locally equivalent graph H and a vertex $v \in V(H)$ such that H - v is prime.

A refined form of Proposition 17 appears in Geelen's thesis [12], which is freely available online. The reader who has not already encountered Proposition 17 is encouraged to read Geelen's account, as the result is stronger and the proof does not require isotropic systems.

Proposition 18 [12, Corollary 5.10] If G is prime and |V(G)| > 5 then either G has a vertex v such that G - v is prime, or G has a degree-2 vertex v such that $G^v - v$ is prime.

5 Step 1 of the proof: uniqueness

As noted in Section 2, it is easy to see that every circle graph is a Naji graph; the interesting part of Naji's theorem is the converse. According to Proposition 11 it suffices to prove the converse for prime Naji graphs. The first step of our proof is the following uniqueness result.

Theorem 19 Let G be a prime Naji graph with $|V(G)| \ge 5$, and let β_0 be any particular Naji solution of G. Then every other Naji solution of G is

$$\beta_0 + \sum_{s \in S} \delta(s) \text{ or } \beta_0 + \rho + \sum_{s \in S} \delta(s)$$

for some subset $S \subseteq V(G)$.

Proof. We leave it to the reader to write down the Naji equations for the graph C_5 and verify that the solution space is of dimension 6. According to Proposition 7, the dimension of the subspace spanned by $\{\rho\} \cup \{\delta(v) \mid v \in V(C_5)\}$ is also 6, so the theorem holds for C_5 . According to Bouchet [2, Lemma 3.1] every prime graph of order 5 is locally equivalent to C_5 , so Proposition 16 tells us that the theorem holds for all prime graphs of order 5.

We proceed using induction on |V(G)| > 5. By Propositions 15 and 17, without loss of generality we may replace G with a locally equivalent graph so that there is a vertex $v \in V(G)$ such that G - v is prime. Let β_1 be some Naji solution for G that is not the same as β_0 . Then β_0 and β_1 define Naji solutions for G - v by restriction, and the inductive hypothesis asserts that $\beta_1|(G - v)$ is

$$(\beta_0 + \sum_{s \in S} \delta(s))|(G - v) \text{ or } (\beta_0 + \rho + \sum_{s \in S} \delta(s))|(G - v)$$

for some subset $S \subseteq V(G - v)$. Replacing β_1 with $\beta_1 + \sum_S \delta(s)$ or $\beta_1 + \rho + \sum_S \delta(s)$, we may presume that $\beta_1 | (G - v) = \beta_0 | (G - v)$. That is,

$$\beta_1(x,y) = \beta_0(x,y) \text{ whenever } v \notin \{x,y\}.$$
(*)

The rest of the proof involves a detailed analysis of the structure of G. We partition V(G - v) into four sets.

- $A = \{a \in V(G v) \mid \beta_0(a, v) = \beta_1(a, v) \text{ and } \beta_0(v, a) = \beta_1(v, a)\}$
- $B = \{b \in V(G v) \mid \beta_0(b, v) = \beta_1(b, v) \text{ and } \beta_0(v, b) \neq \beta_1(v, b)\}$
- $C = \{x \in V(G v) \mid \beta_0(c, v) \neq \beta_1(c, v) \text{ and } \beta_0(v, c) = \beta_1(v, c)\}$
- $D = \{x \in V(G v) \mid \beta_0(d, v) \neq \beta_1(d, v) \text{ and } \beta_0(v, d) \neq \beta_1(v, d)\}$

Claim 1. Both $B \cap N(v) = \emptyset$ and $C \cap N(v) = \emptyset$.

proof: The Naji equations require $\beta_i(v, x) \neq \beta_i(x, v)$ for $i \in \{0, 1\}$ when vx is an edge, and both inequalities cannot hold if $x \in B \cup C$.

Claim 2. Either $A \cap N(v) = \emptyset$ or $C = \emptyset$.

proof: Suppose $a \in A \cap N(v)$ and $c \in C$. If $ac \notin E(G)$ then the Naji equations require $\beta_i(c, a) = \beta_i(c, v)$ for i = 0 and 1. This is not possible, as $\beta_0(c, v) \neq \beta_1(c, v)$ by the definition of C and $\beta_0(c, a) = \beta_1(c, a)$ by (*). Hence $ac \in E(G)$. Then the Naji equations require

$$\beta_i(a,c) + \beta_i(a,v) + \beta_i(v,c) + \beta_i(c,v) = 1$$

for $i \in \{0, 1\}$. Both equations cannot be true as $\beta_0(c, v) \neq \beta_1(c, v)$ and the other terms are all equal.

Claim 3. Either $D \cap N(v) = \emptyset$ or $D - N(v) = \emptyset$.

proof: Suppose $x \in D \cap N(v)$ and $y \in D - N(v)$. If $xy \in E(G)$ then as $vy \notin E(G)$, the Naji equations require

$$\beta_i(x,y) + \beta_i(x,v) + \beta_i(v,y) + \beta_i(y,v) = 1$$

for $i \in \{0,1\}$. Both equations cannot be true as $\beta_0(x,y) = \beta_1(x,y)$ by (*), and the other terms are all unequal. Hence $xy \notin E(G)$, so $\beta_i(y,x) = \beta_i(y,v)$ for $i \in \{0,1\}$. Both equations cannot be true as $\beta_0(y,x) = \beta_1(y,x)$ by (*) but the definition of D requires $\beta_0(y,v) \neq \beta_1(y,v)$.

Claim 4. No edge connects $(A \cup C) - N(v)$ to $(B \cup D) - N(v)$.

proof: Suppose $x \in (A \cup C) - N(v)$ is adjacent to $y \in (B \cup D) - N(v)$. Then the Naji equations require $\beta_i(v, x) = \beta_i(v, y)$ for $i \in \{0, 1\}$, but both equations cannot be true as $\beta_0(v, x) = \beta_1(v, x)$ and $\beta_0(v, y) \neq \beta_1(v, y)$.

Claim 5. Suppose $b \in B$ and $x \notin B$ are neighbors. Then $x \in D$.

proof: If $x \in A \cap N(v)$ then as $b \notin N(v)$, the Naji equations require

$$\beta_i(v, x) + \beta_i(b, x) + \beta_i(b, v) + \beta_i(v, b) = 1$$

for $i \in \{0,1\}$. This is impossible as $\beta_0(v,b) \neq \beta_1(v,b)$ and the other terms of the two equations are all equal. Claim 4 now implies that $x \in D$.

Claim 6. If $a \in A \cap N(v)$ then a is adjacent to every element of D, and the other neighbors of a all lie in A.

proof: Suppose $a \in A \cap N(v)$ is not adjacent to $d \in D$. If $d \notin N(v)$ then the Naji equations require $\beta_i(d, a) = \beta_i(d, v)$ for $i \in \{0, 1\}$, an impossibility as $\beta_0(d, a) = \beta_1(d, a)$ by (*) and $\beta_0(d, v) \neq \beta_1(d, v)$ by the definition of D. If $d \in N(v)$ then the Naji equations require

$$\beta_i(v, d) + \beta_i(v, a) + \beta_i(a, d) + \beta_i(d, a) = 1$$

for $i \in \{0, 1\}$, an impossibility as $\beta_0(v, d) \neq \beta_1(v, d)$ and the other terms of the two equations are all equal.

For the second assertion, observe that claim 2 tells us that $C = \emptyset$ and claim 5 tells us that no $b \in B$ is a neighbor of a.

Claim 7. If $c \in C$ then c is adjacent to every element of $D \cap N(v)$, and the other neighbors of c all lie in $(A \cup C) - N(v)$.

proof: If x is a neighbor of c then claim 2 implies that $x \notin A \cap N(v)$, and claim 4 implies that $x \notin (B \cup D) - N(v)$. Hence $x \in ((A \cup C) - N(v)) \cup (D \cap N(v))$. If $d \in D \cap N(v)$ and $cd \notin E(G)$ then the Naji equations require $\beta_i(c, d) = \beta_i(c, v)$ for $i \in \{0, 1\}$; but this is impossible as $\beta_0(c, d) = \beta_1(c, d)$ by (*) and $\beta_0(c, v) \neq \beta_1(c, v)$ by the definition of C.

Claim 8. If $a \in A - N(v)$ then the neighbors of a all lie in $A \cup C$.

proof: Claim 4 implies that the neighbors of a all lie in $A \cup C \cup (D \cap N(v))$, so it suffices to verify that no neighbor of a lies in $D \cap N(v)$. Suppose instead that $d \in D \cap N(v)$ is a neighbor of a. The Naji equations require

$$\beta_i(v,d) + \beta_i(a,d) + \beta_i(a,v) + \beta_i(v,a) = 1$$

for $i \in \{0,1\}$. This is impossible as $\beta_0(v,d) \neq \beta_1(v,d)$ and the other terms of the two equations are all equal.

Claims 2 and 3 yield four cases.

Case 1. $C \neq \emptyset$ and $A \cap N(v) = \emptyset = D \cap N(v)$. In this case $N(v) = \emptyset$, an impossibility as a prime graph cannot have an isolated vertex.

Case 2. $C \neq \emptyset$ and $A \cap N(v) = \emptyset = D - N(v)$. In this case D = N(v), the elements of C are all adjacent to all the elements of D, the other neighbors of elements of C all lie in $A \cup C$, the neighbors of elements of A all lie in $A \cup C$, and the neighbors of elements of B all lie in $B \cup D$. As $(A \cup C, B \cup D \cup \{v\})$ cannot be a split of G, either $|A \cup C| \leq 1$ or $B \cup D = \emptyset$. The latter is impossible as it would leave v isolated. As $C \neq \emptyset$, $|A \cup C| \leq 1$ implies $A = \emptyset$ and |C| = 1. Hence the lone $c \in C$ has N(c) = D = N(v); but this is impossible as $(B \cup D, \{c, v\})$ would be a split of G.

Case 3. $C = \emptyset$ and $D - N(v) = \emptyset$. In this case all elements of $A \cap N(v)$ are adjacent to all elements of $D \cap N(v)$, the neighbors of elements of B all lie in $B \cup D$, and the neighbors of elements of A - N(v) all lie in A. As neither $(A \cup \{v\}, B \cup D)$ nor $(A, B \cup D \cup \{v\})$ is a split of G, either $A = \emptyset$ or $B \cup D = \emptyset$. If $A = \emptyset = C$ then D = N(v) and B = V(G - v) - N(v); hence $\beta_1 = \beta_0 + \delta(v)$. If $B \cup D = \emptyset = C$ then $\beta_1 = \beta_0$.

Case 4. $C = \emptyset$ and $D \cap N(v) = \emptyset$. In this case $N(v) \subseteq A$, the neighbors of elements of A - N(v) all lie in A, every element of N(v) is adjacent to every element of D, and the neighbors of elements of B all lie in $B \cup D$. As $(A \cup \{v\}, B \cup D)$ cannot be a split of G, it follows that |A| = 0 or $|B \cup D| \leq 1$. If |A| = 0 then v is isolated, an impossibility in a prime graph. If |B| = 1 then the lone $b \in B$ is isolated, another impossibility. If $B = C = \emptyset \neq D$ then there is a lone $d \in D$, with N(d) = N(v). But that too is impossible, as $(A, \{d, v\})$ would be a split of G. Consequently $B = C = D = \emptyset$, so $\beta_1 = \beta_0$.

A corollary of Theorem 19 describes the relationship between the Naji solutions of G and those of G - v, in case both graphs are prime.

Corollary 20 Let G be a prime Naji graph with $|V(G)| \ge 6$, and suppose G-v is also prime. Then restriction defines a 2-to-1 surjection

{*Naji solutions of G*} \rightarrow {*Naji solutions of G* - v}.

Proof. If β is any Naji solution of G then certainly the restriction $\beta|(G-v)$ is a Naji solution of G-v. As the ρ and $\delta(x)$ vectors of G restrict to those of G-v (with the exception that $\delta(v)$ restricts to 0) Theorem 19 guarantees that restriction defines a surjection. To verify that the surjection is 2-to-1, i.e., every Naji solution of G-v corresponds to precisely two Naji solutions of G, note first that every Naji solution of G-v corresponds to at least two different Naji solutions of G; there must be one, as restriction is surjective, and then there is another obtained by adding $\delta(v)$. Then note that Proposition 7 implies that there are twice as many Naji solutions for G as there are for G-v.

Another corollary is the following result of Bouchet [2].

Corollary 21 [2] Let G be a prime circle graph. Then there is only one double occurrence word W with $G = \mathcal{I}(W)$, up to cyclic permutation and reversal.

Proof. Let W and W' be double occurrence words with $\mathcal{I}(W) = \mathcal{I}(W') = G$. Theorem 19 tells us that W and W' have orientations whose corresponding Naji solutions are the same. Choose any $v \in V(G)$, and cyclically permute W and W'so they both begin with v^{out} . As v is not isolated in G, the second letter in W is some $w \neq v$. Adding $\delta(w)$ to both Naji solutions if necessary, we may presume that this second letter of W is w^{out} . Then $\beta(x, v) = \beta(x, w) \ \forall x \notin \{v, w\}$; as Wand W' provide the same Naji solution, Proposition 9 guarantees that v^{out} and w^{out} are consecutive in W' too. It is possible that w^{out} is the last letter in W', rather than the second; if so, reverse and cyclically permute W' so that both W and W' are in the form $v^{out}w^{out}$... Repeat this process to verify that W and W' must have the same third letter, then the same fourth letter, and so on.

Before proceeding we should mention that the appearance of Corollary 21 here is no coincidence. Our proof of Naji's theorem follows the outline of the argument given by Bouchet in justifying his circle graph recognition algorithm [2]. As we are about to see, though, the second part is considerably more difficult for us. The second part of Bouchet's algorithm used simple "brute force" (as he described it on p.253 of [2]) to check all possible double occurrence words for a prime graph G, knowing that the essentially unique double occurrence word for a prime G - v. Our job is more difficult, as we must prove that a prime Naji graph arises from a double occurrence word.

6 Step 2 of the proof: building a word

In this section we complete the proof of Naji's theorem. We begin with a technical observation.

Lemma 22 Let G be a Naji graph with an edge e = vw. Suppose G and G - eshare a Naji solution β . Then $\beta(x, v) \neq \beta(x, w) \ \forall x \in (N(v)\Delta N(w)) - \{v, w\}$, and $\beta(x, v) = \beta(x, w) \ \forall x \notin N(v)\Delta N(w)$. (Here Δ denotes the symmetric difference.)

Proof. Suppose $x \in (N(v)\Delta N(w)) - \{v, w\}$; renaming v and w if necessary, we may presume that $vx \in E(G)$ and $wx \notin E(G)$. Then the Naji equations for G require

$$\beta(x, v) + \beta(w, v) + \beta(x, w) + \beta(w, x) = 1$$

while the Naji equations for G-e require $\beta(w, v) = \beta(w, x)$. If $x \notin N(v) \cup N(w)$ then the Naji equations for G require $\beta(x, v) = \beta(x, w)$. If $x \in N(v) \cap N(w)$ then the Naji equations for G-e require

$$\beta(x,v) + \beta(x,w) + \beta(v,w) + \beta(w,v) = 1$$

while the Naji equations for G require $\beta(v, w) + \beta(w, v) = 1$.

Corollary 23 Let G be a Naji graph with an edge e = vw. Suppose G and G - e share a Naji solution β . Then G and G - e have Naji solutions β_1 and β_2 (respectively) such that (a) the only difference between β_1 and β_2 is that $\beta_1(v, w) = \beta_2(w, v)$ and $\beta_1(w, v) = \beta_2(v, w)$ and (b) $\beta_1(x, v) = \beta_1(x, w)$ $\forall x \notin \{v, w\}$.

Proof. Consider the Naji solutions $\beta_1 = \beta + \delta_G(v) + \delta_G(w)$ and $\beta_2 = \beta + \delta_{G-e}(v) + \delta_{G-e}(w)$.

Theorem 24 Let G be a prime Naji graph with $|V(G)| \ge 5$. Then there is a double occurrence word W with $\mathcal{I}(W) = G$.

Proof. Again, we begin with $C_5 = \mathcal{I}(bacbdcedae)$. Every other prime graph of order 5 is locally equivalent to C_5 [2], and hence is also a circle graph.

If |V(G)| = 6 then Proposition 17 tells us that after replacing G with a locally equivalent graph, we may presume that G has a vertex v such that G-v is prime. Then G-v is locally equivalent to C_5 , so after further local complementation of G (if necessary) we may presume that $G-v = C_5 = \mathcal{I}(bacbdcedae)$. Every proper subset of $\{a, b, c, d, e\}$ can be achieved as an interlacement neighborhood of v in a double occurrence word obtained by inserting two appearances of v into bacbdcedae: for instance vbavcbdcedae, bvacvbdcedae, vbacvbdcedae, bvacvbdcedae, and bvacbdvcedae provide v with the interlacement neighborhoods $\{a, b\}, \{a, c\}, \{a, b, c\}, \{a, b, d\}, \text{ and } \{a, b, c, d\}$ respectively. The interlacement neighborhood $\{a, b, c, d, e\}$ cannot be achieved in this way; we leave it to the reader to verify that the resulting graph, the wheel graph W_5 , is not a Naji graph.

We proceed using induction on |V(G)| > 6. Observe that Proposition 11 tells us that our inductive hypothesis is that *all* Naji graphs smaller than G are circle graphs (not just the prime ones). Proposition 17 tells us that after local complementation, we may presume that G has a vertex v such that G - v is

prime, and similarly some graph H locally equivalent to G - v has a vertex w such that H - w is prime. By applying the local complementations needed to obtain H from G - v to G before deleting v, we may presume simply that G - v and G - v - w are both prime.

We distinguish three cases.

Case 1. G - w is prime.

With Corollary 21, the inductive hypothesis guarantees that up to cyclic permutation and reversal, there is a unique double occurrence word W with $\mathcal{I}(W) = G - v - w$, and there are unique locations to insert two appearances of v and two appearances of w into W so as to obtain double occurrence words whose interlacement graphs are G - w and G - v. Using cyclic permutations, we may presume that if we insert both v and w into W then v appears first, and if v and w are not interlaced then the second appearance of v precedes the first appearance of w. That is, W = ABCD and either W' = vAvBwCwD or W'' = vAwBvCwD correctly describes both G - v and G - w through interlacement. If W' or W'' correctly describes G, we are done.

Otherwise, either W' or W'' correctly describes both G - v and G - w, but does not succeed in describing G; the only way this can happen is that interlacement misrepresents the adjacency of v and w.

Suppose W' correctly describes G - v and G - w, but does not succeed in describing G. That is, if e = vw then $e \in E(G)$ and $\mathcal{I}(W') = G - e$. If B or D is empty then v and w appear consecutively in W'; we may simply interchange their consecutive appearances to obtain a double occurrence word whose interlacement graph is G. A and C cannot be empty as neither v nor wcan be isolated, so we proceed with the assumption that A, B, C and D are all nonempty. We aim for a contradiction.

Let β be any Naji solution for G. Theorem 19 tells us that there is an orientation of W corresponding to the Naji solution $\beta|(G-v-w)$ of G-v-w, and extensions of this orientation to G-v and G-w (i.e., in/out designations of the appearances of v and w in W') such that the resulting orientations of double occurrence words correspond to the Naji solutions $\beta|(G-v)$ and $\beta|(G-w)$ of G-v and G-w. The only possible differences between β and the Naji solution β' of $\mathcal{I}(W')$ corresponding to the resulting orientation of W' involve the values $\beta(v, w)$ and $\beta(w, v)$.

We claim that it is possible to choose β so that $\beta = \beta'$. To verify the claim, suppose we have a β with $\beta(v,w) \neq \beta'(v,w)$ and $\beta(w,v) = \beta'(w,v)$. Consider $\hat{\beta} = \beta + \delta_G(w)$. As $\beta|(G-w) = \hat{\beta}|(G-w)$, β and $\hat{\beta}$ result in the same orientation of W, and the same orientation of the word obtained from W by inserting v. The restriction of $\delta_G(w)$ to G-v is $\delta_{G-v}(w)$, of course, and the effect of adding $\delta_{G-v}(w)$ on the corresponding orientation is to interchange the appearances of w^{in} and w^{out} . Notice that interchanging the appearances of w^{in} and w^{out} in W' = vAvBwCwD has the effect that $(\hat{\beta})'(v,w) = \beta'(v,w)$ and $(\hat{\beta})'(w,v) \neq \beta'(w,v)$. As $\hat{\beta}(v,w) \neq \beta(v,w)$ and $\hat{\beta}(w,v) \neq \beta(w,v)$, it follows that $\hat{\beta}(v,w) = (\hat{\beta})'(v,w)$ and $\hat{\beta}(w,v) = (\hat{\beta})'(w,w)$ and $\hat{\beta}(w,v) = \beta(v,w)$ and $\beta(w,v) \neq \beta'(w,v)$ then $\hat{\beta} = \beta + \delta_G(v)$ will have $\hat{\beta}(v,w) = (\hat{\beta})'(v,w)$ and $\hat{\beta}(w,v) = (\hat{\beta})'(w,v);$ and if $\beta(v,w) \neq \beta'(v,w)$ and $\beta(w,v) \neq \beta'(w,v)$ then $\hat{\beta} = \beta + \delta_G(v) + \delta_G(w)$ will have $\hat{\beta}(v,w) = (\hat{\beta})'(v,w)$ and $\hat{\beta}(w,v) = (\hat{\beta})'(w,v).$

Having verified the claim, we now know that G and G - e share a Naji solution, with e the edge vw. Corollary 23 tells us that consequently, $G - e = \mathcal{I}(W')$ has a Naji solution β_2 with $\beta_2(x,v) = \beta_2(x,w) \ \forall x \notin \{v,w\}$. As G is prime, it has no cutpoint. Consequently G - e is connected and we may cite Proposition 9 to conclude that v and w appear consecutively in W. But this contradicts the assumption that B and D are both nonempty.

Suppose now that W'' = vAwBvCwD correctly describes G - v and G - w, but does not succeed in describing G. That is, if e = vw then $e \in E(\mathcal{I}(W''))$ and $G = \mathcal{I}(W'') - e$. If any one of A, B, C, D is empty then v and w appear consecutively in W''; we may interchange consecutive appearances of v and w to obtain a double occurrence word whose interlacement graph is G. We proceed with the assumption that A, B, C and D are all nonempty, and derive a contradiction.

As before, any Naji solution β for G leads to an orientation of W'' with the property that the corresponding Naji solution β'' of $\mathcal{I}(W'')$ can only differ from β in the values $\beta(v, w)$ and $\beta(w, v)$. We claim again that it is possible to choose β so that $\beta'' = \beta$. If we have a β such that $\beta(v, w) \neq \beta''(v, w)$ and $\beta(w, v) = \beta''(w, v)$ then again, we consider $\hat{\beta} = \beta + \delta_G(w)$. And again, β and $\hat{\beta}$ result in the same orientation of the word obtained from W by inserting v. But now when we interchange w^{in} and w^{out} in W'' = vAwBvCwD we have $(\hat{\beta})''(v, w) \neq \beta''(v, w)$ and $(\hat{\beta})''(w, v) \neq \beta''(w, v)$. As $\hat{\beta}(v, w) = \beta(v, w)$ and $\hat{\beta}(w, v) \neq \beta(w, v)$, though, it follows again that $\hat{\beta}(v, w) = (\hat{\beta})''(v, w)$ and $\hat{\beta}(w, v) = (\hat{\beta})''(w, v)$. As before, the possibility that $\beta(w, v) \neq \beta''(w, v)$ is handled by using $\delta_G(v)$. The claim allows us to use Corollary 23 and Proposition 9 to derive a contradiction.

Case 2. G - w is not connected.

This case cannot occur as the prime graph G cannot have a cutpoint.

Case 3. G - w is connected and not prime. (This is the most delicate case.) Let (X, Y) be a split of G - w with $v \in X$. If |X| > 2 then (X - v, Y) is a split of G - w - v, an impossibility as G - w - v is prime. Consequently |X| = 2; let $X = \{v, x\}$. If v and x are not adjacent we may pick any $y \in N(v) = N(x)$, and replace G with the local complement G^y . If v and x are adjacent and $N(v) \cap Y \neq \emptyset$ we may replace G with the local complement G^x . After these replacements we see that we may assume that the degree of v in G - w is 1. As G is prime, the degree of v in G cannot be 1; hence $vw \in E(G)$, and $N(v) = \{w, x\}$.

Replacing G with G^v if necessary, we may presume that $wx \in E(G)$. According to Corollary 6, G has a Naji solution β with $\beta(w, v) = \beta(x, v) = 0$ and $\beta(u, v) = 1 \quad \forall u \notin \{v, w, x\}$. We have no further need for the hypothesis that G - v - w is prime, so it does no harm to assume that $\beta(w, x) = 1$; if $\beta(w, x) = 0$, we simply interchange the names of w and x. Our job, then, is to produce a double occurrence word W with $\mathcal{I}(W) = G$, under the assumptions that G is a prime Naji graph, G - v is a prime circle graph, $N(v) = \{w, x\}$, $\beta(w,v) = \beta(x,v) = 0, \ \beta(w,x) = 1 \ \text{and} \ \beta(u,v) = 1 \ \forall u \notin \{v,w,x\}.$

According to Corollary 21, up to cyclic permutation and reversal there is a unique double occurrence word W whose interlacement graph is G-v. Theorem 19 tells us that W can be oriented so that the corresponding Naji solution is the restriction $\beta|(G-v)$. Cyclically permute W so the first letter is w^{in} . As w and x are interlaced and $\beta(w, x) = 1$,

$$W = w^{in}Ax^{in}Bw^{out}Cx^{out}D$$

for some subwords A, B, C and D. We will use pairs of letters to designate subsets of V(G) in the natural way: AB denotes the set of vertices that appear once in A and once in B, CC denotes the set of vertices that appear twice and C and so on. Our aim is to prove that there must be locations in W where we can place two appearances of v so that v is interlaced with w and x, but not with any other vertex. If any of A, B, C, D is empty then w and x appear consecutively in W and we can accomplish our aim by replacing a subword wxor xw with vwxv or vxwv. Consequently we may proceed with the assumption that none of A, B, C, D is empty.

We begin with a sequence of claims. During the discussion of the claims we will often use the fact that if $y, z \in V(G) - \{v, w, x\}$ the Naji equations require $\beta(v, y) = \beta(v, z)$ if yz is an edge, or there is a path from y to z in G - v - w - x. Claim 1. $AC = \emptyset$.

proof: Suppose $a \in AC$. If a^{in} appears in A then $\beta(a, w) = 0$ and $\beta(a, x) = 1$; if a^{in} appears in C these values are reversed. Either way, $\beta(a, w) \neq \beta(a, x)$. As $av \notin E(G)$ and $aw, ax, vw, vx \in E(G)$, the Naji equations of G require

$$\beta(a, w) + \beta(v, w) + \beta(a, v) + \beta(v, a) = 1 \text{ and}$$

$$\beta(a, x) + \beta(v, x) + \beta(a, v) + \beta(v, a) = 1.$$

It cannot be that both equations hold as the first terms are unequal and the other terms are all equal. \blacksquare

Claim 2. If $y \in AA \cup AB \cup BB \cup BC \cup CC \cup CD \cup DD$ then y^{in} precedes y^{out} , and if $y \in AD$ then y^{out} precedes y^{in} .

proof: If $y \in AA \cup AB \cup BB \cup CC \cup CD \cup DD$ then y is not interlaced with w, so the Naji equations require $\beta(y, w) = \beta(y, v) = 1$; this in turn requires that y^{in} precedes y^{out} . If $y \in BC$ then y is not interlaced with x, so $\beta(y, x) = \beta(y, v) = 1$ and again this requires that y^{in} precede y^{out} . If $y \in AD$, instead, then $\beta(y, x) = \beta(y, v) = 1$ implies that y^{in} appears in D.

Claim 3. If $a \in AA \cup AB \cup AD$ then $\beta(v, a) = 0$.

proof: If $a \in AB$ then claim 2 implies that $\beta(a, x) = 1$. As $a \in N(x) - N(v)$, the Naji equations require

$$1 = \beta(a, x) + \beta(v, x) + \beta(a, v) + \beta(v, a) = 1 + 1 + 1 + \beta(v, a).$$

Similarly, if $a \in AD$ then $\beta(a, w) = 1$ and $a \in N(w) - N(v)$, so the Naji equations require

$$1 = \beta(a, w) + \beta(v, w) + \beta(a, v) + \beta(v, a) = 1 + 1 + 1 + \beta(v, a).$$

If $a \in AA$ then as G - v is connected, some path in G - v must lead from a to some vertex not in AA. As $AC = \emptyset$, the definition of interlacement makes it clear that the first vertex on this path that is not in AA is in AB or AD; as $\beta(v, y) = 0$ for every such vertex y, and no vertex on the path neighbors v, the Naji equations require that $\beta(v, a) = 0$.

Claim 4. If $c \in BC \cup CC \cup CD$ then $\beta(v, c) = 1$.

proof: The proof is closely analogous to that of claim 3. \blacksquare

Claim 5. In the B portion of W, all vertices from AB precede all vertices from BC.

proof: If $a \in AB$ and $c \in BC$ were interlaced, a Naji equation would require that $\beta(v, a) = \beta(v, c)$; but this would contradict claims 3 and 4.

Claim 6. In the D portion of W, all vertices from CD precede all vertices from AD.

proof: If $c \in CD$ and $a \in AD$ were interlaced, a Naji equation would require that $\beta(v, a) = \beta(v, c)$; but this would contradict claims 3 and 4.

Observe that claim 5 tells us we can partition the *B* portion of *W* as $B_0B_1B_2$ in such a way that all vertices from *AB* appear in B_0 , the last letter in B_0 is a vertex from *AB*, all vertices from *BC* appear in B_2 and the first letter in B_2 is a vertex from *BC*. Claim 6 tells us that we can partition the *D* portion of *W* as $D_0D_1D_2$ in such a way that all vertices from *CD* appear in D_0 , the last letter in D_0 is a vertex from *CD*, all vertices from *AD* appear in D_2 and the first letter in D_2 is a vertex from *AD*. (Some of the subwords B_i, D_i may be empty.) Consequently we have

$$W = w^{in} A x^{in} B_0 B_1 B_2 w^{out} C x^{out} D_0 D_1 D_2.$$

Claim 7. If y appears in B_0 or D_2 then $\beta(v, y) = 0$, and if y appears in B_2 or D_0 then $\beta(v, y) = 1$.

proof: Consider a vertex y that appears in B_0 . If $y \in AB$ then $\beta(v, y) = 0$ by claim 3. If $y \in BD$ then y is interlaced with the vertex $a \in AB$ that appears at the end of B_0 , so $\beta(v, y) = \beta(v, a) = 0$. The same argument applies if $y \in BB$ appears only once in B_0 . If $y \in BB$ appears twice in B_0 , then as G is connected, some path must lead from y to a vertex z that appears only once in B_0 . Consider such a path of shortest length; then all the vertices on the path before z are, like y, elements of BB that appear twice in B_0 . Then $\beta(v, z) = 0$ by the earlier parts of the argument, and the Naji equations require that $\beta(v, y) = \beta(v, z)$. Similar arguments apply in B_2 , D_0 and D_2 .

Notice that claim 7 implies that vertices from B_0 or D_2 cannot appear in B_2 or D_0 , and vice versa. Consequently if B_1 and D_1 are both empty then the word

$$w^{in}Ax^{in}B_0vB_2w^{out}Cx^{out}D_0vD_2$$

has G as its interlacement graph. Our aim is to show that if B_1 and D_1 are not empty, they have "centers" where we can insert the desired appearances of v.

To locate these centers we repartition B and D. Let $B = B^1 B^2 B^3 \dots B^k$ in such a way that each B^i is nonempty and the value of $\beta(v, -)$ is constant on each B^i , with $\beta(v, -)$ changing when we pass from B^i to B^{i+1} . The fact that *B* is nonempty tells us that $k \ge 1$. Claim 7 tells us that B^1 contains B_0 and B^k contains B_2 . (N.b. B_0 or B_2 might be empty.) Partition D as $D^1...D^{\ell}$ in a similar way.

Claim 8. For each i, there is a vertex that appears precisely once in B^i . Similarly, in each D^j some vertex appears exactly once.

proof: If every vertex that appears in B^i appears twice in B^i , then no vertex that appears in B^i is interlaced with any vertex that does not appear in B^i . This is impossible, as B^i is not empty and G is connected. The same observation applies to D^j .

Claim 9. If $i \neq j$ then no vertex appears in both B^i and B^j , and no vertex appears in both D^i and D^j .

proof: Suppose i < j, a vertex y appears in both B^i and B^j , and j-i is as small as possible. Claim 8 tells us that there is a vertex z, which appears precisely once in B^{i+1} . The minimality of j-i guarantees that the other appearance of z is outside the subword $B^i B^{i+1} \dots B^j$, so y and z are interlaced. The Naji equations then require $\beta(v, y) = \beta(v, z)$, contradicting the definition of $B^1 B^2 B^3 \dots B^k$, which guarantees $\beta(v, y) \neq \beta(v, z)$. The same argument applies to D^i and D^j .

Claim 10. Suppose y appears precisely once in B^i . If i = 1 then $y \in AB \cup BD$, if 1 < i < k then $y \in BD$, and if i = k then $y \in BC \cup BD$. Similarly, if z appears precisely once in D^j then j = 1 implies $z \in BD \cup CD$, $1 < j < \ell$ implies $z \in BD$, and $j = \ell$ implies $z \in AD \cup BD$.

proof: Claim 9 tells us that $y \notin BB$, as every element of BB appears twice in the same one of $B^1, ..., B^k$. The assertion regarding B follows because all appearances in B of elements of AB occur in B_0 , which is a subword of B^1 ; and all appearances in B of elements of BC occur in B_2 , which is a subword of B^k . The assertion regarding D is verified in the same way.

Claim 11. Neither k > 2 nor $\ell > 2$ is possible.

proof: Suppose k > 2. Let Y^1 , Y^2 , $Y^3 \subseteq V(G)$ be the subsets consisting of vertices that appear precisely once in B^1 , B^2 and B^3 respectively. Claims 8 and 9 tell us that Y^1 , Y^2 , and Y^3 are nonempty and pairwise disjoint, and claim 10 guarantees that $Y^2 \subseteq BD$.

Subclaim 11a. All the second appearances of elements of Y^2 appear in the same D^j .

proof: If $y, y' \in Y^2$ appear in D^i and D^j with i < j then consider a vertex z that appears once in D^{i+1} ; we have $\beta(v, z) \neq \beta(v, y) = \beta(v, y')$. Claim 10 tells us that $z \in BD$, so the other appearance of z is in B; if z appears in B^1 it is interlaced with y', and if z does not appear in B^1 then it is interlaced with y. Either way we have a contradiction as the Naji equations require that interlaced elements of BD have the same $\beta(v, -)$ value.

We now let τ denote the index of the particular D^j that includes all the second appearances of elements of Y^2 . No element of Y^2 can be interlaced with an element of $Y^1 \cup Y^3$, as the value of $\beta(v, -)$ on Y^2 is different from the value on $Y^1 \cup Y^3$. Consequently if $y \in Y^1$ then the other appearance of y must either occur in D after D^{τ} (if $y \in BD$) or in A (if $y \in AD$). Also, if $y \in Y^3$ then the other appearance of y must occur before D^{τ} , either in D (if $y \in BD$) or in C (if $y \in BC$).

Subclaim 11b. Every vertex that appears precisely once in D^{τ} also appears in Y^2 .

proof: Suppose d appears once in D^{τ} and the other appearance of d is not in Y^2 . Claim 9 tells us that the other appearance of d is not in D. If the other appearance of d is in A, then $d \in AD$ so $\beta(v, d) = 0$ by claim 3. Then $\beta(v,y) = 0 \ \forall y \in Y^2$ and $\beta(v,y') = 1 \ \forall y' \in Y^1$; consequently no vertex that appears in Y^1 is an element of AD, so every vertex y' that appears in Y^1 is an element of BD. According to the paragraph before the statement of this subclaim, every y' that appears in Y¹ appears in D after D^{τ} ; it follows that d is interlaced with every such y'. But this is impossible because the $\beta(v, -)$ values do not match. A similar line of reasoning applies if the other appearance of d is in C: $\beta(v,d) = 1$ by claim 4, so $\beta(v,y) = 1 \ \forall y \in Y^2$ and $\beta(v,y') = 0$ $\forall y' \in Y^3$; hence no element of Y^3 lies in BC, so every element of Y^3 lies in BD. It follows that every element of Y^3 appears in D before D^{τ} , by the paragraph before the statement of this subclaim; but then every such element is interlaced with d, and again the $\beta(v, -)$ values prohibit this. Consequently d must appear in B. But then d appears after Y^3 in B and also after Y^3 in D, an impossibility because d cannot be interlaced with any element of Y^3 .

Now consider the subwords Y^2 and D^{τ} of W. Subclaims 11a and 11b tell us that if X is the set of vertices of G - v that appear outside Y^2D^{τ} and Y is the set of vertices that appear within Y^2D^{τ} , then $X \cap Y = \emptyset$. G - v is prime, so (X, Y) cannot be a split. As X contains w and x along with all the vertices that appear in Y^1 and Y^3 , |X| > 2; hence |Y| = 1. That is, Y^2 and D^{τ} are both of length 1, and mention the same vertex.

Subclaim 11c. No vertex of G - v - w - x is interlaced with the vertex y that appears in Y^2 and D^{τ} .

proof: Suppose z is interlaced with y. Then $z \notin AA \cup CC$. If $z \in BB \cup DD$ then claim 9 tells us that z appears twice in Y^2 or D^{τ} , an impossibility as no vertex other than y appears in either Y^2 or D^{τ} . If $z \in AB$ then $z \in B^1$, so $\beta(v,z) \neq \beta(v,y)$; this is not possible if $yz \in E(G)$. If $z \in AD$ then z appears in A before the appearance of y in B, so z also appears before y in D; as the vertices of AD all appear in D^{ℓ} it follows that $\tau = \ell$. But then both y and z appear in D^{τ} , contradicting the fact that only y appears in D^{τ} . If $z \in BC$ then the appearance of z in C precedes the appearance of y in D, so the appearance of z in B must precede the appearance of y in B; but then z appears in B^1 so $\beta(v,z) \neq \beta(v,y)$, an impossibility if y and z are neighbors in G. If $z \in BD$ then $\beta(v, z) = \beta(v, y)$, so z appears in some B^{2i} with i > 1. Consider a vertex b that appears once in B^3 . To avoid being interlaced with z, b must appear after z in D. To avoid being interlaced with y, b must appear before y in D. Hence z appears before y in D; but this cannot be the case as z appears after y in B and $yz \in E(G)$. The only remaining possibility is $z \in CD$. Such a z would have to appear in or after $D^{\tau+2}$, in order to be interlaced with y. Suppose b appears once in B_3 ; then b must appear in D before D^{τ} , to avoid being interlaced with y. Consequently b and z are interlaced, an impossibility

as $\beta(v, z) = \beta(v, y) \neq \beta(v, b)$.

Subclaim 11c completes the proof that k > 2 is impossible, for it implies that $N(y) = \{w, x\}$; this in turn implies that $\{v, y\}$ is a split of G.

The assertion that $\ell>2$ is impossible can be proven in the same way, so we are done with claim 11. \blacksquare

Claim 12. At least one of AB, AD is not empty, and at least one of BC, CD is not empty. Consequently, $k + \ell \geq 2$.

proof: If $AB = AD = \emptyset$ then as A is not empty, it must be that $AA \neq \emptyset$. But no edge of G can connect a vertex of AA to a vertex outside AA, contradicting the fact that G is connected. Hence at least one of AB, AD is not empty. If $AB \neq \emptyset$ then $k \geq 1$, as the vertices of AB all appear in B^1 ; and if $AD \neq \emptyset$ then $\ell \geq 1$, as the vertices of AD all appear in D^{ℓ} . Similarly, at least one of BC, CD is not empty, so at least one of B^k, D^1 is not empty. Finally, no vertex of $AB \cup AD$ can appear in the same set B^i or D^j as a vertex of $BC \cup CD$, because the $\beta(v, -)$ values do not match.

At this point we change notation slightly. If k = 2 then we let $B(i) = B^i$ for $i \in \{1, 2\}$. If k = 1 and the value of $\beta(v, -)$ on B^1 is 0, we let $B(1) = B^1$, and we let B(2) denote the empty word. If k = 1 and the value of $\beta(v, -)$ on B^1 is 1, we let $B(2) = B^1$ and we let B(1) denote the empty word. Similarly, we define $D(1) = D^1$ and $D(2) = D^2$ if $\ell = 2$, and if $\ell = 1$ we define D(1) and D(2) so that one is empty, the other is D^1 and the value of $\beta(v, -)$ on D(i) is $i \pmod{2}$. We now have

$$W = w^{in} A x^{in} B(1) B(2) w^{out} C x^{out} D(1) D(2),$$

where up to two of B(1), B(2), D(1), D(2) may be empty.

Claim 13. The value of $\beta(v, -)$ is 0 on B(1) and D(2), and 1 on B(2) and D(1).

proof: If $AB \neq \emptyset$ then the claim is true for B(1), as every vertex $a \in AB$ appears in B(1) and has $\beta(v, a) = 0$. Necessarily then the claim is also true for B(2), as the values of of $\beta(v, -)$ on B(1) and B(2) are different. Similarly, if $AD \neq \emptyset$ then the claim is true for D(1) and D(2), as every $a \in AD$ appears in D(2) and has $\beta(v, a) = 0$. The same reasoning shows that the claim holds in B if $BC \neq \emptyset$, and the claim holds in D if $CD \neq \emptyset$.

Claim 12 now assures us that claim 13 holds in at least one of B and D; suppose it holds in B. If the claim does not hold in D then $D(1) = D^1$ and $D(2) = D^2$ are both nonempty, the value of $\beta(v, -)$ on D(1) is 0, and the value of $\beta(v, -)$ on D(2) is 1. Let d_1 and d_2 be vertices that appear precisely once in D(1) and D(2), respectively. Then $d_1, d_2 \notin DD$, of course, and also the values of $\beta(v, d_1)$ and $\beta(v, d_2)$ indicate that $d_1 \notin CD$ and $d_2 \notin AD$. Consequently $d_1 \in AD \cup BD$ and $d_2 \in BD \cup CD$. As $\beta(v, d_1) \neq \beta(v, d_2)$, d_1 and d_2 are not interlaced; d_1 precedes d_2 in D, so d_2 must precede d_1 outside D. This is impossible if $d_1 \in AD$ or $d_2 \in CD$, so it must be that $d_1, d_2 \in BD$. But then the values of $\beta(v, d_i)$ indicate that d_1 appears in B(1) and d_2 appears in B(2), so d_2 does not precede d_1 outside D. By contradiction, we conclude that if claim 13 holds in B it also holds in D. The converse is justified in the same way. Claim 14. G is the interlacement graph of the double occurrence word

$$W' = w^{in} A x^{in} B(1) v B(2) w^{out} C x^{out} D(1) v D(2)$$

proof: A vertex that appears in A has $\beta(v, a) = 0$, by claims 1 and 3, so it may appear twice in A, or once in A and once in B(1), or once in A and once in D(2). In any case it is not interlaced with v in W'. Similarly a vertex that appears in C may appear again in C, or appear in B(2) or D(1); in any case it is not interlaced with v. A vertex b that appears in B and D must appear either in B(1) and D(2) (if the value of $\beta(v, b)$ is 0) or in B(2) and D(1) (if the value of $\beta(v, b)$ is 1); again, neither case allows it to be interlaced with v. Finally, an element of BB or DD is not interlaced with v. Consequently w and x are the only vertices interlaced with v in W'.

7 Bipartite graphs

Bipartite circle graphs are special for two reasons, both connected with planarity. One special property is geometric: bipartite circle graphs correspond to planar 4-regular graphs [18]. (All circle graphs correspond to 4-regular graphs, as a double occurrence word naturally gives rise to an Euler circuit in a 4-regular graph.) Another special property is matroidal: a bipartite graph with adjacency matrix A is a circle graph if and only if the binary matroid represented by $\begin{pmatrix} I & A \end{pmatrix}$ is planar [10]; here I is an identity matrix. (This matroid is the direct sum of a pair of mutually dual matroids, so it is planar if and only if it is either graphic or cographic.)

At the end of [13], Geelen and Gerards deduce an algebraic characterization of planar matroids from their characterization of graphic matroids. The following theorem provides a bridge between their result and Naji's theorem.

Theorem 25 Let G be a bipartite graph with vertex classes V_1 and V_2 . Then G is a circle graph if and only if this system of equations has a solution over GF(2).

(a) If v, w, x are three different elements of the same vertex class and $N(v) \cap N(w) \not\subseteq N(x)$, then $\beta(x, v) = \beta(x, w)$.

(b) If v, w, x are three different elements of the same vertex class and $N(v) \cap N(w) \cap N(x) \neq \emptyset$, then

$$\beta(v,w) + \beta(w,v) + \beta(v,x) + \beta(x,v) + \beta(w,x) + \beta(x,w) = 1$$

Proof. The equations mentioned in the statement follow directly from the Naji equations. For (a), note that if $y \in N(v) \cap N(w) - N(x)$, then the Naji equations require $\beta(x, v) = \beta(x, y)$ and $\beta(x, y) = \beta(x, w)$. For (b), note that if $y \in N(v) \cap N(w) \cap N(x)$ we may add together the following Naji equations.

$$\begin{split} \beta(y,v) + \beta(y,w) + \beta(v,w) + \beta(w,v) &= 1\\ \beta(y,v) + \beta(y,x) + \beta(v,x) + \beta(x,v) &= 1\\ \beta(y,w) + \beta(y,x) + \beta(w,x) + \beta(x,w) &= 1 \end{split}$$

For the converse, suppose the equations mentioned in the statement of this theorem have a solution β . Note that the equations require only that $\beta(v, w)$ be defined when v and w are elements of the same vertex class. In order to build a Naji solution we must define values of $\beta(v, w)$ when v and w are not elements of the same vertex class. We may presume that G is connected, as Naji solutions extend directly from the components of a disconnected graph to a Naji solution for the whole graph.

Suppose $v \in V_1$, $w \in V_2$ and $vw \notin E(G)$. As G is connected, there are $v' \in V_1$ and $w' \in V_2$ such that $v'w, vw' \in E(G)$. Define $\beta(v, w) = \beta(v, v')$ and $\beta(w, v) = \beta(w, w')$. The equations of part (a) of the statement guarantee that these values are well defined. Moreover, these definitions satisfy all the Naji equations listed under (b) in Definition 2.

We index the elements of V_1 and V_2 , $V_1 = \{v_1, ..., v_a\}$ and $V_2 = \{w_1, ..., w_b\}$, in such a way that $v_1w_1 \in E(G)$ and for i > 1,

$$N(v_i) \cap \{w_1, ..., w_{i-1}\} \neq \emptyset \neq N(w_i) \cap \{v_1, ..., v_i\}.$$

One way to construct such an indexing recursively is to find a leaf v of a spanning tree T for G, find an indexing of the specified type for T - v, and then list v as v_a or w_b according to whether $v \in V_1$ or $v \in V_2$.

To define the values of $\beta(v, w)$ with $vw \in E(G)$, begin by defining $\beta(v_1, w_1) = 0$ and $\beta(w_1, v_1) = 1$. If i > 1 and $v_1w_i \in E(G)$, define $\beta(v_1, w_i) = \beta(v_1, w_1) + \beta(w_i, w_1) + \beta(w_1, w_i) + 1$ and $\beta(w_i, v_1) = 1 + \beta(v_1, w_i)$. Interchange the letters v and w to define $\beta(w_1, v_i)$ and $\beta(v_i, w_1)$ if i > 1 and $w_1v_i \in E(G)$. It is easy to check that all the Naji equations involving v_1 or w_1 are satisfied. Suppose $i_0 > 1$ and all values of $\beta(v_i, w_j)$ and $\beta(w_j, v_i)$ have been defined when $i < i_0$ or $j < i_0$, in such a way that all Naji equations are satisfied. By hypothesis, v_{i_0} has a neighbor w_{j_0} with $j_0 < i_0$. If $j > i_0$ and $v_{i_0}w_j \in E(G)$, define $\beta(v_{i_0}, w_j) = \beta(v_{i_0}, w_{j_0}) + \beta(w_{j_0}, w_j) + 1$; equation (b) of the statement guarantees that this definition is independent of the choice of a particular $w_{j_0} \in N(v_{i_0})$. Also define $\beta(w_j, v_{i_0}) = 1 + \beta(v_{i_0}, w_j)$. The equations of the statement imply that all Naji equations involving v_{i_0} are satisfied. The values of $\beta(w_{i_0}, v_j)$ and $\beta(v_j, w_{i_0})$ when $j > i_0$ and $v_jw_{i_0} \in E(G)$ are defined in the same way, mutatis mutandi.

We should mention that a different way to reformulate Naji's theorem for bipartite graphs was given by Bouchet [5].

8 Permutation graphs

Here is a familiar definition; see [15] for instance.

Definition 26 Let π be a permutation of $\{1, ..., n\}$. Then the corresponding permutation graph has vertices 1, ..., n, with an edge ij whenever i < j and $\pi(i) > \pi(j)$.

Naji's theorem leads to the following algebraic characterization of permutation graphs. **Theorem 27** A simple graph G is a permutation graph if and only if this system of equations has a solution over GF(2).

(a) If v and w are two distinct vertices then $\beta(v, w) + \beta(w, v) = 1$.

(b) If v, w, x are three distinct vertices such that $vw, vx \notin E(G)$ and $wx \in E(G)$ then $\beta(v, w) + \beta(v, x) = 0$.

(c) If v, w, x are three distinct vertices such that $vw, vx \in E(G)$ and $wx \notin E(G)$ then $\beta(v, w) + \beta(v, x) = 0$.

Proof. Suppose G is the permutation graph corresponding to the permutation π . Let W be the oriented double occurrence word

$$1^{in}...n^{in}\pi(n)^{out}...\pi(1)^{out}.$$

Then the interlacement graph $\mathcal{I}(W)$ is G, and the Naji solution β corresponding to W satisfies the equations of the statement. For the converse, suppose β satisfies the equations of the statement, and let G + z be the graph obtained from G by adjoining a new vertex z adjacent to all the vertices of G. Extend β by defining $\beta(i, z) \equiv 0$ and $\beta(z, i) \equiv 1$. Then the extended β is a Naji solution for G + z, so G + z is a circle graph. If zW_1zW_2 is a double occurrence word with interlacement graph G + z then each vertex of G must appear once in each W_i , in order to be interlaced with z. Consequently W_1 and W_2 provide a permutation representation of G.

Another way to say the same thing is this: an *n*-vertex simple graph is a permutation graph if and only if it shares a Naji solution with K_n .

References

- A. Bouchet, Caractérisation des symboles croisés de genre nul, C. R. Acad. Sci. Paris Sér. A-B 274 (1972), A724-A727.
- [2] A. Bouchet, Reducing prime graphs and recognizing circle graphs, Combinatorica 7 (1987), 243-254.
- [3] A. Bouchet, Graphic presentation of isotropic systems, J. Combin. Theory Ser. B 45 (1988), 58-76.
- [4] A. Bouchet, Circle graph obstructions, J. Combin. Theory Ser. B 60 (1994), 107-144.
- [5] A. Bouchet, Bipartite graphs that are not circle graphs, Ann. Inst. Fourier (Grenoble) 49 (1999), 809–814.
- [6] H. R. Brahana, Systems of circuits on two-dimensional manifolds, Ann. Math. 23 (1921), 144-168.
- [7] M. Cohn and A. Lempel, Cycle decomposition by disjoint transpositions, J. Combin. Theory Ser. A 13 (1972), 83-89.

- [8] W. H. Cunningham, Decomposition of directed graphs, SIAM J. Alg. Disc. Meth. 3 (1982), 214-228.
- [9] S. Even and A. Itai, Queues, stacks, and graphs, in: Theory of machines and computations (Proc. Internat. Sympos., Technion, Haifa, 1971), pp. 71–86. Academic Press, New York, 1971.
- [10] H. de Fraysseix, Local complementation and interlacement graphs, Discrete Math. 33 (1981), 29-35.
- [11] E. Gasse, A proof of a circle graph characterization, Discrete Math. 173 (1997), 223-238.
- [12] J. F. Geelen, Matchings, matroids, and unimodular matrices (Thesis), Waterloo, 1995.
- [13] J. Geelen and B. Gerards, Characterizing graphic matroids by a system of linear equations, J. Combin. Theory Ser. B 103 (2013), 642-646.
- [14] E. Gioan, C. Paul, M. Tedder and D. Corneil, Practical and efficient circle graph recognition, Algorithmica 69 (2014), 759-788.
- [15] M. C. Golumbic, Algorithmic graph theory and perfect graphs, Academic Press, New York, 1980.
- [16] W. Naji, Graphes de cordes: une caracterisation et ses applications (Thèse), Grenoble, 1985.
- [17] W. Naji, Reconnaissance des graphes de cordes, Discrete Math. 54 (1985), 329-337.
- [18] R. C. Read and P. Rosenstiehl, On the Gauss crossing problem, in: Combinatorics (Proc. Fifth Hungarian Colloq., Keszthely, 1976), Vol. II, Colloq. Math. Soc. János Bolyai, 18, North-Holland, Amsterdam-New York, 1978, pp. 843-876.
- [19] J. Spinrad, Recognition of circle graphs, J. Algorithms 16 (1994), 264–282.
- [20] B. Zelinka, The graph of the system of chords of a given circle, Mat.-Fyz. Časopis Sloven. Akad. Vied 15 (1965), 273–279.