

**THE AUTOMORPHISM GROUPS OF ENRIQUES
SURFACES COVERED BY SYMMETRIC QUARTIC
SURFACES**

SHIGERU MUKAI AND HISANORI OHASHI

Dedicated to Prof. Robert Lazarsfeld on his 60th birthday

ABSTRACT. Let S be the (minimal) Enriques surface obtained from the symmetric quartic surface $(\sum_{i<j} x_i x_j)^2 = kx_1 x_2 x_3 x_4$ in \mathbb{P}^3 with $k \neq 0, 4, 36$, by taking quotient of the Cremona action $(x_i) \mapsto (1/x_i)$. The automorphism group of S is a semi-direct product of a free product \mathcal{F} of four involutions and the symmetric group \mathfrak{S}_4 . Up to action of \mathcal{F} , there are exactly 29 elliptic pencils on S .

The automorphism groups of very general Enriques surfaces, namely those corresponding to very general points in moduli, were computed in Barth-Peters[1] as an explicitly described infinite arithmetic group. Also many authors [3, 1, 9, 5] studied Enriques surfaces with only finitely many automorphisms. The article [1] also includes an example whose automorphism group is infinite but still virtually abelian group. In this paper we give a concrete example of an Enriques surface whose automorphism group is not virtually abelian. Moreover, the automorphism group is explicitly described in terms of generators and relations. See also Remark 5.

We work over any algebraically closed field whose characteristic is not two. Let us introduce the quartic surface with parameters k and l ,

$$(1) \quad \overline{X}: \{s_2^2 = ks_4 + ls_1 s_3\} \subset \mathbb{P}^3,$$

where s_d are the fundamental symmetric polynomials of degree d in the homogeneous coordinates x_1, \dots, x_4 . It is singular at the four coordinate points $(1 : 0 : 0 : 0), \dots, (0 : 0 : 0 : 1)$ and has an action of the symmetric group \mathfrak{S}_4 . It also admits the action of the standard Cremona transformation

$$\varepsilon: (x_1 : \dots : x_4) \mapsto \left(\frac{1}{x_1} : \dots : \frac{1}{x_4} \right)$$

which commutes with \mathfrak{S}_4 . After taking the minimal resolution X , the quotient surface $S = X/\varepsilon$ becomes an Enriques surface, whenever \overline{X} avoids the eight fixed points $(\pm 1 : \pm 1 : \pm 1 : 1)$ of ε . This condition is equivalent to $k + 16l \neq 36, k \neq 4$ and $4l + k \neq 0$.

2000 *Mathematics Subject Classification.* 14J28, 20F55.

Supported in part by the JSPS Grant-in-Aid for Scientific Research (B) 22340007, (S) 19104001, (S) 22224001, (S)25220701, (A) 22244003, for Exploratory Research 20654004 and for Young Scientists (B) 23740010.

The projection from one of four coordinate points exhibits X as a double cover of the projective plane \mathbb{P}^2 . The associated covering involution commutes with ε and defines an involution of the Enriques surface S . In this way we obtain four involutions σ_i ($i = 1, \dots, 4$). The action of \mathfrak{S}_4 also descends to S . Therefore, by mapping the generators of C_2^{*4} to σ_i , we obtain a group homomorphism

$$(2) \quad \mathfrak{S}_4 \times (C_2^{*4}) \rightarrow \text{Aut}(S),$$

where \mathfrak{S}_4 acts on the free product as permutation of the four factors.

In this paper we study the automorphism group and elliptic fibrations of S in the case $l = 0$. Our main result is as follows.

Theorem 1. (=Theorem 3.7) *In the equation (1), let $l = 0$ and $k \neq 0, 4, 36$. Then (2) is an isomorphism. Namely $\text{Aut}(S)$ is isomorphic to the semi-direct product of the free product \mathcal{F} of four involutions σ_i ($i = 1, \dots, 4$) and the symmetric group \mathfrak{S}_4 .*

In the proof of this theorem, we also obtain the following results on elliptic pencils and smooth rational curves. Let S be as in Theorem 1.

Theorem 2. (= Theorem 3.4) *Up to the action of the free product $\mathcal{F} \simeq C_2^{*4}$, there are exactly 29 elliptic pencils on S . They are classified into five types and the main properties are as in the following table.*

	singular fibers	Mordell-Weil rank	number
1)	$\tilde{E}_7 + \tilde{A}_1$	0	12
2)	$\tilde{E}_6 + \tilde{A}_2$	0	4
3)	$\tilde{D}_6 + \tilde{A}_1$	1	6
4)	$\tilde{A}_7 + \tilde{A}_1$	0	3
5)	$2\tilde{A}_5 + \tilde{A}_2 + \tilde{A}_1$	0	4

Here $2\tilde{A}_5$ denotes the multiple fiber and the Mordell-Weil rank stands for that of its Jacobian fibration.

Theorem 3. (= Theorem 3.3) *Up to the action of the free product $\mathcal{F} \simeq C_2^{*4}$, there are exactly sixteen smooth rational curves on S . They are represented by the curves in the configuration $10A + 6B$ (see below).*

The proof of Theorem 1 uses some sixteen smooth rational curves on S and the fact that four involutions $\sigma_1, \dots, \sigma_4$ are numerically reflective. First using the four singularities of type D_4 and four tropes on \overline{X} , we find ten smooth rational curves on S with the dual graph as in Figure 1 (Section 1). We call it the $10A$ configuration.

Also by looking at some other plane sections, we find further six smooth rational curves on S with the dual graph as in Figure 2. This is called the $6B$ configuration.

We denote by $NS(S)_f$ the Néron-Severi lattice of S modulo torsion. The action of involutions $\sigma_1, \dots, \sigma_4$ on $NS(S)_f$ is the reflection in (-2) classes




FIGURE 1. The 10A configuration.




FIGURE 2. The 6B configuration

$G_1, \dots, G_4 \in NS(S)_f$, respectively. For instance, the class G_1 is $E_2 + E_{23} + E_3 + E_{34} + E_4 + E_{24} - E_1$ in terms of Figure 1 (Proposition 2.1). The dual graph of these four (-2) classes is the complete graph in four vertices with doubled edges. This is what is called the 4C configuration.

We can check that the twenty (-2) -classes E_i, E_{ij}, F_{ij}, G_i define a convex polyhedron whose Coxeter diagram satisfies the Vinberg's condition [12]. Namely the subgroup $W(10A + 6B + 4C)$ generated by reflections in these twenty classes has finite index in the orthogonal group $O(NS(S)_f)$. In fact, the limit of our Enriques surfaces as $k \rightarrow \infty$ is of type V in Kondo[5] (see Remark 2.3), and our diagram coincides with Kondo's. Although in his case the classes G_1, \dots, G_4 were also represented by smooth rational curves, in our case they appear just as the *center* of the reflective involutions $\sigma_1, \dots, \sigma_4$ and are not effective (Corollary 2.2).

To prove our Theorem 1, we divide the generators of $W(10A + 6B + 4C)$ into two parts, those coming from $10A + 6B$ and those from $4C$. By a lemma of Vinberg[11], $W(10A + 6B + 4C)$ is the semi-direct product $W(4C) \ltimes \overline{N}(W(10A + 6B))$, where \overline{N} denotes the normal closure. Since the whole $10A + 6B + 4C$ configuration has only \mathfrak{S}_4 -symmetry, we obtain our Theorem 1 and the others (Sections 3).

Remark 4. There are some interesting cases in $l \neq 0$, too.

(1) When $(k-4)(l-4) = 16$, the surface \overline{X} is Kummer's quartic surface $\text{Km}(J(C))$ written in Hutchinson's form. It has 16 nodes. Our four involutions σ_i are called *projections*. As is shown in [7], S is an Enriques surface of Hutchinson-Göpel type and the four involutions are numerically reflective. Especially in the case $(k, l) = (-4, 2)$, the hyperelliptic curve C branches over the vertices of regular octahedron and the equation of \overline{X} becomes

$$(x_1^2 x_2^2 + x_3^2 x_4^2) + (x_1^2 x_3^2 + x_2^2 x_4^2) + (x_1^2 x_4^2 + x_2^2 x_3^2) + 2x_1 x_2 x_3 x_4 = 0.$$

This is the case of the octahedral Enriques surface [8] and S is isomorphic to the normalization of the singular sextic surface

$$x_1^2 + x_2^2 + x_3^2 + x_4^2 + \sqrt{-1} \left(\frac{1}{x_1^2} + \frac{1}{x_2^2} + \frac{1}{x_3^2} + \frac{1}{x_4^2} \right) x_1 x_2 x_3 x_4 = 0.$$

In these cases we know that there exist automorphisms on S induced from X other than projections, namely some switches and correlations. The automorphism group of the octahedral Enriques surface will be discussed elsewhere.

(2) The quartic surface $\overline{X} : ks_4 + ls_1 s_3 = 0$ is the Hessian of the cubic surface

$$k(x_1^3 + x_2^3 + x_3^3 + x_4^3) + l(x_1 + x_2 + x_3 + x_4)^3 = 0.$$

The case $(k : l) = (1 : -1)$ is most symmetric among this one-parameter family. In this special case, the Enriques surface $S = X/\varepsilon$ is of type VI in Kondo[5] and the automorphism group is isomorphic to \mathfrak{S}_5 . In particular, the homomorphism (2) is neither injective nor surjective.

Remark 5. In terms of virtual cohomological dimensions of discrete groups [10], our example can be located in the following way. The virtual cohomological dimension is equal to 0 for finite groups. On the other extreme, the discrete group $\text{Aut}(S)$ for very general Enriques surfaces S has the virtual cohomological dimension 8. See [2]. In our case, the automorphism group has virtual cohomological dimension 1.

1. SMOOTH RATIONAL CURVES

Under the condition $l = 0$, the equation (1) becomes

$$(3) \quad \overline{X} : (x_1 x_2 + x_1 x_3 + x_1 x_4 + x_2 x_3 + x_2 x_4 + x_3 x_4)^2 = k x_1 x_2 x_3 x_4.$$

This surface has four rational double points of type D_4 at the four coordinate points $(1 : 0 : 0 : 0), \dots, (0 : 0 : 0 : 1)$ and by taking the quotient of the minimal resolution X by the standard Cremona involution

$$\varepsilon : (x_1 : \dots : x_4) \mapsto \left(\frac{1}{x_1} : \dots : \frac{1}{x_4} \right),$$

we obtain an Enriques surface $S = X/\varepsilon$. We begin with the study of the configuration of smooth rational curves on the surfaces.

The desingularization X has sixteen smooth rational curves as the exceptional curves of the four D_4 singularities. Also each coordinate plane cuts




FIGURE 3. The quartic surface with four D_4 singularities.

the quartic doubly along a conic, which defines a smooth rational curve on X . They are called *tropes*. The configuration of these twenty curves is as in Figure 3, which depicts the dual graph. Black vertices come from the singularities and white ones are tropes.

The standard Cremona involution ε acts on Figure 3 by the point symmetry. Therefore the Enriques surface S has ten smooth rational curves whose dual graph is the one in Figure 1. In what follows, we call these ten curves on S the $10A$ configuration. The indexing is given as follows. Since a vertex of the tetrahedron corresponds to two curves on X , namely the trope $\{x_i = 0\}$ and the central component of the exceptional curves at $(0 : \cdots : 1 : \cdots : 0)$ (the i -th coordinate is 1), we denote the curve at the vertex by E_i ($i = 1, \dots, 4$). Also if a vertex at the middle of an edge is connected to two vertices, say E_i and E_j , then we denote the curve by E_{ij} . This is the first configuration of smooth rational curves on S of our interest. It is convenient to note that the ten curves $\{E_i, E_{ij}\}$ generate $NS(S)_f$ over the rationals; the Gram matrix of these curves has determinant -64 .


Next let us consider the six plane sections by $\{x_i + x_j = 0\}$ ($i = 1, \dots, 4$). In the equation (3), we see that each plane section decomposes into two conics which are disjoint on X and exchanged by ε . Thus we obtain further six smooth rational curves on S , naturally indexed as F_{ij} . The intersection relation between these curves is as in Figure 2. We call it the $6B$ configuration. Moreover, we can clarify the intersection relations between the configurations as follows.

$$(E_k, F_{ij}) = 0; \quad (E_{kl}, F_{ij}) = \begin{cases} 2 & \text{if } \{k, l\} = \{i, j\}, \\ 0 & \text{otherwise.} \end{cases}$$

The configuration of sixteen curves thus obtained is denoted by $10A + 6B$.

2. NUMERICALLY REFLECTIVE INVOLUTIONS

The quartic surface (3) can be exhibited as a double cover of \mathbb{P}^2 by the projection from one of the coordinate points, say $(0 : 0 : 0 : 1)$. The branch

FIGURE 4. The branch sextic B

$B \subset \mathbb{P}^2$ is the sextic plane curve defined by

$$(4) \quad x_1 x_2 x_3 \left(4(x_1 + x_2 + x_3) \left(\frac{1}{x_1} + \frac{1}{x_2} + \frac{1}{x_3} \right) x_1 x_2 x_3 - k x_1 x_2 x_3 \right) = 0.$$

It is the union of the coordinate triangle $\{x_1 x_2 x_3 = 0\}$ and the cubic curve

$$(5) \quad C: 4(x_1 + x_2 + x_3) \left(\frac{1}{x_1} + \frac{1}{x_2} + \frac{1}{x_3} \right) - k = 0,$$

which is invariant under the Cremona transformation $(x_i) \mapsto (1/x_i)$ of \mathbb{P}^2 . See Figure 4. In this double plane picture, the twenty rational curves in Figure 3 can be seen as the twelve rational curves above the three triple points of B , three rational curves above the three nodes of B , three tropes as the inverse image of the coordinate triangle and some components of inverse images of the curves $L: \{x_1 + x_2 + x_3 = 0\}$ and $Q: \{\frac{1}{x_1} + \frac{1}{x_2} + \frac{1}{x_3} = 0\}$. (We note that the line L must pass through the three simple intersection points of C with the triangle in Figure 4, although it is not visible.)

The covering transformation of this double cover $X \rightarrow \mathbb{P}^2$ is called the *projection*. It is an anti-symplectic involution acting on X . It stabilizes all the curves above the branch curve B (including the ones above the singularities of B). In particular, in Figure 3, if E_4'' is the trope $\{x_4 = 0\}$, then the projection stabilizes all the curves except for E_4'' and its antipodal E_4' (coming from the singularity at $(0 : 0 : 0 : 1)$). It is easy to determine the fixed curves of the projection and it consists of six rational curves (vertices of the cube except for E_4' and E_4'') and the inverse image of the elliptic curve C .

Since the projection commutes with the Cremona involution of \mathbb{P}^3 , we obtain an involution of the Enriques surface S . It is denoted by σ_4 , where the index is in accordance to the center of the projection $(0 : 0 : 0 : 1)$.

Proposition 2.1. *The involution $\sigma_4 \in \text{Aut}(S)$ is numerically reflective. Moreover, its action on the Néron-Severi lattice $NS(S)_f$ is the reflection in the divisor $G_4 = E_1 + E_{12} + E_2 + E_{23} + E_3 + E_{13} - E_4$ of self-intersection (-2) . In Figure 1, the six positive components in G_4 are just the cycle of curves disjoint from E_4 .*

Proof. From our description of fixed curves of the projection as above, we see that σ_4 preserves all the curves E_i and E_{ij} except for E_4 . Compare Figures 1 and 3.

Consider the elliptic fibration $f: S \rightarrow \mathbb{P}^1$ defined by the divisor $2D_f = 2(E_1 + E_{12} + E_2 + E_{23} + E_3 + E_{13})$. It gives the multiple fiber of type ${}_2I_6$ in Kodaira's notation. From Figure 1, we see that the curve E_4 sits inside a reducible fiber which we denote by D' . In comparison with Figure 4, f corresponds to the pencil \mathcal{L} of cubics on \mathbb{P}^2 spanned by the triangle $\{x_1x_2x_3 = 0\}$ and the cubic curve C of (5). Thus we see that the multiple fibers of f are exactly the transform of the triangle, which is nothing but the divisor $2D_f$ of type ${}_2I_6$, and the transform of C , namely some irreducible fiber of type ${}_2I_0$. On the other hand, the cubic

$$(6) \quad C_\infty := L + Q \in \mathcal{L}$$

corresponds to the reducible fiber of f which contains E_4 . Therefore the fiber $D' = E_4 + B$ is of Dynkin type \tilde{A}_1 and is not multiple. (More precisely, it is of type III in characteristic three and otherwise I_2 .) Since the Cremona involution of \mathbb{P}^2 interchanges L and Q , we see that σ_4 interchanges E_4 and B .

From the linear equivalence $E_4 + B \sim 2(E_1 + E_{12} + E_2 + E_{23} + E_3 + E_{13})$, we see that the action is

$$\sigma_4: E_4 \mapsto B = 2(E_1 + E_{12} + E_2 + E_{23} + E_3 + E_{13}) - E_4.$$

By taking the first paragraph into account, we see that σ_4 is numerically reflective and acts on $NS(S)_f$ by the reflection in the divisor

$$G_4 = E_1 + E_{12} + E_2 + E_{23} + E_3 + E_{13} - E_4.$$

□

By symmetry, we obtain divisors $G_i (i = 1, \dots, 4)$ which describe the numerically reflective involutions σ_i in a similar manner. We see that $(G_i, G_j) = 2$ for $i \neq j$ so that the intersection diagram associated to divisors G_1, \dots, G_4 is the complete graph in four vertices with all edges doubled. In what follows we denote this configuration by $4C$.

We note that the automorphism σ_i sends G_i to its negative. It implies the following corollary.

Corollary 2.2. *The numerical classes of G_i are not effective.*

We can compute the intersections of G_i and the $10A + 6B$ configuration. We have the following.

$$(G_i, E_j) = \begin{cases} 2 & \text{if } i = j, \\ 0 & \text{otherwise,} \end{cases} \quad (G_i, E_{kl}) = 0, \quad (G_i, F_{kl}) = \begin{cases} 2 & \text{if } i \notin \{k, l\} \\ 0 & \text{if } i \in \{k, l\}. \end{cases}$$

Remark 2.3. The limit of our quartic surface \overline{X} in (3) as $k \rightarrow \infty$ is the double \mathbb{P}^2 with branch the union of the coordinate triangle and the reducible cubic C_∞ in (6). Hence the limit of our Enriques surfaces is of type V in

Kondo[5]. (See [6] also.) In this limit our divisor class G_4 becomes effective and corresponds to the new singular point coming from the intersection $L \cap Q$ in (6). Thus G_4 can be regarded as the vanishing cycle of this specialization $k \rightarrow \infty$. Furthermore the numerically reflective involution σ_4 becomes numerically trivial in this limit. More precisely, the limit of its graph as $k \rightarrow \infty$ is the union of that of the limit involution and the product $C_4 \times C_4$, where C_4 is the unique (-2) -curve representing G_4 in the limit Enriques surface.

3. PROOF OF THE THEOREMS

In the previous two sections, we obtained sixteen smooth rational curves with the configuration $10A + 6B$ and four numerically reflective involutions σ_i whose centers G_i have the configuration $4C$. We begin with the consideration of the natural representation $r: \text{Aut}(S) \rightarrow O(NS(S)_f)$.

Proposition 3.1. *The homomorphism r is injective, namely there are no nontrivial numerically trivial automorphisms on S .*

Proof. Let g be a numerically trivial automorphism of S , which is tame by virtue of Dolgachev[4]. It preserves each (-2) -curves, in particular, each in the $10A + 6B$ configuration. The curves E_1, \dots, E_4 in Figure 1 must be pointwise fixed, since $\text{Aut}(\mathbb{P}^1)$ is sharply triply transitive and since each E_i has three distinct intersections with its neighbors.

We again focus on the elliptic fibration $f: S \rightarrow \mathbb{P}^1$ defined by $D_f = E_1 + E_{12} + E_2 + E_{23} + E_3 + E_{13}$ as in Proposition 2.1. We saw that $E_4 + B = E_4 + \sigma_4(E_4)$ is a non-multiple fiber of f . Therefore the bisections E_{14}, E_{24}, E_{34} of f must intersect B . By a suitable choice of a bisection $C_f \in \{E_{14}, E_{24}, E_{34}\}$, we can assume that C_f does not pass through the intersection $E_4 \cap B$. Then, since g preserves all (-2) -curves, the curve C_f has three distinct fixed points $E_4 \cap C_f$, $B \cap C_f$ and $E_i \cap C_f$, where E_i is the another vertex of the edge containing C_f in Figure 1. It follows that g fixes C_f pointwise, hence the singular curve $E_4 + C_f$, too. It follows that $g = \text{id}_S$, since g is tame and of finite order. \square

In what follows we denote the hyperbolic lattice $NS(S)_f$ by L . Let us denote by $O'(L)$ the group of integral isometries whose \mathbb{R} -extensions preserve the positive cone of $L \otimes \mathbb{R}$. We denote by Λ the 9-dimensional Lobachevsky space associated to the positive cone. Then $O'(L)$ acts on Λ as a discrete group of motions. We refer the readers to [12] for the theory of discrete groups generated by reflections acting on Lobachevsky spaces.

We let

$$P^c = \{\mathbb{R}_+ x \in PS(L) \mid (x, E) \geq 0 \text{ for all } E \in \{E_i, E_{ij}, F_{ij}, G_i\}\}$$

be the convex polyhedron defined by the twenty roots from the $10A + 6B + 4C$ configuration in the projective sphere $PS(L) = (L - \{0\})/\mathbb{R}_+$ (see [12, Section 2]). We have seen that every intersection number of distinct two divisors

in $10A + 6B + 4C$ is in between 0 and 2, hence the Coxeter diagram associated to these twenty roots has no dotted lines nor Lanner's subdiagrams. Also by an easy check of the $10A + 6B + 4C$ configuration, we have the following.

Lemma 3.2. *The Coxeter diagram of the polyhedron $P = P^c \cap \Lambda$ has exactly 29 parabolic subdiagrams of maximal rank 8. They are as follows.*

	the subdiagram	number	10A	6B	4C
1)	$\tilde{E}_7 + \tilde{A}_1$	12	8	1	1
2)	$\tilde{E}_6 + \tilde{A}_2$	4	7	3	0
3)	$\tilde{D}_6 + \tilde{A}_1 + \tilde{A}_1$	6	8	1	2
4)	$\tilde{A}_7 + \tilde{A}_1$	3	8	2	0
5)	$\tilde{A}_5 + \tilde{A}_2 + \tilde{A}_1$	4	7	3	1

Here the each column 10A, 6B, 4C shows the number of vertices used from the configuration.

It is easy to check that every connected parabolic subdiagram is a connected component of some parabolic subdiagram of rank 8, using the previous table. By Theorem 2.6 bis of [12], we see that P has finite volume and we obtain $P^c \subset \bar{\Lambda}$. This polyhedron gives the fundamental domain of the associated discrete reflection group generated by twenty reflections in the twenty roots $\{E_i, E_{ij}, F_{ij}, G_i\}$, which we denote by $W = W(10A + 6B + 4C)$. Algebra-geometrically, sixteen of the generators are the Picard-Lefschetz transformations in (-2) -curves in the $10A + 6B$ configuration and the rest four are the involutions σ_i ($i = 1, \dots, 4$) corresponding to $4C$. As an abstract group, we see that W has the structure of a Coxeter group whose fundamental relations are given by the Coxeter diagram (see [12]) of P . We note that the quasi-polarization (namely a nef and big divisor)

$$H = \sum_i E_i + \sum_{i < j} E_{ij}$$

defines an element $\mathbb{R}_+ H$ in P .

Now let $W(4C)$ be the subgroup of W generated by four reflections in G_i . Via the homomorphism $r: \text{Aut}(S) \rightarrow O(NS(S)_f)$, the subgroup $\mathcal{F} \subset \text{Aut}(S)$ generated by the four numerically reflective involutions σ_i is mapped onto this Coxeter subgroup $W(4C) \simeq C_2^{*4}$. It follows that $\mathcal{F} \simeq W(4C)$. Let $W(10A + 6B)$ be the subgroup generated by sixteen reflections in E_i, E_{ij} and F_{ij} and let $\overline{N}(W(10A + 6B))$ be the minimal normal subgroup of W which contains $W(10A + 6B)$. Since the intersection numbers between elements of $4C$ and $10A + 6B$ are all even, by [11, Proposition], we have the exact sequence

$$(7) \quad 1 \longrightarrow \overline{N}(W(10A + 6B)) \longrightarrow W \longrightarrow W(4C) \longrightarrow 1.$$

The kernel is exactly the subgroup generated by the conjugates

$$\{\sigma g \sigma^{-1} \mid \sigma \in W(4C), g \text{ a generator of } W(10A + 6B)\}.$$

We have the corresponding geometric consequence as follows.

Theorem 3.3. *There are exactly sixteen smooth rational curves on S up to the action of \mathcal{F} .*

Proof. Let E be a smooth rational curve on S . We consider the orbit $\mathcal{F}.E$. Since the divisor H above is nef, we can choose $E_0 \in \mathcal{F}.E$ such that the degree (E_0, H) is minimal. We shall show that E_0 is one of sixteen curves in $10A + 6B$.

In fact, by the automorphism σ_i , we have

$$(E_0, H) \leq (\sigma_i(E_0), H) = (E_0, H) + (E_0, G_i)(G_i, H),$$

and $0 \leq (E_0, G_i)$ for all i . Suppose that E_0 intersects non-negatively to all the sixteen curves in $10A + 6B$. Then we have $\mathbb{R}_+E_0 \in P^c$. But from $P^c \subset \overline{\Lambda}$, we obtain $(E_0^2) \geq 0$, which is a contradiction. Hence E_0 is negative on some curve in $10A + 6B$ and we see that E_0 is one of them.

Next let us show that two distinct curves E, E' in the $10A + 6B$ configuration are inequivalent under \mathcal{F} . For the six curves E_{ij} from $10A$, we have $(E_{ij}, G_k) = 0$ for all $i < j$ and k . Therefore, by an easy induction, we see that the sextuple $((E_{ij}, E))_{1 \leq i < j \leq 4}$ consisting of intersection numbers is an invariant of the orbit $\mathcal{F}.E$. Suppose that $(E_{ij}, E) = (E_{ij}, E')$ for all $i < j$. Since E and E' both are in the $10A + 6B$ configuration, we see easily that $E = E'$. This shows that the orbits $\mathcal{F}.E$ and $\mathcal{F}.E'$ are disjoint. \square

In another words, the group $\overline{N}(W(10A + 6B))$ is nothing but the Weyl group of S generated by Picard-Lefschetz reflections in all (-2) -curves. We can proceed to elliptic pencils.

Theorem 3.4. *There are exactly 29 elliptic pencils on S up to the action of \mathcal{F} . Their properties are as in the table of Theorem 2.*

Proof. Let $2f$ be a fiber class of an elliptic pencil on S . As before, we choose an element $f_0 \in \mathcal{F}.f$ such that the degree (f_0, H) is minimal. We have

$$(f_0, H) \leq (\sigma_i(f_0), H) = (f_0, H) + (f_0, G_i)(G_i, H),$$

hence $(f_0, G_i) \geq 0$ for all i . Moreover since f_0 is nef we have $(f_0, E) \geq 0$ for all E in the $10A + 6B$ configuration. Therefore $f_0 \in P^c$. This shows that f_0 corresponds to one of maximal parabolic subdiagrams classified in Lemma 3.2.

Conversely, we can construct 29 elliptic pencils from the 29 subdiagrams in Lemma 3.2 as follows. The two types 2) and 4) in the lemma are easiest since they do not contain a class in $4C$. The elliptic pencils of types 2) and 4) have singular fibers of type $\tilde{E}_6 + \tilde{A}_2$ and $\tilde{A}_7 + \tilde{A}_1$, respectively as in the case of [5, Table 2].

In the case of type 1) (resp. 5)), one component of the parabolic subdiagram is an \tilde{A}_1 consisting of a (-2) -curve E in $6B$ (resp. $10A$) and (-2) -class G in $4C$. Moreover, the sum $E+G$ is a half of \tilde{E}_7 (resp. \tilde{A}_2). Hence $E+\sigma(E)$ is a non-multiple fiber of type \tilde{A}_1 since it is linearly equivalent to $2(E+G)$, where σ is the reflection in G .

In the case of type 3), one component is an \tilde{A}_1 consisting of two classes G and G' in $4C$. But the other two components consisting of (-2) -curves. Therefore, the Mordell-Weil group is of rank one since neither G or G' is effective. (The composite $\sigma\sigma'$ of two reflections in G and G' is the translation by a generator of the Mordell-Weil group.)

That these 29 pencils are inequivalent under \mathcal{F} follows from the previous result for rational curves. \square

To study the image of the representation $r: \text{Aut}(S) \rightarrow O(NS(S)_f)$, we need some lemma. We denote by $4A'$ the set of four roots $\{E_i\}$ and by $6A''$ the set $\{E_{ij}\}$. Recall that by Theorem 3.3, all (-2) -curves on S are in the \mathcal{F} -orbit of the three sets $4A'$, $6A''$ and $6B$.

Lemma 3.5. *Let τ be any automorphism of S . Then τ preserves each of the three orbits of rational curves $\mathcal{F}.(4A')$, $\mathcal{F}.(6A'')$ and $\mathcal{F}.(6B)$.*

Proof. Any automorphism τ permutes smooth rational curves on S , hence induces a symmetry of the dual graph of the set of rational curves. Thus, for the proof, it suffices to give a characterization of each orbit in terms of this infinite graph. We use the (full) subgraphs which are isomorphic to the dual graph of reducible fibers of elliptic fibrations.

Consider a vertex v in $\mathcal{F}.(6B)$. Then there exists a subgraph of fiber type I_3 passing through v . Conversely, if for a vertex v there is a subgraph of fiber type I_3 , by Theorem 3.4, it is equivalent to a vertex in $6B$ under \mathcal{F} . Thus the vertices in $\mathcal{F}.(6B)$ are characterized by the property that there exists a subgraph of fiber type I_3 passing through them.

Similarly, vertices in $\mathcal{F}.(10A)$ are characterized by subgraphs of type I_8 . Moreover, the vertices v in $\mathcal{F}.(4A')$ are characterized by the property that there exists a subgraph of type IV^* which has v as its end. In the opposite way, vertices in $\mathcal{F}.(6A'')$ are those which does not have such IV^* subgraphs. Thus the three orbits are all characterized and τ preserves these orbits. \square

Corollary 3.6. *The set of six curves $\{E_{ij}\}$ is preserved under any automorphism.*

Proof. In fact, for any E_{kl} and any σ_i we have $\sigma_i(E_{kl}) = E_{kl}$. Hence $\mathcal{F}.(6A'') = \{E_{ij}\}$. \square

Recall that S has the action by \mathfrak{S}_4 from the symmetry of the defining equation of \bar{X} . Explicitly, it acts on the curves in $10A + 6B$ configuration by the permutation of indices. For involutions σ_i , the same holds true if we regard the action as taking conjugates. It is easy to see that this group \mathfrak{S}_4 can be identified with the symmetry group $\text{Sym}(P)$ of the polyhedron

$P \subset \Lambda$ via r . We can also regard this group as acting on the reflection group W and the exact sequence (7) is preserved under this action. In particular, $W(4C)$ and $\text{Sym}(P)$ generate a group isomorphic to $\mathfrak{S}_4 \times C_2^{*4}$.

Theorem 3.7. *The representation r induces an isomorphism of $\text{Aut}(S)$ onto the group generated by $W(4C)$ and $\text{Sym}(P)$, hence we obtain $\text{Aut}(S) \simeq \mathfrak{S}_4 \times C_2^{*4}$.*

Proof. Since r maps \mathcal{F} onto $W(4C)$, the image of r includes the groups $W(4C)$ and $\text{Sym}(P)$.

Conversely let us pick up an arbitrary automorphism τ . We consider the image $\tau(H)$ of H . We use the elliptic fibration defined by the divisor $f = H - E_{12} - E_{34}$ of type I_8 . By Theorem 3.4, the image $\tau(f)$ is equivalent to one of three elliptic pencils described in item 4) under \mathcal{F} . Moreover, since \mathfrak{S}_4 acts transitively on these three pencils, we can assume that $\tau(f) = f$ by composing τ with some elements of \mathcal{F} and \mathfrak{S}_4 . Thus we have $\tau(H) = f + \tau(E_{12}) + \tau(E_{34})$. By the previous corollary, $\tau(E_{12})$ and $\tau(E_{34})$ are in the set $\{E_{ij}\}$. By an easy check of intersection numbers, we see that $\tau(E_{12}) + \tau(E_{34}) = E_{12} + E_{34}$. In particular we obtain $\tau(H) = H$ as divisors. Since any permutation of the $10A$ configuration can be induced from the automorphism group \mathfrak{S}_4 , this shows that the image of r is contained in the group generated by $W(4C)$ and $\text{Sym}(P)$. \square

REFERENCES

- [1] W. Barth and C. Peters, Automorphisms of Enriques surfaces, *Invent. Math.*, **73** (1983), 383–411.
- [2] A. Borel and J. P. Serre, Corners and arithmetic groups, *Comment. Math. Helv.*, **48** (1973), 436–491.
- [3] I. Dolgachev, On automorphisms of Enriques surfaces, *Invent. Math.*, **76** (1984), 163–177.
- [4] I. Dolgachev, Numerical trivial automorphisms of Enriques surfaces in arbitrary characteristic, in *Arithmetic and Geometry of K3 surfaces and Calabi-Yau threefolds*, Fields Institute Communications **67**, 2013, pp. 267–283.
- [5] S. Kondo, Enriques surfaces with finite automorphism groups, *Japan. J. Math.*, **12** (1986), 191–282.
- [6] S. Mukai, Numerically trivial involutions of Kummer type of an Enriques surface, *Kyoto J. Math.*, **50** (2010), 889–902.
- [7] S. Mukai, Kummer’s quartics and numerically reflective involutions of Enriques surfaces, *J. Math. Soc. Japan*, **64** (2012), 231–246.
- [8] S. Mukai and H. Ohashi, Enriques surfaces of Hutchinson-Göpel type and Mathieu automorphisms, in *Arithmetic and Geometry of K3 surfaces and Calabi-Yau threefolds*, Fields Institute Communications **67**, 2013, pp. 429–454.
- [9] V. V. Nikulin, On a description of the automorphism groups of Enriques surfaces, *Soviet Math. Dokl.*, **30** (1984), 282–285.
- [10] J. P. Serre, Cohomologie des groupes discrets, *Prospects in mathematics (Proc. Symp., Princeton Univ., Princeton, N.J., 1970)*, pp. 77–169. *Ann. of Math. Studies*, No. 70, Princeton Univ. Press, Princeton, N.J., 1971.
- [11] E. B. Vinberg, The two most algebraic $K3$ surfaces, *Math. Ann.*, **265** (1983), 1–21.

- [12] E. B. Vinberg, Some arithmetical discrete groups in Lobačevskii spaces, in *Discrete Subgroups of Lie Groups and Appl. to Moduli (Bombay 1973)*, Oxford University Press, 1975, pp. 323–348.

RESEARCH INSTITUTE FOR MATHEMATICAL SCIENCES, KYOTO UNIVERSITY, KYOTO
606-8502, JAPAN

E-mail address: mukai@kurims.kyoto-u.ac.jp

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND TECHNOLOGY, TOKYO
UNIVERSITY OF SCIENCE, 2641 YAMAZAKI, NODA, CHIBA 278-8510, JAPAN

E-mail address: ohashi@ma.noda.tus.ac.jp, ohashi.hisanori@gmail.com