Lyapunov exponents and strong exponential tails for some contact Anosov flows

BY LUCHEZAR STOYANOV

Abstract. For the time-one map f of a contact Anosov flow on a compact Riemann manifold M, satisfying a certain regularity condition, we show that given a Gibbs measure on M, a sufficiently large Pesin regular set P_0 and an arbitrary $\delta \in (0, 1)$, there exist positive constants C and c such that for any integer $n \ge 1$, the measure of the set of those $x \in M$ with $f^k(x) \notin P_0$ for at least δn values of $k = 0, 1, \ldots, n-1$ does not exceed Ce^{-cn} .

1 Introduction

Let $\phi_t : M \longrightarrow M$ be a C^2 Anosov flow on a C^2 compact Riemann manifold M, and let $f = \phi_1$ be its *time-one map*.

It follows from a well-know result of Oseledets ([Os]; see also [BP] or [R]) that there exists a Borel subset \mathcal{L}_0 of M, which has full measure with respect to any f-invariant Borel probability measure on M, such that for every $x \in \mathcal{L}_0$ there exists a df-invariant decomposition

$$T_x M = E_1(x) \oplus E_2(x) \oplus \ldots \oplus E_{k(x)}(x)$$

and numbers $\chi_1(x) < \chi_2(x) < \ldots < \chi_{k(x)}(x)$, called Lyapunov characteristic exponents, such that:

(a) $\lim_{|n|\to\infty} \frac{1}{n} \log \|df_x^n(v)\| = \chi_i(x) \text{ for all } v \in E_i(x) \setminus \{0\} \text{ and all } i = 1, \dots, k(x).$

(b) For every $\epsilon > 0$ there exists a Borel function $A_{\epsilon} : \mathcal{L}_0 \longrightarrow (1, \infty)$, such that

$$\frac{\|v\|}{A_{\epsilon}(x) e^{|n|\epsilon}} \le \frac{\|df_x^n(v)\|}{e^{n\chi_i(x)}} \le A_{\epsilon}(x) e^{|n|\epsilon} \|v\| \quad , \quad v \in E_i(x) \ , \ n \in \mathbb{Z},$$
(1.1)

for all $x \in \mathcal{L}_0$ and all $i = 1, \ldots, k(x)$, and

$$e^{-\epsilon} \le \frac{A_{\epsilon}(f(x))}{A_{\epsilon}(x)} \le e^{\epsilon} \quad , \quad x \in \mathcal{L}_0.$$
 (1.2)

(c) For all $x \in \mathcal{L}_0$ and all disjoint non-empty subsets I, I' of $\{1, \ldots, k(x)\}$ the smallest angle between non-zero vectors in $E_I(x) = \bigoplus_{i \in I} E_i(x)$ and $E_{I'}(x)$ is $\geq \frac{1}{A_e(x)}$.

(d) If m is an ergodic f-invariant Borel probability measure on M, then the functions k(x) and $\chi_i(x)$ are constant m-almost everywhere.

A function A_{ϵ} satisfying (1.2) is called an ϵ -slow varying function.

Let Φ be a Hölder continuous real-valued function on M and let **m** be the *Gibbs measure* generated by Φ on M (see e.g. [P2] or [PP]). Then **m** is ergodic, so there exists a subset \mathcal{L}'_0 of \mathcal{L}_0 with $\mathbf{m}(\mathcal{L}'_0) = 1$ such that the functions $k(x) = k_0$ and $\chi_i(x) = \chi_i$ are constant on \mathcal{L}'_0 .

It follows e.g. from the arguments in Sect. 3 in [PS], that for every $\epsilon > 0$ there exist constants $\epsilon' \in (0, \epsilon], p \ge 1$ and $\nu > 0$ such that $r_{\epsilon}(x) = \frac{\nu}{(A_{\epsilon'}(x))^p}$ determines an ϵ -slowly varying function on \mathcal{L}'_0 which defines a Lyapunov regular neighbourhood for every $x \in \mathcal{L}'_0$, i.e. for each $x \in \mathcal{L}'_0$ there exists a Lyapunov chart on $B(x, r_{\epsilon}(x))$. On these charts one has estimates of the iterations of the non-linear map f similar to these in (1.1).

It is known that in general the complement of the set \mathcal{L}_0 (and therefore that of \mathcal{L}'_0) can be topologically very large (see [BSau], [BS] or [PSa] for some interesting examples). The regularity functions A_{ϵ} and r_{ϵ} are in general only measurable. The so called *Pesin regular sets*

$$\mathcal{R}_{\ell} = \{ x \in \mathcal{L}'_0 : A_{\epsilon}(x) \le \ell , \ r_{\epsilon}(x) \ge 1/\ell \} \quad , \quad \ell \ge 1,$$

and their closures are of particular importance since on such sets uniform estimates involving Lyapunov exponents are available (see e.g. [P1], [BP], [KM], [LY1], [LY2], [PS], [BPS]). However it seems there is little information in the literature about the measures of the sets \mathcal{R}_{ℓ} and 'how quickly' they fill in \mathcal{L}'_0 , even in the case of uniformly hyperbolic systems.

Let **m** be an ergodic f-invariant Borel probability measure on M, and let k_0 be so that $k(x) = k_0$ for **m**-almost all x. Consider the distributions

$$E^{(i)}(x) = E_1(x) \oplus E_2(x) \oplus \ldots \oplus E_i(x) \quad , \quad \widetilde{E}^{(i)}(x) = E_{i+1}(x) \oplus \ldots \oplus E_{k_0}(x).$$

We will say that $E^{(i)}(x)$ is uniformly continuous in \mathcal{L}'_0 if the map $\mathcal{L}'_0 \ni x \mapsto E^{(i)}(x)$ is uniformly continuous with respect to the natural distance between distributions with the same dimension (see e.g. Sect. 2.3 in [BP]).

Given a Lyapunov regularity function A_{ϵ} , an integer $p \geq 0$ and a constant $\delta \in (0,1)$ set

$$T_p = \{ x \in \mathcal{L}'_0 : A_\epsilon(x) \le e^p \},\$$

and denote by $\Gamma_n = \Gamma_n(\epsilon, \delta, p)$ the set of all $x \in \mathcal{L}'_0$ with $f^k(x) \notin T_p$ for at least δn values of $k = 0, 1, \ldots, n-1$.

In this paper we prove the following.

Theorem 1.1. Let $\phi_t : M \longrightarrow M$ be a C^2 contact Anosov flow on a C^2 compact Riemann manifold M, let Φ be a Hölder continuous real-valued function on M and let \mathbf{m} be the Gibbs measure generated by Φ on M. Assume in addition that for every $i = 1, \ldots, k_0 - 1$ the distributions $E^{(i)}(x)$ and $\widetilde{E}^{(i)}(x)$ are uniformly continuous in \mathcal{L}'_0 . Then for every $\epsilon > 0$ there exists a Lyapunov ϵ -regularity function A_{ϵ} satisfying (1.1) and (1.2) such that the following are satisfied:

(a) (Exponential Tails) There exist constants $C = C(\epsilon) > 0$, $c = c(\epsilon) > 0$ and $p_0 = p_0(\epsilon) > 0$ such that

$$\mathsf{m}\left(M\setminus \bigcup_{k=0}^{n-1}f^{-k}(T_{p_0})\right) \le C e^{-cn}$$

for every integer $n \geq 0$.

(b) (Strong Exponential Tails) For every $\delta \in (0,1)$ there exist constants $p_0 = p_0(\epsilon, \delta) > 0$, $C = C(\epsilon, \delta) > 0$ and $c = c(\epsilon, \delta) > 0$ such that

$$\mathsf{m}\left(\Gamma_n(\epsilon,\delta,p_0)\right) \le C \, e^{-cn}$$

for every integer $n \ge 0$.

Clearly, taking the constant p_0 sufficiently large, we can make $m(T_{p_0})$ arbitrarily close to 1. The assumption that the flow is contact is used essentially in Sect. 6 below.

As we mentioned earlier, for every $p \ge 0$ there exists a constant $r_0 > 0$ (depending on ϵ and p) such that $r_{\epsilon}(x) \ge r_0$ for every $x \in T_p$. That is, for every $x \in T_p$ there exists a Lyapunov chart on $B(x, r_0)$.

The distributions $E^{(i)}(x)$ and $\tilde{E}^{(i)}(x)$ are uniformly continuous (in fact, Hölder continuous) for example when the flow has only three Lyapunov exponents $\chi_1 < \chi_2 = 0 < \chi_3$, or more generally, when for every $i = 1, \ldots, k_0 - 1$ there exist constants λ_i , μ_i and C > 0 such that $\chi_i < \lambda_i < \mu_i < \chi_{i+1}, ||d\phi_t(u)|| \le C \lambda_i^t ||u||$ for all $u \in E^{(i)}(x)$ and $t \ge 0$, and $||d\phi_t(u)|| \ge \frac{1}{C} \mu_i^t ||u||$ for all $u \in \tilde{E}^{(i)}(x)$ and $t \ge 0$ (see e.g. Ch. 3 in [P3]). Thus, the conclusions in Theorem 1.1 always hold for contact Anosov flows on 3-dimensional manifolds. As mentioned above, a more general immediate consequence of Theorem 1.1 is the following.

Corollary 1.2. Let $\phi_t : M \longrightarrow M$ be a C^2 contact Anosov flow on a C^2 compact Riemann manifold M, let Φ be a Hölder continuous real-valued function on M and let \mathfrak{m} be the Gibbs measure generated by Φ on M. Assume that k(x) = 3 for almost all $x \in \mathcal{L}_0$. Then for every $\epsilon > 0$ there exists a Lyapunov ϵ -regularity function A_{ϵ} satisfying (1.1) and (1.2) such that both (a) and (b) in Theorem 1.1 are satisfied.

In this paper we restrict ourselves to contact Anosov flows, however using slight modifications of the arguments in Sects. 3-5 below, results similar to Theorem 1.1 above can be proved for contact Anosov diffeomorphisms¹.

The motivation for the above result came from investigations on spectral properties of the so called Ruelle transfer operators. Attempts to obtain exponentially small estimates of integrals involving iterations of a certain kind of 'contraction operators' (see [D], [St]) naturally lead to estimates of 'tails' of the kind considered in Theorem 1.1. On the other hand, studies on decay of correlations² using the so called Young towers ([Y2]) also involve assumptions on the measures of the tails, and usually exponential tails are associated with exponential decay of correlations. Such assumptions appear naturally also in studies on large deviations (see e.g. [MN] and the references there).

Sect. 2 below contains some basic definitions. In Sect. 3 we introduce a 'regularity function' $R_{\epsilon}(x)$ (and $\tilde{R}_{\epsilon}(x)$ related to it), changing slightly the approach of Simic in [Sim]. Although this is most likely not a slowly varying function, it turns out that const $(\sqrt{\epsilon}A_{2\epsilon}(x))^{1/3} \leq R_{\epsilon}(x) \leq A_{\epsilon}(x)$ for some Lyapunov ϵ -regularity function $A_{\epsilon}(x)$ satisfying (1.1) and (1.2). This is proved in Sect. 6. Consequently, it is enough to prove the analogue of Theorem 1.1 replacing $A_{\epsilon}(x)$ by $R_{\epsilon}(x)$. We study the latter (or rather, the related one $\tilde{R}_{\epsilon}(x)$) in Sects. 4 and 5 using an idea³ from [Sim] and the classical large deviation principle (see [OP], [Ki] or [Y1]). Theorem 1.1 is proved in Sect. 4 as a consequence of a similar result about the function $\tilde{R}_{\epsilon}(x)$.

Acknowledgement. Thanks are due to Sebastian Gouëzel who pointed out to an error in a previous version of the paper.

2 Preliminaries

Throughout this paper M denotes a C^2 compact Riemann manifold, and $\phi_t : M \longrightarrow M$ $(t \in \mathbb{R})$ a C^2 Anosov flow on M. That is, for some constants C > 0 and $0 < \lambda < 1$ there exists a $d\phi_t$ -invariant decomposition $T_x M = E^0(x) \oplus E^u(x) \oplus E^s(x)$ of $T_x M$ $(x \in M)$ into a direct sum of non-zero linear subspaces, where $E^0(x)$ is the one-dimensional subspace determined by the direction of the flow at x, $||d\phi_t(u)|| \leq C \lambda^t ||u||$ for all $u \in E^s(x)$ and $t \geq 0$, and $||d\phi_t(u)|| \leq C \lambda^{-t} ||u||$ for all $u \in E^u(x)$ and $t \leq 0$. For $x \in M$ and a sufficiently small $\epsilon > 0$ let

$$\begin{split} W^s_{\epsilon}(x) &= \{ y \in M : d(\phi_t(x), \phi_t(y)) \le \epsilon \text{ for all } t \ge 0 \ , \ d(\phi_t(x), \phi_t(y)) \to_{t \to \infty} 0 \ \} \ , \\ W^u_{\epsilon}(x) &= \{ y \in M : d(\phi_t(x), \phi_t(y)) \le \epsilon \text{ for all } t \le 0 \ , \ d(\phi_t(x), \phi_t(y)) \to_{t \to -\infty} 0 \ \} \end{split}$$

¹That case is in fact easier.

²See [L], [BaL] and the references there for general information on this topic.

 $^{^{3}}$ We use the idea briefly mentioned at the end of Sect. 1 in [Sim] apparently suggested by the referee of that paper.

be the (strong) stable and unstable manifolds of size ϵ . Then $E^u(x) = T_x W^u_{\epsilon}(x)$ and $E^s(x) = T_x W^s_{\epsilon}(x)$.

The flow ϕ_t is called *contact* if dim(M) = 2n + 1 for some $n \ge 1$ and there exists a C^2 flowinvariant one-form ω on M such that $\omega \land (d\omega)^n \ne 0$ on M. It is well-known that the Lyapunov spectrum of a contact flow is symmetric, i.e. for each $i = 1, \ldots, k_0$ there exists $j = 1, \ldots, k_0$ with $\chi_j = -\chi_i$.

It follows from the hyperbolicity of the flow on M that if $\epsilon_0 > 0$ is sufficiently small, there exists $\epsilon > 0$ such that if $x, y \in M$ and $d(x, y) < \epsilon$, then $W^s_{\epsilon_0}(x)$ and $\phi_{[-\epsilon_0,\epsilon_0]}(W^u_{\epsilon_0}(y))$ intersect at exactly one point [x, y] (cf. [KH]). That is, there exists a unique $t \in [-\epsilon_0, \epsilon_0]$ such that $\phi_t([x, y]) \in W^u_{\epsilon_0}(y)$.

Let $\mathcal{R} = \{R_i\}_{i=1}^{k_0}$ be a Markov family consisting of rectangles R_i , each contained in a submanifold D_i of M of codimension one (see [B]). Assuming that $\epsilon > 0$ is sufficiently small, the projection $\operatorname{pr}_{D_i} : \phi_{[-\epsilon,\epsilon]}(D_i) \longrightarrow D_i$ along the flow is well-defined and smooth. Given $x, y \in D_i$, set $\langle x, y \rangle_{D_i} = \operatorname{pr}_{D_i}([x, y])$. A subset R_i of D_i is called a *rectangle* if $\langle x, y \rangle_{D_i} \in R_i$ for all $x, y \in R_i$. For any $x \in R_i$ define the stable and unstable leaves through x in R_i by $W_{R_i}^s(x) = \operatorname{pr}_{D_i}(W_{\epsilon}^s(x) \cap \phi_{[-\epsilon,\epsilon]}(D_i)) \cap R_i$ and $W_{R_i}^u(x) = \operatorname{pr}_{D_i}(W_{\epsilon}^u(x) \cap \phi_{[-\epsilon,\epsilon]}(D_i)) \cap R_i$. We may assume that each R_i has the form $R_i = \langle U_i, S_i \rangle_{D_i} = \{\langle x, y \rangle_{D_i} : x \in U_i, y \in S_i\}$, where $U_i \subset W_{\epsilon}^u(z_i)$ and $S_i \subset W_{\epsilon}^s(z_i)$, respectively, for some $z_i \in M$. Set $R = \bigcup_{i=1}^{k_0} R_i$. The corresponding *Poincaré map* $\mathcal{P} : R \longrightarrow R$ is defined by $\mathcal{P}(x) = \phi_{\tau(x)}(x) \in R$, where $\tau(x) > 0$ is the smallest positive time with $\phi_{\tau(x)}(x) \in R$. The function τ is the *first return time* associated with \mathcal{R} .

From now on we will assume that $\mathcal{R} = \{R_i\}_{i=1}^{k_0}$ is a fixed Markov family for ϕ_t of size $\chi < \epsilon_0/4 < 1/4$. Denote by \widehat{R} the core of R, i.e. the set $x \in R$ such that $\mathcal{P}^m(x) \in \operatorname{Int}(R) = \bigcup_{i=1}^k \operatorname{Int}(R_i)$ for all $m \in \mathbb{Z}$. It is well-known (see [B]) that \widehat{R} is a residual subset of R and has full measure with respect to any Gibbs measure on R. In general τ is not continuous on R, however it is Hölder on \widehat{R} when considered with respect to an appropriate metric d_{θ} , see below. The same applies to $\mathcal{P} : \widehat{R} \longrightarrow \widehat{R}$. It is well-known ([BR]) that the Markov family \mathcal{R} can be chosen so that τ is non-lattice. From now on we will assume that \mathcal{R} is chosen in this way.

Let $\mathcal{A} = (\mathcal{A}_{ij})_{i,j=1}^k$ be the matrix given by $\mathcal{A}_{ij} = 1$ if $\mathcal{P}(\operatorname{Int}(R_i)) \cap \operatorname{Int}(R_j) \neq \emptyset$ and $\mathcal{A}_{ij} = 0$ otherwise. Consider the symbol space

$$\Sigma_A = \{ (i_j)_{j=-\infty}^{\infty} : 1 \le i_j \le k_0, A_{i_j \ i_{j+1}} = 1 \text{ for all } j \},\$$

with the product topology and the shift map $\sigma : \Sigma_A \longrightarrow \Sigma_A$ given by $\sigma((i_j)) = ((i'_j))$, where $i'_j = i_{j+1}$ for all j. Given $0 < \theta < 1$, consider the metric d_θ on Σ_A defined by $d_\theta(\xi, \eta) = 0$ if $\xi = \eta$ and $d_\theta(\xi, \eta) = \theta^m$ if $\xi_i = \eta_i$ for |i| < m and m is maximal with this property. There is a natural map $\mathcal{W} : \Sigma_A \longrightarrow R$ such that $\mathcal{W} \circ \sigma = \mathcal{P} \circ \mathcal{W}$. In general \mathcal{W} is not one-to-one, however it is a bijection between $\mathcal{W}^{-1}(\widehat{R})$ and \widehat{R} . Choosing $\theta \in (0, 1)$ appropriately, the map $\mathcal{W} : \Sigma_A \longrightarrow R$ is Lipschitz when Σ_A is endowed with d_θ and R with the metric induced by the Riemann metric on M. In what follows we assume $\theta \in (0, 1)$ is a fixed constant with this property, and we will consider \widehat{R} with the metric induced by d_θ via \mathcal{W} .

3 Lyapunov regularity functions

Throughout we assume that ϕ_t is a C^2 contact Anosov flow on M. Let $\Phi: M \longrightarrow \mathbb{R}$ be a fixed Hölder continuous function on M and let m be the *Gibbs measure* determined by Φ . Let \mathcal{L}'_0 be a subset of \mathcal{L}_0 of full m -measure such that $k(x) = k_0$ and $\chi_i(x)$ are constant for $x \in \mathcal{L}'_0$ for all $i = 1, \ldots, k_0$. Removing a set of measure zero, we may assume that $\mathcal{L}'_0 \subset \widehat{R}$. Take $\epsilon > 0$ so small that $\chi_j \notin [\chi_i - 4\epsilon, \chi_i + 4\epsilon]$ whenever $i \neq j$, and set

$$R_{\epsilon}^{+}(x) = \max_{1 \le i \le k_0} \sup_{n \ge 0} \frac{\|(df_x^n)_{|E_i}\|}{e^{(\chi_i + \epsilon)n}},$$
(3.1)

$$R_{\epsilon}^{-}(x) = \max_{1 \le i \le k_0} \sup_{n \ge 0} \frac{\|(df_x^{-n})_{|E_i}\|}{e^{(\chi_i - \epsilon)(-n)}},$$
(3.2)

and $R_{\epsilon}(x) = \max\{R_{\epsilon}^{-}(x), R_{\epsilon}^{+}(x)\}.$

Note. Since $\frac{1}{n}\log \|(df_x^n)_{|E_i}\| \to \chi_i$ as $n \to \infty$, we have $\frac{1}{n}\log \|(df_x^n)_{|E_i}\| < \chi_i + \epsilon$ for large n, so (3.1) exists. Similarly, $\frac{1}{-n}\log \|(df_x^{-n})_{|E_i}\| \to \chi_i$ as $n \to \infty$, so $\frac{1}{-n}\log \|(df_x^{-n})_{|E_i}\| > \chi_i - \epsilon$ for large n. This gives $\log \|(df_x^{-n})_{|E_i}\| < (\chi_i - \epsilon)(-n)$ for large n, and so (3.2) exists.

From the above definitions it is clear that

$$\frac{\|df_x^n(v)\|}{e^{n\chi_i}} \le R_\epsilon(x) e^{|n|\epsilon} \|v\| \quad , \quad v \in E_i(x) \ , \ n \in \mathbb{Z},$$

$$(3.3)$$

for all $x \in \mathcal{L}'_0$ and all $i = 1, ..., k_0$, and moreover for every Lyapunov ϵ -regularity function $A_{\epsilon}(x)$ with (1.1) we have $R_{\epsilon}(x) \leq A_{\epsilon}(x)$ for all $x \in \mathcal{L}'_0$ and $\epsilon > 0$. It is not clear whether $R_{\epsilon}(x)$ is a slowly varying function, i.e. whether it satisfies (1.2), possibly with ϵ replaced by some $\delta > 0$ related to ϵ . However we have the following.

Proposition 3.1. Let $\phi_t : M \longrightarrow M$ be a contact C^2 Anosov flow, and let Φ , \mathfrak{m} and \mathcal{L}'_0 be as above. For every $\epsilon > 0$ there exists an ϵ -slowly varying function A_{ϵ} satisfying (1.1) and (1.2) such that

$$\frac{\epsilon^{1/6}}{C} (A_{2\epsilon}(x))^{1/3} \le R_{\epsilon}(x) \le A_{\epsilon}(x)$$
(3.4)

for almost all $x \in \mathcal{L}'_0$, where $C = C_0^{2/3} (2k_0)^{1/6} > 1$.

Notice that $\frac{\epsilon^{1/6}}{C}(A_{2\epsilon}(x))^{1/3}$ is also an ϵ -slowly varying function.

We prove this proposition in Sect. 6 below. The assumption that the flow is contact is used in an essential way there.

We will also need the functions

$$\widetilde{R}_{\epsilon}^{+}(x) = \max_{1 \le i \le k_{0}} \sup_{n \ge 0} \frac{\|(df_{x}^{n})_{|E^{(i)}}\|}{e^{(\chi_{i} + \epsilon)n}} \quad , \quad \widetilde{R}_{\epsilon}^{-}(x) = \max_{1 \le i \le k_{0}} \sup_{n \ge 0} \frac{\|(df_{x}^{-n})_{|\widetilde{E}^{(i)}}\|}{e^{(\chi_{i} - \epsilon)(-n)}},$$

and $\widetilde{R}_{\epsilon}(x) = \max\{\widetilde{R}_{\epsilon}^{-}(x), \widetilde{R}_{\epsilon}^{+}(x)\}$. Clearly $R_{\epsilon}(x) \leq \widetilde{R}_{\epsilon}(x)$. In Sect. 6 below we will show that there exists a constant $K(\epsilon) \geq 1$ such that

$$\widetilde{R}_{\epsilon}(x) \le K(\epsilon) \left(R_{\epsilon}(x)\right)^2 \tag{3.5}$$

for all $x \in \mathcal{L}'_0$.

Remark 3.2. Proposition 3.1 and (3.5) show that sets of the form $\{x \in \mathcal{L}'_0 : A_{\epsilon}(x) \leq e^p\}$ are easily related to sets of the form $\{x \in \mathcal{L}'_0 : \widetilde{R}_{\epsilon'}(x) \leq e^{p'}\}$ (with $\epsilon' = \epsilon$ or 3ϵ), so it is enough to prove an analogue of Theorem 1.1 replacing the function A_{ϵ} by $C\widetilde{R}_{\epsilon}$ for some constant C > 0.

4 Reductions

Let ϕ_t be a C^2 Anosov flow on M, and let Φ , m and $\mathcal{L}'_0 \subset \mathcal{L}_0$ be as in Sect. 3. In this section and also the next one we do not assume that the flow is contact. However, as in Theorem 1.1, we will assume that the bundles $E^{(i)}(x)$ and $\tilde{E}^{(i)}(x)$ are uniformly continuous on \mathcal{L}'_0 .

Let $\mathcal{R} = \{R_i\}_{i=1}^{k_0}$ be a Markov family for ϕ_t as in Sect. 2, and let $\tau : R = \bigcup_{i=1}^{k_0} R_i \longrightarrow [0, 1/4]$ and $\mathcal{P} : R \longrightarrow R$ be the corresponding first return map and the Poincaré map. Fix constants $0 < \tau_0 < \hat{\tau}_0 < 1/4$ so that $\tau_0 \leq \tau(x) \leq \hat{\tau}_0$ for all $x \in R$. There is a well-defined projection $\pi : M \longrightarrow R$ defined by $\pi(y) = x$, where $x \in R, y = \phi_s(x)$ for some $s \in [0, \tau(x))$ and $s \geq 0$ is the smallest number with this property.

The Gibbs measure \mathbf{m} induces a *Gibbs measure* μ on R (with respect to the Poincaré map \mathcal{P}) for the function

$$F(x) = \int_0^{\tau(x)} \Phi(\phi_s(x)) \, ds \quad , \quad x \in R.$$

The function F is Hölder on \widehat{R} (with respect to the metric d_{θ}). For every continuous function H on M we have (see e.g. [PP])

$$\int_{M} H \, d\mathbf{m} = \frac{\int_{R} \left(\int_{0}^{\tau(x)} H(\phi_s(x)) \, ds \right) d\mu(x)}{\int_{R} \tau \, d\mu}.\tag{4.1}$$

By Birkhoff's Theorem,

$$\lim_{n \to \infty} \frac{\tau_n(x)}{n} = \tau^{\dagger} = \int_R \tau \, d\mu \tag{4.2}$$

for μ -almost all $x \in \mathcal{L}'_0 \cap R$. Here $\tau_n(x) = \tau(x) + \tau(\mathcal{P}(x)) + \ldots + \tau(\mathcal{P}^{n-1}(x))$. By the choice of τ we have $\tau^{\dagger} \leq 1/4$.

Fix a constant $C_1 \ge 1$ so that

$$C_1 \ge \frac{k_0}{\tau^{\dagger}} \sup\{ |\log \| d\phi_t(x) \| | : x \in M , \ 0 \le t \le 1 \},\$$

and $C_1 \geq 2 \max_{1 \leq i \leq k_0} |\chi_i|$. Then fix arbitrary constants

$$0 < \epsilon \le \frac{1}{8} \min\{|\chi_i| : 1 \le i \le k_0, \chi_i \ne 0\} \quad , \quad \delta \in (0,1) \quad , \quad C_0 \ge \frac{100C_1}{\epsilon \tau_0}, \tag{4.3}$$

and set

$$\delta_0 = \frac{\delta}{2k_0}.\tag{4.4}$$

We will study in details the regularity function $\widetilde{R}^+_{\epsilon}(x)$, In a similar way one can deal with $\widetilde{R}^-_{\epsilon}(x)$; one just needs to replace ϕ_t by ϕ_{-t} . Clearly $\widetilde{R}_{\epsilon}(x) = \max_{1 \le i \le k_0} \widetilde{R}^+_{\epsilon,i}(x)$, where

$$\widetilde{R}^+_{\epsilon,i}(x) = \sup_{n \ge 0} \frac{\|(df^n_x)_{|E^{(i)}}\|}{e^{(\chi_i + \epsilon)n}}.$$

In this section, and the next section as well, we will restrict our attention to an arbitrary fixed $i = 1, ..., k_0$, and the corresponding bundle $E = E^{(i)}$. Set $\chi = \chi_i$ and $B_{\epsilon}(x) = \tilde{R}^+_{\epsilon,i}(x)$ for brevity. Thus,

$$B_{\epsilon}(x) = \sup_{n \in \mathbb{N}} \frac{\|d\phi_n(x)|_E\|}{e^{(\chi+\epsilon)n}} \quad , \quad x \in \mathcal{L}'_0, \tag{4.5}$$

where N is the set of non-negative integers. We will compare $B_{\epsilon}(x)$ with the function

$$C_{\epsilon}(x) = \sup_{k>0} \frac{\|d\phi_{\tau_k(x)}(x)|_E\|}{e^{(\chi+\epsilon)\tau_k(x)}} \quad , \quad x \in \mathcal{L}'_0 \cap R.$$

Lemma 4.1. (a) There exists a constant $S_1 = S_1(\epsilon) > 0$ such that

$$B_{\epsilon}(y) \le S_1 C_{\epsilon}(x) \tag{4.6}$$

for all $y \in \mathcal{L}'_0$, $x = \pi(y)$. Moreover we can take $S_1 > 0$ so that (4.6) holds for any $y \in \mathcal{L}$ and $x \in R$ with $y = \phi_t(x)$ for some $t \in [-2, 2]$.

(b) For any integer $s_0 \ge 1$ there exists a constant $S_2 = S_2(\epsilon, s_0) \ge 1$ such that $C_{\epsilon}(\mathcal{P}^{-j}(x)) \le 1$ $S_2 C_{\epsilon}(x)$ for any $x \in \mathcal{L}'_0 \cap R$ and any $j = 0, 1, \ldots, s_0$.

Proof. Set $a_1 = 2(|\chi| + \epsilon)$ and $a_2 = \sup_{|s| \le 2} ||d\phi_s|| < \infty$. (a) Let $y \in \mathcal{L}'_0$; set $x = \pi(y) \in \mathcal{L}'_0 \cap R$. Then $y = \phi_s(x)$ for some $s \in [0, \tau(x))$. We have $B_{\epsilon}(y) = \frac{||d\phi_t(x)|_E||}{e^{(\chi+\epsilon)t}}$ for some integer $t \ge 0$. Denote by $k \ge 0$ the maximal integer so that $\tau_k(x) \le t + s$; then $t + s < \tau_{k+1}(x)$. Thus, $t \ge \tau_k(x) - s \ge \tau_k(x) - \hat{\tau}_0$. Also,

$$\begin{aligned} \|d\phi_t(y)|_E\| &= \|d\phi_{-s+(t+s)}(\phi_s(x))|_E\| \le \|d\phi_{-s}(\phi_s(x))|_E\| \cdot \|d\phi_{t+s}(x)|_E\| \\ &\le a_2 \|d\phi_{\tau_k(x)+(t+s-\tau_k(x))}(x)|_E\| \le a_2^2 \|d\phi_{\tau_k(x)}(x)|_E\|. \end{aligned}$$

Thus,

$$B_{\epsilon}(y) = \frac{\|d\phi_t(y)_{|E}\|}{e^{(\chi+\epsilon)t}} \le \frac{a_2^2 \|d\phi_{\tau_k(x)}(x)_{|E}\|}{e^{(\chi+\epsilon)(\tau_k(x)-\hat{\tau}_0)}} \le e^{a_1} a_2^2 \frac{\|d\phi_{\tau_k(x)}(x)_{|E}\|}{e^{(\chi+\epsilon)\tau_k(x)}}.$$

This shows that $B_{\epsilon}(y) \leq \text{Const } C_{\epsilon}(x)$. In a similar way one proves that $C_{\epsilon}(x) \leq \text{Const } B_{\epsilon}(y)$.

The more general case when $y = \phi_t(x)$ for some $t \in [-2, 2]$ follows similarly.

(b) Take $S_2 = \max_{0 \le j \le s_0} \sup_{z \in R} \frac{\|d\phi_{\tau_j(z)}(z)\|}{e^{(\chi+\epsilon)\tau_j(z)}}$. Given $x \in \mathcal{L}'_0 \cap R$ and $j = 0, 1, \dots, s_0$, set $y = \mathcal{P}^{-j}(x)$. For any integer $n \in [0, s_0]$ clearly $\frac{\|d\phi_{\tau_n(y)}(y)|_E\|}{e^{(\chi+\epsilon)\tau_n(y)}} \leq S_2 \leq S_2 C_{\epsilon}(x)$. Let $n > s_0$. Then n = j + k for some integer k > 0, and therefore

$$\frac{|d\phi_{\tau_n(y)}(y)_{|E}||}{e^{(\chi+\epsilon)\tau_n(y)}} = \frac{\|d\phi_{\tau_{j+k}(y)}(y)_{|E}\|}{e^{(\chi+\epsilon)\tau_{j+k}(y)}} = \frac{\|d\phi_{\tau_j(y)+\tau_k(x)}(y)_{|E}\|}{e^{(\chi+\epsilon)(\tau_j(y)+\tau_k(x))}} \\ \leq \frac{\|d\phi_{\tau_j(y)}(y)_{|E}\|}{e^{(\chi+\epsilon)\tau_j(y)}} \cdot \frac{\|d\phi_{\tau_k(x)}(x)_{|E}\|}{e^{(\chi+\epsilon)\tau_k(x)}} \le S_2 C_{\epsilon}(x)$$

Thus, $C_{\epsilon}(y) \leq S_2 C_{\epsilon}(x)$.

Next, consider the functions

$$G^{(n)}(x) = \log \|d\phi_{\tau_n(x)}(x)|_E\|$$

for $x \in \mathcal{L}'_0 \cap R$ and any $n \ge 0$. Clearly, $|G^{(1)}(x)| \le \text{Const} < \infty$ for all $x \in \mathcal{L}'_0 \cap R$, and

$$\lim_{n \to \infty} \frac{G^{(n)}(x)}{\tau^{\dagger} n} = \lim_{n \to \infty} \frac{G^{(n)}(x)}{\tau_n(x)} \frac{\tau_n(x)}{\tau^{\dagger} n} = \chi$$
(4.7)

for μ -almost all $x \in R$. Here we used the fact that $\lim_{t\to\infty} \frac{\log \|d\phi_t(x)\|_E}{t} = \chi$ for μ -almost all $x \in R$.

The functions $G^{(n)}$ form a sub-additive sequence with respect to the map $\mathcal{P}: R \longrightarrow R$. Indeed, for any $n, m \ge 0$ and any $x \in \mathcal{L}'_0 \cap R$ we have

$$\begin{aligned} G^{(n+m)}(x) &= \log \|d\phi_{\tau_{n+m}(x)}(x)|_{E}\| = \log \|d\phi_{\tau_{n}(x)+\tau_{m}(\mathcal{P}^{n}(x))}(x)|_{E}\| \\ &\leq \log \|d\phi_{\tau_{n}(x)}(x)|_{E}\| + \log \|d\phi_{\tau_{m}(\mathcal{P}^{n}(x))}(\mathcal{P}^{n}(x))|_{E}\| = G^{(n)}(x) + G^{(m)}(\mathcal{P}^{n}(x)). \end{aligned}$$

By Kingsman's Ergodic Theorem ([Kin]) there exists the limit

$$L = \lim_{n \to \infty} \frac{1}{n} \int_{R} G^{(n)}(x) \, d\mu(x) = \inf_{n} \frac{1}{n} \int_{R} G^{(n)}(x) \, d\mu(x), \tag{4.8}$$

and moreover $\lim_{n\to\infty} \frac{G^{(n)}(x)}{n} = L$ for μ -almost all $x \in \mathcal{L}'_0 \cap R$. Now (4.7) shows that $L = \tau^{\dagger} \chi$. **Fix a subset** \mathcal{L} of $\mathcal{L}'_0 \cap R$ with $\mu(\mathcal{L}) = 1$ such that (4.2) and (4.7) hold for all $x \in \mathcal{L}$. Set $\widetilde{\mathcal{L}} = \pi^{-1}(\mathcal{L}) \subset \mathcal{L}'_0$; then $\widetilde{\mathcal{L}} \cap R = \mathcal{L}$.

By Egorov's Theorem, there exists a compact subset K_0 of \mathcal{L} such that

$$\mu(K_0) > 1 - \frac{\delta_0}{C_0},\tag{4.9}$$

and $\frac{\tau_n(x)}{\tau^{\dagger}n} \to 1$ and $\frac{G^{(n)}(x)}{\tau^{\dagger}n} \to \chi$ uniformly on K_0 . Thus, there exists an integer $n_0 \ge 1$ such that

$$\left|\frac{\tau_n(x)}{\tau^{\dagger}n} - 1\right| < \frac{\delta_0}{C_0} \quad , \quad x \in K_0 \; , \tag{4.10}$$

$$\left|\frac{G^{(n)}(x)}{\tau^{\dagger}n} - \chi\right| < \frac{\delta_0}{C_0} \quad , \quad x \in K_0 \; , \tag{4.11}$$

and, using (4.8),

$$\chi \le \frac{1}{\tau^{\dagger} n} \int_{R} G^{(n)}(x) \, d\mu(x) < \chi + \frac{\delta_{0}}{C_{0}} \tag{4.12}$$

for all integers $n \geq n_0$.

Fix an arbitrary integer $s_0 = s_0(\epsilon, \delta_0) \ge n_0$; this will stay fixed throughout the whole Sects. 4 and 5. Then (4.10), (4.11) and (4.12) hold with n replaced by s_0 . Consider the transformation

 $T = \mathcal{P}^{s_0} : R \longrightarrow R$

preserving the measure μ , and the measurable functions

$$u(x) = \frac{G^{(s_0)}(x)}{\tau^{\dagger} s_0} = \frac{1}{\tau^{\dagger} s_0} \log \| d\phi_{\tau_{s_0}(x)}(x)_{|E} \| \quad , \quad x \in \mathcal{L},$$

and

$$g(x) = \frac{\tau_{s_0}(x)}{\tau^{\dagger} s_0} \quad , \quad x \in \mathcal{L}$$

Clearly u and g are bounded, $||u||_{\infty} \leq C_1$ and $\tau_0/\tau^{\dagger} \leq g(x) \leq \hat{\tau}_0/\tau^{\dagger}$ for all $x \in R$. For any integer $n \geq 1$ and $x \in \mathcal{L}$ set

$$u^{n}(x) = u(x) + u(T(x)) + \ldots + u(T^{n-1}(x))$$

(Notice that a superscript is used here when we deal with orbits with respect to $T = \mathcal{P}^{s_0}$; unlike the subscript used for orbits with respect to \mathcal{P} .)

It follows from our assumptions that u is uniformly continuous on \mathcal{L}'_0 . For the integral of u, (4.12) implies

$$\chi \le \int_R u \, d\mu < \chi + \frac{\delta_0}{C_0}.\tag{4.13}$$

Lemma 4.2. There exist a Lipschitz⁴ function $\psi : R \longrightarrow \mathbb{R}$ with respect to the metric d_{θ} such

⁴Hölder continuity is enough.

that $\|\psi\|_{\infty} \leq 2C_1$ and $u(x) \leq \psi(x) \leq u(x) + \delta_0/C_0$ for μ -almost all $x \in R$.

Proof. Since u is uniformly continuous on \mathcal{L}'_0 (with respect to the metric d_{θ}), it has a continuous extension to the whole of R. We will denote this extension by u again, and we still have $||u||_{\infty} \leq C_1$.

Take $a = \frac{\delta_0}{2C_0}$ and consider the function u + a on R. On any given rectangle R_i this function is continuous both with respect to the metric d_{θ} and the Riemann metric. Using the manifold structure of the disk D_i containing R_i and a standard regularization procedure, we find a Lipschitz (with respect to the Riemann metric) function ψ on R_i such that $\|\psi - (u+a)\|_{\infty} < a$ on R_i . Then ψ is Lipschitz on R with respect to d_{θ} and $\psi \geq u$. Moreover, $\|\psi - (u+a)\|_{\infty} < a$ and $\|u\|_{\infty} \leq C_1$ imply $\|\psi\|_{\infty} \leq 2C_1$.

Fix a function ψ with the properties in Lemma 4.2. It follows from (4.13) that

$$\chi \le \tilde{\chi} = \int_R \psi \, d\mu < \chi + \frac{2\delta_0}{C_0}.\tag{4.14}$$

Changing slightly ψ if necessary, we may assume that ψ is not cohomologous to a constant. Next, set

$$D_{\epsilon}(x) = \sup_{m>0} \frac{e^{\psi^m(x)}}{e^{(\tilde{\chi}+\epsilon)g^m(x)}}.$$
(4.15)

By Birkhoff's Theorem, this is well-defined for μ -almost all $x \in \mathcal{L}$. Removing a set of measure zero from \mathcal{L} if necessary (and thus shrinking the compact set K_0 a bit), we will assume that $D_{\epsilon}(x)$ is defined for all $x \in \mathcal{L}$.

Lemma 4.3. There exists a constant $S_3 = S_3(\epsilon, \delta_0) > 0$ such that $C_{2\epsilon}(x) \leq S_3 (D_{\epsilon}(x))^{\tau^{\dagger} s_0}$ for all $x \in \mathcal{L}$.

Proof. Set $a_3 = \sup_{0 \le t \le s_0 \hat{\tau}_0} \log \|d\phi_t(x)\|$. Then $G^{(\ell)}(x) \le a_3$ for all $0 \le \ell \le s_0$ and all $x \in \mathcal{L}$. Given $x \in \mathcal{L}$, there exists $k \in \mathbb{N}$ such that $C_{2\epsilon}(x) = \frac{\|d\phi_{\tau_k(x)}(x)|_E\|}{e^{(\chi+2\epsilon)\tau_k(x)}}$. Let $k = ms_0 + \ell, 0 \le \ell < s_0$. Then, setting $y = \mathcal{P}^{ms_0}(x)$ and using $u \leq \psi$, we get

$$\log \|d\phi_{\tau_k(x)}(x)|_E\| = G^{(ms_0+\ell)}(x) \le G^{(s_0)}(x) + G^{(s_0)}(\mathcal{P}^{s_0}(x)) + \ldots + G^{(s_0)}(\mathcal{P}^{(m-1)s_0}(x)) + G^{(\ell)}(y)$$

$$\le \tau^{\dagger} s_0(u^m(x) + a_3) \le \tau^{\dagger} s_0(\psi^m(x) + a_3).$$

Since $\tau_k(x) \ge \tau_{ms_0}(x)$ and $\tilde{\chi} < \chi + \epsilon$ by (4.14) and the choice of the constants, it follows that

$$\frac{\|d\phi_{\tau_k(x)}(x)_{|E}\|}{e^{(\chi+2\epsilon)\tau_k(x)}} \leq \frac{(e^{a_3} e^{\psi^m(x)})^{\tau^{\dagger}s_0}}{e^{(\chi+2\epsilon)\tau_{ms_0}(x)}} \leq S_3 \left(\frac{e^{\psi^m(x)}}{e^{(\tilde{\chi}+\epsilon)\frac{\tau_{ms_0}(x)}{\tau^{\dagger}s_0}}}\right)^{\tau^{\dagger}s_0} \leq S_3 \left(D_{\epsilon}(x)\right)^{\tau^{\dagger}s_0},$$

since $\frac{\tau_{ms_0}(x)}{s_0\tau^{\dagger}} = g^m(x)$. This is true for all $k \ge 0$, so $C_{2\epsilon}(x) \le S_3 (D_{\epsilon}(x))^{\tau^{\dagger}s_0}$.

For any $p \in \mathbb{R}$ set

$$Q_p(\epsilon) = \{ x \in \mathcal{L} \cap R : D_{\epsilon}(x) \le e^p \},\$$

and, given $\hat{\delta} \in (0, 1]$, denote by $\Xi_m(\epsilon, \hat{\delta})$ the set of those $x \in \mathcal{L}$ such that

$$\sharp \left\{ k : 0 \le k < m , \, T^k(x) \notin Q_0(\epsilon) \right\} \ge \hat{\delta} \, m.$$

The central result in this paper is the following.

Theorem 4.4. (Strong Exponential Tails) Let $\epsilon > 0$, $\delta \in (0,1)$ and δ_0 be as in the beginning of Sect. 4, let the integer $s_0 = s_0(\epsilon, \delta_0) \ge 1$ be chosen as above, and let $T = \mathcal{P}^{s_0}$. For every $\hat{\delta} \in [\delta_0, 1]$ there exist constants $C_2 = C_2(\epsilon, \hat{\delta}) > 0$ and $c_2 = c_2(\epsilon, \hat{\delta}) > 0$ such that

$$\mu\left(\Xi_m(\epsilon,\hat{\delta})\right) \le C_2 e^{-c_2 m}.$$
(4.16)

In particular,

$$\mu\left(R\setminus \bigcup_{k=0}^{m-1}T^{-k}(Q_0(\epsilon))\right) \le C_2 e^{-c_2 m}$$

for every integer $m \geq 1$.

We prove this theorem in the next section. We will now derive Theorem 1.1 from it.

Proof of Theorem 1.1. Recall that $\delta_0 = \frac{\delta}{2k_0}$ by (4.4). Take $\hat{\delta} = \delta_0$. Let $p_i \in \mathbb{R}$ be so that $S_i = e^{p_i}$ (i = 1, 2, 3), where S_1 , S_2 and S_3 are the constants from Lemmas 4.1 and 4.3.

Given an integer $m \ge 0$, set $n = ms_0$ and let Y_n be the set of those $x \in \mathcal{L}$ such that

 $\sharp\{j: 0 \le j < n , C_{2\epsilon}(\mathcal{P}^j(x)) > e^{p_2 + p_3}\} \ge \delta_0 n.$

Set $\Xi_m = \Xi_m(\epsilon, \delta_0)$ for brevity. We will now prove that

$$Y_n \subset \Xi_m. \tag{4.17}$$

Let $x \in Y_n$. Then $C_{2\epsilon}(\mathcal{P}^j(x)) \leq e^{p_2+p_3}$ for less that $(1-\delta_0)n$ values of $j=0,1,\ldots,n-1$. By Lemma 4.3, $C_{2\epsilon}(\mathcal{P}^j(x)) \leq S_3(D_{\epsilon}(\mathcal{P}^j(x)))^{\tau^{\dagger}s_0} = e^{p_3}(D_{\epsilon}(\mathcal{P}^j(x)))^{\tau^{\dagger}s_0}$, so $D_{\epsilon}(\mathcal{P}^j(x)) \leq 1$ implies $C_{2\epsilon}(\mathcal{P}^j(x)) \leq e^{p_3}$. Assume for a moment that $x \notin \Xi_m$. Then $T^k(x) \in Q_0(\epsilon)$ for at least $(1-\delta_0)m$ values of $k = 0, 1, \ldots, m-1$. In other words $D_{\epsilon}(\mathcal{P}^{ks_0}(x)) \leq 1$ for at least $(1-\delta_0)m$ values of k, and so $C_{2\epsilon}(\mathcal{P}^{ks_0}(x)) \leq e^{p_3}$ for at least $(1-\delta_0)m$ values of $k = 0, 1, \ldots, m-1$. For any such k, Lemma 4.1(b) implies that $C_{2\epsilon}(\mathcal{P}^{ks_0-j}(x)) \leq S_2C_{2\epsilon}(\mathcal{P}^{ks_0}(x)) \leq e^{p_2+p_3}$ for all $j = 0, 1, \ldots, s_0 - 1$. Hence $C_{2\epsilon}(\mathcal{P}^r(x)) \leq e^{p_2+p_3}$ for at least $s_0(1-\delta_0)m = (1-\delta_0)n$ values of $r = 0, 1, \ldots, n-1$. In other words, $C_{2\epsilon}(\mathcal{P}^r(x)) > e^{p_2+p_3}$ for less than $\delta_0 n$ values of $r = 0, 1, \ldots, n-1$, which is a contradiction with $x \in Y_n$.

This proves (4.17). Combining the latter with (4.16) gives

$$\mu(Y_n) \le C_2 e^{-c_2 n/s_0}. \tag{4.18}$$

Setting $\widetilde{Y}_n = \pi^{-1}(Y_n) = \{ y \in \widetilde{\mathcal{L}} : \pi(y) \in Y_n \}$, it follows from (4.1) that

$$\mathsf{m}(\widetilde{Y}_{n}) = \frac{1}{\tau^{\dagger}} \int_{R} \int_{0}^{\tau(y)} \chi_{\widetilde{Y}_{n}}(\phi_{s}(y)) \, ds \, d\mu = \frac{1}{\tau^{\dagger}} \int_{R} \tau(y) \chi_{Y_{n}}(y) \, d\mu \le \frac{\hat{\tau}_{0}}{\tau^{\dagger}} \, \mu(Y_{n}). \tag{4.19}$$

Next, set $k = [n/\tau_0]$, and let X_k be the set of those $y \in \widetilde{\mathcal{L}}$ such that

 $\sharp \{r: 0 \le r < k \;,\; B_{2\epsilon}(f^r(y)) > e^{p_1 + p_2 + p_3} \} \ge \delta_0 k.$

We will now prove that

$$X_k \subset Y_n. \tag{4.20}$$

Given $y \in X_k$, set $x = \pi(y) \in \mathcal{L}$; then $y = \phi_s(x)$ for some $s \in [0, \tau(x))$. We have to show that $x \in Y_n$. Now $y \in X_k$ means that $B_{2\epsilon}(f^r(x)) \leq e^{p_1+p_2+p_3}$ for less then $(1-\delta_0)k$ values of $r = 0, 1, \ldots k - 1$. If $C_{2\epsilon}(\mathcal{P}^j(x)) \leq e^{p_2+p_3}$ for some $j = 0, 1, \ldots, n-1$, then for any integer r with

$$\tau_j(x) \le r < \tau_j(x) + 1 \tag{4.21}$$

we have $f^r(y) = \phi_t(\mathcal{P}^j(x))$ for some $t \in [-2, 2]$, so by Lemma 4.1(a) we have $B_{2\epsilon}(f^r(y)) \leq S_1C_{2\epsilon}(\mathcal{P}^j(x)) \leq e^{p_1+p_2+p_3}$. Assume for a moment that $x \notin Y_n$, i.e. $C_{2\epsilon}(\mathcal{P}^j(x)) \leq e^{p_2+p_3}$ for at least $(1 - \delta_0)n$ values of $j = 0, 1, \ldots, n - 1$. Notice that not more than $1/\tau_0$ values of j will produce the same r with (4.24), so there exist at least $(1 - \delta_0)n/\tau_0 \geq (1 - \delta_0)k$ values of r with $B_{2\epsilon}(f^r(y)) \leq e^{p_1+p_2+p_3}$. In other words, $B_{2\epsilon}(f^r(y)) > e^{p_1+p_2+p_3}$ for less than $\delta_0 k$ values of $r = 0, 1, \ldots, k - 1$, a contradiction with $y \in X_k$.

This proves (4.20). Combining (4.20), (4.19) and (4.18) we get

$$\mathsf{m}(X_k) \le \mathsf{m}(\widetilde{Y}_n) \le \frac{\hat{\tau}_0}{\tau^{\dagger}} \, \mu(Y_n) \le \frac{C_2 \hat{\tau}_0}{\tau^{\dagger}} \, e^{-c_2 n/s_0} \le \frac{C_2 \hat{\tau}_0}{\tau^{\dagger}} \, e^{-c_2 k \tau_0/s_0} = C_3 e^{-c_3 k \tau_0}$$

To complete the proof of Theorem 1.1, set $p_0 = p_1 + p_2 + p_3$. Denote by Θ_k the set of those $x \in \widetilde{\mathcal{L}}$ such that $\widetilde{R}_{2\epsilon}(f^r(x)) > e^{p_0}$ for at least δk values of $r = 0, 1, \ldots, k - 1$. We will prove an exponentially small estimate of $\mathfrak{m}(\Theta_k)$. Combining this with Proposition 3.1, in particular, using the left-hand-side inequality in (3.4), and (3.5) as well, will prove Theorem 1.1.

Denote by $Y_k^{+,i}$ the set of those $x \in \widetilde{\mathcal{L}}$ so that $\widetilde{R}_{2\epsilon,i}^+(f^r(x)) > e^{p_0}$ for at least $\delta_0 k$ values of $r = 0, 1, \ldots, k - 1$. Define $Y_k^{-,i}$ in a similar way replacing $\widetilde{R}_{2\epsilon,i}^+$ by $\widetilde{R}_{2\epsilon,i}^-$. The above argument shows that $\mathsf{m}(Y_k^{+,i}) \leq C_3 e^{-c_3 k}$ for all $i = 1, \ldots, k_0$. Using Theorem 4.4 for the flow ϕ_{-t} and replacing the map $f = \phi_1$ by $f' = \phi_{-1}$, the above argument shows that $\mathsf{m}(Y_k^{-,i}) \leq C_3 e^{-c_3 k}$. So, for the set $Y_k = \bigcup_{i=1}^{k_0} (Y_k^{+,i} \cup Y_k^{-,i})$ we get $\mathsf{m}(Y_k) \leq C e^{-ck}$ for some constants C > 0 and c > 0.

On the other hand, $\Theta_k \subset Y_k$. Indeed, let $x \in \Theta_k$. Then $\widetilde{R}_{2\epsilon}(f^r(x)) > e^{p_0}$ for at least δk values of $r = 0, 1, \ldots, k - 1$. Since $\widetilde{R}_{2\epsilon}(y) = \max_{1 \le i \le k_0} \max\{\widetilde{R}^+_{2\epsilon,i}(y), \widetilde{R}^-_{2\epsilon,i}(y)\}$, it follows that there exists $i = 1, \ldots, k_0$ such that either $\widetilde{R}^+_{2\epsilon,i}(f^r(x)) > e^{p_0}$ for at least $\frac{\delta k}{2k_0} = \delta_0 k$ values of $r = 0, 1, \ldots, k - 1$ or $\widetilde{R}^-_{2\epsilon,i}(f^r(x)) > e^{p_0}$ for at least $\frac{\delta k}{2k_0} = \delta_0 k$ values of $r = 0, 1, \ldots, k - 1$. That is $x \in Y_k^{+,i} \cup Y_k^{-,i} \subset Y_k$. Hence $\Theta_k \subset Y_k$ and therefore $\mathsf{m}(\Theta_k) \le Ce^{-ck}$.

5 Proof of Theorem 4.4

Throughout we work under the assumptions in the beginning of Sect. 4 and will use the notation from Sect. 4. Again, the constants $\epsilon > 0$, $\delta > 0$, $\delta_0 > 0$ and $s_0 = s_0(\epsilon, \delta_0)$ will be fixed as in Sect.4 and so will be the compact set K_0 and the functions u, g and ψ . As before $T = \mathcal{P}^{s_0} : R \longrightarrow R$.

Setting

$$A_m(x) = A_m^{(\epsilon)}(x) = \frac{e^{\psi^m(x)}}{e^{(\tilde{\chi} + \epsilon)g^m(x)}} \quad , \quad x \in \mathbb{R},$$

we have

$$D_{\epsilon}(x) = \sup_{m \ge 0} A_m(x) \ge 1$$

for all $x \in \mathcal{L}$. For such x (modifying an idea in [Sim]) define

$$k_{\epsilon}(x) = \min\left\{k \ge 0 : \psi^k(x) - (\tilde{\chi} + \epsilon)g^k(x) = \log D_{\epsilon}(x)\right\}.$$

This is well-defined for μ -almost all $x \in \mathcal{L}$, since $\int_B g \, d\mu = 1$ and $\int_B \psi \, d\mu = \tilde{\chi}$, so

$$\lim_{k \to \infty} \frac{\psi^k(x)}{g^k(x)} = \lim_{k \to \infty} \frac{\psi^k(x)}{k} \cdot \frac{k}{g^k(x)} = \tilde{\chi}$$

for μ -almost all $x \in \mathcal{L}$. Denote by \mathcal{L}' the set of those $x \in \mathcal{L}$ for which $k_{\epsilon}(x)$ is well defined; then $\mu(\mathcal{L}') = 1$.

For any $n, k \ge 0$ we have $g^{n+k}(x) = g^k(x) + g^n(T^k(x))$ and also $\psi^{n+k}(x) = \psi^k(x) + \psi^n(T^k(x))$, therefore

$$A_{n+k}(x) = A_k(x) A_n(T^k(x)).$$
(5.1)

Using this inductively one derives that for every $x \in R$ and every integer $k \ge 1$ we have

$$A_k(x) = A_1(x)A_1(T(x))\dots A_1(T^{k-1}(x)).$$
(5.2)

Next, consider the sets

$$\Omega_m(\epsilon) = \{ x \in \mathcal{L}' : k_\epsilon(x) > m \}.$$

Lemma 5.1. There exist constants C > 0 and c > 0 such that

$$\mu\left(\Omega_m(\epsilon)\right) \le C \, e^{-cm} \tag{5.3}$$

for all integers $m \geq 1$.

Proof of Lemma 5.1. Let $x \in \Omega_m(\epsilon)$. Then for $k = k_{\epsilon}(x)$ we have k > m and

$$\psi^k(x) - (\tilde{\chi} + \epsilon)g^k(x) = \log D_\epsilon(x) \ge 0.$$
(5.4)

Thus, $\psi^k(x)/k - (\tilde{\chi} + \epsilon)g^k(x)/k \ge 0$, so

$$\left[\frac{\psi^k(x)}{k} - \tilde{\chi}\right] + \tilde{\chi} \left[1 - \frac{g^k(x)}{k}\right] \ge \epsilon \frac{g^k(x)}{k} \ge \frac{\epsilon \tau_0}{\tau^{\dagger}}.$$

Here we used $g^k(x) = \frac{\tau_{ks_0}(x)}{s_0\tau^{\dagger}} \ge \frac{k\tau_0}{\tau^{\dagger}}$. First assume $\chi = \chi_i \neq 0$; then (4.14) implies $|\tilde{\chi}| \ge |\chi_i|/2 > 0$. Using the above, we either have

$$\frac{\psi^k(x)}{k} - \tilde{\chi} > \frac{\epsilon \tau_0}{2\tau^\dagger},\tag{5.5}$$

or

$$\left|1 - \frac{g^k(x)}{k}\right| > \frac{\epsilon \tau_0}{2|\tilde{\chi}|\tau^{\dagger}}.$$
(5.6)

Let Δ'_k be the set of those $x \in \Omega_m(\epsilon)$ for which (5.5) holds, and let Δ''_k be the set of those $x \in \Omega_m(\epsilon)$ for which (5.6) holds.

We can now use the classical Large Deviation Principle for the Hölder continuous functions ψ and q on \widehat{R} (and the fact that (\widehat{R}, T) is naturally isomorphic, up to a set of μ -measure zero, to a subshift of finite type⁵ – see [OP], [Ki] or [Y1]. It follows from it that there exist constants⁶ C' > 0 and c' > 0, independent of k and m, such that $\mu(\Delta'_k) \leq C'e^{-c'k}$ and $\mu(\Delta''_k) \leq C'e^{-c'k}$. Since $\Omega_m(\epsilon) \subset \bigcup_{k=m+1}^{\infty} (\Delta'_k \cup \Delta''_k)$, it follows that $\mu(\Omega_m(\epsilon)) \leq Ce^{-cm}$ for some constants C, c > 0independent of m.

When $\chi = \chi_i = 0$, (4.14) implies $0 \le \tilde{\chi} < \frac{2C_1\delta_0}{C_0}$, so (5.4) gives

$$\left[\frac{\psi^k(x)}{k} - \tilde{\chi}\right] \ge \epsilon \, \frac{g^k(x)}{k} + \tilde{\chi} \, \frac{g^k(x)}{k} - \tilde{\chi} \ge \epsilon \, \frac{g^k(x)}{k} - \frac{8C_1\delta_0}{C_0} \ge \frac{\epsilon\tau_0}{2\tau^{\dagger}},$$

and then we can proceed as above. \blacksquare

⁵This natural isomorphism sends both ψ and $g = \frac{1}{s_0 \tau^{\dagger}} \tau_{s_0}$ to Hölder continuous functions with respect to the metric d_{θ} on the shift space. Neither ψ nor g is cohomologous to a constant.

⁶Notice that the rate function involved depends on $T = \mathcal{P}^{s_0}$ and therefore on ϵ and δ_0 .

The following lemma will be used later.

Lemma 5.2. Let $x \in \mathcal{L}'$ and let $k = k_{\epsilon}(x)$. Then:

- (a) $T^k(x) \in Q_0(\epsilon)$.
- (b) $k_{\epsilon}(x)$ coincides with the smallest integer $\ell \geq 0$ such that $T^{\ell}(x) \in Q_0(\epsilon)$.

Proof. (a) From the definition of $k_{\epsilon}(x)$ it follows that $D_{\epsilon}(x) = A_{k_{\epsilon}(x)}(x) \ge A_{n+k_{\epsilon}(x)}(x)$ for all $n \ge 0$. Using this and (5.1) with $k = k_{\epsilon}(x)$, we get $A_n(T^k(x)) \le 1$ for all $n \ge 0$, so $D_{\epsilon}(T^k(x)) \le 1$. Thus, $T^k(x) \in Q_0(\epsilon)$.

(b) Set $k = k_{\epsilon}(x)$. Assume that there exists $\ell = 0, 1, \ldots, k-1$ with $T^{\ell}(x) \in Q_0(\epsilon)$. Then $A_n(T^{\ell}(x)) \leq 1$ for all $n \geq 0$, so by (5.1), $A_{n+\ell}(x) = A_{\ell}(x)A_n(T^{\ell}(x)) \leq A_{\ell}(x)$ for all integers $n \geq 0$. In particular, $A_k(x) = A_{(k-\ell)+\ell}(x) \leq A_{\ell}(x)$, which is a contradiction with the choice of $k = k_{\epsilon}(x)$.

The above already implies exponentially small tails.

Lemma 5.3. There exist constants C > 0 and c > 0 such that

$$\mu\left(R\setminus \bigcup_{k=0}^{m-1}T^{-k}(Q_0(\epsilon))\right) \le Ce^{-cm}$$

for all $m \geq 1$.

Proof. Let $x \in \mathcal{L}'$ be such that $x \notin \Omega_m(\epsilon)$ for some $m \ge 1$, i.e. $k_{\epsilon}(x) < m$. Then by Lemma 5.2(a), $T^k(x) \in Q_0(\epsilon)$, so $x \in T^{-k}(Q_0(\epsilon))$ for some k < m. This shows that

$$\mathcal{L}' \setminus \Omega_m(\epsilon) \subset \bigcup_{k=0}^{m-1} T^{-k}(Q_0(\epsilon)) \quad , \text{ i.e.} \quad \mathcal{L}' \setminus \bigcup_{k=0}^{m-1} T^{-k}(Q_0(\epsilon)) \subset \Omega_m(\epsilon),$$

and by (5.3), $\mu\left(R \setminus \bigcup_{k=0}^{m-1} T^{-k}(Q_0(\epsilon))\right) \leq Ce^{-cm}$, which proves the Lemma.

Next, since the functions ψ and g are continuous on \widehat{R} (with respect to the chosen metric d_{θ}), the set

$$F = \{x \in \widehat{R} : \psi(x) - (\widetilde{\chi} + \epsilon/2)g(x) \le 0\}$$
(5.7)

is closed in \widehat{R} . Take $\eta = \frac{\delta_0}{C_0}$ and consider the open set

$$V_{\eta} = \{ x \in \widehat{R} : \psi(x) - (\widetilde{\chi} + \epsilon/2)g(x) < \eta \}.$$

Clearly, $F \subset V_{\eta}$, so there exists a Hölder continuous (with respect to d_{θ}) function $\varphi : \widehat{R} \longrightarrow [0, 1]$ with $\varphi = 1$ on F and $\varphi = 0$ on $\widehat{R} \setminus V_{\eta}$. Fix φ with this property, and consider the Hölder continuous function

$$h = \left(\psi - (\tilde{\chi} + \epsilon/2)g\right)\varphi$$

Notice that

$$\left| \int_{\widehat{R}} h \, d\mu + \frac{\epsilon}{2} \right| < \frac{8C_1 \delta_0}{C_0}. \tag{5.8}$$

Indeed, it follows from (4.10), (4.11) and (4.14) that for $x \in K_0$ we have

$$|(\psi(x) - (\tilde{\chi} + \epsilon/2)g(x)) + \epsilon/2| \leq |\psi(x) - \tilde{\chi}| + |\tilde{\chi}| |1 - g(x)| + \frac{\epsilon}{2}|1 - g(x)| \leq \frac{5C_1\delta_0}{C_0}.$$

Since $\frac{5C_1\delta_0}{C_0} < \frac{\epsilon}{2}$ by (4.3), it follows that $\psi(x) - (\tilde{\chi} + \epsilon/2)g(x) < 0$ on K_0 , so $K_0 \subset F$ and therefore $\varphi = 1$ on K_0 . Combining the above with (4.9) gives

$$\left| \int_{\widehat{R}} h \, d\mu + \frac{\epsilon}{2} \right| \le \left| \int_{K_0} (h + \epsilon/2) \, d\mu \right| + (\|h\|_{\infty} + \epsilon/2) \mu(\widehat{R} \setminus K_0) \le \frac{5C_1 \delta_0}{C_0} + \frac{3C_1 \delta_0}{C_0} < \frac{8C_1 \delta_0}{C_0} + \frac{3C_1 \delta_0}{C_0} < \frac{8C_1 \delta_$$

since $||h||_{\infty} \le ||\psi||_{\infty} + C_1 \le 3C_1$. This proves (5.8).

Apart from the above notice that

$$\left[\psi(y) - (\tilde{\chi} + \epsilon/2)g(y)\right]\chi_F(y) \ge h(y) - \eta \tag{5.9}$$

for all $y \in \mathcal{L}'$. Here χ_F is the *characteristic function* of F. Indeed, if $y \in F$, then $\varphi(y) = 1$, so $\psi(y) - (\tilde{\chi} + \epsilon/2)g(y) = h(y)$ and (5.9) holds trivially. Let $x \notin F$; then the left-hand-side of (5.9) is 0. If moreover $x \notin V_\eta$, then $\varphi(y) = 0$, so h(y) = 0, and (5.9) becomes $0 \ge -\eta$ which is obviously true. Finally, assume $y \in V_\eta \setminus F$. By the definitions of F and V_η we have $0 < \psi(y) - (\tilde{\chi} + \epsilon/2)g(y) < \eta$. Multiplying this by $\varphi(y) \in [0, 1]$ we get $0 \le h(y) = (\psi(y) - (\tilde{\chi} + \epsilon/2)g(y))\varphi(y) \le \eta\varphi(y) \le \eta$. That is, $0 \ge h(y) - \eta$, so (5.9) holds again.

The following lemma proves Theorem 4.4. Recall the set $\Xi_m(\epsilon, \hat{\delta})$ defined just before Theorem 4.4.

Lemma 5.4. Let $\hat{\delta} \in [\delta_0, 1]$. There exist constants $C = C(\epsilon, \hat{\delta}) \ge 1$ and $c = c(\epsilon, \hat{\delta}) > 0$ such that $\mu\left(\Xi_n(\epsilon, \hat{\delta})\right) \le Ce^{-cn}$ for any integer $n \ge 1$.

Proof. Apart from the functions $A_m(x) = A_m^{(\epsilon)}(x)$ defined in the beginning of Sect. 5 we will also use the functions $A_m^{(\epsilon/2)}(x)$. Clearly

$$A_m^{(\epsilon/2)}(x) = \frac{e^{\psi^m(x)}}{e^{(\tilde{\chi} + \epsilon/2)g^m(x)}} = A_m^{(\epsilon)}(x)e^{\epsilon g^m(x)/2} \ge A_m^{(\epsilon)}(x)e^{m\epsilon\tau_0/(2\tau^{\dagger})} \ge A_m^{(\epsilon)}(x)$$
(5.10)

for all $x \in \mathcal{L}'$ and all integers $m \ge 0$.

Notice that $Q_0(\epsilon/2) \subset Q_0(\epsilon)$. Indeed, if $y \in Q_0(\epsilon/2)$, then $A_m^{(\epsilon/2)}(y) \leq 1$ for all $m \geq 0$, and by (5.10), $A_m^{(\epsilon)}(y) \leq A_m^{(\epsilon/2)}(y) \leq 1$ for all $m \geq 0$, so $y \in Q_0(\epsilon)$. Moreover, for any $y \in Q_0(\epsilon/2)$ we have $A_1^{(\epsilon/2)}(y) \leq 1$, that is $\psi(y) - (\tilde{\chi} + \epsilon/2)g(y) \leq 0$, so $y \in F$, the set defined by (5.7). Thus, $Q_0(\epsilon/2) \subset F$. However in general $Q_0(\epsilon)$ is not contained in F.

Next, setting $\delta_1 = \delta_0/C_0$, it follows from (5.3) that for any integer $n \ge 1$ we have $\mu(\Omega_{[\delta_1 n]}(\epsilon) \le Ce^{-c(\delta_1 n-1)})$, so

$$\mu\left(\cup_{j=0}^{n}T^{-j}(\Omega_{[\delta_{1}n]}(\epsilon))\right) \leq (n+1)Ce^{-c(\delta_{1}n-1)}.$$

Setting

$$\Omega^{(n)} = \bigcup_{j=0}^{n} T^{-j}(\Omega_{[\delta_1 n]}(\epsilon)),$$

it follows from the above that there exist constants $C_4 = C_4(\epsilon, \delta_0) > 0$ and $c_4 = c_4(\epsilon, \delta_0) > 0$ with

$$\mu(\Omega^{(n)}) \le C_4 e^{-c_4 n} \tag{5.11}$$

for all integers $n \ge 1$. Clearly, for every $x \in \mathcal{L}' \setminus \Omega^{(n)}$ we have $k_{\epsilon}(T^j(x)) < \delta_1 n = \frac{\delta_0 n}{C_0}$ for all $j = 0, 1, \ldots, n$.

Next, fix for a moment an arbitrary $x \in \Xi_n \setminus \Omega^{(n)}$. We will now construct a sequence of points x_i on the *orbit*

$$\mathcal{O}_n(x) = \{x, T(x), \dots, T^{n-1}(x)\}$$

and certain integers k_i , t_i using the set $Q_0(\epsilon)$. We will do the construction carefully and in all details, although some of the details will not be used later.

If $x \notin Q_0(\epsilon)$, set $x_0 = x$ and $t_0 = 0$. If $x \in Q_0(\epsilon)$ let $t_0 \ge 1$ be the largest integer such that $T^j(x) \in Q_0(\epsilon)$ for all $j = 0, 1, \ldots, t_0 - 1$. Then set $x_0 = T^{t_0}(x) \notin Q_0(\epsilon)$ and $k_0 = k_{\epsilon}(x_0) \ge 1$. Notice that $k_0 < \delta_1 n$.

By Lemma 5.2(a) we have $T^{k_0}(x_0) \in Q_0(\epsilon)$. Next, let $t_1 \ge 1$ be the largest integer such that $T^{k_0+j}(x_0) \in Q_0(\epsilon)$ for all $j = 0, 1, \ldots, t_1 - 1$. Then set $x_1 = T^{k_0+t_1}(x_0) \notin Q_0(\epsilon)$ and $k_1 = k_{\epsilon}(x_1) \ge 1$.

By induction we construct a sequence of points $x_0, x_1, \ldots, x_{s-1}$ and positive integers $t_0, t_1, \ldots, t_{s-1}$ and $k_0, k_1, \ldots, k_{s-1}$ such that $x_{i+1} = T^{k_i+t_{i+1}}(x_i)$ for all $i = 0, 1, \ldots, s-2$, $k_i = k_{\epsilon}(x_i)$ for all $i = 0, 1, \ldots, s-1$, and $t_{i+1} \ge 1$ is the maximal integer such that $T^{k_i+j}(x_i) \in Q_0(\epsilon)$ for all $j = 0, 1, \ldots, t_{i+1} - 1$. Thus, $x_i \notin Q_0(\epsilon)$, so $k_i \ge 1$ for all $i = 0, 1, \ldots, s-1$.

For s we choose the maximal integer with

$$\sum_{i=0}^{s-1} t_i + \sum_{i=0}^{s-1} k_i \le n.$$
(5.12)

If we have equality in (5.12), set $t_s = 0$, $k_s = 0$ and $x_s = T^{k_{s-1}}(x_{s-1})$. If we have a strict inequality in (5.12), denote by t_s the largest positive integer such that $T^{k_{s-1}+j}(x_{s-1}) \in Q_0(\epsilon)$ for all $j = 0, 1, \ldots, t_s - 1$ and

$$\sum_{i=0}^{s} t_i + \sum_{i=0}^{s-1} k_i \le n.$$
(5.13)

Set $x_s = T^{k_{s-1}+t_s}(x_{s-1})$. If there is an equality in (5.13), set $k_s = 0$. If there is a strict inequality in (5.13), take $k_s \ge 0$ so that

$$\sum_{i=0}^{s} t_i + \sum_{i=s}^{s} k_i = n.$$
(5.14)

It follows from the above construction that (5.14) always holds. Moreover, $k_i > 0$ for $i = 0, 1, \ldots, s - 1$, while for i = s we may have $k_s = 0$ and $x_s \in Q_0(\epsilon)$. However, $k_s < \delta_1 n$ always holds, since $x \notin \Omega^{(n)}$. Hence:

(i) $T^{k_i+j}(x_i) \in Q_0(\epsilon)$ for all $j = 0, 1..., t_{i+1} - 1, i = 0, 1, ..., s - 1$,

ii)
$$k_i = k_{\epsilon}(x_i) \ge 1$$
 for all $i = 0, 1, \dots, s - 1$, and $0 \le k_s \le k_{\epsilon}(x_s) < \delta_1 n$.

Thus, the above construction is such that the orbits $\mathcal{O}_{k_i}(x_i)$ are disjoint, have no common points with $Q_0(\epsilon)$, and

$$\mathcal{O}_n(x) \setminus Q_0(\epsilon) = \bigcup_{i=0}^s \mathcal{O}_{k_i}(x_i)$$

In particular, it follows from it and the choice of ψ in Lemma 4.2 that

$$|\log A_{k_s}^{(\epsilon/2)}(x_s)| = |\psi^{k_s}(x_s) - (\tilde{\chi} + \epsilon/2)g^{k_s}(x_s)| \le k_s(\|\psi\|_{\infty} + 2C_1) \le 4C_1k_s < 4C_1\delta_1n \le \frac{\epsilon\tau_0}{4\tau^{\dagger}}\hat{\delta}n,$$

using (4.3) and $\delta_1 = \delta_0 / C_0 \le \hat{\delta} / C_0$.

Since $x \in \Xi_n = \Xi_n(\epsilon, \hat{\delta})$, it follows from (i) and (ii) that

$$\sum_{i=0}^{s} k_i \ge \hat{\delta}n. \tag{5.15}$$

Next, it follows from (5.2) that for any $i = 0, 1, \ldots, s$ we have

$$\log A_{k_i}^{(\epsilon/2)}(x_i) = \sum_{q=0}^{k_i-1} \log A_1^{(\epsilon/2)}(T^q(x_i)).$$
(5.16)

Similarly,

$$\psi^n(x) - (\tilde{\chi} + \epsilon/2)g^n(x) = \log A_n^{(\epsilon/2)}(x) = \sum_{q=0}^{n-1} \log A_1^{(\epsilon/2)}(T^q(x))$$

and it follows from the above construction that

$$\psi^{n}(x) - (\tilde{\chi} + \epsilon/2)g^{n}(x) = \sum_{i=0}^{s} \log A_{k_{i}}^{(\epsilon/2)}(x_{i}) + \sum_{i=0}^{s-1} \log A_{t_{i}}^{(\epsilon/2)}(T^{k_{i}}(x_{i}))$$
$$= \sum_{i=0}^{s} \log A_{k_{i}}^{(\epsilon/2)}(x_{i}) + \sum_{T^{q}(x) \in Q_{0}(\epsilon)}^{n-1} \log A_{1}^{(\epsilon/2)}(T^{q}(x)). \quad (5.17)$$

By (ii) we have $A_{k_i}^{(\epsilon)}(x_i) > 1$ for i = 0, 1, ..., s-1. This and (5.10) imply $A_{k_i}^{(\epsilon/2)}(x_i) > e^{\epsilon g^{k_i}(x_i)/2}$ and so $\log A_{k_i}^{(\epsilon/2)}(x_i) \geq \frac{\epsilon \tau_0}{2\tau^{\dagger}} k_i$ for all i < s. Combining this with the above, (5.15) and (5.17) give

$$\psi^{n}(x) - (\tilde{\chi} + \epsilon/2)g^{n}(x) \ge \frac{\epsilon\tau_{0}}{2\tau^{\dagger}}\hat{\delta}n - |\log A_{k_{s}}^{(\epsilon)}(x_{s})| + \sum_{\substack{q=0\\T^{q}(x)\in Q_{0}(\epsilon)}}^{n-1} A_{1}^{(\epsilon/2)}(T^{q}(x)) \ge \frac{\epsilon\tau_{0}\hat{\delta}}{4\tau^{\dagger}}n + (I) + (II),$$

where (I) is the sum of the positive terms of the sum $\sum_{\substack{q=0\\T^q(x)\in Q_0(\epsilon)}}^{n-1} A_1^{(\epsilon/2)}(T^q(x))$ and (II) is the sum

of the other terms in this sum. Then $(I) \ge 0$, and, recalling the set F defined by (5.7),

$$(II) = \sum_{\substack{q=0\\T^q(x)\in F\cap Q_0(\epsilon)}}^{n-1} \log A_1^{(\epsilon/2)}(T^q(x)) \ge \sum_{\substack{q=0\\T^q(x)\in F}}^{n-1} \log A_1^{(\epsilon/2)}(T^q(x)).$$

(The terms appearing in the sum in the right-hand-side that do not appear in the sum in the middle are all non-positive.) Hence

$$\begin{split} \psi^{n}(x) - (\tilde{\chi} + \epsilon/2)g^{n}(x) &\geq \frac{\epsilon\tau_{0}\hat{\delta}}{4\tau^{\dagger}}n + \sum_{\substack{q=0\\T^{q}(x)\in F}}^{n-1}\log A_{1}^{(\epsilon/2)}(T^{q}(x)) \\ &= \frac{\epsilon\tau_{0}\hat{\delta}}{4\tau^{\dagger}}n + \sum_{q=0}^{n-1}[\psi(T^{q}(x)) - (\tilde{\chi} + \epsilon/2)g(T^{q}(x))]\chi_{F}(T^{q}(x)) \\ &= \frac{\epsilon\tau_{0}\hat{\delta}}{4\tau^{\dagger}}n + \gamma^{n}(x), \end{split}$$

where $\gamma(y) = [\psi(y) - (\tilde{\chi} + \epsilon/2)g(y)]\chi_F(y)$. On the other hand (5.9) implies $\gamma^n(x) \ge h^n(x) - n\eta$. Using this and $\eta = \frac{\delta_0}{C_0}$, it follows that

$$\psi^n(x) - (\tilde{\chi} + \epsilon/2)g^n(x) \ge \frac{\epsilon\tau_0\hat{\delta}}{4\tau^{\dagger}}n + h^n(x) - \frac{\delta_0}{C_0}n.$$

For the integral $H = \int_{\widehat{R}} h \, d\mu$ it follows from (5.8) that $H + \frac{\epsilon}{2} > -\frac{8C_1\delta_0}{C_0}$. So, we can now rewrite the above as

$$\left[\frac{\psi^n(x)}{n} - \tilde{\chi}\right] + \tilde{\chi} \left[1 - \frac{g^n(x)}{n}\right] + \frac{\epsilon}{2} \left[1 - \frac{g^n(x)}{n}\right] + \left[H - \frac{h^n(x)}{n}\right] \ge H + \frac{\epsilon}{2} + \frac{\epsilon\tau_0\hat{\delta}}{4\tau^{\dagger}} - \frac{\delta_0}{C_0} \\ \ge \frac{\epsilon\tau_0\hat{\delta}}{4\tau^{\dagger}} - \frac{9C_1\delta_0}{C_0} \ge \frac{\epsilon\tau_0\hat{\delta}}{8\tau^{\dagger}},$$

since $\hat{\delta} \ge \delta_0$ and $C_0 \ge \frac{100C_1\tau^{\dagger}}{\epsilon\tau_0}$. Thus,

$$\left[\frac{\psi^n(x)}{n} - \tilde{\chi}\right] + \tilde{\chi} \left[1 - \frac{g^n(x)}{n}\right] + \frac{\epsilon}{2} \left[1 - \frac{g^n(x)}{n}\right] + \left[H - \frac{h^n(x)}{n}\right] \ge \frac{\epsilon \tau_0 \hat{\delta}}{8\tau^{\dagger}}$$

As in the proof of Lemma 5.1, assume first that $\chi = \chi_i \neq 0$. Then the above argument shows that $\Xi_n \setminus \Omega^{(n)} \subset \Xi'_n \cup \Xi''_n \cup \Xi'''_n \cup \Xi'''_n$, where Ξ'_n is the set of those $x \in \mathcal{L}' \setminus \Omega^{(n)}$ such that $\frac{\psi^n(x)}{n} - \tilde{\chi} > \frac{\epsilon\tau_0\hat{\delta}}{32\tau^{\dagger}}$, Ξ''_n is the set of those $x \in \mathcal{L}' \setminus \Omega^{(n)}$ such that $\left|1 - \frac{g^n(x)}{n}\right| > \frac{\epsilon\tau_0\hat{\delta}}{32|\chi|\tau^{\dagger}}$, Ξ'''_n is the set of those $x \in \mathcal{L}' \setminus \Omega^{(n)}$ such that $\left|1 - \frac{g^n(x)}{n}\right| > \frac{\tau_0\hat{\delta}}{16\tau^{\dagger}}$, and Ξ''''_n is the set of those $x \in \mathcal{L}' \setminus \Omega^{(n)}$ such that $\left|H - \frac{h^n(x)}{n}\right| > \frac{\epsilon\tau_0\hat{\delta}}{32\tau^{\dagger}}$.

As in the proof of Lemma 5.1, it now follows from the classical Large Deviation Principle (see [OP], [Ki] or [Y1]) that there exist constants $C' = C'(\epsilon, \hat{\delta}) > 0$ and $c' = c(\epsilon, \hat{\delta}) > 0$ such that $\mu(\Xi_n \setminus \Omega^{(n)}) \leq C' e^{-c'n}$. By (5.11), $\mu(\Omega^{(n)}) \leq C_4 e^{-c_4 n}$, so there exist constants $C \geq 1$ and c > 0 such that $\mu(\Xi_n) \leq C e^{-cn}$ for all n.

When $\chi = \chi_i = 0$, a slightly different argument (as in the proof of Lemma 5.1) shows that again there exist constants $C \ge 1$ and c > 0 such that $\mu(\Xi_n) \le C e^{-cn}$ for all n.

6 Proof of Proposition 2.1

Throughout we assume that ϕ_t is a C^2 contact Anosov flow on M with a C^2 invariant contact form ω . Then the two-form $d\omega$ is C^1 , so there exists a constant $C_0 > 0$ such that

$$|d\omega_x(u,v)| \le C_0 ||u|| ||v|| \quad , \quad u,v \in T_x M \,, \, x \in M,$$
(6.1)

and for every $x \in M$ and every $u \in E^s(x)$ (or $u \in E^u(x)$) with ||u|| = 1 there exists $v \in E^u(x)$ (reps. $v \in E^s(x)$) with ||v|| = 1 such that $d\omega_x(u, v) \ge 1/C_0$.

Fix for a moment $\epsilon > 0$. We will use the notation from Sect. 3. Following general procedures (see Theorem S.2.10 in [KM]) for all $x \in \mathcal{L}'_0$ and $i = 1, \ldots, k_0$ consider the inner product

$$\langle u, v \rangle'_{x,i} = \sum_{m \in \mathbb{Z}} \langle df_x^m(u), df_x^m(v) \rangle e^{-2m\chi_i - 4\epsilon |m|} \quad , \quad u, v \in E_i(x),$$

where $\langle \cdot, \cdot \rangle$ is the inner product on $T_x M$ determined by the Riemann metric. Then, given $u = u_1 + \ldots + u_{k_0}$, $v = v_1 + \ldots + v_{k_0} \in T_x M$ with $u_i, v_i \in E_i(x)$ for all *i*, define

$$\langle u, v \rangle'_x = \sum_{i=1}^{k_0} \langle u_i, v_i \rangle'_{x,i} \quad , \quad ||u||'_x = \sqrt{\langle u, u \rangle'_x}.$$

Notice that for $u \in E_i(x)$ we have

$$\begin{aligned} \langle u, u \rangle_{x,i}' &= \sum_{m \ge 0} \|df_x^m(u)\|^2 \, e^{-2m\chi_i - 4\epsilon m} + \sum_{m < 0} \|df_x^m(u)\|^2 \, e^{-2m\chi_i + 4\epsilon m} \\ &\leq \sum_{m \ge 0} (R_{\epsilon}^+(x) e^{m\chi_i + m\epsilon} \|u\|)^2 \, e^{-2m\chi_i - 4\epsilon m} + \sum_{m < 0} (R_{\epsilon}^-(x) e^{m\chi_i - m\epsilon} \|u\|)^2 \, e^{-2m\chi_i + 4\epsilon m} \\ &= (R_{\epsilon}(x)\|u\|)^2 \left[\sum_{m \ge 0} e^{-2m\epsilon} + \sum_{m < 0} e^{2m\epsilon} \right] \le \frac{1 + e^{-2\epsilon}}{1 - e^{-2\epsilon}} (R_{\epsilon}(x)\|u\|)^2 \le \frac{2}{\epsilon} (R_{\epsilon}(x)\|u\|)^2. \end{aligned}$$

Thus, $||u||'_{x,i} \leq \sqrt{2/\epsilon} R_{\epsilon}(x) ||u||$. Since the subspaces $E_i(x)$ are mutually orthogonal with respect to $\langle \cdot, \cdot \rangle'_x$, it follows that, setting $K_0 = K_0(\epsilon) = 2\sqrt{k_0/\epsilon}$, we have

$$\|u\|'_{x} \le K_{0}(\epsilon) R_{\epsilon}(x) \|u\| \quad , \quad u \in T_{x}M \ , \ x \in \mathcal{L}'_{0}.$$

$$(6.2)$$

In particular, if $u = u_1 + \ldots + u_{k_0}$ with $u_i \in E_i(x)$ for all *i*, then

$$||u_i|| \le ||u_i||'_{x,i} \le ||u||'_x \le K_0(\epsilon) R_\epsilon(x) ||u|| \quad , \quad i = 1, \dots, k_0.$$
(6.3)

Next, for every $i = 1, \ldots, k_0, x \in \mathcal{L}'_0$ and $\epsilon > 0$ set

$$\overline{A}_{i}(x,\epsilon) = \sup\left\{\frac{\|df_{x}^{n+k}(v)\| e^{-(\chi_{i}+\epsilon)n-\epsilon|k|}}{\|df_{x}^{k}(v)\|} : n \ge 0, k \in \mathbb{Z}, v \in E_{i} \setminus \{0\}\right\},$$
$$\widetilde{A}_{i}(x,\epsilon) = \sup\left\{\frac{\|df_{x}^{n+k}(v)\| e^{-(\chi_{i}-\epsilon)n-\epsilon|k|}}{\|df_{x}^{k}(v)\|} : n \le 0, k \in \mathbb{Z}, v \in E_{i} \setminus \{0\}\right\},$$

and

$$A_{\epsilon}(x) = \max_{1 \le i \le k_0} \max\{\overline{A}_i(x,\epsilon), \widetilde{A}_i(x,\epsilon)\}.$$

It follows from the argument in the proof of Proposition 1 in [FHY] that $\overline{A}_i(x,\epsilon)$ and $\widetilde{A}_i(x,\epsilon)$ are finite for all i, so $A_{\epsilon}(x)$ is well-defined for all $x \in \mathcal{L}'_0$, and moreover

$$\frac{\|df_x^n(v)\|}{e^{n\chi_i}} \le A_\epsilon(x) e^{|n|\epsilon} \|v\| \quad , \quad v \in E_i(x) \; , \; n \in \mathbb{Z},$$

$$(6.4)$$

for all $x \in \mathcal{L}'_0$ and all $i = 1, \ldots, k_0$. It now follows from (3.1) and (3.2) that

 $R_{\epsilon}(x) \leq A_{\epsilon}(x)$

for all $\epsilon > 0$ and all $x \in \mathcal{L}'_0$. It is easy to see that $\overline{A}_i(x, \epsilon)$ and $\widetilde{A}_i(x, \epsilon)$ are ϵ -slowly varying functions for all i, so $A_{\epsilon}(x)$ is also an ϵ -slowly varying function.

Next, set $K_1(\epsilon) = C_0^2 K_0(\epsilon)$. We will prove that

$$\frac{1}{K_1(\epsilon) (R_\epsilon(x))^2 e^{m\epsilon}} \le \frac{\|df_x^m(v)\|}{e^{m\chi_i}} \quad , \quad m \ge 0 \; , \; v \in E_i(x) \; , \; \|v\| = 1, \tag{6.5}$$

for all $x \in \mathcal{L}'_0$ and $i = 1, \ldots, k_0$. Given $v \in E_i(x)$ with ||v|| = 1, there exists $w \in T_x M$ with ||w|| = 1 such that $d\omega_x(v, w) \ge 1/C_0$. Let $w = w_1 + \ldots + w_{k_0}$, where $w_k \in E_k(x)$ for all k. Since the flow is contact, there exists j with $\chi_j = -\chi_i$. It is now easy to see that $d\omega_x(v, w) = d\omega_x(v, w_j)$. Indeed, setting $x_m = f^m(x)$, for $\chi_k < \chi_j = -\chi_i$ we have $\chi_k < \chi_j - 3\epsilon$ by the choice of ϵ , so using (6.1) we get

$$\begin{aligned} |d\omega_x(v, w_k)| &= |d\omega_{x_m}(df_x^m(v), df_x^m(w_k))| \le C_0 ||df_x^m(v)|| \, ||df_x^m(w_k)|| \\ &\le C_0 R_{\epsilon}^2(x) ||v|| \, ||w_k|| \, e^{m\chi_i + m\epsilon} e^{m\chi_k + m\epsilon} \to 0 \end{aligned}$$

as $m \to \infty$. Thus, $d\omega_x(v, w_k) = 0$. In a similar way one deals with the case $\chi_k > \chi_j$, considering $m \to -\infty$.

Hence for any $m \ge 0$ we have

$$\frac{1}{C_0} \le d\omega_x(v, w_j) = d\omega_{f^m(x)}(df_x^m(v), df_x^m(w_j)) \le C_0 \|df_x^m(v)\| \|df_x^m(w_j)\|.$$

This, (3.1) and (6.3) imply

$$\begin{aligned} \frac{\|df_x^m(v)\|}{e^{m\chi_i}} &\geq \frac{1}{C_0^2 \|df_x^m(w_j)\|e^{m\chi_i}} \geq \frac{1}{C_0^2 R_{\epsilon}^+(x) e^{m\chi_j + m\epsilon} \|w_j\|e^{m\chi_i}} \\ &\geq \frac{1}{C_0^2 R_{\epsilon}^+(x) e^{m\epsilon} K_0(\epsilon) R_{\epsilon}(x) \|w\|} \geq \frac{1}{C_0^2 K_0(\epsilon) (R_{\epsilon}(x))^2 e^{m\epsilon}}, \end{aligned}$$

which proves (6.5). In a similar way one shows that

$$\frac{1}{K_1(\epsilon) (R_\epsilon(x))^2 e^{m\epsilon}} \le \frac{\|df_x^{-m}(v)\|}{e^{-m\chi_i}} \quad , \quad m \ge 0 \; , \; v \in E_i(x) \; , \; \|v\| = 1, \tag{6.6}$$

for all $x \in \mathcal{L}'_0$ and $i = 1, \ldots, k_0$.

Next, we will prove that

$$\overline{A}_i(x,\epsilon) \le K_1(\epsilon/2)(R_{\epsilon/2}(x))^3 \tag{6.7}$$

for all $x \in \mathcal{L}'_0$ and $i = 1, ..., k_0$. Fix x and i for a moment. Given $v \in E_i(x) \setminus \{0\}$, $n \ge 0$ and $k \in \mathbb{Z}$, we will consider 3 cases.

Case 1. $k \ge 0$; then $n + k \ge 0$, too. Using (3.1) and (6.5) with ϵ replaced by $\epsilon/2$, we get

$$\frac{\|df_x^{n+k}(v)\| e^{-(\chi_i+\epsilon)n-\epsilon|k|}}{\|df_x^k(v)\|} \leq \frac{R_{\epsilon/2}(x)e^{(n+k)(\chi_i+\epsilon/2)}\|v\| e^{-(\chi_i+\epsilon)n-\epsilon|k|}}{\frac{e^{k\chi_i}\|v\|}{K_1(\epsilon/2)(R_{\epsilon/2}(x))^2e^{k\epsilon/2}}} \\
= K_1(\epsilon/2)(R_{\epsilon/2}(x))^3 e^{(n+k)\epsilon/2}e^{-n\epsilon-k\epsilon}e^{k\epsilon/2} \\
\leq K_1(\epsilon/2)(R_{\epsilon/2}(x))^3.$$

Case 2. k < 0 and $n + k \ge 0$. Then (3.1) and (6.6) imply

$$\frac{\|df_x^{n+k}(v)\| e^{-(\chi_i+\epsilon)n-\epsilon|k|}}{\|df_x^k(v)\|} \leq \frac{R_{\epsilon/2}(x)e^{(n+k)(\chi_i+\epsilon/2)}\|v\| e^{-(\chi_i+\epsilon)n+\epsilon k}}{\frac{e^{k\chi_i}\|v\|}{K_1(\epsilon/2)(R_{\epsilon/2}(x))^2e^{-k\epsilon/2}}} \\ = K_1(\epsilon/2)(R_{\epsilon/2}(x))^3e^{(n+k)\epsilon/2}e^{-n\epsilon+k\epsilon}e^{-k\epsilon/2} \\ \leq K_1(\epsilon/2)(R_{\epsilon/2}(x))^3.$$

Case 3. k < 0 and n + k < 0. Then (3.2) and (6.6) imply

$$\frac{\|df_x^{n+k}(v)\| e^{-(\chi_i+\epsilon)n-\epsilon|k|}}{\|df_x^k(v)\|} \leq \frac{R_{\epsilon/2}(x)e^{(n+k)(\chi_i-\epsilon/2)}\|v\| e^{-(\chi_i+\epsilon)n+\epsilon k}}{\frac{e^{k\chi_i}\|v\|}{K_1(\epsilon/2)(R_{\epsilon/2}(x))^2e^{-k\epsilon/2}}} \\ = K_1(\epsilon/2)(R_{\epsilon/2}(x))^3 e^{-(n+k)\epsilon/2}e^{-n\epsilon+k\epsilon}e^{-k\epsilon/2} \\ \leq K_1(\epsilon/2)(R_{\epsilon/2}(x))^3.$$

This proves (6.7). In a similar way we will now prove

$$\widetilde{A}_i(x,\epsilon) \le K_1(\epsilon/2)(R_{\epsilon/2}(x))^3 \tag{6.8}$$

for all $x \in \mathcal{L}'_0$ and $i = 1, ..., k_0$. Fix x and i for a moment and let $v \in E_i(x) \setminus \{0\}$, $n \leq 0$ and $k \in \mathbb{Z}$. Again we will consider 3 cases.

Case 4. $k \ge 0$ and $n + k \ge 0$. Using (3.1) and (6.5), we get

$$\frac{\|df_x^{n+k}(v)\| \ e^{-(\chi_i - \epsilon)n - \epsilon|k|}}{\|df_x^k(v)\|} \leq \frac{R_{\epsilon/2}(x)e^{(n+k)(\chi_i + \epsilon/2)}\|v\| \ e^{-(\chi_i - \epsilon)n - \epsilon k}}{\frac{e^{k\chi_i}\|v\|}{K_1(\epsilon/2)(R_{\epsilon/2}(x))^2 e^{k\epsilon/2}}}$$
$$= K_1(\epsilon/2)(R_{\epsilon/2}(x))^3 e^{(n+k)\epsilon/2} e^{n\epsilon - k\epsilon} e^{k\epsilon/2}$$
$$\leq K_1(\epsilon/2)(R_{\epsilon/2}(x))^3.$$

Case 5. $k \ge 0$ and n + k < 0. Then

$$\frac{\|df_x^{n+k}(v)\| e^{-(\chi_i - \epsilon)n - \epsilon|k|}}{\|df_x^k(v)\|} \leq \frac{R_{\epsilon/2}(x)e^{(n+k)(\chi_i - \epsilon/2)}\|v\| e^{-(\chi_i - \epsilon)n - \epsilon k}}{\frac{e^{k\chi_i}\|v\|}{K_1(\epsilon/2)(R_{\epsilon/2}(x))^2 e^{k\epsilon/2}}} \\ = K_1(\epsilon/2)(R_{\epsilon/2}(x))^3 e^{-(n+k)\epsilon/2} e^{n\epsilon - k\epsilon} e^{k\epsilon/2} \\ \leq K_1(\epsilon/2)(R_{\epsilon/2}(x))^3.$$

Case 6. k < 0, and so n + k < 0. Then

$$\frac{\|df_x^{n+k}(v)\| e^{-(\chi_i - \epsilon)n - \epsilon|k|}}{\|df_x^k(v)\|} \leq \frac{R_{\epsilon/2}(x)e^{(n+k)(\chi_i - \epsilon/2)}\|v\| e^{-(\chi_i - \epsilon)n + \epsilon k}}{\frac{e^{k\chi_i}\|v\|}{K_1(\epsilon/2)(R_{\epsilon/2}(x))^2 e^{-k\epsilon/2}}}$$
$$= K_1(\epsilon/2)(R_{\epsilon/2}(x))^3 e^{-(n+k)\epsilon/2} e^{n\epsilon + k\epsilon} e^{-k\epsilon/2}$$
$$\leq K_1(\epsilon/2)(R_{\epsilon/2}(x))^3.$$

This proves (6.8). Combining the latter with (6.7) gives $A_{\epsilon}(x) \leq K_1(\epsilon/2)(R_{\epsilon/2}(x))^3$, and therefore

$$K_2(\epsilon)(A_{2\epsilon}(x))^{1/3} \le R_{\epsilon}(x) \le A_{\epsilon}(x) \quad , \quad x \in \mathcal{L}'_0,$$

where $K_2(\epsilon) = \frac{\epsilon^{1/6}}{C_0^{2/3}(2k_0)^{1/6}}$.

Proof of (3.5). Given $i = 1, ..., k_0$ and $x \in \mathcal{L}'_0$, there exist an integer $n \ge 0$ and $v \in E^{(i)}$ with ||v|| = 1 such that $\widetilde{R}^+_{\epsilon,i}(x) = \frac{||df_x^n(v)||}{e^{(\chi_i+\epsilon)}}$. Let $v = \sum_{k=1}^i v_k$ with $v_k \in E_k(x)$ for all k. Then by (6.3), $||v_k|| \le K_0(\epsilon)R_\epsilon(x)||v|| \le K_0(\epsilon)R_\epsilon(x)$ for all k. Using this and $\chi_i \ge \chi_k$ for $k \le i$ yields

$$\begin{aligned} \widetilde{R}_{\epsilon,i}^{+}(x) &= \frac{\|df_{x}^{n}(v)\|}{e^{(\chi_{i}+\epsilon)n}} \leq \frac{\sum_{k=1}^{i} \|df_{x}^{n}(v_{k})\|}{e^{(\chi_{i}+\epsilon)n}} \leq \sum_{k=1}^{i} \frac{\|df_{x}^{n}(v_{k})\|}{e^{(\chi_{k}+\epsilon)n}} \\ &\leq \sum_{k=1}^{i} \frac{\|(df_{x}^{n})|_{E_{k}}\| \|v_{k}\|}{e^{(\chi_{k}+\epsilon)n}} \leq \sum_{k=1}^{i} R_{\epsilon,k}^{+}(x) K_{0}(\epsilon)R_{\epsilon}(x) \leq k_{0}K_{0}(\epsilon) (R_{\epsilon}(x))^{2}. \end{aligned}$$

This is true for all i, so $\widetilde{R}_{\epsilon}^+(x) \leq k_0 K_0(\epsilon) (R_{\epsilon}(x))^2$. In a similar way, considering the flow ϕ_{-t} and the map $f^{-1} = \phi_{-1}$ one shows that $\widetilde{R}_{\epsilon}^-(x) \leq k_0 K_0(\epsilon) (R_{\epsilon}(x))^2$. This proves (3.5).

References

- [BaL] V. Baladi and C. Liverani, Exponential decay of correlations for piecewise cone hyperbolic contact flows, Comm. Math. Phys. 314 (2012), 689-773.
- [BP] L. Barreira and Ya. Pesin, Lyapunov exponents and smooth ergodic theory, University Lecture Series 23, American Mathematical Society, Providence, RI, 2001.
- [FHY] A. Fathi, M.R.Herman and J.C.Yoccoz, A proof of Pesin's stable manifold theorem, Lect. Notes in Math. Springer, Vol. 1007 (1983), 177-215.
- [BPS] L. Barreira, Ya. Pesin and J. Schmeling, Dimension and product structure of hyperbolic measures, Ann. of Math. 149 (1999), 755-783.
- [BSau] L. Barreira, B. Saussol, Multifractal analysis of hyperbolic flows, Comm. Math. Phys. 214 (2000), 339-371.
- [BS] L. Barreira, J. Schmeling, Sets of "non-typical" points have full topological entropy and full Hausdorff dimension, Israel J. Math. 116 (2000), 29-70.
- [B] R. Bowen, Symbolic dynamics for hyperbolic flows, Amer. J. Math. 95 (1973), 429-460.
- [BR] R. Bowen and D. Ruelle, The ergodic theory of Axiom A flows, Invent. Math. 29, 181-202 (1975)
- [D] D. Dolgopyat, On decay of correlations in Anosov flows, Ann. of Math. 147 (1998), 357-390.
- [KH] A. Katok and B. Hasselblatt, Introduction to the Modern Theory of Dynamical Systems, Cambridge Univ. Press, Cambridge 1995.
- [KM] A. Katok and L. Mendoza, Dynamical systems with nonuniformly hyperbolic behaviour, Appendix S in [KH].
- [Ki] Yu. Kifer, Large deviations in dynamical systems and stochastic processes, Trans. Amer. Math. Soc. 321 (1990), 505-524.
- [Kin] J.F.C. Kingsman, The ergodic theory of sub-additive stochastic processes, J. Royal. Stat. Soc. B 30 (1968), 499-510.
- [LY1] F. Ledrappier and L.-S. Young, The metric theory of diffeomorphisms: Part I: Characterization of measures satisfying Pesin's entropy formula, Ann. of Math. 122 (1985), 540-574.
- [LY2] F. Ledrappier and L.-S. Young, The metric theory of diffeomorphisms: Part II: Relations between entropy, exponents and dimension, Ann. of Math. 122 (1985), 509-539.
- [L] C. Liverani, On contact Anosov flows, Ann. of Math. 159 (2004), 1275-1312.
- [MN] I. Melbourne and M. Nicol, Large deviations for non-uniformly hyperbolic systems, Trans. Amer. Math. Soc. 360 (2008), 6661-6676.
- [OP] S. Orey and S. Pelikan, Deviation of trajectory averages and the defect in Pesin's formula for Anosov diffeomorphisms, Trans. Amer. Math. Soc. 315 (1989), 741-753.
- [Os] V. I. Oseledets, A multiplicative ergodic theorem. Lyapunov characteristic numbers for dynamical systems, Trans. Moscow Math. Soc. 19 (1968), 197-221.
- [PP] W. Parry and M. Pollicott, Zeta functions and the periodic orbit structure of hyperbolic dynamics, Astérisque 187-188, (1990).
- [P1] Ya. Pesin, Characteristic exponents and smooth ergodic theory, Russian Mathematical Surveys 32 (1977), 55-114.
- [P2] Ya. Pesin, Dimension theory in dynamical systems: Contemporary views and applications, The University of Chicago Press, Chicago, 1997.
- [P3] Ya. Pesin, Lectures on partial hyperbolicity and stable ergodicity, Zürich Lectures in Advanced Math., Zürich 2004.
- [PSa] Ya. Pesin, V. Sadovskaya, Multifractal analysis of conformal Axiom A flows, Comm. Math. Phys. 216 (2001), 277-312.
- [PS] C. Pugh and M. Shub, Ergodic attractors, Trans. Amer. Math. Soc. 312 (1989), 1-54.
- [R] D. Ruelle, Ergodic theory of differentiable dynamical systems, Publ. Math. de l'IHES 50 (1979), 27-58.
- [Sim] S. N. Simic, Oseldets regularity functions for Anosov flows, Comm. Math. Phys. 305 (2011), 1-21.
- [St] L. Stoyanov, Spectra of Ruelle transfer operators for Axiom A flows, Nonlinearity 24 (2011), 1089-1120.
- [Y1] L.-S. Young, Large deviations in dynamical systems, Trans. Amer. Math. Soc. 318 (1990), 525-543.
- [Y2] L.-S. Young, Statistical properties of systems with some hyperbolicity including certain billiards, Ann. Math. 147 (1998) 585-650.

UNIVERSITY OF WESTERN AUSTRALIA, CRAWLEY WA 6009, AUSTRALIA *E-mail address:* luchezar.stoyanov@uwa.edu.au