

ON THE GROWTH RATE OF IDEAL COXETER GROUPS IN HYPERBOLIC 3-SPACE

YOHEI KOMORI AND TOMOSHIGE YUKITA

ABSTRACT. We study the set \mathcal{G} of growth rates of ideal Coxeter groups in hyperbolic 3-space which consists of real algebraic integers greater than 1. We show that (1) \mathcal{G} is unbounded above while it has the minimum, (2) any element of \mathcal{G} is a Perron number, and (3) growth rates of ideal Coxeter groups with n generators are located in the closed interval $[n - 3, n - 1]$.

1. INTRODUCTION

The upper half space $\mathbb{H}^3 = \{x = (x_1, x_2, x_3) \in \mathbb{R}^3 \mid x_3 > 0\}$ with the metric $|dx|/x_3$ is a model of hyperbolic 3-space, so called the *upper half space model*. The Euclidean plane $\mathbb{E}^2 = \{x = (x_1, x_2, x_3) \in \mathbb{R}^3 \mid x_3 = 0\}$ and the point at infinity ∞ consist of the boundary at infinity $\partial\mathbb{H}^3$ of \mathbb{H}^3 . A subset $B \subset \mathbb{H}^3$ is called a *hyperplane* of \mathbb{H}^3 if it is a Euclidean hemisphere or a half plane orthogonal to \mathbb{E}^2 . When we restrict the hyperbolic metric $|dx|/x_3$ of \mathbb{H}^3 to B , it becomes a model of hyperbolic plane. We define a *polytope* as a closed domain P of \mathbb{H}^3 which can be written as the intersection of finitely many closed half spaces H_B bounded by hyperplanes B , say $P = \bigcap H_B$. In this presentation of P , we assume that $F_B = P \cap B$ is a hyperbolic polygon of B . F_B is called a *facet* of P , and B is called the *supporting hyperplane* of F_B . If the intersection of two facets F_{B_1} and F_{B_2} of P consists of a geodesic segment, it is called an *edge* of P , while if the intersection $\bigcap F_B$ of more than two facets is a point, then it is called a *vertex* of P . If F_{B_1} and F_{B_2} intersect only at a point of the boundary $\partial\mathbb{H}^3$ of \mathbb{H}^3 , it is called an *ideal vertex* of P . A polytope P is called *ideal* if all of its vertices are ideal. Related to Jakob Steiner's problem on the combinatorial characterization of polytopes inscribed in the two-sphere S^2 , ideal polytopes in hyperbolic 3-space \mathbb{H}^3 has been studied extensively [2, 8].

A *horosphere* Σ of \mathbb{H}^3 based at $v \in \partial\mathbb{H}^3$ is defined by a Euclidean sphere in \mathbb{H}^3 tangent to \mathbb{E}^2 at v when $v \in \mathbb{E}^2$, or a Euclidean plane in \mathbb{H}^3 parallel to \mathbb{E}^2 when $v = \infty$. When we restrict the hyperbolic metric of \mathbb{H}^3 to Σ , it becomes a model of Euclidean plane. Let $v \in \partial\mathbb{H}^3$ be an ideal vertex of a polytope P in \mathbb{H}^3 and Σ be a horosphere of \mathbb{H}^3 based at v such that Σ meets just the facets of P incident with v . Then the *vertex link* $L(v) := P \cap \Sigma$ of v in P is a Euclidean convex polygon in the horosphere Σ . If F_{B_1} and F_{B_2} are adjacent facets of P incident with v , then the Euclidean dihedral angle between $F_{B_1} \cap \Sigma$ and $F_{B_2} \cap \Sigma$ in Σ is equal to the hyperbolic dihedral angle between the supporting hyperplanes B_1 and B_2 in \mathbb{H}^3 (cf. [8, Theorem 6.4.5]).

2010 *Mathematics Subject Classification*. Primary 20F55, Secondary 20F65.

Key words and phrases. Coxeter group; growth function; growth rate; Perron number.

In this paper we consider ideal *Coxeter* polytopes P , which mean that the dihedral angles of edges of P are submultiples of π . Since any Euclidean Coxeter polygon is a rectangle or a triangle with dihedral angles $(\pi/2, \pi/3, \pi/6)$, $(\pi/2, \pi/4, \pi/4)$ or $(\pi/3, \pi/3, \pi/3)$, we see that the dihedral angles of an ideal Coxeter polytope must be $\pi/2, \pi/3, \pi/4$ or $\pi/6$.

Any Coxeter polytope P is a fundamental domain of the discrete group Γ generated by the set S consisting of the reflections with respects to its facets. We call (Γ, S) the *Coxeter system* associated to P . In this situation we can define the *word length* $\ell_S(x)$ of $x \in \Gamma$ with respect to S by the smallest integer $k \geq 0$ for which there exist $s_1, s_2, \dots, s_k \in S$ such that $x = s_1 s_2 \cdots s_k$. The *growth function* $f_S(t)$ of (Γ, S) is the formal power series $\sum_{k=0}^{\infty} a_k t^k$ where a_k is the number of elements $g \in \Gamma$ satisfying $\ell_S(g) = k$. It is known that the *growth rate* of (Γ, S) , $\tau = \limsup_{k \rightarrow \infty} \sqrt[k]{a_k}$ is bigger than 1 ([3]) and less than or equal to the cardinality $|S|$ of S from the definition. By means of Cauchy-Hadamard formula, the radius of convergence R of $f_S(t)$ is the reciprocal of τ , i.e. $1/|S| \leq R < 1$. In practice the growth function $f_S(t)$ which is analytic on $|t| < R$ extends to a rational function $P(t)/Q(t)$ on \mathbb{C} by analytic continuation where $P(t), Q(t) \in \mathbb{Z}[t]$ are relatively prime. There are formulas due to Solomon and Steinberg to calculate the rational function $P(t)/Q(t)$ from the data of finite Coxeter subgroups of (Γ, S) ([11, 12]. See also [4]).

Theorem 1. (Solomon's formula)

The growth function $f_S(t)$ of an irreducible finite Coxeter group (Γ, S) can be written as $f_S(t) = [m_1+1, m_2+1, \dots, m_k+1]$ where $[n] = 1+t+\dots+t^{n-1}$, $[m, n] = [m][n]$, etc. and $\{m_1, m_2, \dots, m_k\}$ is the set of exponents of (Γ, S) .

Theorem 2. (Steinberg's formula)

Let (Γ, S) be a hyperbolic Coxeter group. Let us denote the Coxeter subgroup of (Γ, S) generated by the subset $T \subseteq S$ by (Γ_T, T) , and denote its growth function by $f_T(t)$. Set $\mathcal{F} = \{T \subseteq S : \Gamma_T \text{ is finite}\}$. Then

$$\frac{1}{f_S(t^{-1})} = \sum_{T \in \mathcal{F}} \frac{(-1)^{|T|}}{f_T(t)}.$$

In this case, $t = R$ is a pole of $f_S(t) = P(t)/Q(t)$. Hence R is a real zero of the denominator $Q(t)$ closest to the origin $0 \in \mathbb{C}$ of all zeros of $Q(t)$. Solomon's formula implies that $P(0) = 1$. Hence $a_0 = 1$ means that $Q(0) = 1$. Therefore $\tau > 1$, the reciprocal of R , becomes a real algebraic integer whose conjugates have moduli less than or equal to the modulus of τ . If $t = R$ is the unique zero of $Q(t)$ with the smallest modulus, then $\tau > 1$ is a real algebraic integer whose conjugates have moduli less than the modulus of τ : such a real algebraic integer is called a *Perron number*.

The following result is a criterion for growth rates to be Perron numbers.

Proposition 1. ([6], Lemma1)

Consider the following polynomial of degree $n \geq 2$

$$g(t) = \sum_{k=1}^n a_k t^k - 1,$$

where a_k is a non-negative integer. We also assume that the greatest common divisor of $\{k \in \mathbb{N} \mid a_k \neq 0\}$ is 1. Then there is a real number r_0 , $0 < r_0 < 1$ which is the unique zero of $g(t)$ having the smallest absolute value of all zeros of $g(t)$.

In this paper we study the set \mathcal{G} of growth rates of ideal Coxeter polytopes in hyperbolic 3-space. We will show that (1) \mathcal{G} is unbounded above while it has the minimum, (2) any element of \mathcal{G} is a Perron number, and (3) growth rates of ideal Coxeter polytopes with n facets are located in the closed interval $[n - 3, n - 1]$.

2. IDEAL COXETER POLYTOPES WITH 4 OR 5 FACETS IN \mathbb{H}^3

Let p, q, r and s be the number of edges with dihedral angles $\pi/2$, $\pi/3$, $\pi/4$, and $\pi/6$ of an ideal Coxeter polytope P in \mathbb{H}^3 . By Andreev theorem [1], we can classify ideal Coxeter polytopes with 4 or 5 facets, and calculate the growth functions $f_S(t)$ of P by means of Steinberg's formula and also growth rates, see Table 1. Every denominator polynomial has a form $(t - 1)H(t)$ and all coefficients of $H(t)$ satisfies the condition of Proposition 1, so that the growth rates of ideal Coxeter polytopes with 4 or 5 facets are Perron numbers.

(p, q, r, s)	combinatorial type	denominator polynomial	growth rate
$(2, 2, 0, 2)$	simplex	$(t - 1)(3t^5 + t^4 + t^3 + t^2 + t - 1)$	2.03074
$(2, 0, 4, 0)$	simplex	$(t - 1)(3t^3 + t^2 + t - 1)$	2.13040
$(0, 6, 0, 0)$	simplex	$(t - 1)(3t^2 + t - 1)$	2.30277
$(4, 2, 0, 2)$	pyramid	$(t - 1)(4t^5 + t^4 + 2t^3 + t^2 + 2t - 1)$	2.74738
$(4, 0, 4, 0)$	pyramid	$(t - 1)(4t^3 + t^2 + 2t - 1)$	2.84547
$(2, 5, 0, 2)$	prism	$(t - 1)(5t^5 + 2t^4 + t^3 + 3t^2 + 2t - 1)$	3.16205

TABLE 1.




FIGURE 1.

After glueing m copies of the ideal Coxeter pyramid with $p = r = 4$ along their sides successively, we can construct a hyperbolic ideal Coxeter polytope P with $n = m + 4$ facets. In Figure 1 we are looking at the ideal Coxeter polytope P with 8 facets from the point at infinity ∞ , which consists of 4 copies of ideal Coxeter pyramid with $p = r = 4$ whose apexes are located at ∞ ; squares represent bases of pyramids and disks are supporting hyperplanes of these bases. The growth function of P has the following denominator polynomial

$$(t - 1)H(t) = (t - 1)(2(n - 3)t^3 + (n - 4)t^2 + (n - 3)t - 1).$$

Then all coefficients of $H(t)$ except its constant term are non-negative. Therefore we can apply Proposition 1 to conclude that the growth rate of P is a Perron

number. Moreover $H(t)$ has a unique zero on the unit interval $[0, 1]$ and the following inequalities hold:

$$H\left(\frac{1}{n-3}\right) = \frac{n-2}{(n-3)^2} > 0, \quad H\left(\frac{1}{n-1}\right) = \frac{-n^2+n-4}{(n-1)^3} < 0.$$

They imply that the growth rate τ of P with n facets satisfies

$$n-3 \leq \tau \leq n-1,$$

which will be generalized in the next section. In particular

Proposition 2. *The set \mathcal{G} of growth rates of 3-dimensional hyperbolic ideal Coxeter polytopes is unbounded above.*

3. THE GROWTH RATES OF IDEAL COXETER POLYTOPES IN \mathbb{H}^3

Recall that p, q, r and s be the number of edges with dihedral angles $\pi/2, \pi/3, \pi/4,$ and $\pi/6$ of an ideal Coxeter polytope P in \mathbb{H}^3 . By means of Steinberg's formula, we can calculate the growth function $f_S(t)$ of P as

$$1/f_S(1/t) = 1 - n/[2] + p/[2, 2] + q/[2, 3] + r/[2, 4] + s/[2, 6],$$

where $[2, 3] = [2][3]$, etc. It can be rewritten as

$$\begin{aligned} 1/f_S(t) &= 1 - nt/[2] + pt^2/[2, 2] + qt^3/[2, 3] + rt^4/[2, 4] + st^6/[2, 6] \\ &= \frac{1}{[2, 2, 3, 4, 6]}G(t), \end{aligned}$$

where

$$G(t) = [2, 2, 3, 4, 6] - nt[2, 3, 4, 6] + pt^2[3, 4, 6] + qt^3[2, 4, 6] + rt^4[2, 3, 6] + st^6[2, 3, 4].$$

Proposition 3. *Put $a = p/2, b = q/3, c = r/4, d = s/6$. Then*

$$(1) \quad a + b + c + d = n - 2.$$

Proof. By a result of Serre ([10]. See also [4])

$$G(1) = [2, 3, 4, 6](1)(2 - n + p/2 + q/3 + r/4 + s/6) = 0$$

□

By using this equality (1) we represent $H(t) = G(t)/(t-1)$ as

$$\begin{aligned} H(t) &= -[2, 3, 4, 6] + at[3, 4, 6] + bt(2t+1)[2, 4, 6] \\ &\quad + ct(3t^2+2t+1)[2, 3, 6] + dt(5t^4+4t^3+3t^2+2t+1)[2, 3, 4] \\ &= -1 + (-4+a+b+c+d)t + (-9+3a+5b+5c+5d)t^2 \\ &\quad + (-15+6a+11b+14c+14d)t^3 + (-20+9a+17b+25c+29d)t^4 \\ &\quad + (-23+11a+22b+33c+49d)t^5 + (-23+12a+24b+36c+66d)t^6 \\ &\quad + (-20+11a+23b+35c+71d)t^7 + (-15+9a+19b+31c+61d)t^8 \\ &\quad + (-9+6a+13b+22c+40d)t^9 + (-4+3a+7b+11c+19d)t^{10} \\ &\quad + (-1+a+2b+3c+5d)t^{11}. \end{aligned}$$

From this formula we have the following result (see also [7], Theorem 3).

Theorem 3. *The growth rates of ideal Coxeter polytopes in \mathbb{H}^3 are Perron numbers.*

Proof. When n the number of facets satisfies $n \geq 6$, the equality (1) of Proposition 3 implies $a + b + c + d = n - 2 \geq 4$. Then all coefficients of $H(t)$ except its constant term are non-negative. Hence Proposition 1 implies the assertion. For $n = 4, 5$, this claim was already proved in the previous section. \square

Moreover the equality (1) induces the following two functions $H_1(t)$ and $H_2(t)$ satisfying $H_1(t) \leq H(t) \leq H_2(t)$ for any $t > 0$:

$$\begin{aligned} H_1(t) &= -1 + (-4 + (n-2)t + (-9 + 3(n-2))t^2 + (-15 + 6(n-2))t^3 \\ &\quad + (-20 + 9(n-2))t^4 + (-23 + 11(n-2))t^5 + (-23 + 12(n-2))t^6 \\ &\quad + (-20 + 11(n-2))t^7 + (-15 + 9(n-2))t^8 + (-9 + 6(n-2))t^9 \\ &\quad + (-4 + 3(n-2))t^{10} + (-1 + (n-2))t^{11} \\ &= (1+t)^2(-1-3t+nt)(1+t^2)(1-t+t^2)(1+t+t^2)^2, \end{aligned}$$

$$\begin{aligned} H_2(t) &= -1 + (-4 + (n-2)t + (-9 + 5(n-2))t^2 + (-15 + 14(n-2))t^3 \\ &\quad + (-20 + 29(n-2))t^4 + (-23 + 49(n-2))t^5 + (-23 + 66(n-2))t^6 \\ &\quad + (-20 + 71(n-2))t^7 + (-15 + 61(n-2))t^8 + (-9 + 40(n-2))t^9 \\ &\quad + (-4 + 19(n-2))t^{10} + (-1 + 5(n-2))t^{11} \\ &= (1+t)^2(1+t^2)(1+t+t^2) \\ &\quad (-1-3t+nt-5t^2+2nt^2-7t^3+3nt^3-9t^4+4nt^4-11t^5+5nt^5). \end{aligned}$$

Now we assume that $n \geq 6$. Then all coefficients of $H_1(t)$ and $H_2(t)$ except their constant terms are non-negative so that they have unique zeros in $(0, \infty)$. The following inequalities

$$H_1\left(\frac{1}{n-3}\right) = 0, \quad H_2\left(\frac{1}{n-1}\right) = -\frac{6}{(n-1)^5} < 0$$

guarantee that the zero of $H(t)$ is located in $[\frac{1}{n-1}, \frac{1}{n-3}]$. Combining with the similar result for $n = 4, 5$ in the previous section, we have the following theorem which is our main result.

Theorem 4. *The growth rate τ of an ideal Coxeter polytope with n facets in \mathbb{H}^3 satisfies*

$$(2) \quad n - 3 \leq \tau \leq n - 1.$$

Corollary 1. *An ideal Coxeter polytope P with n facets in \mathbb{H}^3 is right-angled if and only if its growth rate τ is equal to $n - 3$.*

Proof. The factor $H(t)$ of the denominator polynomial $G(t) = (t-1)H(t)$ of the growth function of P is equal to $H_1(t)$ if and only if $b = c = d = 0$, which means that all dihedral angles are $\pi/2$. \square

From the inequality (2), we see that the growth rate τ of an ideal Coxeter polytope with n facets with $n \geq 6$ satisfies $\tau \geq 3$. Therefore combining with the result of growth rates for $n = 4, 5$ shown in the previous section, we also have the following corollary (see also [7], Theorem 4).

Corollary 2. *The minimum of the growth rates of 3-dimensional hyperbolic ideal Coxeter polytopes is $0.492432^{-1} = 2.03074$, which is uniquely realized by the ideal simplex with $p = q = s = 2$.*

4. ACKNOWLEDGEMENT

The authors thank Jun Nonaka for explaining his paper [7].

REFERENCES

- [1] E. M. Andreev, On convex polyhedra of finite volume in Lobachevskij space, *Mat. Sb.*, Nov. Ser. **83** (1970), 256-260. English transl.: *Math. USSR, Sb.* **12** (1971), 255- 259.
- [2] F. Gueritaud, On an Elementary Proof of Rivin’s Characterization of Convex Ideal Hyperbolic Polyhedra by their Dihedral Angles, *Geometriae Dedicata* 108 (2004), 111–124.
- [3] P. de la Harpe, Groupes de Coxeter infinis non affines, *Exposition. Math* 5 (1987), 91–96.
- [4] G. J. Heckman, The volume of hyperbolic Coxeter polytopes of even dimension, *Indag. Math. (N.S.)* 6 (1995), 189-196.
- [5] J. E. Humphreys, *Reflection groups and Coxeter groups*, Cambridge Studies in Advanced Mathematics, **29**, Cambridge Univ. Press, Cambridge, 1990.
- [6] Y. Komori and Y. Umemoto, On the growth of hyperbolic 3-dimensional generalized simplex reflection groups, *Proc. Japan Acad. Ser. A Math. Sci.* Volume **88**, Number 4 (2012), 62–65.
- [7] J. Nonaka, The growth rates of ideal Coxeter polyhedra in hyperbolic 3-space, arXiv:1504.06718.
- [8] I. Rivin, A characterization of ideal polyhedra in hyperbolic 3-space, *Ann. of Math.* 143 (1996), 5–70.
- [9] J. G. Ratcliffe, *Foundations of hyperbolic manifolds*, Grad. Texts in Math., **149**, Springer, New York, 1994.
- [10] J. P. Serre, Cohomologie des groupes discrets, *Prospects Math.*, *Ann. Math. Stud.* 70, 77-169, 1971.
- [11] L. Solomon, The orders of the finite Chevalley groups, *J. Algebra* **3**(1966), 376–393.
- [12] R. Steinberg, *Endomorphisms of linear algebraic groups*, *Memoirs of the American Mathematical Society*, No. 80, Amer. Math. Soc., Providence, RI, 1968.

DEPARTMENT OF MATHEMATICS, SCHOOL OF EDUCATION, WASEDA UNIVERSITY, NISHI-WASEDA 1-6-1, SHINJUKU, TOKYO 169-8050, JAPAN
E-mail address: ykomori@waseda.jp

DEPARTMENT OF MATHEMATICS, SCHOOL OF EDUCATION, WASEDA UNIVERSITY, NISHI-WASEDA 1-6-1, SHINJUKU, TOKYO 169-8050, JAPAN
E-mail address: yshigetomo@suou.waseda.jp