# Saddle point inflation in string-inspired theory

Yuta Hamada,<sup>\*</sup> Hikaru Kawai,<sup>†</sup> and Kiyoharu Kawana<sup>‡</sup>

Department of Physics, Kyoto University, Kyoto 606-8502, Japan

June 2, 2021

#### Abstract

The observed value of the Higgs mass indicates the possibility that there is no supersymmetry below the Planck scale and that the Higgs can play the role of the inflaton. We examine the general structure of the saddle point inflation in string-inspired theory without supersymmetry. We point out that the string scale is fixed to be around the GUT scale  $\sim 10^{16}$  GeV in order to realize successful inflation. We find that the inflaton can be naturally identified with the Higgs field.

<sup>\*</sup>E-mail: hamada@gauge.scphys.kyoto-u.ac.jp

<sup>&</sup>lt;sup>†</sup>E-mail: hkawai@gauge.scphys.kyoto-u.ac.jp

<sup>&</sup>lt;sup>‡</sup>E-mail: kiyokawa@gauge.scphys.kyoto-u.ac.jp

# 1 Introduction

The recently observed particle at ATLAS [1] and CMS [2] experiments at the Large Hadron Collider (LHC) is consistent with the Standard Model (SM) Higgs with the mass around 125 GeV. Up to now, there has been observed no significant deviation from the SM nor a hint of new physics. Once the Higgs mass is determined, we have fixed all the parameters in the SM and can extrapolate it up to its ultraviolet (UV) cutoff scale. In particular, the quadratically divergent bare Higgs mass is found to be suppressed when the UV cutoff is at around the Planck scale [3], see also Ref. [4]. Furthermore, the quartic Higgs coupling becomes tiny at the same time, see e.g Refs. [3, 5]. This opens up the possibilities of identifying the Higgs field as the inflaton [6, 7], and of the absence of supersymmetry below the Planck scale. Although non-supersymmetric vacua are ubiquitous in string theory [8, 9], their phenomenology has not been well studied. It becomes important to explore the phenomenology starting from non-supersymmetric theory.

In this letter, we consider the saddle point inflation scenario starting in string-inspired theory without supersymmetry. The potential is generated perturtabatively in contrast to the supersymmetric case where the potential comes only non-perturbatively. Then, we calculate the cosmological parameters by assuming that the potential is tuned in such a way that the first n derivatives vanish at some point. The predicted cosmological parameters are consistent with the recent Planck 2015 result [11]. Furthermore, we can estimate the order of the string scale from the height of the potential that is roughly given by the string scale to the fourth multiplied by the rather small ten-dimensional one-loop factor.

To realize the saddle point, some amount of the fine-tuning is needed. This fine-tuning would be achieved by some principles which are beyond the ordinary local field theory, e.g. the multiple point criticality principle [13] and the maximum entropy principle [14].

This letter is organized as follows. In the next section, we consider the potential that has a saddle point where the first n derivatives vanish. Then we calculate the cosmological parameters of the model. In Sec. 3, we estimate the stirng scale in the case of the non-supersymmetric heterotic-like string model. In Sec. 4, we summarize our result.

# 2 Saddle point inflation and observables

We start with a general potential V as a function of an inflaton field  $\varphi$ . We will discuss the possibility of identifying it as the SM Higgs in the next section.

We expand the potential around the saddle point  $\varphi_c$  as  $\varphi = \varphi_c + \delta \varphi$ :

$$V = \sum_{n=0}^{\infty} \frac{V_c^{(n)}}{n!} \delta \varphi^n = V_c + V_c' \,\delta \varphi + \frac{V_c''}{2} \,\delta \varphi^2 + \frac{V_c'''}{3!} \,\delta \varphi^3 + \cdots \,. \tag{1}$$

We assume that the first  $n \geq 2$  derivatives vanish at  $\varphi_c$ :

$$V'_c = V''_c = \dots = V_c^{(n)} = 0.$$
 (2)

Here, we also assume  $V_c^{(n+1)} > 0$  (< 0) for even (odd) n so that  $\varphi$  rolls down from  $\varphi_c$  towards 0. This is because we are going to identify  $\varphi$  as the Higgs field.

The slow roll parameters around the saddle point are obtained as  $^1$ 

$$\epsilon := \frac{M_P^2}{2} \left(\frac{V'}{V}\right)^2 = \frac{M_P^2 \left(V_c^{(n+1)}\right)^2}{2(n!)^2 V_c^2} \,\delta\varphi^{2n} + O\left(\delta\varphi^{2n+1}\right),\tag{3}$$

$$\eta := M_P^2 \frac{V''}{V} = \frac{M_P^2 V_c^{(n+1)}}{(n-1)! V_c} \delta \varphi^{n-1} + O(\delta \varphi^n), \qquad (4)$$

$$\zeta^{2} := M_{P}^{4} \frac{V'''V'}{V^{2}} = \frac{M_{P}^{4} \left( V_{c}^{(\mathsf{n}+1)} \right)^{2}}{(\mathsf{n}-2)! \, \mathsf{n}! \, V_{c}^{2}} \, \delta\varphi^{2\mathsf{n}-2} + O\left(\delta\varphi^{2\mathsf{n}-1}\right). \tag{5}$$

We see that  $\epsilon \ll |\eta|, \zeta^2$  for  $\delta \varphi \ll M_P$ . The inflation ends when  $\epsilon$  becomes of order unity, and we define its end point by  $\epsilon(\delta \varphi_{end}) = 1$  to get

$$(\delta\varphi_{\rm end})^{\mathsf{n}} \simeq \frac{\sqrt{2}\,\mathsf{n}!\,V_c}{M_P V_c^{(\mathsf{n}+1)}}.\tag{6}$$

The e-folding number N from a given stage of the inflation  $\varphi = \varphi_c + \delta \varphi$  to its end  $\varphi_{\text{end}} = \varphi_c + \delta \varphi_{\text{end}}$  is <sup>2</sup>

$$N = \int_{\varphi_{\text{end}}}^{\varphi} \frac{\mathrm{d}\varphi}{M_P^2} \frac{V}{V'} = \frac{\mathsf{n}!}{(\mathsf{n}-1)} \frac{V_c}{M_P^2 V_c^{(\mathsf{n}+1)}} \left[ \frac{1}{(\delta\varphi_{\text{end}})^{\mathsf{n}-1}} - \frac{1}{(\delta\varphi)^{\mathsf{n}-1}} \right] \\ \simeq \frac{\mathsf{n}!}{(\mathsf{n}-1)} \frac{V_c}{M_P^2 \left| V_c^{(\mathsf{n}+1)} \right| \left| \delta\varphi \right|^{\mathsf{n}-1}}, \tag{7}$$

<sup>1</sup> If  $\varphi_c$  is the only mass scale of theory as the model we will consider in the next section, we have  $V_c^{(n+2)}/V_c^{(n+1)} \sim \varphi_c^{-1}$ . Therefore, the condition for the validity of neglecting the higher order terms is

$$\frac{V_c^{(n+2)}\delta\varphi^{n+2}}{(n+2)!} \left/ \frac{V_c^{(n+1)}\delta\varphi^{n+1}}{(n+1)!} \sim \frac{\delta\varphi}{\varphi_c} \ll 1.$$

 $^2$  The problem about the initial condition can be avoided by considering the eternal inflation scenario at the saddle point [7].

where we have assumed  $|\delta\varphi_{\rm end}| \gg |\delta\varphi|$  in the last step. From Eqs. (3)(4)(5)(7), we obtain

$$\epsilon = \frac{1}{2M_P^2} \left[ \frac{\mathbf{n}!}{\left(\mathbf{n}-1\right)^{\mathbf{n}}} \frac{1}{N^{\mathbf{n}}} \frac{V_c}{M_P^2 \left| V_c^{(\mathbf{n}+1)} \right|} \right]^{\frac{2}{\mathbf{n}-1}}, \quad \eta = -\frac{\mathbf{n}}{\left(\mathbf{n}-1\right)N}, \quad \zeta^2 = \frac{\mathbf{n}}{\left(\mathbf{n}-1\right)N^2}.$$
(8)

The cosmological observables, namely the scalar perturbation  $A_s$ , spectral index  $n_s$ , tensor-to-scalar ratio r, and running index  $dn_s/d \ln k^{-3}$ 

$$A_s = \frac{V}{24\pi^2 \epsilon M_P^4},\tag{9}$$

$$n_s = 1 - 6\epsilon + 2\eta \simeq 1 - \frac{2\mathsf{n}}{(\mathsf{n} - 1)N},$$
 (10)

$$r = 16\epsilon,\tag{11}$$

$$\frac{\mathrm{d}n_s}{\mathrm{d}\ln k} = -16\epsilon\eta + 24\epsilon^2 + 2\zeta^2 \simeq 2\frac{\mathsf{n}}{(\mathsf{n}-1)N^2},\tag{12}$$

are constrained by the Plnack 2015 data [11]

$$A_s \simeq 2.2 \times 10^{-9}, \quad 0.954 < n_s < 0.980, \quad r < 0.168, \quad -0.03 < \frac{\mathrm{d}n_s}{\mathrm{d}\ln k} < 0.007,$$
(13)

at the 95% CL.<sup>4</sup> The e-folding number

$$N_* = 62 - \ln\left(\frac{10^{16} \,\text{GeV}}{V_{\text{end}}^{1/4}}\right) \simeq 64 + \frac{1}{4}\ln\epsilon \tag{14}$$

corresponds to the stage of inflation observed by the Planck experiment. We note that this model gives a concave potential ,  $\eta < 0$ , which is favored by the recent Planck data.<sup>5</sup>

# 3 Saddle point inflation in string-inspired theory

In this section, we consider the saddle point inflation in the non-supersymmetric heterotic-like string model. Here we assume that the tree level potential of the

<sup>&</sup>lt;sup>3</sup> It appears that these quantities change their values discretely with n. This is because n is the number of fine-tunings. However, if we take the next order term into account, we can explicitly check that the limit of  $V_c^{(n+1)} \rightarrow 0$  continuously connects the case n to n + 1. Thus we can have fractional n effectively.

<sup>&</sup>lt;sup>4</sup> To give the most conservative bound, here we employ the constraint from the Planck TT+lowP data.

 $<sup>^{5}</sup>$  We thank the referee for pointing out this point.

inflaton is absent. This is realized if the inflaton comes from the extra component of the gauge field/metric, for example. Then the dominant contribution to the potential is the one loop correction, which is suppressed compared to the string scale by the loop factor:

$$\int \frac{d^d k}{(2\pi)^d} = \frac{S_{d-1}}{2(2\pi)^d} \int dk^2 (k^2)^{\frac{d}{2}-1} \sim \frac{S_{d-1}}{2(2\pi)^d} M_s^d.$$
 (15)

For d = 10, we obtain the following numerical value

$$C_{\text{loop}} \equiv \frac{S_{d-1}}{2(2\pi)^d} = \frac{2\pi^5}{\Gamma(5)} \frac{1}{2(2\pi)^{10}} \simeq 1.3 \times 10^{-7}.$$
 (16)

In fact, the 10 dimensional cosmological constant of  $SO(16) \times SO(16)$  heterotic string theory [12] is calculated as

$$\Lambda_{SO(16)\times SO(16)} \simeq 3.9 \times 10^{-6} M_s^{10}.$$
 (17)

Because we assume that the tree potential of the inflaton vanishes, the effective action below the string scale becomes

$$S = \frac{M_s^8}{g_s^2} \int d^{10}x \sqrt{g} A(\chi) \mathcal{R} + \frac{M_s^8}{g_s^2} \int d^{10}x \sqrt{g} B(\chi) (\partial \chi)^2 + C_{\text{loop}} M_s^{10} \int d^{10}x \sqrt{g} V(\chi) + \cdots$$
  
$$= \frac{M_s^8}{g_s^2} V_6 \int d^4x \sqrt{g} A(\chi) \mathcal{R} + \frac{M_s^8}{g_s^2} V_6 \int d^4x \sqrt{g} B(\chi) (\partial \chi)^2 + C_{\text{loop}} M_s^{10} V_6 \int d^4 \sqrt{g} V(\chi) + \cdots$$
  
(18)

Here  $\chi$  is the dimensionless inflaton field,  $g_s$  is the string coupling,  $V(\chi)$  is the one loop potential, and  $V_6$  is the compactification volume. Because  $M_s$  is the only mass scale of the theory,  $A(\chi)$ ,  $B(\chi)$  and  $V(\chi)$  should be functions of order one

$$A(\chi) = a_0 + a_2 g_s^2 \chi^2 + \cdots, \quad B(\chi) = b_0 + b_2 g_s^2 \chi^2 + \cdots, \quad V(\chi) = v_0 + v_2 g_s^2 \chi^2 + \cdots,$$
(19)

with  $a_i$ 's,  $b_i$ 's and  $v_i$ 's being order one constants. Next let us move to the Einstein frame. Namely, we redefine the metric in such a way that  $A(\chi)$  becomes 1. In the Einstein frame, we have

$$S = M_P^2 \int d^4x \sqrt{g} \mathcal{R} + M_P^2 \int d^4x \sqrt{g} C(\chi) (\partial \chi)^2 + C_{\text{loop}} g_s^2 M_P^2 M_s^2 \int d^4x \sqrt{g} U(\chi).$$
(20)

Here

$$M_P^2 = \frac{M_s^2}{g_s^2} (M_s^6 V_6), \quad C(\chi) = c_0 + c_2 g_s^2 \chi^2 + \cdots, \quad U = u_0 + u_2 g_s^2 \chi^2 + \cdots,$$
(21)

where  $c_i$ 's and  $u_i$ 's are order one constants. In terms of the dimensionless canonical field  $\phi$ , the action becomes

$$S = M_P^2 \int d^4x \sqrt{g} \,\mathcal{R} + M_P^2 \int d^4x \sqrt{g} \,(\partial\phi)^2 + C_{\text{loop}} g_s^2 M_P^2 M_s^2 \int d^4x \sqrt{g} \,W\left(\phi\right),\tag{22}$$

where  $W(\phi)$  is a function of order one.

The argument so far is quite general. In the following, we assume that the potential has a saddle point where the first n derivatives vanish as in Sec. 2. This may happen by some mechanism beyond the ordinary local field theory such as the multiple point criticality principle [13] and the maximum entropy principle [14]. Here, we take

$$W(\phi) = W_0 \left( 1 - \left( 1 - \frac{\phi}{\phi_c} \right)^{\mathsf{n}+1} \right), \tag{23}$$

as a simple possibility. We expect that  $\phi_c$  is the order one quantity. In terms of the canonical field  $\varphi = M_P \phi$ , the potential  $V(\varphi)$  becomes

$$V(\varphi) = C_{\text{loop}} g_s^2 M_P^2 M_s^2 \times W\left(\frac{\varphi}{M_P}\right), \qquad (24)$$

Then, from Eq. (8), we get

$$\epsilon = \frac{1}{2M_P^2} \left[ \frac{\mathsf{n}!}{(\mathsf{n}-1)^{\mathsf{n}}} \frac{1}{N^{\mathsf{n}}} \frac{V_c}{M_P^2 \left| V_c^{(\mathsf{n}+1)} \right|} \right]^{\frac{2}{\mathsf{n}-1}} \\ = \frac{1}{2} \left[ \frac{\mathsf{n}!}{(\mathsf{n}-1)^{\mathsf{n}}} \frac{1}{N^{\mathsf{n}}} \frac{\phi_c^{\mathsf{n}+1}}{(\mathsf{n}+1)!} \right]^{\frac{2}{\mathsf{n}-1}}.$$
(25)

Furthermore, Eq. (25) and the COBE normalization Eq. (9) fix the value of  $V_c = C_{\text{loop}} g_s^2 M_P^2 M_s^2 W_0$ , from which we can obtain the string scale. In Table 1, we present the predictions of the cosmological parameters taking  $C_{\text{loop}} = 10^{-7}$ , N = 60. From this table, we can see that  $n \ge 4$  is favored by the current observation Eq. (13). The tensor to scalar ratio is very small compared to the current limit provided that  $\phi_c$  is of order one. As we vary n from 2 to 6,  $g_s M_s$  takes from  $4 \times 10^{14} \text{ GeV}$  to  $3 \times 10^{15} \text{ GeV}$  for  $W_0 = 1$ . If  $g_s$  takes  $\mathcal{O}(0.1)$ , the result indicates that the  $M_s$  is around the GUT scale  $\sim 10^{16} \text{ GeV}$ .

Finally let us discuss the possibility of identifying the inflaton as the SM Higgs. The recent analysis shows that the Higgs potential for the large values of the Higgs field h is roughly given by

$$V_{\rm SM} \sim 10^{-6} h^4,$$
 (26)

n	$n_s$	$\epsilon$	$V_c/M_P^4$	$g_s M_s \sqrt{W_0}/M_P$
2	$0.933\cdots$	$4.3\times10^{-9}\phi_c^6$	$2.2 \times 10^{-15} \phi_c^6$	$1.5 \times 10^{-4}$
3	0.95	$7.2\times10^{-8}\phi_c^4$	$3.8 \times 10^{-14} \phi_c^4$	$6.1 \times 10^{-4}$
4	$0.955\cdots$	$1.7 \times 10^{-7} \phi_c^{10/3}$	$8.6 \times 10^{-14} \phi_c^{10/3}$	$9.3 \times 10^{-4}$
5	$0.95833\cdots$	$2.3\times10^{-7}\phi_c^3$	$1.2 \times 10^{-13} \phi_c^3$	$1.1 \times 10^{-3}$
6	0.96	$2.6 \times 10^{-7} \phi_c^{14/5}$	$1.4 \times 10^{-13} \phi_c^{14/5}$	$1.2 \times 10^{-3}$

Table 1: The predictions of cosmological parameters and the string scale for  $N = 60, C_{\text{loop}} = 10^{-7}$ .



Figure 1: Matching between Eq. (24) and Eq. (26).

in the SM [3, 5] and its simple extensions [15] when the top mass is around 171–172 GeV. We examine whether  $V_{\rm SM}$  can be connected to the potential V in Eq. (24) under the assumption that  $\varphi$  is identified as h. In Fig. 1, Eqs. (24)(26) are plotted. Here we take  $n = 4, \phi_c = 1, W_0 = 1$  as an example. One can see that two lines are crossed at around  $\varphi \simeq 10^{16}$  GeV, which we call  $\varphi_0$ . We interpret this as the indication that the potential is given by the SM at lower energies, and becomes stringy, Eq. (24), above the string scale  $\sim 10^{16}$  GeV. We also show  $\varphi_0$  as a function of  $\varphi_c = \phi_c M_P$  in Fig. 2.  $\varphi_0$  takes the order of  $10^{16}$  GeV for  $\varphi_c = \mathcal{O}(M_P)$ .

### 4 Summary

We have examined the possibility of the saddle point inflation in the context of non-supersymmetric string theory, which is ubiquitous and becomes more



Figure 2:  $\varphi_0$  as a function of  $\varphi_c$ .  $\varphi_0$  is the value of  $\varphi$  for which  $V_{\rm SM}$  equals V.

realistic in light of the recent LHC result. Contrary to supersymmetric theory, the potential is generated perturbatively. We have assumed that the potential of the inflaton is identically zero at the tree level, and it is radiatively generated by the loop effect. We have estimated the string scale that realizes a successful inflation assuming that the potential is tuned so that it has a saddle point where first n derivatives vanish. Interestingly, the string scale becomes around the GUT scale,  $\sim 10^{16}$ GeV, if the string coupling is  $\mathcal{O}(0.1)$ . Furthermore, we have found that it is reasonable to identify the inflaton as the Higgs field. It is interesting that, in addition to the LHC results, the scale of the inflation supports non-supersymmetric string theory.

### Acknowledgement

We thank Kin-ya Oda for useful discussions. This work is supported by the Grant-in-Aid for Japan Society for the Promotion of Science (JSPS) Fellows No.251107 (YH) and No.271771 (KK). H.K.s work is supported in part by the Grant-in-Aid for Scientific Research No. 22540277.

# References

[1] G. Aad et al. [ATLAS Collaboration], "Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC," Phys. Lett. B **716** (2012) 1 [arXiv:1207.7214 [hep-ex]].

- [2] S. Chatrchyan et al. [CMS Collaboration], "Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC," Phys. Lett. B 716 (2012) 30 [arXiv:1207.7235 [hep-ex]].
- [3] Y. Hamada, H. Kawai and K. y. Oda, "Bare Higgs mass at Planck scale," Phys. Rev. D 87 (2013) 5, 053009 [Phys. Rev. D 89 (2014) 5, 059901] [arXiv:1210.2538 [hep-ph]].
  F. Jegerlehner, "The Standard model as a low-energy effective theory: what is triggering the Higgs mechanism?," Acta Phys. Polon. B 45 (2014) 6, 1167 [arXiv:1304.7813 [hep-ph]].
  F. Jegerlehner, "The hierarchy problem of the electroweak Standard Model revisited," arXiv:1305.6652 [hep-ph].
  Y. Hamada, H. Kawai and K. y. Oda, "Bare Higgs mass and potential at
- [4] M. S. Al-sarhi, I. Jack and D. R. T. Jones, "Quadratic divergences in gauge theories," Z. Phys. C 55 (1992) 283.
  D. R. T. Jones, "Comment on Bare Higgs mass at Planck scale," Phys. Rev. D 88 (2013) 9, 098301 [arXiv:1309.7335 [hep-ph]].

ultraviolet cutoff," arXiv:1305.7055 [hep-ph].

 [5] M. Holthausen, K. S. Lim and M. Lindner, "Planck scale Boundary Conditions and the Higgs Mass," JHEP 1202 (2012) 037 [arXiv:1112.2415 [hep-ph]];

F. Bezrukov, M. Y. Kalmykov, B. A. Kniehl and M. Shaposhnikov, *"Higgs boson mass and new physics,"* JHEP **1210** (2012) 140 [arXiv:1205.2893 [hep-ph]].

G. Degrassi, S. Di Vita, J. Elias-Miro, J. R. Espinosa, G. F. Giudice,
G. Isidori and A. Strumia, "Higgs mass and vacuum stability in the Standard Model at NNLO," JHEP 1208 (2012) 098 [arXiv:1205.6497 [hep-ph]].
S. Alekhin, A. Djouadi and S. Moch, "The top quark and Higgs boson masses and the stability of the electroweak vacuum," Phys. Lett. B 716 (2012) 214 [arXiv:1207.0980 [hep-ph]].

I. Masina, "The Higgs boson and Top quark masses as tests of Electroweak Vacuum Stability," Phys. Rev. D 87 (2013) 5, 053001 [arXiv:1209.0393 [hep-ph]].

[6] F. L. Bezrukov and M. Shaposhnikov, "The Standard Model Higgs boson as the inflaton," Phys. Lett. B 659 (2008) 703 [arXiv:0710.3755 [hep-th]]. A. Salvio, "Higgs Inflation at NNLO after the Boson Discovery," Phys. Lett. B 727 (2013) 234 [arXiv:1308.2244 [hep-ph]].

Y. Hamada, H. Kawai and K. y. Oda, *"Minimal Higgs inflation,"* PTEP **2014** (2014) 023B02 [arXiv:1308.6651 [hep-ph]].

Y. Hamada, H. Kawai, K. y. Oda and S. C. Park, *"Higgs Inflation is Still Alive after the Results from BICEP2,"* Phys. Rev. Lett. **112** (2014) 24, 241301 [arXiv:1403.5043 [hep-ph]].

F. Bezrukov and M. Shaposhnikov, "Higgs inflation at the critical point,"

Phys. Lett. B **734** (2014) 249 [arXiv:1403.6078 [hep-ph]].

Y. Hamada, H. Kawai, K. y. Oda and S. C. Park, *"Higgs inflation from Standard Model criticality,"* Phys. Rev. D **91** (2015) 5, 053008 [arXiv:1408.4864 [hep-ph]].

Y. Hamada, K. y. Oda and F. Takahashi, *"Topological Higgs inflation: Origin of Standard Model criticality,"* Phys. Rev. D **90** (2014) 9, 097301 [arXiv:1408.5556 [hep-ph]].

F. Bezrukov, J. Rubio and M. Shaposhnikov, "Living beyond the edge: Higgs inflation and vacuum metastability," arXiv:1412.3811 [hep-ph].

Y. Hamada, "Standard Model criticality and Higgs inflation," arXiv:1504.04246 [hep-ph].

A. Salvio and A. Mazumdar, "Classical and Quantum Initial Conditions for Higgs Inflation," arXiv:1506.07520 [hep-ph].

- [7] Y. Hamada, H. Kawai and K. y. Oda, "Eternal Higgs inflation and cosmological constant problem," arXiv:1501.04455 [hep-ph].
- [8] A. Ashoorioon, H. Firouzjahi and M. M. Sheikh-Jabbari, "M-flation: Inflation From Matrix Valued Scalar Fields," JCAP 0906 (2009) 018 [arXiv:0903.1481 [hep-th]].

M. Blaszczyk, S. Groot Nibbelink, O. Loukas and S. Ramos-Sanchez, "Non-supersymmetric heterotic model building," JHEP **1410** (2014) 119 [arXiv:1407.6362 [hep-th]].

Y. Hamada, T. Kobayashi and S. Uemura, "Standard Model-like D-brane models and gauge couplings," arXiv:1409.2740 [hep-th].

S. Abel, K. R. Dienes and E. Mavroudi, *"Towards a Non-Supersymmetric String Phenomenology,"* arXiv:1502.03087 [hep-th].

S. G. Nibbelink, "Model building with the non-supersymmetric heterotic  $SO(16) \times SO(16)$  string," arXiv:1502.03604 [hep-th].

- [9] H. Kawai, D. C. Lewellen and S. H. H. Tye, "Construction of Four-Dimensional Fermionic String Models," Phys. Rev. Lett. 57 (1986) 1832
  [Phys. Rev. Lett. 58 (1987) 429].
  H. Kawai, D. C. Lewellen and S. H. H. Tye, "Construction of Fermionic String Models in Four-Dimensions," Nucl. Phys. B 288 (1987) 1.
  W. Lerche, D. Lust and A. N. Schellekens, "Chiral Four-Dimensional Heterotic Strings from Selfdual Lattices," Nucl. Phys. B 287 (1987) 477.
  I. Antoniadis, C. P. Bachas and C. Kounnas, "Four-Dimensional Superstrings," Nucl. Phys. B 289 (1987) 87.
- [10] C. Ford, I. Jack and D. R. T. Jones, "The Standard model effective potential at two loops," Nucl. Phys. B 387 (1992) 373 [Erratum-ibid. B 504 (1997) 551] [hep-ph/0111190].
- [11] P. A. R. Ade et al. [Planck Collaboration], "Planck 2015 results. XX. Constraints on inflation," arXiv:1502.02114 [astro-ph.CO].

- [12] L. J. Dixon and J. A. Harvey, "String Theories in Ten-Dimensions Without Space-Time Supersymmetry," Nucl. Phys. B 274 (1986) 93.
  L. Alvarez-Gaume, P. H. Ginsparg, G. W. Moore and C. Vafa, "An O(16) x O(16) Heterotic String," Phys. Lett. B 171 (1986) 155.
  H. Itoyama and T. R. Taylor, "Supersymmetry Restoration in the Compactified O(16) × O(16)-prime Heterotic String Theory," Phys. Lett. B 186 (1987) 129.
- [13] C. D. Froggatt and H. B. Nielsen, "Standard model criticality prediction: Top mass 173 ± 5-GeV and Higgs mass 135 +- 9-GeV," Phys. Lett. B 368 (1996) 96 [hep-ph/9511371].
  H. B. Nielsen, "PREdicted the Higgs Mass," arXiv:1212.5716 [hep-ph].
- [14] H. Kawai and T. Okada, "Solving the Naturalness Problem by Baby Universes in the Lorentzian Multiverse," Prog. Theor. Phys. 127 (2012) 689 [arXiv:1110.2303 [hep-th]].
  H. Kawai, "Low energy effective action of quantum gravity and the naturalness problem," Int. J. Mod. Phys. A 28 (2013) 1340001.
  Y. Hamada, H. Kawai and K. Kawana, "Evidence of the Big Fix," Int. J. Mod. Phys. A 29 (2014) 17, 1450099 [arXiv:1405.1310 [hep-ph]].
  K. Kawana, "Reconsideration of the Coleman's Baby Universe," arXiv:1405.2743 [hep-th].
  Y. Hamada, H. Kawai and K. Kawana, "Weak Scale From the Maximum Entropy Principle," PTEP 2015 (2015) 3, 033B06 [arXiv:1409.6508 [hep-ph]]

*Entropy Principle*," PTEP **2015** (2015) 3, 033B06 [arXiv:1409.6508 [hep-ph]].

[15] C. S. Chen and Y. Tang, "Vacuum stability, neutrinos, and dark matter," JHEP 1204 (2012) 019 [arXiv:1202.5717 [hep-ph]].

N. Haba, K. Kaneta and R. Takahashi, "Planck scale boundary conditions in the standard model with singlet scalar dark matter," JHEP **1404** (2014) 029 [arXiv:1312.2089 [hep-ph]].

Y. Hamada, H. Kawai and K. y. Oda, "Predictions on mass of Higgs portal scalar dark matter from Higgs inflation and flat potential," JHEP 1407 (2014) 026 [arXiv:1404.6141 [hep-ph]].

N. Khan and S. Rakshit, "Study of electroweak vacuum metastability with a singlet scalar dark matter," Phys. Rev. D **90**, no. 11, 113008 (2014) [arXiv:1407.6015 [hep-ph]].

K. Kawana, "Multiple Point Principle of the Standard Model with Scalar Singlet Dark Matter and Right Handed Neutrinos," PTEP **2015** (2015) 2, 023B04 [arXiv:1411.2097 [hep-ph]].

K. Kawana, "Criticality and Inflation of the Gauged B-L Model," arXiv:1501.04482 [hep-ph].

Y. Hamada, K. Kawana and K. Tsumura, "Landau pole in the Standard Model with weakly interacting scalar fields," Phys. Lett. B **747** (2015) 238 [arXiv:1505.01721 [hep-ph]].

N. Khan and S. Rakshit, "Constraints on inert dark matter from metastability of electroweak vacuum," arXiv:1503.03085 [hep-ph].
Y. Hamada and K. Kawana, "Vanishing Higgs Potential in Minimal Dark Matter Models," arXiv:1506.06553 [hep-ph].