SOBOLEV EMBEDDING OF A SPHERE CONTAINING AN ARBITRARY CANTOR SET IN THE IMAGE

PIOTR HAJLASZ AND XIAODAN ZHOU

ABSTRACT. We construct a large class of pathological n -dimensional topological spheres in \mathbb{R}^{n+1} by showing that for any Cantor set $C \subset \mathbb{R}^{n+1}$ there is a topological embedding $f: \mathbb{S}^n \to \mathbb{R}^{n+1}$ of the Sobolev class $W^{1,n}$ whose image contains the Cantor set C.

1. INTRODUCTION

In 1924 J. W. Alexander [1], constructed a homeomorphism $f : \mathbb{S}^2 \to f(\mathbb{S}^2) \subset \mathbb{R}^3$ so that the unbounded component of $\mathbb{R}^3 \setminus f(\mathbb{S}^2)$ is not simply connected. In particular it is not homeomorphic to the complement of the standard ball in \mathbb{R}^3 . This famous construction known as the Alexander horned sphere can be easily generalized to higher dimensions. The aim of this paper is to show that a large class of pathological topological n -dimensional spheres, including the Alexander horned sphere, can be realized as the images of Sobolev $W^{1,n}$ homeomorphisms, each of which being a smooth diffeomorphism outside of a Cantor set in \mathbb{S}^n of Hausdorff dimension zero. For a precise statement, see Theorem 1.1. Here and in what follows \mathbb{S}^n will denote the standard unit sphere in \mathbb{R}^{n+1} .

The following classical result [27, Theorem 30.3] provides a characterization of spaces that are homeomorphic to the ternary Cantor set: A metric space is homeomorphic to the ternary Cantor set if and only if it is compact, totally disconnected and has no isolated points. Recall that the space is *totally disconnected* if the only non-empty connected subsets are one-point sets. In what follows by a *Cantor set* we will mean any subset of Euclidean space that is homeomorphic to the ternary Cantor set.

Recall also that the *Sobolev space* $W^{1,p}$ consists of functions in L^p whose distributional gradient is in L^p . By $f \in W^{1,p}(\mathbb{S}^n,\mathbb{R}^{n+1})$ we will mean that the components of the mapping $f : \mathbb{S}^n \to \mathbb{R}^{n+1}$ are in $W^{1,p}$. For more details regarding Sobolev spaces see [2, 8].

Throughout the paper by an *embedding* we will mean a homeomorphism onto the image i.e., $f: X \to Y$ is an embedding if $f: X \to f(X)$ is a homeomorphism. In the literature such an embedding is often called a topological embedding. The main result of the paper reads as follows.

Theorem 1.1. For any Cantor set $C \subset \mathbb{R}^{n+1}$, $n \geq 2$, there is an embedding $f : \mathbb{S}^n \to \mathbb{R}^{n+1}$ such that

²⁰¹⁰ Mathematics Subject Classification. 46E35, 57A50.

Key words and phrases. Sobolev mappings, Alexander horned sphere.

P.H. was supported by NSF grant DMS-1161425.

- (a) $f \in W^{1,n}(\mathbb{S}^n,\mathbb{R}^{n+1}),$
- (b) $C \subset f(\mathbb{S}^n)$,
- (c) $f^{-1}(C)$ ⊂ \mathbb{S}^n is a Cantor set of Hausdorff dimension zero,
- (d) f is a smooth diffeomorphism in $\mathbb{S}^n \setminus f^{-1}(C)$.

Remark 1.2. Our construction resembles that of the Alexander horned sphere and it will be clear that it can be used to construct a version of the Alexander horned sphere $f: \mathbb{S}^n \to f(\mathbb{S}^n) \subset \mathbb{R}^{n+1}$ so that $f \in W^{1,n}$ and f is a smooth diffeomorphism outside a Cantor set of Hausdorff dimension zero.

Remark 1.3. A similar technique to the one used in the proof of Theorem 1.1 has also been employed in a variety of different settings [3, 4, 6, 9, 10, 11, 14, 15, 16, 17, 19, 20, 24, 25, 26].

Remark 1.4. According to Theorem 1.1 we can construct a topological sphere in \mathbb{R}^3 that is $W^{1,2}$ homeomorphic to \mathbb{S}^2 and that contains Antoine's necklace. Antoine's necklace is a Cantor set in \mathbb{R}^3 whose complement is not simply connected. Hence the unbounded component of $\mathbb{R}^3 \setminus f(\mathbb{S}^2)$ is also not simply connected. This gives a different example with the same topological consequences as those of the Alexander horned sphere. In fact, using results of Sher [22] we will show that there are uncountably many "essentially different" examples. For a precise statement see Theorem 1.7.

Remark 1.5. One cannot in general demand the function constructed in the theorem to be in $W^{1,p}, p > n$. Indeed, if $f \in W^{1,p}(\mathbb{S}^n, \mathbb{R}^{n+1})$, then the image $f(\mathbb{S}^n)$ has finite *n*-dimension Hausdorff measure, but a Cantor set in \mathbb{R}^{n+1} may have positive $(n+1)$ -dimensional measure and in that case it cannot be contained in the image of f. The fact that the image $f(\mathbb{S}^n)$ has finite n-dimensional measure follows from the area formula and the integrability of the Jacobian of f. The fact that the area formula is satisfied for mappings $f \in W^{1,p}(\mathbb{S}^n,\mathbb{R}^{n+1})$ with $p > n$ is well known and follows from the following observations. The area formula is true for Lipschitz mappings [8, Theorem 3.3.2]. The domain \mathbb{S}^n is the union of countably many sets on which f is Lipschitz continuous [8, Section 6.6.3] plus a set of measure zero. On Lipschitz pieces the area formula is satisfied. Since the mapping f maps sets of measure zero to sets of *n*-dimensional Hausdorff measure zero, $[12,$ Theorem 4.2, the area formula is in fact true for f. The proof presented in $[12,$ Theorem 4.2 is in the case of mappings into \mathbb{R}^n , but the same proof works in the case of mappings into \mathbb{R}^{n+1} .

Remark 1.6. It is well known, [12, Theorem 4.9], that any homeomorphism $f : \Omega \to$ $f(\Omega) \subset \mathbb{R}^n$ of class $W^{1,n}$, where $\Omega \subset \mathbb{R}^n$ is open, has the Lusin property, i.e. it maps sets of measure zero to sets of measure zero. Reshetnyak [21] observed that this is no longer true for embeddings $f \in W^{1,n}(\mathbb{S}^n,\mathbb{R}^m)$, when $m > n \geq 2$. In his example he considered $n = 2$ and $m = 3$, see also [7, Example 5.1]. Later Väisälä [23] generalized it to any $n > 2$ and $m > n$. In the constructions of Reshetnyak and Väisälä a set of measure zero is mapped to a set of positive n-dimensional Hausdorff measure. Theorem 1.1 also provides an example of this type. Indeed, if a Cantor set $C \subset \mathbb{R}^{n+1}$ has positive $(n + 1)$ dimensional measure, then the embedding $f \in W^{1,n}(\mathbb{S}^n,\mathbb{R}^{n+1})$ from Theorem 1.1 maps the set $f^{-1}(C)$ of Hausdorff dimension zero onto the set $C \subset f(\mathbb{S}^n)$ of positive $(n+1)$ dimensional measure. Actually, in a context of the Lebesgue area a similar example has already been constructed by Besicovitch [3, 4], but Besicovitch did not consider the Sobolev regularity of the mapping. The construction of Besicovitch is very different from that of Reshetnyak and Väisäla and it is more related to ours. While Besicovitch's construction

deals with a particular Cantor set, we deal with any Cantor set and we prove that the resulting mapping f belongs to the Sobolev space $W^{1,n}$.

We say that two embeddings $f, g : \mathbb{S}^n \to \mathbb{R}^{n+1}$, are *equivalent* if there is a homeomorphism $h: \mathbb{R}^{n+1} \to \mathbb{R}^{n+1}$ such that $h(f(\mathbb{S}^n)) = g(\mathbb{S}^n)$.

The second main result of the paper reads as follows.

Theorem 1.7. There are uncountably many embeddings $f : \mathbb{S}^2 \to \mathbb{R}^3$ of class $W^{1,2}(\mathbb{S}^2, \mathbb{R}^3)$ which are not equivalent.

The theorem can be generalized to higher dimensions, but we consider the case $n = 2$ only because our proof is based on a result of Sher [22] about Cantor sets in \mathbb{R}^3 . Generalization of Theorem 1.7 to $n \geq 3$ would require a generalization of Sher's result to higher dimensions. Since this would be a work of purely technical nature with predicted answer, we do not find it particularly interesting.

The paper is organized as follows. Sections 2 and 3 contain preliminary material needed in the proof of Theorem 1.1. In Section 2 we introduce some terminology regarding Cantor sets and we construct a Cantor tree. In Section 3 we define surfaces, called tentacles, around smooth curves. Finally in Sections 4 and 5 we prove Theorems 1.1 and 1.7 respectively.

Acknowledgements. The authors would like to thank the referee for valuable comments that led to an improvement of the paper.

2. Cantor sets and trees

2.1. **Ternary Cantor set.** The ternary Cantor set will be denoted by \mathfrak{C} . It is constructed by removing the middle third of the unit interval [0, 1], and then successively deleting the middle third of each resulting subinterval. Denote by \mathfrak{I}^k all binary numbers $i_1 \ldots i_k$ such that $i_j \in \{0,1\}$ for $j = 1,2,\ldots,k$, and by \mathfrak{I}^{∞} all binary infinite sequences $i_1 i_2 \ldots$ Clearly, the ternary Cantor set $\mathfrak C$ can be written as

$$
\mathfrak{C} = \bigcap_{k=1}^{\infty} \bigcup_{i_1...i_k \in \mathfrak{I}^k} I_{i_1...i_k},
$$

where $I_{i_1...i_k}$ is one of the 2^k closed intervals in the k-th level of the construction of the Cantor set \mathfrak{C} ; the binary number $i_1 \ldots i_k$ denotes the position of this interval: If $i_k = 0$, it is the left subinterval of $I_{i_1...i_{k-1}}$, otherwise it is the right subinterval.

We also have that for any $k \in \mathbb{N}$

$$
\mathfrak{C} = \bigcup_{i_1 \ldots i_k \in \mathfrak{I}^k} \mathfrak{C}_{i_1 \ldots i_k}
$$

where $\mathfrak{C}_{i_1...i_k} = \mathfrak{C} \cap I_{i_1...i_k}$.

Note that points in the Cantor set $\mathfrak C$ can be uniquely encoded by infinite binary sequences. Indeed, if $\mathfrak{i} = i_1 i_2 \ldots \in \mathfrak{I}^{\infty}$, then

$$
\{\mathfrak{c}_i\}=\bigcap_{k=1}^{\infty}I_{i_1...i_k}=\bigcap_{k=1}^{\infty}\mathfrak{C}_{i_1...i_k}
$$

consists of a single point $c_i \in \mathfrak{C}$ and

$$
\mathfrak{C}=\bigcup_{i\in\mathfrak{I}^\infty}\{ \mathfrak{c}_i\}.
$$

2.2. Cantor trees. Let $C \subset \mathbb{R}^{n+1}$ be a Cantor set and let $f : \mathfrak{C} \to C$ be a homeomorphism. We will write $C_{i_1...i_k} = f(\mathfrak{C}_{i_1...i_k})$ and $c_i = f(\mathfrak{c}_i)$ for $i \in \mathfrak{I}^{\infty}$. Since the mapping f is uniformly continuous,

(2.1)
$$
\max_{i_1...i_k \in \mathfrak{I}^k} (\text{diam } C_{i_1...i_k}) \to 0 \text{ as } k \to \infty.
$$

For each k and each $i_1 \ldots i_k \in \mathfrak{I}^k$ we select a point $A_{i_1 \ldots i_k}$ such that

- \bullet The point $A_{i_1 \ldots i_k}$ does not belong to $C,$
- The distance of the point $A_{i_1...i_k}$ to $C_{i_1...i_k}$ is less than 2^{-k} ,
- $A_{i_1...i_k} \neq A_{j_1...j_\ell}$ if $i_1...i_k \neq j_1...j_\ell$.

It is easy to see that if $i = i_1 i_2 ... \in \mathfrak{I}^{\infty}$, then

$$
A_{i_1...i_k} \to c_{\mathfrak{i}} = f(\mathfrak{c}_{\mathfrak{i}}) \quad \text{as } k \to \infty.
$$

Indeed, $c_i \in C_{i_1...i_k}$ so

$$
|A_{i_1\ldots i_k} - c_i| < 2^{-k} + \operatorname{diam} C_{i_1\ldots i_k} \to 0 \quad \text{as } k \to \infty.
$$

Now we are ready to build a *Cantor tree* by adding branches $J_{i_1...i_ki_{k+1}}$ connecting $A_{i_1...i_ki_k}$ to $A_{i_1...i_ki_{k+1}}$. The precise construction goes as follows.

By translating the coordinate system we may assume that the distance between the origin in \mathbb{R}^{n+1} and the Cantor set C is greater than 100 (that is way too much, but there is nothing wrong with being generous).

Let J_0 and J_1 be smooth Jordan arcs (i.e., smoothly embedded arcs without selfintersections) of unit speed (i.e., parametrized by arc-length) connecting the origin 0 to the points A_0 and A_1 respectively. We also assume that

- The curve J_0 does not intersect with the curve J_1 (except for the common endpoint 0).
- The curves J_0 and J_1 avoid the Cantor set C.
- The curves J_0 and J_1 meet the unit ball $\mathbb{B}^n(0,1) \subset \mathbb{R}^n \times \{0\} \subset \mathbb{R}^{n+1}$ lying in the hyperplane of the first n coordinates only at the origin and both curves exit $\mathbb{B}^n(0,1)$ on the same side of $\mathbb{B}^n(0,1)$ in \mathbb{R}^{n+1} .

The curves J_0 and J_1 will be called *branches of order* 1.

A simple topological observation is needed here. While the topological structure of a Cantor set C inside \mathbb{R}^{n+1} may be very complicated (think of Antoine's necklace), no

Cantor set can separate open sets in \mathbb{R}^{n+1} . Indeed, by [13, Corollary 2 of Theorem IV 3] compact sets separating open sets in \mathbb{R}^{n+1} must have topological dimension at least n but the topological dimension of a Cantor set is 0, [13, Example II 3]. Hence we can connect points in the complement of a Cantor set by smooth Jordan arcs that avoid the Cantor set. Moreover we can construct such an arc in a way that it is arbitrarily close to the line segment connecting the endpoints.

Recall that the ε -neighborhood of a set A is the set of all points whose distance to the set A is less than ε .

Suppose that we have already constructed all branches $J_{i_1...i_k}$ of order $k \geq 1$. The construction of the branches of order $k+1$ goes as follows. $\{J_{i_1...i_{k+1}}\}$ is a family of 2^{k+1} curves such that

- $J_{i_1...i_{k+1}}$ is a smooth Jordan arc parametrized by arc-length that connects $A_{i_1...i_k}$ (an endpoint of the branch $J_{i_1...i_k}$) to $A_{i_1...i_ki_{k+1}}$.
- The curves $J_{i_1...i_{k+1}}$ do not intersect with the Cantor set C, they do not intersect with each other (except for the common endpoints) and they do not intersect with previously constructed branches of orders less than or equal to k (except for the common endpoints).
- The image of the curve $J_{i_1...i_{k+1}}$ is contained in the 2^{-k}-neighborhood of the line segment $A_{i_1...i_k}A_{i_1...i_{k+1}}$.
- The angle between the branch $J_{i_1...i_k}$ and each of the emerging branches $J_{i_1...i_k0}$ and $J_{i_1...i_k1}$ at the point $A_{i_1...i_k}$ where the curves meet is larger than $\pi/2$.

The reason why we require the last condition about the angles is far from being clear at the moment, but it will be clarified in Section 2.3.

In what follows, depending on the situation, $J_{i_1...i_k}$ will denote either the curve (a map from an interval to \mathbb{R}^{n+1} or its image (a subset of \mathbb{R}^{n+1}), but it will always be clear from the context what interpretation we use.

A Cantor tree is the closure of the union of all branches

$$
T = \overline{\bigcup_{k=1}^{\infty} \bigcup_{i_1...i_k \in \mathfrak{I}^k} J_{i_1...i_k}}.
$$

We also define T_k to be the tree with branches of orders less than or equal to k removed. Formally

$$
T_k = T \setminus B_k
$$
 where $B_k = \bigcup_{s=1}^k \bigcup_{i_1...i_s \in \mathfrak{I}^s} J_{i_1...i_s}.$

Note that the set T_k is not closed – it does not contain the endpoints $A_{i_1...i_k}$.

The branches $J_{i_1...i_{k+1}}$ are very close to the sets $C_{i_1...i_k}$ in the following sense.

Lemma 2.1. A branch $J_{i_1...i_{k+1}}$ is contained in the 2^{-k+2} + diam $C_{i_1...i_k}$ neighborhood of $C_{i_1...i_k}$.

Proof. A branch $J_{i_1...i_{k+1}}$ connects the points $A_{i_1...i_k}$ and $A_{i_1...i_{k+1}}$. The distance of $A_{i_1...i_k}$ and $A_{i_1...i_{k+1}}$ to the set $C_{i_1...i_k}$ is less than 2^{-k} (because $C_{i_1...i_{k+1}} \subset C_{i_1...i_k}$). Hence

$$
|A_{i_1...i_k} - A_{i_1...i_{k+1}}| < 2 \cdot 2^{-k} + \text{diam } C_{i_1...i_k}
$$

so by the triangle inequality the line segment $\overline{A_{i_1...i_k}A_{i_1...i_{k+1}}}$ is contained in the 3 · 2^{-k} + diam $C_{i_1...i_k}$ neighborhood of $C_{i_1...i_k}$. Since $J_{i_1...i_{k+1}}$ is contained in the 2^{-k} neighborhood of the line segment, the lemma follows.

Corollary 2.2. T_k is contained in the

$$
\varepsilon_k := 2^{-k+2} + \max_{i_1 \dots i_k \in \mathcal{I}^k} (\text{diam } C_{i_1 \dots i_k})
$$

neighborhood of the Cantor set C and $\varepsilon_k \to 0$ as $k \to \infty$.

Proof. Indeed, the branches of T_k are of the form $J_{i_1...i_{s+1}}$, $s \geq k$. Each such branch is contained in the 2^{-s+2} + diam $C_{i_1...i_s}$ neighborhood of $C_{i_1...i_s} \subset C$. Since $s \geq k$ and $C_{i_1...i_s} \subset C_{i_1...i_k}$ we have

$$
2^{-s+2} + \text{diam}\, C_{i_1...i_s} \le 2^{-k+2} + \text{diam}\, C_{i_1...i_k}
$$

so T_k is contained in the ε_k neighborhood of C. The fact that $\varepsilon_k \to 0$ follows from (2.1).
The proof is complete The proof is complete.

Since the sets B_k are compact and their complements $T_k = T \setminus B_k$ are in close proximity of C by Corollary 2.2, it easily follows that

$$
T = C \cup \bigcup_{k=1}^{\infty} \bigcup_{i_1...i_l \in \mathfrak{I}^k} J_{i_i...i_k}.
$$

2.3. What is it for? The idea of the proof of Theorem 1.1 is to build a surface that looks very similar to the tree T with one dimensional branches $J_{i_1...i_k}$ of the tree T replaced by smooth thin surfaces built around the curves $J_{i_1...i_k}$; such surfaces will be called *tentacles*. The parametric surface $f : \mathbb{S}^n \to \mathbb{R}^{n+1}$ will be constructed as a limit of smooth surfaces $f_k: \mathbb{S}^n \to \mathbb{R}^{n+1}$. This sequence will be defined by induction. In the step k we replace all branches $J_{i_1...i_k}$ by smooth surfaces – tentacles. Such surfaces will be very close to the branches $J_{i_1...i_k}$ and they will pass through the endpoints $A_{i_1...i_k}$. The only place where the set T_k gets close to the branch $J_{i_1...i_k}$ is the endpoint $A_{i_1...i_k}$ where the two branches $J_{i_1...i_k0}$ and $J_{i_1...i_k}$ of the set T_k emerge. The surface around $J_{i_1...i_k}$, and passing through the point $A_{i_1...i_k}$ will be orthogonal to the curve $J_{i_1...i_k}$ at the point $A_{i_1...i_k}$. Since the branches $J_{i_1...i_k0}$ and $J_{i_1...i_k}$ emerging from that point form angles larger than $\pi/2$ with $J_{i_1...i_k}$ the surface will not intersect the branches $J_{i_1...i_k0}$ and $J_{i_1...i_k1}$. By making the surfaces around $J_{i_1...i_k}$ thin enough we can make them disjoint from the set T_k (note that $A_{i_1...i_k}$ does not belong to T_k).

3. Sobolev tentacles

It is well known and easy to prove that $\eta(x) = \log |\log |x|| \in W^{1,n}(\mathbb{B}^n(0, e^{-1}))$. Define the truncation of η between levels s and t, $0 < s < t < \infty$ by

$$
\eta_s^t(x) = \begin{cases} t-s & \text{if } \eta(x) \ge t, \\ \eta(x) - s & \text{if } s \le \eta(x) \le t \\ 0 & \text{if } \eta(x) \le s. \end{cases}
$$

Fix an arbitrary $\tau > 0$. For every $\delta > 0$ there is a sufficiently large s such that $\widetilde{\eta}_{\delta,\tau} := \eta_s^{s+\tau}$ is a Lipschitz function on \mathbb{R}^n with the properties:

$$
\mathrm{supp}\,\widetilde{\eta}_{\delta,\tau}\subset\mathbb{B}^n(0,\delta/2),
$$

 $0 \leq \tilde{\eta}_{\delta,\tau} \leq \tau$ and $\tilde{\eta}_{\delta,\tau} = \tau$ in a neighborhood $\mathbb{B}(0,\delta')$ of 0, Z $\int_{\mathbb{R}^n} |\nabla \widetilde \eta_{\delta,\tau} |^n < \delta^n.$

The function $\widetilde{\eta}_{\delta,\tau}$ is not smooth because it is defined as a truncation, however, mollifying $\widetilde{\eta}_{\delta,\tau}$ gives a smooth function, denoted by $\eta_{\delta,\tau}$, with the same properties as those of $\widetilde{\eta}_{\delta,\tau}$ listed above. In particular

(3.1)
$$
\int_{\mathbb{R}^n} |\nabla \eta_{\delta,\tau}|^n < \delta^n.
$$

The graph of $\eta_{\delta,\tau}$ restricted to the ball $\overline{\mathbb{B}}^n(0,\delta)$ is contained in the cylinder

(3.2)
$$
\{(x_1,\ldots,x_{n+1})|x_1^2+\ldots+x_n^2\leq \delta^2,\ 0\leq x_{n+1}\leq \tau\},\
$$

and it forms a slim "tower" around the x_{n+1} -axis. The function $\eta_{\delta,\tau}$ equals zero in the annulus $\mathbb{B}^n(0,\delta) \setminus \mathbb{B}^n(0,\delta/2)$ and equals τ in the ball $\mathbb{B}^n(0,\delta')$.

Consider now a smooth Jordan arc $\gamma : [-1, \tau + 1] \to \mathbb{R}^{n+1}$ parametrized by arc-length. We want to construct a smooth mapping $\gamma_{\delta} : \mathbb{B}^n(0,\delta) \to \mathbb{R}^{n+1}$ whose image will be a smooth, thin, tentacle-shaped surface around the curve $\gamma|_{[0,\tau]}$. To do this we will apply a diffeomorphism Φ mapping the cylinder (3.2) onto a neighborhood of the image of the curve γ. The tentacle-like surface will be the image of the graph of $η_{δ,τ}$ under the diffeomorphism Φ.

The construction of a diffeomorphism Φ follows a standard procedure. Let

$$
v_1, \ldots, v_n : [-1, \tau + 1] \to T\mathbb{R}^{n+1}
$$

be a smooth orthonormal basis in the orthogonal complement of the tangent space to the curve γ , i.e., for every $t \in [-1, \tau + 1]$, $\langle v_1(t), \ldots, v_n(t), \gamma'(t) \rangle$ is a positively oriented orthonormal basis of $T_{\gamma(t)} \mathbb{R}^{n+1}$. Now we define

$$
\Phi(x_1, \dots, x_{n+1}) = \gamma(x_{n+1}) + \sum_{i=1}^n x_i v_i(x_{n+1}) \quad \text{for } x \in \mathbb{R}^{n+1} \text{ with } -1 \le x_{n+1} \le \tau + 1.
$$

Clearly, Φ is smooth and its Jacobian equals 1 along the x_{n+1} axis, $-1 < x_{n+1} < \tau + 1$. Hence Φ is a diffeomorphism in a neighborhood of any point on the x_{n+1} -axis, $-1 < x_{n+1}$ $\tau+1$. Using compactness of the image of the curve γ it easily follows that there is a $\delta_0 > 0$

8 P. HAJŁASZ AND X. ZHOU

such that for all $0 < \delta < \delta_0$, Φ is a diffeomorphism in an open neighborhood of the cylinder (3.2). Now we define

$$
\gamma_{\delta}: \overline{\mathbb{B}}^n(0,\delta) \to \mathbb{R}^{n+1}, \quad \gamma_{\delta}(x_1,\ldots,x_n) = \Phi(x_1,\ldots,x_n,\eta_{\delta,\tau}(x_1,\ldots,x_n)).
$$

Since

$$
\frac{\partial \gamma_{\delta}}{\partial x_{i}} = \frac{\partial \Phi}{\partial x_{i}} + \frac{\partial \Phi}{\partial x_{n+1}} \frac{\partial \eta_{\delta, \tau}}{\partial x_{i}}
$$

it follows that

$$
|D\gamma_\delta| \leq \sqrt{n} \|D\Phi\|_\infty (1 + |\nabla \eta_{\delta,\tau}|),
$$

where $||D\Phi||_{\infty}$ is the supremum of the Hilbert-Schmidt norms $|D\Phi|$ over the cylinder (3.2). Hence using (3.1), for every $\varepsilon > 0$ we can find $\delta > 0$ so small that

(3.3)
$$
\int_{\mathbb{B}^n(0,\delta)} |D\gamma_\delta|^n \leq C(n) \|D\Phi\|_\infty^n \delta^n < \varepsilon.
$$

Observe that δ depends on γ (because $||D\Phi||_{\infty}$ depends on γ).

The tentacle γ_{δ} maps the annulus $\overline{\mathbb{B}}^n(0,\delta) \setminus \mathbb{B}^n(0,\delta/2)$ onto the isometric annulus in the hyperplane orthogonal to γ at $\gamma(0)$. Indeed, for $x \in \overline{\mathbb{B}}^n(0,\delta) \setminus \mathbb{B}^n(0,\delta/2)$, $\eta_{\delta,\tau}(x) = 0$ and hence

$$
\gamma_{\delta}(x) = \Phi(x_1, ..., x_n, 0) = \gamma(0) + \sum_{i=1}^{n} x_i v_i(0)
$$

is an affine isometry. For a similar reason γ_δ maps the ball $\overline{\mathbb{B}}^n(0,\delta')$ onto the isometric ball in the hyperplane orthogonal to γ at $\gamma(\tau)$. Finally for $x \in \overline{\mathbb{B}}^n(0, \delta/2) \setminus \mathbb{B}^n(0, \delta')$, γ_{δ} creates a smooth thin surface around the curve γ that connects the annulus at $\gamma(0)$ with the ball at $\gamma(\tau)$.

Composition with a translation allows us to define a tentacle $\gamma_{\delta} : \overline{\mathbb{B}}^n(p,\delta) \to \mathbb{R}^{n+1}$ centered at any point $p \in \mathbb{R}^n$.

4. Proof of Theorem 1.1

We will construct the Sobolev embedded surface $f : \mathbb{S}^n \to \mathbb{R}^{n+1}$ as a limit of smooth embedded surfaces $f_k: \mathbb{S}^n \to \mathbb{R}^{n+1}$.

By replacing \mathbb{S}^n with a diffeomorphic submanifold (still denoted by \mathbb{S}^n) we may assume that it contains the unit ball

$$
\mathbb{B}^n = \mathbb{B}^n(0,1) \subset \mathbb{R}^n \times \{0\} \subset \mathbb{R}^{n+1}.
$$

lying in the hyperplane of the first n coordinates.

Since the distance of the Cantor set C to the origin is larger than 100, the only parts of the Cantor tree T that are close to \mathbb{B}^n are the branches J_0 and J_1 that connect the origin to A_0 and A_1 . Since the branches meet \mathbb{B}^n only at the origin and leave \mathbb{B}^n on the same side of \mathbb{B}^n , we can assume that \mathbb{S}^n meets T only at the origin.

Now we will describe the construction of f_1 . As indicated in Section 2.3 we want to grow two tentacles from \mathbb{S}^n near the branches J_0 and J_1 all the way to points A_0 and A_1 , but we want to make sure that the tentacles do not touch the set T_1 .

SOBOLEV EMBEDDING 9

To do this we choose two distinct points $p_0, p_1 \in \mathbb{B}^n$ close to the origin and modify the curves J_0 and J_1 only near the origin, so that the modified Jordan arcs γ^0 and γ^1 emerge from the points p_0 and p_1 instead of the origin, and they are orthogonal to \mathbb{B}^n at the points p_0 and p_1 . The curves γ^0 and γ^1 quickly meet with J_0 and J_1 and from the points where they meet they coincide with J_0 and J_1 , so all non-intersection properties of the curves are preserved. Since the curves are modified only at their beginnings, outside that place they are identical with J_0 and J_1 .

Next, we find $\delta_1 > 0$ so small that the balls $\overline{\mathbb{B}}^n(p_0, \delta_1)$ and $\overline{\mathbb{B}}^n(p_1, \delta_1)$ are disjoint and contained in \mathbb{B}^n and that there are disjoint tentacles

$$
\gamma_{\delta_1}^i : \overline{\mathbb{B}}^n(p_i, \delta_1) \to \mathbb{R}^{n+1} \text{ for } i = 0, 1
$$

along the curves γ^0 and γ^1 such that

$$
\int_{\mathbb{B}(p_i,\delta_1)} |D\gamma^i_{\delta_1}|^n < 4^{-n} \quad \text{for } i = 0, 1.
$$

Observe that $\gamma_{\delta_1}^i(p_i) = A_i$ for $i = 0, 1$.

Note that the images of small balls $\mathbb{B}^n(p_0, \delta'_1)$ and $\mathbb{B}^n(p_1, \delta'_1)$ are isometric balls, orthogonal to the curves γ^0 and γ^1 (and hence to the curves J_0 and J_1) at the endpoints A_0 and A_1 . Since the branches J_{00} , J_{01} form angles larger than $\pi/2$ with the curve γ^0 at A_0 and the branches J_{10} , J_{11} form angles larger than $\pi/2$ with the curve γ^1 at A_1 we may guarantee, by making the tentacles sufficiently thin, that they are disjoint from the set T_1 (observe that the endpoints A_0 and A_1 do not belong to T_1).

Also each annulus $\overline{\mathbb{B}}^n(p_i, \delta_1) \setminus \mathbb{B}^n(p_i, \delta_1/2)$ for $i = 0, 1$ is mapped isometrically by $\gamma_{\delta_1}^i$ onto an annulus centered at $\gamma^{i}(0) = p_{i}$ in the hyperplane orthogonal to γ^{i} at $\gamma^{i}(0) = p_{i}$. Since the curve γ^i is orthogonal to \mathbb{B}^n at $\gamma^i(0) = p_i$, the annulus $\overline{\mathbb{B}}^n(p_i, \delta_1) \setminus \mathbb{B}^n(p_i, \delta_1/2)$ is mapped isometrically onto itself. By choosing an appropriate orthonormal frame $v_1^i, \ldots v_n^i$ in the definition of $\gamma_{\delta_1}^i$ we may assume that $\gamma_{\delta_1}^i$ is the identity in the annulus. This guarantees that the mapping

$$
f_1(x) = \begin{cases} \gamma_{\delta_1}^0(x) & \text{if} \\ \gamma_{\delta_1}^1(x) & \text{if} \\ x & \text{if} \end{cases} \quad \begin{array}{l} x \in \overline{\mathbb{B}}^n(p_0, \delta_1), \\ x \in \overline{\mathbb{B}}^n(p_1, \delta_1), \\ (\mathbb{B}^n(p_0, \delta_1) \cup \mathbb{B}^n(p_1, \delta_1)) \end{array}
$$

is continuous and hence smooth. The construction guarantees also that f_1 is a smooth embedding of \mathbb{S}^n into \mathbb{R}^{n+1} with the image that is disjoint from T_1 .

The mapping f_1 maps the small balls $\overline{\mathbb{B}}^n(p_0, \delta'_1)$ and $\overline{\mathbb{B}}^n(p_1, \delta'_1)$ onto isometric balls centered at A_0 and A_1 respectively with $f_1(p_i) = A_i$ for $i = 0, 1$. Now the mapping f_2 will be obtained from f_1 by adding four more tentacles: from the ball $f_1(\mathbb{B}^n(p_0, \delta'_1))$ centered at A_0 there will be two tentacles connecting this ball to the points A_{00} and A_{01} and from the ball at $f_1(\mathbb{B}^n(p_1, \delta'_1))$ centered at A_1 there will be two tentacles connecting this ball to the points A_{10} and A_{11} . More precisely the inductive step is described as follows.

Suppose that we have already constructed a mapping $f_k, k \geq 1$ such that

• f_k is a smooth embedding of \mathbb{S}^n into \mathbb{R}^{n+1} whose image is disjoint from T_k .

• There are 2^k disjoint balls $\overline{\mathbb{B}}^n(p_{i_1...i_k}, \delta_k) \subset \mathbb{B}^n$ and 2^k tentacles

 $\gamma^{i_1...i_k}_{\delta_k}$ $\delta_k^{i_1...i_k} : \overline{\mathbb{B}}^n(p_{i_1...i_k}, \delta_k) \to \mathbb{R}^{n+1}$

for $i_1 \dots i_k \in \mathfrak{I}^k$ such that

- $\Diamond \gamma^{i_1...i_k}_{\delta_k}$ $\delta_k^{i_1...i_k}(p_{i_1...i_k})=A_{i_1...i_k}.$
- \Diamond The image of $\gamma_{\delta_k}^{i_1...i_k}$ $\delta_k^{i_1...i_k}$ is in the 2^{-k} neighborhood of the curve $J_{i_1...i_k}$.
- \diamond We have

(4.1)
$$
\int_{\mathbb{B}^n(p_{i_1...i_k},\delta_k)} |D\gamma_{\delta_k}^{i_1...i_k}|^n dx < 4^{-nk}.
$$

• The mapping f_k satisfies

$$
f_k = \begin{cases} \gamma_{\delta_k}^{i_1 \dots i_k} & \text{in } \mathbb{B}^n(p_{i_1 \dots i_k}, \delta_k) \text{ for } i_1 \dots i_k \in \mathfrak{I}^k, \\ f_{k-1} & \text{in } \mathbb{S}^n \setminus \bigcup_{i_1 \dots i_k \in \mathfrak{I}^k} \mathbb{B}^n(p_{i_1 \dots i_k}, \delta_k). \end{cases}
$$

Observe that $f_k(p_{i_1...i_k}) = A_{i_1...i_k}$ and that for some small $0 < \delta'_k < \delta_k$, f_k maps balls $\overline{\mathbb{B}}^n(p_{i_1...i_k},\delta'_k)$ onto isometric balls centered at $A_{i_1...i_k}$.

Now we will describe the construction of the mapping f_{k+1} .

For each $i_1 \dots i_k \in \mathfrak{I}^k$ we choose two points

$$
p_{i_1...i_k0}, p_{i_1...i_k1} \in \mathbb{B}^n(p_{i_1...i_k}, \delta'_k)
$$

and modify the curves $J_{i_1...i_k0}$ and $J_{i_1...i_k1}$ to $\gamma^{i_1...i_k0}$ and $\gamma^{i_1...i_k1}$ in a pretty similar way as we did for the curves γ^0 and γ^1 : the new curves $\gamma^{i_1...i_k0}$ and $\gamma^{i_1...i_k1}$ emerge from the points $f_k(p_{i_1...i_k0})$ and $f_k(p_{i_1...i_k1})$, they are orthogonal to the ball $f_k(\mathbb{B}^n(p_{i_1...i_k}, \delta'_k))$ at these points and then they quickly meet and coincide with $J_{i_1...i_k0}$ and $J_{i_1...i_k1}$.

We find $\delta_{k+1} > 0$ so small that

• The balls

$$
\overline{\mathbb{B}}^n(p_{i_1\ldots i_k0}, \delta_{k+1}), \overline{\mathbb{B}}^n(p_{i_1\ldots i_k1}, \delta_{k+1}) \subset \mathbb{B}^n(p_{i_1\ldots i_k}, \delta'_k)
$$

are disjoint.

• There are tentacles

$$
\gamma_{\delta_{k+1}}^{i_1\ldots i_k i} : \overline{\mathbb{B}}^n(p_{i_1\ldots i_k i}, \delta_{k+1}) \to \mathbb{R}^{n+1} \text{ for } i = 0, 1
$$

such that

 $\Diamond \ \gamma^{i_1...i_ki}_{\delta_{k+1}}$ $\delta_{k+1}^{i_1...i_ki}(p_{i_1...i_ki})=A_{i_1...i_ki}.$

- \Diamond The image of $\gamma^{i_1...i_k}_{\delta_{k+1}}$ $\delta_{\delta_{k+1}}^{i_1...i_k}$ is in the $2^{-(k+1)}$ neighborhood of the curve $J_{i_1...i_k}$.
- \Diamond The tentacles do not intersect and they avoid the set T_{k+1}

 \diamond We have

$$
\int_{\mathbb{B}^n(p_{i_1\ldots i_k i},\delta_{k+1})} |D\gamma_{\delta_{k+1}}^{i_1\ldots i_k i}|^n\,dx<4^{-n(k+1)}.
$$

The condition about the distance of the tentacle to the curve $J_{i_1...i_k}$ can be easily guaranteed, because the curve $\gamma^{i_1...i_k i}$ can be arbitrarily close to $J_{i_1...i_k i}$ and the tentacle can be arbitrarily thin.

Because the curves $\gamma^{i_1...i_k i}$, $i=0,1$, are orthogonal to the balls $f_k(\mathbb{B}^n(p_{i_1...i_k},\delta'_k))$ at the points $\gamma^{i_1...i_k i}(0) = f_k(p_{i_1...i_k i}),$ by choosing appropriate orthonormal frames in the definition of $\gamma_{\delta_{k+1}}^{i_1...i_k}$ $\delta_{\delta_{k+1}}^{i_1...i_k}$ we may guarantee one more condition

$$
\bullet \ \gamma_{\delta_{k+1}}^{i_1...i_k i} = f_k \text{ in } \overline{\mathbb{B}}^n(p_{i_1...i_k i}, \delta_{k+1}) \setminus \mathbb{B}^n(p_{i_1...i_k i}, \delta_{k+1}/2).
$$

We are using here the fact that both $\gamma_{\delta_{k+1}}^{i_1...i_k}$ $\delta_{\delta_{k+1}}^{i_1...i_k}$ and f_k are isometries in that annulus. Now we define

$$
f_{k+1} = \begin{cases} \gamma_{\delta_k}^{i_1 \dots i_k i_{k+1}} & \text{in } \mathbb{B}^n(p_{i_1 \dots i_{k+1}}, \delta_{k+1}) \text{ for } i_1 \dots i_{k+1} \in \mathfrak{I}^{k+1}, \\ f_k & \text{in } \mathbb{S}^n \setminus \bigcup_{i_1 \dots i_{k+1} \in \mathfrak{I}^{k+1}} \mathbb{B}^n(p_{i_1 \dots i_{k+1}}, \delta_{k+1}). \end{cases}
$$

As before f_{k+1} is a smooth embedding of \mathbb{S}^n into \mathbb{R}^{n+1} whose image is disjoint from T_{k+1} .

Let

$$
W_k = \bigcup_{i_1...i_k \in \mathfrak{I}^k} \overline{\mathbb{B}}^n(p_{i_1...i_k}, \delta_k).
$$

Clearly, W_k is a decreasing sequence of compact sets and

$$
E:=\bigcap_{k=1}^\infty W_k
$$

is a Cantor set $E \subset \mathbb{B}^n \subset \mathbb{S}^n$. By making the sequence δ_k converge to zero sufficiently fast we may guarantee that the Hausdorff dimension of E equals zero. Similarly as in the case of the ternary Cantor set, points in the set E can be encoded by infinite binary sequences For $\mathfrak{i} = i_1 i_1 \ldots \in \mathfrak{I}^{\infty}$ we define

$$
\{e_i\} = \bigcap_{k=1}^{\infty} \overline{\mathbb{B}}^n(p_{i_1...i_k,\delta_k}) \quad \text{so} \quad E = \bigcup_{i \in \mathfrak{I}^{\infty}} \{e_i\}.
$$

Now we define

$$
f(x) = \begin{cases} x & \text{if } x \in \mathbb{S}^n \setminus W_1, \\ f_k(x) & \text{if } x \in W_k \setminus W_{k+1}, k = 1, 2, \dots \\ c_i & \text{if } x = e_i \in E, i \in \mathfrak{I}^\infty. \end{cases}
$$

Recall that $c_i = f(c_i)$ is a point of the Cantor set C.

Lemma 4.1. $f = f_k$ when restricted to $\mathbb{S}^n \setminus W_{k+1}$.

 $($

Proof. The lemma can be easily proved by induction. Let $f_0 = id$. By the definition of f, $f = f_0$ in $\mathbb{S}^n \setminus W_1$. Suppose now that $f = f_k$ in $\mathbb{S}^n \setminus W_{k+1}$. According to the construction of $f_{k+1}, f_{k+1} = f_k$ in $\mathbb{S}^n \setminus W_{k+1}$, but the definition of f yields $f = f_{k+1}$ in $W_{k+1} \setminus W_{k+2}$ so $f = f_{k+1}$ in

$$
\mathbb{S}^n \setminus W_{k+1}) \cup (W_{k+1} \setminus W_{k+2}) = \mathbb{S}^n \setminus W_{k+2}.
$$

This proves the lemma.

Since each of the mappings f_k is a smooth embedding whose image does not intersect with the Cantor set C it follows from the lemma that f restricted to the open set $\mathbb{S}^n \setminus E$ is a smooth embedding and $f(\mathbb{S}^n \setminus E) \cap C = \emptyset$. In the remaining Cantor set E, the mapping f is defined as a bijection that maps E onto C. Therefore the mapping f is one-to-one in \mathbb{S}^n and $C \subset f(\mathbb{S}^n)$.

12 P. HAJLASZ AND X. ZHOU

It remains to prove that f is continuous and that $f \in W^{1,n}$.

First we will prove that $f \in W^{1,n}$. The mapping f is bounded and hence its components are in L^n . Since the mapping f is smooth outside the Cantor set E of Hausdorff dimension zero, according to the characterization of the Sobolev space by absolute continuity on lines, [8, Section 4.9.2], it suffices to show that the classical derivative of f defined outside of E (and hence a.e. in \mathbb{S}^n) belongs to $L^n(\mathbb{S}^n)$. We have

$$
\int_{\mathbb{S}^n} |Df|^n = \int_{\mathbb{S}^n \backslash E} |Df|^n = \int_{\mathbb{S}^n \backslash W_1} |Df|^n + \sum_{k=1}^{\infty} \int_{W_k \backslash W_{k+1}} |Df_k|^n
$$

$$
\leq \int_{\mathbb{S}^n \backslash W_1} |Df|^n + \sum_{k=1}^{\infty} \int_{W_k} |Df_k|^n.
$$

Since $f(x) = x$ in $\mathbb{S}^n \setminus W_1$, we do not have to worry about the first term on the right hand side and it remains to estimate the infinite sum.

Note that $f_k = \gamma_{\delta_k}^{i_1...i_k}$ $\overline{\mathbb{B}}^n(p_{i_1...i_k},\delta_k)$ so (4.1) yields \int \int $\frac{1}{2}$ \int

$$
\int_{\mathbb{B}^n(p_{i_1\ldots i_k},\delta_k)} |Df_k|^n = \int_{\mathbb{B}^n(p_{i_1\ldots i_k},\delta_k)} |D\gamma_{\delta_k}^{i_1\ldots i_k}|^n < 4^{-nk}.
$$

Hence

$$
\int_{W_k} |Df_k|^n = \sum_{i_1...i_k \in \Im^k} \int_{\overline{\mathbb{B}}^n(p_{i_1...i_k}, \delta_k)} |Df_k|^n < 2^k \cdot 4^{-nk} < 2^{-nk}
$$

so

$$
\sum_{k=1}^{\infty} \int_{W_k} |Df_k|^n < \sum_{k=1}^{\infty} 2^{-nk} < \infty.
$$

This completes the proof that $f \in W^{1,n}$.

Remark 4.2. Replacing the estimate in (4.1) by $\varepsilon 4^{-nk}$ one can easily modify the construction so that the mapping f will have an arbitrarily small Sobolev norm $W^{1,n}$.

It remains to prove that f is continuous. We will need

Lemma 4.3. $f(\overline{\mathbb{B}}^n(p_{i_1...i_k}, \delta_k))$ is contained in the

$$
r_k := 2^{-k+4} + \text{diam } C_{i_1...i_{k-1}}
$$

neighborhood of $C_{i_1...i_{k-1}}$.

Proof. If $x \in \overline{\mathbb{B}}^n(p_{i_1...i_k}, \delta_k) \cap E$, then $x = e_i$ for some $i \in \mathfrak{I}^\infty$ with the first k binary digits equal i_1, \ldots, i_k i.e., $i = i_1 \ldots i_k \ldots$ Hence

$$
f(x) = f(e_i) = c_i = c_{i_1...i_k...} \in C_{i_1...i_{k-1}}.
$$

If $x \in \overline{\mathbb{B}}^n(p_{i_1...i_k}, \delta_k) \setminus E$, then there is $s \geq k$ such that

$$
x \in \overline{\mathbb{B}}^n(p_{i_1 \dots i_k i_{k+1} \dots i_s}, \delta_s) \setminus W_{s+1} \subset W_s \setminus W_{s+1}
$$

for some binary numbers i_{k+1}, \ldots, i_s .

Since $f = f_s$ in $W_s \setminus W_{s+1}$ and $f_s = \gamma_{\delta_s}^{i_1...i_s}$ $\overline{\mathbb{B}}^n(p_{i_1...i_s}, \delta_s)$ we conclude that $f(x) =$ $\gamma^{i_1...i_s}_{\delta_s}$ $\delta_s^{i_1...i_s}(x)$.

It remains to show that the image of $\gamma_{\delta_s}^{i_1...i_s}$ $\delta_s^{i_1...i_s}$ is contained in the r_k neighborhood of $C_{i_1...i_{k-1}}.$

By Lemma 2.1, $J_{i_1...i_s}$ is in the 2^{-s+3} + diam $C_{i_1...i_{s-1}}$ neighborhood of $C_{i_1...i_{s-1}}$. Also the image of $\gamma_{\delta_{s}}^{i_{1}...i_{s}}$ $\delta_s^{i_1...i_s}$ is contained in the 2^{-s} neighborhood of $J_{i_1...i_s}$ so the image of $\gamma_{\delta_s}^{i_1...i_s}$ $\frac{i_{1}\dots i_{s}}{\delta_{s}}$ is contained in the

$$
2^{-s+4} + \operatorname{diam} C_{i_1 \dots i_{s-1}}
$$

neighborhood of $C_{i_1...i_{s-1}}$. Since $C_{i_1...i_{s-1}} \subset C_{i_1...i_{k-1}}$ and

$$
2^{-s+4} + \operatorname{diam} C_{i_1 \dots i_{s-1}} \le 2^{-k+4} + \operatorname{diam} C_{i_1 \dots i_{k-1}}
$$

the lemma follows.

Now we are ready to complete the proof of continuity of f . Clearly, f is continuous on $\mathbb{S}^n \backslash E$ so it remains to prove its continuity on the Cantor set E. Let $e_i \in E$, $i = i_1 i_2 ... \in \mathfrak{I}^\infty$. Since $f(e_i) = c_i$ we need to show that for any $\varepsilon > 0$ there is $\delta > 0$ such that if $|x - e_i| < \delta$, then $|f(x) - c_i| < \varepsilon$.

Let $\varepsilon > 0$ be given. Let k be so large that

$$
2^{-k+4} + 2 \operatorname{diam} C_{i_1 \dots i_{k-1}} < \varepsilon
$$

and let $\delta > 0$ be so small that

$$
\mathbb{B}^n(e_i,\delta) \subset \mathbb{B}^n(p_{i_1...i_k},\delta_k).
$$

If $|x-e_i| < \delta$, then $x \in \overline{\mathbb{B}}^n(p_{i_1...i_k}, \delta_k)$ so by Lemma 4.3, $f(x)$ belongs to the r_k neighborhood of the set $C_{i_1...i_{k-1}}$. Since $c_i \in C_{i_1...i_{k-1}}$, the distance $|f(x) - c_i|$ is less than

$$
r_k + \text{diam } C_{i_1 \dots i_{k-1}} = 2^{-k+4} + 2 \text{ diam } C_{i_1 \dots i_{k-1}} < \varepsilon.
$$

The proof is complete. \Box

5. Proof of Theorem 1.7

Antoine's necklace A is a Cantor set that is constructed iteratively as follows: Inside a solid torus A_0 in \mathbb{R}^3 (iteration 0) we construct a chain A_1 (iteration 1) of linked solid tori so that the chain cannot be contracted to a point inside the torus A_0 . A_1 is a subset of \mathbb{R}^3 , the union of the linked tori. Iteration $n+1$ is obtained from the iteration n by constructing a chain of tori inside each of the tori of A_n (i.e. inside each of the connected components of A_n). Again A_{n+1} is a subset of \mathbb{R}^3 – the union of all tori in this step of construction. We also assume that the maximum of the diameters of tori in iteration n converges to zero as *n* approaches to infinity. Antoine's necklace is the intersection $A = \bigcap_{n=0}^{\infty} A_n$. For more details we refer to [22].

Antoine's necklace has the following properties:

- $\mathbb{R}^3 \setminus A$ is not simply connected.
- For any $x \in A$ and any $r > 0$, $A \cap \mathbb{B}^3(x,r)$ contains Antoine's necklace.

The first property is well known [18, Chapter 18] while the second one is quite obvious: $\mathbb{B}^{3}(x,r)$ contains one of the tori T of one of the iterations (actually infinitely many of such tori) and $T \cap A$ is also Antoine's necklace because of the iterative nature of the procedure.

We also need the following observation.

Lemma 5.1. Let A be Antoine's necklace and let M be a smooth 2-dimensional surface in \mathbb{R}^3 . Then $A \cap M$ is contained in the closure of $A \setminus M$, $A \cap M \subset \overline{A \setminus M}$.

Proof. Suppose to the contrary that for some $x \in A \cap M$, and some $r > 0$ we have $\mathbb{B}^3(x,r) \cap (A \setminus M) = \emptyset$. By taking $r > 0$ sufficiently small we can assume that $\mathbb{B}^3(x,r) \cap M$ is diffeomorphic to a disc. More precisely, there is a diffeomorphism Φ of \mathbb{R}^3 which maps $\mathbb{B}^3(x,r) \cap M$ onto the ball $\mathbb{B}^2(0,2)$ in the xy-coordinate plane. Note that by the second property listed above $\mathbb{B}^3(x,r) \cap A$ contains Antoine's necklace denoted by \tilde{A} . Since $\tilde{A} \cap$ $(A \setminus M) = \emptyset$ we have $\tilde{A} \subset \mathbb{B}^3(x,r) \cap M$ so $\Phi(\tilde{A}) \subset \mathbb{B}^2(0,2)$ and $(\mathbb{B}^2(0,2) \times \mathbb{R}) \setminus \Phi(\tilde{A})$ is not simply connected. By $[18,$ Theorem 13.7 p.93 there is a homeomorphism h of the ball $\mathbb{B}^2(0,2)$ onto itself in such that $\Phi(\tilde{A})$ is mapped onto the standard ternary Cantor set $\mathfrak C$ on the x-axis. This homeomorphism can be trivially extended to a homeomorphism of $\mathbb{B}^2(0,2) \times \mathbb{R}$ by letting $H(x,y,z) = (h(x,y),z)$. Clearly the complement of $H(\Phi(\tilde{A}))$ in $\mathbb{B}^2(0,2) \times \mathbb{R}$ is not simply connected. On the other hand since $H(\Phi(\tilde{A})) = \mathfrak{C}$, the complement of this set is simply connected in $\mathbb{B}^2(0,2) \times \mathbb{R}$ which is a contradiction. \square

The key argument in our proof is the following result of Sher [22, Corollary 1] that we state as a lemma.

Lemma 5.2. There is an uncountable family of Antoine's necklaces $\{A_i\}_{i\in I}$ such that for any $i, j \in I$, $i \neq j$ there is no homeomorphism $h : \mathbb{R}^3 \to \mathbb{R}^3$ with the property that $h(A_i) = A_j.$

For each of the sets A_i , let $f_i : \mathbb{S}^2 \to \mathbb{R}^3$ be an embedding as in Theorem 1.1 with the property that $A_i \subset f_i(\mathbb{S}^2)$ and $f_i(\mathbb{S}^2) \setminus A_i$ is a smooth surface (but not closed). It remains to prove that for $i \neq j$ there is no homeomorphism $h : \mathbb{R}^3 \to \mathbb{R}^3$ such that $h(f_i(\mathbb{S}^2)) = f_j(\mathbb{S}^2)$. Suppose to the contrary that such a homeomorphism h exists. We will show that $h(A_i) = A_j$ which is a contradiction with Lemma 5.2.

Clearly,

$$
h(A_i) = (h(A_i) \cap A_j) \cup (h(A_i) \cap (f_j(\mathbb{S}^2) \setminus A_j)).
$$

Since $f_j(\mathbb{S}^2) \setminus A_j$ is a smooth surface, Lemma 5.1 yields

$$
h(A_i) \cap (f_j(\mathbb{S}^2) \setminus A_j) \subset \overline{h(A_i) \setminus (f_j(\mathbb{S}^2) \setminus A_j)} = \overline{h(A_i) \cap A_j} \subset A_j
$$

so $h(A_i) \cap (f_j(\mathbb{S}^2) \setminus A_j) = \emptyset$ and hence $h(A_i) \subset A_j$. Applying the same argument to h^{-1} we obtain that $h^{-1}(A_j) \subset A_i$ so $A_j \subset h(A_i)$ and hence $h(A_i) = A_j$. The proof is complete. \Box

REFERENCES

[1] Alexander, J. W.: An example of a simply connected surface bounding a region which is not simply connected. Proc. Nat. Acad. Sci., 10 (1924), 8-10.

SOBOLEV EMBEDDING 15

- [2] AUBIN, T.: Nonlinear analysis on manifolds. Monge-Ampère equations. Grundlehren der Mathematischen Wissenschaften, 252. Springer-Verlag, New York, 1982.
- [3] Besicovitch, A. S.: Parametric surfaces. Bull. Amer. Math. Soc. 56 (1950) 288–296.
- [4] BESICOVITCH, A. S.: On the definition and value of the area of a surface. *Quart. J. Math. Oxford* Ser. 16 (1945) 86–102.
- [5] Bojarski, B., Iwaniec, T.: Analytical foundations of the theory of quasiconformal mappings in R ⁿ. Ann. Acad. Sci. Fenn. Ser. A I Math. 8 (1983), 257–324.
- [6] Cesari, L.: Sulle trasformazioni continue. Ann. Mat. Pura Appl. 21 (1942), 157–188.
- [7] CSÖRNYEI, M., HENCL, S., MALÝ, J.: Homeomorphisms in the Sobolev space $W^{1,n-1}$. J. Reine Angew. Math. 644 (2010), 221–235.
- [8] Evans, L. C., Gariepy, R. F.: Measure theory and fine properties of functions. Studies in Advanced Mathematics. CRC Press, Boca Raton, FL, 1992.
- [9] GOLDSTEIN, P., HAJ LASZ, P.: A measure and orientation preserving homeomorphism of a cube with Jacobian equal −1 almost everywhere. arXiv:1510.05575, 2015.
- [10] HAJ LASZ, P., TYSON, J. T.: Sobolev Peano cubes. *Michigan Math. J.* 56 (2008), 687–702.
- [11] HAJLASZ, P., TYSON, J.: Lipschitz and Hölder Peano cubes and highly regular surjections between Carnot groups. Preprint.
- [12] HENCL, S., KOSKELA, P.: Lectures on mappings of finite distortion. Lecture Notes in Mathematics, 2096. Springer, Cham, 2014.
- [13] Hurewicz, W., Wallman, H.: Dimension Theory. Princeton Mathematical Series, v. 4. Princeton University Press, Princeton, N. J., 1941.
- [14] Iwaniec, T., Martin, G.: Geometric function theory and non-linear analysis. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, New York, 2001.
- [15] Iwaniec, T., Martin, G.: Squeezing the Sierpinski sponge. Studia Math. 149 (2002), 133–145.
- [16] KAUFMAN, R.: A singular map of a cube onto a square. *J. Differential Geom.* 14 (1979), 593–594 (1981).
- [17] MALÝ. J., MARTIO, O.: Lusin's condition (N) and mappings of the class $W^{1,n}$. J. Reine Angew. Math. 458 (1995), 19–36.
- [18] Moise, E. E.: Geometric topology in dimensions 2 and 3. Graduate Texts in Mathematics, Vol. 47. Springer-Verlag, New York-Heidelberg, 1977.
- [19] PONOMAREV, S. P.: On the N-property of homeomorphisms of the class W_p^1 . Sibirsk. Mat. Zh. 28 (1987), 140–148 (in Russian).
- [20] PONOMAREV, S. P.: An example of an $ACTL^p$ homeomorphism that is not absolutely continuous in the sense of Banach. Dokl. Akad. Nauk SSSR 201 (1971), 1053–1054 (in Russian).
- [21] RESHETNYAK, YU. G.: The N condition for spatial mappings of the class $W^{1,n}_{loc}$. Sibirsk. Mat. Zh. 28 (1987), 149–153 (in Russian).
- [22] SHER, R. B.: Concerning wild Cantor sets in E^3 . Proc. Amer. Math. Soc. 19 (1968) 1195-1200.
- [23] VÄISÄLÄ, J.: Quasiconformal maps and positive boundary measure. Analysis 9 (1989), 205–216.
- [24] Wenger, S., Young, R.: Lipschitz homotopy groups of the Heisenberg groups. Geom. Funct. Anal. 24 (2014), 387–402.
- [25] WILDRICK, K., ZÜRCHER, T.: Space filling with metric measure spaces. *Math. Z.* 270 (2012), 103– 131.
- [26] WILDRICK, K., ZÜRCHER, T.: Peano cubes with derivatives in a Lorentz space. Illinois J. Math. 53 (2009), 365–378.
- [27] WILLARD, S.: General topology. Addison-Wesley Publishing Co., 1970.

Piotr Hajlasz, Department of Mathematics, University of Pittsburgh, Pittsburgh, PA 15260, USA, hajlasz@pitt.edu

Xiaodan Zhou, Department of Mathematics, University of Pittsburgh, Pittsburgh, PA 15260, USA, xiz78@pitt.edu