

Tight uniform continuity bounds for quantum entropies: conditional entropy, relative entropy distance and energy constraints

Andreas Winter^{1,*}

¹*ICREA & Física Teòrica: Informació i Fenòmens Quàntics,
Universitat Autònoma de Barcelona, ES-08193 Bellaterra (Barcelona), Spain*

We present a bouquet of continuity bounds for quantum entropies, falling broadly into two classes: First, a tight analysis of the Alicki-Fannes continuity bounds for the conditional von Neumann entropy, reaching almost the best possible form that depends only on the system dimension and the trace distance of the states. Almost the same proof can be used to derive similar continuity bounds for the relative entropy distance from a convex set of states or positive operators.

Second, analogous continuity bounds for the von Neumann entropy and conditional entropy in infinite dimensional systems under an energy constraint, most importantly systems of multiple quantum harmonic oscillators. While without an energy bound the entropy is discontinuous, it is well-known to be continuous on states of bounded energy. However, a quantitative statement to that effect seems not to have been known. Here, under some regularity assumptions on the Hamiltonian, we find that, quite intuitively, the Gibbs entropy at the given energy roughly takes the role of the Hilbert space dimension in the finite-dimensional Fannes inequality.

I. INTRODUCTION

On finite dimensional systems, the von Neumann entropy is continuous, but this becomes useful only once one has explicit continuity bounds, most significantly the one due to Fannes [9], the sharpest form of which is the following:

Lemma 1 (Audenaert [2], Petz [21]) *For states ρ and σ on a Hilbert space H of dimension $d = |H| < \infty$, if $\frac{1}{2}\|\rho - \sigma\|_1 \leq \epsilon \leq 1$, then*

$$|S(\rho) - S(\sigma)| \leq \epsilon \log(d-1) + h(\epsilon),$$

with $h(x) = H(x, 1-x) = -x \log x - (1-x) \log(1-x)$ the binary entropy.

We include a short proof for self-containedness, and also because it deserves to be known better. It seems that it was first found by Petz [21, Thm. 3.8], who attributes its central idea to Csiszár.

Proof. We begin with the classical case of two probability distributions p and q on the same ground set of d elements. It is well known that one can find two random variables, $X \sim p$ and $Y \sim q$, with $\Pr\{X \neq Y\} = \frac{1}{2}\|p - q\|_1 \leq \epsilon$ (and this is the minimum probability over all such random variables). This is an elementary special case of the Kantorovich-Rubinshtein dual formula for the Wasserstein distance in the case of the trivial metric $d(x, y) = 1$ for all $x \neq y$ and $d(x, x) = 0$ (cf. the broad survey [3]). Then, by the monotonicity of the Shannon entropy under taking marginals and Fano's inequality (see [5]),

$$\begin{aligned} H(X) - H(Y) &\leq H(XY) - H(Y) \\ &= H(X|Y) \leq \epsilon \log(d-1) + h(\epsilon), \end{aligned}$$

and likewise for $H(Y) - H(X)$.

Reduction of the quantum case to the classical case: W.l.o.g. $S(\rho) \leq S(\sigma)$, and consider the dephasing operation E in the eigenbasis of ρ , which maps ρ to itself, a diagonal matrix with a probability distribution p along the diagonal, and σ to $E(\sigma)$, a diagonal matrix with a probability distribution q along the diagonal. Hence

$$H(p) = S(\rho) \leq S(\sigma) \leq S(E(\sigma)) = H(q).$$

At the same time, $\|p - q\|_1 = \|E(\rho) - E(\sigma)\|_1 \leq \|\rho - \sigma\|_1$, and so, using the classical case,

$$|S(\rho) - S(\sigma)| \leq H(q) - H(p) \leq \epsilon \log(d-1) + h(\epsilon).$$

Note that the inequality is tight for every ϵ and d , for example choosing $\sigma = |0\rangle\langle 0|$ and $\rho = (1-\epsilon)|0\rangle\langle 0| + \frac{\epsilon}{d-1}(\mathbb{1} - |0\rangle\langle 0|)$. \square

We are interested in bounds of the above form, i.e. only referring to the trace distance of the states and some general global parameter specifying the system, for a number of entropic quantities, starting with the conditional von Neumann entropy, relative entropy distances from certain sets, etc, which have numerous applications in quantum information theory and quantum statistical physics. Furthermore, and perhaps even more urgently, in situations of infinite dimensional Hilbert spaces, where the above form of the Fannes inequality becomes trivial.

The rest of the paper is structured as follows: in Section II we present and prove an almost tight version of Lemma 1 for the conditional entropy (originally due to Alicki and Fannes [1]), then in Section III we generalize the principle behind our proof to a family of relative entropy distance measures from a convex set. In Section IV we expand the methodology of the first part of the paper to infinite dimensional systems, where Fannes-type continuity bounds are obtained under an energy constraint for a broad class of Hamiltonians, and specifically for quantum harmonic oscillators.

* andreas.winter@uab.cat

II. CONDITIONAL ENTROPY

Alicki and Fannes [1] proved an extension of the Fannes inequality for the conditional entropy

$$S(A|B)_\rho = S(\rho^{AB}) - S(\rho^B),$$

defined for states ρ on a bipartite (tensor product) Hilbert space $A \otimes B$. While a double application of Lemma 1 would yield such a bound involving both the dimensions of A and B , Alicki and Fannes show that if $\|\rho - \sigma\|_1 \leq \epsilon \leq 1$, then

$$|S(A|B)_\rho - S(A|B)_\sigma| \leq 4\epsilon \log |A| + 2h(\epsilon).$$

In particular, this form is independent of the dimension of B , which might even be infinite. Note that for classical, Shannon, conditional entropy, an inequality like the above can be obtained from Lemma 1 by convex combination.

The Alicki-Fannes inequality has several applications in quantum information theory, from the proof of asymptotic continuity of entanglement measures — most notably squashed entanglement [4] and conditional entanglement of mutual information (CEMI) [35] —, to the continuity of quantum channel capacities [16], and on to the recent discussion of approximately degradable channels [28].

We present a simple proof of the Alicki-Fannes inequality that yields the following stronger version.

Lemma 2 *For states ρ and σ on a Hilbert space $A \otimes B$, if $\frac{1}{2}\|\rho - \sigma\|_1 \leq \epsilon \leq 1$, then*

$$|S(A|B)_\rho - S(A|B)_\sigma| \leq 2\epsilon \log |A| + (1 + \epsilon)h\left(\frac{\epsilon}{1 + \epsilon}\right).$$

Proof. The right hand side is monotonic in ϵ , hence we may assume $\frac{1}{2}\|\rho - \sigma\|_1 = \epsilon$. Let $\epsilon\Delta = (\rho - \sigma)_+$ be the positive part of $\rho - \sigma$. Note that because of the traceless of this difference and its trace norm being 2ϵ , Δ is a bona fide state. Furthermore,

$$\begin{aligned} \rho &= \sigma + (\rho - \sigma) \\ &\leq \sigma + \epsilon\Delta \\ &= (1 + \epsilon)\left(\frac{1}{1 + \epsilon}\sigma + \frac{\epsilon}{1 + \epsilon}\Delta\right) \\ &=: (1 + \epsilon)\omega. \end{aligned}$$

By letting $\epsilon\Delta' := (1 + \epsilon)\omega - \rho$, we obtain another state Δ' , such that

$$\omega = \frac{1}{1 + \epsilon}\sigma + \frac{\epsilon}{1 + \epsilon}\Delta = \frac{1}{1 + \epsilon}\rho + \frac{\epsilon}{1 + \epsilon}\Delta'. \quad (1)$$

This is a slightly optimized version of the trick in the proof of Alicki and Fannes [1].

Now, we use the following well-known variational characterization of the conditional entropy:

$$-S(A|B)_\omega = \min_{\xi} D(\omega^{AB} \| \mathbb{1}^A \otimes \xi^B).$$

Choosing an optimal state ξ for ω (which is $\xi = \omega^B$), we have, from Eq. (1),

$$\begin{aligned} S(A|B)_\omega &= -D(\omega^{AB} \| \mathbb{1}^A \otimes \xi^B) \\ &= S(\omega) + \text{Tr } \omega \log \xi^B \\ &\leq h\left(\frac{\epsilon}{1 + \epsilon}\right) + \frac{1}{1 + \epsilon}S(\rho) + \frac{\epsilon}{1 + \epsilon}S(\Delta') \\ &\quad + \frac{1}{1 + \epsilon} \text{Tr } \rho \log \xi^B + \frac{\epsilon}{1 + \epsilon} \text{Tr } \Delta' \log \xi^B \\ &= h\left(\frac{\epsilon}{1 + \epsilon}\right) - \frac{1}{1 + \epsilon}D(\rho \| \mathbb{1} \otimes \xi) - \frac{\epsilon}{1 + \epsilon}D(\Delta' \| \mathbb{1} \otimes \xi) \\ &\leq h\left(\frac{\epsilon}{1 + \epsilon}\right) + \frac{1}{1 + \epsilon}S(A|B)_\rho + \frac{\epsilon}{1 + \epsilon}S(A|B)_{\Delta'}. \end{aligned}$$

Using the other decomposition in Eq. (1), the concavity of the conditional entropy (equivalent to strong subadditivity of the von Neumann entropy) [17] gives

$$S(A|B)_\omega \geq \frac{1}{1 + \epsilon}S(A|B)_\sigma + \frac{\epsilon}{1 + \epsilon}S(A|B)_{\Delta}.$$

Putting these two bounds together and multiplying by $1 + \epsilon$, we arrive at

$$\begin{aligned} S(A|B)_\sigma - S(A|B)_\rho &\leq \epsilon(S(A|B)_{\Delta'} - S(A|B)_{\Delta}) \\ &\quad + (1 + \epsilon)h\left(\frac{\epsilon}{1 + \epsilon}\right). \end{aligned}$$

The proof is concluded observing that the conditional entropy of any state is bounded between $-\log |A|$ and $+\log |A|$. \square

Remark 3 Lemma 2 is almost best possible, as we can see by considering the example of $\sigma^{AB} = \Phi_d$, the maximally entangled state on $A = B = \mathbb{C}^d$, and $\rho^{AB} = (1 - \epsilon)\Phi_d + \frac{\epsilon}{d^2 - 1}(\mathbb{1} - \Phi_d)$. Clearly, $\frac{1}{2}\|\rho - \sigma\|_1 = \epsilon$, while

$$\begin{aligned} S(A|B)_\rho - S(A|B)_\sigma &= (\epsilon \log(d^2 - 1) + h(\epsilon) - \log d) - (-\log d) \\ &= 2\epsilon \log d + h(\epsilon) - O\left(\frac{\epsilon}{d^2}\right). \end{aligned}$$

This asymptotically matches Lemma 2 for large d and small ϵ .

III. RELATIVE ENTROPY DISTANCES

The same method employed in Lemma 2 can be used to derive asymptotic continuity bounds for the relative entropy distance with respect to any closed convex set C of states, or more generally positive semidefinite operators, on a Hilbert space H , cf. [30]),

$$D_C(\rho) = \min_{\gamma \in C} D(\rho \| \gamma). \quad (2)$$

Unlike [30], C has to contain only at least one full-rank state, so that D_C is guaranteed to be finite; in addition, C should be bounded, so that D_C is bounded from below.

Lemma 4 For a closed, convex and bounded set C of positive semidefinite operators, containing at least one of full rank, let

$$\kappa := \sup_{\tau, \tau'} D_C(\tau) - D_C(\tau')$$

be the largest variation of D_C . Then, for any two states ρ and σ with $\frac{1}{2}\|\rho - \sigma\|_1 \leq \epsilon$,

$$|D_C(\rho) - D_C(\sigma)| \leq \epsilon \kappa + (1 + \epsilon) h\left(\frac{\epsilon}{1 + \epsilon}\right). \quad (3)$$

Proof. The only modification with respect to the proof of Lemma 2 is that we replace the invocation of concavity of the conditional entropy by the joint convexity of the relative entropy, which makes D_C a convex functional.

Namely, with ω as in Eq. (1), we have on the one hand,

$$D_C(\omega) \leq \frac{1}{1 + \epsilon} D_C(\sigma) + \frac{\epsilon}{1 + \epsilon} D_C(\Delta).$$

On the other hand, with an optimal $\gamma \in C$,

$$\begin{aligned} D_C(\omega) &= D(\omega \parallel \gamma) \\ &= -S(\omega) - \text{Tr } \omega \log \gamma \\ &\geq -h\left(\frac{\epsilon}{1 + \epsilon}\right) - \frac{1}{1 + \epsilon} S(\rho) - \frac{\epsilon}{1 + \epsilon} S(\Delta') \\ &\quad - \frac{1}{1 + \epsilon} \text{Tr } \rho \log \gamma - \frac{\epsilon}{1 + \epsilon} \text{Tr } \Delta' \log \gamma \\ &= -h\left(\frac{\epsilon}{1 + \epsilon}\right) + \frac{1}{1 + \epsilon} D(\rho \parallel \gamma) + \frac{\epsilon}{1 + \epsilon} D(\Delta' \parallel \gamma) \\ &\geq -h\left(\frac{\epsilon}{1 + \epsilon}\right) + \frac{1}{1 + \epsilon} D_C(\rho) + \frac{\epsilon}{1 + \epsilon} D_C(\Delta'). \end{aligned}$$

Putting these two inequalities together yields the claim of the lemma. \square

In particular, for the relative entropy of entanglement of a bipartite system $A \otimes B$, denoting the smaller of the two dimensions by d (cf. Donald and Horodecki [7]), we have:

Corollary 5 For any two states ρ and σ , $\frac{1}{2}\|\rho - \sigma\|_1 \leq \epsilon$ implies

$$|E_R(\rho) - E_R(\sigma)| \leq \epsilon \log d + (1 + \epsilon) h\left(\frac{\epsilon}{1 + \epsilon}\right). \quad \square$$

Note that this bound only depends on the smaller of the two dimensions, in contrast to [7]; in particular, it applies even in the case that one of the two Hilbert spaces is infinite dimensional.

Again, in Lemma 4 and Corollary 5, the constant in the linear term (proportional to ϵ) is best possible, as we see by taking two states maximizing the difference $D_C(\rho) - D_C(\sigma)$, i.e. attaining κ , since $\frac{1}{2}\|\rho - \sigma\|_1 \leq 1 =: \epsilon$.

Remark 6 Lemma 4 improves upon similar-looking general bounds by Synak-Radtke and Horodecki [30], which were subsequently optimized by Mosonyi and Hiai [18, Prop. VI.1]. The latter paper also explains lucidly (in Sec. VI) that the coefficient $\frac{1}{1 + \epsilon}$ in the convex decomposition of ω in two ways, into ρ and Δ' and into σ and Δ , is optimal, and gives a nice geometric interpretation of ω as a max-relative entropy center of ρ and σ (cf. [14]). Thus, at least following the same strategy one cannot improve the bound any more.

Remark 7 It would be interesting to lift the restriction that C has to be a convex set: Natural examples are the case that C is the set of all product states in a bipartite (multipartite) system, in which case D_C becomes the quantum mutual information (multi-information); or the case that C is the closure of the set of all Gibbs states for a suitable Hamiltonian operator H ,

$$C = \overline{\left\{ \frac{1}{\text{Tr } e^{-\beta H}} e^{-\beta H} : \beta > 0 \right\}}.$$

Both examples have in common that C is an exponential family (or the closure of one); it is known that at least in some cases D_C is continuous, but counterexamples of discontinuous behaviour are known [33].

IV. BOUNDED ENERGY

If the Hilbert space in the Fannes inequality (Lemma 1) has infinite dimension, or likewise A in the Alicki-Fannes inequality (Lemma 2), then the bound becomes trivial: the right hand side is infinite. This is completely natural, since the entropy is not even continuous, and these Fannes-type bounds imply a sort of uniform continuity. Continuity is restored, however, when restricting to states of finite energy, for instance of a quantum harmonic oscillator [32], see also [8] and [23] for more recent results and excellent surveys on the status of continuity of the entropy. Shirokov [24] has developed an approach to prove (local) continuity of entropic quantities, based on certain finite entropy assumptions, in which he uses Alicki-Fannes inequalities on finite approximations.

Uniform bounds are still out of the question, but what we shall show here is that the Fannes and Alicki-Fannes inequalities discussed above have satisfying analogues, with a dependence on the energy of the states rather than the Hilbert space dimension.

Abstractly, our setting is this: Consider a Hamiltonian H on a infinite dimensional separable Hilbert space A . If there is another system B and we consider bipartite states and conditional entropy, we implicitly assume trivial Hamiltonian on B , i.e. global Hamiltonian $H = H^A \otimes \mathbb{1}^B$. We shall need a number of assumptions on H , to start with that it has discrete spectrum and that it is bounded from below; for normalization purposes we fix the ground state energy of H to be 0. The mathematically precise assumption is the following.

Gibbs Hypothesis. For every $\beta > 0$, $Z(\beta) := \text{Tr} e^{-\beta H} < \infty$, so that $\frac{1}{Z(\beta)} e^{-\beta H}$ is a bona fide state, which we demand to have finite entropy. In this case, for every energy E in the spectrum of H , the (unique) maximizer of the entropy $S(\rho)$ subject to $\text{Tr} \rho H \leq E$ is of this form:

$$\gamma(E) = \frac{1}{Z(\beta(E))} e^{-\beta(E)H},$$

where $\beta = \beta(E)$ is decreasing with E and is the solution to the equation

$$\text{Tr} e^{-\beta H} (H - E) = 0.$$

The entropy in this case is given by

$$S(\gamma(E)) = \log Z + \beta(E)(\log e)E.$$

This implies that the spectrum is unbounded above, and that the energy levels cannot become “too dense” with growing energy value.

Let us immediately draw some conclusions from these assumptions.

Proposition 8 *For a Hamiltonian H satisfying the Gibbs Hypothesis, $S(\gamma(E))$ is a strictly increasing, strictly concave function of the energy E .*

Proof. It is clear from the maximum entropy characterization of $\gamma(E)$ that the entropy as a function of E must be non-decreasing; it is unbounded by looking at the formula for the entropy in terms of $\log Z$.

Furthermore, for energies E_1 and E_2 , and $0 \leq p \leq 1$,

$$\text{Tr}(p\gamma(E_1) + (1-p)\gamma(E_2))H \leq pE_1 + (1-p)E_2 =: E,$$

and so concavity follows:

$$\begin{aligned} S(\gamma(E)) &\geq S(p\gamma(E_1) + (1-p)\gamma(E_2)) \\ &\geq pS(\gamma(E_1)) + (1-p)S(\gamma(E_2)). \end{aligned} \quad (4)$$

From this it follows that $S(\gamma(E))$ is strictly increasing, because otherwise $S(\gamma(E_1)) = S(\gamma(E_2))$ for some $E_1 < E_2$, but then $S(\gamma(E_2)) < S(\gamma(E_3))$ for some $E_2 < E_3$, since the entropy grows to infinity as $E \rightarrow \infty$, contradicting concavity.

But this means that for $E_1 \neq E_2$, necessarily $\gamma(E_1) \neq \gamma(E_2)$, and so by the strict concavity of the von Neumann entropy, we have strict inequality in the second line of Eq. (4) for $0 < p < 1$. \square

Corollary 9 *If H satisfies the Gibbs Hypothesis, then for any $\delta > 0$,*

$$\sup_{0 < \lambda \leq \delta} \lambda S(\gamma(E/\lambda)) = \delta S(\gamma(E/\delta)).$$

Proof. The right hand side is clearly attained by letting $\lambda = \delta$. To prove “ \leq ” for any admissible λ , observe that by concavity (Proposition 8),

$$S(\gamma(tF)) \geq tS(\gamma(F)) + (1-t)S(\gamma(0)) \geq tS(\gamma(F)).$$

Letting $t = \frac{\lambda}{\delta} \leq 1$ and $F = \frac{E}{\lambda}$ concludes the proof. \square

We start with an easy-to-prove continuity bound for the entropy, inspired by the proof of Lemma 1, though for the conditional entropy we shall have to resort to a different argument.

Proposition 10 *Let ρ and σ be states on the same Hilbert space A . Then, there exist a state ω on $A \otimes A =: A_1 A_2$ such that $\omega^{A_1} = \rho$, $\omega^{A_2} = \sigma$, and*

$$\|\omega\|_\infty \geq 1 - \frac{1}{2}\|\rho - \sigma\|_1.$$

Proof. Choose spectral decompositions

$$\begin{aligned} \rho &= \sum_i r_i |e_i\rangle\langle e_i|, \\ \sigma &= \sum_i s_i |f_i\rangle\langle f_i|, \end{aligned}$$

of the two states, with $r_1 \geq r_2 \geq \dots$ and $s_1 \geq s_2 \geq \dots$; then, the ℓ^1 -distance between the probability vectors (r_i) and (s_i) is not larger than the trace distance between ρ and σ :

$$\|\rho - \sigma\|_1 \geq \|(r_i) - (s_i)\|_1 =: 2\epsilon.$$

(This is known as Mirsky’s inequality [13, Cor. 7.4.9.3].) Defining a vector

$$|\phi\rangle := \sum_i \sqrt{\min\{r_i, s_i\}} |e_i\rangle^{A_1} |f_i\rangle^{A_2}$$

in $A_1 A_2$, we clearly have $\text{Tr} |\phi\rangle\langle\phi| = 1 - \epsilon$, and $\phi^{A_1} \leq \rho$, $\phi^{A_2} \leq \sigma$, thus we can write

$$\rho = |\phi\rangle\langle\phi|^{A_1} + \epsilon\Delta_1, \quad \sigma = |\phi\rangle\langle\phi|^{A_2} + \epsilon\Delta_2,$$

with bona fide states Δ_1 and Δ_2 .

The proof is concluded by checking that the definition $\omega := |\phi\rangle\langle\phi| + \epsilon\Delta_1 \otimes \Delta_2$ satisfies all requirements. \square

Lemma 11 *Let the Hamiltonian H on A satisfying the Gibbs Hypothesis. Then for any two states ρ and σ on A with $\text{Tr} \rho H, \text{Tr} \sigma H \leq E$ and $\frac{1}{2}\|\rho - \sigma\|_1 \leq \epsilon \leq 1$,*

$$|S(\rho) - S(\sigma)| \leq 2\epsilon S(\gamma(E/\epsilon)) + h(\epsilon).$$

Proof. Pick a state ω on $A_1 A_2$, according to Proposition 10: $\omega^{A_1} = \rho$, $\omega^{A_2} = \sigma$, and with largest eigenvalue $\geq 1 - \epsilon$, meaning that we can write

$$\omega = (1 - \epsilon)|\psi\rangle\langle\psi| + \epsilon\omega',$$

with a pure state $|\psi\rangle$ (the normalized vector $|\phi\rangle$ from the proof of Proposition 10) and some other state ω' . Hence,

$$\begin{aligned} |S(\rho) - S(\sigma)| &= |S(\omega^{A_1}) - S(\omega^{A_2})| \\ &\leq S(\omega^{A_1 A_2}) \\ &\leq \epsilon S(\omega') + h(\epsilon) \\ &\leq 2\epsilon S(\gamma(E/\epsilon)) + h(\epsilon). \end{aligned}$$

Here, we have first used the marginals of ω , then in the second line the Araki-Lieb ‘‘triangle’’ inequality, in the third line strong subadditivity, and in the last step the maximum entropy principle, noting that with respect to the Hamiltonian $H^{A_1} \otimes \mathbb{1}^{A_2} + \mathbb{1}^{A_1} \otimes H^{A_2}$, ω has energy $\leq 2E$, and so the energy of ω' is bounded by $2E/\epsilon$. For the last line, observe that the Gibbs state at energy $2E/\epsilon$ of the composite system is $\gamma(E/\epsilon)^{\otimes 2}$. \square

The following two general bounds lack perhaps the simple elegance of Lemma 11, but they turn out to be more flexible, and stronger in certain regimes.

Meta-Lemma 12 (Entropy) *For a Hamiltonian H on A satisfying the Gibbs Hypothesis and any two states ρ and σ with $\text{Tr } \rho H, \text{Tr } \sigma H \leq E$, $\frac{1}{2}\|\rho - \sigma\|_1 \leq \epsilon < \epsilon' \leq 1$, and $\delta = \frac{\epsilon' - \epsilon}{1 + \epsilon'}$,*

$$|S(\rho) - S(\sigma)| \leq (\epsilon' + 2\delta)S(\gamma(E/\delta)) + h(\epsilon') + h(\delta).$$

Meta-Lemma 13 (Conditional entropy) *For states ρ and σ on the bipartite system $A \otimes B$ and otherwise the same assumption as before,*

$$\begin{aligned} |S(A|B)_\rho - S(A|B)_\sigma| &\leq (2\epsilon' + 4\delta)S(\gamma(E/\delta)) \\ &\quad + (1 + \epsilon')h\left(\frac{\epsilon'}{1 + \epsilon'}\right) + h(\delta). \end{aligned}$$

To interpret these bounds, we remark that in a certain sense they show that the Gibbs entropy at the cutoff energy E/ϵ (E/δ) takes on the role of the logarithm of the dimension in the finite dimensional case. Before we launch into their proof, let us introduce some notation: Define the energy cutoff projectors

$$P_{\leq} := \sum_{0 \leq E_n \leq E/\delta} |n\rangle\langle n|, \quad P_{>} := \mathbb{1} - P_{\leq},$$

where $|n\rangle$ is the eigenvector of eigenvalue E_n of the Hamiltonian H . We shall also consider the pinching map

$$\mathcal{T}(\xi) = P_{\leq} \xi P_{\leq} + P_{>} \xi P_{>},$$

which is a unital channel, as well as its action on the original ρ and σ :

$$\begin{aligned} \mathcal{T}(\rho) &=: (1 - \lambda)\rho_{\leq} + \lambda\rho_{>}, \\ \mathcal{T}(\sigma) &=: (1 - \mu)\sigma_{\leq} + \mu\sigma_{>}. \end{aligned}$$

Note that because H commutes with the action of \mathcal{T} , we have $\text{Tr } \xi H = \text{Tr } \mathcal{T}(\xi)H$, and so the energy bound E applies also to $\mathcal{T}(\rho)$ and $\mathcal{T}(\sigma)$. Hence,

$$\lambda \leq \delta, \quad \lambda \text{Tr } \rho_{>} H \leq E, \quad \mu \leq \delta, \quad \mu \text{Tr } \sigma_{>} H \leq E. \quad (5)$$

Our strategy will be to relate $S(\rho)$ to $S(\rho_{\leq})$ (and the same for σ and σ_{\leq}) via entropy inequalities, including concavity, similar to the first part of the paper, and then apply the usual Fannes (Alicki-Fannes) inequalities to ρ_{\leq} and σ_{\leq} .

Proof of Lemma 12. First of all, by concavity of the entropy (monotonicity under unital cptp maps),

$$\begin{aligned} S(\rho) &\leq S(\mathcal{T}(\rho)) \\ &= h(\lambda) + (1 - \lambda)S(\rho_{\leq}) + \lambda S(\rho_{>}). \end{aligned} \quad (6)$$

Now, by Eq. (5), the maximum entropy principle and Corollary 9,

$$\lambda S(\rho_{>}) \leq \lambda S(\gamma(E/\lambda)) \leq \delta S(\gamma(E/\delta)).$$

Thus, from Eq. (6), observing $\delta \leq \frac{1}{2}$, we get

$$S(\rho) \leq S(\rho_{\leq}) + h(\delta) + \delta S(\gamma(E/\delta)), \quad (7)$$

and likewise for σ .

Second, we have

$$S(\sigma) \geq (1 - \mu)S(\sigma_{\leq}) + \mu S(\sigma_{>}). \quad (8)$$

To see this, we think of the action of \mathcal{T} as a binary measurement on the system A , which we can implement coherently with two ancilla qubits X and X' ,

$$|\varphi\rangle \mapsto (P_{\leq}|\varphi\rangle)^A |00\rangle^{XX'} + (P_{>}|\varphi\rangle)^A |11\rangle^{XX'}.$$

Applying this to σ , we have by unitary invariance and the Araki-Lieb triangle inequality,

$$\begin{aligned} S(\sigma) &= S(AXX') \geq S(AX) - S(X') \\ &= S(AX) - S(X) \\ &= S(A|X) = (1 - \mu)S(\sigma_{\leq}) + \mu S(\sigma_{>}). \end{aligned}$$

Thus, using that the energy of σ_{\leq} is at most E/δ by construction, and so $S(\sigma_{\leq}) \leq S(\gamma(E/\delta))$,

$$S(\sigma) \geq (1 - \mu)S(\sigma_{\leq}) \geq S(\sigma_{\leq}) - \delta S(\gamma(E/\delta)). \quad (9)$$

Third, by definitions, contractivity of the trace norm and triangle inequality,

$$\begin{aligned} 2\epsilon &\geq \|\rho - \sigma\|_1 \\ &\geq \|P_{\leq} \rho P_{\leq} - P_{\leq} \sigma P_{\leq}\|_1 \\ &= \|(1 - \lambda)\rho_{\leq} - (1 - \mu)\sigma_{\leq}\|_1 \\ &= \|(1 - \delta)(\rho_{\leq} - \sigma_{\leq}) + (\delta - \lambda)\rho_{\leq} + (\mu - \delta)\sigma_{\leq}\|_1 \\ &\geq (1 - \delta)\|\rho_{\leq} - \sigma_{\leq}\|_1 - 2\delta, \end{aligned}$$

and so

$$\frac{1}{2} \|\rho_{\leq} - \sigma_{\leq}\|_1 \leq \frac{\epsilon + \delta}{1 - \delta} = \epsilon'. \quad (10)$$

Hence by the Fannes inequality in the form of Lemma 1,

$$\begin{aligned} |S(\rho_{\leq}) - S(\sigma_{\leq})| &\leq \epsilon' \log \text{Tr } P_{\leq} + h(\epsilon') \\ &\leq \epsilon' S(\gamma(E/\delta)) + h(\epsilon'). \end{aligned} \quad (11)$$

The latter inequality holds because the state $\frac{1}{\text{Tr } P_{\leq}} P_{\leq}$ clearly has energy bounded by E/δ , and so cannot have entropy larger than the Gibbs state.

With these three elements we can conclude the proof: W.l.o.g. $S(\rho) \geq S(\sigma)$, and so from Eqs. (7), (9) and (11),

$$\begin{aligned} S(\rho) - S(\sigma) &\leq S(\rho_{\leq}) - S(\sigma_{\leq}) + h(\delta) + 2\delta S(\gamma(E/\delta)) \\ &\leq (\epsilon' + 2\delta)S(\gamma(E/\delta)) + h(\epsilon') + h(\delta), \end{aligned}$$

as advertised. \square

Proof of Lemma 13. It is very similar to the previous one, only that we have to be a bit more careful in some details, as the conditional entropy can be negative.

The first step goes through almost unchanged, with the map $\mathcal{T} \otimes \text{id}_B$, since the conditional entropy is concave as well (equivalent to strong subadditivity) [17]:

$$\begin{aligned} S(A|B)_{\rho} &\leq S(A|B)_{\mathcal{T}(\rho)} \\ &= h(\lambda) + (1 - \lambda)S(A|B)_{\rho_{\leq}} + \lambda S(A|B)_{\rho_{>}}. \end{aligned}$$

The remainder term $\lambda S(A|B)_{\rho_{>}}$ is upper bounded by $\lambda S(\rho_{>}^A)$ (again by strong subadditivity), hence the upper bound $\lambda S(\gamma(E/\lambda))$ still applies. The only change is due to the fact that the conditional entropy can be negative. However, for any bipartite state ξ^{AB} ,

$$-S(\xi^A) \leq S(A|B)_{\xi} \leq S(\xi^A).$$

Here, the right hand inequality is strong subadditivity that we have used before; introducing a purification $|\varphi\rangle^{ABC}$ of the state, we have $-S(A|B)_{\varphi} = S(A|C)_{\varphi} \leq S(\xi^A)$, which is the left hand inequality. Thus,

$$(1 - \lambda)S(A|B)_{\rho_{\leq}} \leq S(A|B)_{\rho_{\leq}} + \delta S(\gamma(E/\delta)).$$

Altogether,

$$S(A|B)_{\rho} \leq S(A|B)_{\rho_{\leq}} + 2\delta S(\gamma(E/\delta)) + h(\delta). \quad (12)$$

Also the second step requires only minor modifications, since the Araki-Lieb triangle inequality generalizes to

$$\begin{aligned} S(AXX'|B) &\geq S(AX|B) - S(X'|B) \\ &= S(AX|B) - S(X|B) = S(A|BX) \end{aligned}$$

(again, strong subadditivity). Once more, since conditional entropies can be negative, we have to be more careful with remainder terms and get

$$S(A|B)_{\sigma} \geq S(A|B)_{\sigma_{\leq}} - 2\delta S(\gamma(E/\delta)). \quad (13)$$

In the third step, the trace norm estimate (10) goes through unchanged, and then we apply the Alicki-Fannes inequality in the form of Lemma 2:

$$\begin{aligned} |S(A|B)_{\rho_{\leq}} - S(A|B)_{\sigma_{\leq}}| &\leq 2\epsilon' \log \text{Tr } P_{\leq} + (1 + \epsilon') h\left(\frac{\epsilon'}{1 + \epsilon'}\right) \\ &\leq 2\epsilon' S(\gamma(E/\delta)) + (1 + \epsilon') h\left(\frac{\epsilon'}{1 + \epsilon'}\right). \end{aligned}$$

Putting this together with Eqs. (12) and (13), assuming w.l.o.g. that $S(A|B)_{\rho} \geq S(A|B)_{\sigma}$, we obtain

$$\begin{aligned} S(A|B)_{\rho} - S(A|B)_{\sigma} &\leq S(A|B)_{\rho_{\leq}} - S(A|B)_{\sigma_{\leq}} \\ &\quad + h(\delta) + 4\delta S(\gamma(E/\delta)) \\ &\leq (2\epsilon' + 4\delta)S(\gamma(E/\delta)) \\ &\quad + (1 + \epsilon') h\left(\frac{\epsilon'}{1 + \epsilon'}\right) + h(\delta), \end{aligned}$$

and we are done. \square

The bounds of Lemmas 12 and 13 are very general, and it may not be immediately apparent how useful they are. We now specialize them to the important case of a collection of ℓ quantum harmonic oscillators, where we shall see that the bounds are not only meaningful, but asymptotically tight. The Hamiltonian is

$$H = \sum_{i=1}^{\ell} \hbar\omega_i a_i^{\dagger} a_i, \quad (14)$$

where ω_i is the native frequency of the i -th oscillator and a_i is its annihilation (aka lowering) operator (see e.g. [15] or [31]). Note that we chose the slightly unusual energy convention such that the ground state has energy 0, rather than $\sum_i \frac{1}{2} \hbar\omega_i$, to be able to apply directly our above results. In the case of a single mode, and choosing units such that $\hbar\omega_1 = 1$, the Hamiltonian simply becomes the number operator N . In that case, it is well-known that

$$\begin{aligned} S(\gamma(N)) &= g(N) := (N + 1) \log(N + 1) - N \log N \\ &\leq \log(N + 1) + \log e. \end{aligned}$$

Crucially, and in accordance with Proposition 8, g is a concave, monotone increasing function of N .

In the general case of Eq. (14), $\gamma(E) = \bigotimes_{i=1}^{\ell} \gamma_i(E_i)$, with $E = \sum_i E_i$ and where $\gamma_i(E_i)$ is the Gibbs state of the i -th mode with energy E_i . Maximizing the entropy,

$$S\left(\bigotimes_{i=1}^{\ell} \gamma_i(E_i)\right) = \sum_{i=1}^{\ell} g\left(\frac{E_i}{\hbar\omega_i}\right),$$

over all allocations of the total energy over the ℓ modes leads to a transcendental equation, but we do not need to solve it as we only want an upper bound, via $g(N) \leq$

$\log(N + 1) + \log e$. By a straightforward Lagrange multiplier calculation we see that the optimum is to divide the energy equally among the modes:

$$\begin{aligned} S(\gamma(E)) &\leq \max \sum_{i=1}^{\ell} \left[\log \left(\frac{E_i}{\hbar\omega_i} + 1 \right) + \log e \right] \\ &= (\log e)\ell + \sum_{i=1}^{\ell} \log \left(\frac{\bar{E}}{\hbar\omega_i} + 1 \right), \end{aligned} \quad (15)$$

with $E =: \ell\bar{E}$.

By using this upper bound in Lemmas 12 and 13, for $\delta = \alpha\epsilon(1 - \epsilon)$, with a parameter α between 0 and $\frac{1}{2}$, and introducing

$$\tilde{h}(x) := \begin{cases} h(x) & \text{for } x \leq \frac{1}{2}, \\ 1 & \text{for } x \geq \frac{1}{2}, \end{cases}$$

we obtain directly the following:

Lemma 14 *Consider two states ρ and σ of the ℓ -oscillator system (14), whose energies are bounded $\text{Tr } \rho H, \text{Tr } \sigma H \leq E = \ell\bar{E}$. Then, $\frac{1}{2}\|\rho - \sigma\|_1 \leq \epsilon < 1$ implies*

$$\begin{aligned} &|S(\rho) - S(\sigma)| \\ &\leq \epsilon \left(\frac{1 + \alpha}{1 - \alpha} + 2\alpha \right) \left[\sum_{i=1}^{\ell} \log \left(\frac{\bar{E}}{\hbar\omega_i} + 1 \right) + \ell \log \frac{e}{\alpha(1 - \epsilon)} \right] \\ &\quad + (\ell + 2) \left(\frac{1 + \alpha}{1 - \alpha} + 2\alpha \right) \tilde{h} \left(\frac{1 + \alpha}{1 - \alpha} \epsilon \right). \end{aligned}$$

If the states live on a system composed of the ℓ oscillators (A) and another system B, then

$$\begin{aligned} &|S(A|B)_\rho - S(A|B)_\sigma| \\ &\leq 2\epsilon \left(\frac{1 + \alpha}{1 - \alpha} + 2\alpha \right) \left[\sum_{i=1}^{\ell} \log \left(\frac{\bar{E}}{\hbar\omega_i} + 1 \right) + \ell \log \frac{e}{\alpha(1 - \epsilon)} \right] \\ &\quad + (2\ell + 3) \left(\frac{1 + \alpha}{1 - \alpha} + 2\alpha \right) \tilde{h} \left(\frac{1 + \alpha}{1 - \alpha} \epsilon \right). \quad \square \end{aligned}$$

Remark 15 For each fixed $\epsilon \leq 1$, we can make α arbitrarily small, and then for large energy $E \gg \sum_i \hbar\omega_i$, the bounds of Lemma 14 are asymptotically tight, in the sense that apart from the additive offset terms, the factor multiplying ϵ (2ϵ , resp.) cannot be smaller than

$$S(\gamma(E)) \approx \sum_{i=1}^{\ell} \log \left(\frac{\bar{E}}{\hbar\omega_i} + 1 \right).$$

This can be seen in the entropy case by comparing the vacuum state $\rho = |0\rangle\langle 0|^{\otimes \ell}$ of all ℓ modes with the state $\sigma = (1 - \epsilon)|0\rangle\langle 0|^{\otimes \ell} + \epsilon\gamma(E)$; in the conditional entropy case, take ρ to be a purification of the Gibbs state $\gamma(E)$ on $A \otimes B$, and $\sigma = (1 - \epsilon)\rho + \epsilon\gamma(E)^A \otimes \tau^B$ with an arbitrary state τ on B .

V. CONCLUSIONS

Using entropy inequalities, specifically concavity, we improved the appearance of the Alicki-Fannes inequality for the conditional von Neumann entropy to an almost tight form. It would be nice to know the ultimately best form among all formulas that depend only on the dimension of the Hilbert space and the trace distance, but we have to leave this as an open problem to the interested reader. In particular, it would be curious to find the optimal form of Proposition 10,

$$\frac{1}{2}\|\rho - \sigma\|_1 \geq \min_{\omega^{A_1}=\rho, \omega^{A_2}=\sigma} (1 - \|\omega\|_\infty)$$

which may be regarded as a quantum state analogue of the trick used in the proof of Lemma 1,

$$\frac{1}{2}\|p - q\|_1 = \min_{X \sim p, Y \sim q} \Pr\{X \neq Y\}.$$

Furthermore, are there versions of these statements that would allow for alternative proofs or tighter versions of Lemmas 2 and 13 for the conditional entropy?

The same principle lead to the apparently first uniform continuity bounds of the entropy and conditional on infinite dimensional Hilbert spaces under a bound on the expected energy (or, for that matter, bounded expectation of any sufficiently well-behaved Hermitian operator). In the case of a system of harmonic oscillators, we have seen that the bound is, in a certain sense, asymptotically tight, even though here we are much farther away from a universally optimal form.

The Fannes and Alicki-Fannes inequalities already are known to have many applications in quantum information theory. These include the continuity of certain entanglement measures such as entanglement of formation [19], relative entropy of entanglement [7], entanglement cost, squashed entanglement [4] and conditional entanglement of mutual information [35], and of various quantum channel capacities [16]. In fact, we invariably get explicit continuity bounds in terms of the trace distance of the state or diamond norm distance of the channels, resp. While in many applications it is of minor interest to have the optimal form of the bound (for example when ϵ goes to 0), it pays off to have a tighter bound than [1], in the setting of approximately degradable channels [28]. Using our new bound and a treatment of approximately degradable channel capacity modelled on [6], the bounds of Sutter *et al.* [28] on the regularized quantum and private capacity in terms of the single-letter coherent information can be improved significantly.

The infinite dimensional versions of these entropy bounds under an energy constraint are awaiting applications, though it seems clear that explicit bounds on the continuity and asymptotic continuity of entanglement measures [8, 25] and channel capacities [10–12, 22] in infinite dimension should be among the first, as well as the extension of the approximate degradability theory of Sutter *et al.* to Bosonic channels [29].

ACKNOWLEDGMENTS

Thanks to David Sutter and Volkher Scholz for stimulating discussions, and to Nihat Ay and Milan Mosonyi for comments on general relative entropy distances. The hospitality of the Banff International Research Station (BIRS) during the workshop “Beyond IID in Information Theory” (5-10 July 2016) is gratefully acknowledged,

where Volkher Scholz and David Sutter posed the derivation of infinite dimensional Fannes type inequalities as an open problem, and where the present work was initiated.

The author’s work was supported by the EU (STREP “RAQUEL”), the ERC (AdG “IRQUAT”), the Spanish MINECO (grants FIS2008-01236 & FIS2013-40627-P) with the support of FEDER funds, as well as by the Generalitat de Catalunya CIRIT, project 2014-SGR-966.

-
- [1] R. Alicki and M. Fannes, “Continuity of quantum conditional information”, *J. Phys. A: Math. Gen.* **37**:L55-L57 (2004).
- [2] K.M.R. Audenaert, “A sharp continuity estimate for the von Neumann entropy”, *J. Math. Phys. A: Math. Theor.* **40**(28):8127-8136 (2007).
- [3] V.I. Bogachev and A.V. Kolesnikov, “The Monge-Kantorovich problem: achievements, connections, and perspectives”, *Russian Math. Surveys* **67**(5):785-890 (2012).
- [4] M. Christandl and A. Winter, “‘Squashed entanglement’ – An additive entanglement measure”, *J. Math. Phys.* **45**(3):829-840 (2003).
- [5] T.M. Cover and J.A. Thomas, *Elements of Information Theory*, 2nd ed., John Wiley & Sons, 2006.
- [6] I. Devetak and P.W. Shor, “The capacity of a quantum channel for simultaneous transmission of classical and quantum information”, *Commun. Math. Phys.* **256**:287-303 (2005).
- [7] M.J. Donald and M. Horodecki, “Continuity of relative entropy of entanglement”, *Phys. Lett. A* **264**:257-260 (1999).
- [8] J. Eisert, C. Simon, and M.B. Plenio, “On the quantification of entanglement in infinite-dimensional quantum systems”, *J. Phys. A: Math. Gen.* **35**(17):3911-3923 (2002).
- [9] M. Fannes, “A continuity property of the entropy density for spin lattice systems”, *Commun. Math. Phys.* **31**:291-294 (1973).
- [10] A.S. Holevo, “On Quantum Communication Channels with Constrained Inputs”, arXiv:quant-ph/9705054 (1997).
- [11] A.S. Holevo, “Entanglement-Assisted Capacities of Constrained Quantum Channels”, *Theory Probab. Appl.* **48**(2):243-255 (2006).
- [12] A.S. Holevo and M.E. Shirokov, “Continuous Ensembles and the Capacity of Infinite-Dimensional Quantum Channels”, *Theory Probab. Appl.* **50**(1):86-98 (2006).
- [13] R. Horn and Ch. Johnson, *Matrix analysis*, Cambridge University Press, 2nd edition, 2013.
- [14] G. Kimura, T. Miyadera, and H. Imai, “Optimal state discrimination in general probabilistic theories”, *Phys. Rev. A* **79**:062306 (2009).
- [15] P. Kok and B.W. Lovett, *Introduction to Optical Quantum Information Processing*, Cambridge University Press, 2010.
- [16] D. Leung and G. Smith, “Continuity of quantum channel capacities”, *Commun. Math. Phys.* **292**(1):201-215 (2009).
- [17] E.H. Lieb and M.-B. Ruskai, “Proof of the strong subadditivity of quantum-mechanical entropy”, *J. Math. Phys.* **14**(12):1938-1941 (1973).
- [18] M. Mosonyi and F. Hiai, “On the Quantum Rényi Relative Entropies and Related Capacity Formulas”, *IEEE Trans. Inf. Theory* **57**(4):2474-2487 (2011).
- [19] M.A. Nielsen, “Continuity bounds for entanglement”, *Phys. Rev. A* **61**:064301 (2000).
- [20] Y. Ouyang, “Channel covariance, twirling, contraction, and some upper bounds on the quantum capacity”, *Quantum Inf. Comput.* **14**(11&12):917-936 (2014); arXiv[quant-ph]:1106.2337v6.
- [21] D. Petz, *Quantum Information Theory and Quantum Statistics*, Springer Verlag, Berlin Heidelberg, 2008.
- [22] M.E. Shirokov and A.S. Holevo, “On Approximation of Infinite-Dimensional Quantum Channels”, *Probl. Inf. Transm.* **44**(2):73-90 (2008).
- [23] M.E. Shirokov, “Continuity of the von Neumann entropy”, *Commun. Math. Phys.* **296**(3):625-654 (2010).
- [24] M.E. Shirokov, “Measures of quantum correlations in infinite-dimensional systems”, arXiv[quant-ph]:1506.06377 (2015).
- [25] M.E. Shirokov, “Squashed entanglement in infinite dimensions”, arXiv[quant-ph]:1507.08964 (2015).
- [26] G. Smith, J.A. Smolin, and A. Winter, “The Quantum Capacity With Symmetric Side Channels”, *IEEE Trans. Inf. Theory* **54**(9):4208-4217 (2008).
- [27] G. Smith, “Private classical capacity with a symmetric side channel and its application to quantum cryptography”, *Phys. Rev. A* **78**:022306 (2008).
- [28] D. Sutter, V.B. Scholz, and R. Renner, “Approximate Degradable Quantum Channels”, arXiv[quant-ph]:1412.0980 (2014).
- [29] D. Sutter, V.B. Scholz, R. Renner, and A. Winter, in preparation.
- [30] B. Synak-Radtke and M. Horodecki, “On asymptotic continuity of functions of quantum states”, arXiv:quant-ph/0507126 (2005).
- [31] C. Weedbrook, S. Pirandola, R. García-Patrón, N.J. Cerf, T.C. Ralph, J.H. Shapiro, and S. Lloyd, “Gaussian quantum information”, *Rev. Mod. Phys.* **84**(2):621-669 (2012).
- [32] A. Wehrl, “General properties of entropy”, *Rev. Mod. Phys.* **50**(2):221-260 (1978).
- [33] S. Weis and A. Knauf, “Entropy distance: new quantum phenomena”, *J. Math. Phys.* **53**:102206 (2012).
- [34] M.M. Wilde, *Quantum Information Theory*, Cambridge University Press, 2013.
- [35] D. Yang, M. Horodecki, and Z.D. Wang, “An Additive and Operational Entanglement Measure: Conditional Entanglement of Mutual Information”, *Phys. Rev. Lett.* **101**:140501 (2008); arXiv:quant-ph/0701149.