

Perfect Secrecy

under Deep Random assumption

Thibault de Valroger ^(*)

Abstract

We present a new idea to design perfectly secure information exchange protocol, based on a new form of randomness, called « Deep Randomness », generated in such a way that probability distribution of the output signal is made unknowledgeable for an observer. By limiting, thanks to Deep Randomness, the capacity of the opponent observer to perform Bayesian inference over public information to estimate private information, we can design protocols, beyond Shannon limit, enabling two legitimate partners, sharing originally no common private information, to exchange secret information with accuracy as close as desired from perfection, and knowledge as close as desired from zero by any unlimitedly powered opponent. We introduce the Deep Randomness assumption, based on Prior Probability theory developed by Carnap, Cox and Jaynes. Deep Random assumption gives a rigorous rule to perform bayesian inference under uncertainty situation. We also present a workable example of protocol with proof of perfect secrecy under Deep Random assumption, and we present a method to generate Deep Randomness from classical computation resources.

Key words. Perfect secrecy, Deep Random, advantage distillation, secret key agreement, unconditional security, quantum resistant

Modern cryptography mostly relies on mathematical problems commonly trusted as very difficult to solve, such as large integer factorization or discrete logarithm, belonging to complexity theory. No certainty exist on the actual difficulty of those problems, not even the truth of the famous $P \neq NP$ conjecture, and, in addition, most of them are not resistant to quantum computing, which should make them useless in the next decades. Some other methods, rather based on information theory, have been developed since early 90's. Those methods relies on hypothesis about the opponent (such as « memory bounded » adversary [6]) or about the communication channel (such as « independent noisy channels » [5]) ; unfortunately, even if they have been proven Information Theoretically secure under given hypothesis, none of those hypothesis are easy to ensure in practice. At last, some other methods based on physical theories like quantum indetermination [3] or chaos generation have been described and experimented, but they are complex to implement, and, again, relies on solid but not proven and still partly understood theories.

Considering this theoretically unsatisfying situation, we propose to explore a new path, where Information Theoretical security against unlimited opponent can be reached, without relying on any

() See contact and information about the author at last page*

assumption about the opponent, that is supposed to have unlimited computation and storage power, nor about the communication channel, that is supposed to be perfectly public, accessible and equivalent for any playing party (legitimate partners and opponents). The considered opponent is passive, which means that it does not interfere actively in the communication by suppressing, adding or modifying information exchanged between the legitimate partners; it just has a full access to it. Active opponent could be considered by adding authentication schemes, but this is not the core objective of the present work.

I. Introduction

Definition of “Information Theoretically Secure” Security Property

The most common Security Property to study Information Theoretically secure protocols is Perfect Secrecy, introduced by Claude Shannon [1]. This Security Property is defined as the condition $P(M|E) = P(M)$, where M designates the private information or “Message” in the protocol, although E designates the public information or “Encrypted information” in the protocol. As a security property, Perfect Secrecy has two drawbacks: (i) it is absolute and thus doesn’t give rule to measure the secrecy as soon as $H(M) > H(M|E)$ (where $H(\cdot)$ designates Shannon’s entropy), (ii) it does not explicitly differentiate the point of view of the opponent and the point of view of the legitimate receiver, and thus gives no rule to compare them in terms of capacity to access the private information as soon as the above inequality is strict.

In a very general view, in order to define the security property that we aim at satisfying, we consider here communication protocols that can be modeled as proposed by Csiszàr and Körner [13]. The legitimate partners are A and B , and ξ is the passive opponent. The purpose of the protocol is for A and B to reach common knowledge, with least possible errors of an output digital information, and to ensure that the passive opponent can learn the lesser possible information about that output digital information. By convention, the reference output digital information is the one reached by A called the Secret Common Output information and denoted X . For that reason B will be called the legitimate receiver, even if the protocol is not designed as a one way transmission. X is a discrete random variable taking values over a finite alphabet \mathcal{X} . A and B communicate over a Discrete Memoryless Channel (DMC) called main channel and B generates its best version, denoted Y , of the Secret Common Output information. Y is a discrete random variable taking values over a finite alphabet \mathcal{Y} (in most cases $\mathcal{Y} = \mathcal{X}$). The opponent ξ receives exact or altered copy (depending on the security model) of all information exchanged on the main channel over another DMC called side channel or wire-tap channel, and then generates its best version of the Secret Common Output information denoted Z . Z is a discrete random variable taking values over a finite alphabet \mathcal{Z} (in most cases $\mathcal{Z} = \mathcal{X}$). The alphabets $\mathcal{X}, \mathcal{Y}, \mathcal{Z}$ take values in a measurable sample space F . ξ ’s objective is to obtain the best possible knowledge of X . The protocol is a predefined and public communication routine involving public parameters U ; it is then denoted $\mathcal{P}(U)$. Each stakeholder in the protocol (A, B, ξ) is an entity capable to Generate random bit strings, Publish bit strings on its DMC (ξ is however supposed passive), Read bit strings published on its DMC, Store bit strings, Make computation on bit strings.

Then, we define that the protocol is “Information Theoretically Secure” if, $\forall \epsilon > 0$, there exists a value of the parameter U such that:

$$H(X|Z) - H(X|Y) \geq (1 - \epsilon)H(X) \quad (SP)$$

It is easy to see that this Security Property tends to Perfect Secrecy when $\epsilon \rightarrow 0$. Indeed, for the above inequality to hold when $\epsilon = 0$, it is necessary that (i) $H(X|Y) = 0$ which means that the protocol is errorless, and (ii) that $H(X|Z) = H(X)$ which means that $P(X|Z) = P(X)$. Csiszàr and Körner have shown that the secrecy capacity C_s of the protocol is lower-bounded by:

$$C_s \geq \max_{P_X} (H(X|Z) - H(X|Y))$$

which explains the rationale for the choice of (SP) above to characterize the security property. The literature also often reference this property under the name ‘Unconditional Security’, even if that naming may appear a bit improper due to the fact that some constraint \mathcal{C} is always applied on the stakeholders or on the communication channels, in order to overcome Shannon’s impossibility theorem (see next section).

The constraint \mathcal{C} applied to the side channel compared to the main channel may vary with the model. For instance, in *Partially Independent Channels* model introduced by Maurer in [5], the constraint is that the opponent receives the public information over a noisy side BSC (Binary Symmetric Channel) whose noise is at least partially statistically independent of the noise applying on the main BSC. It is then shown that a protocol can be designed to reach (SP) under that constraint, even if the opponent has unlimited computation and storage power. Another example is BB84, the first Quantum Key Distribution protocol proposed in [3], in which the constraint is that for each digit transmitted thanks to the polarization of a photon, the opponent has to choose a filter among 2 possible ones to measure the polarization wearing the digital information, and if the opponent makes the wrong choice, it obtains a random result rather than the actual result. This constraint legitimately comes from two properties of quantum systems: (i) the Heisenberg uncertainty principle and (ii) the no-cloning theorem. It is then shown as well that BB84 can reach (SP) under that constraint, even if the opponent has unlimited computation and storage power.

In this article, the constraint of the Security Model is called “Deep Random Assumption”. It is detailed, together with the Security Model, in the next sections. It relies on the fact that the probability distribution used by each legitimate partner to generate private information is made ‘unknown’ for any other party (including the other partner and of course the opponent). We rigorously explain what we mean by ‘unknown’ probability distribution in the section presenting the Deep Random Assumption hereafter. We present in Section III a protocol that satisfies (SC) under Deep Random Assumption, even if the opponent has unlimited computation and storage power. Section III also presents the comparison and the similarities between our protocol and both BB84 and Maurer’s protocol.

Shannon and the need for Bayesian inference

Shannon, in [1], established his famous impossibility result. Shannon defines perfect secrecy of a secrecy system, as its ability to equal the probability of the clear message $P(M)$ and the conditional probability $P(M|E)$ of the clear message knowing the encrypted message. In the case where the encryption system is using a shared secret key K with a public transformation procedure to transform the clear message into the encrypted message, Shannon establishes that perfect secrecy can only be obtained if $H(K) \geq H(M)$.

It is a common belief in the cryptologic community that, in cases where the legitimate partners initially share no secret information (which we can write $H(K) = 0$), the result of Shannon thus means that it is impossible for them to exchange a perfectly secret (or almost perfectly secret) bit of information. The support for that belief is that, in the absence of key entropy, the conditional

expectation $E[M|E]$, that is the best possible estimation of M knowing E , is completely and equally known by all the parties (legitimate receiver and opponent), as :

$$E[M|E] = \sum_m m \frac{P(E|m)P(m)}{\sum_{m'} P(E|m')P(m')}$$

and thus, that the legitimate receiver cannot gain any advantage over the opponent when he tries to estimate the secret clear message from the public encrypted message. This reasoning however supposes that all the parties have a full knowledge of the distribution $P(M)$, enabling them to perform the above Bayesian inference to estimate M from E .

Shannon himself warned the reader of [1] to that regard, but considered that this assumption is fairly reasonable (let's remember that computers were almost not yet existing when he wrote his article):

*« There are a number of difficult epistemological questions connected with the theory of secrecy, or in fact with any theory which involves questions of probability (particularly a priori probabilities, Bayes' theorem, etc.) when applied to a physical situation. Treated abstractly, probability theory can be put on a rigorous logical basis with the modern measure theory approach. As applied to a physical situation, however, especially when "subjective" probabilities and unrepeatable experiments are concerned, there are many questions of logical validity. For example, in the approach to secrecy made here, **a priori probabilities of various keys and messages are assumed known by the enemy cryptographer.** »*

The model of security that we develop in this article, by enabling the legitimate partners to use a specific form of randomness where the a priori probabilities of the messages cannot be efficiently known by the opponent, puts this opponent in a situation where the above reasoning based on Bayesian inference no longer stands.

It is remarkable that in his considerations about Perfect Secrecy, and in his proof of the impossibility result, Shannon only considers errorless protocols, meaning protocols such that $H(E|M, K) = H(M|E, K) = 0$. As soon as the protocol is not errorless, and assuming that the protocol has no prior key arrangement ($H(K) = 0$), then the impossibility result is intuitively interpreted as the fact that the legitimate partner cannot get a strictly "lesser" error rate than the opponent; "lesser" meaning as measured in the measurable sample space. We give in [13] a relation between secrecy measurement based on a distance in the sample space and secrecy measurement based on conditional entropy, as proposed by Csiszàr and Körner, when the sample space is a subset of the fairly general Euclidean space \mathbb{R}^n . For non-trivial message distribution, the function $S(E)$ that sets the most probable message given the public information is determined as the unique solution to

$$\text{Inf}_S \|S(E) - M\|$$

That unique solution is of course $E[M|E]$, the conditional expectation of the secret information (generically called by Shannon "message") knowing the public information (generically called by Shannon "cryptogram"). The determination of $E[M|E]$ requires knowledge of the distribution $P(M)$. As soon as the opponent doesn't know the distribution $P(M)$, he cannot determine the unique optimal solution $S(E)$. This does not mean of course that the error rate of the legitimate receiver will become automatically lesser than the error rate of the opponent, but it explains why the proof of Shannon cannot any longer be trivially applied in such cases.

Prior probabilities theory

Before presenting the Deep Random assumption, it is needed to introduce Prior probability theory.

The art of prior probabilities consists in assigning probabilities to a random event in a context of partial or complete uncertainty regarding the probability distribution governing that random event. The first author who has rigorously considered this question is Laplace [10], proposing the famous *principle of insufficient reason* by which, if an observer does not know the prior probability of occurrence of 2 events, he should consider them as equally likely. In other words, if a random variable X can take several values v_1, \dots, v_n , and if no information regarding the prior probabilities $P(X = v_i)$ is available for the observer, he should assign them $P(X = v_i) = 1/n$ in any attempt to produce inference from an experiment of X .

Several authors observed that this principle can lead to conclusion contradicting the common sense in certain cases where some knowledge is available to the observer but not reflected in the assignment principle.

If we denote $\mathfrak{I}_<$ the set of all prior information available to observer regarding the probability distribution of a certain random variable ('prior' meaning before having observed any experiment of that variable), and $\mathfrak{I}_>$ any public information available regarding an experiment of X , it is then possible to define the set of possible distributions that are compatible with the information $\mathfrak{I} \triangleq \mathfrak{I}_< \cup \mathfrak{I}_>$ regarding an experiment of X ; we denote this set of possible distributions as:

$$\Omega_{\mathfrak{I}}$$

The goal of Prior probability theory is to provide tools enabling to make rigorous inference reasoning in a context of partial knowledge of probability distributions. A key idea for that purpose is to consider groups of transformation, applicable to the sample space of a random variable X , that do not change the global perception of the observer. In other words, for any transformation τ of such group, the observer has no information enabling him to privilege $\varphi_{\mathfrak{I}}(v) \triangleq P(X = v|\mathfrak{I})$ rather than $\varphi_{\mathfrak{I}} \circ \tau(v) = P(X = \tau(v)|\mathfrak{I})$ as the actual distribution. This idea has been developed by Jaynes [7], in order to avoid the incoherence brought in some cases by the imprudent application of Laplace principle.

We will consider only finite groups of transformation, because one manipulates only discrete and bounded objects in digital communications. We define the acceptable groups G as the ones fulfilling the 2 conditions below:

(C1) Stability - For any distribution $\Phi \in \Omega_{\mathfrak{I}}$, and for any transformation $\tau \in G$, then $\Phi \circ \tau \in \Omega_{\mathfrak{I}}$

(C2) Convexity - Any distribution that is invariant by action of G does belong to $\Omega_{\mathfrak{I}}$

It can be noted that the set of distributions that are invariant by action of G is exactly:

$$R_{\mathfrak{I}}(G) \triangleq \left\{ \frac{1}{|G|} \sum_{\tau \in G} \Phi \circ \tau \mid \forall \Phi \in \Omega_{\mathfrak{I}} \right\}$$

The condition (C2) is justified by the fact that in the absence of information enabling the observer to privilege Φ from $\Phi \circ \tau$, he should choose equivalently one or the other distribution, but then of course the average distribution $\frac{1}{|G|} \sum_{\tau \in G} \Phi \circ \tau$ should still belong to the set $\Omega_{\mathfrak{I}}$ of possible distributions knowing \mathfrak{I} .

The set of acceptable groups as defined above is denoted:

$$\Gamma_{\mathfrak{S}}$$

Let's consider some examples.

Example 1: we consider a 6-sided dice. We are informed that the distribution governing the probability to draw a given side is altered, but we have no information of what that distribution actually is, and we have no available information regarding an experiment. We want nevertheless to assign an a priori probability distribution for the draw of dice. In this very simple case, it seems quite reasonable to assign an a priori probability of $1/6$ to each side. A more rigorous argument to justify this decision, based on the above, is the following: let's consider G the finite group of transformation that let the dice unchanged, this group G is well known, it is generated by the 3 axis 90° rotations, and has 24 elements. It is clear here that $G \in \Gamma_{\mathfrak{S}}$. It is also clear that, by considering a given distribution (p_1, \dots, p_6) , the a priori information available to the observer gives him no ground to privilege (p_1, \dots, p_6) rather than $(p_{g(1)}, \dots, p_{g(6)})$ for any $g \in G$, and therefore the distribution should be of the form:

$$\left\{ \left(\frac{1}{|G|} \sum_{g \in G} p_{g(i)} \right)_{i \in \{1, \dots, 6\}} \right\}_{\{(p_1, \dots, p_6)\}}$$

It is easy to compute that whatever is (p_1, \dots, p_6) ,

$$\frac{1}{|G|} \sum_{g \in G} p_{g(i)} = \frac{1}{6} \sum_{i=1}^6 p_i = \frac{1}{6}$$

and therefore in this trivial example, the Laplace principle applies nicely.

Example 2: let's now suppose that we have the result J of a draw of the dice. Then the symmetry disappears and the opponent may want to assign, as an extreme, a probability

$$p_j = \Gamma_j(j) \triangleq \begin{cases} 1 & \text{if } j = J \\ 0 & \text{otherwise} \end{cases}$$

One could argue that in a given probabilistic situation, there may exist several groups of transformation in $\Gamma_{\mathfrak{S}}$, and in that case, the choice of a given such group may appear arbitrary to assign the probability distribution. Although that oddness is not a problem for our purpose, we can solve it when 2 reasonable conditions are fulfilled: (i) we only consider finite sample space (note that objects in digital communication theory are bounded and discrete), and (ii) we assume that $\Omega_{\mathfrak{S}}$ is convex (which can be ensured by design of the Deep Random source). Under those 2 conditions, all groups of transformation applying on the sample space are sub-groups of the finite (large) group of permutations \mathfrak{S} applying on all the possible states, and if 2 groups of transformations G and G' applying on the sample space are in $\Gamma_{\mathfrak{S}}$, it is easy to see that $G \vee G'$ is still a finite sub-group of \mathfrak{S} , it contains G and G' and it is still in $\Gamma_{\mathfrak{S}}$. Consequently, for any observer having the same knowledge \mathfrak{S} of the distribution, there exists a unique maximal sub-group $T_{\mathfrak{S}} \in \Gamma_{\mathfrak{S}}$, and this one should be ideally applied to obtain a maximal restriction of the set of distribution (because it is easy to check that if $G' \subset G$, then $R_{\mathfrak{S}}(G) \subset$

$R_{\mathfrak{S}}(G')$). We can remark also that, if the uniform distribution is in $\Omega_{\mathfrak{S}}$, then $\bigcap_{G \in \Gamma_{\mathfrak{S}}} R_{\mathfrak{S}}(G) \neq \emptyset$ because $R_{\mathfrak{S}}(G)$ then always contains the uniform distribution.

However, it is not necessary for our purpose to be able to determine such unique maximal sub-group if it exists.

Another objection could be that the definition of $\Omega_{\mathfrak{S}}$ may appear too ‘black or white’ (a distribution belongs or does not belong to $\Omega_{\mathfrak{S}}$), although real situations may not be so contrasted. This objection is not applicable to our purpose because in the case of Deep Random Generation, that we will introduce in section IV, we can ensure by design that distributions do belong to a specific set $\Omega_{\mathfrak{S}}$. But otherwise, generally speaking, it is somehow preferable to draft the condition (C1) as:

(C1) For any distribution Φ , and for any transformation τ in such group, the observer has no information enabling him to privilege Φ rather than $\Phi \circ \tau$ as the actual distribution.

Deep Random assumption

We can now introduce and rigorously define the Deep Random assumption. The Deep Random assumption imposes a rule enabling to make rigorous inference reasoning about an observer ξ in a context where that observer ξ only has partial knowledge of a probability distribution.

Let X be a private random variable with values in a measurable set F . Keeping the notations of the previous section, we still denote $\mathfrak{S}_{<}$ the set of all prior information available to observer regarding the probability distribution of X (‘prior’ meaning before having observed any experiment of that variable), and $\mathfrak{S}_{>}$ any public information available regarding an experiment of X . Typically, X represents a secret information that ξ has to determine with the highest possible accuracy, and $\mathfrak{S}_{>}$ represents the public information available to ξ (like for instance the information exchanged during a communication protocol as introduced in the following sections).

For any group G of transformations applying on the sample space F , we denote by $\Omega_{\mathfrak{S}}(G)$ the set of all possible conditional expectations when the distribution of X courses $R_{\mathfrak{S}}(G)$. In other words:

$$\Omega_{\mathfrak{S}}(G) \triangleq \{Z(\mathfrak{S}) \triangleq E[X|\mathfrak{S}] | \forall \varphi_{\mathfrak{S}} \in R_{\mathfrak{S}}(G)\}$$

Or also:

$$\Omega_{\mathfrak{S}}(G) = \left\{ Z(\mathfrak{S}) = \int_F v \varphi_{\mathfrak{S}}(v) dv | \forall \varphi_{\mathfrak{S}} \in R_{\mathfrak{S}}(G) \right\}$$

The **Deep Random assumption** prescribes that, if $G \in \Gamma_{\mathfrak{S}}$, the strategy Z_{ξ} of the opponent observer ξ , in order to estimate X from the public information \mathfrak{S} , should be chosen by the opponent observer ξ within the restricted set of strategies:

$$Z_{\xi} \in \Omega_{\mathfrak{S}}(G) \tag{A}$$

The Deep Random assumption can thus be seen as a way to restrict the possibilities of ξ to choose his strategy in order estimate the private information X from his knowledge of the public information \mathfrak{S} . It is a fully reasonable assumption as exposed in the previous section presenting Prior probability theory, because the assigned prior distribution should remain stable by action of a transformation that let the distribution uncertainty unchanged.

(A) suggests of course that Z_ξ should eventually be picked in $\bigcap_{G \in \Gamma_\xi} \Omega_\xi(G)$ (that equals to $\Omega_\xi(T_\xi)$ when T_ξ exists), but it is enough for our purpose to find at least one group of transformation with which one can apply efficiently the Deep Random assumption to the a protocol in order to measure an advantage distilled by the legitimate partners compared to the opponent.

The security model

In our security model, we consider secrecy protocols being specific case of the Csiszàr and Körner model, where all the information exchanged by the legitimate partners A and B over the main channel are fully available to the passive opponent ξ . ξ has unlimited computation and storage power. A and B share initially no private information.

Beyond being capable to Generate random bit strings, Publish bit strings on the main channel, Read published bit strings from the main channel, Store bit strings, and Make computation on bit strings, the legitimate partners A and B are also capable to generate bit strings with Deep Randomness, by using their Deep Random Generator (DRG). A DRG is a random generator that produces an output with a probability distribution that is made unknown, under the meaning given in previous sections, to an external observer. We present in Section IV how a DRG can be implemented with classical computing resources.

ξ is capable to Read all published bit strings from the main channel, Store bit strings, and Make computation on bit strings, with unlimited computation and storage power. But when ξ desires to infer a private information generated by A 's DRG (or by B 's DRG) from public information, he can only do it in respect of the Deep Random assumption (A) presented in previous section. This assumption creates a 'virtual' side channel for the opponent conditioning the optimal information Z he can obtain to estimate the Secret Common Output information X .


This assumption is fully reasonable, as established in the former sections, under the condition that the DRG of A and the DRG of B can actually produce distributions that are truly undistinguishable and unpredictable among a set $R_\xi(G)$.

The protocols that we consider obey the Kerckhoffs's principle by the fact that their specifications are entirely public.

Information Theoretically Secure Protocols (ITSP)

The main purpose of this work is to introduce how to design an « Information Theoretically Secure Protocol » (ITSP) under Deep Random assumption, and to introduce how to generate Deep Random from classical computing resources.

We consider protocols as per the Security Model presented in previous section,



in which A (resp. B) requests its Deep Random Generator (DRG) to obtain an experiment u (resp. v) of a private random variable with hidden probability distribution ; A (resp. B) publishes the set of information i (resp. j) on the public channel along the protocol. A computes the Secret Common Output information $X(u, i, j)$, with value in measurable space F . B computes its estimation $Y(v, i, j)$ of the Secret Common Output information, also with value in F . In this model, $\mathfrak{S}_<$ is the public information available regarding the DRG of A or B (that are supposed to have the same design), and $\mathfrak{S}_> = \{i, j\}$ is the set of information published by the partners along the execution of the protocol.

The DRG of A is run by A completely privately and independently of B , and reversely the DRG of B is run by B completely privately and independently of A . The 2 DRG are thus not in any way secretly correlated, as one of the assumptions of the security model is that A and B share initially no private information.

The eavesdropping opponent ξ who has a full access to the public information, computes its own estimation $Z(\mathfrak{S})$ (that we will also shortly denote $Z(i, j)$). Z is called ‘strategy’ of the opponent.

As introduced in the ‘Deep Random assumption’ section, $\mathfrak{S}_<$ designates the public information available about a DRG. We assume here that $\mathfrak{S}_<$ is the same for both DRG of A and B , meaning that they have the same conception.

From the Deep Random assumption, for any group G in $\Gamma_{\mathfrak{S}}$, the set of optimal strategies for the opponent can be restricted to:

$$\Omega_{\mathfrak{S}}(G, \mathcal{P}) = \{Z(\mathfrak{S}) = E[X|\mathfrak{S}] | \forall (\Phi(u), \Phi'(v)) \in R_{\mathfrak{S}}(G) \times R_{\mathfrak{S}}(G)\}$$

Then, the protocol \mathcal{P} is a ITSP if it verifies the following property (P):

$$\exists G \in \Gamma_{\mathfrak{S}}, \alpha > 0 | \forall Z \in \Omega_{\mathfrak{S}}(G, \mathcal{P}): H(X|Z) - H(X|Y) \geq \alpha \quad (P)$$

It has been shown ([4], [5]) that when a protocol satisfies $H(X|Z) - H(X|Y) \geq \alpha$, it can be complemented with Reconciliation and Privacy Amplification techniques to satisfy the targeted security property (SP) introduced in the first section with ϵ as small as desired.

We will also impose a second condition for the definition of an ITSP \mathcal{P} ; that condition ensures that one can implement a DRG suitable for \mathcal{P} thanks to a recursive and continuous generation algorithm, as presented in section IV and introduced below. The second condition is the following:

$$\exists \alpha > 0 | \forall Z(\mathfrak{Z}), \exists \Phi(u), \Phi'(v) | H(X|Z) - H(X|Y) \geq \alpha \quad (P')$$

The condition (P') means that for any strategy $Z(\mathfrak{Z})$, even outside of $\Omega_{\mathfrak{Z}}(G, \mathcal{P})$, there exists some distributions $\Phi(u), \Phi'(v)$ of the private information of A and B , potentially outside of any $R_{\mathfrak{Z}}(G)$, such that the condition $H(X|Z) - H(X|Y) \geq \alpha$ is satisfied. The interest of that condition is that (i) it is not referring to the Deep Random Assumption, and therefore (ii) it enables to build a Deep Random Generator based on a recursive and continuous generation algorithm proceeding as follows:

Deep Random Generation

The basic idea is to use infinite incrementing counters as primary source of entropy, and then to run across those counters one or several Cantor' style diagonal recursive constructing process. Such a DRG, based on continuous and recursive constructing process, can only be designed in association with a given ITSP. At each step $m + 1$, the generator emulates the ITSP internally and picks (through classical randomness) a new couple of distributions $\Phi(u), \Phi'(v)$ that defeats the best possible strategy knowing all the past distributions for $t \leq m$, which is always possible for an ITSP as given by condition (P') . The source of secret entropy is the current values of the infinite counters (several can run in parallel) of the continuous recursive process, together with the classical random that is used at each step to pick a defeating distribution. Such entropy must of course be at least equal to $\log|\mathcal{X}|$ in order to approach perfect secrecy as close as desired.

A detailed presentation of a DRG using a recursive method as introduced above can be found in Section IV.

An example of ITSP is given in Section III, as well as a comparison between that example and well known protocols like Maurer's broadcast satellite scenario and Bennett and Brassard Quantum Key Agreement protocol BB84.

In Section II, we present the formalism of Bernoulli random variables that we use to design our protocol; notations are defined and first considerations are discussed; Section II can be skipped for a conceptual, non-technical, reading of this article.

In Section III, we present our protocol, an example of ITSP, and we present the main theorem that establishes Information Theoretical Security of the protocol ; a high level argumentation of the proof is given in Section III, while a detailed and fully rigorous proof can be found in Annex I, with a set of preparatory technical results;

In Section IV, we present some methods using classical computing resources to design a Deep Random Generator;

In Section V, we draw conclusions about the present work and open on further questions to progress towards applications to modern cryptography.

In Annex, we give the details of the proofs of the technical results introduced in Section III.

II. Notations and first considerations

The protocol that we will present in Section III uses Bernoulli random variables, which present the advantage to enable Reconciliation thanks to the law of large numbers.

Let's define some notations. Considering $x = (x_1, \dots, x_n)$ and $y = (y_1, \dots, y_n)$ some parameter vectors in $[0,1]^n$ and $i = (i_1, \dots, i_n)$ and $j = (j_1, \dots, j_n)$ some experiment vectors in $\{0,1\}^n$, $l, r \in \mathbb{N}_n^*$ two integers, and $\theta \in [0,1]$, we define :

$x \cdot y$ (resp. $i \cdot j$) the scalar product of x and y (resp. i and j)

$$|x| \triangleq \sum_{s=1}^n x_s ; |i| \triangleq \sum_{s=1}^n i_s$$

$$i \subset j \Leftrightarrow \{s | i_s = 1\} \subset \{s | j_s = 1\}$$

$$\bar{i} \triangleq (u_1, \dots, u_n) \mid u_s = 1 \Leftrightarrow i_s = 0$$

$$i \setminus j \triangleq (u_1, \dots, u_n) \mid u_s = 1 \Leftrightarrow (i_s = 1) \& (j_s = 0)$$

$$i \cap j \triangleq (u_1, \dots, u_n) \mid u_s = 1 \Leftrightarrow (i_s = 1) \& (j_s = 1)$$

$$i \cup j \triangleq (u_1, \dots, u_n) \mid u_s = 1 \Leftrightarrow (i_s = 1) \text{ or } (j_s = 1)$$

$$\chi_i(x) \triangleq \prod_{s=1}^n (i_s x_s + (1 - i_s)(1 - x_s))$$

that represents probability of obtaining i in a draw of a Bernoulli random variable with parameter vector x .

$$\Pi_i(x) \triangleq \prod_{s=1}^n (i_s x_s + (1 - i_s))$$

$$\psi_{i,r}(x) \triangleq \sum_{j | j \cdot i = r} \chi_j(x) ; \psi_l(x) = \psi_{(1, \dots, 1), l}(x) \triangleq \sum_{j | |j| = l} \chi_j(x)$$

$$\beta_{l,r}(\theta) \triangleq \binom{r}{l} \theta^l (1 - \theta)^{r-l}$$

that represents the Bernoulli coefficient of parameter θ .

We will also manipulate permutation operators over vectors. For $\sigma \in \mathfrak{S}_n$, we write $\text{supp}(\sigma) = \ker(\sigma - id_{\mathfrak{S}_n}) = \{i, \sigma(i) \neq i\}$ and $|\sigma| = \text{card}(\text{supp}(\sigma))$. The permutation of a vector is the following linear application:

$$\forall \sigma \in \mathfrak{S}_n, \sigma(x) \triangleq (x_{\sigma(1)}, \dots, x_{\sigma(n)}) \text{ where } \mathfrak{S}_n \text{ represents the symmetric group}$$

The calculations about Bernoulli random variable take place in the vector space of multinomials with n variables and local degree 1, noted $\mathcal{M}_1(n)$, whose $\{\Pi_i(x)\}_{i \in \{0,1\}^n}$ and $\{\chi_i(x)\}_{i \in \{0,1\}^n}$ are basis. It is easy to justify that $\{\chi_i(x)\}_{i \in \{0,1\}^n}$ is a basis by noting the inversion formula:

$$\sum_{i \in \{0,1\}^n} \chi_i(x) = 1 \Rightarrow \Pi_j(x) = \sum_{i \supset j} \chi_i(x)$$

To ease the manipulation of upper bounds, we will use the notation $\triangleleft (M)$ for any quantity with value in \mathbb{R} and whose absolute value is bounded by M .

If A and B are two partners, owning private parameter vectors x/k and y/k respectively (where k is a real number parameter > 1), and generating vectors i and j respectively through Bernoulli random variables from x/k and y/k respectively, we consider the respective random variables V_A and V_B such that :

$$V_A = \frac{x \cdot j}{n} ; V_B = \frac{i \cdot y}{n}$$

By direct calculation:

$$\begin{aligned} E[V_A|x, y] &= E[V_B|x, y] = \frac{x \cdot y}{nk} \\ \text{Var}(V_A|x, y)^2 &= E[(V_A - E[V_A])^2|x, y] = \frac{1}{nk} \left(\frac{1}{n} \sum_{r=1}^n x_r^2 y_r \left(1 - \frac{y_r}{k}\right) \right) \\ \text{Var}(V_B|x, y)^2 &= E[(V_B - E[V_B])^2|x, y] = \frac{1}{nk} \left(\frac{1}{n} \sum_{r=1}^n y_r^2 x_r \left(1 - \frac{x_r}{k}\right) \right) \\ E[(V_B - V_A)^2|x, y] &= \frac{1}{n^2 k} \sum_{r=1}^n x_r y_r \left(x_r + y_r - \frac{2x_r y_r}{k} \right) \leq \frac{2}{nk} \end{aligned} \quad (\text{II.0})$$

The opponent ξ has access only to (i, j) and elaborates a strategy $V_\xi(i, j)$ in order to estimate V_A as accurately as possible. If ξ wants to obtain the equality

$$E[V_\xi|x, y] = E[V_A|x, y] \quad \forall x, y$$

one can easily determine that the only possible strategy satisfying the above is:

$$V_{\xi,0}(i, j) = \frac{ki \cdot j}{n}$$

On the other hand, we observe that the distance from $V_{\xi,0}$ to V_A is greater than $E[(V_B - V_A)^2]$ when $k > 3$:

$$E[(V_{\xi,0} - V_A)^2|x, y] = \frac{1}{n} \left(\frac{1}{n} \sum_{r=1}^n x_r y_r \left(1 - \frac{x_r}{k}\right) \right) \geq \frac{k-1}{2} E[(V_B - V_A)^2|x, y]$$

which means that if ξ wants to evaluate V_A from the public information (i, j) with exactly the same first moment $E[V_\xi|x, y] = E[V_A|x, y] \forall x, y$, then it is necessarily less accurate than the legitimate partners.

Is it so simple ?

Unfortunately it is not, because the opponent has no obligation to estimate V_A with a strategy having exactly the same first moment than V_A . As soon as the opponent knows the probability distributions used respectively by A and B to independently generate respectively x and y , he can generate a strategy $V_\xi^*(i, j)$ from the public information (i, j) , by using Bayesian inference. That strategy this time verifies:

$$E \left[(V_\xi^*(i, j) - V_A)^2 \right] \leq E[(V_B - V_A)^2]$$

For a proof, see Proposition 11 in Annex I.

$V_\xi^*(i, j)$ is of course the conditional expectation $E[V_A|i, j]$.

If Φ and Φ' are the probability distribution of x and y respectively, the optimal strategy $V_\xi^*(i, j)$ is expressed by:

$$V_\xi^*(i, j) = E[V_A|i, j] = \frac{\int_{x \in [0,1]^n} \frac{x \cdot j}{nk} \chi_i \left(\frac{x}{k} \right) \Phi(x) dx}{\int_{x \in [0,1]^n} \chi_i \left(\frac{x}{k} \right) \Phi(x) dx}$$

that clearly depends on the knowledge of Φ .

DRG's Independence Phenomenon

Let's give a useful example of a strategy that is in average more efficient than $V_{\xi,0}$ (the strategy equalizing the first moments as seen above), without being dependent of Φ , when the parameter vectors are chosen independently by the legitimate partners. Let's assume that Φ and Φ' are picked independently by A and B . We can then suppose that Φ and Φ' are symmetric (Φ is symmetric meaning that $\Phi \circ \sigma = \Phi \forall \sigma \in \mathfrak{S}_n$) because the choice of Φ is not more probable than the choice of $\Phi \circ \sigma$ for any permutation $\sigma \in \mathfrak{S}_n$. In that case, the strategy:

$$V_{\xi,1}(i, j) \triangleq \frac{k|i||j|}{n^2}$$

is more efficient than $V_{\xi,0}$ and is of the same accuracy than V_B to estimate V_A ; indeed by direct computation we get:

$$E \left[(V_{\xi,1} - V_A)^2 \right] \leq \frac{2}{nk} \tag{II.1}$$

This efficient strategy takes advantage of the symmetry resulting from the independence between the choices of the legitimate partners for their respective distributions. We will see in section III that the Independence Phenomenon is an important problem to overcome when designing an ITSP.

III. Presentation of the protocol and proof of perfect secrecy

In this Section we present an example of ITSP. Besides being hidden to the opponent, we will see that the probability distribution used by each legitimate partner also need to have specific properties in order to prevent the opponent to evaluate V_A with the same accuracy than the legitimate partners by using the Independence Phenomenon introduced in Section II.

When Bernoulli random variable are considered through quadratic evaluation, the probability distribution Φ of a legitimate partner can be considered through its quadratic matrix:

$$M_\Phi(u, v) = \int_{[0,1]^n} x_u x_v \Phi(x) dx$$

M_Φ is a symmetric matrix with elements in $[0,1]$. Let S_n stand for the set of the subsets I of $\{1, \dots, n\}$ with size $n/2$; we define $\|\cdot\|_c$ the c-norm by:

$$\forall I \in S_n, c_I(M_\Phi) = \frac{4}{n^2} \sum_{u,v \in I \times \bar{I}} M_\Phi(u, v) ; \|M_\Phi\|_c = \max_{I \in S_n} |c_I(M_\Phi)|$$

$\|\cdot\|_c$ is a norm (see Proposition 10) on the subspace of the matrix without their main diagonal, for any $n > 4$.

We associate to any distribution quadratic matrix M_Φ the matrix $\overline{M_\Phi}$ defined by :

$$M_\Phi \mapsto \overline{M_\Phi} = m_\Phi \begin{pmatrix} 0 & \dots & 1 \\ \vdots & \ddots & \vdots \\ 1 & \dots & 0 \end{pmatrix} \text{ where } m_\Phi = \frac{1}{n(n-1)} \sum_{u \neq v} M(u, v)$$

Then we define the following probability distribution set:

$$\zeta(\alpha) \triangleq \{\Phi \mid \|M_\Phi - \overline{M_\Phi}\|_c \geq \sqrt{\alpha}\}$$

That public information regarding the distributions constitutes in the protocol presented hereafter the prior information $\mathfrak{S}_<$ as defined in Section I.

We introduce the « tidied form » of a probability distribution $\Phi \in \zeta(\alpha)$. For any Φ , there exist a (not necessarily unique) permutation σ_Φ in \mathfrak{S}_n such that:

$$\sum_{u,v \in I_0 \times \bar{I}_0} M_{\Phi \circ \sigma_\Phi}(u, v) = \min_{\sigma \in \mathfrak{S}_n} \left(\sum_{u,v \in I_0 \times \bar{I}_0} M_{\Phi \circ \sigma}(u, v) \right)$$

Where

$$I_0 \triangleq \{1, \dots, n/2\}$$

$\Phi \circ \sigma_\Phi$ is then called a tidied form of Φ . σ_Φ is called a tidying permutation of Φ . Minimizing the above criteria is an efficient way to ensure that $\Phi \circ \sigma_\Phi$ is driven away from its symmetric projection

$$\bar{\Phi} \triangleq \frac{1}{n!} \sum_{\sigma \in \mathfrak{S}_n} \Phi \circ \sigma$$

Tidying permutations enable to ‘synchronize’ two distributions (see Proposition 9 in Annex), in the sense that the permuted distributions $\Phi \circ \sigma_\Phi$ and $\Phi' \circ \sigma_{\Phi'}$ avoid the DRG’s Independence Phenomenon introduced in Section II:

$$E \left[(V_{\xi,1} - V_A)^2 \right]_{\Phi \circ \sigma_\Phi, \Phi' \circ \sigma_{\Phi'}} \gg E[(V_B - V_A)^2]_{\Phi \circ \sigma_\Phi, \Phi' \circ \sigma_{\Phi'}}$$

Now we can explicit our example of ITSP protocol. It will be demonstrated in Theorem 1 and Theorem 2 that the presented ITSP fulfills the 2 conditions (P) and (P’) defining an ITSP. The protocol doesn’t have purpose to be optimal in terms of implementation. We will discuss quantitative aspects in Section V, and in further works.

Here are the steps of the proposed ITSP:

A and B are the legitimate partners of the Security Model presented in Section I. The steps of the protocol $\mathcal{P}(\alpha, n, k, K, L)$ are the followings:

Step 1 – Deep Random Generation: A and B pick independantly the respective probability distributions Φ and $\Phi' \in \zeta(\alpha)$, so that Φ (resp. Φ') is secret (under Deep Random assumption) for any observer other than A (resp. B) beholding all the published information. A draws the parameter vector $x \in \{0,1\}^n$ from Φ . B draws the parameter vector $y \in \{0,1\}^n$ from Φ' .

Step 2 – Degradation: $k > 1$ is a public Degradation parameter; A generates a Bernoulli experiment vectors $i \in \{0,1\}^n$ from the parameter vector $\frac{x}{k}$. A publishes i . B generates a Bernoulli experiment vectors $j \in \{0,1\}^n$ from the parameter vector $\frac{y}{k}$. B publishes j .

Step 3 – Dispersion: A and B also pick a second probability distribution respectively Ψ and $\Psi' \in \zeta(\alpha)$ such that it is also secret (under Deep Random assumption) for any observer other than A (resp. B). Ψ is selected also such that $\int_{|x| \in [k|i| - \sqrt{n}, k|i| + \sqrt{n}]} \Psi(x) dx \geq \frac{1}{2\sqrt{n}}$ in order to ensure that $|i|$ is not an unlikely value for $\approx \left\lfloor \frac{x}{k} \right\rfloor$ (same for Ψ' by replacing x by y and i by j). Ψ (resp. Ψ') is used to scramble the publication of the tidying permutation of A (resp. B). A (resp. B) calculates a permutation $\sigma_d[i]$ (resp. $\sigma'_d[j]$) representing the reverse of the most likely tidying permutation on Ψ (resp. Ψ') to produce i (resp. j). In other words, with i , $\sigma_d[i]$ realizes :

$$\max_{\sigma \in \mathfrak{S}_n} \int_x P(i|x) \Psi \circ \sigma_\Psi \circ \sigma^{-1}(x) dx$$

Then A (resp. B) draws a boolean $b \in \{0,1\}$ (resp. b') and publishes in a random order $(\mu_1, \mu_2) = t^b(\sigma_d[i], \sigma_\Phi)$, (resp. $(\mu'_1, \mu'_2) = t^{b'}(\sigma'_d[j], \sigma_{\Phi'})$) where t represents the transposition of elements in a couple.

(we remark that it is possible to publish only $(\mu_1(I_0), \mu_2(I_0))$ and $(\mu'_1(I_0), \mu'_2(I_0))$ rather than (μ_1, μ_2) and (μ'_1, μ'_2) , which leads to quantity of bits of $4n$ instead of $4n \log n$).

Step 4 – Synchronization: A (resp. B) chooses randomly σ_A (resp. σ_B) among (μ'_1, μ'_2) (resp. (μ_1, μ_2)).

Step 5 – Decorrelation: A computes $V_A = \frac{\sigma_{\Phi}^{-1}(x) \cdot \sigma_A^{-1}(j)}{n}$, B computes $V_B = \frac{\sigma_B^{-1}(i) \cdot \sigma_{\Phi'}^{-1}(y)}{n}$. V_A and V_B are then transformed respectively by A and B in binary output thanks to the sampling method:

$$\tilde{e}_A = \left\lfloor \frac{(V_A - \rho)\sqrt{nk}}{K} \right\rfloor \bmod 2, \tilde{e}_B = \left\lfloor \frac{(V_B - \rho)\sqrt{nk}}{K} \right\rfloor \bmod 2$$

(where ρ is a translation parameter randomly and publicly picked at each instance of the protocol, following the assumption of Proposition 15). The multiplicative factor K is chosen such that:

$$\frac{1}{\sqrt{nk}} \ll \frac{K}{\sqrt{nk}} \ll \frac{1}{\sqrt{n}}$$

(and following the assumption of Proposition 16). At this stage the protocol can then be seen as a broadcast model with 2 Binary Symmetric Channels (BSC), one between A and B and one between A and ξ who computes a certain \tilde{e}_{ξ} , called ξ 's strategy. It is shown in Theorem 1 that those 2 BSC are partially independent, which enable to create Advantage Distillation as shown in [5] [14].

Step 5' – Advantage Distillation: by applying error correcting techniques with code words of length L between A and B , as introduced in [5], we show in Theorem 1 that we can then create advantage for B compared to ξ in the binary flows resulting from the error correcting code.

Step 6: classical Information Reconciliation and Privacy Amplification (IRPA) techniques then lead to get accuracy as close as desired from perfection between estimations of legitimate partners, and knowledge as close as desired from zero by any unlimited opponent, as shown in [4].

The choice of the parameters (α, n, k, K, L) will be discussed in proof of main Theorem. They are set to make steps 5, 5' and 6 possible, which is the main achievement of this work.

The protocol can be heuristically analyzed as follows:

Regarding the legitimate partners, when B picks $\sigma_B = \sigma_{\Phi}$ and A picks $\sigma_A = \sigma_{\Phi'}$ (1/4 of cases), the choice of σ_A and σ_B remain independant from i, j , so that i and j remain draws of independent Bernoulli random variables, then allowing to apply Chernoff-style bounds with accuracy $O\left(\frac{1}{\sqrt{nk}}\right)$

$(E \left[\left(V_A |_{\sigma_A = \sigma_{\Phi'}} - V_B |_{\sigma_B = \sigma_{\Phi}} \right)^2 \right]^{1/2} = O\left(\frac{1}{\sqrt{nk}}\right)$ as shown in inequality (II.0) in Section II). When B picks $\sigma_B = \sigma_d[i]$ or A picks $\sigma_A = \sigma'_d[j]$, this is no longer true and V_B or V_A become erratic, which will lead to error detection by error correcting code at step 5'.

The below table summarizes that situation:

	B picks $\sigma_B = \sigma_\Phi$ among (μ_1, μ_2)	B picks $\sigma_B = \sigma_d[i]$ among (μ_1, μ_2)
A picks $\sigma_A = \sigma_{\Phi'}$ among (μ'_1, μ'_2)	<p>A and B respective estimations are close in $\sim 100\%$ of cases, and thus both obtain accurate estimation of the combined shared secret in $\sim 100\%$ of cases.</p> <p>ξ cannot make accurate estimation of \tilde{e}_A in at least $\sim 25\%$ of cases (if ξ tries to have a strategy depending on $(\mu_1, \mu_2, \mu'_1, \mu'_2)$, then $(\sigma_d[i], \sigma'_d[j])$ is indistinguishable from $(\sigma_\Phi, \sigma_{\Phi'})$ and is thus picked by ξ in 25% of cases.</p>	A and B respective estimations are not close which leads to error detection and finally discarding.
A picks $\sigma_A = \sigma'_d[j]$ among (μ'_1, μ'_2)	A and B respective estimations are not close which leads to error detection and finally discarding.	A and B respective estimations are not close which leads to error detection and finally discarding.

This is a heuristic reasoning, and we must rather consider most general strategies $\tilde{e}_\xi(i, j, \mu_1, \mu_2, \mu'_1, \mu'_2)$ and write the probability equations with the appropriate group transform, under Deep Random assumption. But this table explains why we create partial independence between the BSC and consequently then an advantage for the legitimate partners compared to the opponent, bearing in mind that (μ_1, μ_2) (resp. (μ'_1, μ'_2)) are absolutely undistinguishable knowing i (resp. j), due to the fact that the distributions Φ and Ψ (resp. Φ' and Ψ') are unknown and thus also absolutely undistinguishable by ξ .

Below we explain the role of each step:

The Degradation transformations $x \mapsto \frac{x}{k}$ and $y \mapsto \frac{y}{k}$ with $k > 1$ at step 2 are necessary so that $\sigma_d[i](I_0) \neq \sigma_\Phi(I_0)$ (resp. $\sigma'_d[i](I_0) \neq \sigma_{\Phi'}(I_0)$). If $k = 1$, then $i = x$ and $j = y$ and this would cause $\sigma_d[i](I_0) \approx \sigma_\Phi(I_0)$ (resp. $\sigma'_d[i](I_0) \approx \sigma_{\Phi'}(I_0)$).

The Deep Random Generation at step 1 prevents the use of Bayesian inference based on the knowledge of the probability distribution. In particular, if Φ, Ψ (resp. Φ', Ψ') were not managed by Deep Randomness, ξ would be able discriminate σ_Φ among (μ_1, μ_2) (resp. $\sigma_{\Phi'}$ among (μ'_1, μ'_2)) by Bayesian inference.

Dispersion step 3 mixes σ_Φ within (μ_1, μ_2) with another permutation $\sigma_d[i]$ (and $\sigma_{\Phi'}$ within (μ'_1, μ'_2) with another permutation $\sigma'_d[j]$). i is entirely determined by $|i|$ and a permutation, which explains the constraint and transformation applied on Ψ in step 3 to make σ_Φ and $\sigma_d[i]$ indistinguishable knowing i (same with $\sigma'_d[j]$, $\sigma_{\Phi'}$, and j). $\sigma_d[i]$ is (1) undistinguishable from σ_Φ knowing $\mathfrak{S}_> = \{i, j, (\mu_1, \mu_2), (\mu'_1, \mu'_2)\}$, and (2) manages to make the estimation of ξ unefficient as shown in

the proof of Theorem 1. That undistinguishability can be expressed by the symmetry Σ_1 associated to the group $\{\text{Id}, t\}$ (t being the transposition of pairs) applied to (μ_1, μ_2) and (μ'_1, μ'_2) . Then, Φ_A designating the distribution potentially chosen by A in the opponent's perspective, for ξ willing to evaluate V_A knowing the public information $\mathfrak{F}_>$, Σ_1 results into the fact that the probability distribution of $P(x, (\sigma_{\Phi_A}, \sigma_A) | \mathfrak{F})$ can be restricted under Deep Random Assumption to the set:

$$R_{\mathfrak{F}}(\Sigma_1) = \left\{ \frac{1}{2} \left(\frac{\chi_i \left(\frac{x}{k} \right) \Phi(u) dx}{\int_u \chi_i \left(\frac{u}{k} \right) \Phi(u) du} \cdot \delta_{\sigma_{\Phi}}(\sigma_{\Phi_A}) + \frac{\chi_i \left(\frac{x}{k} \right) \Psi(u) dx}{\int_u \chi_i \left(\frac{u}{k} \right) \Psi(u) du} \cdot \delta_{\sigma_{d[i]}}(\sigma_{\Phi_A}) \right) \cdot \left(\frac{1}{2} \left(\delta_{\sigma_{\Phi'}}(\sigma_A) + \delta_{\sigma'_{d[j]}}(\sigma_A) \right) \right) \mid \forall \Phi, \Psi \in \zeta(\alpha) \right\}$$

(where δ_v represents the Dirac distribution non-null at v). The combination of the 2 second terms in the sums above gives, with probability 1/4, the cases where ξ is misled in his attempt to estimate V_A with accuracy $O\left(\frac{1}{\sqrt{nk}}\right)$. See proof of Theorem 1 for the fully rigorous analysis.

The synchronization step 4 is designed to overcome the independence between the choices of the distributions of A and B . It needs that the distributions to have special properties ($\in \zeta(\alpha)$) in order to efficiently play their role. It is efficient in 1/4 of cases (when B picks $\sigma_B = \sigma_{\Phi}$ and A picks $\sigma_A = \sigma_{\Phi'}$, which we will call 'favorable cases'). If σ_{Φ} and $\sigma_{\Phi'}$ were chosen randomly by A and B instead of being the respective synchronization permutations of Φ and Φ' , then ξ could, even without knowing Φ and Φ' , use $V_{\xi,1}(i, j) \triangleq \frac{k|i||j|}{n^2}$ in order to estimate say V_A with accuracy $O\left(\frac{1}{\sqrt{nk}}\right)$, in which case $\tilde{e}_{\xi} = \left\lceil \frac{(V_{\xi,1} - \rho)\sqrt{nk}}{K} \right\rceil$ would be as close to \tilde{e}_A as \tilde{e}_B in favorable cases (see inequality (II. 1) in Section II).

The step 5 is called Decorrelation because at this step, thanks to the Deep Random Assumption, we have managed to create a protocol that can be equivalently modeled by a broadcast communication over 2 partially independent BSC, as shown in Theorem 1. Consequently it is possible to apply error correcting techniques to create Advantage Distillation as established in [5].

Theorem 1.

Under the Deep Random assumption, $\mathcal{P}(\alpha, n, k, K, L)$ satisfies condition (P)

The details of the proof are given in Annex. The strategy of the proof is to restrict the strategy of the opponent to $\Omega_{\mathfrak{F}}(\Sigma_1)$ (actually to an even more restricted set $\Omega_{\mathfrak{F}}(\Sigma_1 \times \Sigma_2 \times \Sigma_3)$) resulting from 2 other symmetries Σ_2 and Σ_3) thanks to the Deep Random assumption, and then to establish via Lemma 3 that in the protocol modeled as 3-segments independent BSC (as per Proposition 12), the opponent segment has an error probability $\epsilon_{\xi} > 0$. Then, thanks to error correcting method, an Advantage can be distilled. Finally, Information Reconciliation and Privacy Amplification techniques enable to satisfy (P) and thus even (SP) with ϵ as small as desired.

We also have with a very similar proof (also found in Annex):

Theorem 2.

$\mathcal{P}(\alpha, n, k, K, L)$ satisfies condition (P')

Parallel with BB84

The protocol \mathcal{P} presents some interesting similarities with the Quantum Key Distribution protocol BB84 presented by Bennet and Brassard in 1984 [3].

Let's first briefly remind the principles of BB84. A is secretly polarizing photons; for each photon, the polarization is chosen randomly by A between $\{0^\circ, 90^\circ\}$ (right polarization) or between $\{45^\circ, 135^\circ\}$ (inclined polarization); the choice between right polarization type or inclined polarization type is also made randomly by A . B can measure the polarization by choosing a right filter (that will give the good answer if the polarization is right), or an inclined filter (that will give the good answer if the polarization is inclined). If B chooses the wrong filter, the result of the measurement is random. The exact same situation applies for the opponent ξ .

The polarization in BB84 can be compared to the Dispersion step in \mathcal{P} where 2 indistinguishable permutations, associated to a same Bernoulli vector, are published.

The choice of filter by B (or E) in BB84, imposed by the Heisenberg uncertainty principle, can be compared to the Synchronization step in \mathcal{P} where B (or E) has no other choice, as per Deep Random Assumption, than choosing randomly one of the 2 permutations because they are indistinguishable knowing the Bernoulli vector.

In case of wrong choice of filter, the measurement step in BB84 will result into a random result for the measuring party. Similarly, the choice of the wrong permutation (σ_d instead of σ_Φ) in the Decorrelation step, can be shown (see Annex 1 for the detailed proof of Theorem 1) to result into erratic variable.

Then the protocols differ because BB84 can more easily exploit the partial independence created between choices of B and E thanks to the no-cloning theorem. Indeed, B can publicly disclose its filter choice so that A can publicly discard the wrong choices; ξ will not be able to exploit this information because it cannot redo another measurement on a 'copy' of the polarized photon. Contrariwise, \mathcal{P} only manipulates conventional digital information, that can be copied and accessed anytime by the unlimited opponent. Therefore, more sophisticated error correcting methods are needed to discard the cases where the legitimate partners have made unaligned choices of permutations.

Parallel with Maurer's broadcast satellite protocol

We have already mentioned the comparison between \mathcal{P} and the various protocols introduced by Maurer in [5], where the partners take advantage of partial independence noise between the main channel and the wire-tap channel. In particular, Maurer considers a scenario where A , B and ξ all receive a digit flow \widetilde{e}_A , \widetilde{e}_B and \widetilde{e}_ξ outputting from a broadcast satellite; each flow transmission can be modeled by a BSC with partially independent error variables. If the flow received by A is taken as a reference, then we denote ε_B and ε_ξ as the respective error variables of B 's channel and ξ 's channel, compared to the flow received by A : $\varepsilon_B = \widetilde{e}_A \oplus \widetilde{e}_B$, $\varepsilon_\xi = \widetilde{e}_A \oplus \widetilde{e}_\xi$.

A broadcast BSC channel $BSC(\varepsilon_B, \varepsilon_\xi)$ with 1 sender A and 2 receivers B and ξ , having respective error variables ε_B and ε_ξ , as shown in Fig. 1,




Fig. 1

can always be modeled (classical, see Proposition 12 in Annex I for a proof) as a cascaded broadcast channel $CBCS(x, y, z)$ with 3 independent BSC, 1 common attached to the sender, with error probability x , and 2 cascaded with the sender's one, each attached to one of the receivers, with respective error probability y and z , as shown in Fig. 2




Fig. 2

Partial independence between the BSC in a general $BSC(\varepsilon_B, \varepsilon_\xi)$ model, means that $z > 0$ in the equivalent cascaded model.

Maurer shown that in such case, it is possible for the legitimate partners to distill an Information Theoretically secure Secret Common Output information, thanks to error correcting and Privacy Amplifications techniques.

In case of \mathcal{P} , we can, after the Decorrelation step 5, use the following very basic technique to transform V_A and V_B in an intermediate binary flow: we can typically sample a value in $[0,1]$ like V_A or V_B with a gauge being a multiple of the variance $E \left[\left(V_A |_{\sigma_A = \sigma_\Phi} - V_B |_{\sigma_B = \sigma_\Phi} \right)^2 \right]^{1/2} = O \left(\frac{1}{\sqrt{nk}} \right)$. The multiplicative factor K is chosen such that:

$$\frac{1}{\sqrt{nk}} \ll \frac{K}{\sqrt{nk}} \ll \frac{1}{\sqrt{n}}$$

and therefore, each experiment of the protocol can lead respectively A , B , and ξ to distill an intermediate digit defined by :

$$\tilde{e}_A = \left\lfloor \frac{(V_A - \rho)\sqrt{nk}}{K} \right\rfloor \bmod 2, \tilde{e}_B = \left\lfloor \frac{(V_B - \rho)\sqrt{nk}}{K} \right\rfloor \bmod 2, \tilde{e}_\xi = \left\lfloor \frac{(V_\xi - \rho)\sqrt{nk}}{K} \right\rfloor \bmod 2$$

where ρ a translation parameter. We can easily see that the channels are BSC by remarking that changing ρ in $\rho + \frac{K}{\sqrt{nk}}$ inverses the digits and is independent of V_A , V_B , and V_ξ . With this transformation, the steps 1 to 5 of \mathcal{P} lead to a situation equivalent to the one of Maurer's scenario, as explained in the heuristic table presented above. It is shown in particular in Theorem 1 (see Annex I)

that the BSC channels are partially independent under the definition given above, and therefore the same error correcting and Privacy Amplifications techniques can be applied to obtain Information Theoretical Security for \mathcal{P} . The final digit flows after having applied those techniques are denoted $X = e_A, Y = e_B$ and $Z = e_{\xi}$.

However, it remains to ensure that the error correcting method does not bring additional information to the opponent due to the choice of code words. Indeed, contrariwise the protocol presented in [5], in our case the opponent can try the various possible combinations of choices for σ_A and σ_B , and compare them in terms of matching a code word. The design of the error correcting method should then ensure that the indistinguishability for the opponent between μ_1 and μ_2 or between μ'_1 and μ'_2 is maintained after step 5'. An example of such design is given in the proof of Theorem 1 in Annex I.

IV. Discussion about Deep Random generation

In the present Section, we discuss methods to generate Deep Random from a computing source. It may appear as a non-sense to desire to generate Deep Random from a deterministic computable program. In the real world, even a computer may access sources of randomness whose probability distribution is at least partly unknown, but it doesn't mean that we can build from it Deep Random reliable for cryptographic applications.

Generation process and strategies

We present a theoretically valid method to generate Deep Random from a computing source. That method mainly relies on infinite counters $\in \mathbb{N}^+$ as unpredictable randomness source, associated with a Cantor' style diagonal constructing argument.

If one doesn't know the date of beginning and the speed of an infinite counter, no probability distribution can be even approximated about the value of the counter at a given time, because of the unlimited nature of a counter. If performed in a physical computing source, the actual speed of the counter is impacted by all external tasks of the processor, for which no probability distribution can be estimated; the only thing that an opponent can do is estimate a rough upper bound of that speed, which, as we will see, is far from being enough to breach the Deep Randomness of the proposed method, and thus to break the secrecy of a protocol such as the one presented in Section III.

We have introduced in Section I the principle of continuous and recursive constructing process, designed in association with a given ITSP. At each step $m + 1$, the generator emulates the ITSP internally and picks (through classical randomness) a new couple of distributions $\Phi(u), \Phi'(v)$ that defeats the best possible strategy knowing all the past distributions for $t \leq m$, which is always possible for an ITSP as given by condition (P') . In this Section, we will present rigorously a method based on that principle. Nevertheless, due to the sophistication of the protocol \mathcal{P} presented in Section III, we will apply the method, for more simplicity, to the intermediate variable $E[V_A|x, y] = \frac{x \cdot y}{nk}$.

We will denote:

$$\Omega_{\#} \triangleq \left\{ \omega \in [0,1]^{2^{2n}} \mid \omega(\sigma(i), \sigma(j)) = \omega(i, j) , \forall \sigma \in \mathfrak{S}_n \right\}$$

and:

$$\langle \omega, \Phi \rangle \triangleq E \left[\left(\omega_{i,j} - \frac{x \cdot y}{nk} \right)^2 \right] = \int_{x,y \in [0,1]^n} \sum_{i,j \in \{0,1\}^n} \left(\omega_{i,j} - \frac{x \cdot y}{nk} \right)^2 \chi_i \left(\frac{x}{k} \right) \chi_j \left(\frac{y}{k} \right) \Phi(x) \Phi(y) dx dy$$

Corollary 2.

Let Φ be a sleeked distribution (under the meaning of Proposition 5 in Annex) in $\zeta(\alpha)$, then:

- (i) $M_{\Phi} \cdot M_{\Phi} - M_{\Phi} \cdot \overline{M_{\Phi}} \geq \alpha n^2 + O(n)$ and $\Delta_0(\Phi, \Phi) \geq \left(\frac{\alpha}{4} - O\left(\frac{1}{n}\right) \right)^2$ which means in other words that $\Phi \in \zeta(\alpha)$ is always self-synchronized (the operator $\Delta_0(\Phi, \Phi)$ is defined in Proposition 9 in Annex)
- (ii) For any strategy ω , there exist $\sigma_{\omega} \in \mathfrak{S}_n$ such that $\langle \omega, \Phi \circ \sigma_{\omega} \rangle \geq \frac{c}{n}$
- (iii) For any strategy ω , $\frac{1}{n!} \sum_{\sigma \in \mathfrak{S}_n} \langle \omega, \Phi \circ \sigma \rangle \geq \frac{c}{n}$

■

For (i), by considering the quadratic matrix M_Φ as a vector in a $n(n+1)/2$ dimension linear space, from Proposition 8 (whose proof is given in Annex) and Lemma 1, we have :

$$\begin{aligned} M_\Phi \cdot M_\Phi - M_\Phi \cdot \overline{M_\Phi} &\geq M_\Phi \cdot M_\Phi - \min_{\rho \in \underline{\mathfrak{S}}_{\frac{n(n+1)}{2}}} (M_\Phi \cdot \rho(M_\Phi)) + O(n) \\ &\geq \max_{\sigma \in \mathfrak{S}_n} |M_\Phi \cdot M_\Phi - M_\Phi \cdot \sigma(M_\Phi)| + O(n) \geq \alpha n^2 + O(n) \end{aligned}$$

And thus, with similar reasoning than in Proposition 9, we obtain:

$$\Delta_0(\Phi, \Phi) \geq \left(\frac{\alpha}{4} - O\left(\frac{1}{n}\right) \right)^2$$

(iii) is an immediate consequence of lemma 3:

$$\frac{1}{n!} \sum_{\sigma \in \mathfrak{S}_n} \langle \{\omega_{i,j}\}, \Phi \circ \sigma \rangle = \frac{1}{n!} \sum_{\sigma \in \mathfrak{S}_n} \langle \{\omega_{\sigma(i),\sigma(j)}\}, \Phi \rangle \geq \left\langle \frac{1}{n!} \sum_{\sigma \in \mathfrak{S}_n} \omega_{\sigma(i),\sigma(j)} \right\rangle, \Phi$$

And because $\frac{1}{n!} \sum_{\sigma \in \mathfrak{S}_n} \omega_{\sigma(i),\sigma(j)} \in \Omega_\#$, the right hand term of the above inequality can be lower bounded by $\frac{c}{n}$ as stated by Lemma 3.

And (ii) is an immediate consequence of (iii) as the average summing is lesser than the maximum element.

■

Let's present first a heuristic argument. With an infinite counter privately executed by a DRG, the moments m and $m+1$ are indistinguishable for the opponent ξ . If a set Ω_m of winning strategies at the moment m exists for ξ , then for any probability distribution Φ :

$$\frac{1}{|\Omega_m|} \sum_{\omega \in \Omega_m} \langle \omega, \Phi \rangle \geq \left\langle \frac{1}{|\Omega_m|} \sum_{\omega \in \Omega_m} \omega, \Phi \right\rangle$$

and thus, by choosing at moment $m+1$ the probability distribution Φ_{m+1} such that :

$$\left\langle \frac{1}{|\Omega_m|} \sum_{\omega \in \Omega_m} \omega, \Phi_{m+1} \right\rangle \geq \frac{c}{n}$$

(which is always possible as stated by Corollary 2) the DRG guarantees that no absolute winning strategy exist, because moment of observation cannot be determined by opponent as rather being m or $m+1$.

This heuristic argument does not explain how to practically build a Deep Random generator with classical computing resources, but it introduces the diagonal constructing process inspired from non-collaborative games' theory. Of course, no Nash' style equilibrium can be found as the DRG is continuously deviating its distribution strategy to avoid forecast from opponent.

Another interesting remark at this stage is that, in Corollary 2, we highlighted low entropy subsets of probability distributions (typically $\{\Phi \circ \sigma | \forall \sigma \in \mathfrak{S}_n\}$) in which a distribution can always be found to scramble a given strategy ω . Such subsets have entropy of order $O(n \ln n)$ although the whole set of possible probability distributions has an entropy of order $O(2^n)$, which is not manageable by classical computing resources.

Let's now explicit a constructing process, inspired from the above heuristic argument.

Let's first consider the two following recursive constructing process :

Process 1 :

Φ_0 is a distribution in $\zeta(\alpha)$; at step m :

- i) $\hat{\omega}_m$ is performing a minimum value of $\min_{\omega} \langle \omega, \Phi_m \rangle$
- ii) Ψ_{m+1} is chosen in $\zeta(\alpha)$
- iii) σ_{m+1} is chosen such that $\langle \hat{\omega}_m, \Psi_{m+1} \circ \sigma_{m+1} \rangle \geq \frac{c}{n}$ which is always possible as per Corollary 2
- iv) $\Phi_{m+1} = \Psi_{m+1} \circ \sigma_{m+1}$

Process 2 :

Φ_0 is a distribution in $\zeta(\alpha)$; at step m :

- i) $\hat{\omega}_m$ is performing a minimum value of $\min_{\omega} \langle \omega, \frac{1}{m} \sum_{s=1}^m \Phi_s \rangle$
- ii) Ψ_{m+1} is chosen in $\zeta(\alpha)$
- iii) σ_{m+1} is chosen such that $\langle \hat{\omega}_m, \Psi_{m+1} \circ \sigma_{m+1} \rangle \geq \frac{c}{n}$ which is always possible as per Corollary 2
- iv) $\Phi_{m+1} = \Psi_{m+1} \circ \sigma_{m+1}$

The process 1 involves fast variations, but, if the choice of Ψ_{m+1} and σ_{m+1} is deterministic at each step m , it can have short period (typically period of 2), which makes it unsecure. The process 2 has no period but its variations are slowing down when m is increasing. We can justify that the process 2 has no period by seeing that, if it would have a period, then $\frac{1}{m} \sum_{s=1}^m \Phi_s$ would converge, which would contradict the fact that $\langle \hat{\omega}_m, \Phi_{m+1} \rangle \geq \frac{c}{n}$.

Thus, by combining the two process, we get a sequence with both fast variations and no periodic behavior. A method to combine both process is the following :

The Recursive Generation Process :

Φ_0 is a distribution in $\zeta(\alpha)$; at step m :

- i) $\hat{\omega}_m$ is performing a minimum value of $\min_{\omega} \langle \omega, \Phi_m \rangle$
- ii) $\hat{\omega}'_m$ is performing a minimum value of $\min_{\omega} \langle \omega, \frac{1}{m} \sum_{s=1}^m \Phi_s \rangle$
- iii) Ψ_{m+1} and Ψ'_{m+1} are chosen in $\zeta(\alpha)$
- iv) σ_{m+1} is chosen such that $\langle \hat{\omega}_m, \Psi_{m+1} \circ \sigma_{m+1} \rangle \geq \frac{c}{n}$ which is always possible as per Corollary 2

- v) σ'_{m+1} is chosen such that $\langle \hat{\omega}'_m, \Psi'_{m+1} \circ \sigma'_{m+1} \rangle \geq \frac{c}{n}$ which is always possible as per Corollary 2
- vi) $\Phi_{m+1} = \alpha_{m+1} \Psi_{m+1} \circ \sigma_{m+1} + (1 - \alpha_{m+1}) \Psi'_{m+1} \circ \sigma'_{m+1}$ where $\alpha_{m+1} \in [0,1]$ can be picked randomly

Such a process is secure « against the past », even if the choices at each step are deterministic, but it is not secure « against the future » if the choices at each step are deterministic. Being secure « against the future » means that if the opponent runs the recursive process on its own and is « in advance » compared to the legitimate partner, it still cannot obtain knowledge about the current value of Φ_m . So, for the process to remain secure also against the future, it is necessary that the choices at each step involve randomness (but not necessarily Deep Randomness) with maximum possible entropy among $\zeta(\alpha)$.

It is also important that the DRG is capable to pick a distribution within the widest possible sub-set of $\zeta(\alpha)$, otherwise, any restriction in such possible ‘picking set’ would add prior information for the opponent about the possible distribution, and would therefore reduce $\mathfrak{S}_\zeta \supseteq \zeta(\alpha)$.

Let’s first notice that the set of possible quadratic matrix is (if Φ is restricted over $\{0,1\}^n$) the convex envelop of all matrix in the set:

$$\{\sigma(S_r) | \sigma \in \mathfrak{S}_n, r \in \mathbb{N}_n\}$$

where $S_r = \begin{pmatrix} C_r & 0 \\ 0 & 0 \end{pmatrix}$, with the notation of Proposition 9 and Lemma 1. We can easily calculate that, for r not too close from 0 or:

$$\|S_r - \bar{S}_r\|_c = \begin{cases} \frac{r(r-1)}{n(n-1)} & \Leftrightarrow r < \frac{n}{2} \\ \frac{(n-r)(n-r-1)}{n(n-1)} & \Leftrightarrow r \geq \frac{n}{2} \end{cases}$$

and therefore to determine if $\delta_x \in \zeta(\alpha)$.

It is possible to combine several distributions to create a more elaborated one still being in $\zeta(\alpha)$. A combination method is for instance:

$$\{\Psi_1, \dots, \Psi_S\} \mapsto \frac{1}{S} \sum_{s=1}^S \Psi_s \circ \sigma_{\Psi_s} \circ \mu$$

We can summarize the above findings in the following Deep Random generation process. A DRG is running both:

- a recursive and continuous generation process in which $S \geq 1$ continuous sequences $\{\Phi[p]_m\}_{p \in \mathbb{N}_S^*, m \in \mathbb{N}}$ are running in parallel according to a Recursive Generation Process presented above. The initial distribution of each sequence is picked among a large predefined list of distributions in $\zeta(\alpha)$, that can contain for instance all the Dirac distributions associated to the $\{0,1\}^n$ vectors, plus other more elaborated distributions that could be determined. The largest is this initial list, the better is the quality of Deep Random. At each step of a given sequence, $\Psi[p]_{m+1}$ and $\Psi'[p]_{m+1}$ (as per the notations used in description of the Recursive Generation Process given above) can be chosen among the initial list or among the current

values of the other sequences; it can also be decided (over random decision) to update the current value by a combination of all the current values of the sequences using the Internal Combination Process presented above. The quality of the Deep Random depends on the variety of the initial list and also on the increasing number of steps (rounds) performed in each sequences. An argument about the minimum number of steps will be discussed below; once the minimum number of steps evaluated for the generator has been reached, it enters into its « maturity » period.

- a process of selection upon request in which the DRG can at any moment select and combine some current values among the S sequences in order to elect a distribution to be used for the execution of a secure communication (typically through the protocol presented in Section 0).

Finally the elected distribution Φ must also be transformed by the permutative sleeking transformation introduced in Lemma 3 so that it can be used safely in protocol \mathcal{P} .

Practically, Deep Random is of the highest quality if generators can benefit from the most possible time, entropy and calculation power. And on the other hand, sources of Deep Random can be combined to produce other sequences that can become themselves sources. This means that certain entities can be « specialized » in the network to generate centrally highest quality of Deep Random sources (we will talk about level 1 Deep Random generators), and then can deliver those sources to DRG's owner which then use them as seeds to derive their own local sequences for the purpose of their secure communications (we will talk about level 2 Deep Random generators).

When such a collaboration is occurring, the seed must of course be securely transmitted from a level 1 generator to the level 2 generator. To that extend, the asymmetry of the protocol \mathcal{P} presented in Section III can there be used in an adapted way ; indeed, in \mathcal{P} , the party called A needs to generate more Deep Randomness. The role of A should then be played by a level 1 generator, whereas the role of B should be played by the level 2 generator.

To close this discussion, let's give, to open ideas, a simple example of elaborated initial seed Φ_0 ; we suggest the sleeked distribution given by :

$$P\left(\sum_{s=1}^{n/2} x_s = t\right) = P\left(\sum_{s=n/2+1}^n x_s = t'\right) = \frac{2}{n}, \forall t, t' \in \{0, \dots, n/2\}$$

Its quadratic matrix is:

$$\begin{pmatrix} 1/2 & \dots & 1/3 & \left| & 1/4 & \dots & 1/4 \right. \\ \vdots & \ddots & \vdots & \left| & \vdots & & \vdots \right. \\ 1/3 & \dots & 1/2 & \left| & 1/4 & \dots & 1/4 \right. \\ \hline 1/4 & \dots & 1/4 & \left| & 1/2 & \dots & 1/3 \right. \\ \vdots & & \vdots & \left| & \vdots & \ddots & \vdots \right. \\ 1/4 & \dots & 1/4 & \left| & 1/3 & \dots & 1/2 \right. \end{pmatrix}$$

And it verifies:

$$\|M_{\Phi_0} - \overline{M_{\Phi_0}}\|_c = \frac{1}{12} + o\left(\frac{1}{n}\right)$$

More elaborated distributions, belonging to wide subsets, can be imagined by replacing $\sum_{s=1}^{n/2} x_s$ by other families of algebraic forms.

Argument about the minimum number of steps to reach the maturity period of the Recursive Generation Process

We consider a convex and compact subset Ω of possible strategies for the opponent (for any subset, its closed convex envelop corresponds to that characteristic). We consider in the followings the sequence of probability distributions $\{\Phi_m\}_{m \in \mathbb{N}^*}$ constructed as follows :

$$\Phi_1 = \Phi$$

where Φ is a sleeked distribution in $\zeta(\alpha)$, called the seed of the sequence ; $\hat{\omega}_m$ is performing a minimum value in :

$$\min_{\omega \in \Omega} \frac{1}{m} \sum_{s=1}^m \langle \omega, \Phi_s \rangle$$

Φ_{m+1} is chosen such that:

$$\langle \hat{\omega}_m, \Phi_{m+1} \rangle \geq \frac{C}{n}$$

Diagonal Sequence Property.

There exists two constants C' and C'' such that, for any N verifying $\frac{N}{\ln(N)} \geq C' \dim(\Omega)$,

$$\min_{\omega \in \Omega} \frac{1}{N} \sum_{s=1}^N \langle \omega, \Phi_s \rangle \geq \frac{C''}{n}$$

■

More precisely, we chose Φ_{m+1} as :

$$\Phi_{m+1} = \frac{1}{2} (1 + \Phi \circ \sigma_m) \tag{IV.1}$$

where σ_m is a permutation such that $\langle \hat{\omega}_m, \Phi \circ \sigma_m \rangle \geq \frac{C}{n}$.

Let's set :

$$\hat{Q}_m(\omega - \hat{\omega}_m) \triangleq \frac{1}{m} \sum_{s=1}^m \langle \omega, \Phi_m \rangle - \frac{1}{m} \sum_{s=1}^m \langle \hat{\omega}_m, \Phi_m \rangle$$

$$\hat{\Sigma}_m \triangleq \frac{1}{m} \sum_{s=1}^m \langle \hat{\omega}_m, \Phi_s \rangle$$

$$\langle \omega, \Phi_m \rangle = Q_m(\omega - \omega_m) + \theta_m$$

\hat{Q}_m and Q_m are positive quadratic forms over $\omega \in \Omega$ derived by respective distributions $\frac{1}{m} \sum_{s=1}^m \Phi_s(x) \Phi_s(y)$ and $\Phi_m(x) \Phi_m(y)$. (IV. 1) ensures that \hat{Q}_m and Q_m are non-degenerate in Ω . By considering a basis that is simultaneously \hat{Q}_m -orthonormal and Q_m -orthogonal, we obtain the 2 following relations (remind that $\hat{\omega}_m$ is the minimum over the quadratic form \hat{Q}_m):

$$\frac{1}{m} \sum_{s=1}^m \langle \omega, \Phi_s \rangle - \frac{1}{m} \sum_{s=1}^m \langle \hat{\omega}_m, \Phi_s \rangle = \|\omega - \hat{\omega}_m\|_{\hat{Q}_m}^2$$

$$|\langle \omega, \Phi_{m+1} \rangle - \langle \hat{\omega}_m, \Phi_{m+1} \rangle| \leq 2\sqrt{\dim(\Omega)} \|\omega - \hat{\omega}_m\|_{\hat{Q}_m}$$

We have then:

$$\begin{aligned} & \frac{1}{m+1} \sum_{s=1}^{m+1} \langle \omega, \Phi_s \rangle \\ &= \frac{m}{m+1} \left(\hat{\Sigma}_m + \left(\frac{1}{m} \sum_{s=1}^m \langle \omega, \Phi_s \rangle - \frac{1}{m} \sum_{s=1}^m \langle \hat{\omega}_m, \Phi_s \rangle \right) \right) \\ &+ \frac{1}{m+1} (\langle \omega, \Phi_{m+1} \rangle - \langle \hat{\omega}_m, \Phi_{m+1} \rangle) + \frac{1}{m+1} \langle \hat{\omega}_m, \Phi_{m+1} \rangle \end{aligned} \quad (IV. 2)$$

Applying the 2 above relations to (IV. 2), we get:

$$\begin{aligned} \hat{\Sigma}_{m+1} &\geq \frac{m}{m+1} \hat{\Sigma}_m + \frac{m}{m+1} \|\omega - \hat{\omega}_m\|_{\hat{\Omega}}^2 - \frac{2\sqrt{\dim(\Omega)}}{m+1} \|\omega - \hat{\omega}_m\|_{\hat{\Omega}} + \frac{1}{m+1} \frac{C}{4n} \\ &\geq \frac{m}{m+1} \hat{\Sigma}_m + \frac{1}{m+1} \frac{C}{4n} - \frac{\dim(\Omega)}{m(m+1)} \end{aligned}$$

where the last inequality is obtained by taking the minimum of $\frac{m}{m+1} X^2 - \frac{2\sqrt{\dim(\Omega)}}{m+1} X$ over X .

Let's consider the recurring sequence:

$$(m+1)X_{m+1} = mX_m + \frac{C}{4n} - \frac{\dim(\Omega)}{m}$$

$$X_1 = \hat{\Sigma}_1$$

By recurring argument, $\hat{\Sigma}_m \geq X_m$ and thus:

$$\hat{\Sigma}_m \geq \frac{C}{4n} - \dim(\Omega) O\left(\frac{\ln(m)}{m}\right)$$

which achieves the proof. ■

If we consider first Ω as the full set of strategies (e.g. $\left[0, \frac{1}{k}\right]^{2^{2n}}$), the dimension is 2^{2n} and thus the constructing process must be iterated exponential times to reach positive lower bound. On the other hand, we make the excessive assumption that the opponent has a full knowledge of the distributions chosen by the DRG at each step m (of the sequences $\{\Phi_s\}_{s \leq m}$) to be able to build optimal $\hat{\omega}_m$. In practice, if we consider the protocol \mathcal{P} introduced in Section III, we can see that ξ has never more knowledge about a distribution than a draw of 1 degraded vector i , which means that we can restrict the set of strategies to a subset Ω^* with dimension $O(n)$, and thus the recommended number of iterations for a recursive constructing process before being able to trustedly pick a distribution is $O(n \ln(n))$. This result also means that the generated sequence is $O(k)$ -undistinguishable under the definition given in Section I, for for $\frac{N}{\ln(N)} \geq C' \dim(\Omega)$.

With the same reasoning than the one of Diagonal Sequence Property, we can obtain a more general result:

Generalized Diagonal Sequence Property.

There exists two constants $C'(\alpha)$ and $C''(\alpha)$ such that, for any N verifying $\frac{N}{\ln(N)} \geq C' \dim(\Omega)$, and for any $\{\Phi_1, \dots, \Phi_N\} \in \zeta(\alpha)$ and $\{\Phi'_1, \dots, \Phi'_N\} \in \zeta(\alpha)$, and any permutation σ_s synchronizing Φ_s and Φ'_s :

$$\min_{\omega \in \Omega} \frac{1}{N} \sum_{s=1}^N \langle \omega, \Phi_s, \Phi'_s \circ \sigma_s \rangle \geq \frac{C''}{n}$$

This result also means that when $\{\Phi_1, \dots, \Phi_N\}$ and $\{\Phi'_1, \dots, \Phi'_N\}$ are generated by DRGs, from the principle of objectivity explained in Section I, an opponent should choose the same strategy at each step s , and therefore the above inequality implies that whatever is that strategy, in average we have $\langle \omega, \Phi_s, \Phi'_s \circ \sigma_s \rangle \gg_k E[(V_A - V_B)^2]$; which gives Advantage Distillation for $\frac{N}{\ln(N)} \geq C' \dim(\Omega)$.

V. Conclusion and further directions

We proved that Information Theoretical Security could be reached under Deep Random assumption.

Let's remark that when the alphabets \mathcal{X} , \mathcal{Y} and \mathcal{Z} introduced in Section I are all equal to $\{0,1\}$, and if each digit issued by the protocol is managed independently of the others, then the Security Property:

$$R^*(\mathcal{P}, Z) \triangleq H(X|Z) - H(X|Y) \geq (1 - \epsilon)H(X) \quad (SP)$$

introduced in Section I can equivalently be written as: whatever are $\epsilon, \epsilon' > 0$, there exists a set of parameters U for the protocol \mathcal{P} such that the two conditions above are verified:

- (i) $2 \left(\min(P(e_A \neq e_B), 1 - P(e_A \neq e_B)) \right) \leq \epsilon$
- (ii) $2 \left(\max \left(P(e_\xi = e_B), 1 - P(e_\xi = e_B) \right) - \frac{1}{2} \right) \leq \epsilon'$

where the opponent ξ is passive, with unlimited access to the public information and with unlimited power of storage and computation; e_A, e_B, e_ξ being estimations of an average digit of information exchanged between A and B through \mathcal{P} respectively from A, B, ξ .

The obtained result, apparently contradicting Shannon's pessimistic theorem, is possible thanks to the nature of Deep Randomness, that prevents the use of Bayesian inference from public information, because not only draws but also probability distributions themselves are unknown to the beholding opponent ξ .

We also proposed a method to generate Deep Randomness (Section IV) from classical computation resources, thanks to recursive and continuously executed algorithm, based on Cantor' style diagonal constructing method.

In this work, our main objective was to expose this new idea and to prove the existence of a working protocol. But of course, once done, the next question is: if such protocol exists, what is the best one? In cryptology and communication sciences, there exist many criteria of quality for a protocol. We will concentrate here on the most critical one for practical implementation which is the burden of exchanged data needed to obtain an Information Theoretically secure digit of information.

Considering a ITSP \mathcal{P} , we define

$$Q_{X,Y}(\mathcal{P})$$

as the quantity of digit exchanged by A and B on the main channel through the protocol \mathcal{P} so that A and B can collect their respective value of X and Y .

The question raised above comes down to determine the maximum:

$$C = \sup_{\mathcal{P}, G \in \Gamma_{\mathfrak{S}}} \left(\frac{\inf_{Z \in \Omega_{\mathfrak{S}}(G, \mathcal{P})} R^*(\mathcal{P}, Z)}{Q_{X,Y}(\mathcal{P})} \right)$$

The quantity C is called the Cryptologic Limit. It is then a constant in $[0,1]$. The Theorem 1 can then be formulated simply as:

Theorem:

Under the Deep Random assumption, $C > 0$

The research of the Cryptologic Limit opens, to my opinion, an interesting field of investigations. Empirical research of specifically designed protocols may give rapidly interesting improvements to the presently proposed example; estimation of performance can be easily done by simulation rather than painful theoretical examination of upper-bounds. But the theoretical full resolution of the cryptologic limit question will need new ideas.

I hope this challenge will create enthusiasm for the largest possible number of curious minds interested in cryptology, information theory and communication sciences.

References

- [1] C. E. Shannon, « Communication theory of secrecy systems », Bell Syst. Tech. J., Vol. 28, pp. 656-715, Oct. 1949
- [2] A. N. Kolmogorov, « On Tables of Random Numbers », Sankhya. Indian Journal of Statistics A, 25(4) :369-376
- [3] C. H. Bennet and G. Brassard, « Quantum cryptography and its application to provable secure key expansion, public-key distribution and coin-tossing », Proc. IEEE International Conference on Computers, Systems and Signal Processing, Bangalore, India, pp. 175-179, Dec. 1984
- [4] C. H. Bennet, G. Brassard and J.-M. Robert, « Privacy Amplification by Public Discussion », SIAM J. COMPUT., Vol. 17, No. 2, Apr. 1988
- [5] U. M. Maurer, « Secret Key Agreement by Public Discussion from Common Information », IEEE Transactions on Information Theory, Vol. 39, No. 3, May 1993
- [6] C. Cachin and U. M. Maurer, « Unconditional Security Against Memory-Bounded Adversaries », Proceeding of CRYPTO '97, Lecture Notes in Computer Science, Springer, 1997
- [7] E. T. Jaynes, « Prior Probabilities », IEEE Transactions On Systems Science and Cybernetics, vol. sec-4, no. 3, 1968, pp. 227-241
- [8] R. T. Cox, « Probability, frequency and reasonable expectation », American Journal of Physics 17 (1946) 1-13
- [9] C. E. Shannon, « A Mathematical Theory of Communication », The Bell System Technical Journal, Vol. 27, pp. 379–423, 623–656, July, October, 1948
- [10] P. S. de Laplace, « A philosophical essay on Probabilities », Paris 1814
- [11] R. Carnap, « The Continuum of Inductive Methods », University of Chicago Press, 1952
- [12] I. J. Good, « Maximum entropy for hypothesis formulation », Ann. Math. Statist., 34, pp. 911-930, 1963
- [13] T. de Valroger, “A relation between conditional entropy and conditional expectation to evaluate secrecy systems” Arxiv 2017: <https://arxiv.org/abs/1709.01550>
- [14] U. Maurer, S. Wolf, “Unconditionally Secure Key Agreement and the Intrinsic Conditional Information”, IEEE transactions on information theory, vol. 45, no. 2, March 1999

Who is the author ?

I have been an engineer in computer science for 20 years. My professional activities in private sector are related to IT Security and digital trust, but have no relation with my personal research activity in the domain of cryptology. If you are interested in the topics introduced in this article, please feel free to establish first contact at tdevalroger@gmail.com

VI. Annex: proof of technical results

The Annex contains the proofs of Theorem 1 and 2, showing that the protocol presented in Section III is an ITSP. They are preceded by technical preliminary results:

- Propositions 1,2,3,4 and 11 deal with basic Bernoulli variable and Chernoff bound style results. Proposition 5 introduces Permutative sleeking transform and gives basic property. Propositions 6,7 and 8 are simple useful inequalities. Propositions 9 and 10 give useful results on c-norm and tidying permutations. Propositions 12, 13, 15 and 16 give an analysis of Broadcast BSC model. Proposition 14 justifies restricted form for the opponent strategy.
- Then we show 4 technical lemmas. Lemma 1 and 1' give important criteria to synchronize 2 distributions. Lemma 2 give reversing law of Permutative sleeking transform and applies it a category of lower-bounds. Lemma 3 is the core technical result that shows how the opponent is misled when choosing $\sigma_d[i]$ and $\sigma'_d[j]$ from the published information of the protocol.

Proposition 1.

For all $i \in \{0,1\}^n$ and $l \in \mathbb{N}_n^*$, stands the relation : $\psi_{i,l}\left(\frac{x}{k}\right) = \sum_{r=l}^{|i|} \beta_{l,r} \left(\frac{1}{k}\right) \psi_{i,r}(x)$

■ Let's start by expressing the multinomial $\chi_i\left(\frac{x}{k}\right)$ in the basis $\{\chi_i(x)\}_{i \in \{0,1\}^n}$:

$$\begin{aligned}
 \chi_j\left(\frac{x}{k}\right) &= \frac{1}{k^{|j|}} \sum_{u \supset j} \frac{(-1)^{|u \setminus j|}}{k^{|u \setminus j|}} \Pi_u(x) \\
 &= \frac{1}{k^{|j|}} \sum_{u \supset j} \frac{(-1)^{|u \setminus j|}}{k^{|u \setminus j|}} \sum_{u' \supset u} \chi_{u'}(x) \\
 &= (-1)^{|j|} \sum_{u \supset j} \left(\sum_{j \subset v \subset u} \left(-\frac{1}{k}\right)^{|v|} \right) \chi_u(x) \\
 &= (-1)^{|j|} \sum_{u \supset j} \left(\sum_{r=|j|}^{|u|} \binom{|u|-|j|}{r-|j|} \left(-\frac{1}{k}\right)^r \right) \chi_u(x) \\
 &= \left(\frac{1}{k}\right)^{|j|} \sum_{u \supset j} \left(1 - \frac{1}{k}\right)^{|u|-|j|} \chi_u(x)
 \end{aligned}$$

We thus have :

$$\begin{aligned}
 \psi_{i,l}\left(\frac{x}{k}\right) &= \sum_{j|i, j=l} \left(\frac{1}{k}\right)^{|j|} \sum_{s \supset j} \left(1 - \frac{1}{k}\right)^{|s|-|j|} \chi_s(x) \\
 &= \sum_{s|i, s \geq l} \left(\sum_{\substack{j|i \\ j \subset s}} \left(\frac{1}{k}\right)^{|j|} \left(1 - \frac{1}{k}\right)^{|s|-|j|} \right) \chi_s(x) \\
 &= \sum_{s|i, s \geq l} \left(1 - \frac{1}{k}\right)^{|s|} \left(\sum_{\substack{j|i \\ j \subset s}} \left(\frac{1}{k-1}\right)^{|j|} \right) \chi_s(x)
 \end{aligned}$$

Then :

$$\sum_{\substack{j|i \\ j \subset s}} \left(\frac{1}{k-1}\right)^{|j|} = \sum_{\substack{j|i \\ j \subset s}} \left(\frac{1}{k-1}\right)^{l+j \cap (s \setminus i)} = \sum_{j \subset (s \setminus i)} \sum_{\substack{j' \subset (s \cap i) \\ |j'|=l}} \left(\frac{1}{k-1}\right)^{l+|j'|}$$

$$\begin{aligned}
&= \left(\frac{1}{k-1}\right)^l \binom{|s \cap i|}{l} \sum_{j \subset (s \setminus i)} \left(\frac{1}{k}\right)^{|j|} = \left(\frac{1}{k-1}\right)^l \binom{|s \cap i|}{l} \sum_{t=0}^{|s \setminus i|} \left(\frac{1}{k-1}\right)^t \binom{|s \setminus i|}{t} \\
&= \left(\frac{1}{k-1}\right)^l \binom{|s \cap i|}{l} \left(\frac{k}{k-1}\right)^{|s| - i \cdot s}
\end{aligned}$$

And finally,

$$\psi_{i,l} \left(\frac{x}{k}\right) = \sum_{s \mid i \cdot s \geq l} \left(\frac{1}{k-1}\right)^l \binom{|s \cap i|}{l} \left(\frac{k-1}{k}\right)^{i \cdot s} \chi_s(x) = \sum_{r=l}^{|i|} \beta_{l,r} \left(\frac{1}{k}\right) \psi_{i,r}(x) \quad \blacksquare$$

It is well known that Bernoulli coefficients can be approximated by a Gaussian distribution $N(u, v) \triangleq \frac{1}{\sqrt{2\pi v}} e^{-\frac{u^2}{2v}}$. Proposition 2 gives an upper bound for infinite branch, while Proposition 4 gives a « central limit » style approximation.

Proposition 2.

For all Δ , we have the upper bound : $\beta_{l,kl+\Delta} \left(\frac{1}{k}\right) \leq e^{-\frac{\Delta^2}{2k^2l}}$

■ We can write the quotient :

$$\begin{aligned}
\frac{\beta_{l,kl+\Delta} \left(\frac{1}{k}\right)}{\beta_{l,kl} \left(\frac{1}{k}\right)} &= \prod_{s=1}^{\Delta} \frac{(k-1)(kl+s)}{k((k-1)l+s)} = \prod_{s=1}^{\Delta} \frac{1 + \frac{(k-1)s}{k(k-1)l}}{1 + \frac{ks}{k(k-1)l}} \\
&= \exp \left(\sum_{s=1}^{\Delta} \left(\ln \left(1 + \frac{(k-1)s}{k(k-1)l} \right) - \ln \left(1 + \frac{ks}{k(k-1)l} \right) \right) \right)
\end{aligned}$$

Considering that $f(x) = \ln(1+x)$ is 1-lipschitz over \mathbb{R}^+ and increasing, we then conclude easily :

$$\frac{\beta_{l,kl+\Delta} \left(\frac{1}{k}\right)}{\beta_{l,kl} \left(\frac{1}{k}\right)} \leq \exp \left(- \sum_{s=1}^{\Delta} \frac{s}{k(k-1)l} \right) \leq e^{-\frac{\Delta^2}{2k^2l}} \quad \blacksquare$$

As a consequence of Propositions 1 & 2, we obtain a Chernoff-style bound for the legitimate partners evaluation gap:

Proposition 3.

Let a be a positive integer such that $a < \frac{x \cdot y}{k}$, then we have the following upper bound for the legitimate partner evaluation gap :

$$P(|V_A - V_B| \geq 2a|x, y) \leq 2ne^{-\frac{na^2}{2E[V_A]}}$$

■ For more simplicity, we will suppose in the following that $x, y \in \{0,1\}^n$ rather than $[0,1]^n$, we then have :

$$\begin{aligned} P(|V_A - E[V_A]| \geq a|x, y) &= \sum_{|r-nE[V_A]| \geq na} \sum_{x,j=r}^j \chi_j\left(\frac{y}{k}\right) \\ &= \sum_{|r-nE[V_A]| \geq na} \sum_{s=r}^n \beta_{r,s}\left(\frac{1}{k}\right) \psi_{x,s}(y) \text{ from Proposition 1.} \end{aligned}$$

and noticing that $\psi_{x,s}(y) = \begin{cases} 1 & \text{if } x \cdot y = s \\ 0 & \text{else} \end{cases}$:

$$= \sum_{|r-nE[V_A]| \geq na} \beta_{r,x \cdot y}\left(\frac{1}{k}\right) \leq ne^{-\frac{na^2}{2E[V_A]}} \text{ from Proposition 2.}$$

(notice also that $\left(\left(r \leq \frac{x \cdot y}{k} - na\right) \text{ or } \left(r \geq \frac{x \cdot y}{k} + na\right)\right) \Rightarrow ((x \cdot y \leq kr - kna) \text{ or } (x \cdot y \geq kr + kna))$). We conclude by writing that :

$$P(|V_A - V_B| \geq 2a|x, y) \leq P(|V_A - E[V_A]| \geq a|x, y) + P(|V_B - E[V_B]| \geq a|x, y)$$

■

The classical Chernoff bound results are not efficient at neighborhood of zero (which, in our context, is due to the increasing of the degradation parameter k), this is why we need to adapt them.

Proposition 4.

$\forall u, |u| \leq u_m \ll \theta w$, there exists two constants C_1, C_2 such that :

$$\beta_{w\theta+u,w}(\theta) = N(u, w\sigma^2) \left(1 + \ll \left(C_1 \frac{u_m}{\theta w}\right)\right) e^{-\ll \left(C_2 \frac{u_m^3}{w^2 \theta^2}\right)} \text{ with variance } \sigma^2 \triangleq \theta(1 - \theta).$$

■ We can consider $\beta_{w\theta+u,w}(\theta)$ as continuous in u by extending $n!$ to $\Gamma(n)$. By Stirling formula, there exists C such that :

$$\Gamma(x) = \sqrt{2\pi} x^{x+\frac{1}{2}} e^{-x} \left(1 + \frac{C(x)}{x}\right) \text{ with } C(x) \leq C$$

On the other hand, we have, by Taylor's inequality, for any $|y| \leq y_m < 1$:

$$\left| -\ln(1-y) - \left(y + \frac{y^2}{2}\right) \right| \leq \frac{y_m^3}{3(1-y_m)^3}$$

We then develop $\beta_{w\theta+u,w}(\theta)$ using the two above inequalities, which gives, by noting $= w\theta + u$:

$$\beta_{w\theta+u,w}(\theta) = \sqrt{\frac{w}{2\pi r(w-r)} \frac{1 + \frac{C(w)}{w}}{\left(1 + \frac{C(r)}{r}\right) \left(1 + \frac{C(w-r)}{w-r}\right)}} \exp\left(-\frac{u^2}{2w\theta(1-\theta)}\right) + \frac{u^3}{2w^2} \left(\frac{1}{\theta^2} - \frac{1}{(1-\theta)^2}\right) + o\left(\theta w \frac{u_m^3}{w^3 \theta^3}\right)$$

There exists two constants C_1, C_2 such that :

$$\sqrt{\frac{w}{2\pi r(w-r)} \frac{1 + \frac{C(w)}{w}}{\left(1 + \frac{C(r)}{r}\right) \left(1 + \frac{C(w-r)}{w-r}\right)}} = \sqrt{\frac{1}{2\pi w\theta(1-\theta)}} \left(1 + o\left(C_1 \frac{u_m}{\theta w}\right)\right)$$

$$\exp\left(\frac{u^3}{2w^2} \left(\frac{1}{\theta^2} - \frac{1}{(1-\theta)^2}\right) + o\left(\theta w \frac{u_m^3}{w^3 \theta^3}\right)\right) = e^{o\left(C_2 \frac{u_m^3}{w^2 \theta^2}\right)}$$

which ends the proof. ■

We will need, in order to demonstrate the main « Synchronized Strategy lemma » (Lemma 3), to ensure that the probability distributions are « sufficiently regular », meaning that they avoid brutal variation.

To that extent, we will use a prior transformation of a probability distribution, called « permutative sleeking ». A permutative sleeking kernel γ is a function $\mathbb{N}_n^* \rightarrow [0,1]$ (note that it is impossible that $|\sigma| = 1$ and thus the component for 1 can be ignored) that verifies :

$$\sum_{\sigma \in \mathfrak{S}_n} \gamma(|\sigma|) = 1$$

Then, for a given probability distribution Φ , the permutative sleeking transformation of Φ is :

$$\Phi \mapsto T_\gamma[\Phi] \triangleq \sum_{\sigma \in \mathfrak{S}_n} \gamma(|\sigma|) \Phi \circ \sigma$$

A simple property of permutative sleeking transformations is the following :

Proposition 5.

The space of permutative sleeking transformations, fitted with composition operation, is stable.

■ Let's first remark that, for any $\sigma, \mu \in \mathfrak{S}_n$:

$$|\sigma| = |\mu \circ \sigma \circ \mu^{-1}|$$

Indeed, if $i \in \mu(\text{supp}(\sigma))$, then i can uniquely be written $i = \mu(j)$ with $j \in \text{supp}(\sigma)$. Then $\mu \circ \sigma \circ \mu^{-1}(i) = \mu \circ \sigma(j) \neq i$ because $\sigma(j) \neq j$, which implies that $|\sigma| \leq |\mu \circ \sigma \circ \mu^{-1}|$. Reversely, if $i \notin \mu(\text{supp}(\sigma))$, then i can uniquely be written $i = \mu(j')$ with $j' \notin \text{supp}(\sigma)$, which implies that $\sigma(j') = j'$ and following $\mu \circ \sigma \circ \mu^{-1}(i) = \mu \circ \sigma(j') = \mu(j') = i$; thus $n - |\sigma| \leq n - |\mu \circ \sigma \circ \mu^{-1}|$, and finally $|\sigma| = |\mu \circ \sigma \circ \mu^{-1}|$.

In addition, it is obvious that $\mu \circ \sigma \circ \mu^{-1} = \text{Id}_{\mathfrak{S}_n} \Leftrightarrow \sigma = \text{Id}_{\mathfrak{S}_n}$, and thus by cardinality $\{\sigma \mid |\sigma| = p\} = \{\mu \circ \sigma \circ \mu^{-1} \mid |\sigma| = p\} \forall \mu \in \mathfrak{S}_n$.

We can then write the composition of two permutative sleeking kernels :

$$T_\gamma \circ T_\delta[\Phi] = \sum_{\sigma \in \mathfrak{S}_n} \gamma(|\sigma|) \sum_{\mu \in \mathfrak{S}_n} \delta(|\mu|) \Phi \circ \sigma \circ \mu = \sum_{s \in \mathfrak{S}_n} \left(\sum_{\substack{\sigma, \mu \in \mathfrak{S}_n \\ \sigma \circ \mu = s}} \gamma(|\sigma|) \delta(|\mu|) \right) \Phi \circ s$$

Let's now show that the coefficient depends only on $|s|$:

$$\sum_{\substack{\sigma, \mu \in \mathfrak{S}_n \\ \sigma \circ \mu = s}} \gamma(|\sigma|) \delta(|\mu|) = \sum_{\sigma \in \mathfrak{S}_n} \gamma(|\sigma|) \delta(|\sigma^{-1} \circ s|)$$

Then for any $\mu \in \mathfrak{S}_n$:

$$\begin{aligned} \sum_{\sigma \in \mathfrak{S}_n} \gamma(|\sigma|) \delta(|\sigma^{-1} \circ \mu \circ s \circ \mu^{-1}|) &= \sum_{\sigma \in \mathfrak{S}_n} \gamma(|\sigma|) \delta(|\mu^{-1} \circ \sigma^{-1} \circ \mu \circ s|) \\ &= \sum_{\sigma \in \mathfrak{S}_n} \gamma(|\mu \circ \sigma \circ \mu^{-1}|) \delta(|\mu^{-1} \circ \sigma^{-1} \circ \mu \circ s|) = \sum_{\sigma' \in \mathfrak{S}_n} \gamma(|\sigma'|) \delta(|\sigma'^{-1} \circ s|) \end{aligned}$$

with the change of variable $\sigma' = \mu \circ \sigma \circ \mu^{-1}$, that courses the set $\{\sigma' \mid |\sigma'| = |\sigma|\}$ as seen above. This shows that the coefficient only depends on $|s|$. In addition, obviously :

$$\sum_{s \in \mathfrak{S}_n} \sum_{\sigma \in \mathfrak{S}_n} \gamma(|\sigma|) \delta(|\sigma^{-1} \circ s|) = \left(\sum_{\sigma \in \mathfrak{S}_n} \gamma(|\sigma|) \right) \left(\sum_{s' \in \mathfrak{S}_n} \gamma(|s'|) \right) = 1$$

which means that $T_\gamma \circ T_\delta$ is still a permutative sleeking transformation. ■

The following is a very simple result for quadratic upper bounds :

Proposition 6.

Let f and $g : I \rightarrow \mathbb{R}$ be two square-integrable functions, $\alpha \in]0,1[$, and ϕ be a probability distribution function over I , with $\int_I f(x)^2 \phi(x) dx \geq A$ and $\int_I g(x)^2 \phi(x) dx \leq a \leq A \left(\frac{\alpha}{1-\alpha} \right)^2$, then :

$$\int_I (\alpha f(x) + (1-\alpha)g(x))^2 \phi(x) dx \geq \alpha^2 A \left(1 - \frac{1-\alpha}{\alpha} \sqrt{\frac{a}{A}} \right)^2$$

■ Obvious from Cauchy-Schwarz inequality :

$$\begin{aligned}
& \int_I (\alpha f(x) + (1 - \alpha)g(x))^2 \phi(x) dx \\
& \geq \alpha^2 \int_I f(x)^2 \phi(x) dx + (1 - \alpha)^2 \int_I g(x)^2 \phi(x) dx - 2\alpha(1 \\
& \quad - \alpha) \int_I |f(x)||g(x)|\phi(x) dx \\
& \geq \alpha^2 \int_I f(x)^2 \phi(x) dx + (1 - \alpha)^2 \int_I g(x)^2 \phi(x) dx - 2\alpha(1 \\
& \quad - \alpha) \left(\int_I f(x)^2 \phi(x) dx \right)^{1/2} \left(\int_I g(x)^2 \phi(x) dx \right)^{1/2}
\end{aligned}$$

■

Proposition 7.

With $a, b \in \mathbb{R}$ and $\delta > 0$, let $f : [a - \delta, b + \delta] \rightarrow \mathbb{R}^+$ having only one extremum on $[a - \delta, b + \delta]$ and such that this extremum is a maximum, reached for c . If :

$$|c - b| \geq D \text{ and } |c - a| \geq D$$

then $S(\delta_1, \delta_2) \triangleq \sum_{n=a+\delta_1}^{b+\delta_2} f(n)$ verifies, for $|\delta_1|, |\delta_2| < \delta : \frac{S(\delta_1, \delta_2)}{S(0)} \leq 1 + \frac{\delta}{D}$.

■ We can suppose that a, b, c are integers, which does not change the above argument. The worst case to upper bound $S(\delta_1, \delta_2)$ is $\delta_1 \leq 0, \delta_2 \geq 0$:

$$\begin{aligned}
S(\delta_1, \delta_2) & \leq S(0) + \sum_{n=a-|\delta_1|}^a f(n) + \sum_{n=b}^{b+|\delta_2|} f(n) \leq S(0) + |\delta_1|f(a) + |\delta_2|f(b) \\
& \leq S(0) + |\delta_1| \frac{\sum_{n=a}^c f(n)}{D} + |\delta_2| \frac{\sum_{n=c}^b f(n)}{D} \\
& \leq \left(1 + \frac{|\delta_1|}{D}\right) \sum_{n=a}^c f(n) + \left(1 + \frac{|\delta_2|}{D}\right) \sum_{n=c}^b f(n) \leq S(0) \left(1 + \frac{\max(|\delta_1|, |\delta_2|)}{D}\right) \\
& \leq S(0) \left(1 + \frac{\delta}{D}\right)
\end{aligned}$$

■

The following general results on vectors of positive real numbers will provide useful bounds related to distribution's quadratic matrix in Section III.

Proposition 8.

Let x be a vector in \mathbb{R}^{+n} , we have :

- (i) $\sum_{s=1}^n x_s^2 \geq \sum_{s=1}^n x_s x_{\sigma(s)}, \forall \sigma \in \mathfrak{S}_n$
- (ii) If $x_1 \geq \dots \geq x_n$ then $\sum_{s=1}^n x_s x_{n-s} \leq \sum_{s=1}^n x_s x_{\sigma(s)}, \forall \sigma \in \mathfrak{S}_n$

$$(iii) \quad \forall \sigma \in \mathfrak{S}_n, \quad \frac{1}{2n} \left(\sum_{s=1}^n x_s^2 + \sum_{s=1}^n x_s x_{\sigma(s)} \right) \geq \frac{n}{n-1} \left(\frac{1}{n} \sum_{s=1}^n x_s \right)^2$$

■ (i) : it is sufficient to prove the result for a cycle, as σ can be decomposed in product of disjoint cycles. This can be done easily by recurring over the length of the cycle. For $n = 2$, it is obvious. Suppose it true for $n - 1 \geq 2$, let σ be a cycle of length n : $\sigma = (s_1, \dots, s_n)$, one can also chose without loss of generality that $x_{s_n} = \min_s(x_s)$. Let's set $(y_1, \dots, y_n) = (x_{s_1}, \dots, x_{s_n})$.

$$\begin{aligned} \sum_{s=1}^n x_s x_{\sigma(s)} &= \sum_{s=1}^n y_s y_{s+1} \leq \sum_{s=1}^{n-1} y_s^2 - (y_1(y_{n-1} - y_n) - y_{n-1}y_n) \\ &\leq \sum_{s=1}^n y_s^2 - (y_1(y_{n-1} - y_n) + y_n^2 - y_{n-1}y_n) \leq \sum_{s=1}^n y_s^2 = \sum_{s=1}^n x_s^2 \end{aligned}$$

(ii) : let's denote $S_\sigma = \sum_{s=1}^n x_s x_{\sigma(s)}$, if $\rho = \sigma \circ \tau_{r,s}$, then $S_\sigma - S_\rho = (x_r - x_s)(x_{\sigma(r)} - x_{\sigma(s)})$. The only permutation σ such that there exists no couple (r, s) such that $r < s$ and $\sigma(r) < \sigma(s)$, is $\sigma = (n, n - 1, \dots, 1)$, and thus :

$$\sum_{s=1}^n x_s x_{n-s} \leq \sum_{s=1}^n x_s x_{\sigma(s)}$$

(iii) : we can suppose without loss of generality that $x_1 \geq \dots \geq x_n$, and, also to ease calculation that n is even. Because of (ii) we have :

$$\sum_{s=1}^n x_s x_{n-s} \leq \frac{1}{n!} \sum_{\sigma \in \mathfrak{S}_n} \left(\sum_{s=1}^n x_{\sigma(s)} x_{\sigma(n-s)} \right) = \sum_{s=1}^n \left(\frac{1}{n!} \sum_{\sigma \in \mathfrak{S}_n} x_{\sigma(s)} x_{\sigma(n-s)} \right)$$

n being even, $s \neq n - s$, we have :

$$\frac{1}{n!} \sum_{\sigma \in \mathfrak{S}_n} x_{\sigma(s)} x_{\sigma(n-s)} = \frac{1}{n(n-1)} \sum_{\substack{r,s \\ r \neq s}} x_r x_s = \frac{1}{n(n-1)} \left(\left(\sum_{s=1}^n x_s \right)^2 - \sum_{s=1}^n x_s^2 \right)$$

By injecting the above in $\frac{1}{2n} \left(\sum_{s=1}^n x_s^2 + \sum_{s=1}^n x_s x_{n-s} \right)$, and using (i), we obtain :

$$\frac{1}{2n} \left(\sum_{s=1}^n x_s^2 + \sum_{s=1}^n x_s x_{\sigma(s)} \right) \geq \frac{1}{2n} \left(\sum_{s=1}^n x_s^2 + \sum_{s=1}^n x_s x_{n-s} \right) \geq \frac{n}{n-1} \left(\frac{1}{n} \sum_{s=1}^n x_s \right)^2$$

■

The quadratic matrix M_Φ introduced in Section I is a symmetric matrix with elements in $[0,1]$. We also notice that it is positive (and even positive-definite if Φ is not a Dirac distribution).

$$\sum_{u,v \in \{1, \dots, n\}} M_\Phi(u, v) X_u X_v = \int_{[0,1]^n} \left(\sum_{r \in \{1, \dots, n\}} x_r X_r \right)^2 \Phi(x) dx \geq 0$$

For such matrix, we define the permutation as, $\forall \sigma \in \mathfrak{S}_n$:

$$\sigma(M_\Phi) = M_{\Phi \circ \sigma} = \{M_\Phi(\sigma(u), \sigma(v))\}_{u,v \in \{1, \dots, n\}}$$

and the scalar product, in the vector space of $n \times n$ matrix:

$$M_\Phi \cdot M_{\Phi'} = \text{Tr}(M_\Phi M_{\Phi'}) = \sum_{u,v \in \{1, \dots, n\}} M_\Phi(u, v) M_{\Phi'}(u, v)$$

Let's consider two probability distributions Φ and Φ' ; we introduce the Permutative Gap between Φ and Φ' defined by:

$$\langle \Phi, \Phi' \rangle = \max_{\sigma \in \mathfrak{S}_n} |M_\Phi \cdot M_{\Phi'} - M_\Phi \cdot \sigma(M_{\Phi'})|$$

In the following, we determine a criteria, that Φ and Φ' only need to match separately, to lower bound their Permutative Gap.

We will also manipulate in the following the below operator for 2 distributions Φ, Φ' :

$$\Delta_0(\Phi, \Phi') \triangleq \int_{x,y \in [0,1]^n} \left(\frac{|x||y|}{n^2} - \frac{x \cdot y}{n} \right)^2 \Phi(x) \Phi'(y) dx dy$$

The above definition is introduced because of the close link existing between $\Delta_0(\Phi, \Phi')$ and $\langle \Phi, \Phi' \rangle$.

This link is shown in Proposition 9 below.

A consequence of Proposition 9 is that it is possible to synchronize two probability distributions chosen in $\zeta(\alpha)$ only by composing one of them by a well chosen permutation.

Proposition 9.

If Φ and $\Phi' \in \zeta(\alpha)$, then : $\max_{\sigma \in \mathfrak{S}_n} \Delta_0(\Phi, \Phi' \circ \sigma) \geq \left(\frac{\langle \Phi, \Phi' \rangle}{4n^2} \right)^2 \geq \left(\frac{\alpha}{4} - \left\langle \frac{1}{n} \right\rangle \right)^2$. Φ and $\Phi' \circ \sigma$ are then said 'synchronized'.

■ Let's simply write :

$$\begin{aligned} \frac{\langle \Phi, \Phi' \rangle}{n^2} &= \left| \int_{x,y \in [0,1]^n} \left(\left(\frac{x \cdot \sigma_{max}(y)}{n} \right)^2 - \left(\frac{x \cdot \sigma_{min}(y)}{n} \right)^2 \right) \Phi(x) \Phi'(y) dx dy \right| \\ &\leq 2 \left(\int_{x,y \in [0,1]^n} \left(\frac{x \cdot \sigma_{max}(y)}{n} - \frac{x \cdot \sigma_{min}(y)}{n} \right)^2 \Phi(x) \Phi'(y) dx dy \right)^{1/2} \\ &\leq 2 \left(2(\Delta_0(\Phi, \Phi' \circ \sigma_{max}) + \Delta_0(\Phi, \Phi' \circ \sigma_{min})) \right)^{1/2} \\ &\leq 4 \left(\frac{\Delta_0(\Phi, \Phi' \circ \sigma_{max}) + \Delta_0(\Phi, \Phi' \circ \sigma_{min})}{2} \right)^{1/2} \end{aligned}$$

Thus :

$$\frac{\Delta_0(\Phi, \Phi' \circ \sigma_{max}) + \Delta_0(\Phi, \Phi' \circ \sigma_{min})}{2} \geq \left(\frac{\langle \Phi, \Phi' \rangle}{4n^2} \right)^2$$

which implies, together with Lemma 1 (see hereafter), that there exists $\sigma \in \mathfrak{S}_n$, such that :

$$\Delta_0(\Phi, \Phi' \circ \sigma) \geq \left(\frac{\langle \Phi, \Phi' \rangle}{4n^2} \right)^2 \geq \left(\frac{\alpha}{4} - \frac{1}{n} \right)^2$$

■

Proposition 10.

$\| \cdot \|_c$ is a norm on the subspace of the matrix without their main diagonal, for any $n > 4$.

■ Let's justify by showing that $\| M \|_c = 0 \Leftrightarrow M(u, v) = 0 \forall u \neq v$:

\Leftarrow is obvious.

For \Rightarrow , we calculate $\frac{n^4}{16} \sum_{I \in \mathfrak{S}_n} c_I(M)^2$:

First we have :

$$\sum_{\substack{I \in \mathfrak{S}_n \\ I \times \bar{I} \ni (u, v), (u', v')}} 1 = \binom{n-2}{\frac{n}{2}-2}$$

$$\sum_{\substack{I \in \mathfrak{S}_n \\ I \times \bar{I} \ni (u, v), (u, v')}} 1 = \binom{n-2}{\frac{n}{2}-1}$$

$$\sum_{\substack{I \in \mathfrak{S}_n \\ I \times \bar{I} \ni (u, v)}} 1 = \binom{n-1}{\frac{n}{2}-1}$$

$$\begin{aligned}
\frac{n^4}{16} \sum_{I \in \mathcal{S}_n} c_I(M)^2 &= \sum_{I \in \mathcal{S}_n} \sum_{(u,v),(u',v') \in I \times \bar{I}} M(u,v)M(u',v') \\
&= \sum_{\substack{(u,v),(u',v') \\ u \neq v, u \neq u' \\ u' \neq v', v \neq v'}} \left(\sum_{I \in \mathcal{S}_n} \mathbb{1}_{I \times \bar{I} \ni (u,v),(u',v')} \right) M(u,v)M(u',v') \\
&\quad + 2 \sum_{\substack{u,v,v' \\ u \neq v, u \neq v' \\ v \neq v'}} \left(\sum_{I \in \mathcal{S}_n} \mathbb{1}_{I \times \bar{I} \ni (u,v),(u,v')} \right) M(u,v)M(u,v') + \sum_{\substack{u,v \\ u \neq v}} \left(\sum_{I \in \mathcal{S}_n} \mathbb{1}_{I \times \bar{I} \ni (u,v)} \right) M(u,v)^2 \\
&= \left(\frac{n-2}{\frac{n}{2}-2} \right) \left(\sum_{\substack{u,v \\ u \neq v}} M(u,v) \right)^2 + 2 \left(\left(\frac{n-2}{\frac{n}{2}-1} \right) - \left(\frac{n-2}{\frac{n}{2}-2} \right) \right) \sum_u \left(\sum_{\substack{v \\ v \neq u}} M(u,v) \right)^2 \\
&\quad + \left(\left(\frac{n-1}{\frac{n}{2}-1} \right) + \left(\frac{n-2}{\frac{n}{2}-2} \right) - 2 \left(\frac{n-2}{\frac{n}{2}-1} \right) \right) \sum_{\substack{u,v \\ u \neq v}} M(u,v)^2
\end{aligned}$$

We conclude by noticing that :

$$\left(\frac{n-2}{\frac{n}{2}-1} \right) - \left(\frac{n-2}{\frac{n}{2}-2} \right) > 0 \text{ and } \left(\frac{n-1}{\frac{n}{2}-1} \right) + \left(\frac{n-2}{\frac{n}{2}-2} \right) - 2 \left(\frac{n-2}{\frac{n}{2}-1} \right) = \left(\frac{n-2}{\frac{n}{2}-1} \right) \frac{n-4}{n} > 0 \text{ for } n > 4. \quad \blacksquare$$

Proposition 11.

With the notations of Section II, we have : $\inf_{\omega} E[(\omega_{i,j} - V_B)^2] \leq E[(V_A - V_B)^2]$.

■ we write $E[(V_A - V_B)^2]$ and simply apply Schwarz inequality :

$$\begin{aligned}
E[(V_A - V_B)^2] &= \int_{x,y \in [0,1]^n} \sum_{i,j \in \{0,1\}^n} \left(\frac{x \cdot j}{n} - \frac{i \cdot y}{n} \right)^2 \chi_i \left(\frac{x}{k} \right) \chi_j \left(\frac{y}{k} \right) \Phi(x) \Phi'(y) dx dy \\
&= \int_y \sum_{i,j} \left[\int_x \left(\frac{x \cdot j}{n} - \frac{i \cdot y}{n} \right)^2 \chi_i \left(\frac{x}{k} \right) \Phi(x) dx \right] \chi_j \left(\frac{y}{k} \right) \Phi'(y) dy \\
&\geq \int_y \sum_{i,j} \left(\int_x \chi_i \left(\frac{x}{k} \right) \Phi(x) dx \right) \left(\frac{\int_x \frac{x \cdot j}{n} \chi_i \left(\frac{x}{k} \right) \Phi(x) dx}{\int_x \chi_i \left(\frac{x}{k} \right) \Phi(x) dx} - \frac{i \cdot y}{n} \right)^2 \chi_j \left(\frac{y}{k} \right) \Phi'(y) dy
\end{aligned}$$

By remarking that $\frac{\int_x \frac{x \cdot j}{n} \chi_i \left(\frac{x}{k} \right) \Phi(x) dx}{\int_x \chi_i \left(\frac{x}{k} \right) \Phi(x) dx}$ is of the form $\omega(i,j)$ the result follows. ■

A classical result that we give here is that a broadcast BSC channel with 1 sender and 2 receivers:




Fig 1.

can always be modeled by a broadcast channel with independent BSC channels as shown below:




Fig. 2

We denote:

$$\varphi(x, y) \triangleq x(1 - y) + y(1 - x) = x + y - 2xy$$

$$\phi(x, y, z) \triangleq \begin{pmatrix} \varphi(x, y) \\ \varphi(x, z) \\ \varphi(y, z) \end{pmatrix}$$

and:

$$\mathcal{F} \triangleq \left\{ X, Y, Z \in [0, 1/2[^3 \left| \begin{array}{l} (1 - 2X)(1 - 2Y) \leq (1 - 2Z) \\ (1 - 2X)(1 - 2Z) \leq (1 - 2Y) \\ (1 - 2Y)(1 - 2Z) \leq (1 - 2X) \end{array} \right. \right\}$$

Proposition 12.

A broadcast BSC channel $BSC(\varepsilon_B, \varepsilon_E)$ with 1 sender A and 2 receivers B and E, having respective error variables ε_B and ε_E , (i) can be modeled as a cascaded broadcast channel $CBSC(x, y, z)$ with 3 independent BSC, one common attached to the sender, with error probability x , and two cascaded with the sender's one, each attached to one of the receivers, with respective error probability y and z . (ii) If $(P(\varepsilon_B = 1), P(\varepsilon_E = 1), P(\varepsilon_B \neq \varepsilon_E)) \in \mathcal{F}$ then the solution is unique and also belongs to $[0, 1/2[^3$. (iii) The modelization transform is a diffeomorphism from \mathcal{F} in $\phi^{-1}(\mathcal{F}) \in [0, 1/2[^3$. (iv) if there exists $\alpha < 1/2$, such that $P(\varepsilon_B = 1) \leq \alpha$, $P(\varepsilon_E = 1) \leq \alpha$ and $P(\varepsilon_B \neq \varepsilon_E) \leq \alpha$, then there exists $\beta > 0$ such that $|J_\phi| \leq \beta$ and $|J_{\phi^{-1}}| \leq \beta$

■ the probability distribution governing initial broadcast as shown in Fig. 1 is fully determined by:

$$X = P(\varepsilon_B = 1)$$

$$Y = P(\varepsilon_E = 1)$$

$$Z = P(\varepsilon_B \neq \varepsilon_E)$$

Indeed the distribution $P_{\varepsilon_B, \varepsilon_E}$ is fully defined by the relations below:

$$\begin{pmatrix} Y & -X \\ Y & X \end{pmatrix} \begin{pmatrix} P(\varepsilon_B = 0 | \varepsilon_E = 1) \\ P(\varepsilon_E = 0 | \varepsilon_B = 1) \end{pmatrix} = \begin{pmatrix} 0 \\ Z \end{pmatrix}$$

$$\begin{pmatrix} 1 - Y & -1 + X \\ 1 - Y & 1 - X \end{pmatrix} \begin{pmatrix} P(\varepsilon_B = 1 | \varepsilon_E = 0) \\ P(\varepsilon_E = 1 | \varepsilon_B = 0) \end{pmatrix} = \begin{pmatrix} 0 \\ Z \end{pmatrix}$$

If (x, y, z) denotes the error probabilities of the 3 independent BSC in the model shown in Fig. 2, then in such model, (X, Y, Z) is given by:

$$\begin{pmatrix} X \\ Y \\ Z \end{pmatrix} = \phi(x, y, z)$$

We first observe that we can exclude $x = 1/2$, $y = 1/2$, and $z = 1/2$ because it would then lead one of the X, Y, Z to be equal to $1/2$. We can then apply the following reverse formula:

$$x = \frac{X - y}{1 - 2y}$$

$$z = \frac{Z - y}{1 - 2y}$$

$$Y = \frac{X - y}{1 - 2y} + \frac{Z - y}{1 - 2y} - 2 \frac{X - y}{1 - 2y} \frac{Z - y}{1 - 2y}$$

Which gives a unique solution in $[0, 1/2[$ for y in terms of X, Y, Z (and similarly also x and z by circular permutation) as:

$$y = \frac{1 - \sqrt{\Delta(X, Y, Z)}}{2}$$

$$\Delta(X, Y, Z) = \frac{(1 - 2X)(1 - 2Z)}{1 - 2Y} > 0$$

As $(X, Y, Z) \in \mathcal{F}$, it is verified also that $\Delta(X, Y, Z) \leq 1$, a necessary and sufficient condition to have $y \geq 0$. We can compute the Jacobian determinant of ϕ as being:

$$J_\phi(x, y, z) = 2(1 - 2x)(1 - 2y)(1 - 2z) > 0$$

The inversion formula above ensures injectivity of ϕ^{-1} from $[0, 1/2]^3$ in \mathcal{F} , together with the fact that Jacobian is never zero provide that ϕ^{-1} is a diffeomorphism from $[0, 1/2]^3$ in $\phi^{-1}(\mathcal{F}) \in [0, 1/2]^3$. Assertion (iv) comes with the fact that $1 - 2X, 1 - 2Y, 1 - 2Z$ can be lower-bounded when $\alpha < 1/2$.

■

We give below the justification that, in a $BSC(\varepsilon_B, \varepsilon_E)$ broadcast model, the assumption that $(X, Y, Z) \in \mathcal{F}$ is legitimate. We denote by e_A, e_B, e_E the versions of respectively A, B, E for the considered bit, then $\varepsilon_B = e_A \oplus e_B$, $\varepsilon_E = e_A \oplus e_E$, and $\varepsilon_B \oplus \varepsilon_E = e_B \oplus e_E$.

- 1) If the 3 components X, Y, Z are $> 1/2$, then for instance B inverse e_B and then we will land to case 2 where only one component is $> 1/2$.
- 2) If only one component is $> 1/2$: if it is Y then it means that we already have an advantage for the legitimate partners A and B ; if it is X , A can reverse its bit e_A and it becomes Y the only component $> 1/2$ which again leads to Advantage for the partners; if it is Z , both A and B reversing their bits e_A and e_B , and again it becomes Y the only component $> 1/2$ which again leads to Advantage for the partners.
- 3) If two components over three are $> 1/2$: if it is X , A can reverse its bit e_A and the three components become $< 1/2$; if it is Z , B can reverse its bit e_B and the three components become again $< 1/2$; if it is Y both A and B reversing their bits e_A and e_B , and again the three components become $< 1/2$. Items 1, 2 and 3 prove that one can restrict $(X, Y, Z) \in [0, 1/2]^3$.
- 4) A and B can alternatively change their role, and thus in average, $X = Z$. With that equality, it is clear that the two conditions of \mathcal{F}

$$(1 - 2Y)(1 - 2Z) \leq (1 - 2X)$$

$$(1 - 2X)(1 - 2Y) \leq (1 - 2Z)$$

are satisfied

- 5) Finally, if the third condition of \mathcal{F} was not satisfied, e.g. $(1 - 2X)(1 - 2Z) > (1 - 2Y)$, then this would lead to an Advantage for the partners because

$$\min((1 - 2X), (1 - 2Z)) \geq (1 - 2X)(1 - 2Z)$$

and thus an Advantage would exist for both e_A and e_B over e_E ($Y > X$ and $Y > Z$).

We show below that the repetition error correcting code used at step 5^o of the protocol (and proposed in [5]) produces Advantage Distillation in a $CBSC(x, y, z)$ model. We consider the specific case where $x = y$ that is sufficient for our purpose.

Proposition 13.

In a broadcast BSC channel $BSC(\varepsilon_B, \varepsilon_E)$, modeled as a $CBSC(x, y, z)$ model, where we apply a repetition error correcting code of length L (an even integer), we have (with notation of Proposition 12):

- (i) *The error rate ratio is given by:*

$$\frac{P_{err}(B)}{P_{err}(E)} = \frac{\varphi(x, y)^L}{\sum_{l>\frac{L}{2}} \binom{L}{l} p_l(x, y, z)}$$

- (ii) *In the case where $x = y$, the ratio is upper-bounded by:*

$$\frac{P_{err}(B)}{P_{err}(E)} \leq \frac{x^2 + (1 - z)(1 - 2x)}{x^2 + z(1 - 2x)} \frac{C\sqrt{L}}{\left(1 + z(1 - z) \frac{(1 - 2x)^2}{x^2(1 - x)^2}\right)^{\frac{L}{2}}}$$

Where

$$p_l(x, y, z) = (x(1 - y) + z(y - x))^l (y(1 - x) - z(y - x))^{L-l} + (xy + z(1 - y - x))^l ((1 - y)(1 - x) - z(1 - y - x))^{L-l}$$

■ (i) if we denote P_{acc} the probability that B accepts the code word sent by A , and $(\delta_A, \delta_B, \delta_E)$ the respective binary independent error variables then

$$\begin{aligned}
P_{err}(B) &= P((\delta_A \oplus \delta_B)_L = (1)_L | B \text{ accepts}) = \frac{P((\delta_A \oplus \delta_B)_L = (1)_L)}{P_{acc}} = \frac{\varphi(x, y)^L}{P_{acc}} \\
P_{err}(E) &= P\left(|(\delta_A \oplus \delta_B)_L| > \frac{L}{2} | B \text{ accepts}\right) \\
&= \frac{P\left(\left(|(\delta_A \oplus \delta_B)_L| > \frac{L}{2}\right) \& \left(|(\delta_A \oplus \delta_B)_L| = L\right)\right) \parallel \left(\left(|(\delta_A \oplus \delta_B)_L| > \frac{L}{2}\right) \& \left(|(\delta_A \oplus \delta_B)_L| = 0\right)\right)}{P_{acc}} \\
&= \frac{1}{P_{acc}} \sum_{|I| > \frac{L}{2}} \left(P((\delta_A \neq \delta_E)_I) P((\delta_A = \delta_E)_{\bar{I}}) P((\delta_A \neq \delta_B)_I) P((\delta_A = \delta_B)_{\bar{I}}) \right) \\
&\quad + P((\delta_A \neq \delta_E)_I) P((\delta_A = \delta_E)_{\bar{I}}) P((\delta_A = \delta_B)_I) P((\delta_A = \delta_B)_{\bar{I}}) = \frac{1}{P_{acc}} \sum_{|I| > \frac{L}{2}} \binom{L}{I} p_I(x, y, z)
\end{aligned}$$

(ii) we can lower bound $p_I(x, y, z)$ by:

$$\begin{aligned}
p_I(x, y, z) &\geq (xy + z(1 - y - x))^{\frac{L}{2}+1} ((1 - y)(1 - x) - z(1 - y - x))^{\frac{L}{2}-1} \\
&= \frac{xy + z(1 - y - x)}{(1 - y)(1 - x) - z(1 - y - x)} (x(1 - y))^{\frac{L}{2}} (y(1 - x))^{\frac{L}{2}} \left(1 + z \frac{1 - x - y}{xy}\right)^{\frac{L}{2}} \left(1 - z \frac{1 - x - y}{(1 - x)(1 - y)}\right)^{\frac{L}{2}}
\end{aligned}$$

With $x = y$, that lower bound becomes

$$\binom{L}{L/2} p_{L/2}(x, x, z) \geq \frac{1}{2^n} \binom{L}{L/2} \varphi(x, x)^L \frac{x^2 + z(1 - 2x)}{x^2 + (1 - z)(1 - 2x)} \left(1 + z(1 - z) \frac{(1 - 2x)^2}{x^2(1 - x)^2}\right)^{\frac{L}{2}}$$

As it is well known that for a certain constant C

$$\frac{1}{2^n} \binom{L}{L/2} \geq \frac{1}{C\sqrt{L}}$$

we get the result. ■

The following proposition justifies the form of the canonical strategy in the case where the distributions $\Sigma(\Phi, I)$ and $\Sigma(\Phi', J)$ are symmetric within 2 respective index subsets (I, \bar{I}) and (J, \bar{J}) . We denote \mathfrak{S}_I is the sub-group of \mathfrak{S}_n : $\mathfrak{S}_I \triangleq \{\sigma \in \mathfrak{S}_n | \forall u \in I, \sigma(u) \in I\}$. we define by $|I|$ the cardinality of I . The notations $\Sigma(\Phi, I)$ and $\Sigma(\Phi', J)$ are justified by the fact that such distributions symmetric within the respective index subsets (I, \bar{I}) and (J, \bar{J}) can be generated from any pair of distributions Φ and Φ' by performing the following process: In the choice of Φ by A 's Deep Random Generator, all happen as if A would perform the sequence below:

$$\Phi \circ \sigma_\Phi: x'' \xrightarrow{\sigma \in \mathfrak{S}_{I_0}} x' \xrightarrow{\sigma_I} x$$

At the first step x'' is generated by $\Phi \circ \sigma_\Phi$ (synchronized on I_0), at second step $x' = \sigma^{-1}(x'')$ corresponds to a mixing within I_0 , and at third step, σ_I , designating any permutation sending I_0 in I , produces the final x . The resulting distribution to be considered is denoted:

$$\Sigma(\Phi, I) \triangleq \frac{1}{((n/2)!)^2} \sum_{\sigma \in \mathfrak{S}_{I_0}} \Phi \circ \sigma_\Phi \circ \sigma \circ \sigma_I^{-1}$$

The notation $\Sigma(\Phi, I)$ is justified by the fact that the above sum does not depend on which σ_I is chosen. In case where $I = \sigma_\Phi(I_0)$, we simply denote:

$$\Sigma(\Phi) \triangleq \Sigma(\Phi, \sigma_\Phi(I_0))$$

In the following, $x|_I$ designates the restriction of the vector x to its components having their index within I . We denote, for i and j being $\{0,1\}$ vectors in I :

$$f_I(i, j) \triangleq \left(\frac{|I| - |i| - |j| + i \cdot j}{(|I| - |i|)(|I| - |j|)} (k-1)^2 |i||j| + \frac{|i| - i \cdot j}{(|I| - |j|)} (k-1)|j| + \frac{|j| - i \cdot j}{(|I| - |i|)} (k-1)|i| + i \cdot j \right)$$

Proposition 14.

If the distributions Φ and Φ' are symmetric within 2 respective index subsets (I, \bar{I}) and (J, \bar{J}) , if they are transformed by a sufficiently regular permutative sleeking transform (as also assumed in further lemma 3), and if we assume $|i|, |j| \in [1, n/k]$, we have the equalities below. Under Deep Random assumption, if Φ (resp. Φ') is undistinguishable from any member of $R_{\mathfrak{S}}(\mathfrak{S}_{\sigma_\Phi(I_0)})$ (resp. $R_{\mathfrak{S}}(\mathfrak{S}_{\sigma_{\Phi'}(I_0)})$) by the opponent, the equalities below should be assume for the opponent

$$\begin{aligned} E \left[\frac{\sigma_\Phi^{-1}(x) \cdot \sigma_{\Phi'}^{-1}(y)}{nk} \mid i, j \right] &= E \left[\frac{\sigma_\Phi^{-1}(x) \cdot \sigma_{\Phi'}^{-1}(y)}{nk} \mid i, j \right]_{\Sigma(\Phi), \Sigma(\Phi')} \\ &= \frac{1}{nk} \left(f_{I_0}(\sigma_\Phi^{-1}(i)|_{I_0}, \sigma_{\Phi'}^{-1}(j)|_{I_0}) + f_{\bar{I}_0}(\sigma_\Phi^{-1}(i)|_{\bar{I}_0}, \sigma_{\Phi'}^{-1}(j)|_{\bar{I}_0}) \right) + O \left(\sqrt{\frac{\ln n}{nk}} \right) \\ &= \frac{2k}{n^2} \left((\sigma_\Phi^{-1}(i) \cdot I_0)(\sigma_{\Phi'}^{-1}(j) \cdot I_0) + (\sigma_\Phi^{-1}(i) \cdot \bar{I}_0)(\sigma_{\Phi'}^{-1}(j) \cdot \bar{I}_0) \right) + O \left(\frac{1}{k^2} \right) \end{aligned}$$

■ The first equality comes from the Deep Random Assumption. It is sufficient to prove the second and third equality. From Bayes:

$$\begin{aligned} E \left[\frac{\sigma_\Phi^{-1}(x) \cdot \sigma_{\Phi'}^{-1}(y)}{nk} \mid i, j \right]_{\Sigma(\Phi), \Sigma(\Phi')} \\ &= \frac{1}{nk} \frac{\sum_{\sigma, \mu \in \mathfrak{S}_{I_0}} \int_{x, y \in \{0,1\}^n} x \cdot y \chi_{\sigma_\Phi^{-1}(i)} \left(\frac{x}{k} \right) \chi_{\sigma_{\Phi'}^{-1}(j)} \left(\frac{y}{k} \right) \Phi \circ \sigma_\Phi \circ \sigma(x) \Phi' \circ \sigma_{\Phi'} \circ \mu(y) dx dy}{\sum_{\sigma, \mu \in \mathfrak{S}_{I_0}} \int_{x, y \in \{0,1\}^n} \chi_{\sigma_\Phi^{-1}(i)} \left(\frac{x}{k} \right) \chi_{\sigma_{\Phi'}^{-1}(j)} \left(\frac{y}{k} \right) \Phi \circ \sigma_\Phi \circ \sigma(x) \Phi' \circ \sigma_{\Phi'} \circ \mu(y) dx dy} \end{aligned}$$

In all the following, in order to simplify notations, we will generically compute

$$E \left[\frac{x \cdot y}{nk} \mid i, j \right]_{\Sigma(\Phi), \Sigma(\Phi')}$$

with assumption that $\sigma_\Phi = \sigma_{\Phi'} = \text{Id}_{\mathfrak{S}_n}$. The above equality shows how to transpose the generic computation to the desired result. We denote the restricted distributions:

$$\begin{aligned}\Sigma_i(\Phi)(x|_{I_0}) &\triangleq \left(\int_{x|_{\bar{I}_0}} \chi_{i|\bar{I}_0} \left(\frac{x|_{\bar{I}_0}}{k} \right) \Sigma(\Phi)(x) dx |_{\bar{I}_0} \right) / \left(\int_x \chi_{i|\bar{I}_0} \left(\frac{x|_{\bar{I}_0}}{k} \right) \Sigma(\Phi)(x) dx \right) \\ \overline{\Sigma_i(\Phi)}(x|_{\bar{I}_0}) &\triangleq \left(\int_{x|_{I_0}} \chi_{i|I_0} \left(\frac{x|_{I_0}}{k} \right) \Sigma(\Phi)(x) dx |_{I_0} \right) / \left(\int_x \chi_{i|I_0} \left(\frac{x|_{I_0}}{k} \right) \Sigma(\Phi)(x) dx \right)\end{aligned}$$

We see that $\Sigma_i(\Phi)(x|_{I_0})$ and $\overline{\Sigma_i(\Phi)}(x|_{\bar{I}_0})$ are distributions that are symmetric over respectively $x|_{I_0}$ and $x|_{\bar{I}_0}$, and their dependency on i are respectively restricted to only $i|_{\bar{I}_0}$ and $i|_{I_0}$. By splitting the sum of the scalar product $x \cdot y$ in $\sum_{t \in I_0} x_t y_t + \sum_{t \in \bar{I}_0} x_t y_t$, we obtain:

$$\begin{aligned}E \left[\frac{x \cdot y}{nk} | i, j \right]_{\Sigma(\Phi), \Sigma(\Phi')} &= \frac{1}{nk} \frac{\int_{x|_{I_0}, y|_{I_0} \in [0,1]^n x|_{I_0} \cdot y|_{I_0} \chi_{i|I_0} \left(\frac{x|_{I_0}}{k} \right) \chi_{j|I_0} \left(\frac{y|_{I_0}}{k} \right) \Sigma_i(\Phi)(x|_{I_0}) \Sigma_j(\Phi')(y|_{I_0}) dx|_{I_0} dy|_{I_0}}{\int_{x|_{I_0}, y|_{I_0} \in [0,1]^n \chi_{i|I_0} \left(\frac{x|_{I_0}}{k} \right) \chi_{j|I_0} \left(\frac{y|_{I_0}}{k} \right) \Sigma_i(\Phi)(x|_{I_0}) \Sigma_j(\Phi')(y|_{I_0}) dx|_{I_0} dy|_{I_0}} \\ &+ \frac{1}{nk} \frac{\int_{x|_{\bar{I}_0}, y|_{\bar{I}_0} \in [0,1]^n x|_{\bar{I}_0} \cdot y|_{\bar{I}_0} \chi_{i|\bar{I}_0} \left(\frac{x|_{\bar{I}_0}}{k} \right) \chi_{j|\bar{I}_0} \left(\frac{y|_{\bar{I}_0}}{k} \right) \overline{\Sigma_i(\Phi)}(x|_{\bar{I}_0}) \overline{\Sigma_j(\Phi')}(y|_{\bar{I}_0}) dx|_{\bar{I}_0} dy|_{\bar{I}_0}}{\int_{x|_{\bar{I}_0}, y|_{\bar{I}_0} \in [0,1]^n \chi_{i|\bar{I}_0} \left(\frac{x|_{\bar{I}_0}}{k} \right) \chi_{j|\bar{I}_0} \left(\frac{y|_{\bar{I}_0}}{k} \right) \overline{\Sigma_i(\Phi)}(x|_{\bar{I}_0}) \overline{\Sigma_j(\Phi')}(y|_{\bar{I}_0}) dx|_{\bar{I}_0} dy|_{\bar{I}_0}}\end{aligned} \quad (p14.1)$$

We can restrict ourselves to study generically:

$$\frac{1}{nk} \frac{\int_{x, y \in \{0,1\}^n x \cdot y \chi_i \left(\frac{x}{k} \right) \chi_j \left(\frac{y}{k} \right) \bar{\Phi}(x) \bar{\Phi}'(y) dx dy}{\int_{x, y \in \{0,1\}^n \chi_i \left(\frac{x}{k} \right) \chi_j \left(\frac{y}{k} \right) \bar{\Phi}(x) \bar{\Phi}'(y) dx dy}$$

where $\bar{\Phi}$ and $\bar{\Phi}'$ are symmetric distribution over $\{0,1\}^n$. Let's denote

$$J_1 = \bar{i} \cap \bar{j}, J_2 = i \cap \bar{j}, J_3 = \bar{i} \cap j, J_4 = i \cap j$$

By considering $\mathfrak{S}_{i,j}$ the subgroup of \mathfrak{S}_n whose permutations let J_1, J_2, J_3, J_4 , stable, $\bar{\Phi}$ and $\bar{\Phi}'$ are obviously invariant by action of $\mathfrak{S}_{i,j}$, and thus we can write:

$$\begin{aligned}& \frac{1}{nk} \frac{\int_{x, y \in \{0,1\}^n x \cdot y \chi_i \left(\frac{x}{k} \right) \chi_j \left(\frac{y}{k} \right) \bar{\Phi}(x) \bar{\Phi}'(y) dx dy}{\int_{x, y \in \{0,1\}^n \chi_i \left(\frac{x}{k} \right) \chi_j \left(\frac{y}{k} \right) \bar{\Phi}(x) \bar{\Phi}'(y) dx dy} \\ &= \frac{1}{nk} \sum_{s=1}^4 \sum_{t \in J_s} \left(\frac{\int_{x \in \{0,1\}^n x_t \chi_i \left(\frac{x}{k} \right) \bar{\Phi}(x) dx}{\int_{x \in \{0,1\}^n \chi_i \left(\frac{x}{k} \right) \bar{\Phi}(x) dx} \right) \left(\frac{\int_{y \in \{0,1\}^n y_t \chi_j \left(\frac{y}{k} \right) \bar{\Phi}'(y) dy}{\int_{y \in \{0,1\}^n \chi_j \left(\frac{y}{k} \right) \bar{\Phi}'(y) dy} \right) \\ &= \frac{1}{nk} \sum_{s=1}^4 \frac{1}{|J_s|} \left(\frac{\int_{x \in \{0,1\}^n x \cdot J_s \chi_i \left(\frac{x}{k} \right) \bar{\Phi}(x) dx}{\int_{x \in \{0,1\}^n \chi_i \left(\frac{x}{k} \right) \bar{\Phi}(x) dx} \right) \left(\frac{\int_{y \in \{0,1\}^n y \cdot J_s \chi_j \left(\frac{y}{k} \right) \bar{\Phi}'(y) dy}{\int_{y \in \{0,1\}^n \chi_j \left(\frac{y}{k} \right) \bar{\Phi}'(y) dy} \right)\end{aligned} \quad (p14.2)$$

Let's study the first bracket. For $s \in \{2,4\}$, due to the fact that $x_t \in \{0,1\}$:

$$\frac{\int_{x \in \{0,1\}^n} x \cdot J_s \chi_i \left(\frac{x}{k} \right) \bar{\Phi}(x) dx}{\int_{x \in \{0,1\}^n} \chi_i \left(\frac{x}{k} \right) \bar{\Phi}(x) dx} = |J_s|$$

For $s \in \{1,3\}$:

$$\begin{aligned} \frac{\int_{x \in \{0,1\}^n} x \cdot J_s \chi_i \left(\frac{x}{k} \right) \bar{\Phi}(x) dx}{\int_{x \in \{0,1\}^n} \chi_i \left(\frac{x}{k} \right) \bar{\Phi}(x) dx} &= \frac{|J_s| \int_{x \in \{0,1\}^n} x \cdot \bar{\chi}_i \left(\frac{x}{k} \right) \bar{\Phi}(x) dx}{|i| \int_{x \in \{0,1\}^n} \chi_i \left(\frac{x}{k} \right) \bar{\Phi}(x) dx} \\ &= \frac{|J_s|}{n - |i|} \left(\frac{\int_{x \in \{0,1\}^n} |x| \chi_i \left(\frac{x}{k} \right) \bar{\Phi}(x) dx}{\int_{x \in \{0,1\}^n} \chi_i \left(\frac{x}{k} \right) \bar{\Phi}(x) dx} - |i| \right) \end{aligned}$$

We denote $\forall u \in \mathbb{N}_n$, $P_u \triangleq \int_{|x|=u} \bar{\Phi}(x) dx$. Then let's study:

$$\begin{aligned} \frac{\int_{x \in \{0,1\}^n} |x| \chi_i \left(\frac{x}{k} \right) \bar{\Phi}(x) dx}{\int_{x \in \{0,1\}^n} \chi_i \left(\frac{x}{k} \right) \bar{\Phi}(x) dx} &= \frac{\int_{x \in \{0,1\}^n} |x| \psi_{|i|} \left(\frac{x}{k} \right) \bar{\Phi}(x) dx}{\int_{x \in \{0,1\}^n} \psi_{|i|} \left(\frac{x}{k} \right) \bar{\Phi}(x) dx} \\ &= \frac{\int_{x \in \{0,1\}^n} |x| \sum_{\Delta=(1-k)|i|}^{n/2-k|i|} \beta_{|i|,k|i|+\Delta} \left(\frac{1}{k} \right) \bar{\Phi}(x) dx}{\int_{x \in \{0,1\}^n} \sum_{\Delta=(1-k)|i|}^{n/2-k|i|} \beta_{|i|,k|i|+\Delta} \left(\frac{1}{k} \right) \bar{\Phi}(x) dx} \\ &= \frac{\sum_{\Delta=(1-k)|i|}^{n/2-k|i|} (k|i| + \Delta) \beta_{|i|,k|i|+\Delta} \left(\frac{1}{k} \right) P_{k|i|+\Delta}}{\sum_{\Delta=(1-k)|i|}^{n/2-k|i|} \beta_{|i|,k|i|+\Delta} \left(\frac{1}{k} \right) P_{k|i|+\Delta}} \end{aligned}$$

K being a constant that we will calibrate appropriately. We also denote:

$$\begin{aligned} U_K &\triangleq \{r \in \mathbb{Z} | \Delta^2 \leq 2Kk^2|i|\} \\ \bar{U}_K &\triangleq \{(1-k)|i|, \dots, n/2 - k|i|\} \setminus U_K \end{aligned}$$

Thanks to Proposition 1, we can upper-bound the difference:

$$\begin{aligned} \left| \frac{\int_{x \in \{0,1\}^n} (|x| - k|i|) \chi_i \left(\frac{x}{k} \right) \bar{\Phi}(x) dx}{\int_{x \in \{0,1\}^n} \chi_i \left(\frac{x}{k} \right) \bar{\Phi}(x) dx} \right| &\leq \frac{\sum_{\Delta=(1-k)|i|}^{n/2-k|i|} |\Delta| \beta_{|i|,k|i|+\Delta} \left(\frac{1}{k} \right) P_{k|i|+\Delta}}{\sum_{\Delta=(1-k)|i|}^{n/2-k|i|} \beta_{|i|,k|i|+\Delta} \left(\frac{1}{k} \right) P_{k|i|+\Delta}} \\ &\leq \frac{\sum_{\Delta \in U_K} |\Delta| \beta_{|i|,k|i|+\Delta} \left(\frac{1}{k} \right) P_{k|i|+\Delta} + \sum_{\Delta \in \bar{U}_K} |\Delta| \beta_{|i|,k|i|+\Delta} \left(\frac{1}{k} \right) P_{k|i|+\Delta}}{\sum_{\Delta \in U_K} \beta_{|i|,k|i|+\Delta} \left(\frac{1}{k} \right) P_{k|i|+\Delta}} \\ &\leq \frac{\sqrt{2Kk^2|i|} \sum_{\Delta \in U_K} \beta_{|i|,k|i|+\Delta} \left(\frac{1}{k} \right) P_{k|i|+\Delta} + n^2 e^{-K}}{\sum_{\Delta \in U_K} \beta_{|i|,k|i|+\Delta} \left(\frac{1}{k} \right) P_{k|i|+\Delta}} \\ &\leq 2\sqrt{2Kk^2|i|} \end{aligned} \tag{p14.3}$$

if K is calibrated such that:

$$\frac{n^2}{2} e^{-K} \leq \sqrt{2Kk^2|i|} \sum_{\Delta \in U_K} \beta_{|i|,k|i|+\Delta} \left(\frac{1}{k}\right) P_{k|i|+\Delta} \quad (C)$$

This is always possible as soon as $|i| \geq 1$; and when Φ (resp. Φ') are transformed by a sufficiently regular permutative sleeking transform, there exist a constant γ such that:

$$P_{k|i|} \geq \frac{\gamma}{n}$$

Therefore, there exist a constant $C_0 > 0$ such that

$$\sum_{\Delta \in U_K} \beta_{|i|,k|i|+\Delta} \left(\frac{1}{k}\right) P_{k|i|+\Delta} \geq \beta_{|i|,k|i|} \left(\frac{1}{k}\right) P_{k|i|} \geq \frac{C_0}{n\sqrt{n}}$$

and thus (C) implies that $K = O(\ln n)$.

By injecting the upper-bounds obtained at (p14.3) in (p14.2) and then the result in (p14.1), we finally get the expected result. ■

It is not difficult to use the same method to obtain the same equalities with $E \left[\frac{\sigma_{\Phi}^{-1}(x) \cdot \sigma_{\Phi'}^{-1}(j)}{n} | i, j \right]_{\Sigma(\Phi), \Sigma(\Phi')}$ and $E \left[\frac{\sigma_{\Phi}^{-1}(i) \cdot \sigma_{\Phi'}^{-1}(y)}{n} | i, j \right]_{\Sigma(\Phi), \Sigma(\Phi')}$. Therefore the above proposition, combined with the following one, will be used in the proof of main Theorem to exhibit an approximated modelization of the protocol \mathcal{P} as a cascaded broadcast independent channel $CBSC(x, y, z)$.

The following propositions is useful to establish that the sampling method used in protocol \mathcal{P} introduced in Section 0 enable to modelize \mathcal{P} as cascaded broadcast channel $CBSC(x, y, z)$ with 3 independent BSC. We denote Γ the sampling function:

$$\Gamma_{\theta}(x) \triangleq \left\lfloor \frac{x}{\theta} \right\rfloor \bmod 2$$

where θ is a real number such that $0 < \theta \ll 1$.

Proposition 15.

Let U and V be 2 independent random variables with values in $[0,1]$. Let also ρ be a random variable independent of U and V , and equi-distributed over $[0,2\theta[$. Then the variable $e_V = \Gamma_{\theta}(U + V + \rho) \oplus \Gamma_{\theta}(U + \rho)$ is independent of the variable $e_U = \Gamma_{\theta}(U + \rho)$.

■ Let's denote $f(u)$ and $g(v)$ the respective distributions of U and V over $[0,1]$. We also denote the set:

$$\Gamma_{\theta}^+ \triangleq \bigcup_{p \in \mathbb{Z}} [(2p - 1)\theta, 2p\theta[$$

$$\Gamma_{\theta}^- \triangleq \bigcup_{p \in \mathbb{Z}} [2p\theta, (2p + 1)\theta[$$

In order to establish independence between e_U and e_V , we can write $P((e_U = 1) \& (e_V = 1))$ and $P(e_V = 1)$:

$$P((e_U = 1) \& (e_V = 1)) = \int_{\rho=0}^{2\theta} \int_{u,v=0}^1 P((u + \rho \in \Gamma_{\theta}^+) \& (u + v + \rho \in \Gamma_{\theta}^+)) f(u)g(v)dudvd\rho$$

$$P(e_V = 1) = \int_{\rho=0}^{2\theta} \int_{u,v=0}^1 P((u + \rho \in \Gamma_{\theta}^+) \& (u + v + \rho \in \Gamma_{\theta}^+)) f(u)g(v)dudvd\rho$$

$$+ \int_{\rho=0}^{2\theta} \int_{u,v=0}^1 P((u + \rho \in \Gamma_{\theta}^-) \& (u + v + \rho \in \Gamma_{\theta}^-)) f(u)g(v)dudvd\rho$$

By doing the change of variable $\rho' = (\rho - \theta) \bmod 2\theta$ on the second term of the right side of the above equality, we get

$$P(e_V = 1) = 2P((e_U = 1) \& (e_V = 1))$$

And of course we have $\frac{1}{2} = P(e_U = 1)$, which means eventually that

$$P((e_U = 1) \& (e_V = 1)) = P(e_U = 1)P(e_V = 1)$$

■

Proposition 16.

Let U and V be 2 independent random variables with values in $[0,1]$. Let $\theta_0 > 0$, $\lambda \gg 1$ be two real positive numbers, and θ be a random variable, independent of U and V , and distributed according to the distribution $g(\theta)d\theta = \frac{\lambda\theta_0}{(\lambda-1)\theta^2} d\theta$ over $\theta \in [\theta_0, \lambda\theta_0]$. Let also ρ be a random variable independent of U and V , and equi-distributed over $[0, \theta]$. We have:

- (i) If $|U - V| < \theta$, then $E[\Gamma_{\theta}(U - \rho) \oplus \Gamma_{\theta}(V - \rho)|U, V, \theta] = \frac{|U-V|}{\theta}$
- (ii) In all cases, $E[\Gamma_{\theta}(U - \rho) \oplus \Gamma_{\theta}(V - \rho)|U, V] \leq \frac{1}{2} + \frac{1}{2(\lambda-1)}$

■ (i): ρ can be considered as equidistributed over $[\min(U, V), \min(U, V) + \theta[$ because $\Gamma_{\theta}(U - \rho) \oplus \Gamma_{\theta}(V - \rho)$ is unchanged by translation of θ . Then the result is obvious by observing that ρ need to be within $[\min(U, V), \max(U, V)[$ so that $\Gamma_{\theta}(U - \rho) \oplus \Gamma_{\theta}(V - \rho) = 1$.

(ii): We can write:

$$E[\Gamma_{\theta}(U - \rho) \oplus \Gamma_{\theta}(V - \rho)|U, V] = \int_{\theta_0}^{\lambda\theta_0} E[\Gamma_{\theta}(U - \rho) \oplus \Gamma_{\theta}(V - \rho)|U, V, \theta]g(\theta)d\theta$$

$$= \int_{\theta_0}^{\lambda\theta_0} \left(\frac{|U - V|}{\theta} - \left\lfloor \frac{|U - V|}{\theta} \right\rfloor \right) g(\theta)d\theta = \frac{1}{|U - V|} \frac{\lambda\theta_0}{(\lambda - 1)} \int_{\frac{|U-V|}{\lambda\theta_0}}^{\frac{|U-V|}{\theta_0}} (u - [u])du$$

with the change of variable $u = \frac{|U-V|}{\theta}$. We remark, by considering the surface covered by the function $f(u) \triangleq u - [u]$ over $\left[\frac{|U-V|}{\lambda\theta_0}, \frac{|U-V|}{\theta_0}\right]$, that:

$$\frac{\theta_0}{|U - V|} \int_0^{\frac{|U-V|}{\theta_0}} f(u) du \leq \frac{1}{2}$$

which gives an upper-bound for $\frac{\theta_0}{|U-V|} \int_{\frac{|U-V|}{\lambda\theta_0}}^{\frac{|U-V|}{\theta_0}} (u - \lfloor u \rfloor) du$, and finally:

$$E[\Gamma_\theta(U - \rho) \oplus \Gamma_\theta(V - \rho) | U, V] \leq \frac{\lambda}{(\lambda - 1)} \frac{1}{2} = \frac{1}{2} + \frac{1}{2(\lambda - 1)}$$

■

We can now explicit the Separable Criteria lemma. This lemma is a useful tool to build probability distributions that are not sensitive to the Independence Phenomenon introduced in Section I.

Lemma 1. (Separable Criteria lemma)

Let M_Φ and $M_{\Phi'}$ be two symmetric matrix associated respectively to the probability distributions Φ and Φ' , we then have the inequality :

$$\langle \Phi, \Phi' \rangle \geq (4n) + n^2 \frac{n-1}{n-2} \|M_\Phi - \overline{M_\Phi}\|_c \|M_{\Phi'} - \overline{M_{\Phi'}}\|_c$$

■ It is first clear that within the group \mathfrak{S}_n :

$$\langle \Phi, \Phi' \rangle = \max_{\sigma, \mu \in \mathfrak{S}_n} |M_\Phi \cdot M_{\Phi'} - \sigma(M_\Phi) \cdot \mu(M_{\Phi'})|$$

For $I \in S_n$, we note $\mathfrak{S}_I = \{\sigma \in \mathfrak{S}_n | \forall u \in I, \sigma(u) \in I\}$; \mathfrak{S}_I is a subgroup of \mathfrak{S}_n , \bar{I} is also stable by action of any element of \mathfrak{S}_I , and $|\mathfrak{S}_I| = \left(\frac{n}{2}!\right)^2$. Let's fix σ_0 and $\mu_0 \in \mathfrak{S}_n$ and consider the sub-sum :

$$\Delta_{I,J}(\Phi, \Phi') = \left(\frac{n}{2}!\right)^{-4} \sum_{\sigma, \mu \in \mathfrak{S}_I \times \mathfrak{S}_J} \sigma_0 \circ \sigma(M_\Phi) \cdot \mu_0 \circ \mu(M_{\Phi'})$$

We observe the 3 following stability properties :

(i) for any $(u, v) \in I^2, u \neq v$, $\{\sigma \in \mathfrak{S}_I | \sigma(u) = u' \text{ \& } \sigma(v) = v'\}_{(u', v') \in I^2, u' \neq v'}$ is an equi-partition of \mathfrak{S}_I , which then implies :

$$\frac{1}{|\mathfrak{S}_I|} \sum_{\sigma \in \mathfrak{S}_I} M_\Phi(\sigma(u), \sigma(v)) = \frac{4}{n(n-2)} \sum_{\substack{r, s \in I \\ r \neq s}} M_\Phi(r, s) = s_I(\Phi)$$

(ii) for any $(u, v) \in I \times \bar{I}, \{\sigma \in \mathfrak{S}_I | \sigma(u) = u' \text{ \& } \sigma(v) = v'\}_{(u', v') \in I \times \bar{I}}$ is an equi-partition of \mathfrak{S}_I , which then implies :

$$\frac{1}{|\mathfrak{S}_I|} \sum_{\sigma \in \mathfrak{S}_I} M_\Phi(\sigma(u), \sigma(v)) = \frac{4}{n^2} \sum_{r, s \in I \times \bar{I}} M_\Phi(r, s) = c_I(\Phi)$$

(iii) for any $u \in I$, $\{\sigma \in \mathfrak{S}_I \mid \sigma(u) = u'\}_{u' \in I}$ is an equi-partition of \mathfrak{S}_I , which then implies :

$$\frac{1}{|\mathfrak{S}_I|} \sum_{\sigma \in \mathfrak{S}_I} M_{\Phi}(\sigma(u), \sigma(u)) = \frac{2}{n} \sum_{r \in I} M_{\Phi}(r, r) = d_I(\Phi)$$

By choosing σ_0 such that $\sigma_0(I) = \{1, \dots, n/2\}$ and μ_0 such that $\mu_0(J) = \{1, \dots, n/2\}$, we deduce from above the following equality, written with block matrix :

$$\sigma_0 \left(\left(\frac{n}{2}! \right)^{-2} \sum_{\sigma \in \mathfrak{S}_I} \sigma(M_{\Phi}) \right) = \begin{pmatrix} S_I(\Phi) & C_I(\Phi) \\ C_I(\Phi) & S_I(\Phi) \end{pmatrix}$$

where $S_I(\Phi) = \begin{pmatrix} d_I(\Phi) & \cdots & s_I(\Phi) \\ \vdots & \ddots & \vdots \\ s_I(\Phi) & \cdots & d_I(\Phi) \end{pmatrix}$ and $C_I(\Phi) = \begin{pmatrix} c_I(\Phi) & \cdots & c_I(\Phi) \\ \vdots & & \vdots \\ c_I(\Phi) & \cdots & c_I(\Phi) \end{pmatrix}$. And the same stands for

μ_0, J . In the following, we will note $C_n = \begin{pmatrix} 1 & \cdots & 1 \\ \vdots & & \vdots \\ 1 & \cdots & 1 \end{pmatrix}$.

By summing the elements of the matrix globally or by blocks, we get :

$$s_I(\Phi) + s_I(\Phi) = 4 \frac{n-1}{n-2} m_{\Phi} - \frac{2n}{n-2} c_I(\Phi) \quad (l1.1)$$

which enables us to express $\Delta_{I,J}(\Phi, \Phi')$ as follows :

$$\begin{aligned} \Delta_{I,J}(\Phi, \Phi') &= \frac{n}{2} \left[(d_I(\Phi) - s_I(\Phi)) (d_J(\Phi') - s_J(\Phi')) + (d_{\bar{I}}(\Phi) - s_{\bar{I}}(\Phi)) (d_{\bar{J}}(\Phi') - s_{\bar{J}}(\Phi')) \right. \\ &\quad + (d_I(\Phi) - s_I(\Phi)) s_{\bar{J}}(\Phi') + (d_J(\Phi') - s_J(\Phi')) s_I(\Phi) + (d_{\bar{I}}(\Phi) - s_{\bar{I}}(\Phi)) s_{\bar{J}}(\Phi') \\ &\quad \left. + (d_{\bar{J}}(\Phi') - s_{\bar{J}}(\Phi')) s_{\bar{I}}(\Phi) \right] + \frac{n^2}{4} [s_I(\Phi) s_J(\Phi') + s_{\bar{I}}(\Phi) s_{\bar{J}}(\Phi') + 2c_I(\Phi) c_J(\Phi')] = \\ &\triangleleft (n) + \frac{n^2}{4} [s_I(\Phi) s_J(\Phi') + s_{\bar{I}}(\Phi) s_{\bar{J}}(\Phi') + 2c_I(\Phi) c_J(\Phi')] \quad (l1.2) \end{aligned}$$

By symmetry of M_{Φ} and $M_{\Phi'}$ we indeed get $c_I(\Phi) = c_{\bar{I}}(\Phi)$ and $c_J(\Phi') = c_{\bar{J}}(\Phi')$.

Let's chose $I^+(\Phi)$ that achieves the maximum of $c_I(\Phi)$ over S_n , and $I^-(\Phi)$ that achieves the minimum. We then set, for $\xi, \xi' \in \{+, -\}$:

$$\Delta_{\xi, \xi'}(\Phi, \Phi') = \Delta_{I^{\xi}(\Phi), I^{\xi'}(\Phi')}(\Phi, \Phi') + \Delta_{I^{\xi}(\Phi), \overline{I^{\xi'}(\Phi')}}(\Phi, \Phi')$$

By direct calculation from (l1.2) :

$$\begin{aligned} \Delta_{\xi, \xi'}(\Phi, \Phi') &= \triangleleft (2n) \\ &\quad + \frac{n^2}{4} \left[\left(s_{I^{\xi}(\Phi)}(\Phi) + s_{\overline{I^{\xi}(\Phi)}}(\Phi) \right) \left(s_{I^{\xi'}(\Phi')}(\Phi') + s_{\overline{I^{\xi'}(\Phi')}}(\Phi') \right) \right. \\ &\quad \left. + 4c_{I^{\xi}(\Phi)}(\Phi) c_{I^{\xi'}(\Phi')}(\Phi') \right] \end{aligned}$$

and by applying (l1.1) :

$$|\Delta_{+,+}(\Phi, \Phi') - \Delta_{+,-}(\Phi, \Phi') - \Delta_{-,+}(\Phi, \Phi') + \Delta_{-,-}(\Phi, \Phi')| = \\ \leq (8n) + 2n^2 \frac{n-1}{n-2} |c_{I^+(\Phi)}(\Phi) - c_{I^-(\Phi)}(\Phi)| |c_{I^+(\Phi')}(\Phi') - c_{I^-(\Phi')}(\Phi')|$$

with obviously $|c_{I^+(\Phi)}(\Phi) - c_{I^-(\Phi)}(\Phi)| \geq \|M_\Phi - \overline{M}_\Phi\|_c$

And at last, on the other hand :

$$|\Delta_{+,+}(\Phi, \Phi') - \Delta_{+,-}(\Phi, \Phi') - \Delta_{-,+}(\Phi, \Phi') + \Delta_{-,-}(\Phi, \Phi')| \leq 2 \max_{I, J, I', J' \in \mathcal{S}_n} |\Delta_{I, J}(\Phi, \Phi') - \Delta_{I', J'}(\Phi, \Phi')| \\ \leq 2 \max_{\sigma, \mu \in \mathfrak{S}_n} |M_\Phi \cdot M_{\Phi'} - \sigma(M_\Phi) \cdot \mu(M_{\Phi'})| \leq 2\langle \Phi, \Phi' \rangle$$

which concludes the proof. ■

The following Lemma gives a relationship between tidying and synchronization.

Lemma 1' (Synchronized Distributions Lemma).

For Φ and Φ' two tidied distribution in $\zeta(\alpha)$, we have :

$$E[\Delta_0(\Phi, \Phi' \circ \tau) | |I_0 \cap \tau(I_0)| = r] = \frac{1}{n^2} (n - 4r)^2 (m_\Phi - c_\Phi)(m'_{\Phi'} - c'_{\Phi'}) + O\left(\frac{1}{n}\right)$$

■

For a given distribution Φ , we denote :

$$\Sigma_{I_0}(\Phi) \triangleq \frac{1}{|\mathfrak{S}_{I_0}|} \sum_{\tau \in \mathfrak{S}_{I_0}} \Phi \circ \tau$$

We also denote, for $L \in \mathcal{S}_n$:

$$\mathfrak{S}_{I, L} = \{\sigma \in \mathfrak{S}_n | \forall u \in I, \sigma(u) \in L\}$$

We can initially remark that, for any given $I \in \mathcal{S}_n$, the set $\{\mathfrak{S}_{I, L}\}_{L \in \mathcal{S}_n}$ is an equi-partition of \mathfrak{S}_n . From this, we can deduce that $|\mathfrak{S}_{I, L}|$ only depends on $|I \cap L|$. In particular, we can thus set :

$$\mathcal{L}_0 \triangleq \{L | |I_0 \cap L| = r\}$$

Let's express $E[\Delta_0(\Phi, \Phi' \circ \tau) | |I_0 \cap \tau(I_0)| = r]$ in terms of $\mathfrak{S}_{I_0, L}$

$$E[\Delta_0(\Phi, \Phi' \circ \tau) | |I_0 \cap \tau(I_0)| = r] = \frac{1}{|\mathcal{L}_0|} \sum_{L \in \mathcal{L}_0} \frac{1}{|\mathfrak{S}_{I_0}|^2} \frac{1}{|\mathfrak{S}_{I_0, \tau(I_0)}|} \sum_{\sigma, \mu \in \mathfrak{S}_{I_0}} \sum_{\tau \in \mathfrak{S}_{I_0, L}} \Delta_0(\Phi \circ \sigma, \Phi' \circ \mu \circ \tau) \\ = \frac{1}{|\mathcal{L}_0|} \sum_{L \in \mathcal{L}_0} \frac{1}{|\mathfrak{S}_{I_0, \tau(I_0)}|} \sum_{\tau \in \mathfrak{S}_{I_0, L}} \Delta_0(\Sigma_{I_0}(\Phi), \Sigma_{I_0}(\Phi') \circ \tau) \quad (l1'.1)$$

The quadratic matrix of $\Sigma_{I_0}(\Phi)$ has the form :

$$\begin{pmatrix} S_\Phi & C_\Phi \\ C_\Phi & \bar{S}_\Phi \end{pmatrix}$$

Where :

$$S_{\Phi} \triangleq \begin{pmatrix} d_{\Phi} & \cdots & s_{\Phi} \\ \vdots & \ddots & \vdots \\ s_{\Phi} & \cdots & d_{\Phi} \end{pmatrix}, \bar{S}_{\Phi} \triangleq \begin{pmatrix} \bar{d}_{\Phi} & \cdots & \bar{s}_{\Phi} \\ \vdots & \ddots & \vdots \\ \bar{s}_{\Phi} & \cdots & \bar{d}_{\Phi} \end{pmatrix} \text{ and } C_{\Phi} \triangleq \begin{pmatrix} c_{\Phi} & \cdots & c_{\Phi} \\ \vdots & & \vdots \\ c_{\Phi} & \cdots & c_{\Phi} \end{pmatrix},$$

$$s_{\Phi} \triangleq \frac{1}{|\mathfrak{S}_{I_0}|} \sum_{\sigma \in \mathfrak{S}_{I_0}} M_{\Phi}(\sigma(u), \sigma(v)) = \frac{4}{n(n-2)} \sum_{\substack{r,s \in I_0 \\ r \neq s}} M_{\Phi}(r, s)$$

$$\bar{s}_{\Phi} \triangleq \frac{1}{|\mathfrak{S}_{\bar{I}_0}|} \sum_{\sigma \in \mathfrak{S}_{\bar{I}_0}} M_{\Phi}(\sigma(u), \sigma(v)) = \frac{4}{n(n-2)} \sum_{\substack{r,s \in \bar{I}_0 \\ r \neq s}} M_{\Phi}(r, s)$$

$$d_{\Phi} \triangleq \frac{1}{|\mathfrak{S}_{I_0}|} \sum_{\sigma \in \mathfrak{S}_{I_0}} M_{\Phi}(\sigma(u), \sigma(u)) = \frac{2}{n} \sum_{r \in I_0} M_{\Phi}(r, r)$$

$$\bar{d}_{\Phi} \triangleq \frac{1}{|\mathfrak{S}_{\bar{I}_0}|} \sum_{\sigma \in \mathfrak{S}_{\bar{I}_0}} M_{\Phi}(\sigma(u), \sigma(u)) = \frac{2}{n} \sum_{r \in \bar{I}_0} M_{\Phi}(r, r)$$

Then, starting from the general relation :

$$\Delta_0(\Phi, \Phi') = \frac{1}{n^4} \sum_{i,j,u,v} M_{\Phi}(i, u) M_{\Phi'}(j, v) - \frac{2}{n^3} \sum_{i,j,u} M_{\Phi}(i, u) M_{\Phi'}(j, u) + \frac{1}{n^2} \sum_{i,j} M_{\Phi}(i, j) M_{\Phi'}(i, j)$$

and using the above matrix forms, we can calculate (calculations are a bit long and are not presented here in details) :

$$\Delta_0(\Sigma_{I_0}(\Phi), \Sigma_{I_0}(\Phi') \circ \tau) = \frac{1}{n^2} (n - 4r)^2 (m_{\Phi} - c_{\Phi})(m'_{\Phi} - c'_{\Phi}) + O\left(\frac{1}{n}\right)$$

And because this only depends on r , we can conclude from (l1'. 1). ■

The following lemma gives a useful calculation method when using distributions transformed by permutative sleeking. It applies the method for a specific variable, but we will rely on the same method to establish other similar equalities involving such distributions.

Lemma 2.

Let Φ and Φ' be two probability distributions, and γ be a permutative sleeking kernel, then we have :

$$\begin{aligned} \Delta_{\gamma}(\Phi, \Phi') &\triangleq \sum_{\sigma \in \mathfrak{S}_n} \gamma(|\sigma|) \int_{x,y \in [0,1]^n} (x \cdot y - x \cdot \sigma(y))^2 \Phi(x) \Phi'(y) dx dy \\ &= \sum_{s=0}^n \gamma(s) A_n^s \left(\frac{s^2}{e} \Delta_0(\Phi, \Phi') + O(n) \right) \end{aligned}$$

■ For I a subset of \mathbb{N}_n we define by $|I|$ the cardinality of I and \mathcal{S}_I the subgroup of \mathfrak{S}_n including permutations for which $\text{supp}(\sigma) \subset I$. Let's set :

$$f(\sigma) = \int_{x,y \in [0,1]^n} (x \cdot y - x \cdot \sigma(y))^2 \Phi(x) \Phi'(y) dx dy$$

$$F(s) = \sum_{\substack{\sigma \in \mathfrak{S}_n \\ |\sigma|=s}} f(\sigma)$$

$$G(s) = \sum_{\substack{I \\ |I|=s}} \sum_{\sigma \in \mathcal{S}_I} f(\sigma)$$

We first establish a relationship between G and F :

$$G(s) = \sum_{\substack{\sigma \in \mathfrak{S}_n \\ |\sigma| \leq s}} \left(\sum_{\substack{I \\ |I|=s \\ I \supset \text{supp}(\sigma)}} 1 \right) f(\sigma) = \sum_{\substack{\sigma \in \mathfrak{S}_n \\ |\sigma| \leq s}} \binom{n-|\sigma|}{s-|\sigma|} f(\sigma) = \sum_{u=0}^s \binom{n-u}{s-u} F(u)$$

This formula can be reversed by using the classical matrix inverse :

$$\left(\begin{array}{ccc} \dots & \dots & 0 \\ \vdots & \ddots & \vdots \\ \binom{i}{j} & \dots & \dots \end{array} \right)_{i,j \in \mathbb{N}_n}^{-1} = \left(\begin{array}{ccc} \dots & \dots & 0 \\ \vdots & \ddots & \vdots \\ (-1)^{i+j} \binom{i}{j} & \dots & \dots \end{array} \right)_{i,j \in \mathbb{N}_n}$$

(which can be easily seen by considering for instance the inverse of the $\mathbb{R}_n[X]$ endomorphism $P(X) \mapsto P(X+1)$).

By setting $j' = n-j$ and $i' = n-i$, we have $\binom{n-j}{i-j} = \binom{j'}{j'-i'} = \binom{j'}{i'}$, and thus :

$$\left(\begin{array}{ccc} \dots & \dots & 0 \\ \vdots & \ddots & \vdots \\ \binom{n-j}{i-j} & \dots & \dots \end{array} \right)_{i,j \in \mathbb{N}_n}^{-1} = \left(\begin{array}{ccc} \dots & \dots & 0 \\ \vdots & \ddots & \vdots \\ (-1)^{i+j} \binom{n-j}{i-j} & \dots & \dots \end{array} \right)_{i,j \in \mathbb{N}_n}$$

from which we take the reverse formula :

$$F(s) = \sum_{u=0}^s (-1)^{u+s} \binom{n-u}{s-u} G(u)$$

Thanks to the above, we can write Δ_γ depending on $G(u)$ as follows:

$$\begin{aligned}
\Delta_\gamma &= \sum_{\sigma \in \mathfrak{S}_n} \gamma(|\sigma|) f(\sigma) = \sum_{s=0}^n \gamma(s) \sum_{\substack{\sigma \in \mathfrak{S}_n \\ |\sigma|=s}} f(\sigma) = \sum_{s=0}^n \gamma(s) F(s) \\
&= \sum_{s=0}^n \gamma(s) \sum_{u=0}^s (-1)^{u+s} \binom{n-u}{s-u} G(u)
\end{aligned} \tag{I2.1}$$

Let's now calculate $G(u)$ by using the properties of \mathcal{S}_I as a group (in the following, C, C', C'' are constants); we will use the notation $\Phi_{r,t} = M_\Phi(r, t)$ to simplify the writings :

$$\begin{aligned}
G(u) &= \sum_{\substack{I \\ |I|=s}} \sum_{\sigma \in \mathcal{S}_I} \left(\sum_{r,t \in \mathbb{N}_n^*} (\Phi_{r,t} \Phi'_{r,t} - 2\Phi_{r,t} \Phi'_{r,\sigma(t)} + \Phi_{r,t} \Phi'_{\sigma(r),\sigma(t)}) \right) \\
&= \sum_{\substack{I \\ |I|=s}} \left(u! \sum_{r,t \in \mathbb{N}_n^*} \Phi_{r,t} \Phi'_{r,t} - 2(u-1)! \sum_{r,t,t' \in I} \Phi_{r,t} \Phi'_{r,t'} - 2u! \sum_{r,t \in \mathbb{N}_n^* \times \bar{I}} \Phi_{r,t} \Phi'_{r,t} \right. \\
&\quad \left. + (u-2)! \sum_{r,r',t,t' \in I} \Phi_{r,t} \Phi'_{r',t'} + u! \sum_{r,t \in \bar{I}} \Phi_{r,t} \Phi'_{r,t} + \ll (Cnu!) \right) \\
&= u! \binom{n-2}{u-2} \sum_{r,t \in \mathbb{N}_n^*} \Phi_{r,t} \Phi'_{r,t} + (u-2)! \binom{n-4}{u-4} \sum_{r,r',t,t' \in \mathbb{N}_n^*} \Phi_{r,t} \Phi'_{r',t'} \\
&\quad - 2(u-1)! \binom{n-3}{u-3} \sum_{r,t,t' \in \mathbb{N}_n^*} \Phi_{r,t} \Phi'_{r,t'} + \ll (C''nu! \binom{n}{u}) \\
&= A_n^u (u^2 \Delta_0(\Phi, \Phi') + \ll (C''n))
\end{aligned}$$

In addition, we have :

$$\sum_{u=0}^s (-1)^{u+s} u^2 A_n^u \binom{n-u}{n-s} = A_n^s \sum_{u=0}^s (-1)^{u+s} \frac{u^2}{(s-u)!} = A_n^s \left(\frac{s^2}{e} + \frac{2s}{e} + O(1) \right)$$

and :

$$\left| \sum_{u=0}^s (-1)^{u+s} A_n^u \binom{n-u}{n-s} \ll (C''n) \right| \leq C''n \sum_{u=0}^s A_n^u \binom{n-u}{n-s} \leq eC''nA_n^s$$

By injecting the calculation of $G(u)$ in (I2.1), we can conclude the proof with the two above relations. ■

An immediate consequence of lemma 2 is that, if the probability distributions Φ and $\Phi' \in \zeta(\alpha)$ are synchronized, then stands the following inequality :

$$\Delta_\gamma(\Phi, \Phi') \geq \sum_{s=0}^n \gamma(s) A_n^s \left(\frac{s^2}{e} \left(\frac{\alpha}{4} - \ll \left(\frac{1}{n} \right) \right)^2 + O(n) \right)$$

We will also use the following corollary, that relies on the same calculation technique than presented in Lemma 2.

The « border polynomial » is defined by :

$$P_{u,v}(X) \triangleq X \left(\frac{u}{n} - X \right) \left(\frac{v}{n} - X \right) \left(1 - \frac{u}{n} - \frac{v}{n} + X \right)$$

Corollary 1 (treatment of border conditions).

For Φ and Φ' two synchronized probability distributions in $\zeta(\alpha)$, there exists a constant $K(\alpha) > 0$ (independent from Φ and Φ') and a permutative sleeking transformation δ , depending on α but not on Φ and Φ' , such that :

$$\begin{aligned} & \sum_{\sigma \in \mathbb{G}_n} \int_{x,y \in [0,1]^n} P_{|x|,|y|} \left(\frac{x \cdot y}{n} \right) P_{|x|,|y|} \left(\frac{x \cdot \sigma(y)}{n} \right) \left(\frac{x \cdot y}{n} - \frac{x \cdot \sigma(y)}{n} \right)^2 \gamma(|\sigma|) \Phi(x) T_\delta[\Phi'](y) dx dy \\ & \geq K(\alpha) \sum_{r=0}^n \frac{r^2}{n^2} A_n^r \gamma(r) - \frac{1}{\sqrt{e}} \sum_{r=0}^n \frac{r^3}{n^3} A_n^r \gamma(r) - o\left(\frac{1}{\sqrt{n}}\right) \end{aligned}$$

■

By using the same calculation technique than in Lemma 2 (we won't present the details here), we obtain more generally that for any polynomial P with degree p , and $t \in [0,1]$, there exists a constant $K_p > 0$ such that :

$$\frac{1}{A_n^{nt}} \sum_{\substack{\sigma \in \mathbb{G}_n \\ |\sigma|=nt}} P \left(\frac{x \cdot \sigma(y)}{n} \right) = \frac{1}{e} P \left(t \frac{|x||y|}{n^2} + (1-t) \frac{x \cdot y}{n} \right) + \triangleleft \left(\frac{K_p \|P\|_\infty}{n} \right) \quad (c1.1)$$

We have obviously (because $|P_{|x|,|y|}'(X)| \leq 4$ over $[0,1]$) :

$$P_{|x|,|y|} \left(\frac{x \cdot \sigma(y)}{n} \right) = P_{|x|,|y|} \left(\frac{x \cdot y}{n} \right) + \triangleleft \left(4 \left| \frac{x \cdot y}{n} - \frac{x \cdot \sigma(y)}{n} \right| \right)$$

Let's apply a first time (c1.1) to :

$$\begin{aligned} & \frac{1}{A_n^{nt}} \sum_{\substack{\sigma \in \mathbb{G}_n \\ |\sigma|=nt}} \int_{x,y \in [0,1]^n} P_{|x|,|y|} \left(\frac{x \cdot y}{n} \right) P_{|x|,|y|} \left(\frac{x \cdot \sigma(y)}{n} \right) \left(\frac{x \cdot y}{n} - \frac{x \cdot \sigma(y)}{n} \right)^2 \Phi(x) \Phi'(y) dx dy \\ & \geq \frac{1}{A_n^{nt}} \sum_{\substack{\sigma \in \mathbb{G}_n \\ |\sigma|=nt}} \int_{x,y \in [0,1]^n} P_{|x|,|y|} \left(\frac{x \cdot y}{n} \right)^2 \left(\frac{x \cdot y}{n} - \frac{x \cdot \sigma(y)}{n} \right)^2 \Phi(x) \Phi'(y) dx dy \\ & \quad - 4 \left(\frac{1}{A_n^{nt}} \sum_{\substack{\sigma \in \mathbb{G}_n \\ |\sigma|=nt}} \int_{x,y \in [0,1]^n} \left(\frac{x \cdot y}{n} - \frac{x \cdot \sigma(y)}{n} \right)^6 \Phi(x) \Phi'(y) dx dy \right)^{1/2} \\ & \geq \frac{t^2}{e} \int_{x,y \in [0,1]^n} P_{|x|,|y|} \left(\frac{x \cdot y}{n} \right)^2 \left(\frac{|x||y|}{n^2} - \frac{x \cdot y}{n} \right)^2 \Phi(x) \Phi'(y) dx dy + \\ & \triangleleft \left(\frac{2K_2}{n} \right) - 4 \left(\frac{t^6}{e} + \triangleleft \left(\frac{15K_6}{n} \right) \right)^{1/2} \end{aligned} \quad (c1.2)$$

Let's consider the probability distribution, obtained by the following « Dirac » permutative sleeking transformation :

$$\Phi'_l = \frac{1}{A_n^{nl}} \sum_{\substack{\mu \in \mathcal{C}_n \\ |\mu|=nl}} \Phi' \circ \mu$$

It can be easily remarked that :

$$P_{|x|,|y|} \left(t \frac{|x||y|}{n^2} + (1-t) \frac{x \cdot y}{n} \right) \geq t^4 \left(\frac{|x||y|}{n} \frac{1}{n} \left(1 - \frac{|x|}{n} \right) \left(1 - \frac{|y|}{n} \right) \right)^2$$

By replacing Φ' by Φ'_l in (c1.2) and applying a second time (c1.1) to the first term of the right side of the inequality, with the above we obtain :

$$\begin{aligned} & \int_{x,y \in [0,1]^n} P_{|x|,|y|} \left(\frac{x \cdot y}{n} \right)^2 \left(\frac{|x||y|}{n^2} - \frac{x \cdot y}{n} \right)^2 \Phi(x) \Phi'_l(y) dx dy \\ & \geq \frac{1}{A_n^{nl}} \sum_{\substack{\mu \in \mathcal{C}_n \\ |\mu|=nl}} \int_{x,y \in [0,1]^n} P_{|x|,|y|} \left(\frac{x \cdot \mu(y)}{n} \right)^2 \left[\left(\frac{|x||y|}{n^2} - \frac{x \cdot y}{n} \right)^2 - 2l \right] \Phi(x) \Phi'(y) dx dy \\ & \geq \frac{l^8}{e} \int_{x,y \in [0,1]^n} \left(\frac{|x||y|}{n} \frac{1}{n} \left(1 - \frac{|x|}{n} \right) \left(1 - \frac{|y|}{n} \right) \right)^4 \left[\left(\frac{|x||y|}{n^2} - \frac{x \cdot y}{n} \right)^2 - 2l \right] \Phi(x) \Phi'(y) dx dy + \ll \left(\frac{28K_8}{n} \right) \end{aligned} \quad (c1.3)$$

In addition, as $\left(\frac{|x||y|}{n^2} - \frac{x \cdot y}{n} \right)^2$ is unchanged by transformations $x \mapsto \bar{x}$ and $y \mapsto \bar{y}$ we can write :

$$\left(\frac{|x||y|}{n^2} - \frac{x \cdot y}{n} \right)^2 \leq \min \left(\left(\frac{|x||y|}{n^2} \right)^2, \left(\left(1 - \frac{|x|}{n} \right) \left(1 - \frac{|y|}{n} \right) \right)^2 \right) \leq \frac{|x||y|}{n} \frac{1}{n} \left(1 - \frac{|x|}{n} \right) \left(1 - \frac{|y|}{n} \right)$$

And therefore we can go on (c1.3) with Proposition 9 and the above :

$$\begin{aligned} & \int_{x,y \in [0,1]^n} P_{|x|,|y|} \left(\frac{x \cdot y}{n} \right)^2 \left(\frac{|x||y|}{n^2} - \frac{x \cdot y}{n} \right)^2 \Phi(x) \Phi'_l(y) dx dy \\ & \geq \frac{l^8}{e} \int_{x,y \in [0,1]^n} \left[\left(\frac{|x||y|}{n^2} - \frac{x \cdot y}{n} \right)^{10} - 2l \right] \Phi(x) \Phi'(y) dx dy + \ll \left(\frac{28K_8}{n} \right) \\ & \geq \frac{l^8}{e} [\Delta_0(\Phi, \Phi')^5 - 2l] + \ll \left(\frac{28K_8}{n} \right) \geq \frac{l^8}{e} \left[\alpha^{10} - 2l \right] + O\left(\frac{1}{n}\right) \end{aligned}$$

We can take the maximum over $l \in [0,1]$ which is obtained for $l_0 = \frac{\alpha^{10}}{9 \cdot 2^{18}}$:

$$\int_{x,y \in [0,1]^n} P_{|x|,|y|} \left(\frac{x \cdot y}{n} \right)^2 \left(\frac{|x||y|}{n^2} - \frac{x \cdot y}{n} \right)^2 \Phi(x) \Phi'_{l_0}(y) dx dy \geq eK(\alpha) + O\left(\frac{1}{n}\right)$$

By injecting the above in (c1.2) and remarking that $\sqrt{a+b} \leq \sqrt{a} + \sqrt{b}, \forall a, b > 0$, we obtain the result. The permutative sleeking transformation $T[\Phi'] = \frac{1}{A_n^{nl_0}} \sum_{\substack{\mu \in \mathfrak{S}_n \\ |\mu|=nl_0}} \Phi' \circ \mu$ is indeed only depending on α .

■

One can remark that the set of strategies $\Omega_{\#}$ introduced in Section IV can be equivalently defined as :

$$\Omega_{\#} = \{\omega \in [0,1]^{2^{2n}} \mid \omega_{i,j} = f(|i|, |j|, i.j), \forall f: \mathbb{N}_n^3 \mapsto [0,1]\}$$

Let's justify this equivalence :

■ it is clear that any strategy of the form $f(|i|, |j|, i.j)$ is in $\Omega_{\#}$. Let ω be in $\Omega_{\#}$, and let's consider (i, j) and (i', j') such that :

$$\begin{cases} |i| = |i'| \\ |j| = |j'| \\ i.j = i'.j' \end{cases}$$

Then, as $\{i \setminus j, i \cap j, j \setminus i, \overline{i \cup j}\}$ and $\{i' \setminus j', i' \cap j', j' \setminus i', \overline{i' \cup j'}\}$ are isomorphic partitions of \mathbb{N}_n^* , there exists $\sigma \in \mathfrak{S}_n$ such that :

$$\begin{cases} \sigma(i \setminus j) = \sigma(i' \setminus j') \\ \sigma(i \cap j) = \sigma(i' \cap j') \\ \sigma(j \setminus i) = \sigma(j' \setminus i') \\ \sigma(\overline{i \cup j}) = \sigma(\overline{i' \cup j'}) \end{cases}$$

and thus $\omega_{i',j'} = \omega_{\sigma(i),\sigma(j)} = \omega_{i,j}$, which means that $\omega_{i,j}$ only depends on $|i|, |j|, i.j$

■

Lemma 3. (Synchronized Strategy lemma)

Let ω be a strategy in $\Omega_{\#}$, and Φ and Φ' be two probability distributions in $\zeta(\alpha)$ with at least one of them transformed by permutative sleeking (with a permutative sleeking transformation that will be explicited), say $\Phi' = T[\Phi''']$, then for σ performing a synchronization of Φ and Φ''' , there exists a constant $C(\alpha)$ such that :

$$E \left[\left(\omega_{i,j} - \frac{x.y}{nk} \right)^2 \right]_{\Phi, T[\Phi''' \circ \sigma]} \geq \frac{C(\alpha)}{n}$$

■ The proof of lemma 3 is a bit long, we will sometimes give shortened explanations to avoid to present exhaustively some painful calculations. We divide the proof in 3 parts. In part 1, we will reduce the general Bernoulli distributions to simpler multi-Gaussian form. In part 2, we will discard the dependence in $\omega_{i,j}$. And in part 3, we will show that the resultant form from part 2 is not submitted to rapid variation in $x.y$ when $|x|$ and $|y|$ are fixed. In the conclusive part, we will deal with border conditions.

In the following, we will suppose Φ''' synchronized to Φ and thus we will denote the probability distribution directly Φ''' instead of $\Phi''' \circ \sigma$, and thus $\Phi' = T[\Phi''']$ instead of $\Phi' = T[\Phi''' \circ \sigma]$. T will be explicited in the course of the proof. We will also restrict the probability distributions to the generation of experiment vectors in $\{0,1\}^n$ instead of $[0,1]^n$ because it eases the proof, and also because anyway, they are easier to manipulate with computers.

We will denote in the followings:

$$\langle \omega, \Phi, \Phi' \rangle \triangleq E \left[\left(\omega_{i,j} - \frac{x \cdot y}{nk} \right)^2 \right]_{\Phi, \Phi'}$$

Part 1.

Let's first set the following quantities, for $\{u, v, w\} \in \mathbb{N}_n^3$:

$$\mathcal{J}(u) \triangleq \{x \in \mathbb{N}_n \mid |x| = u\}$$

$$\mathcal{K}(u, v, w) \triangleq \{x, y \in \mathbb{N}_n \mid |x| = u; |y| = v; x \cdot y = w\}$$

$$\widehat{\mathcal{K}}(u, v, w) \triangleq \{x, y \in \mathbb{N}_n \mid |x| \geq u; |y| \geq v; x \cdot y \geq w\}$$

$$P(u, v, w) \triangleq \int_{\mathcal{K}(u, v, w)} \Phi(x) \Phi'(y) dx dy$$

$$\psi_{u, v, w}(x, y) \triangleq \sum_{i, j \in \mathcal{K}(u, v, w)} \chi_i(x) \chi_j(y)$$

$\{\psi_{r, s, t}\}$ is a linearly independant set that generates a vector subspace ; by considering dimension and because for any $\sigma \in \mathfrak{S}_n$,

$$\psi_{r, s, t}(\sigma(x), \sigma(y)) = \psi_{r, s, t}(x, y)$$


this subspace is thus isomorphic to the vector subspace of the multinomials :

$$\left\{ \sum_{i, j \in \{0,1\}^n} \omega_{i,j} \chi_i(x) \chi_j(y) \mid \omega \in \Omega_{\#} \right\}$$

This implies that the multinomial $\psi_{r, s, t} \left(\frac{x}{k}, \frac{y}{k} \right)$ can be written as linear combination of $\{\psi_{u, v, w}(x, y)\}_{u, v, w}$. Let's then write this combination, with same method than seen in Proposition 1 :

$$\begin{aligned} \sum_{i, j \in \mathcal{K}(r, s, t)} \chi_i \left(\frac{x}{k} \right) \chi_j \left(\frac{y}{k} \right) &= \sum_{i, j \in \mathcal{K}(r, s, t)} \frac{1}{k^{r+s}} \sum_{\substack{l \supseteq i \\ l' \supseteq j}} \left(1 - \frac{1}{k} \right)^{|l|+|l'|-r-s} \chi_l(x) \chi_{l'}(y) \\ &= \sum_{l, l' \in \widehat{\mathcal{K}}(r, s, t)} \frac{1}{k^{r+s}} \left(\sum_{\substack{i \subseteq l \\ j \subseteq l'}} 1 \right) \left(1 - \frac{1}{k} \right)^{|l|+|l'|-r-s} \chi_l(x) \chi_{l'}(y) \end{aligned}$$

The drawing below :



helps to understand the following expression of the above summing between brackets, where u, v, w stand for $u = |l|, v = |l'|, w = l \cdot l'$:

$$\sum_{\substack{i,j \in \mathcal{K}(r,s,t) \\ i \subset l \\ j \subset l'}} 1 = \binom{w}{t} \sum_{q=0}^{\min(r-t, w-t, v-s)} \binom{w-t}{q} \binom{u-w}{r-t-q} \binom{v-t-q}{s-t}$$

Thanks to which we obtain :

$$\begin{aligned} & \frac{1}{k^{r+s}} \left(\sum_{\substack{i,j \in \mathcal{K}(r,s,t) \\ i \subset l \\ j \subset l'}} 1 \right) \left(1 - \frac{1}{k}\right)^{|l|+|l'|-r-s} \\ &= \beta_{t,w} \left(\frac{1}{k^2}\right) \sum_{q=0}^{\min(r-t, w-t, v-s)} \beta_{q,w-t} \left(\frac{1}{k+1}\right) \beta_{s-t, v-t-q} \left(\frac{1}{k}\right) \beta_{r-t-q, u-w} \left(\frac{1}{k}\right) \\ &= \beta_{t,w} \left(\frac{1}{k^2}\right) \sum_{q=t}^{\min(r,w, v-s+t)} \beta_{q-t, w-t} \left(\frac{1}{k+1}\right) \beta_{s-t, v-q} \left(\frac{1}{k}\right) \beta_{r-q, u-w} \left(\frac{1}{k}\right) \end{aligned} \quad (I3.1)$$

We remark that, at distance $O\left(\sqrt{\frac{n}{k}}\right)$ in the neighborhood of :

$$\begin{cases} r = u/k \\ s = v/k \\ t = w/k^2 \end{cases}$$

the three maxima of the Gaussian filters under the sum in (I3.1) are reached for an identical value of $q_m = \frac{w}{k}$ because

$$\frac{q_m - t}{w - t} = \frac{w/k - w/k^2}{w - w/k^2} = \frac{1}{k+1}$$

And by also remarking that :

$$\beta_{s-t, v-q_m} \left(\frac{1}{k}\right) = (1 + o(1)) \beta_{s-t + \frac{q_m}{k}, v} \left(\frac{1}{k}\right)$$

we see that the width of the Gaussian filter $\beta_{q-t, w-t} \left(\frac{1}{k+1} \right)$ in q is smaller than the width of the two other filters. Thus, by approximating the summing on the narrowest of the two other filters, we obtain that there exists a constant $C_1 > 0$ such that :

$$(I3.1) = \triangleright (C_1) \beta_{t,w} \left(\frac{1}{k^2} \right) \beta_{s-t, v-q_m} \left(\frac{1}{k} \right) \beta_{r-q_m, u-w} \left(\frac{1}{k} \right) \sum_{q=t}^r \beta_{q-t, w-t} \left(\frac{1}{k+1} \right)$$

And because $t \ll \frac{w}{k} \leq \frac{u}{k}$ in the considered neighborhood, we have :

$$\sum_{q=t}^r \beta_{q-t, w-t} \left(\frac{1}{k+1} \right) \sim 1$$

Thus finally there exists a constant $C_2 > 0$ such that :

$$(I3.1) = \triangleright (C_2) \beta_{t,w} \left(\frac{1}{k^2} \right) \beta_{s,v} \left(\frac{1}{k} \right) \beta_{r,u} \left(\frac{1}{k} \right)$$

We saw that, as $\omega \in \Omega_{\#}$, it can be written as $\omega_{i,j} = f(|i|, |j|, i, j)$, thus, in the followings, we will write $\omega_{i,j} = \omega_{r,s}(t)$. It follows from the above that :

$$\begin{aligned} \langle \omega, \Phi, \Phi' \rangle = \triangleright (C_2) \sum_{u,v \in \mathbb{N}_n} \sum_{r,s \in \mathbb{N}_n} \left(\sum_{w \in \mathbb{N}_n} \left(\sum_{t=0}^w (\omega_{r,s}(t) - \frac{w}{nk})^2 \beta_{t,w} \left(\frac{1}{k^2} \right) \right) P(u, v, w) \right) \beta_{s,v} \left(\frac{1}{k} \right) \beta_{r,u} \left(\frac{1}{k} \right) \end{aligned} \quad (I3.2)$$

which ends Part 1.

Part 2.

From Proposition 4, $\beta_{\frac{w}{k^2}+u, w} \left(\frac{1}{k^2} \right)$ can be approximated (with $\sigma^2 = \frac{1}{k^2} \left(1 - \frac{1}{k^2} \right)$) by

$$\triangleright (C'') N(u, w\sigma^2)$$

under the condition that $|u| \leq \left(\frac{w}{k^2} \right)^{2/3}$. On the other hand, from Proposition 2, if $|u| > \mathcal{K}(\alpha) \sqrt{\frac{w}{k^2}}$ for $\mathcal{K}(\alpha)$ sufficiently large, the remains of the binomial distribution can be kept bounded by $\frac{C(\alpha)}{2n}$. Thus, supposing that $n \gg k^2$, it is possible to ease calculations to replace the binomial distributions by gaussian distributions in (I3.2), with a multiplicative lower bound.

Let's now consider :

$$F_{r,s,u,v} \triangleq \int_{w \in \mathbb{R}} \left(\int_{t \in \mathbb{R}} \left(\omega_{r,s}(t) - \frac{w}{nk} \right)^2 N \left(t - \frac{w}{k^2}, w\sigma^2 \right) dt \right) P(u, v, w) dw$$

where $\omega_{r,s}(t)$ is extended to \mathbb{R} being 0 outside $[0, n]$ (note also that $P(u, v, w) = 0$ if $w \notin [0, n]$). The above argument means that there exists a constant $C_3 > 0$ such that :

$$\langle \omega, \Phi, \Phi' \rangle = \triangleright (C_3) \sum_{u,v \in \mathbb{N}_n} \sum_{r,s \in \mathbb{N}_n} F_{r,s,u,v} \beta_{s,v} \left(\frac{1}{k} \right) \beta_{r,u} \left(\frac{1}{k} \right) \quad (13.2')$$

We now study $F_{r,s,u,v}$ with objective to discard the dependence in $\omega_{r,s}(t)$.

On one hand, if V and V' are two random variables with identical probability distribution, and K a constant, we have the obvious relations :

$$E[(V - K)^2] = E[(V - E[V])^2] + (E[V] - K)^2 = \frac{1}{2} E[(V - V')^2] + (E[V] - K)^2 \geq \frac{1}{2} E[(V - V')^2]$$

which, applied to $F_{r,s,u,v}$ gives :

$$F_{r,s,u,v} \geq \int_{w \in \mathbb{R}} \left(\int_{t, \delta \in \mathbb{R}} \left(\omega_{r,s}(t + \delta) - \omega_{r,s}(t) \right)^2 N \left(t + \delta - \frac{w}{k^2}, w\sigma^2 \right) N \left(t - \frac{w}{k^2}, w\sigma^2 \right) dt d\delta \right) P(u, v, w) dw \quad (13.3)$$

We observe that :

$$N \left(t + \delta - \frac{w}{k^2}, w\sigma^2 \right) N \left(t - \frac{w}{k^2}, w\sigma^2 \right) = N(\delta, 2w\sigma^2) N \left(t + \frac{\delta}{2} - \frac{w}{k^2}, \frac{w\sigma^2}{2} \right)$$

We use the change of variable :

$$\begin{pmatrix} w \\ \delta \end{pmatrix} \mapsto \begin{pmatrix} 1 & -k^2/2 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} w \\ \delta \end{pmatrix}$$

whose Jacobian is 1. Let's set, for $\in]0,1[$:

$$J(\delta, \lambda) = \{w | (w - k^2|\delta| \geq 2\lambda n) \& (n \geq w)\}$$

We have then, for $\in J(\delta, \lambda)$:

$$N \left(\delta, 2 \left(w + \frac{k^2\delta}{2} \right) \sigma^2 \right) \geq \sqrt{\frac{2\lambda n}{w + \frac{k^2\delta}{2}}} N(\delta, 4\lambda n\sigma^2) \geq \sqrt{\lambda} N(\delta, 4\lambda n\sigma^2)$$

$$N \left(t - \frac{w}{k^2}, \frac{1}{2} \left(w + \frac{k^2\delta}{2} \right) \sigma^2 \right) \geq \sqrt{\frac{2\lambda n}{w + \frac{k^2\delta}{2}}} N \left(t - \frac{w}{k^2}, \lambda n\sigma^2 \right) \geq \sqrt{\lambda} N \left(t - \frac{w}{k^2}, \lambda n\sigma^2 \right)$$

Applying the variable change to (13.3) together with the above Gaussian relations, we obtain :

$$F_{r,s,u,v} \geq \lambda \int_{\delta \in \mathbb{R}} N(\delta, 4\lambda n\sigma^2) \int_{w \in J(\delta, \lambda)} \left(\int_{t \in \mathbb{R}} \left(\omega_{r,s}(t + \delta) - \omega_{r,s}(t) \right)^2 N \left(t - \frac{w}{k^2}, \lambda n\sigma^2 \right) dt \right) P \left(u, v, w + \frac{k^2\delta}{2} \right) dw d\delta \quad (13.4)$$

On the other hand, by applying the change of variable, for any :

$$\begin{pmatrix} w \\ t \end{pmatrix} \mapsto \begin{pmatrix} w - k^2 \delta \\ t + \delta \end{pmatrix}$$

we obtain the two equalities for $F_{r,s,u,v}$:

$$\begin{aligned} F_{r,s,u,v} &= \int_{w \in \mathbb{R}} \left(\int_{t \in \mathbb{R}} \left(\omega_{r,s}(t + \delta) - \frac{w}{nk} - \frac{k\delta}{n} \right)^2 N \left(t - \frac{w}{k^2}, (w + k^2 \delta) \sigma^2 \right) dt \right) P(u, v, w + k^2 \delta) dw \\ &= \int_{\delta \in \mathbb{R}} N(\delta, 4\lambda n \sigma^2) \int_{w \in \mathbb{R}} \left(\int_{t \in \mathbb{R}} \left(\omega_{r,s}(t + \delta) - \frac{w}{nk} - \frac{k\delta}{n} \right)^2 N \left(t - \frac{w}{k^2}, (w + k^2 \delta) \sigma^2 \right) dt \right) P(u, v, w + k^2 \delta) dw d\delta \end{aligned} \quad (I3.5)$$

And trivially :

$$\begin{aligned} F_{r,s,u,v} &= \int_{\delta \in \mathbb{R}} N(\delta, 4\lambda n \sigma^2) \int_{w \in \mathbb{R}} \left(\int_{t \in \mathbb{R}} \left(\omega_{r,s}(t) - \frac{w}{nk} \right)^2 N \left(t - \frac{w}{k^2}, w \sigma^2 \right) dt \right) P(u, v, w) dw d\delta \end{aligned} \quad (I3.6)$$

As before we lower bound the involved Gaussian distributions, for $\in J(\delta, \lambda)$:

$$\begin{aligned} N \left(t - \frac{w}{k^2}, (w + k^2 \delta) \sigma^2 \right) &\geq \sqrt{\frac{\lambda n}{w + k^2 \delta}} N \left(t - \frac{w}{k^2}, \lambda n \sigma^2 \right) \geq \sqrt{\frac{\lambda}{2}} N \left(t - \frac{w}{k^2}, \lambda n \sigma^2 \right) \\ N \left(t - \frac{w}{k^2}, w \sigma^2 \right) &\geq \sqrt{\frac{\lambda n}{w}} N \left(t - \frac{w}{k^2}, \lambda n \sigma^2 \right) \geq \sqrt{\lambda} N \left(t - \frac{w}{k^2}, \lambda n \sigma^2 \right) \end{aligned}$$

(I3.5) and (I3.6) are thus lower-bounded in respectively :

$$\begin{aligned} F_{r,s,u,v} &\geq \sqrt{\frac{\lambda}{2}} \int_{\delta \in \mathbb{R}} N(\delta, 4\lambda n \sigma^2) \int_{w \in J(\delta, \lambda)} \left(\int_{t \in \mathbb{R}} \left(\omega_{r,s}(t + \delta) - \frac{w}{nk} - \frac{k\delta}{n} \right)^2 N \left(t - \frac{w}{k^2}, \lambda n \sigma^2 \right) dt \right) P(u, v, w + k^2 \delta) dw d\delta \end{aligned} \quad (I3.7)$$

$$\begin{aligned} F_{r,s,u,v} &\geq \sqrt{\lambda} \int_{\delta \in \mathbb{R}} N(\delta, 4\lambda n \sigma^2) \int_{w \in J(\delta, \lambda)} \left(\int_{t \in \mathbb{R}} \left(\omega_{r,s}(t) - \frac{w}{nk} \right)^2 N \left(t - \frac{w}{k^2}, \lambda n \sigma^2 \right) dt \right) P(u, v, w) dw d\delta \end{aligned} \quad (I3.8)$$

For any X, Y and $a, b > 0$, we have :

$$aX^2 + bY^2 \geq \frac{ab}{(a+b)} (X - Y)^2 \geq \min(a, b) (X - Y)^2$$

By applying this to (I3.7) and (I3.8) we obtain :

$$F_{r,s,u,v} \geq \sqrt{\frac{\lambda}{2}} \int_{\delta \in \mathbb{R}} N(\delta, 4\lambda n \sigma^2) \int_{w \in J(\delta, \lambda)} \left(\int_{t \in \mathbb{R}} \left(\omega_{r,s}(t + \delta) - \omega_{r,s}(t) - \frac{k\delta}{n} \right)^2 N\left(t - \frac{w}{k^2}, \lambda n \sigma^2\right) dt \right) \min(P(u, v, w), P(u, v, w + k^2\delta)) dw d\delta \quad (I3.9)$$

Then we apply a second time the inequality, this time to (I3.4) and (I3.9) to finally discard ω in the lower bound of $F_{r,s,u,v}$:

$$F_{r,s,u,v} \geq \min\left(\lambda, \sqrt{\frac{\lambda}{2}}\right) \int_{\delta \in \mathbb{R}} \frac{k^2 \delta^2}{n^2} N(\delta, 4\lambda n \sigma^2) \int_{w \in J(\delta, \lambda)} \min\left(P(u, v, w), P\left(u, v, w + \frac{k^2 \delta}{2}\right), P(u, v, w + k^2\delta)\right) dw d\delta \quad (I3.10)$$

which ends Part 2.

Part 3.

In this last but not least part, we will lower bound the minimum appearing in (I3.10), by showing that $P(u, v, w)$ is slowly varying in w , slowly compared to a variation of $k^2\delta$, when probability distributions Φ and Φ' verify the separable criteria (e.g. are in $\zeta(\alpha)$), with at least one of them, say Φ' , being sufficiently sleek, as stated in the hypothesis of the lemma.

Φ' being result of a permutative sleeking transformation, there exists a probability distribution $\Phi'' \in \zeta(\alpha)$ and a permutative sleeking kernel γ (that we will carefully choose) such that :

$$\Phi'(y) = \sum_{\sigma \in \mathfrak{S}_n} \gamma(|\sigma|) \Phi'' \circ \sigma(y)$$

And then (restricting x, y to $\{0,1\}^n$ instead of $[0,1]^n$) :

$$\begin{aligned} P(u, v, w) &= \sum_{x, y \in \mathcal{K}(u, v, w)} \Phi(x) \sum_{\sigma \in \mathfrak{S}_n} \gamma(|\sigma|) \Phi'' \circ \sigma(y) = \sum_{x, z \in J(u) \times J(v)} \left(\sum_{\substack{\sigma \in \mathfrak{S}_n \\ \sigma(x).z=w}} \gamma(|\sigma|) \right) \Phi(x) \Phi''(z) \\ &= \sum_{x, z \in J(u) \times J(v)} \left(\sum_{l=0}^n \gamma(l) \sum_{\substack{\sigma \in \mathfrak{S}_n \\ \sigma(x).z=w \\ |\sigma|=l}} 1 \right) \Phi(x) \Phi''(z) \\ &= \sum_{x, z \in J(u) \times J(v)} \left(\sum_{l=0}^n \gamma(l) \sum_{\substack{x' \in J(u) \\ x'.z=w}} \left(\sum_{\substack{\sigma \in \mathfrak{S}_n \\ \sigma(x)=x' \\ |\sigma|=l}} 1 \right) \right) \Phi(x) \Phi''(z) \quad (I3.11) \end{aligned}$$

It is easy to verify that, for any $\sigma, \mu \in \mathfrak{S}_n$:

$$|\sigma| = |\mu \circ \sigma \circ \mu^{-1}|$$

as shown in the proof of Proposition 5. We set :

$$\theta_u(l, \|x - x'\|^2) = \sum_{\substack{\sigma \in \mathfrak{S}_n \\ \sigma(x)=x' \\ |\sigma|=l}} 1, \quad \forall x, x' \in \mathcal{J}(u)$$


This writing can be justified thanks to the above by observing that, $\forall \mu \in \mathfrak{S}_n$:

$$\sum_{\substack{\sigma \in \mathfrak{S}_n \\ \sigma \circ \mu(x)=\mu(x') \\ |\sigma|=l}} 1 = \sum_{\substack{\sigma \in \mathfrak{S}_n \\ \mu^{-1} \circ \sigma \circ \mu(x)=x' \\ |\mu^{-1} \circ \sigma \circ \mu|=l}} 1 = \sum_{\substack{\sigma' \in \mathfrak{S}_n \\ \sigma'(x)=x' \\ |\sigma'|=l}} 1$$

$\sigma \mapsto \mu^{-1} \circ \sigma \circ \mu$ being a bijective transformation in \mathfrak{S}_n . The explicit calculation of $\theta_u(l, s)$ can be obtained but is not necessary for the proof. We can then write (I3.11) as follows :

$$P(u, v, w) = \sum_{x, z \in \mathcal{J}(u) \times \mathcal{J}(v)} \left(\sum_{l=0}^n \gamma(l) \sum_{s=0}^l \theta_u(l, s) \left(\sum_{\substack{x' \in \mathcal{J}(u) \\ x'.z=w \\ \|x-x'\|^2=s}} 1 \right) \right) \Phi(x) \Phi''(z) \quad (I3.12)$$

The quantity in the internal brackets of the above can be calculated by considering the drawing below :



x' can be considered as partitionned in zones a, b, c, d , with the following constraints :

$$\begin{cases} 0 \leq a = w - b \leq v - x.z \\ 0 \leq c = u - \frac{s}{2} - b \leq u - x.z \\ 0 \leq d = b + \frac{s}{2} - w \leq n - u - v + x.z \\ 0 \leq b \leq x.z \end{cases}$$

Let's denote :

$$b^-(v, w, x.z, s) \triangleq \max\left(0, w - v + x.z, x.z - \frac{s}{2}, w - \frac{s}{2}\right)$$

$$b^+(u, v, w, x.z, s) \triangleq \min\left(x.z, w, u - \frac{s}{2}, n - u - v + x.z + w - \frac{s}{2}\right)$$

$$F_{u,v,w,x.z,s}(b) \triangleq \binom{v-x.z}{w-b} \binom{x.z}{b} \binom{u-x.z}{u-s/2-b} \binom{n-u-v+x.z}{b+s/2-w}$$

Then, the counting of the quantity in the internal brackets of (I3.12) gives, with the help of the above drawing :

$$M(u, v, w, x, z, s) \triangleq \sum_{\substack{x' \in J(u) \\ x'.z=w \\ \|x-x'\|^2=s}} 1 = \sum_{b=b^-(v,w,x,z,s)}^{b^+(u,v,w,x,z,s)} F_{u,v,w,x,z,s}(b)$$

It can be shown with no particular difficulty that $b^-(v, w, x, z, s) \leq b^+(u, v, w, x, z, s)$. The boundaries being non empty imply that :

$$\max(x.z - s/2, w - s/2) \leq \min(x.z, w) \Rightarrow s/2 \geq |w - x.z| \quad (I3.13)$$

Let's set also :

$$K_\gamma(u, s) \triangleq \sum_{l=s}^n \gamma(l) \theta_u(l, s)$$

$$\begin{aligned} \Delta_\gamma(\Phi, \Phi'') &= E[(w - x.z)^2] = \sum_{\sigma \in \mathbb{G}_n} \gamma(|\sigma|) \int_{x,y \in [0,1]^n} (x.y - x.\sigma(y))^2 \Phi(x) \Phi''(y) dx dy \\ &= \sum_{s=0}^n \gamma(s) A_n^s \left(\frac{s^2}{e} \Delta_0(\Phi, \Phi'') + O(n) \right) \end{aligned}$$

The last relation coming from Lemma 2.

$P(u, v, w)$ can be expressed with those notations :

$$P(u, v, w) = \sum_{x,z \in J(u) \times J(v)} \sum_{s=0}^n K_\gamma(u, s) M(u, v, w, x, z, s) \Phi(x) \Phi''(z)$$

Injecting this relation into (I3.10), and denoting :

$$M_p = M_p(u, v, w, x, z, s) \triangleq M\left(u, v, w + p \frac{k^2 \delta}{2}, x, z, s\right), \quad p \in \mathbb{N}$$

we obtain the intermediary lower bound (from (I3.2') and (I3.10)) :

$$\begin{aligned} \langle \omega, \Phi, \Phi' \rangle &\geq C_3 \min\left(\lambda, \sqrt{\frac{\lambda}{2}}\right) \sum_{\substack{u,v,w \in \mathbb{N}_n \\ w \in \hat{J}(\lambda)}} \sum_{x,z \in J(u) \times J(v)} \sum_{s=0}^n K_\gamma(u, s) \left(\int_{|\delta| \leq \lambda n / k^2} \frac{k^2 \delta^2}{n^2} N(\delta, \right. \\ &\quad \left. 4\lambda n \sigma^2) \min(M_0, M_1, M_2) d\delta \right) \Phi(x) \Phi''(z) \end{aligned} \quad (I3.14)$$

where $\hat{J}(\lambda) = \{w | w \geq 3\lambda n\}$; indeed, if $|\delta| \leq \lambda n / k^2$, then $J(\delta, \lambda) \supset \hat{J}(\lambda)$.

We will now concentrate on $\min(M_0, M_1, M_2)$, to discard the minimum with lower bound by a constant under the hypothesis of the Lemma.

We set the following conditions under simplex form (their role will appear later) :

$$\begin{aligned}
& \left. \begin{aligned}
& P_0(\lambda) = \{w | w \geq 3\lambda n\} & (1) \\
& P_1(\theta) = \{w, x, z | (w - x.z)^2 \geq \theta^2 \Delta_\gamma\} & (2) \\
& P_2(\varepsilon) = \left\{ u, v, w, x, z, s \left| \left(x.z - w - \frac{s}{2} \left(\frac{x.z}{u} - \frac{v - x.z}{n - u} \right) \right)^2 \leq \varepsilon \Delta_\gamma \right. \right\} & (3) \\
& P_3(\theta') = \left\{ u, v, w, x, z \left| \min \left(P_{u,v} \left(\frac{w}{n} \right), P_{u,v} \left(\frac{x.z}{n} \right) \right) \geq \theta' \right. \right\} & (4) \\
& P_4(n\varepsilon', \theta'') = \left\{ u, v, w, x, z \left| \eta \left(u, v, x.z, \frac{n\varepsilon'}{2} \right) \geq \theta'' \right. \right\} & (5)
\end{aligned} \right\} (\text{II})
\end{aligned}$$

where :

$$\eta(u, v, x.z, s/2) \triangleq \min \left(\frac{(u - s/2)(u - x.z)x.z}{u^3}, \frac{(n - u - s/2)(v - x.z)(n - u - v + x.z)}{(n - u)^3} \right)$$

From (l3.14), we can write the weaker lower bound :

$$\begin{aligned}
\langle \omega, \Phi, \Phi' \rangle \geq & C_3 \min \left(\lambda, \sqrt{\frac{\lambda}{2}} \right) \sum_{\substack{u, v, w \in \mathbb{N}_n \\ x, z \in \mathcal{J}(u) \times \mathcal{J}(v) \\ P_0, P_1, P_2, P_3, P_4}} \sum_{s=0}^n K_\gamma(u, s) \left(\int_{|\delta| \leq \lambda n / k^2} \frac{k^2 \delta^2}{n^2} N(\delta, \right. \\
& \left. 4\lambda n \sigma^2) \min(M_0, M_1, M_2) d\delta \right) \Phi(x) \Phi''(z) & (l3.15)
\end{aligned}$$

The condition P_3 comes from the examination of the distance between the unique pick of $F_{u,v,w,x,z,s}(b)$ and the boundaries of the summing $b^-(v, w, x.z, s)$ and $b^+(u, v, w, x.z, s)$. The extrema of F verify (with error lesser than a unit) :

$$\frac{(w - b)(x.z - b)(u - b - \frac{s}{2})(n - u - v + x.z + w - b - \frac{s}{2})}{b(v - x.z - w + b)(b + \frac{s}{2} - x.z)(b + \frac{s}{2} - w)} = 1 \quad (l3.16)$$

obtained by writing $\frac{F_{u,v,w,x,z,s}(b+1)}{F_{u,v,w,x,z,s}(b)} = 1$.

The boundaries of the summing M imply that :

$$b \in \left[w - \frac{s}{2}, w \right] \cap \left[x.z - \frac{s}{2}, x.z \right]$$

One observes that there exists only one unique zero of (l3.16) within the boundaries. Indeed, the numerator is a decreasing function within the boundaries, because its four monomials are positive and decreasing within. On the other hand the denominator is increasing because its four monomials are positive and increasing within. Thus there exists a unique zero. In addition, that zero is a maximum of F because F is increasing at righthand neighborhood of the lower boundary b^- (the fraction is there $+\infty > 1$).

Let b_0 be the maximum. It can be shown without any particular difficulty (we won't present the useless details here) that :

$$\min_{\varepsilon \in \{+, -\}} d(b_0, b^\varepsilon) \geq \frac{s}{16} \min \left(R_{u,v}(w, x.z), R_{u,v}(x.z, w), R_{u,v}^{-1}(w, x.z), R_{u,v}^{-1}(x.z, w) \right)$$

where :

$$R_{u,v}(X, Y) \triangleq \frac{(u - Y)(n - u - v + X)}{Y(v - X)}$$

It is easy to remark that :

$$P_3(\theta') \Rightarrow \min \left(R_{u,v}(w, x, z), R_{u,v}(x, z, w), R_{u,v}^{-1}(w, x, z), R_{u,v}^{-1}(x, z, w) \right) \geq \theta'$$

The strategy for examining $\min(M_0, M_1, M_2)$ is to put F under the shape :

$$F(b) = G(b - \rho w)H(w)(1 + o(1))$$

where G is a gaussian distribution with ρ independant from b and w , and H is a gaussian distribution whose filter's width is large compared to $k^2\delta \sim k\sqrt{\lambda n}$. Indeed then H varies slowly by translation of length $k^2\delta$ and a variation of w in G causes an indicial translation of length $\sim \rho k^2\delta$ in the summing in M , and this length can be shown as remaining small compared to $\min_{\epsilon \in \{+, -\}} d(b_0, b^\epsilon)$.

To that extent, we consider the following grouping in :

$$F_{u,v,w,x,z,s}(b) = \left(\binom{x.z}{b} \binom{u - x.z}{u - s/2 - b} \right) \left(\binom{v - x.z}{w - b} \binom{n - u - v + x.z}{b + s/2 - w} \right)$$

Let's first remark that, for any integers m, n, p, q, t , such that :

$$\max(-m + p, -q) \leq t \leq \min(p, n)$$

we have, $\forall \zeta \in]0, 1[$:

$$\begin{aligned} \binom{m}{p-t} \binom{n}{q+t} &= \frac{\zeta^{p-t} (1-\zeta)^{m-p+t} \zeta^{q+t} (1-\zeta)^{m-q-t}}{\zeta^{p+q} (1-\zeta)^{m+n-(p+q)}} \binom{m}{p-t} \binom{n}{q+t} \\ &= \frac{1}{\zeta^{p+q} (1-\zeta)^{m+n-(p+q)}} \beta_{p-t, m}(\zeta) \beta_{q+t, n}(\zeta) \end{aligned}$$

We choose ζ , in such a manner that the picks in t of the above binomial distributions coincide in a common pick in t_0 . This leads to :

$$\begin{cases} p - t_0 = \alpha m \\ q + t_0 = \alpha n \end{cases}$$

which results in $\zeta = \frac{p+q}{m+n}$ and $t_0 = \frac{np-mq}{m+n}$.

Then, from Proposition 4 at the neighborhood of t_0 we get :

$$\binom{m}{p-t} \binom{n}{q+t} \sim \frac{1}{2\pi \zeta^{p+q+1} (1-\zeta)^{m+n-(p+q)-1} \sqrt{mn}} e^{-\frac{m+n}{2\zeta(1-\zeta)mn} \left(t - \frac{np-mq}{m+n}\right)^2}$$

We apply this approximation to both grouping in F , and we denote b_1 and b_2 the respective picks of both grouping. We have :

$$b_1 - b_2 = x.z - w - \frac{s}{2} \left(\frac{x.z}{u} - \frac{v - x.z}{n - u} \right)$$

We determine now the permutative sleeking kernel as follows :

$$\gamma(l) \triangleq \frac{1}{A_n^l} \Gamma(\varepsilon' n, l)$$

where $\Gamma(a, b) = \begin{cases} 1 & \text{if } a = b \\ 0 & \text{if } a \neq b \end{cases}$. By using the same technique than in Lemma 2 to perform the calculations, we obtain (calculations are a bit painful but present no higher difficulty than in Lemma 2, it is required that $x, z \in \{0,1\}^n$ to get the result) that there exists a constant $C_4 > 0$ such that :

$$E \left[\left(x.z - w - \frac{s}{2} \left(\frac{x.z}{u} - \frac{v - x.z}{n - u} \right) \right)^2 \right] \leq C_4 \varepsilon'^2 n \quad (13.17)$$

This important relation justifies the choice of condition P_2 , that thus implies :

$$(b_1 - b_2)^2 \leq \varepsilon \Delta_\gamma$$

ε is chosen such that :

$$C_4 \varepsilon'^2 n \ll \varepsilon \Delta_\gamma \ll \theta \theta'' \sqrt{\Delta_\gamma} \quad (13.17')$$

which can be made possible (see discussion on parameters at the end of the proof), and enables us to obtain, from Proposition 4, the simultaneous validity of the Gaussian approximations for the two grouping within F . Indeed, the filter's width for them are respectively :

$$\begin{cases} \frac{s(u - s/2)(u - x.z)x.z}{u^3} & (\triangleq a^{-1}) \\ \frac{s(n - u - s/2)(v - x.z)(n - u - v + x.z)}{(n - u)^3} & (\triangleq a'^{-1}) \end{cases}$$

that are both of superior to $\theta \theta'' \sqrt{\Delta_\gamma}$ under condition $P_1(\theta)$ and $P_4(\theta'')$.

We then use the identity :

$$a(t - c)^2 + a'(t - c')^2 = (a + a') \left(t - \frac{ac + a'c'}{a + a'} \right)^2 + \frac{aa'(c - c')^2}{a + a'}$$

to reconstruct the two Gaussian under the shape :

$$G(b - \rho w)H(w)$$

We find :

$$\rho = \left(1 + \frac{u^3(n - u - s/2)(v - x.z)(n - u - v + x.z)}{(n - u)^3(u - s/2)(u - x.z)x.z} \right)^{-1}$$

In addition, the filter's width of H equals to :

$$\begin{aligned} (a^{-1} + a'^{-1}) &= s \left(\frac{(u - s/2)(u - x.z)x.z}{u^3} + \frac{(n - u - s/2)(v - x.z)(n - u - v + x.z)}{(n - u)^3} \right) \\ &\geq 2\eta(u, v, x, z, s/2)\theta \sqrt{\Delta_\gamma} \end{aligned}$$

due to (l3.13) and (for last inequality above) under condition P_1 .

As $(c - c')^2 = (b_1 - b_2)^2 \leq \varepsilon \Delta_\gamma \ll \theta \theta'' \sqrt{\Delta_\gamma}$, we have :

$$H(w) = H(w, u, v, x, z, s) = H_0(u, v, x, z, s) e^{-\frac{aa'(c-c')^2}{a+a'}} \in \left[H_0 e^{-\frac{1}{2}}, H_0 \right]$$

When we apply the indicial translation of length $\sim \rho k^2 \delta \sim \rho k \sqrt{\lambda n}$ in the summing in M (corresponding to the variation of w in M_0, M_1, M_2), it remains very small compared the distance from the pick to the boundaries, indeed, under P_1 and P_3 (and considering (l3.13) and (l3.16')) :

$$\min_{\varepsilon \in \{+, -\}} d(b_0, b^\varepsilon) \geq \frac{1}{8} \theta \theta' \sqrt{\Delta_\gamma} \triangleq D$$

We must choose ε' so that :

$$k\sqrt{\lambda n} \ll \theta \theta' \sqrt{\Delta_\gamma} \quad (l3.17'')$$

and remaining compatible with (l3.17').

From Proposition 7 applied to the sub-part G of $F_{u,v,w,x,z,s}(b)$ within the boundaries of the summing in M we obtain, for $\varepsilon \in \{0, 1, 2\}$:

$$\sum_{b=b^-(w+p\frac{k^2\delta}{2})}^{b^+(w+p\frac{k^2\delta}{2})} G\left(b - \rho\left(w + p\frac{k^2\delta}{2}\right)\right) \geq \left(1 - \frac{(1+\rho)k^2\delta}{D}\right) \sum_{b=b^-(w)}^{b^+(w)} G(b - \rho w)$$

And finally :

$$M_p \geq e^{-\frac{1}{2}} \left(1 - \frac{(1+\rho)k^2\delta}{D}\right) M_0 \geq e^{-\frac{1}{2}} \left(1 - \frac{2k^2\delta}{D}\right) M_0$$

which also means that :

$$\min(M_0, M_1, M_2) \geq e^{-\frac{1}{2}} \left(1 - \frac{2k^2\delta}{D}\right) M_0 \quad (l3.18)$$

We note that, by choice of kernel γ , we have necessarily $s \leq n\varepsilon'$ (because otherwise $K_\gamma(u, s) = 0$), and thus $P_4(n\varepsilon', \theta'') \Rightarrow P_4(s, \theta'')$.

This ends Part 3.

In order to finish the proof, we will now deal with the conditions P_0, \dots, P_4 and discuss the values of the parameters $\varepsilon, \varepsilon', \theta, \theta', \theta'', \lambda$.

First, we can inject (I3.18) in (I3.15) to obtain :

$$\begin{aligned} \langle \omega, \Phi, \Phi' \rangle \geq & C_3 \min \left(\lambda, \sqrt{\frac{\lambda}{2}} \right) e^{-\frac{1}{2}} \left(\int_{|\delta| \leq \lambda n / k^2} \left(1 - \frac{2k^2 \delta}{D} \right) \frac{k^2 \delta^2}{n^2} N(\delta, \right. \\ & \left. 4\lambda n \sigma^2) d\delta \right) \sum_{\substack{u, v, w \in \mathbb{N}_n \\ x, z \in \mathcal{J}(u) \times \mathcal{J}(v) \\ P_0, P_1, P_2, P_3, P_4}} \sum_{s=0}^n K_\gamma(u, s) M(u, v, w, x, z, s) \Phi(x) \Phi''(z) \end{aligned} \quad (I3.19)$$

We have :

$$\begin{aligned} & \sum_{\substack{u, v, w \in \mathbb{N}_n \\ x, z \in \mathcal{J}(u) \times \mathcal{J}(v) \\ P_0, P_1, P_2, P_3, P_4}} \sum_{s=0}^n K_\gamma(u, s) M(u, v, w, x, z, s) \Phi(x) \Phi''(z) \\ \geq & \sum_{\substack{u, v, w \in \mathbb{N}_n \\ x, z \in \mathcal{J}(u) \times \mathcal{J}(v) \\ P_0, P_1, P_3, P_4}} \sum_{s=0}^n K_\gamma(u, s) M(u, v, w, x, z, s) \Phi(x) \Phi''(z) \\ - & \sum_{\substack{u, v, w \in \mathbb{N}_n \\ x, z \in \mathcal{J}(u) \times \mathcal{J}(v) \\ \overline{P_2}}} \sum_{s=0}^n K_\gamma(u, s) M(u, v, w, x, z, s) \Phi(x) \Phi''(z) \end{aligned} \quad (I3.19')$$

We can thus deal with P_0, P_1, P_3, P_4 all together on one hand, and P_2 (that is the only one of the 5 conditions depending on s) on the other hand. As already explained above, the choice of kernel γ gives :

$$K_\gamma(u, s) = \begin{cases} \theta_u(n\varepsilon', s) & \text{if } s \leq n\varepsilon' \\ 0 & \text{otherwise} \end{cases}$$

We can first remark the following dependence between the conditions P_0, P_1, P_3, P_4 . As already noticed :

$$P_3(\theta') \Rightarrow \min \left(R_{u,v}(w, x, z), R_{u,v}(x, z, w), R_{u,v}^{-1}(w, x, z), R_{u,v}^{-1}(x, z, w) \right) \geq \theta'$$

In addition, still under $P_3(\theta')$, it can easily be seen that :

$$\eta(u, v, x, z, n\varepsilon') \geq \theta'^2 - \varepsilon'$$

which implies that,

$$\text{if } \theta'^2 - \varepsilon' \geq \theta'', \text{ then } P_3(\theta') \Rightarrow P_4(n\varepsilon', \theta''). \quad (I3.20)$$

At last, under $P_1(\theta)$ and $P_3(\theta')$, $\frac{x, z}{n} \geq \theta' \Rightarrow \frac{w}{n} \geq \theta' - \left| \frac{w}{n} - \frac{x, z}{n} \right| \geq \theta' - \theta \frac{\sqrt{\Delta_\gamma}}{n}$

which implies that,

$$\text{if } \theta' - \theta \frac{\sqrt{\Delta_\gamma}}{n} \geq \lambda, \text{ then } P_1(\theta) \& P_3(\theta') \Rightarrow P_0(\lambda). \quad (I3.20')$$

This means that under conditions (I3.20) and (I3.20') on $\varepsilon', \theta, \theta', \theta'', \lambda$, we have :

$$\begin{aligned} & \sum_{\substack{u,v,w \in \mathbb{N}_n \\ x,z \in \mathcal{J}(u) \times \mathcal{J}(v) \\ P_0, P_1, P_3, P_4}} \sum_{s=0}^n K_\gamma(u, s) M(u, v, w, x, z, s) \Phi(x) \Phi''(z) \\ &= \sum_{\substack{u,v,w \in \mathbb{N}_n \\ x,z \in \mathcal{J}(u) \times \mathcal{J}(v) \\ P_1, P_3}} \sum_{s=0}^n K_\gamma(u, s) M(u, v, w, x, z, s) \Phi(x) \Phi''(z) \\ &= \sum_{\substack{u,v \in \mathbb{N}_n \\ x,z \in \mathcal{J}(u) \times \mathcal{J}(v) \\ P_3}} \sum_{w \in \mathbb{N}_n} \sum_{\sigma \in \mathfrak{S}_n} \gamma(|\sigma|) \Phi(x) \Phi''(z) \\ &\quad (w-x, z)^2 \geq \theta^2 \Delta_\gamma, \sigma(x).z=w \\ &= \sum_{\substack{u,v \in \mathbb{N}_n \\ x,z \in \mathcal{J}(u) \times \mathcal{J}(v) \\ P_3}} \sum_{\sigma \in \mathfrak{S}_n} \gamma(|\sigma|) \Phi(x) \Phi''(z) \\ &\quad (\sigma(x).z-x, z)^2 \geq \theta^2 \Delta_\gamma \\ &\geq \sum_{\substack{u,v \in \mathbb{N}_n \\ x,z \in \mathcal{J}(u) \times \mathcal{J}(v) \\ P_3}} \sum_{\sigma \in \mathfrak{S}_n} \frac{4(\sigma(x).z-x, z)^2}{n^2 \varepsilon'^2} \gamma(|\sigma|) \Phi(x) \Phi''(z) \\ &\geq \sum_{\substack{u,v \in \mathbb{N}_n \\ x,z \in \mathcal{J}(u) \times \mathcal{J}(v) \\ P_3}} \sum_{\sigma \in \mathfrak{S}_n} \min \left(P_{u,v} \left(\frac{\sigma(x).z}{n} \right), P_{u,v} \left(\frac{x, z}{n} \right) \right) \frac{4(\sigma(x).z-x, z)^2}{n^2 \varepsilon'^2} \gamma(|\sigma|) \Phi(x) \Phi''(z) \\ &- \theta' \sum_{\substack{u,v \in \mathbb{N}_n \\ x,z \in \mathcal{J}(u) \times \mathcal{J}(v) \\ P_3}} \sum_{\sigma \in \mathfrak{S}_n} \frac{4(\sigma(x).z-x, z)^2}{n^2 \varepsilon'^2} \gamma(|\sigma|) \Phi(x) \Phi''(z) \\ &\geq \sum_{\substack{u,v \in \mathbb{N}_n \\ x,z \in \mathcal{J}(u) \times \mathcal{J}(v) \\ P_3}} \sum_{\sigma \in \mathfrak{S}_n} P_{u,v} \left(\frac{\sigma(x).z}{n} \right) P_{u,v} \left(\frac{x, z}{n} \right) \frac{4(\sigma(x).z-x, z)^2}{n^2 \varepsilon'^2} \gamma(|\sigma|) \Phi(x) \Phi''(z) - \frac{4\theta^2 \Delta_\gamma}{n^2 \varepsilon'^2} \\ &- \theta' \sum_{\substack{u,v \in \mathbb{N}_n \\ x,z \in \mathcal{J}(u) \times \mathcal{J}(v) \\ P_3}} \sum_{\sigma \in \mathfrak{S}_n} \frac{4(\sigma(x).z-x, z)^2}{n^2 \varepsilon'^2} \gamma(|\sigma|) \Phi(x) \Phi''(z) \end{aligned}$$

where the first inequality comes from the fact that $2|w-x, z| \leq s \leq n\varepsilon'$ as expressed in (I3.13). In the second inequality, the first term can be lower-bounded thanks to Corollary 1, and the third term can be upper bounded by $4\theta'$.

To deal with the first term, we replace Φ'' by Φ''' as follows :

$$\Phi''(y) = \sum_{\sigma \in \mathfrak{S}_n} \delta(|\sigma|) \Phi''' \circ \sigma(y)$$

where δ is the permutative sleeking transformation introduced in Corollary 1.

We thus get then from Corollary 1 :

$$\begin{aligned}
& \sum_{\substack{u,v,w \in \mathbb{N}_n \\ x,z \in \mathcal{J}(u) \times \mathcal{J}(v) \\ P_0, P_1, P_3, P_4}} \sum_{s=0}^n K_\gamma(u, s) M(u, v, w, x, z, s) \Phi(x) \Phi''(z) \\
& \geq K(\alpha) - \frac{\varepsilon'}{\sqrt{e}} - O\left(\frac{1}{\sqrt{n}}\right) - \frac{4\theta^2 \Delta_\gamma}{n^2 \varepsilon'^2} - 4\theta' \tag{I3.21}
\end{aligned}$$

It remains now to upper bound the term conditioned by P_2 as appearing in (I3.19'), this can be easily done thanks to Markov inequality :

$$\begin{aligned}
& \sum_{\substack{u,v,w \in \mathbb{N}_n \\ x,z \in \mathcal{J}(u) \times \mathcal{J}(v) \\ \overline{P_2}}} \sum_{s=0}^n K_\gamma(u, s) M(u, v, w, x, z, s) \Phi(x) \Phi''(z) \\
& = \sum_{\substack{u,v,w \in \mathbb{N}_n \\ x,z \in \mathcal{J}(u) \times \mathcal{J}(v) \\ \overline{P_2}}} \sum_{s=0}^{n\varepsilon'} \theta_u(n\varepsilon', s) M(u, v, w, x, z, s) \Phi(x) \Phi''(z) \\
& \leq \sum_{\substack{u,v,w \in \mathbb{N}_n \\ x,z \in \mathcal{J}(u) \times \mathcal{J}(v) \\ \overline{P_2}}} \sum_{s=0}^{n\varepsilon'} \frac{\left(x, z - w - \frac{s}{2} \left(\frac{x, z}{u} - \frac{v - x, z}{n - u}\right)\right)^2}{\varepsilon \Delta_\gamma} \theta_u(n\varepsilon', s) M(u, v, w, x, z, s) \Phi(x) \Phi''(z) \\
& \leq \frac{C_4 \varepsilon'^2 n}{\varepsilon \Delta_\gamma}
\end{aligned}$$

where last inequality comes from (I3.17) ; we know that $\frac{C_4 \varepsilon'^2 n}{\varepsilon \Delta_\gamma} \ll 1$ from (I3.17').

As long as $\lambda \gg \frac{k^2}{n}$, we can write :

$$\begin{aligned}
\int_{|\delta| \leq \lambda n / k^2} \frac{k^2 \delta^2}{n^2} N(\delta, 4\lambda n \sigma^2) d\delta & = \int_{\mathbb{R}} \frac{k^2 \delta^2}{n^2} N(\delta, 4\lambda n \sigma^2) d\delta - \triangleleft \left(e^{-\frac{\lambda n}{8k^2}} \right) = \frac{4\lambda}{n} \left(1 - \frac{1}{k^2} \right) - \\
& \triangleleft \left(e^{-\frac{\lambda n}{8k^2}} \right)
\end{aligned}$$

Finally, we get from (I3.19), (I3.19'), (I3.21) and the above :

$$\begin{aligned}
\langle \omega, \Phi, \Phi' \rangle & \geq C_3 \min \left(\lambda, \sqrt{\frac{\lambda}{2}} \right) e^{-\frac{1}{2}} \left(1 - \frac{2k\sqrt{\lambda n}}{D} \right) \left(\frac{4\lambda}{n} \left(1 - \frac{1}{k^2} \right) - e^{-\frac{\lambda n}{8k^2}} \right) \left(K(\alpha) - \frac{\varepsilon'}{\sqrt{e}} - O\left(\frac{1}{\sqrt{n}}\right) \right. \\
& \quad \left. - \frac{4\theta^2 \Delta_\gamma}{n^2 \varepsilon'^2} - 4\theta' - \frac{C_4 \varepsilon'^2 n}{\varepsilon \Delta_\gamma} \right) \tag{I3.22}
\end{aligned}$$

And we obtained from Lemma 2 and Corollary 1 the lower/upper bound for Δ_γ :

$$\begin{aligned} \frac{n^2 \varepsilon'^2}{e} + O(n) &\geq \Delta_\gamma = \sum_{s=0}^n \gamma(s) A_n^s \left(\frac{s^2}{e} \Delta_0(\Phi, \Phi'') + O(n) \right) \geq \frac{n^2 \varepsilon'^2}{e} (\Delta_0(\Phi, \Phi''') - 2l_0(\alpha)) + O(n) \\ &\geq \frac{n^2 \varepsilon'^2}{e} \left(\frac{\alpha^2}{16} - 2l_0(\alpha) \right) + O(n) \end{aligned}$$

This inequality (l3.22), that gives the result of Lemma 3, can only stand if the following simplex conditions can be satisfied ((l3.17'), ((l3.17''), (l3.20), (l3.20')) :

$$(S): \begin{cases} \varepsilon' \ll K(\alpha) & (1) \\ \frac{1}{\sqrt{n}} \ll K(\alpha) & (2) \\ \theta^2 \ll K(\alpha) & (3) \\ \theta' \ll K(\alpha) & (4) \\ n^2 \varepsilon'^2 \gg n & (5) \\ \frac{n \varepsilon'^2}{\varepsilon \Delta_\gamma} \ll K(\alpha) & (6) \\ \theta'^2 - \varepsilon' \geq \theta'' & (7) \\ \theta' - \theta \frac{\sqrt{\Delta_\gamma}}{n} \geq \lambda & (8) \\ k\sqrt{\lambda n} \ll \theta \theta' \sqrt{\Delta_\gamma} & (9) \\ \varepsilon \sqrt{\Delta_\gamma} \ll \theta \theta'' \sqrt{\Delta_\gamma} & (10) \end{cases}$$

Replacing Δ_γ by its lower/upper bound above, it can be easily seen that those conditions can be satisfied. $\theta, \theta', \theta''$ can be chosen to satisfy (3), (4) and (7) (with ε' small compared to them), then λ is chosen to satisfy (8), then ε' is chosen sufficiently small so that ε can exist (it is sufficient that $\varepsilon' \ll \theta \theta'' \left(\frac{\alpha^2}{16} - 2l_0(\alpha) \right) K(\alpha)$ by gathering (6) and (10), thanks to which ε can be found in $\frac{1}{n \left(\frac{\alpha^2}{16} - 2l_0(\alpha) \right) K(\alpha)} \ll \varepsilon \ll \frac{\theta \theta''}{n \varepsilon'}$), and finally n is chosen sufficiently large to match (2), (5), (8) and (9).

The permutative sleeking transformation to adopt is (with notation of Corollary 1) :

$$\Phi' = T_\gamma \circ T_\delta[\Phi''']$$

and Proposition 5 confirms that such composition is still a permutative sleeking transformation. ■

Corollary 3.

Under the conditions of Lemma 3, there exists $C_2 > 0$ such that

$$P\left((V_A - \omega)^2 \geq \frac{C(\alpha)}{2n} \left(1 - \sqrt{\frac{1}{kC(\alpha)}}\right)^2\right) \geq \frac{C_2}{\ln n}$$

■ Let's consider the strategy $\tilde{\omega}$ derived from ω , and defined as follows (with chosen $\lambda > C$):

$$\begin{cases} \text{if } \left(\omega - \frac{ki.j}{n}\right)^2 \geq \frac{\lambda}{n} \text{ then } \tilde{\omega} = \frac{ki.j}{n} \triangleq V_0 \\ \text{else } \tilde{\omega} = \omega \end{cases}$$

It is clear that $\tilde{\omega} \in \Omega_{\#}$ as ω also does. Thus, from Lemma 3, we have:

$$E\left[\left(\tilde{\omega} - \frac{x.y}{kn}\right)^2\right] \geq \frac{C}{n}$$

and, as $E\left[\left(V_A - \frac{x.y}{kn}\right)^2\right] = \text{Var}(V_A)^2 \leq \frac{1}{nk}$ as shown in Section II, we have by Proposition 6:

$$E[(\tilde{\omega} - V_A)^2] \geq \frac{C}{n} \left(1 - \sqrt{\frac{1}{kC}}\right)^2 \triangleq \frac{C'}{n} \quad (I.1)$$

Then, by writing:

$$(\tilde{\omega} - V_A)^2 \leq \frac{1}{2}(\tilde{\omega} - V_0)^2 + \frac{1}{2}(V_0 - V_A)^2 \leq \frac{1}{2}\frac{\lambda}{n} + \frac{1}{2}(V_0 - V_A)^2$$

we get:

$$P\left((V_A - \tilde{\omega})^2 \geq \frac{\lambda}{n}\right) \leq P\left(\frac{1}{2}\frac{\lambda}{n} + \frac{1}{2}(V_0 - V_A)^2 \geq \frac{\lambda}{n}\right) = P\left((V_0 - V_A)^2 \geq \frac{\lambda}{n}\right) \quad (I.2)$$

Now we decompose the summing set of $E[(\tilde{\omega} - V_A)^2]$ as partitioned in 3:

$$\begin{aligned} U_1 &\triangleq \left\{ \Sigma \mid (\tilde{\omega} - V_A)^2 < \frac{C'}{2n} \right\} \\ U_2 &\triangleq \left\{ \Sigma \mid (\tilde{\omega} - V_A)^2 \in \left[\frac{C'}{2n}, \frac{\lambda}{n} \right] \right\} \\ U_3 &\triangleq \left\{ \Sigma \mid (\tilde{\omega} - V_A)^2 > \frac{\lambda}{n} \right\} \end{aligned}$$

and we can write, using (I.1) and (I.2):

$$\begin{aligned} \frac{C'}{n} &\leq E[(\tilde{\omega} - V_A)^2] = \int_{U_1} (\tilde{\omega} - V_A)^2 d_\Sigma + \int_{U_2} (\tilde{\omega} - V_A)^2 d_\Sigma + \int_{U_3} (\tilde{\omega} - V_A)^2 d_\Sigma \\ &\leq \frac{C'}{2n} + \frac{\lambda}{n} P\left((V_A - \tilde{\omega})^2 \geq \frac{C'}{2n}\right) + P\left((V_0 - V_A)^2 \geq \frac{\lambda}{n}\right) \end{aligned}$$

which implies:

$$P\left((V_A - \tilde{\omega})^2 \geq \frac{C'}{2n}\right) \geq \frac{C' - 2nP\left((V_0 - V_A)^2 \geq \frac{\lambda}{n}\right)}{2\lambda}$$

Finally, provided that $C' < 2\lambda$, we have the following obvious logical statement:

$$\left\{(V_A - \omega)^2 \leq \frac{C'}{2n}\right\} \Rightarrow \left\{(V_A - \omega)^2 \leq \frac{C'}{2n}\right\}_{\tilde{\omega} = \omega} \Rightarrow \left\{(V_A - \tilde{\omega})^2 \leq \frac{C'}{2n}\right\}$$

and therefore:

$$P\left((V_A - \omega)^2 \leq \frac{C'}{2n}\right) \leq P\left((V_A - \tilde{\omega})^2 \leq \frac{C'}{2n}\right) \leq 1 - \frac{C' - 2nP\left((V_0 - V_A)^2 \geq \frac{\lambda}{n}\right)}{2\lambda}$$

$$P\left((V_A - \omega)^2 \geq \frac{C'}{2n}\right) \geq \frac{C' - 2nP\left((V_0 - V_A)^2 \geq \frac{\lambda}{n}\right)}{2\lambda}$$

We conclude by using a similar upper-bound than the one given in Proposition 3, giving:

$$P\left((V_0 - V_A)^2 \geq \frac{\lambda}{n}\right) \leq ne^{-\frac{\lambda}{2}}$$

with clearly:

$$\sup_{\lambda > c} \left(\frac{C' - 2n^2 e^{-\frac{\lambda}{2}}}{2\lambda}\right) = \frac{C_2}{\ln n}$$

■

Theorem 1.

Under the Deep Random assumption, $\mathcal{P}(\alpha, n, k, K, L)$ satisfies condition (P).

■ In order to achieve the proof, we will establish that the communication model induced by use of protocol \mathcal{P} is equivalent to a model where the legitimate partner B and the opponent ξ receives A 's digit flow through partially independent BSC. Thus, thanks to error correcting methods, such model can enable legitimate partners to create Advantage Distillation as shown in Proposition 13.

We will first determine a restricted form for the opponent's strategy by considering 3 symmetries applicable to the knowledge of ξ and then applying accordingly the Deep Random Assumption with those 3 symmetries.

The first symmetry Σ_1 , and also the most important one, results from the fact that ξ cannot distinguish σ_Φ within (μ_1, μ_2) , and neither $\sigma_{\Phi'}$ within (μ'_1, μ'_2) knowing the public information $\mathfrak{I} = \{i, j, (\mu_1, \mu_2), (\mu'_1, \mu'_2)\}$. But nevertheless, Φ_A designating the distribution potentially chosen by A in the opponent's perspective, if ξ consider $\mu_{t^{b(1)}}$ ($b \in \{0,1\}$) as the choice of A for σ_{Φ_A} then obviously Φ_A should have μ_b as its the synchronization permutation. The first symmetry thus results from the group $\{\text{Id}, t\}$ applied to (μ_1, μ_2) and (μ'_1, μ'_2) . The second symmetry Σ_2 results from the fact that ξ cannot distinguish the actual distribution among all distributions having μ_b as synchronization permutation. The second symmetry thus results from the group \mathfrak{S}_{I_0} applied to (Φ_A, μ_1) or (Φ_A, μ_2) . Then, for ξ willing to evaluate V_A knowing the public information \mathfrak{I} , the above symmetries result into the fact that the probability distribution of $P(x, (\sigma_{\Phi_A}, \sigma_A) | \mathfrak{I})$ can be restricted under Deep Random Assumption to the set:

$$R_{\mathfrak{I}}(\Sigma_1 \times \Sigma_2) = \left\{ \frac{1}{2} \left(\frac{\chi_i \left(\frac{x}{k} \right) \Sigma(\Phi, \sigma_\Phi(I_0))(u) dx}{\int_u \chi_i \left(\frac{u}{k} \right) \Sigma(\Phi, \sigma_\Phi(I_0))(u) du} \cdot \delta_{\sigma_\Phi}(\sigma_{\Phi_A}) + \frac{\chi_i \left(\frac{x}{k} \right) \Sigma(\Psi, \sigma_d[i](I_0))(u) dx}{\int_u \chi_i \left(\frac{u}{k} \right) \Sigma(\Psi, \sigma_d[i](I_0))(u) du} \cdot \delta_{\sigma_d[i]}(\sigma_{\Phi_A}) \right) \cdot \left(\frac{1}{2} \left(\delta_{\sigma_{\Phi'}}(\sigma_A) + \delta_{\sigma'_d[j]}(\sigma_A) \right) \right) \mid \forall \Phi, \Psi \in \zeta(\alpha) \right\}$$

(the notations Σ_I are coming from notations of Proposition 14).

The third symmetry Σ_3 results from the fact that the roles of A and B are symmetric in the protocol, and thus ξ cannot distinguish them. It is given by:

$$R_{\mathfrak{I}}(\Sigma_3) = \{i, (\mu_1, \mu_2) \leftrightarrow j, (\mu'_1, \mu'_2)\}$$

The third symmetry thus results from the group $\{\text{Id}, t\}$ applied to $(\{i, (\mu_1, \mu_2)\}, \{j, (\mu'_1, \mu'_2)\})$. Eventually, combining those 3 symmetries, we can restrict the strategy of ξ to estimate V_A knowing the public information $\{i, j, (\mu_1, \mu_2), (\mu'_1, \mu'_2)\}$, under Deep Random Assumption by:

$$\begin{aligned}
& \Omega_3(\Sigma_1 \times \Sigma_2 \times \Sigma_3) \\
&= \left\{ \frac{1}{4} \int_x \frac{\sigma_\Phi^{-1}(x) \cdot (\sigma_{\Phi'}^{-1}(j) + \sigma'_{d[j]^{-1}}(j)) \chi_i \left(\frac{x}{k}\right) \Sigma(\Phi, \sigma_\Phi(I_0))(x) dx}{2n \int_u \chi_i \left(\frac{u}{k}\right) \Sigma(\Phi, \sigma_\Phi(I_0))(u) du} \right. \\
&+ \frac{1}{4} \int_x \frac{\sigma_d[i]^{-1}(x) \cdot (\sigma_{\Phi'}^{-1}(j) + \sigma'_{d[j]^{-1}}(j)) \chi_i \left(\frac{x}{k}\right) \Sigma(\Psi, \sigma_d[i](I_0))(x) dx}{2n \int_u \chi_i \left(\frac{u}{k}\right) \Sigma(\Psi, \sigma_d[i](I_0))(u) du} \\
&+ \frac{1}{4} \int_y \frac{\sigma_{\Phi'}^{-1}(y) \cdot (\sigma_\Phi^{-1}(i) + \sigma_d[i]^{-1}(i)) \chi_j \left(\frac{y}{k}\right) \Sigma(\Phi, \sigma_{\Phi'}(I_0))(y) dy}{2n \int_u \chi_j \left(\frac{u}{k}\right) \Sigma(\Phi, \sigma_{\Phi'}(I_0))(u) du} \\
&+ \frac{1}{4} \int_y \frac{\sigma'_{d[j]^{-1}}(y) \cdot (\sigma_\Phi^{-1}(i) + \sigma_d[i]^{-1}(i)) \chi_j \left(\frac{y}{k}\right) \Sigma(\Psi, \sigma'_{d[j]}(I_0))(y) dy}{2n \int_u \chi_j \left(\frac{u}{k}\right) \Sigma(\Psi, \sigma'_{d[j]}(I_0))(u) du} \Big|_{\forall \Phi, \Psi} \\
&\in \zeta(\alpha) \Big\} \tag{T1.0}
\end{aligned}$$

Let's consider the 4 situations that happens with a probability $\frac{1}{4}$. We denote the partitioned situations:

$$S = S_0 \triangleq [(\sigma_A = \sigma_{\Phi'}) \& (\sigma_B = \sigma_\Phi)]$$

$$S_1 \triangleq [(\sigma_A = \sigma_{\Phi'}) \& (\sigma_B = \sigma_d[i])]$$

$$S_2 \triangleq [(\sigma_A = \sigma'_{d[j]}) \& (\sigma_B = \sigma_\Phi)]$$

$$S_3 \triangleq [(\sigma_A = \sigma'_{d[j]}) \& (\sigma_B = \sigma_d[i])]$$

We perform the demonstration in 2 steps: in the first step, we will show that there exists a constant C_1 such that

$$E \left[(V_A - V_\xi)^2 | S \right] \geq \frac{C_1}{n}$$

In the second step, we will use the above result together with an error correcting code to show that Advantage can be obtained by the legitimate partners over the opponent.

Step 1:

By denoting also, for $u \in \{0,1,2,3\}$:

$$\omega_0^* \triangleq \frac{1}{2} \left(\int_x \frac{\sigma_\Phi^{-1}(x) \cdot \sigma_{\Phi'}^{-1}(j) \chi_i \left(\frac{x}{k}\right) \Sigma_{\sigma_\Phi(I_0)}(\Phi)(x) dx}{n \int_u \chi_i \left(\frac{u}{k}\right) \Sigma_{\sigma_\Phi(I_0)}(\Phi)(u) du} + \int_y \frac{\sigma_{\Phi'}^{-1}(y) \cdot \sigma_\Phi^{-1}(i) \chi_j \left(\frac{y}{k}\right) \Sigma_{\sigma_{\Phi'}(I_0)}(\Phi)(y) dy}{n \int_u \chi_j \left(\frac{u}{k}\right) \Sigma_{\sigma_{\Phi'}(I_0)}(\Phi)(u) du} \right)$$

$$\omega_1^* \triangleq \frac{1}{2} \left(\int_x \frac{\sigma_\Phi^{-1}(x) \cdot \sigma'_{d[j]^{-1}}(j) \chi_i \left(\frac{x}{k}\right) \Sigma_{\sigma_\Phi(I_0)}(\Phi)(x) dx}{n \int_u \chi_i \left(\frac{u}{k}\right) \Sigma_{\sigma_\Phi(I_0)}(\Phi)(u) du} + \int_y \frac{\sigma_{\Phi'}^{-1}(y) \cdot \sigma_d[i]^{-1}(i) \chi_j \left(\frac{y}{k}\right) \Sigma_{\sigma_{\Phi'}(I_0)}(\Phi)(y) dy}{n \int_u \chi_j \left(\frac{u}{k}\right) \Sigma_{\sigma_{\Phi'}(I_0)}(\Phi)(u) du} \right)$$

$$\omega_2^* \triangleq \frac{1}{2} \left(\int_x \frac{\sigma_d[i]^{-1}(x) \cdot \sigma_{\Phi'}^{-1}(j) \chi_i \left(\frac{x}{k}\right) \Sigma_{\sigma_d[i](I_0)}(\Psi)(x) dx}{n \int_u \chi_i \left(\frac{u}{k}\right) \Sigma_{\sigma_d[i](I_0)}(\Psi)(u) du} + \int_y \frac{\sigma'_{d[j]^{-1}}(y) \cdot \sigma_\Phi^{-1}(i) \chi_j \left(\frac{y}{k}\right) \Sigma_{\sigma'_{d[j]}(I_0)}(\Psi)(y) dy}{n \int_u \chi_j \left(\frac{u}{k}\right) \Sigma_{\sigma'_{d[j]}(I_0)}(\Psi)(u) du} \right)$$

$$\omega_3^* \triangleq \frac{1}{2} \left(\int_x \frac{\sigma_d[i]^{-1}(x) \cdot \sigma'_d[j]^{-1}(j) \chi_i \left(\frac{x}{k} \right) \Sigma_{\sigma_d[i](I_0)}(\Psi)(x) dx}{n \int_u \chi_i \left(\frac{u}{k} \right) \Sigma_{\sigma_d[i](I_0)}(\Psi)(u) du} + \int_y \frac{\sigma'_d[j]^{-1}(y) \cdot \sigma_d[i]^{-1}(i) \chi_j \left(\frac{y}{k} \right) \Sigma_{\sigma'_d[j](I_0)}(\Psi)(y) dy}{n \int_u \chi_j \left(\frac{u}{k} \right) \Sigma_{\sigma'_d[j](I_0)}(\Psi)(u) du} \right)$$

As seen in (T1.0) the optimal strategy $\omega^*(i, j, \mu_1, \mu_2, \mu'_1, \mu'_2)$ can be written:

$$\omega^*(i, j, \mu_1, \mu_2, \mu'_1, \mu'_2) = \frac{1}{4} \sum_{u \in \{0,1,2,3\}} \omega_u^*$$

And thus, from Proposition 6:

$$E[(\omega^* - V_A)^2 | S] \geq \frac{1}{16} E[(\omega_3^* - V_A)^2 | S] \left(1 - \sqrt{\frac{E[(\omega_0^* - V_A)^2 | S]}{E[(\omega_3^* - V_A)^2 | S]}} \right)^2$$

We remark that $\forall \mu \in \mathfrak{S}_n$ $\sigma_d[\mu(i)]^{-1} \circ \mu = \sigma_d[i]^{-1}$, and therefore, $\sigma_d[i]^{-1}(i)$ is stable by action of \mathfrak{S}_n on i . This also means that $\sigma_d[i]^{-1}(i)$ is of the form $\sigma_d[i]^{-1}(i) = f(|i|)$. Similarly, $\sigma'_d[j]^{-1}(j)$ is also stable by action of \mathfrak{S}_n on j , and is therefore also of the same form. We also remark that $\forall \mu \in \mathfrak{S}_n$, $\Sigma(\Psi, \mu(I)) = \Sigma(\Psi, I) \circ \mu^{-1}$ and therefore, thanks to the above:

$$\Sigma(\Psi, \sigma_d[\mu(i)](I_0)) = \Sigma(\Psi, \sigma_d[i](I_0)) \circ \mu^{-1}$$

The considerations above applied to $\omega_3^*(i, j, \sigma_d[i], \sigma'_d[j])$ imply that

$$\forall \sigma, \mu \in \mathfrak{S}_n, \omega_3^*(\sigma(i), \mu(j), \sigma_d[\sigma(i)], \sigma'_d[\mu(j)]) = \omega_3^*(i, j, \sigma_d[i], \sigma'_d[j])$$

Which also means that $\omega_3^*(i, j, \sigma_d[i], \sigma'_d[j])$ is of the form

$$\omega_3^*(i, j, \sigma_d[i], \sigma'_d[j]) = g(|i|, |j|)$$

Then, the exact writing of $E[(\omega_3^* - V_A)^2 | S]$ is:

$$\begin{aligned} & E[(\omega_3^* - V_A)^2 | S] \\ &= \sum_{i,j} \int_{x,y} \left(\omega_3^*(i, j, \sigma_d[i], \sigma'_d[j]) - \frac{\sigma_\Phi^{-1}(x) \cdot \sigma_{\Phi'}^{-1}(j)}{n} \right)^2 \chi_i \left(\frac{x}{k} \right) \chi_j \left(\frac{y}{k} \right) \Phi(x) \Phi'(y) dx dy \\ &= \sum_{i,j} \int_{x,y} \left(\omega_3^*(\sigma_\Phi(i), \sigma_{\Phi'}(j), \sigma_d[\sigma_\Phi(i)], \sigma'_d[\sigma_{\Phi'}(j)]) - \frac{x \cdot j}{n} \right)^2 \chi_i \left(\frac{x}{k} \right) \chi_j \left(\frac{y}{k} \right) \Phi \\ &\quad \circ \sigma_\Phi(x) \Phi' \circ \sigma_{\Phi'}(y) dx dy = \\ &= \sum_{i,j} \int_{x,y} \left(\omega_3^*(i, j, \sigma_d[i], \sigma'_d[j]) - \frac{x \cdot j}{n} \right)^2 \chi_i \left(\frac{x}{k} \right) \chi_j \left(\frac{y}{k} \right) \Phi \circ \sigma_\Phi(x) \Phi' \circ \sigma_{\Phi'}(y) dx dy \\ &= \sum_{i,j} \int_{x,y} \left(g(|i|, |j|) - \frac{x \cdot j}{n} \right)^2 \chi_i \left(\frac{x}{k} \right) \chi_j \left(\frac{y}{k} \right) \Phi \circ \sigma_\Phi(x) \Phi' \circ \sigma_{\Phi'}(y) dx dy \end{aligned}$$

and from there, $\Phi \circ \sigma_\Phi$ and $\Phi' \circ \sigma_{\Phi'}$ being synchronized (see Lemma 1' for a justification), we are in a specific case of the assumptions of Lemma 3, which we can apply to obtain the lower-bound:

$$E[(\omega_3^* - V_A)^2 | S] \geq \frac{C(\alpha)}{n}$$

By observing (for justification, see Proposition 11 for the first inequality and (II.0) for the second one) on the other hand that:

$$E[(\omega_0^* - V_A)^2 | S] \leq E[(V_B - V_A)^2 | S] \leq \frac{2}{nk} \quad (T1.1)$$

we finally get:

$$E[(\omega^* - V_A)^2 | S] \geq \frac{C(\alpha)}{16n} \left(1 - \sqrt{\frac{2}{kC(\alpha)}}\right)^2 \quad (T1.2)$$

The above inequality stands for all $\omega \in \Omega_3(\Sigma_1 \times \Sigma_2 \times \Sigma_3)$, and confirms that the considered situation is as desired for k sufficiently large, and thus ends step 1, with

$$C_1 = \frac{C(\alpha)}{16} \left(1 - \sqrt{\frac{2}{kC(\alpha)}}\right)^2$$

Step 2:

Each experiment of the protocol $\mathcal{P}_{1 \rightarrow 5}$ can lead to a binary encoding of V_A , V_B , and thus also V_ξ , as explained below:

We can typically sample a value in $[0,1]$ like V_A or V_B with a gauge being a multiple of the variance $E[(V_B - V_A)^2 | S]^{1/2} = \ll \left(\sqrt{\frac{2}{nk}}\right)$, the multiplicative factor K is chosen such that :

$$\frac{1}{\sqrt{nk}} \ll \frac{K}{\sqrt{nk}} \ll \frac{1}{\sqrt{n}}$$

and therefore, each experiment of the protocol can lead respectively A and B to distill an intermediate digit defined by:

$$\tilde{e}_A = \left\lfloor \frac{(V_A - \rho)\sqrt{nk}}{K} \right\rfloor \bmod 2, \tilde{e}_B = \left\lfloor \frac{(V_B - \rho)\sqrt{nk}}{K} \right\rfloor \bmod 2, \tilde{e}_\xi = \left\lfloor \frac{(V_\xi - \rho)\sqrt{nk}}{K} \right\rfloor \bmod 2$$

where ρ a translation parameter. If ρ is chosen (publicly by A) such that

$$\inf_{q \in \mathbb{Z}} \left| V_A - \rho - \frac{qK}{\sqrt{nk}} \right| \geq \frac{\lambda K}{\sqrt{nk}} \quad (C1)$$

with $\frac{1}{K} \ll \lambda \leq \frac{1}{2}$ then the inequality (T1.1) combined with Proposition 3, enables us to obtain:

$$P(\tilde{e}_A \neq \tilde{e}_B | S) \leq P\left((V_B - V_A)^2 \geq \frac{\lambda^2 K^2}{nk} | S\right) \leq 2ne^{-\frac{\lambda^2 K^2}{2}} \quad (T1.3)$$

The fact to choose ρ publicly, does not give useful information to ξ when S is satisfied, because of (T1.4) shown below.

On the other hand, by denoting $C^* = C_1 \left(1 - \sqrt{\frac{1}{kC_1}}\right)^2$, one can apply Corollary 3 to get:

$$P\left((\omega^* - V_A)^2 \geq \frac{C^*}{2n} | S\right) \geq \frac{C_2}{\ln n}$$

Because $\frac{C^*}{2n} \gg \frac{K^2}{nk}$, the above inequality implies (to be fully convinced, one can chose the origin of the sampling and the constant K as independent random variables) that:

$$P(\widetilde{e}_A \neq \widetilde{e}_\xi | S) \geq \frac{C_2}{2 \ln n} - O\left(ne^{-\frac{K^2}{2}}\right) \quad (T1.4)$$

The reception of \widetilde{e}_A as \widetilde{e}_B by B , and as \widetilde{e}_ξ by ξ corresponds to a broadcast BSC channel $BSC(\varepsilon_B, \varepsilon_\xi)$ with notation of Proposition 12 (we can easily see that the channels are BSC by remarking that changing ρ in $\rho + \frac{K}{\sqrt{nk}}$ inverses the digits and is independent of V_A, V_B , and V_ξ). From Proposition 12, $BSC(\varepsilon_B, \varepsilon_\xi)$ can be modeled in a $CBSC(\epsilon_A, \epsilon_B, \epsilon_\xi)$ where the 3 independent BSC have binary error variables $(\delta_A, \delta_B, \delta_\xi)$ with respective binary error probability $(\epsilon_A, \epsilon_B, \epsilon_\xi)$. It has been shown by Maurer in [5] that in a $CBSC$ model, if $\epsilon_\xi > 0$ then Advantage Distillation can be achieved thanks to error correcting methods, and consecutively perfect secrecy can be approached as close as desired thanks to Information Reconciliation and Privacy Amplification techniques. The proof is also given in Proposition 13 (ii) in the specific case where $\epsilon_A = \epsilon_B$; and we remark that our protocol fulfill that condition $\epsilon_A = \epsilon_B$ due to the fact that steps 1 to 5 are symmetric from a design perspective, and symmetry Σ_3 introduced earlier ensures that strategy $\omega^* = V_\xi$ (and thus also error variable ε_ξ) is invariant by permuting A and B roles. Let's therefore establish a lower-bound for ϵ_ξ .

We can consider the conditional expectations of $\widetilde{e}_A, \widetilde{e}_B$ and \widetilde{e}_ξ knowing $\mathfrak{S} \triangleq (i, j, \mu_1, \mu_2, \mu'_1, \mu'_2)$. If we write:

$$\begin{aligned} \widetilde{e}_A &= E[\widetilde{e}_A | \mathfrak{S}] \oplus \delta_A^* \\ \widetilde{e}_B &= E[\widetilde{e}_B | \mathfrak{S}] \oplus \delta_B^* \\ \widetilde{e}_\xi &= E[\widetilde{e}_\xi | \mathfrak{S}] \oplus \delta_\xi^* \end{aligned}$$

δ_A^* is independent of $E[\widetilde{e}_A | \mathfrak{S}]$, δ_B^* and δ_ξ^* ; δ_B^* is independent of $E[\widetilde{e}_B | \mathfrak{S}]$, δ_A^* and δ_ξ^* ; δ_ξ^* is independent of $E[\widetilde{e}_\xi | \mathfrak{S}]$, δ_B^* and δ_A^* . $(\delta_A^*, \delta_B^*, \delta_\xi^*)$ have respective error probability $(\epsilon_A^*, \epsilon_B^*, \epsilon_\xi^*)$. Furthermore, by approximating $E[\widetilde{e}_A | \mathfrak{S}]$, $E[\widetilde{e}_B | \mathfrak{S}]$, and $E[\widetilde{e}_\xi | \mathfrak{S}]$ as shown in Proposition 14, we obtain (with notations of Proposition 14) that

$$E[\widetilde{e}_A | \mathfrak{S}] + O\left(\sqrt{\frac{\ln n}{nk}}\right) = E[\widetilde{e}_B | \mathfrak{S}] + O\left(\sqrt{\frac{\ln n}{nk}}\right) = E[\widetilde{e}_\xi | \mathfrak{S}] + O\left(\sqrt{\frac{\ln n}{nk}}\right)$$

We then deduce that the model $(\delta_A^*, \delta_B^*, \delta_\xi^*)$ is a $O\left(\sqrt{\frac{\ln n}{nk}}\right)$ approximation of the $CBSC(\epsilon_A, \epsilon_B, \epsilon_\xi)$ model. Indeed:

$$X = P(\varepsilon_B = 1) = P(\widetilde{e}_A \oplus \widetilde{e}_B = 1) = \varphi(\epsilon_A, \epsilon_B)$$

$$= P(\widetilde{e}_A \oplus \widetilde{e}_B \oplus (E[\widetilde{e}_A | \mathfrak{S}] \oplus E[\widetilde{e}_B | \mathfrak{S}]) = 1) = \varphi(\epsilon_A^*, \epsilon_B^*) + O\left(\sqrt{\frac{\ln n}{nk}}\right)$$

and similarly for Y and Z . We thus have the approximation:

$$\phi(\epsilon_A, \epsilon_B, \epsilon_\xi) = \phi(\epsilon_A^*, \epsilon_B^*, \epsilon_\xi^*) + O\left(\sqrt{\frac{\ln n}{nk}}\right) \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$$

From Proposition 16 (ii), if we pick ρ, λ and $\frac{\lambda K}{\sqrt{nk}} = \theta_0$ as suggested in assumption of Proposition 16, we can apply the result to $U = V_A$ and $V = V_B$ in situation $S_1 \cup S_2 \cup S_3 = \{(\sigma_A, \sigma_B) \neq (\sigma_{\Phi'}, \sigma_\Phi)\}$, and we deduce that

$$P((\epsilon_B = 1) | (\sigma_A, \sigma_B) \neq (\sigma_{\Phi'}, \sigma_\Phi)) \leq \frac{1}{2} + \frac{1}{2(\lambda - 1)}$$

On the other hand, with inequality (T1.3) applied to $P((\epsilon_B = 1) | (\sigma_A, \sigma_B) = (\sigma_{\Phi'}, \sigma_\Phi))$, we finally get the following upper-bounds for $P(\epsilon_B = 1)$:

$$P(\epsilon_B = 1) \leq \frac{3}{8} + \frac{3}{8(\lambda - 1)} + \frac{n}{4} e^{-\frac{\lambda^2 K^2}{2}}$$

We can assume that we have also both upper-bounds:

$$P(\epsilon_\xi = 1) \leq \frac{3}{8} + \frac{3}{8(\lambda - 1)} + \frac{n}{4} e^{-\frac{\lambda^2 K^2}{2}}$$

$$P(\epsilon_\xi \neq \epsilon_B) \leq \frac{3}{8} + \frac{3}{8(\lambda - 1)} + \frac{n}{4} e^{-\frac{\lambda^2 K^2}{2}}$$

otherwise we would already have an advantage of the estimation of A or the estimation of B over the estimation of ξ . Therefore, these upper-bounds enable to apply Lipschitzian approximation authorized by Proposition 12 (iv) for λ sufficiently large:

$$\phi(\epsilon_A, \epsilon_B, \epsilon_\xi) = \phi(\epsilon_A^*, \epsilon_B^*, \epsilon_\xi^*) + O\left(\sqrt{\frac{\ln n}{nk}}\right) \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \Rightarrow \begin{cases} \epsilon_A^* = \epsilon_A + O\left(\sqrt{\frac{\ln n}{nk}}\right) \\ \epsilon_B^* = \epsilon_B + O\left(\sqrt{\frac{\ln n}{nk}}\right) \\ \epsilon_\xi^* = \epsilon_\xi + O\left(\sqrt{\frac{\ln n}{nk}}\right) \end{cases}$$

δ_ξ^* is independent of the main channel, and in particular then of S , δ_A^* , and δ_B^* , and also independent of \mathfrak{S} . This allows us to write

$$P(\widetilde{e}_A \neq \widetilde{e}_\xi | S) = P(\delta_A^* \oplus (E[\widetilde{e}_A | \mathfrak{S}] \oplus E[\widetilde{e}_\xi | \mathfrak{S}]) = 1 | S) (1 - \epsilon_\xi^*) \\ + \epsilon_\xi^* \left(1 - P(\delta_A^* \oplus (E[\widetilde{e}_A | \mathfrak{S}] \oplus E[\widetilde{e}_\xi | \mathfrak{S}]) = 1 | S)\right)$$

$$\Rightarrow P(\widetilde{e}_A \neq \widetilde{e}_\xi | S) = P(\delta_A^* = 1 | S)(1 - \epsilon_\xi^*) + \epsilon_\xi^*(1 - P(\delta_A = 1 | S)) + O\left(\sqrt{\frac{\ln n}{nk}}\right) \quad (T1.5)$$

Furthermore, the same independence argument allow us to also write, $\forall(\tau, \tau') \in \{0,1\}^2$:

$$\begin{aligned} P((\delta_A^* = \tau) \& (\delta_B^* = \tau') \& S) &= P((\delta_A^* = \tau) \& (\sigma_A = \sigma_{\Phi'}))P((\delta_B^* = \tau') \& (\sigma_B = \sigma_\Phi)) \\ \Rightarrow P((\delta_A^* = \tau) \& (\delta_B^* = \tau') | S)P(S) &= P((\delta_A^* = \tau) \& (\sigma_A = \sigma_{\Phi'}) | (\sigma_B = \sigma_\Phi))P((\delta_B^* = \tau') \& (\sigma_B = \sigma_\Phi) | (\sigma_A = \sigma_{\Phi'})) \\ \Rightarrow P((\delta_A^* = \tau) \& (\delta_B^* = \tau') | S)P(S)^2 &= P((\delta_A^* = \tau) \& (\sigma_A = \sigma_{\Phi'}) \& (\sigma_B = \sigma_\Phi))P((\delta_B^* = \tau') \& (\sigma_B = \sigma_\Phi) \& (\sigma_A = \sigma_{\Phi'})) \\ &\Rightarrow P((\delta_A^* = \tau) \& (\delta_B^* = \tau') | S) = P((\delta_A^* = \tau) | S)P((\delta_B^* = \tau') | S) \end{aligned}$$

The above equality implies that:

$$\begin{aligned} P((\delta_A^* = 1) | S) &\leq \frac{P((\epsilon_B \oplus E[\widetilde{e}_A | \mathfrak{S}] \oplus E[\widetilde{e}_B | \mathfrak{S}] = 1) | S)}{(1 - P((\delta_B^* = 1) | S))} \leq 2P((\epsilon_B \oplus E[\widetilde{e}_A | \mathfrak{S}] \oplus E[\widetilde{e}_B | \mathfrak{S}] = 1) | S) \\ &\Rightarrow P((\delta_A^* = 1) | S) \leq 2P((\epsilon_B = 1) | S) + O\left(\sqrt{\frac{\ln n}{nk}}\right) \end{aligned}$$

which we can inject in (T1.5) and combine with (T1.4) and (T1.3) to obtain:

$$\epsilon_\xi^* \geq \frac{C_2}{2 \ln n} - O\left(ne^{-\frac{\lambda^2 K^2}{2}}\right) - O\left(\sqrt{\frac{\ln n}{nk}}\right)$$

and as we have also obtained above that $\epsilon_\xi^* = \epsilon_\xi + O\left(\sqrt{\frac{\ln n}{nk}}\right)$, we can establish a final lower-bound.

That gives the lower-bound fitting Maurer's condition for K, n, k sufficiently large.

However, it remains to justify that Reconciliation technique does not bring additional information to the opponent due to the choice of code words. Indeed, contrariwise the protocol presented in [5], in our case the opponent can try the various possible combinations of choice for σ_A and σ_B , and compare them in terms of matching a code word. To illustrate the issue, let's consider the non-optimal 'bit repeating' error correction method described by Maurer in [5]: the codeword v_A chosen by A can only be $(0,0, \dots, 0)_L$ or $(1,1, \dots, 1)_L$ depending on $e_A = 0$ or $e_A = 1$. B publicly discards all decoded sequence v_B that is not $(0,0, \dots, 0)_L$ or $(1,1, \dots, 1)_L$ and obviously decodes accordingly $e_B = 0$ if $|v_B| = 0$, and $e_B = 1$ if $|v_B| = L$. In such scenario, the opponent will use the strategy below, as justified by Proposition 14 under Deep Random assumption:

$$V_\xi(i, j, \sigma_{\xi,A}, \sigma_{\xi,B}) = \frac{2k \left(\left(\sum_{r \in \sigma_{\xi,B}(I_0)} i_r \right) \left(\sum_{r \in \sigma_{\xi,A}(I_0)} j_r \right) + \left(\sum_{r \in \overline{\sigma_{\xi,B}(I_0)}} i_r \right) \left(\sum_{r \in \overline{\sigma_{\xi,A}(I_0)}} j_r \right) \right)}{n^2}$$

where $\sigma_{\xi,B}$ is a permutation chosen by ξ among (μ_1, μ_2) and $\sigma_{\xi,A}$ is a permutation chosen by ξ among (μ'_1, μ'_2) . It is easy to show that the strategy above is the canonical strategy efficient when $(\sigma_{\xi,A}, \sigma_{\xi,B}) = (\sigma_{\Phi'}, \sigma_{\Phi})$. Then, at each instantiation m of the protocol, we can consider the vector of $\{0,1\}^4$ defined by:

$$T_m = \left\{ \tilde{e}_{\xi}(\mu'_1, \mu_1)_m, \tilde{e}_{\xi}(\mu'_1, \mu_2)_m, \tilde{e}_{\xi}(\mu'_2, \mu_1)_m, \tilde{e}_{\xi}(\mu'_2, \mu_2)_m \right\}$$

When A encodes a code word with a sequence of L instantiations of the protocol, if, for one of the instance m within the sequence, we have $\sigma_A = \sigma_{\Phi'}$ and $T_m = (0,0,0,0)$ (or $T_m = (1,1,1,1)$), then it is clear that the code word is $(0,0, \dots, 0)_L$ (or $(1,1, \dots, 1)_L$). This example shows how, even with Deep Randomness indistinguishability applied locally to each instance, the opponent can still take advantage of an imprudent implementation of the error correcting method to break the secrecy. In order to avoid this flaw, and to maintain the invariance by the group of symmetry in the context of adding the error correction at the end of the protocol, it is needed to ensure that:

$$|T_m| = 2, \quad \forall m$$

Indeed, in the above condition, the opponent has no information in any circumstance to privilege $(0,0, \dots, 0)_L$ or $(1,1, \dots, 1)_L$ as the chosen code word. We can remark that A can easily ensure the above condition in almost all cases: indeed, as soon as the four possible values of $V_{\xi}(i, j, \sigma_{\xi,A}, \sigma_{\xi,B})$ (when $\sigma_{\xi,B}$ courses (μ_1, μ_2) and $\sigma_{\xi,A}$ courses (μ'_1, μ'_2)) are all different, it is possible to find a value of the translation parameter ρ such that the above condition is satisfied. In order to justify this, we can consider the sampling function:

$$f_{\epsilon}(v) = \begin{cases} 0 & \forall v \in [2q + \epsilon, 2q + 1 - \epsilon], \quad q \in \mathbb{Z} \\ 1 & \forall v \in [2q + 1 + \epsilon, 2q + 2 - \epsilon], \quad q \in \mathbb{Z} \\ -\frac{1}{2\epsilon}(v - 2q) + \frac{1}{2} & \forall v \in [2q - \epsilon, 2q + \epsilon], \quad q \in \mathbb{Z} \\ \frac{1}{2\epsilon}(v - 2q - 1) + \frac{1}{2} & \forall v \in [2q + 1 - \epsilon, 2q + 1 + \epsilon], \quad q \in \mathbb{Z} \end{cases}$$

Indeed, it is clear that (i) $f_{\epsilon} \left(\frac{(V_{\xi} - \rho)\sqrt{nk}}{K} \right) \xrightarrow{\epsilon \rightarrow 0} \tilde{e}_{\xi}$, (ii) f_{ϵ} is continuous, and (iii) $f_{\epsilon}(v + 1) = 1 - f_{\epsilon}(v)$.

Therefore, $T_{m,\epsilon}(\rho) = \overline{T_{m,\epsilon} \left(\rho + \frac{K}{\sqrt{nk}} \right)}$, and by continuity, as soon as ϵ is smaller than the smallest gap between two of the four possible values of $V_{\xi}(i, j, \sigma_{\xi,A}, \sigma_{\xi,B})$, it is possible to find a value of ρ_{ϵ} that satisfies

$$|T_{m,\epsilon}(\rho)| = 2 \tag{C2}$$

in any interval with size $\frac{K}{\sqrt{nk}}$. And one can conclude by making $\epsilon \rightarrow 0$ with an accumulation point of ρ_{ϵ} . Due to the (T1.4) bound compared to the (T1.3) bound, there is a fair probability (controlled especially by the parameter λ) that the conditions (C1) and (C2) coexist, even without the dependency of ρ_{ϵ} in $(i, j, \sigma_{\xi,A}, \sigma_{\xi,B})$ introduced by the above method. If ever the 2 conditions don't coexist for a given draw, the instance can just be discarded.

If ever the four possible values of $V_{\xi}(i, j, \sigma_{\xi,A}, \sigma_{\xi,B})$ are not all different, the instance can just be discarded.

This is enough to conclude the proof of the Theorem. ■

Theorem 2.

$\mathcal{P}(\alpha, n, k, K, L)$ satisfies condition (P').

■ If the strategy chosen by the opponent verifies the inequality (T1.2) of Theorem 1, then the result follows. If not, it is sufficient to replace randomly in 50% of cases $(\Phi, \Psi) \rightarrow (\Psi, \Phi)$ and then the proof of Theorem 1 applies equally to the new situation. ■