

A New Property of Hamilton Graphs

Heping JIANG

Rm. 702, Building 12, Luomashi Avenue, Xi Cheng district,
Beijing, China, zip code: 100052

E-mail address: jjhpjhp@gmail.com

Abstract. A Hamilton cycle is a cycle containing every vertex of a graph. A graph is called Hamiltonian if it contains a Hamilton cycle. The Hamilton cycle problem is to find the sufficient and necessary condition that a graph is Hamiltonian. In this paper, we give out some new kind of definitions of the subgraphs and determine the Hamiltonicity of edges according to the existence of the subgraphs in a graph, and then obtain a new property of Hamilton graphs as being a necessary and sufficient condition characterized in the connectivity of the subgraph that induced from the cycle structure of a given graph.

MSC2010 05C45

Keywords Hamilton graphs Hamilton edges cycle-sets subgraphs removable cycles

1. Introduction

Graphs considered throughout this paper are finite, undirected and simple connected graphs. A Hamilton cycle is a cycle containing every vertex of a graph. A graph is called Hamiltonian if it contains a Hamilton cycle. The Hamilton cycle problem is to find the necessary and sufficient condition that a graph is Hamiltonian. Since Eulerian graphs have a simple characterization, the study of the relationship between Hamilton graphs and degree of vertices became a classical method and received numerous results [1]. In the meantime, different from the classical method, a kind of quantitative analysis, S. Goodman and S. Hedetniemi investigated the relationship between subgraphs and the Hamilton graphs and developed a qualitative analysis [2]. From then on the subgraphs have been widely studied [1] [3]. However, the Hamilton cycle problem is still an open problem. In this paper, using the qualitative analysis, we give out some new kind of definitions of the subgraphs and determine the Hamiltonicity of edges according to the existence of the subgraphs in a graph, and then obtain a new property of Hamilton graphs as being a necessary and sufficient condition characterized in the connectivity of the subgraph that induced from the cycle structure of a given graph. Based on this property, we will show a polynomial algorithm for finding a Hamilton cycle in a graph in the other paper.

Let G be a graph with vertex set V and edge set E . We use combination to represent the overlapping of two graphs (or cycles) based on common vertices and edges. A cycle-set is the combination of $V-E+1$ cycles of G such that it covers all vertices and edges of G . The union of all elements in a cycle-set is the graph itself. F_s is a spanning cycle-set, denotes a set of minimum number of cycles covering all vertices in G . In a cycle-set of G , every node corresponding to the vertex in G implies the overlap of vertices of the cycles passing through. So does an edge in G . For brevity, we reserve the

word “vertex” and “edge” to denote the overlapping node and the overlapping edge respectively. Let R be the overlapping number of cycles passing through an edge of G . We use R_i to denote an edge of $R=i$ (i is a natural number). In a cycle-set of G , a boundary vertex is a vertex being incident with two R_1 only, an interior vertex is a vertex being incident with R_2 only, and a cycle is removable implies that it has a unique R_1 whose two end vertices are boundary vertices. We use C_x to denote a removable cycle such that the rest of the cycle-set is non-Hamiltonian when removing it from the cycle-set. A vertex that there are not less than 3 vertices of degree less than or equal to 2 in its neighborhood is denoted by N_3 , N_2 denotes a vertex that there are 2 vertices of degree less than or equal to 2 in its neighborhood, and N_1 denotes a vertex that there are vertices of degree equal to 1. If there are no N_3 , N_2 and N_1 in G , we say G is (N_3, N_2, N_1) -free. R_2 -free denotes there are no R_2 in a cycle-set of G . Sometimes we combine N_3 , N_2 , N_1 and R_2 in one parentheses such as (N_3, N_2, N_1, R_2) -free. We use F_s+C_x to denote recombination of F_s and C_x such that there are no other removable cycles except for C_x . Let \mathbf{F} be a (N_3, N_2, R_2) -free subgraph of F_s+C_x . For terms and notations not defined here see [3] [4].

In this paper we investigate the relationship between C_x and Hamilton cycles in a cycle-set of a graph. Generally speaking, for a given (N_3, N_2, N_1) -free subgraph of a Hamilton graph G , we could obtain a set of cycles by removing some cycles from it, and the union of these cycles yields a Hamilton cycle. But, for a graph that the Hamiltonicity is unknown, if there is a C_x in the removed cycles, then the claim of Hamiltonicity of the given graph is sometimes not true (see **Lemma 2.3.**). Thus, we study the Hamiltonicity of edges in the structure of the cycle-set and develop a new method to determine the Hamiltonicity of the given graph. After excluding the graphs with N_3 and eliminating the paths which a Hamilton cycle must not pass through at N_2 and N_1 (see **Lemma 2.1.**, **Corollary 2.1.**), we obtain a (N_3, N_2, N_1) -free subgraph of G . For a cycle-set of this subgraph, by deleting all the removable cycles arbitrarily in a sequence one by one under the condition of N_3 -free, we can detect C_x out (see **Corollary 2.1.**, **Lemma 2.2.**) and recombine C_x and the subgraph. With the same method, we obtain the F_s+C_x eventually. We show that a connected \mathbf{F} is the necessary and sufficient condition to determine the given graph is Hamiltonian (see **Theorem 2.1.**, **Theorem 2.2.**, **Theorem 2.3.**, and **Corollary 2.2.**).

2. The proofs

Lemma 2.1. *Let G be a graph. For $N_2 \in V(G)$, the edge not being incident with the vertices of degree 1 or 2 in the neighborhood of N_2 is not a path which a Hamilton cycle must pass through.*

Proof. Let G be a graph. For $N_2 \in V(G)$, v denotes a vertex of degree 1 or 2 in the neighborhood of N_2 . Since a Hamiltonian cycle (if has) will enter and leave each vertex in G exactly once, deleting just one edge of v only causes G non-Hamiltonian. In the other words, it means that all the edges not being incident with the vertices of degree 1 or 2 in the neighborhood of N_2 are not paths which a Hamilton cycle must pass through. \square

Corollary 2.1. *Let G be a graph. For $N_3 \in V(G)$, G is non-Hamiltonian if it includes a N_3 .*

Proof. Let v_1, v_2, \dots, v_i ($i \geq 3$) be the vertices of degree 1 or 2 in neighborhood of N_3 ($N_3 \in V(G)$). From **Lemma 2.1.**, since v_1 and v_2 constitute N_2 at vertex N_3 , then the edge N_2v_i ($i \geq 3$) is not a path which a Hamilton cycle (if has) must pass through, and that implies no Hamilton cycles (if has) pass through v_i ($i \geq 3$). Thus, G is non-Hamiltonian. \square

Theorem 2.1. *F_s is Hamiltonian if, and only if, its (N_3, N_2, R_2) -free subgraph is connected.*

Proof. For a spanning cycle-set F_s , induced from a given (N_3, N_2, N_1) -free graph G , we can divide the edges into three kinds: R_1 , R_2 , and R_{3+i} (i is a natural number). By the definition, we know that no pair of cycles is same in a covering of a cycle-set over on the graph, therefore, F_s never consists of only R_2 or R_{3+i} and there have four cases as following: $\{R_1\}$, $\{R_1, R_2\}$, $\{R_1, R_{3+i}\}$, $\{R_1, R_2, R_{3+i}\}$. By the definition, an F_s deleted any one of irremovable cycles is non-Hamiltonian, that implies a R_1 on the irremovable cycles is a path which a Hamilton cycle must pass through. On the other, we know that a R_2 is a public edge of tow cycles, if there are two R_1 on an endvertex of a R_2 , then the R_2 is not a path which a Hamilton cycle must pass through by **Lemma 2.1.**. Thus, the following analysis will base on the fact that a R_1 in an F_s is irremovable and a R_2 is removable.

(i) $\{R_1\}$ If the edges in an F_s consist of R_1 only, it is obviously that the edges in the (N_3, N_2, R_2) -free subgraph consist of R_1 only, too. If the subgraph is disconnected, then there must be a R_2 in F_s . It is clear that the case does not exist. And this means the subgraph is connected. Note that a R_1 in an F_s is a path which a Hamilton cycle must pass through. Hence, the only result under the (N_3, N_2, R_2) -free condition is that the subgraph is a Hamilton cycle.

(ii) $\{R_1, R_2\}$ If the subgraph is disconnected then there must be a vertex of order 2 (as an interior vertex) on a R_2 in the F_s , and then the two R_2 are paths which a Hamilton cycle must pass through. But an F_s is a graph without removable cycles, so there must be two R_1 being adjacent to the endvertex of an R_2 . Therefore, the endvertex is a N_3 . By **Corollary 2.1.**, the F_s is non-Hamiltonian. Since an F_s is a subgraph induced from a (N_3, N_2, N_1) -free graph, then, If the subgraph is connected then the Union of all elements in the (N_3, N_2, R_2) -free subgraph is actually a spanning cycle, that is the F_s is Hamiltonian.

(iii) $\{R_1, R_{3+i}\}$ Without loss of generality, we consider the case that there are $3+i$ irremovable cycles on a R_{3+i} . Under (N_3, N_2, R_2) -free condition, by **Lemma 2.1.**, the two edges that are adjacent to each of the endvertices of the R_{3+i} are only paths which a Hamilton cycle must pass through. So, there is only one result for the (N_3, N_2, R_2) -free subgraph, the subgraph is disconnected that means there are not enough paths in the F_s to satisfy the existence of a Hamilton cycle.

(iv) $\{R_1, R_2, R_{3+i}\}$ We consider mainly the relationship between the location of a R_2 and the connectivity of the subgraph besides that the case of R_1 and R_{3+i} is the same as (iii).

(1) In the case of that every R_2 is a chord of a spanning cycle, deleting a R_2 under (N_3, N_2, R_2) -free condition has no relationship with the connectivity of the subgraph. Hence, the case is the same as (iii).

(2) There are not less than two disjoint R_2 on a spanning cycle. If R_2 are deleted under (N_3, N_2, R_2) -free condition, then the subgraph of the F_s is disconnected. Note that a R_2 in an F_s is removable. Thus, it is obviously that there are not enough paths which a Hamilton cycle must pass through, that is the F_s is non-Hamiltonian.

(3) If anyone of the R_2 is a chord of a cycle in F_s , then, the endvertices of the R_2 will be belong to two cycles respectively after deleting the R_2 , it can only mean one thing that there is a cut vertex in the subgraph of the F_s . Clearly, the case is the same as (iii) also.

As proved above, a connected (N_3, N_2, R_2) -free subgraph is the necessary and sufficient condition of a Hamiltonian F_s . \square

Lemma 2.2. *Let F be a cycles-set of (N_3, N_2) -free graph G . Every R_1 of C_x in G is a path which a Hamilton cycle must pass through.*

Proof. By the definition of N_2 and N_3 , we can easily construct a series of non-Hamiltonian graphs by deleting one irremovable cycle such that all non-Hamiltonian graphs are isomorphic to one of these non-Hamiltonian graphs, showing in the Figure 1.




Figure 1

Therefore, we can construct a recombination of F_s and C_x (the irremovable cycle) showing in the Figure 2 below. It is clear that F_s+C_x is Hamiltonian implies there exists




Figure 2

a path which a Hamilton cycle must pass through in C_x . Since a R_1 is the only edge added into the graph (the doting line in the figure), then a R_1 of C_x in G must be a path which a Hamilton cycle must pass through. \square

Theorem 2.2. F_s+C_x is Hamiltonian if, and only if, F is connected.

Proof. Let G be a (N_3, N_2, N_1) -free graph, $F_s \subseteq G$, $C_x \subseteq G$, F is a (N_3, N_2, R_2) -free subgraph. We first prove a statement that no endvertex of R_1 in F is an isolated vertex. Since G is a (N_3, N_2, N_1) -free graph, then it is impossible for an endvertex existing as an isolated vertex in F_s which induced from a cycle-set of G . Therefore, all R_1 is a path which a Hamilton cycle must pass through in F under the (N_3, N_2, R_2) -free condition induced from F_s+C_x . Hence, the statement is hold.

Since there have only R_1 and R_{3+i} in F , and by **Theorem 2.1.** and **Lemma 2.2.**, all the R_1 in an F_s or C_x is a path which a Hamilton cycle must pass through, then it is enough to consider the case of R_{3+i} in F . Know that there is no isolated vertex to be as an endvertex in F , thus only four cases of the R_{3+i} existing with R_1 in F need considering as following (see the Figure 3),




Figure 3

- Case 1 R_{3+i} is an edge of a cycle;
- Case 2 R_{3+i} is an chord of a cycle;
- Case 3 R_{3+i} is a bridge connected two cycles;
- Case 4 R_{3+i} is a bridge connected three or more cycles.

For Case 3 and Case 4, it is easy to determine, by **Lemma 2.1.**, that R_{3+i} is a path which a Hamilton cycle must pass through so that it must be deleted when inducing F from F_s+C_x , thus, F is disconnected, and then F_s+C_x is non-Hamiltonian clearly. For Case 2, similarly by **Lemma 2.1.**, R_{3+i} must be deleted, so that F is connected, and then F_s+C_x is Hamiltonian. For case 1, since no isolated vertex exists, then there is a cycle pass through every vertex once exactly, and then F is connected, F_s+C_x is Hamiltonian. In summary, we derive that F_s+C_x is Hamiltonian if, and only if, F is connected. Moreover, when adding a vertex (or more) of degree 2 into R_{3+i} , there is no change of the connectivity of F despite of stronger passing necessity to R_{3+i} so that the conclusion above is hold. \square

Lemma 2.3. *Let F be a cycle-set of (N_3, N_2) -free graph G . C_r is all the irremovable cycles except C_x ($C_x \in F$). R_1 in C_r is a path by which a Hamilton cycle must pass in F .*

Proof. After deleting all irremovable cycles, if a spanning cycle-set induced from F is Hamiltonian, then we have two cases to consider in these irremovable cycles: $C_x=0$, $C_x \neq 0$. It is clear that R_1 in C_r is a path which a Hamilton cycle must pass through in F no matter what case to consider. If a spanning cycle-set induced from F is non-Hamiltonian, we still have two cases: $C_x=0$, $C_x \neq 0$. By **Theorem 2.2.**, we know the only reason for F_s+C_x is non-Hamiltonian is that a (N_3, N_2, R_2) -free edge induced subgraph from F_s+C_x is disconnected, and there is no relationship to R_1 in C_r . Hence, the lemma is hold. \square

Theorem 2.3. *Every two spanning cycle-sets in the given graph have the same Hamiltonian property with the given graph.*

Proof. Graph G is a (N_3, N_2) -free graph. Let F_1 and F_2 denote two different labeled cycle-sets in graph G . F_{s1} and F_{s2} denote the spanning cycle-set of F_1 and F_2 respectively. By the definition of a cycle-set, we know that the union of the cycles in a cycle-set of graph G is the graph G itself, so do F_1 and F_2 . Therefore, there exist the same number of paths which a Hamilton cycle must pass through both in F_1 and F_2 . Since every edge in G is correspondent to a value of R in an F , then we can determine the Hamiltonicity of an edge. By **Lemma 2.3.**, we know R_1 of C_r is a path which a Hamilton cycle must not pass through, thus we can derive F_{s1} and F_{s2} by deleting all the irremovable cycles from G , and therefore, we can determine R_2 in F_s is a path which a Hamilton cycle must not pass through also. While by **Lemma 2.2.**, we know R_1 in F_s is a path which a Hamilton cycle must pass through. In addition, there is no change of the number of R_{3+i} when inducing F_s from F . So, all the paths which a Hamilton cycle must pass through in F_1 and F_2 remained in F_{s1} and F_{s2} . Hence, F_{s1} and F_{s2} not only have same number of the paths in graph G but also have the same Hamiltonicity with graph G . That completes the proof. \square

Corollary 2.2. *Graph G is Hamiltonian if, and only if, F_s+C_x is Hamiltonian.*

Proof. By **Theorem 2.3.**, we know that there are no changes of the Hamiltonicity of a cycle-set (F) derived from anyone of the cycle-set of the given graph G , and so does every F_s+C_x induced from graph G . \square

References

- [1] Ronald J. Gould, Advances on the Hamiltonian problem—a survey, *Graphs Combinatorics*, 19 (1) (2003) 7-52.
- [2] S. Goodman, S. Hedetniemi, Sufficient Condition for a graph to be Hamiltonian, *Journal of Combinatorial Theory Ser. B* 16 (1974) 175-180.
- [3] Ronald J. Gould, *Graph Theory*, Dover Publications, Inc., New York, 2012.
- [4] J.A. Bondy, U.S.R. Murty, *Graph Theory with Applications*, Fifth Printing, Elsevier Science Publishing Co., Inc., New York, 1982.