No simple arbitrage for fractional Brownian motion

Rémi Peyre*

August 3, 2015

Abstract

We prove the following result: For $(Z_t)_{t \in \mathbf{R}}$ a fractional Brownian motion with arbitrary Hurst parameter, there does not exist any stopping time τ adapted to the natural filtration of the increments of Z such that, with positive probability, τ a local minimum at right of the trajectory of Z.

1 Introduction

1.1 Context

In this article, we consider a filtered probability space $(\Omega, \mathbb{P}, \mathcal{B}, (\mathcal{B}_t)_{t \in \mathbf{R}})$. Notation ' ω ' will implicitly refer to eventualities of Ω ; we will use it from time to time when needing to make the dependency on the random phenomenon perfectly clear. We consider a (bilateral) Brownian motion $(W_t)_{t \in \mathbf{R}}$ whose increments are adapted to our filtered space, which means, for all $t \in \mathbf{R}$, for all $u \leq 0$, $(W_{t+u} - W_t)$ is \mathcal{B}_t -measurable, while for all $v \geq 0$, $(W_{t+v} - W_t)$ is independent from \mathcal{F}_t .

We fix once for all some arbitrary parameter $H \in (0, 1)$ (so that, in the sequel, "absolute" constants may actually depend on H) such that $H \neq 1/2$; moreover, in all this article, (H-1/2) may be referred to as η . Then we consider the fractional Brownian motion (fBm) $(Z_t)_{t \in \mathbf{R}}$ driven by W with Hurst parameter H, which means that

$$Z_t := C_1 \int_{\mathbf{R}} \left((t-s)_+^{\eta} - (-s)_+^{\eta} \right) dW_s, \tag{1}$$

(with the convention that $0^r = 0 \ \forall r \in \mathbf{R}$), where

$$C_1 := \left(\frac{1}{2H} + \int_0^\infty \left((1+s)^\eta - s^\eta\right)^2 ds\right)^{-1/2}.$$
 (2)

Then, the properties of Z are well known: it is a centred Gaussian process whose increments are adapted to $(\mathcal{B}_t)_t$, with $\operatorname{Var}(Z_t - Z_s) = |t - s|^{2H}$ and $Z_0 = 0$ a.s.; its trajectories are locally $(H - \varepsilon)$ -Hölder with divergence in $O(|t|^{H+\varepsilon})$ at infinity, etc. (see [2, Chap. 2] by example).

Remark 1. Here the integral in the r-h.s. of (1) should be seen as a deterministic integral rather than as an Itô integral. Indeed, integrating by parts, one has:

^{*}CNRS & Institut Élie-Cartan de Lorraine (Univ. Lorraine); remi.peyre@univ-lorraine.fr.

(here in the case t > 0),

$$C_{1}^{-1}Z_{t} = \int_{\mathbf{R}}^{0} \left((t-s)_{+}^{\eta} - (-s)_{+}^{\eta} \right) dW_{s}$$

$$= \int_{-\infty}^{0} \left((t-s)^{\eta} - (-s)^{\eta} \right) dW_{s} + \int_{0}^{t} (t-s)^{\eta} dW_{s}$$

$$= \left[\left((t-s)^{\eta} - (-s)^{\eta} \right) W_{s} \right]_{s=-\infty}^{0} - \eta \int_{-\infty}^{0} \left((t-s)^{\eta-1} - (-s)^{\eta-1} \right) W_{s} ds$$

$$+ \left[(t-s)^{\eta} (W_{s} - W_{t}) \right]_{s=0}^{t} - \eta \int_{0}^{t} (t-s)^{\eta-1} (W_{s} - W_{t}) ds$$

$$= t^{\eta} W_{t} - \eta \int_{-\infty}^{t} \left((t-s)^{\eta-1} - (-s)^{\eta-1}_{+} \right) (W_{s} - \mathbf{1}_{s>0} W_{t}) ds, \quad (3)$$

where all the computations are licit (with absolutely converging integrals) because of the properties of regularity and slow divergence of the (ordinary) Brownian motion.

Remark 2. It has to be stressed that in all this article, actually we are not interested in the values themselves of the processes W and Z, but rather in their increments. This way, the fact that $W_0, Z_0 = 0$ should be considered as a mere convention, completely unessential though convenient.

1.2 Main result

Now we turn to defining the central concept of this article, which we call "arbitrage stopping times":

Definition 3 (Local minimum at right). For $X : \mathbf{R} \to \mathbf{R}$ a (deterministic) trajectory, $t \in \mathbf{R}$, we say that t is a *local minimum at right* (l.m.a.r.) for X when the following holds:

$$\exists \varepsilon > 0 \quad \forall v \in [0, \varepsilon] \quad X_{\tau+v} \geqslant X_{\tau}.$$

$$\tag{4}$$

Definition 4 (Arbitrage time). For $(X_t)_{t \in \mathbf{R}}$ a real random process, for τ a random time, we say that τ is *arbitrage* for Z when there is positive probability that $\tau(\omega)$ is a local minimum at right for $Z(\omega)$.

In this article, our goal will be to prove the following

Theorem 5. In the context of § 1.1, there does not exist any stopping time adapted to $(\mathcal{B}_t)_t$ which would be arbitrage for Z.

Remark 6. As the increments of Z are adapted to the filtration $(\mathcal{B}_t)_t$, obviously in Theorem 5 we may replace that filtration by the filtration generated by the increments of Z.

Remark 7. In this article we are only considering the case $H \neq 1/2$, but Theorem 5 is trivially valid for H = 1/2 too, since then the fBm Z is nothing but the ordinary Brownian motion W itself, for which the result follows immediately from the Markov property and the local properties of oBm.

Stated informally, Theorem 5 means that, if you are discovering the trajectory of a fractional Brownian motion along time, you cannot find a time at which you might foresee that the trajectory would go on upwards. So, this is a kind of very weak "martingale" property for the fBm, showing that the existence of correlations for it does not allow you to make anything yet.

The motivation for Theorem 5 comes from the article [1] by C. Bender, where it is explained that this result would be incompatible with the possibility, for a financial random process undergoing a fractional Brownian motion (or rather an exponential fBm), that it had an opportunity of so-called "simple arbitrage". [cf. [1, Prop. 3.3]]. So, our theorem shows that making an arbitrage on a fBm is necessarily "complicated".

1.3 Outline of the proof

The sequel of this paper is devoted to proving Theorem 5. In § 2, we will see how one can get rid of the notion of stopping time to get Theorem 5 back to a result on the trajectories of the fractional Brownian motion. In § 3, we will make the needed result on fBm's trajectories more precise, by establishing a law of iterated logarithm for some variant of the fBm. Next, an issue will be that we have to control the probability of an event being a union over a continuous infinity of t's: that issue will be handled by § 4, in which we will use regularity estimates on the fBm to get our continuous union back to a finite union. Finally, after all these simplifications it will only remain to prove some estimates on Gaussian vectors, which will be the work of § 5.

Some technical results will be postponed to appendices. In particular, in Appendix A we will compute the precise expression of the "drift operator" appearing in Lemma 15 describing what the law of the fBm becomes when you condition it by a stopping time: this formula, though not actually required to prove our main result, looks indeed intrinsically worthy to be written down to my eyes. Also, in Appendix B we will investigate some basic properties of "thick" subsets of N; and in Appendices C and D we will prove two lemmas on resp. the supremum of Gaussian processes and the inverse of nearly diagonal matrices.

2 Conditional future of the fractional Brownian motion

2.1 Preliminary definitions

To begin with, it will be convenient to set some notation for certain sets of trajectories:

Definition 8 (Sets \mathcal{P} and \mathcal{F}).

- We denote by \mathcal{P} [like "past"] the set of the (deterministic) paths $(X_u)_{u \leq 0}$ such that:
 - 1. $X_0 = 0;$
 - 2. X is locally $(H \varepsilon)$ -Hölder for all $\varepsilon > 0$;

3. $X_u \stackrel{u \to -\infty}{=} O(|u|^{H+\varepsilon})$ for all $\varepsilon > 0$.

• Similarly, we denote by \mathcal{F} [like "future"] the space of the paths $(X_v)_{v \ge 0}$ satisfying the analogues of conditions 1–3 for non-negative times.

Remark 9. With the notation of Definition 8, one has almost-surely that, for all $t \in \mathbf{R}$, $(Z(\omega)_{t+u} - Z(\omega)_t)_{u \leq 0} \in \mathcal{P}$ and $(Z(\omega)_{t+v} - Z(\omega)_t)_{v \geq 0} \in \mathcal{F}$. [cf. [2, Prop. 1.6 & Prop. 2.2.3]].

We also define a certain "drift operator":

Definition 10 ("Drift operator" **D**). Let $\mathbf{D} \colon \mathcal{P} \to \mathcal{F}$ be the linear operator such that, for $X \in \mathcal{P}$:

$$\left(\mathbf{D}X\right)_{v} \coloneqq \int_{-\infty}^{0} K(u, v) X_{u} du, \tag{5}$$

where

$$K(u,v) \coloneqq \frac{\eta}{\Pi(\eta)\Pi(-\eta)} \times \left\{ \eta \int_{-\infty}^{0} (\mathbf{1}_{s>u}\xi_{\eta-1}(s-u,v)\xi_{-\eta-1}(-s,s-u) - \xi_{\eta-1}(-u,v)\xi_{-\eta-1}(-s,-u)) ds - v(v-u)^{\eta-1}(-u)^{-\eta-1} \right\}, \quad (6)$$

where $\Pi(\cdot)$ is Euler's pi function extrapolating the factorial, and where we denote, for $r \in \mathbf{R}$, a, b > 0:

$$\xi_r(a,b) \coloneqq (a+b)^r - a^r. \tag{7}$$

Remark 11. Note that, since $H \in (0, 1)$, the integrals in (6) and (5) do converge (absolutely) indeed, and **D** is well-defined on the whole \mathcal{P} with values in \mathcal{F} .

Remark 12. The equations (5)-(6) defining \mathbf{D} , though interesting as such, shall not play an essential role in this article. What is really important to have in mind is the moral *meaning* of this operator: actually \mathbf{D} was defined so that, informally,

$$(\mathbf{D}X)_v = \mathbb{E}(Z_v | (Z_u)_{u \leqslant 0} = (X_u)_{u \leqslant 0}).$$
(8)

The formal meaning of (8) will be made clear by Lemma 15 below.

Finally we define a process called the "Lévy fractional Brownian motion", which is a kind of unilateral version of the "regular" fBm:

Definition 13 (Lévy fBm). If $(W_v)_{v \ge 0}$ is a (unilateral) ordinary Brownian motion, then the process $(Y_v)_{v \ge 0}$ defined by

$$Y_v \coloneqq C_1 \int_0^v (v-s)^\eta dW_s \tag{9}$$

(interpreted via the same integration by parts trick as in (3)) (and where we recall that C_1 is defined by (2)) is called a *Lévy fractional Brownian motion* (with Hurst parameter H)—or, more accurately, the law of this process (which (9) defines without ambiguity) is called "the law of the Lévy fBm".

Remark 14. From the regularity properties of the oBm, it is easy to check that the trajectories of the Lévy fBm lie in \mathcal{F} a.s..

2.2 Conditioning lemma

Now we can state the key lemma of this section:

Lemma 15. In the context of § 1.1, for τ a stopping time, $((Z_{\tau+v} - Z_{\tau})_{v \ge 0} - \mathbf{D}((Z_{\tau+u} - Z_{\tau})_{u \le 0}))$ is independent of \mathcal{B}_{τ} , and its law is the Lévy fBm.

Remark 16. In other words, Lemma 15 states that, conditionally to \mathcal{B}_{τ} (or, morally, knowing the past trajectory of Z until τ), the law of the future trajectory of Z is equal to a "deterministic" drift term $\mathbf{D}((Z_{\tau+u} - Z_{\tau})_{u \leq 0})$ plus a "random" noise term being a Lévy fBm.

Proof of Lemma 15. As the increments of W are adapted to $(\mathcal{B}_t)_t$, conditionally to \mathcal{B}_{τ} , the past trajectory $(W_{\tau+u}-W_{\tau})_{u\leqslant 0}$ of (the increments of) W is deterministic, while its future trajectory $(W_{\tau+v}-W_{\tau})_{v\geqslant 0}$ still has the unconditioned law of a standard oBm. Therefore, for $t \ge 0$, we split

$$Z_{\tau+t} - Z_{\tau} \stackrel{=}{=} C_1 \int_{s \in \mathbf{R}} \left((\tau + t - s)^{\eta}_{+} - (\tau - s)^{\eta}_{+} \right) dW_s$$

$$\stackrel{=}{\underset{s \leftarrow s - \tau}{=}} C_1 \int_{s \in \mathbf{R}} \left((t - s)^{\eta}_{+} - (-s)^{\eta}_{+} \right) dW_{\tau+s}$$

$$= C_1 \int_{u = -\infty}^0 \left((t - u)^{\eta} - (-u)^{\eta} \right) dW_{\tau+u} + C_1 \int_{v=0}^t (t - v)^{\eta} dW_{\tau+v}, \quad (10)$$

in which the first term is deterministic and given by some function of $(W_{\tau+u} - W_{\tau})_{u \leq 0}$, while the second term (seen as a trajectory indexed by t) has the law of the Lévy fBm indeed.

To end the proof, it remains to show that the aforementioned first term (seen as a trajectory indexed by t) is equal to $\mathbf{D}((Z_{\tau+u} - Z_{\tau})_{u \leq 0})$ indeed. Since this point is actually not needed to prove our main result, we will postpone it to Appendix A.

2.3 Reformulation of the main theorem

Thanks to Lemma 15, we will be able to get a sufficient condition for Theorem 5 in which there are no stopping times any more. For this we need first an *ad hoc* definition:

Definition 17 (Arbitrage path). We say that a deterministic path $(X_u)_{u \leq 0} \in \mathcal{P}$ is *arbitrage*, and we denote " $X \in \mathcal{A}$ ", when, for Y a Lévy fBm:

$$\mathbb{P}(0 \text{ is a local minimum at right for } (\mathbf{D}X + Y(\omega))) > 0.$$
(11)

Now, Theorem 5 will be a consequence of the following

Proposition 18. In the context of § 1.1:

$$\mathbb{P}\big(\exists t \in \mathbf{R} \quad (Z(\omega)_{t+u} - Z(\omega)_t)_{u \leq 0} \in \mathcal{A}\big) = 0.$$
(12)

Proof of Theorem 5 from Proposition 18. Let τ be any stopping time. Then, writing the law of total probability w.r.t. the σ -algebra \mathcal{B}_{τ} :

$$\mathbb{P}(\tau(\omega) \text{ is a local minimum at right for } Z(\omega)) = \mathbb{E}(\mathbb{P}(\tau \text{ is a l.m.a.r. for } Z(\omega') | \mathcal{B}_{\tau})(\omega))^{[1]} = \mathbb{E}(\mathbb{P}(0 \text{ is a l.m.a.r. for } (Z(\omega')_{\tau+v} - Z(\omega')_{\tau})_{v \ge 0} | \mathcal{B}_{\tau})(\omega)) = \mathbb{E}(\mathbb{P}(0 \text{ is a l.m.a.r. for } (\mathbf{D}((Z_{\tau+u} - Z_{\tau})_{u \le 0}) + Y(\omega')))(\omega))) = \mathbb{E}(0 \text{ whenever } ((Z(\omega)_{\tau+u} - Z(\omega)_{\tau})_{u \le 0}) \notin \mathcal{A}) = 0, \quad (13)$$

so that τ is not arbitrage.

So, in the sequel, our new goal will be to prove Proposition 18.

3 Local behaviour of fBm's trajectories

3.1 A law of the iterated logarithm for the Lévy fBm

• In all this article, we denote $[n[:= \mathbf{N} \cap [0, n) = \{0, 1, 2, ..., n-1\}$. A subset $\mathcal{I} \subset \mathbf{N}$ will be said to be thick when it has positive upper asymptotic density:

$$\limsup_{n \to \infty} \frac{|\mathcal{I} \cap \llbracket n \rrbracket|}{n} > 0.$$
(14)

A few basic results on thick subsets of \mathbf{N} are gathered in Appendix B.

The first main result of this section is the following

Lemma 19. Let $r \in (0,1)$ and let \mathcal{I} be a thick subset of \mathbf{N} ; then, for Y a Lévy fBm:

$$\liminf_{\substack{i \in \mathcal{I} \\ i \to \infty}} \frac{Y_{r^i}}{(\log i)^{1/2} r^{Hi}} = -H^{-1/2} \quad a.s..$$
(15)

Remark 20. Actually only the ' \leq ' sense of (15) (which is the harder one) will be needed in this article.

Proof of Lemma 19. It will be convenient in this proof to assume that Y is driven by some oBm W according to (9). Then, for $v \ge 0$, let us define

$$\tilde{Y}_v \coloneqq \int_{rv}^v (v-s)^\eta dW_s,\tag{16}$$

resp.

$$Y'_{v} := Y_{v} - \tilde{Y}_{v} = \int_{0}^{rv} (v - s)^{\eta} dW_{s}.$$
 (17)

First let us study the \tilde{Y}_{r^i} 's. Obviously these random variables are independent, with $Y_{r^i} / r^{Hi} \sim \mathcal{N}(0, (1-r)^{2H} / 2H) \quad \forall i$. Now, using that $\mathbb{P}(\mathcal{N}(0, 1) \leq -x) \geq$

^[1]In this computation I am using notation ' ω ' when referring to an eventuality for the conditional law $\mathbb{P}(\cdot | \mathcal{B}_{\tau})$, while ' ω ' is reserved to eventualities for the unconditioned law.

 $e^{-x^2/2}/2\sqrt{2\pi}x$ for $x \ge 1$,^[2] we get that for *i* large enough: (having fixed some arbitrary small $\varepsilon \in (0, 1)$),

$$\mathbb{P}\big(\tilde{Y}_{r^{i}} / (\log i)^{1/2} r^{Hi} \leqslant -(1-\varepsilon)(1-r)^{H} H^{-1/2}\big) \\
= \mathbb{P}\big(\mathcal{N}(0,1) \leqslant -(1-\varepsilon)\sqrt{2}(\log i)^{1/2}\big) \\
\geqslant i^{-(1-\varepsilon)^{2}} / (1-\varepsilon)4\sqrt{\pi}(\log i)^{1/2} \stackrel{i\to\infty}{=} \Omega(i^{-1}), \quad (18)$$

where " $f(i) = \Omega(g(i))$ " means that g(i) = O(f(i)). As \mathcal{I} is thick, the series $\sum_{i \in \mathcal{I}} i^{-1}$ is divergent (*cf.* Lemma 36 in Appendix B), thus so is

$$\sum_{i\in\mathcal{I}} \mathbb{P}\bigg(\frac{\tilde{Y}_{r^i}}{(\log i)^{1/2} r^{Hi}} \leqslant -(1-\varepsilon)(1-r)^H H^{-1/2}\bigg).$$
(19)

Since the events concerning the different \tilde{Y}_{r^i} 's are independent, it follows by the (second) Borel–Cantelli lemma that almost-surely there are infinitely many $i \in \mathcal{I}$ for which $\tilde{Y}_{r^i} / (\log i)^{1/2} r^{Hi} \leq -(1-\varepsilon)(1-r)^H H^{-1/2}$, so that:

$$\liminf_{\substack{i \in \mathcal{I} \\ i \to \infty}} \frac{Y_{r^i}}{(\log i)^{1/2} r^{Hi}} \leqslant -(1-\varepsilon)(1-r)^H H^{-1/2} \quad \text{a.s.},\tag{20}$$

in which the factor $(1 - \varepsilon)$ may be removed by letting $\varepsilon \to 0$.

Now let us handle the Y'_{ri} 's. One has $Y'_{ri} / r^{Hi} \sim \mathcal{N}(0, 1 - (1 - r)^{2H} / 2H)$; therefore, using that $\mathbb{P}(\mathcal{N}(0, 1) \ge x) \le e^{-x^2/2}$ for all x,^[3] we get that: (having fixed some arbitrary small $\varepsilon > 0$),

$$\mathbb{P}(Y'_{r^{i}} / (\log i)^{1/2} r^{Hi} \ge (1+\varepsilon) (1-(1-r)^{2H})^{1/2} H^{-1/2}) \\
= \mathbb{P}(\mathcal{N}(0,1) \ge (1+\varepsilon) \sqrt{2} (\log i)^{1/2}) \le i^{-(1+\varepsilon)^{2}}. \quad (21)$$

The series $\sum_{i\in\mathcal{I}} i^{-(1+\varepsilon)^2}$ is convergent since $\sum_{i\in\mathbb{N}} i^{-(1+\varepsilon)^2}$ is, thus so is

$$\sum_{i \in \mathcal{I}} \mathbb{P}\left(\frac{Y'_{r^i}}{(\log i)^{1/2} r^{Hi}} \ge (1+\varepsilon) \left(1 - (1-r)^{2H}\right)^{1/2} H^{-1/2}\right).$$
(22)

It follows by the (first) Borel–Cantelli lemma that almost-surely there are only finitely many *i*'s for which $\tilde{Y}_{ri} / (\log i)^{1/2} r^{Hi} \ge (1 + \varepsilon)(1 - (1 - r)^{2H})^{1/2} H^{-1/2}$, so that:

$$\limsup_{\substack{i \in \mathcal{I} \\ i \to \infty}} \frac{Y'_{r^i}}{(\log i)^{1/2} r^{Hi}} \leqslant (1+\varepsilon) \left(1 - (1-r)^{2H}\right)^{1/2} H^{-1/2} \quad \text{a.s.}, \tag{23}$$

in which the factor $(1 + \varepsilon)$ may be removed by letting $\varepsilon \to 0$.

Summing (20) and (23), we get an intermediate result:

Proposition 21. Under the assumptions of Lemma 19, almost-surely:

$$\liminf_{i \in \mathcal{I}} \frac{Y_{r^i}}{(\log i)^{1/2} r^{Hi}} \leqslant -\lambda(r), \tag{24}$$

where

$$\lambda(r) \coloneqq \left((1-r)^H - \left(1 - (1-r)^{2H} \right)^{1/2} \right) H^{-1/2}.$$
(25)

^[2]This is because of convexity of the density $y \mapsto \varphi(y) \coloneqq e^{-y^2/2} / \sqrt{2\pi}$ on $(-\infty, -1]$: from this property you deduce that $\int_{-\infty}^{-x} \varphi(y) dy \ge \varphi(-x)^2 / 2\varphi'(-x) = \varphi(-x) / 2x$.

^[3]This is because $\mathbb{E}(e^{x\mathcal{N}(0,1)}) = e^{x^2/2}$, from which the claimed formula follows by Markov's inequality.

So, now it remains to improve the constant $\lambda(r)$ in (24) into $H^{-1/2}$. For this, we begin with observing that the Lévy fBm is scale invariant with exponent H (by which I mean that for $a \in \mathbf{R}^*_+$, $(Y_{av} / a^H)_{v \ge 0}$ is also a Lévy fBm); therefore, Proposition 21 has the following

Corollary 22. Under the assumptions of Lemma 19, for $a \in \mathbf{R}^*_+$, one has almostsurely:

$$\liminf_{i \in \mathcal{I}} \frac{Y_{ar^i}}{a^H (\log i)^{1/2} r^{Hi}} \leqslant -\lambda(r).$$
(26)

Now let k > 1 be an arbitrary large integer; and take $l \in [k]$ such that $\mathcal{J} := \{j \in \mathbb{N} | kj + l \in \mathcal{I}\}$ is thick—existence of such an l is guaranteed by Lemma 35 in Appendix B. Then one has:

$$\liminf_{i \in \mathcal{I}} \frac{Y_{r^{i}}}{(\log i)^{1/2} r^{H_{i}}} \leq \liminf_{j \in \mathcal{J}} \frac{Y_{r^{kj+l}}}{\left(\log(kj+l)\right)^{1/2} r^{H(kj+l)}} = \liminf_{j \in \mathcal{J}} \frac{Y_{r^{l}(r^{k})^{j}}}{(r^{l})^{H}(\log j)^{1/2} (r^{k})^{H_{j}}}, \quad (27)$$

where in the last equality we used that $\left(\log(kj+l)\right)^{1/2} \stackrel{j\to\infty}{\sim} (\log j)^{1/2}$. But, applying Corollary 22 with ' $r' = r^k$, ' $a' = r^l$ and ' $\mathcal{I}' = \mathcal{J}$, the r-h.s. of (27) is bounded above by $-\lambda(r^k)$; so,

$$\liminf_{i \in \mathcal{I}} \frac{Y_{r^i}}{(\log i)^{1/2} r^{Hi}} \leqslant -\lambda(r^k).$$
(28)

Letting k tend to infinity, $\lambda(r^k)$ tends to $H^{-1/2}$, which proves the ' \leq ' sense of (15).

For the ' \geq ' sense, it is the same reasoning as for deriving (23), just replacing "Y'" by "-Y" and " $(1 - (1 - r)^{2H}) / 2H$ " by "1 / 2H".

3.2 Arbitrage condition as a limit

For all the sequel of this article, we fix some $r \in (0, 1)$ small enough (in a sense to be made precise later); we also fix arbitrarily two parameters $\alpha \in (0, 1)$ and $p \in (0, 1)$. Then we define, for all n > 0:

$$\mathcal{A}_{n} \coloneqq \left\{ (X_{u})_{u \leqslant 0} \in \mathcal{P} \middle| \operatorname{card} \{ i \in \llbracket n \llbracket | (\mathbf{D}X)_{r^{i}} \geqslant \alpha H^{-1/2} (\log i)_{+}^{1/2} r^{Hi} \} \geqslant pn \right\}.$$

$$\tag{29}$$

Then we have the following connection between \mathcal{A} and the \mathcal{A}_n 's:

Lemma 23.

$$\mathcal{A} \subset \liminf_{n \in \mathbf{N}} \mathcal{A}_n. \tag{30}$$

Proof. We prove the contrapositive inclusion. Let $(X_u)_{u \leq 0} \in \mathcal{P}$ be such that $X \notin \liminf\{\mathcal{A}_n\}$, that is, the set $\{n \mid X \notin \mathcal{A}_n\}$ is unbounded; and set

$$\mathcal{I} := \{ i \in \mathbf{N} | \ (\mathbf{D}X)_{r^i} < \alpha H^{-1/2} (\log i)_+^{1/2} r^{Hi} \}.$$
(31)

One has by definition that $|\mathcal{I} \cap [n[|/n \ge 1-p \text{ for all } n \text{ such that } X \notin \mathcal{A}_n;$ as these n are unbounded and 1-p > 0, it follows that \mathcal{I} is thick. Therefore, Lemma 19 gives that for almost-all Lévy fBm $Y(\omega)$, one has that

$$\liminf_{i\in\mathcal{I}}\frac{Y(\omega)_{r^i}}{(\log i)^{1/2}r^{Hi}}\leqslant -H^{-1/2}.$$
(32)

On the other hand, the definition of \mathcal{I} obviously implies that

$$\liminf_{i \in \mathcal{I}} \frac{(\mathbf{D}X)_{r^i}}{r^{Hi} (\log i)^{1/2}} \leqslant \alpha H^{-1/2}.$$
(33)

Summing (32) and (33), it follows that almost-surely:

$$\liminf_{i \in \mathcal{I}} \frac{(\mathbf{D}X + Y(\omega))_{r^i}}{(\log i)^{1/2} r^{Hi}} \leqslant -(1 - \alpha) H^{-1/2} < 0.$$
(34)

But (34) implies that $(\mathbf{D}X + Y(\omega))_{r^i}$ is negative for values of r^i arbitrarily close to 0, so that 0 is almost-surely not a local minimum at right for $(\mathbf{D}X + Y(\omega))$; therefore $X \notin \mathcal{A}$, which is what we wanted.

3.3 Second reformulation of the main theorem

Thanks to the work of this section, we are now able to show that the following result will be a sufficient condition for Proposition 18:

Proposition 24. In the context of \S 1.1:

$$\mathbb{P}\big(\exists t \in [0,1] \quad (Z(\omega)_{t+u} - Z(\omega)_t)_{u \leqslant 0} \in \mathcal{A}_n\big) \xrightarrow{n \to \infty} 0.$$
(35)

Proof of Proposition 18 from Proposition 24. Proposition 23 implies that

$$\{ \omega \in \Omega | \exists t \in [0,1] \quad (Z(\omega)_{t+u} - Z(\omega)_t)_{u \leq 0} \in \mathcal{A} \}$$

$$\subset \liminf_{n \to \infty} \{ \omega | \exists t \in [0,1] \quad (Z(\omega)_{t+u} - Z(\omega)_t)_{u \leq 0} \in \mathcal{A}_n \};$$
(36)

therefore, by the (first) Borel–Cantelli lemma, Proposition 24 yields that

$$\mathbb{P}\big(\exists t \in [0,1] \quad (Z(\omega)_{t+u} - Z(\omega)_t)_{u \leqslant 0} \in \mathcal{A}\big) = 0.$$
(37)

But, since the increments of the fractional Brownian motion are stationary, in (37) we may replace [0,1] by [n, n+1] for all $n \in \mathbb{Z}$; and then, by countable union, we get the wished result (12).

So, in the sequel, our new goal will be to prove Proposition 24.

4 Pathwise control via pointwise control

4.1 A regularity result

One of our issues to prove Proposition 24 is that we have to bound the probability of an event defined as a union for uncountably infinitely many t's. To overcome this issue, we will need a tool to "get rid of the trajectorial aspects" of the problem: this is the work of this section.

First, we need a little notation:

Definition 25 (Processes $\hat{\Gamma}_i$ and variables Γ_i). Within the context of § 1.1, for $i \in \mathbf{N}$, we define the following random process (indexed by $t \in \mathbf{R}$):

$$\hat{\Gamma}_{i}(\omega)_{t} \coloneqq \frac{\mathbf{D}\big((Z(\omega)_{t+u} - Z(\omega)_{t})_{u \leqslant 0}\big)(r^{i})}{r^{Hi}}.$$
(38)

We also define the following random variable:

$$\Gamma_i(\omega) \coloneqq \hat{\Gamma}_i(\omega)_0 = \frac{\mathbf{D}((Z(\omega)_u)_{u \leqslant 0})(r^i)}{r^{Hi}}.$$
(39)

Then, the main result of this section is the following

Lemma 26. In the context of this section, there exist absolute^[4] constants $C_a > 0$, $C_b < \infty$ (whose exact expressions do not matter) such that for all $i \in \mathbf{N}$, for all T > 0:

$$\mathbb{P}\big(\exists t \in [0,T] \quad |(\hat{\Gamma}_i)_t - \Gamma_i| \ge 1\big) \leqslant C_{\mathrm{b}} \exp\big(-C_{\mathrm{a}}(r^i / T)^{2H \wedge 1}\big). \tag{40}$$

Proof. First, since Z is scale invariant with exponent H (and operator **D** preserves that scale invariance), it will be enough to prove Lemma 26 for i = 0; so we will only handle that case. Then the subscript i becomes useless, so we remove it in our notation.

Because of the characterization (8) of **D**, $\hat{\Gamma}_t$ may be written as a function of W:

$$\hat{\Gamma}_t = \int_{-\infty}^t \left((t+1-s)^\eta - (t-s)^\eta \right) dW_s.$$
(41)

That shows that $\hat{\Gamma}$ is a stationary centred Gaussian random process, with

$$\operatorname{Var}(\hat{\Gamma}_{t} - \hat{\Gamma}_{0}) = \int_{\mathbf{R}} \left(\mathbf{1}_{s \leqslant t} (t+1-s)^{\eta} - \mathbf{1}_{s \leqslant 0} (1-s)^{\eta} - (t-s)^{\eta}_{+} + (-s)^{\eta}_{+} \right)^{2} ds$$
$$\stackrel{t \to 0}{=} O(t^{2H \wedge 1}). \quad (42)$$

To go further, we need the following lemma, whose proof is postponed to Appendix C:

Lemma 27. Let $(X_t)_{t \in [0,1]}$ be a centred Gaussian process such that $X_0 = 0$ a.s. and

$$\forall t, s \in [0, 1] \quad \operatorname{Var}(X_t - X_s) \leqslant |t - s|^{2\theta}$$
(43)

for some $\theta \in (0,1]$. Then it is known that, by the Kolmogorov continuity theorem, X has a continuous version. The present lemma states that, for this continuous version, the random variable $||X||(\omega) \coloneqq \sup_{t \in [0,1]} |X(\omega)_t|$ is sub-Gaussian with absolute constants, i.e. there exist constants $C_c(\theta) > 0, C_d(\theta) < \infty$ such that

$$\forall x \ge 0 \quad \mathbb{P}(\|X\| \ge x) \le C_{\mathrm{d}} \exp(-C_{\mathrm{c}} x^2). \tag{44}$$

We apply Lemma 27 in the following way. From (42), one has $\operatorname{Var}(\hat{\Gamma}_t - \hat{\Gamma}_s) \leq C_e |t-s|^{2H\wedge 1}$ for all $t, s \in [0, 1]$, for some $C_e < \infty$. Therefore, for $T \leq 1$, the random process

$$X(\omega)_t \coloneqq (C_{\rm e} T^{2H\wedge 1})^{-1/2} (\hat{\Gamma}(\omega)_{tT} - \Gamma(\omega))$$
(45)

satisfies the assumptions of Lemma 27 with ' θ ' = $H \wedge 1/2$, so that (44) yields:

$$\mathbb{P}\left(\exists t \in [0,T] \quad |(\hat{\Gamma}_i)_t - \Gamma_i| \ge x\right) \le C_{\mathrm{d}} \exp\left(-C_{\mathrm{c}} C_{\mathrm{e}}^{-1} x^2 / T^{2H\wedge 1}\right).$$
(46)

This implies (40) for $T \leq 1$, with constants not depending on T. On the other hand, up to replacing $C_{\rm b}$ by $(e^{C_{\rm a}} \vee C_{\rm b})$, (40) is automatically true for T > 1; so the proof of Lemma 26 is completed.

^[4]Remember that in this article, "absolute" constants may actually depend on H.

4.2 Third reformulation of the main result

Now, we will see how Proposition 26 allows one to find an easier sufficient condition for Proposition 24. First of all, we have to introduce a little notation: in this section, we fix some arbitrary $\alpha' \in (0, \alpha), p' \in (0, p)$, and we define \mathcal{A}'_n by the variant of Equation (29) in which α and p are replaced by resp. α' and p'; also, we fix some arbitrary $\tilde{r} \in (0, r)$, and we set

$$T_n \coloneqq \tilde{r}^n. \tag{47}$$

Now, we introduce the following events of Ω :

Definition 28 (Events A_n , A'_n , \bar{A}_n , \bar{A}_n^k and \bar{A}_n^*).

$$A_n \coloneqq \{\omega \mid (Z(\omega)_u)_{u \le 0} \in \mathcal{A}_n\}; \tag{48}$$

$$A'_{n} \coloneqq \{\omega \mid (Z(\omega)_{u})_{u \leqslant 0} \in \mathcal{A}'_{n}\}; \tag{49}$$

$$\bar{A}_n \coloneqq \{\omega \mid \exists t \in [0, T_n] \quad (Z(\omega)_{t+u} - Z(\omega)_t)_{u \leqslant 0} \in \mathcal{A}_n\};$$
(50)

$$\bar{A}_n^k \coloneqq \{\omega \mid \exists t \in [kT_n, (k+1)T_n] \quad (Z(\omega)_{t+u} - Z(\omega)_t)_{u \leqslant 0} \in \mathcal{A}_n\}; \tag{51}$$

$$\bar{A}_n^* \coloneqq \{\omega \mid \exists t \in [0,1] \quad (Z(\omega)_{t+u} - Z(\omega)_t)_{u \leqslant 0} \in \mathcal{A}_n\}.$$
(52)

Then we claim that

Lemma 29. For n large enough,

$$\omega \in \bar{A}_n \smallsetminus A'_n \quad \Rightarrow \quad \exists i \in [n[\exists t \in [0, T_n]] \quad |\hat{\Gamma}_i(\omega)_t - \Gamma_i(\omega)| \ge 1.$$
 (53)

Proof. Assume that $\omega \in \overline{A}_n \setminus A'_n$. Then the fact that $\omega \in \overline{A}_n$ means that there exists some $t \in [0, T_n]$ such that $(Z_{t+u} - Z_t)_{u \leq 0} \in \mathcal{A}_n$. For such a t, going back to the definitions (29) and (38) of \mathcal{A}_n and $\widehat{\Gamma}_i$, this means that

$$\operatorname{card}\{i \in \llbracket n \llbracket | \ (\hat{\Gamma}_i)_t \geqslant \alpha H^{-1/2} (\log i)_+^{1/2}\} \geqslant pn.$$

$$\tag{54}$$

Similarly, the fact that $\omega \notin A'_n$ means that

card{
$$i \in [n[] | \Gamma_i \ge \alpha' H^{-1/2} (\log i)_+^{1/2}$$
} < $p'n.$ (55)

Therefore, there exist at least (p - p')n indices 'i' such that $(\hat{\Gamma}_i)_t \ge \alpha H^{-1/2} \times (\log i)^{1/2}_+$ while (for the same i) $\Gamma_i < \alpha' H^{-1/2} (\log i)^{1/2}_+$. Necessarily one these indices is $\ge (p - p')n - 1$; thus, for such an *i*, one has:

$$(\hat{\Gamma}_i)_t - \Gamma_i \ge (\alpha - \alpha') H^{-1/2} (\log((p - p')n - 1))^{1/2}.$$
 (56)

But, provided $n \ge (e^{H/(\alpha - \alpha')^2} + 1) / (p - p')$, the r-h.s. of (56) is ≥ 1 ; so in the end we have found $i \in [n[, t \in [0, T_n]]$ such that $|\hat{\Gamma}_i(\omega)_t - \Gamma_i(\omega)| \ge 1$, proving the lemma.

Combining Lemma 29 with Lemma 26, we get that

$$\mathbb{P}(\bar{A}_n \smallsetminus A'_n) \leqslant \sum_{i=0}^{n-1} C_{\mathrm{b}} \exp\left(-C_{\mathrm{a}}(r^i / T_n)^{2H \wedge 1}\right),\tag{57}$$

in which the right-hand side is obviously bounded by

$$nC_{\rm b}\exp\left(-C_{\rm a}(r\,/\,\tilde{r})^{(2H\wedge1)n}\right),\tag{58}$$

which shows that $\mathbb{P}(\bar{A}_n \smallsetminus A'_n)$ decreases superexponentially in n (i.e. faster than any exponential).

Now $\bar{A}_n^* \subset \bigcup_{k \in \llbracket \lceil 1/T_n \rceil \llbracket} \bar{A}_n^k$, where $\mathbb{P}(\bar{A}_n^k) = \mathbb{P}(\bar{A}_n) \forall k$ by translation invariance, so it follows that

$$\mathbb{P}(\bar{A}_n^*) \leqslant \lceil 1/T_n \rceil \mathbb{P}(\bar{A}_n) \leqslant \lceil \tilde{r}^{-n} \rceil \left(\mathbb{P}(A_n') + \mathbb{P}(\bar{A}_n \smallsetminus A_n') \right) = \lceil \tilde{r}^{-n} \rceil \mathbb{P}(A_n') + o(1).$$
(59)

Our goal being to prove that $\mathbb{P}(\bar{A}_n^*) \to 0$ as $n \to \infty$ (that is just re-writing Proposition 24 with the notation of this section), it will be sufficient for that to prove the following

Proposition 30. $\mathbb{P}(A'_n)$ decreases superexponentially in n.

So, as A'_n corresponds to a condition on a finite-dimensional Gaussian vector, we have managed to get completely rid of the trajectorial aspects of the problem! Now our ultimate goal will be to prove Proposition 30.

Remark 31. As the "prime" symbols would be somehow cumbersome, we will drop them in the sequel, thus actually proving the superexponential decrease of $\mathbb{P}(A_n)$. Nevertheless this should not be confusing, as the constraints on α and p (and therefore on A_n) are the same as on α' and p' (and therefore on A'_n).

5 Final computations: controlling a Gaussian vector

5.1 Covariance structure

✓ In this section, for 'X' a symbol and I a discrete set, " \vec{X}_I " will be a shorthand for " $(X_i)_{i \in I}$ ".

So, our goal is to prove the superexponential decay of $\mathbb{P}(A_n)$, which can be re-written as

$$\mathbb{P}(A_n) = \mathbb{P}\left(\operatorname{card}\{i \in [n[| \Gamma_i \ge \alpha H^{-1/2}(\log i)^{1/2}_+\} \ge pn\right)$$
(60)

(where Γ_i was defined by (39)). (60) obviously implies that

$$\mathbb{P}(A_n) \leqslant \sum_{\substack{I \subset \llbracket n \rrbracket \\ |I| \ge pn}} \mathbb{P}\left(\forall i \in I \quad \Gamma_i \geqslant \alpha H^{-1/2} (\log i)_+^{1/2}\right).$$
(61)

As there are only 2^n subsets of $[n[, to prove that <math>\mathbb{P}(A_n)$ decreases superexponentially it is therefore sufficient to prove that

$$\sup_{\substack{I \subset \llbracket n \rrbracket \\ |I| \ge pn}} \mathbb{P} \Big(\forall i \in I \quad \Gamma_i \ge \alpha H^{-1/2} (\log i)_+^{1/2} \Big)$$
(62)

decreases superexponentially.

Now, by (8) one has

$$\Gamma_i(\omega) = r^{-Hi} \int_{-\infty}^0 \left((r^i - s)^\eta - (-s)^\eta \right) dW(\omega)_s;$$
(63)

therefore $\vec{\Gamma}_{\mathbf{N}}$ is a centred Gaussian vector, with:

$$\operatorname{Cov}(\Gamma_{i},\Gamma_{j}) = r^{-(i+j)H} \int_{-\infty}^{0} \left((r^{i}-s)^{\eta} - (-s)^{\eta} \right) \left((r^{j}-s)^{\eta} - (-s)^{\eta} \right) ds$$

$$= r^{-|i-j|H} \int_{-\infty}^{0} \left((r^{|i-j|} - s)^{\eta} - (-s)^{\eta} \right) \left((1-s)^{\eta} - (-s)^{\eta} \right) ds$$

$$\leqslant C_{\mathrm{f}} r^{(1/2 - |\eta|)|i-j|}, \quad (64)$$

for some absolute constant $C_{\rm f} < \infty$. Therefore, provided r was chosen small enough, we have the following control on the covariance matrix of $\vec{\Gamma}_{\rm N}$:

$$\operatorname{Cov}(\Gamma_i, \Gamma_j) = \sigma^2$$
 for $i = j;$ (65)

$$|\operatorname{Cov}(\Gamma_i, \Gamma_j)| \leqslant \sigma^2 \varepsilon^{|i-j|} \qquad \text{for } i \neq j, \tag{66}$$

where $\varepsilon > 0$ is some small parameter which will be fixed later, and where $\sigma := \operatorname{Var}(\Gamma)^{1/2} > 0$ (since $H \neq 1/2$).

5.2 Density estimates

To exploit (65)-(66), we need the following lemma (whose proof is postponed to Appendix D):

Lemma 32. For $n \in \mathbf{N}$, $\varepsilon > 0$, let $\mathbf{A} \coloneqq ((a_{ij}))_{i,j \in [n[]}$ be a square matrix such that $a_{ii} = 1 \ \forall i \ and \ |a_{ij}| \leq \varepsilon^{|i-j|} \ \forall i \neq j$. Then:

$$\det \mathbf{A} \ge \exp(-n\Phi_{\mathrm{g}}(\varepsilon)\varepsilon^{2}),\tag{67}$$

where $\Phi_{g}: (0, \infty) \to [1, \infty]$ is some absolute function (in particular, not depending on n) such that $\Phi_{g}(\varepsilon) \stackrel{\varepsilon \to 0}{\to} 1$ —we will call such a function a quasi-one function. In particular, provided ε is small enough, **A** is invertible. Then the present lemma asserts moreover that, denoting $\mathbf{A}^{-1} \coloneqq ((\mathbf{b}_{ij}))_{i,j}$:

$$|b_{ij}| \leq 2^{|i-j|-1} (\Phi_{\mathbf{h}}(\varepsilon)\varepsilon)^{|i-j|} \qquad \forall i \neq j;$$
(68)

$$|b_{ii} - 1| \leqslant 2\Phi_{\rm i}(\varepsilon)\varepsilon^2 \qquad \qquad \forall i, \tag{69}$$

for Φ_i and Φ_h some other absolute "quasi-one functions".

We apply Lemma 32 to the covariance matrix of $\vec{\Gamma}_{\llbracket n \llbracket}$ (assuming ε was chosen small enough so that Φ_{g} is finite); then, the formula for the density of Gaussian vectors gives that:

$$\frac{d\mathbb{P}(\vec{\Gamma}_{\llbracket n \llbracket} = \vec{\gamma}_{\llbracket n \rrbracket})}{d\gamma} \leqslant \left(\frac{\exp(\Phi_{g}(\varepsilon)\varepsilon^{2})}{2\pi\sigma^{2}}\right)^{n/2} \times \\
\exp\left(\frac{1}{2\sigma^{2}}\left((-1 + 2\Phi_{i}(\varepsilon)\varepsilon^{2})\sum_{i \in \llbracket n \rrbracket}\gamma_{i}^{2} + \frac{1}{2}\sum_{\substack{i,j \in \llbracket n \rrbracket\\i \neq j}}|\gamma_{i}||\gamma_{j}| \times (2\Phi_{h}(\varepsilon)\varepsilon)^{|i-j|}\right)\right). \quad (70)$$

Bounding above $|\gamma_i||\gamma_j|$ by $\frac{1}{2}(\gamma_i^2 + \gamma_j^2)$, that is bounded again by

$$\left(\Phi_{j}(\varepsilon) / 2\pi\sigma^{2}\right)^{n/2} \exp\left(-\frac{1}{2\Phi_{k}(\varepsilon)\sigma^{2}} \sum_{i \in \llbracket n \rrbracket} \gamma_{i}^{2}\right),\tag{71}$$

where

$$\Phi_{\mathbf{j}}(\varepsilon) \coloneqq \exp(\Phi_{\mathbf{g}}(\varepsilon)\varepsilon^2) \tag{72}$$

and

$$\Phi_{\mathbf{k}}(\varepsilon) \coloneqq \left(1 - 2\Phi_{\mathbf{i}}(\varepsilon)\varepsilon^{2} - \frac{1}{2}\sum_{z \in \mathbf{Z}^{*}} (2\Phi_{\mathbf{h}}(\varepsilon)\varepsilon)^{|z|}\right)_{+}^{-1}$$
(73)

are "quasi-one functions" again.

In the sequel we assume that ε was chosen small enough so that $\Phi_j(\varepsilon), \Phi_k(\varepsilon) < \infty$; and we define the following vectorial random variable (which we are actually only interested in through its law):

Definition 33 (Variable $\vec{\Pi}_{N}$). $\vec{\Pi}_{N}$ is a random vector on \mathbf{R}^{N} whose entries are i.i.d. $\mathcal{N}(0, \Phi_{k}(\varepsilon)\sigma^{2})$.

Then, Equation (71) can be rephrased into:

$$\frac{d\mathbb{P}(\vec{\Gamma}_{\llbracket n \rrbracket} = \vec{\gamma}_{\llbracket n \rrbracket})}{d\mathbb{P}(\vec{\Pi}_{\llbracket n \rrbracket} = \vec{\gamma}_{\llbracket n \rrbracket})} \leqslant (\Phi_{j}(\varepsilon)\Phi_{k}(\varepsilon))^{n/2} \quad \text{uniformly in } \vec{\gamma}_{\llbracket n \rrbracket}.$$
(74)

Therefore, for $I \subset [n]$ with $|I| \ge pn$:

$$\mathbb{P}\left(\forall i \in I \quad \Gamma_i \geqslant \alpha H^{-1/2} (\log i)_+^{1/2}\right) \leqslant (\Phi_{\mathbf{j}}(\varepsilon) \Phi_{\mathbf{k}}(\varepsilon))^{n/2} \mathbb{P}\left(\forall i \in I \quad \Pi_i \geqslant \alpha H^{-1/2} (\log i)_+^{1/2}\right).$$
(75)

But

$$\mathbb{P}\left(\forall i \in I \quad \Pi_{i} \ge \alpha H^{-1/2} (\log i)_{+}^{1/2}\right) \\
= \prod_{i \in I} \mathbb{P}\left(\Pi_{i} \ge \alpha H^{-1/2} (\log i)_{+}^{1/2}\right) \\
= \prod_{i \in I} \mathbb{P}\left(\mathcal{N}(0, 1) \ge C_{1} (\log i)_{+}^{1/2}\right) \\
\leqslant \prod_{i \in I} \exp\left(-C_{1}^{2} (\log i)_{+} / 2\right) \\
= \prod_{i \in I} (i \lor 1)^{-C_{1}^{2}/2} \\
\leqslant (|I| - 1)_{+}!^{-C_{1}^{2}/2} \leqslant (\lceil pn \rceil - 1)_{+}!^{-C_{1}^{2}/2} \quad (76)$$

(with $C_{\rm l} \coloneqq \alpha H^{-1/2} / \Phi_{\rm k}(\varepsilon)^{1/2} \sigma$), where the penultimate inequality comes from ordering $I \eqqcolon \{i_0, i_1, \ldots, i_{|I|-1}\}$ with $i_0 < i_1 < \cdots$, and observing that then $i_j \ge j$ for all j, so that $(i_j \lor 1)^{-C_1^2/2} \le (j \lor 1)^{-C_1^2/2}$.

Combining (75) with (76) shows that $\mathbb{P}(\Gamma_i \ge \alpha H^{-1/2}(\log i)^{1/2}_+ \quad \forall i \in I)$ decreases superexponentially in n uniformly in I, which finally proves Proposition 30 and hence Theorem 5.

A Conditional expectation of the fBm

This appendix is devoted to ending the proof of Lemma 15 initiated in § 2.2. At the point we have got to, what remains to do is showing that

$$C_1 \int_{s=-\infty}^{0} \left((v-s)^{\eta} - (-s)^{\eta} \right) dW_{\tau+s}$$
(77)

(seen as a trajectory indexed by $v \in \mathbf{R}_+$) is actually equal to $\mathbf{D}((Z_{\tau+u} - Z_{\tau})_{u \leq 0})$ with \mathbf{D} defined by (5)-(6), where W is the ordinary Brownian motion driving the fBm Z. To alleviate notation, actually we will only prove this result for $\tau \equiv 0$, the original case being the same up to time translation of the increments (hence the informal definition (8) of \mathbf{D}).

The starting point for our computation is the *Pipiras-Taqqu formula*, which says that Equation (1) defining the past increments of Z as a function of the past increments of W has an "inverse" giving back the past increments of W from the past increments of Z:

Proposition 34 ([3, Cor. 1.1]). In the context of § 1.1, one has almost-surely, for all t:

$$W_t = \frac{C_1^{-1}}{\Pi(\eta)\Pi(-\eta)} \int_{\mathbf{R}} \left((t-s)_+^{-\eta} - (-s)_+^{-\eta} \right) dZ_s.$$
(78)

(Recall that $\Pi(\bullet)$ is Euler's pi function extrapolating the factorial).

From now on in this appendix, it will be convenient to shorthand "1 / $\Pi(\eta)\Pi(-\eta)$ " into " C_H ".

So, let us use (78) to get (5)-(6). First, like (1), Equations (77) and (78) have to be interpreted by integrating by parts: for $v \ge 0, s \le 0$, that means resp. that:

(77) =
$$\eta C_1 \int_{-\infty}^0 \xi_{\eta-1}(-s, v) W_s ds;$$
 (79)

$$\frac{W_t}{C_1^{-1}C_H} = \eta \int_{-\infty}^t \xi_{-\eta-1}(t-s,-t)(Z_s-Z_t)ds + \eta \int_t^0 (-s)^{-\eta-1}Z_sds + (-t)^{-\eta}Z_t, \quad (80)$$

where we recall that $\xi_r(a,b) \coloneqq (a+b)^r - a^r$. Hence, (77) is equal to:

$$\eta^2 C_H \int_{s=-\infty}^0 \int_{u=-\infty}^s \xi_{\eta-1}(-s,v)\xi_{-\eta-1}(s-u,-s)(Z_u-Z_s)dsdu$$
(81)

$$+ \eta^2 C_H \int_{s=-\infty}^0 \int_{u=s}^0 \xi_{\eta-1}(-s,v)(-u)^{-\eta-1} Z_u ds du$$
(82)

$$+ \eta C_H \int_{-\infty}^{0} \xi_{\eta-1}(-s,v)(-s)^{-\eta} Z_s ds.$$
(83)

Now we are going to re-write each of the terms (81)-(83) as an integral against $Z_u du$, in order to get (5)-(6). First, Term (83) is already of the wanted form, up to renaming 's' into 'u'. Next, Term (82) simplifies into:

$$(82) = \eta^{2} C_{H} \int_{u=-\infty}^{0} \left(\int_{s=-\infty}^{u} \xi_{\eta-1}(-s,v) ds \right) (-u)^{-\eta-1} Z_{u} du$$
$$= -\eta C_{H} \int_{-\infty}^{0} \left[\xi_{\eta}(-s,v) \right]_{s=-\infty}^{u} (-u)^{-\eta-1} Z_{u} du$$
$$= -\eta C_{H} \int_{-\infty}^{0} \xi_{\eta}(-u,v) (-u)^{-\eta-1} Z_{u} du. \quad (84)$$

Term (81) is the hardest to get into the wanted form, because splitting naively the factor $(Z_u - Z_s)$ would yield divergent integrals. To bypass that problem, we first make a truncation: for ε a small positive number,

$$(81) \approx \eta^2 C_H \int_{s=-\infty}^0 \int_{u=-\infty}^{(1+\varepsilon)s} \xi_{\eta-1}(-s,v) \xi_{-\eta-1}(s-u,-s) (Z_u - Z_s) ds du$$

= $\eta^2 C_H \iint_{\substack{s<0\\u<(1+\varepsilon)s}} \xi_{\eta-1}(-s,v) \xi_{-\eta-1}(s-u,-s) Z_u ds du$ (85)

$$-\eta^2 C_H \iint_{\substack{s<0\\u<(1+\varepsilon)s}} \xi_{\eta-1}(-s,v)\xi_{-\eta-1}(s-u,-s)Z_s ds du.$$
(86)

By the change of variables $(s, u) \leftarrow (u - s, u)$,

$$(85) = \eta^2 C_H \int_{u=-\infty}^0 \left(\int_{s=-\infty}^{\varepsilon u/(1+\varepsilon)} \mathbf{1}_{s>u} \xi_{\eta-1}(s-u,v) \xi_{-\eta-1}(-s,s-u) ds \right) Z_u du;$$
(87)

and by the change of variables $(s, u) \leftarrow (u - s, s)$,

$$(86) = -\eta^{2} C_{H} \int_{u=-\infty}^{0} \left(\int_{s=-\infty}^{\varepsilon u} \xi_{\eta-1}(-u,v) \xi_{-\eta-1}(-s,-u) ds \right) Z_{u} du$$
$$\approx -\eta^{2} C_{H} \int_{u=-\infty}^{0} \left(\int_{s=-\infty}^{\varepsilon u/(1+\varepsilon)} \xi_{\eta-1}(-u,v) \xi_{-\eta-1}(-s,-u) ds \right) Z_{u} du, \quad (88)$$

where by " \approx " we mean that, for all v, the difference between the two members from either side of the ' \approx ' sign tends to 0 as $\varepsilon \to 0$, as one can check by simple estimates. So,

(81)
$$\approx \eta^2 C_H \int_{-\infty}^0 \left(\int_{-\infty}^{\varepsilon u/(1+\varepsilon)} J(v, u, s) ds \right) Z_u du,$$
 (89)

with

$$J(v, u, s) \coloneqq \mathbf{1}_{s>u} \xi_{\eta-1}(s-u, v) \xi_{-\eta-1}(-s, s-u) - \xi_{\eta-1}(-u, v) \xi_{-\eta-1}(-s, -u).$$
(90)

But $\int_{0}^{0} J(v, u, s) ds$ does converge, so, letting ε tend to 0, we get in the end:

(81) =
$$\eta^2 C_H \int_{-\infty}^0 \left(\int_{-\infty}^0 J(v, u, s) ds \right) Z_u du.$$
 (91)

Summing (83), (84) and (91), and observing that $\xi_{\eta-1}(-u,v)(-u)^{-\eta}-\xi_{\eta}(-u,v)(-u)^{-\eta-1} = -v(v-u)^{\eta-1}(-u)^{-\eta-1}$, finally yields Equation (5)-(6).

B On thick subsets of N

Remember that we have defined a subset $\mathcal{I} \subset \mathbf{N}$ to be *thick* when

$$\limsup_{n \to \infty} \frac{|\mathcal{I} \cap \llbracket n \rrbracket|}{n} > 0, \tag{92}$$

where $[n] := \{0, ..., n-1\}$ is the set of the *n* smallest natural integers. In this appendix, we will prove two basic properties of thick subsets of **N** which we used in the body of the article:

Lemma 35. Let \mathcal{I} be a thick set of integers and let k > 1. Then there exists $l \in [\![k]\![$ such that the set

$$\{j \in \mathbf{N} | jk + l \in \mathcal{I}\}$$
(93)

is thick.

Proof. Let \mathcal{I} be a subset of \mathbf{N} ; and for $l \in \llbracket k \rrbracket$, denote the set defined by (93) as \mathcal{J}_l . Then the \mathcal{J}_l 's make a partition of \mathcal{I} ; so, for $n \in \mathbf{N}$:

$$\mathcal{I} \cap \llbracket n \llbracket = \bigsqcup_{l \in \llbracket k \rrbracket} \left((k\mathcal{J}_l + l) \cap \llbracket n \rrbracket \right) = \bigsqcup_{l \in \llbracket k \rrbracket} \left(k \left(\mathcal{J}_l \cap \llbracket \lceil (n-l) / k \rceil \rrbracket \right) + l \right).$$
(94)

Thus,

$$\frac{|\mathcal{I} \cap \llbracket n \rrbracket|}{n} = \sum_{l \in \llbracket k \rrbracket} \frac{|\mathcal{J}_l \cap \llbracket \lceil (n-l) / k \rceil \llbracket |}{n}$$
$$= \sum_{l \in \llbracket k \rrbracket} \frac{\lceil (n-l) / k \rceil}{n} \frac{|\mathcal{J}_l \cap \llbracket \lceil (n-l) / k \rceil \llbracket |}{\lceil (n-l) / k \rceil}; \quad (95)$$

and therefore, letting $n \to \infty$:

$$\limsup_{n \to \infty} \frac{|\mathcal{I} \cap \llbracket n \rrbracket|}{n} \leqslant \sum_{l \in \llbracket k \rrbracket} k^{-1} \limsup_{m \to \infty} \frac{|\mathcal{J}_l \cap \llbracket m \rrbracket|}{m}.$$
(96)

So, for \mathcal{I} to be thick, one at least of the \mathcal{J}_l 's has to be thick, proving the lemma. \Box

Lemma 36. If $\mathcal{I} \subset \mathbf{N}$ is thick, then the series

$$\sum_{i\in\mathcal{I}}i^{-1}\tag{97}$$

is divergent.

Proof. Assume that \mathcal{I} is thick and let

$$p \coloneqq \limsup_{n \to \infty} \frac{|\mathcal{I} \cap \llbracket n \rrbracket|}{n} \tag{98}$$

(which then is positive); and fix arbitrary p > p' > p'' > 0.

Then, there will be arbitrarily large $n \in \mathbf{N}$ such that

$$|\mathcal{I} \cap \llbracket n \llbracket| \ge p'n,\tag{99}$$

so that we can define the following sequence by induction: $n_0 = 1$, and n_{k+1} is the smallest *n* satisfying (99) such that $n_{k+1} \ge n_k / (p' - p'')$.

Now, observe that

$$|\mathcal{I} \cap [n_k, n_{k+1})| \ge |\mathcal{I} \cap [[n_{k+1}[[|-n_k] \ge p'n_{k+1} - (p'-p'')n_{k+1}] = p''n_{k+1}, \quad (100)$$

so that

$$\sum_{\mathcal{I} \cap [n_k, n_{k+1})} i^{-1} \ge |\mathcal{I} \cap [n_k, n_{k+1})| \inf_{i \in \mathcal{I} \cap [n_k, n_{k+1})} i^{-1} \ge p'' n_{k+1} \times n_{k+1}^{-1} = p'', \quad (101)$$

and therefore:

$$\sum_{\substack{i\in\mathcal{I}\\i\geqslant 1}} i^{-1} \geqslant \sum_{k=0}^{\infty} \sum_{\substack{i\in\mathcal{I}\\n_k \le i < n_{k+1}}} i^{-1} \geqslant \sum_{k=0}^{\infty} p'' = \infty,$$
(102)

proving the lemma.

C Explicit estimate for the supremum of Gaussian processes

Proof of Lemma 27. Let $\theta \in (0, 1]$ and let X satisfying the assumptions of the lemma. Then obviously, for the continuous version of X:

$$\|X\|(\omega) \leqslant \sum_{i=0}^{\infty} \sup_{a \in [2^{i}[]} |X(\omega)_{a2^{-i}} - X(\omega)_{(a+1)2^{-i}}|.$$
 (103)

Therefore, for $(\gamma_i)_i$ a sequence of positive real numbers such that $\sum_i \gamma_i = 1$, one has that, for all $x \ge 0$:

$$\mathbb{P}(\|X\| \ge x) \le \sum_{i=0}^{\infty} \mathbb{P}(\sup_{a \in [2^{i}[]} |X_{a2^{-i}} - X_{(a+1)2^{-i}}| \ge \gamma_{i}x)$$
$$\le \sum_{i} 2^{i} \sup_{a} \mathbb{P}(|X_{a2^{-i}} - X_{(a+1)2^{-i}}| \ge \gamma_{i}x). \quad (104)$$

But, uniformly in a,

$$\mathbb{P}(|X_{a2^{-i}} - X_{(a+1)2^{-i}}| \ge \gamma_i x) = \mathbb{P}\big(\mathcal{N}(0,1) \ge \gamma_i x / \operatorname{Var}^{1/2}(X_{a2^{-i}} - X_{(a+1)2^{-i}})\big)$$
$$\leqslant \mathbb{P}\big(\mathcal{N}(0,1) \ge 2^{i\theta}\gamma_i x\big) \le 2\exp\big(-2^{2i\theta-1}(\gamma_i x)^2\big); \quad (105)$$

so, taking $\gamma_i \coloneqq (1 - 2^{-\theta/2}) 2^{-i\theta/2}$:

$$\mathbb{P}(\|X\| \ge x) \le \sum_{i=0}^{\infty} 2^{i+1} \exp\left(-(1-2^{-\theta/2})^2 \times 2^{i\theta-1}x^2\right).$$
(106)

Provided $x \ge 2 / (1 - 2^{-\theta/2})\theta^{1/2} =: C_{\rm m}(\theta)$, one has (bounding $2^{i\theta}$ below by $(1 + i\theta \log 2))$

$$\exp\left(-2^{i\theta}x^2 / 2(1-2^{-\theta/2})^2\right) \leqslant \exp\left(-x^2 / 2(1-2^{-\theta/2})^2 - \log(4)i\right), \quad (107)$$

so that, for $x \ge C_{\rm m}$:

$$\mathbb{P}(\|X\| \ge x) \le \left(\sum_{i=0}^{\infty} 2^{i+1} 4^{-i}\right) \exp\left(-x^2 / 2(1 - 2^{-\theta/2})^2\right) =: 4e^{-C_{\rm c}(\theta)x^2}.$$
(108)

On the other hand, for $x < C_{\rm m}$ one has obviously $\mathbb{P}(||X|| \ge x) \le 1$; so Equation (44) follows with $C_{\rm d} := 4 \lor e^{C_{\rm c}C_{\rm m}^2}$.

D Almost diagonal matrices

Proof of Lemma 32. Consider **A** satisfying the assumptions of the lemma, and denote $\mathbf{I}_n - \mathbf{A} =: \mathbf{H}$. The first part of this proof will consist in deriving estimates on the entries of **H** and its powers. Denote respectively

$$\mathbf{H} \coloneqq ((h_{ij}))_{i,j \in \llbracket n \llbracket}; \tag{109}$$

$$\mathbf{H}^{k} \coloneqq ((h_{ij}^{(k)}))_{i,j \in \llbracket n \rrbracket} \qquad \forall k \ge 0.$$
(110)

Then the assumptions of the lemma ensure that one has $|h_{ii}| = 0 \ \forall i$, resp. $|h_{ij}| \leq \varepsilon^{|i-j|} \ \forall i \neq j$, and hence

$$|h_{ij}^{(k+1)}| \leq \sum_{i' \in [n[]} |h_{ii'}| |h_{i'j}^{(k)}| = \sum_{i' \neq i} \varepsilon^{|i-i'|} |h_{i'j}^{(k)}| \quad \forall i, j \quad \forall k.$$
(111)

That suggests to define by induction:

$$\begin{cases} \mathfrak{h}_{z}^{(0)} \coloneqq \mathbf{1}_{z=0} & \forall z \in \mathbf{Z} \\ \mathfrak{h}_{z}^{(k+1)} \coloneqq \sum_{z' \neq z} \varepsilon^{|z-z'|} \mathfrak{h}_{z'}^{(k)} & \forall z \in \mathbf{Z} \quad \forall k \ge 0, \end{cases}$$
(112)

so that one has

$$|h_{ij}^{(k)}| \leq \mathfrak{h}_{i-j}^{(k)} \quad \forall i, j \quad \forall k.$$
(113)

The interest of having introduced the $\mathfrak{h}_z^{(k)}$'s is that these are easier to bound than the $h_{ij}^{(k)}$'s themselves. To bound the $\mathfrak{h}_z^{(k)}$'s, we begin with observing that one has obviously by induction:

$$\mathfrak{h}_{z}^{(k)} = \sum_{\substack{(0,s_{1},s_{2},\ldots,s_{k-1},z)\\0\neq s_{1},s_{1}\neq s_{2},\ldots,s_{k-1}\neq z}} \varepsilon^{\sum_{i\in\llbracket k\rrbracket}|s_{i}-s_{i+1}|} \\
= \sum_{n\geqslant 0} \operatorname{card}\Big\{(0,s_{1},s_{2},\ldots,s_{k-1},z)\Big| \ s_{i+1}\neq s_{i} \ \forall i \ \text{and} \ \sum_{i}|s_{i}-s_{i+1}| = n\Big\}\varepsilon^{n}.$$
(114)

To bound the cardinality appearing in (114), we observe that a (k + 1)-tuple $(0, s_1, s_2, \ldots, s_{k-1}, z)$ such that $s_{i+1} \neq s_i \forall i$ and $\sum_i |s_i - s_{i+1}| = n$ (we will call such a (k+1)-tuple as *valid*) can be *coded* by a word of *n* symbols from $\{+, +|, -, -|\}$, in the following way: successively, for each *i* we write $(|s_{i+1} - s_i| - 1)$ symbols "sgn $(s_{i+1} - s_i)$ " followed by a symbol "sgn $(s_{i+1} - s_i)$ "—for instance, for k = 5, z = 2, n = 8, one would have

$$(0, 1, 4, 2, 1, 2) \mapsto "+_{|} + + +_{|} - -_{|} -_{|} +_{|}".$$

$$(115)$$

Obviously such a coding in injective. Moreover, for given k, z, n, an *n*-character word may be the image of a valid (k + 1)-tuple only if z and n have the same parity, that the word contains (n + z) / 2 symbols from $\{+, +_{\mid}\}$ vs. (n - z) / 2symbols from $\{-, -_{\mid}\}$, and that k exactly of the n symbols, necessarily including the last one, are from $\{+_{\mid}, -_{\mid}\}$. Henceforth:

$$\operatorname{card}\left\{ (0, s_1, s_2, \dots, s_{k-1}, z) \middle| s_{i+1} \neq s_i \; \forall i \text{ and } \sum_i |s_i - s_{i+1}| = n \right\} \\ \leqslant \mathbf{1}_{2|n-z} \binom{n}{(n-1)(n-2)\binom{n-1}{k-1}}.$$
(116)

In the end, combining (113), (114) and (116):

$$|h_{ij}^{(k)}| \leq \sum_{m \geq 0} \binom{|i-j|+2m}{m} \binom{|i-j|+2m-1}{k-1} \varepsilon^{|i-j|+2m} \quad \forall i, j \quad \forall k.$$
(117)

After these preliminary estimates, let us turn to proving the lemma itself. We begin with the first part, namely, bounding det **A** below. For $\mathbf{X} =: ((x_{ij}))_{i,j}$ an $n \times n$ matrix, denote

$$\|\mathbf{X}\| \coloneqq \sup_{j \in [\![n]\!]} \sum_{i \in [\![n]\!]} |x_{ij}| :$$
(118)

 $\|\cdot\|$ is the operator norm of **X** when seen as an operator from $\ell^1([n])$ into itself, so it is sub-multiplicative. Then, the formula

$$\log(\mathbf{I}_n - \mathbf{H}) = \sum_{k=1}^{\infty} k^{-1} \mathbf{H}^k$$
(119)

converges as soon as $\|\mathbf{H}\| < 1$, then yielding:

$$|\operatorname{tr} \log(\mathbf{I}_{n} - \mathbf{H})| \leq \sum_{k=1}^{\infty} k^{-1} |\operatorname{tr} \mathbf{H}^{k}| \leq 0 + \frac{1}{2} |\operatorname{tr} \mathbf{H}^{2}| + n \sum_{k \geq 3} k^{-1} ||\mathbf{H}^{k}|| \leq \frac{1}{2} |\operatorname{tr} \mathbf{H}^{2}| + n \sum_{k \geq 3} k^{-1} ||\mathbf{H}||^{k}.$$
(120)

But the assumptions on the entries of \mathbf{H} imply that

$$\|\mathbf{H}\| \leqslant \sum_{z \in \mathbf{Z}^*} \varepsilon^{|z|} = \frac{2\varepsilon}{1 - \varepsilon},\tag{121}$$

which is < 1 as soon as $\varepsilon < 1/3$; and on the other hand, we get from (117) that, for all i,

$$|h_{ii}^{(2)}| \leq \sum_{m \geq 1} \binom{2m}{m} (2m-1)\varepsilon^{2m} \leq 2\varepsilon^2 + \sum_{m \geq 2} 2^{2m} (2m-1)\varepsilon^{2m} =: 2\Phi_{\mathbf{n}}(\varepsilon)\varepsilon^2, \quad (122)$$

so that $|\operatorname{tr} \mathbf{H}^2| \leq 2n\Phi_n(\varepsilon)\varepsilon^2$. In the end:

$$\det \mathbf{A} = \det \exp \log(\mathbf{I}_n - \mathbf{H}) = \exp \operatorname{tr} \log(\mathbf{I}_n - \mathbf{H})$$

$$\geq \exp\left(-n\Phi_{\mathbf{n}}(\varepsilon)\varepsilon^2 - n\sum_{k\geq 3} k^{-1} \left(\frac{2\varepsilon}{1-\varepsilon}\right)^k\right) =: \exp(-n\Phi_{\mathbf{g}}(\varepsilon)\varepsilon^2), \quad (123)$$

which is Equation (67).

Now let us handle the second part of the lemma, namely, bounding the entries of $(\mathbf{A}^{-1} - \mathbf{I}_n)$. Provided $\|\mathbf{H}\| < 1$, one has

$$\mathbf{A}^{-1} - \mathbf{I}_n = \sum_{k=1}^{\infty} \mathbf{H}^k, \tag{124}$$

so that

$$|b_{ij} - \mathbf{1}_{i=j}| \leq \sum_{k=1}^{\infty} |h_{ij}^{(k)}| \quad \forall i, j.$$

$$(125)$$

To bound the r-h.s. of (125), we write that, starting from (117):

$$\sum_{k \ge 1} |h_{ij}^{(k)}| \le \sum_{m \ge 0} \binom{|i-j|+2m}{m} \binom{|i-j|+2m}{k-1} \binom{|i-j|+2m-1}{k-1} \varepsilon^{|i-j|+2m} = \sum_{m \ge 0} \binom{|i-j|+2m}{m} \mathbf{1}_{|i-j|+2m \ge 1} 2^{|i-j|+2m-1} \varepsilon^{|i-j|+2m} = \left(\sum_{m \ge 0} \mathbf{1}_{|i-j|+2m \ge 1} \binom{|i-j|+2m}{m} (4\varepsilon^2)^m\right) \times 2^{|i-j|-1} \varepsilon^{|i-j|}.$$
 (126)

But we observe that, for $z \ge 1$, x > 0,

$$\sum_{m \ge 0} {\binom{z+2m}{m}} x^m = 1 + \sum_{m \ge 1} {\binom{z+2m}{m}} x^m \leqslant 1 + \sum_{m \ge 1} \frac{(z+2m)^m}{m!} x^m$$
$$\leqslant 1 + \sum_{m \ge 1} \frac{(2z)^m + (4m)^m}{m!} x^m$$
$$\leqslant 1 + \sum_{m \ge 1} \frac{(2zx)^m}{m!} + \sum_{m \ge 1} (4ex)^m = e^{2zx} + \frac{4ex}{1-4ex}$$
$$\leqslant \left(e^{2x} + \frac{4ex}{1-4ex}\right)^z =: \Phi_0(x)^z, \quad (127)$$

so that here, for $i \neq j$:

$$\sum_{k=1}^{\infty} |h_{i,j}^{(k)}| \leq \Phi_{\rm o}(4\varepsilon^2)^{|i-j|} \times 2^{|i-j|-1}\varepsilon^{|i-j|} =: 2^{|i-j|-1}(\Phi_{\rm h}(\varepsilon)\varepsilon)^{|i-j|},$$
(128)

which is Equation (68).

Equation (69) for the case i = j is derived in the same way as (68), with just a few minor differences at the beginning of the computation: namely, in the l-h.s. of (126), we treat apart the cases "k = 1" (which yields zero here since $h_{ii} = 0$ by assumption) and "k = 2" (which has already been handled by (122)); then all the sequel is the same.

Remark 37. The bounds (67)-(69) of Lemma 32 are optimal at first order. Actually this is much more than needed to prove Theorem 5, and we could have got a sufficient result with a shorter proof; but it seemed interesting to me to state the sharp version of the lemma.

References

- Christian Bender. Simple arbitrage. Ann. Appl. Probab., 22(5):2067–2085, 2012.
- [2] Ivan Nourdin. Selected aspects of fractional Brownian motion, volume 4 of Bocconi & Springer Series. Springer, Milan; Bocconi University Press, Milan, 2012.
- [3] Vladas Pipiras and Murad S. Taqqu. Deconvolution of fractional Brownian motion. J. Time Ser. Anal., 23(4):487–501, 2002.