

# Asynchronous stochastic convex optimization

John C. Duchi<sup>1</sup> Sorathan Chaturapruek<sup>2</sup> Christopher Ré<sup>2</sup>

Stanford University

Departments of <sup>1</sup>Statistics, <sup>1</sup>Electrical Engineering, and <sup>2</sup>Computer Science

{jduchi, sorathan, chrismre}@stanford.edu

## Abstract

We show that asymptotically, completely asynchronous stochastic gradient procedures achieve optimal (even to constant factors) convergence rates for the solution of convex optimization problems under nearly the same conditions required for asymptotic optimality of standard stochastic gradient procedures. Roughly, the noise inherent to the stochastic approximation scheme dominates any noise from asynchrony. We also give empirical evidence demonstrating the strong performance of asynchronous, parallel stochastic optimization schemes, demonstrating that the robustness inherent to stochastic approximation problems allows substantially faster parallel and asynchronous solution methods.

## 1 Introduction

We study a natural asynchronous stochastic gradient method for the solution of minimization problems of the form

$$\text{minimize } f(x) := \mathbb{E}_P[F(x; W)] = \int_{\Omega} F(x; \omega) dP(\omega), \quad (1)$$

where  $x \mapsto F(x; \omega)$  is convex for each  $\omega \in \Omega$ ,  $P$  is a probability distribution on  $\Omega$ , and the vector  $x \in \mathbb{R}^d$ . Stochastic gradient techniques for the solution of problem (1) have a long history in optimization, starting from the early work of Robbins and Monro [29] and continuing on through Ermoliev [12] and Polyak and Juditsky [26] and Nemirovski et al. [23]. The latter two papers show how certain long stepsizes and averaging techniques yield more robust and asymptotically optimal optimization schemes, and we show how their results extend to practical parallel and asynchronous optimization settings.

We consider an extension of previous stochastic gradient methods to a natural family of *asynchronous* gradient methods (see, e.g., the book of Bertsekas and Tsitsiklis [5]), where multiple processors can draw samples from the distribution  $P$  and asynchronously perform updates to a centralized parameter vector  $x$ . Our iterative scheme is based on the HOGWILD! algorithm of Niu et al. [25], which is designed to asynchronously solve certain stochastic optimization problems in multi-core environments, though our analysis and iterations are different. In particular, we study the following procedure, where each processor runs asynchronously and independently of the others, though they maintain a shared iteration counter  $k$ ; each processor performs the following:

- (i) Processor reads current problem data  $x$  and counter  $k$
- (ii) Processor draws a random sample  $W \sim P$ , computes  $g = \nabla F(x; W)$ , and increments a centralized counter  $k$
- (iii) Processor updates  $x \leftarrow x - \alpha_k g$  via sequential updates  $[x]_j = [x]_j - \alpha_k [g]_j$  for each coordinate  $j \in \{1, \dots, d\}$ .

In the iterations (i)–(iii), the scalars  $\alpha_k$  are a non-increasing stepsize sequence.

## 1.1 Main results and outline

The thrust of our results is that because of the noise inherent to the sampling process for  $W$ , the errors introduced by asynchrony in the iterations (i)–(iii) are asymptotically negligible: they do not matter. Even more, we can efficiently construct an  $x$  from the asynchronous process possessing optimal convergence properties and asymptotic variance. This has consequences for solving stochastic optimization problems on multi-core and multi-processor systems; we can leverage parallel computing without performing any synchronization, so that given a machine with  $m$  processors, we can read data and perform updates  $m$  times more quickly than what is possible with a single processor, and the error from reading stale information on  $x$  becomes asymptotically negligible. In Section 2, we make these optimality claims formal, presenting our main convergence theorems about the asynchronous iterations (i)–(iii) for solving the problem (1). Our main result, Theorem 1, gives explicit conditions under which an asynchronous stochastic gradient procedure converges at the optimal rate and with optimal asymptotic variance, and we give applications to specific stochastic optimization problems in Section 2.2. We also provide a more general result (Theorem 2) on the asynchronous solution of more general stochastic operator equations, again demonstrating that asynchrony introduces asymptotically less noise than that inherent in the stochastic problem itself. While we give explicit conditions under which our results hold, we note that roughly all we require is a type of local strong convexity around the optimal point  $x^* = \operatorname{argmin}_x f(x)$ , that the Hessian of  $f$  be positive definite near  $x^*$ , and a Lipschitz (smoothness) condition on the gradients  $\nabla f(x)$ .

In addition to theoretical results, in Section 3 we give empirical results on the power of parallelism and asynchrony in the implementation of stochastic approximation procedures. Our experiments demonstrate two results: first, even in non-asymptotic finite-sample settings, asynchrony introduces little degradation in solution quality, regardless of data sparsity (a common assumption in previous analyses); that is, asynchronously-constructed estimates are statistically efficient. Second, we show that there is some subtlety in implementation of these procedures in real hardware; while increases in parallelism lead to concomitant linear improvements in the speed with which we compute solutions to problem (1), in some cases we require strategies to reduce hardware resource competition between processors to achieve the full benefits of asynchrony.

## 1.2 Related work

Several researchers have provided and analyzed asynchronous algorithms for optimization. The seminal work of Bertsekas and Tsitsiklis [5] provides a comprehensive study both of models of asynchronous computation and analyses of asynchronous numerical algorithms, including coordinate- and gradient-descent methods. The results of theirs relevant to our work are roughly of two types. For non-stochastic problems, they show (roughly) linear convergence of iterative methods assuming the iterations satisfy certain contractive properties, which roughly correspond to variants of diagonal dominance of the Hessian of  $f$  (see, for example [5, Chapters 6.3 and 7.5]). For stochastic problems [5, Chapter 7.8], they show results that have a similar flavor to ours: errors due to asynchrony scale approximately quadratically in the stepsize  $\alpha$ , while gradient information scales linearly with  $\alpha$  so that it dominates other errors. Bertsekas and Tsitsiklis use this error scaling to show that stepsize choices of the form  $\alpha_k \approx 1/k$  guarantee asymptotic convergence under models of asynchrony with bounded delay. They leave open, however, a few interesting questions, namely,

attainable rates of convergence for asynchronous stochastic procedures, the effects of unbounded delays, and what optimality guarantees are possible relative to synchronous executions.

Due to their simplicity and dimension-independent convergence properties, stochastic and non-stochastic gradient methods have become extremely popular for large-scale data analysis and optimization problems (e.g. [23, 24, 8, 7, 6]). Consequently, with the advent of multi-core processing systems, there has been substantial work building on Bertsekas’ and Tsitsiklis’s results. Much of this work shows that asynchrony introduces negligible penalty in rates of convergence for optimization procedures under suitable conditions, such as gradient sparsity, conditioning of the Hessian of  $f$ , or allowable types of asynchrony (none, as we show, are essential). Niu et al. [25] propose the HOGWILD! method and show that under strong sparsity and smoothness assumptions on the data (essentially, that the gradients  $\nabla F(x; W)$  have a vanishing fraction of non-zero entries, that  $f$  is strongly convex, and  $\nabla F(x; \omega)$  is Lipschitz for all  $\omega$ ), convergence guarantees similar to the synchronous case are possible. Agarwal and Duchi [1] showed under restrictive ordering assumptions that some delayed gradient calculations have negligible asymptotic effect. Duchi et al. [10] extended Niu et al.’s results to a dual averaging algorithm that works for non-smooth, non strongly-convex problems, again so long as strong gradient sparsity assumptions hold (roughly, that the probability of an entry of  $\nabla F(x; \omega)$  being non-zero is inversely proportional to the maximum delay of any processor) and delays are bounded. Researchers have also investigated parallel coordinate descent solvers: Richtárik and Takáč [28] and Liu et al. [21] show how certain “separability” properties of an objective function  $f$ —meaning the degree to which different coordinates of  $x$  jointly affect  $f(x)$  (rather than  $f(x)$  depending on each coordinate  $x_j$  independently)—govern convergence rate of parallel coordinate descent methods, the latter focusing on asynchronous schemes. The conditions sufficient for fast or asynchronous convergence of coordinate methods are similar to the diagonal dominance conditions used by Bertsekas and Tsitsiklis [5, Chapter 6.3.2]. Yet, as we show, large-scale stochastic optimization renders many of these problem assumptions unnecessary. In particular, the asynchronous iterations (i)–(iii) retain all optimality properties of synchronous (correct) gradient procedures, even in the face of nearly unbounded delays, and enjoy optimal rates of convergence, even to constant pre-factors.

**Notation** We say a sequence of random variables or vectors  $X_n$  converges in distribution to a random variable  $Z$ , denoted  $X_n \xrightarrow{d} Z$ , if for all bounded continuous functions  $\mathbb{E}[f(X_n)] \rightarrow \mathbb{E}[f(Z)]$ . We say that a sequence of (finite-dimensional) random vectors  $X_n$  converges in  $L_p$  to a random vector  $Z$ , denoted  $X_n \xrightarrow{L_p} Z$ , if there is a norm  $\|\cdot\|$  such that  $\mathbb{E}[\|X_n - Z\|^p] \rightarrow 0$  as  $n \rightarrow \infty$ . This convergence is equivalent for any choice of the norm  $\|\cdot\|$ . We let  $X_n \xrightarrow{p} Z$  denote that  $X_n$  converges in probability to  $Z$ , meaning that  $\mathbb{P}(\|X_n - Z\| > \epsilon) \rightarrow 0$  as  $n \rightarrow \infty$  for any  $\epsilon > 0$ , and  $X_n \xrightarrow{a.s.} c$  denotes almost sure convergence, meaning that  $\mathbb{P}(\lim_n X_n \neq c) = 0$ . The notation  $\mathbf{N}(\mu, \Sigma)$  denotes the multivariate Gaussian with mean  $\mu$  and covariance  $\Sigma$ . We let  $I_{d \times d}$  denote the identity matrix in  $\mathbb{R}^{d \times d}$ , using  $I$  when the dimension is clear from context.

We use standard big- $O$  notation. For (nonnegative) sequences  $a_n$  and  $b_n$ , we let  $a_n \lesssim b_n$  mean there exists a constant  $C < \infty$  such that  $a_n \leq C b_n$  for all  $n$ , and  $a_n \asymp b_n$  means that there exist constants  $0 < c \leq C < \infty$  such that  $c \leq \liminf_n \frac{a_n}{b_n} \leq \limsup_n \frac{a_n}{b_n} \leq C$ . For random vectors  $X_n, Z_n$ , we say  $X_n = O_P(Z_n)$  if for all  $\epsilon > 0$ , there exists  $C < \infty$  such that  $\sup_n \mathbb{P}(\|X_n\| \geq C \|Z_n\|) \leq \epsilon$ , while  $X_n = o_P(Z_n)$  means that for all  $c > 0$ ,  $\limsup_n \mathbb{P}(\|X_n\| \geq c \|Z_n\|) = 0$ .

## 2 Main results

Our main results repose on a few standard assumptions often used for the analysis of stochastic optimization procedures, which we now detail, along with a few necessary definitions. We let  $k$  denote the iteration counter used throughout the asynchronous gradient procedure. Given that we compute  $g = \nabla F(x; W)$  with counter value  $k$  in the iterations (i)–(iii), we let  $x_k$  denote *the* (possibly inconsistent) particular  $x$  used to compute  $g$ , and likewise say that  $g = g_k$ , noting that the update to  $x$  is then performed using  $\alpha_k$ .

With the update (iii), we can give a more explicit formula for  $x_k$  as a function of time  $k$  with a small amount of additional notation. In particular, let  $E^{ki} \in \{0, 1\}^{d \times d}$  be a diagonal matrix whose  $j$ th diagonal entry is 1 if the  $i$ th gradient (i.e. that computed when the iteration counter is  $i$ ) has been incorporated into iterate  $x_k$  and is 0 otherwise. Then the iteration (i)–(iii) and index assignments imply that

$$x_k = - \sum_{i=1}^{k-1} \alpha_i E^{ki} g_i. \quad (2)$$

With this definition of the update matrices  $E^{ki}$ , we then associate a delay value  $M_k$  for each  $k$ , defined by

$$M_k := \min\{l - k : E^{lk} = I_{d \times d}, l \geq k\},$$

or the amount of time required for all updates from the  $k$ th gradient to be incorporated into the central  $x$  vector. Rather than assuming a uniform bound on the delay, throughout, we make the following assumption on the moments of the random variables  $M_k$ .

**Assumption A.** *There exists  $\tau > 2$  and a constant  $M < \infty$  such that*

$$\sup_k \mathbb{E} [M_k^\tau]^{1/\tau} \leq M.$$

Assumption A places our asynchronous iterations (i)–(iii) somewhere between Bertsekas’ and Tsitsiklis’s classification of *totally asynchronous* algorithms [5, Chapter 6], which require only that each processor performs its updates eventually, and *partially asynchronous* algorithms [5, Chapter 7], which specify a uniform bound on any processor’s delay; roughly, we have a quantitative version of total asynchrony. For example, if we know that processors have bounded delays, we may take  $\tau = \infty$  and assume that  $M = \sup_k M_k < \infty$ . In more general cases, however, we can allow infrequent longer delays with, as we shall see, negligible effect on our results except that the allowable stepsizes  $\alpha_i$  are more restricted.

### 2.1 Asynchronous convex optimization

We now present our main theoretical results for solving the stochastic convex problem (1), giving the necessary assumptions on  $f$  and  $F(\cdot; W)$  for our results. Our first assumption roughly states that  $f$  has a unique minimizer  $x^*$ , that  $f$  has a quadratic expansion near the point  $x^*$ , and is smooth (similar assumptions are common [e.g. 26, 5] and are satisfied in our applications).

**Assumption B.** *The function  $f$  has unique minimizer  $x^*$  and is twice continuously differentiable in the neighborhood of  $x^*$  with positive definite Hessian  $H = \nabla^2 f(x^*) \succ 0$ . There is a covariance matrix  $\Sigma \succ 0$  such that*

$$\mathbb{E}[\nabla F(x^*; W) \nabla F(x^*; W)^\top] = \Sigma.$$

Additionally, there exists a constant  $C < \infty$  such that the gradients  $\nabla F(x; W)$  satisfy

$$\mathbb{E}[\|\nabla F(x; W) - \nabla F(x^*; W)\|^2] \leq C \|x - x^*\|^2, \quad \text{all } x \in \mathbb{R}^d. \quad (3)$$

Lastly,  $f$  has  $L$ -Lipschitz continuous gradient, meaning  $\|\nabla f(x) - \nabla f(y)\| \leq L \|x - y\|$  for  $x, y \in \mathbb{R}^d$ .

Assumption B guarantees the uniqueness of the vector  $x^*$  minimizing  $f(x)$  over  $\mathbb{R}^d$  and ensures that  $f$  is well-behaved enough for our asynchronous iteration procedure to introduce negligible noise over a non-asynchronous procedure. In addition to Assumption B, we make one of two additional assumptions. In the first case, we assume that  $f$  is strongly convex:

**Assumption C.** *The function  $f$  is  $\lambda$ -strongly convex over all of  $\mathbb{R}^d$  for some  $\lambda > 0$ , that is,*

$$f(y) \geq f(x) + \langle \nabla f(x), y - x \rangle + \frac{\lambda}{2} \|x - y\|^2 \quad \text{for } x, y \in \mathbb{R}^d. \quad (4)$$

Our alternate assumption is a Lipschitz assumption on  $f$  itself, made by virtue of a second moment bound on  $\nabla F(x; W)$ .

**Assumption C'.** *There exists a constant  $G < \infty$  such that for all  $x \in \mathbb{R}^d$ ,*

$$\mathbb{E}[\|\nabla F(x; W)\|^2] \leq G^2. \quad (5)$$

In Section 2.2 to come, we give examples in which all of these assumptions are satisfied, showing that they are not too restrictive.

With our assumptions in place, we obtain our main theorem.

**Theorem 1.** *Let Assumptions A with moment  $\tau > 2$  and B hold. Let the iterates  $x_k$  be generated by the asynchronous process (i), (ii), (iii) with stepsize choice  $\alpha_k = \alpha k^{-\beta}$ , where  $\beta \in (\frac{1}{2} + \frac{1}{\tau-1}, 1)$  and  $\alpha > 0$ . Then if either of Assumptions C or C' holds, we have  $x_n \xrightarrow{\text{a.s.}} x^*$  and*

$$\frac{1}{\sqrt{n}} \sum_{k=1}^n (x_k - x^*) \overset{d}{\rightsquigarrow} \mathbf{N}(0, H^{-1} \Sigma H^{-1}) = \mathbf{N}(0, (\nabla^2 f(x^*))^{-1} \Sigma (\nabla^2 f(x^*))^{-1}).$$

Before moving to example applications of Theorem 1, we make a few additional remarks on the theorem, its consequences, and its associated conditions. Let  $\bar{x}_n := \frac{1}{n} \sum_{k=1}^n x_k$  for shorthand. First, using the delta method [e.g. 18, Theorem 1.8.12], we can give convergence rates for the function values  $f(\bar{x}_n)$  to  $f(x^*)$ . Specifically, they converge at the optimal rate of  $1/n$ , and we can give explicit constants.

**Corollary 1.** *Let the conditions of Theorem 1 hold. Then*

$$n(f(\bar{x}_n) - f(x^*)) \overset{d}{\rightsquigarrow} \frac{1}{2} \text{tr}[H^{-1} \Sigma] \cdot \chi_1^2,$$

where  $\chi_1^2$  denotes a chi-squared random variable with 1 degree of freedom, and  $H = \nabla^2 f(x^*)$  and  $\Sigma = \mathbb{E}[\nabla F(x^*; W) \nabla F(x^*; W)^\top]$ .

**Proof** Theorem 1 implies  $\bar{x}_n \xrightarrow{a.s.} x^*$ . By a Taylor expansion, we have

$$n(f(\bar{x}_n) - f(x^*)) = n \left[ \langle \nabla f(x^*), \bar{x}_n - x^* \rangle + \frac{1}{2} \langle \bar{x}_n - x^*, \nabla^2 f(x^*)(\bar{x}_n - x^*) \rangle + E_3(\bar{x}_n - x^*) \right],$$

where the error term  $E_3$  satisfies  $|E_3(x - x^*)| \leq r(x) \|x - x^*\|_2^2$  for a function  $r$  satisfying  $r(x) \rightarrow 0$  as  $x \rightarrow x^*$ . As  $\nabla f(x^*) = 0$ , we see that for a remainder  $|r_n| \leq r(\bar{x}_n) \cdot n \|\bar{x}_n - x^*\|_2^2 \xrightarrow{P} 0$ , we have

$$n(f(\bar{x}_n) - f(x^*)) = \frac{1}{2} \left\langle n^{-\frac{1}{2}}(\bar{x}_n - x^*), \nabla^2 f(x^*) n^{-\frac{1}{2}}(\bar{x}_n - x^*) \right\rangle + r_n.$$

By Theorem 1, the first term is asymptotically distributed as  $Z^\top H Z$  for  $Z \sim \mathbf{N}(0, H^{-1} \Sigma H^{-1})$ , and applying Slutsky's theorem [31, Theorem 2.7] and the continuous mapping theorem gives the result.  $\square$

Moreover, the convergence guarantee in Theorem 1 is generally unimprovable even by numerical constants. Recall that we have

$$\sqrt{n}(\bar{x}_n - x^*) \overset{d}{\rightsquigarrow} \mathbf{N}(0, H^{-1} \Sigma H^{-1}).$$

Standard results in asymptotic statistics imply that the rate of convergence and covariance  $H^{-1} \Sigma H^{-1}$  are optimal. Indeed, the Le Cam-Hájek local minimax theorem [17] implies that in standard statistical models, if we define the balls  $B(x, t) = \{x' \in \mathbb{R}^d : \|x - x'\| \leq t\}$ , then

$$\lim_{t \rightarrow \infty} \liminf_{n \rightarrow \infty} \sup_F \left\{ n \mathbb{E}[\|\hat{x}_n - x\|_2^2] : x = \operatorname{argmin}_{x'} \mathbb{E}[F(x'; W)] \in B(x^*, t/\sqrt{n}) \right\} \geq \operatorname{tr}[H^{-1} \Sigma H^{-1}],$$

where the supremum is taken over loss functions  $F$  satisfying Assumptions B and C or C', and  $\hat{x}_n$  is *any* sequence of estimators based on observing a sample  $W_1, \dots, W_n$ . The Le Cam-Hájek convolution theorem [17] and classical calculations with Bahadur efficiency (cf. van der Vaart [31, Chapter 8]) also show that the asymptotic covariance  $H^{-1} \Sigma H^{-1}$  is also generally optimal, meaning that no estimator can converge faster than  $\sqrt{n}$ , and the asymptotic covariance  $S$  of essentially any estimator converging at the rate  $\sqrt{n}$  must satisfy  $S \succeq H^{-1} \Sigma H^{-1}$ .

More concisely, in spite of the asynchrony we allow in the iterations (i)–(iii), we attain the best possible convergence rate. For the decision variables  $x$ , the rate  $n^{-\frac{1}{2}}$  is unimprovable, as is—at least generally—the asymptotic covariance  $H^{-1} \Sigma H^{-1}$ . We also have  $f(\bar{x}_n) - f(x^*) = O_P(n^{-1})$ , which is information-theoretically optimal [22, 2]. So we see that quite literally, the noise inherent to the sampling in stochastic gradient procedures swamps any noise introduced by asynchrony.

## 2.2 Examples

We now give two classical statistical optimization problems to illustrate Theorem 1. We verify that the conditions of the theorem hold for each of the examples, and these show that the conditions of Assumptions B and C or C' are not overly restrictive.

**Linear regression** Standard linear regression problems satisfies the conditions of Assumption C under an additional fourth moment condition. In this case, the data  $\omega = (a, b) \in \mathbb{R}^d \times \mathbb{R}$  and

the objective  $F(x; \omega) = \frac{1}{2}(\langle a, x \rangle - b)^2$ . If we have moment bounds  $\mathbb{E}[\|a\|_2^4] < \infty$ ,  $\mathbb{E}[b^2] < \infty$  and  $H = \mathbb{E}[aa^\top] \succ 0$ , we have  $\nabla^2 f(x^*) = H$ , and

$$\mathbb{E} \left[ \|\nabla F(x; W) - \nabla F(x^*; W)\|_2^2 \right] = \mathbb{E} \left[ \|a\|_2^2 (a^\top (x - x^*))^2 \right] \leq \mathbb{E} \left[ \|a\|_2^4 \right] \|x - x^*\|_2^2,$$

whence the assumptions of Theorem 1 are satisfied. We can give more explicit calculations if we make standard modeling assumptions, for example, that  $b = \langle a, x^* \rangle + \varepsilon$ , where  $\varepsilon$  is an independent mean-zero noise sequence with  $\mathbb{E}[\varepsilon^2] = \sigma^2$ . In this case, the minimizer of  $f(x) = \mathbb{E}[F(x; W)]$  is  $x^*$ , we have  $\langle a, x^* \rangle - b = -\varepsilon$ , and

$$\mathbb{E}[\nabla F(x^*; W) \nabla F(x^*; W)^\top] = \mathbb{E}[(\langle a, x^* \rangle - b)aa^\top(\langle a, x^* \rangle - b)] = \mathbb{E}[aa^\top \varepsilon^2] = \sigma^2 \mathbb{E}[aa^\top] = \sigma^2 H.$$

In particular, the asynchronous iterates satisfy

$$n^{-\frac{1}{2}} \sum_{k=1}^n (x_k - x^*) \overset{d}{\rightsquigarrow} \mathbf{N}(0, \sigma^2 H^{-1}) = \mathbf{N}\left(0, \sigma^2 \mathbb{E}[aa^\top]^{-1}\right).$$

This has the asymptotic variance of the ordinary least squares estimate of  $x^*$ , which is minimax optimal [18, Chapter 5].

**Logistic regression** As long as the data has finite second moment, logistic regression problems satisfy all the conditions of Assumption C' in Theorem 1. In this case we have  $\omega = (a, b) \in \mathbb{R}^d \times \{-1, 1\}$  and instantaneous objective  $F(x; \omega) = \log(1 + \exp(-b \langle a, x \rangle))$ . For fixed  $\omega$ , this function is Lipschitz continuous and has gradient and Hessian

$$\nabla F(x; \omega) = -\frac{1}{1 + \exp(b \langle a, x \rangle)} ba \quad \text{and} \quad \nabla^2 F(x; \omega) = \frac{e^{b \langle a, x \rangle}}{(1 + e^{b \langle a, x \rangle})^2} aa^\top,$$

where  $\nabla F(x; \omega)$  is Lipschitz continuous as  $\|\nabla^2 F(x; (a, b))\| \leq \frac{1}{4} \|a\|_2^2$ . Thus, so long as  $\mathbb{E}[\|a\|_2^2] < \infty$  and  $\mathbb{E}[\nabla^2 F(x^*; W)] \succ 0$  (the latter occurs if  $\mathbb{E}[aa^\top]$  is full rank), logistic regression satisfies the conditions of Theorem 1. In particular, the asynchronous stochastic gradient method achieves optimal convergence guarantees.

### 2.3 Extension to nonlinear problems and variational inequalities

We prove Theorem 1 by way of a more general result on finding the zeros of a residual operator  $R : \mathbb{R}^d \rightarrow \mathbb{R}^d$ , where we only observe noisy views of  $R(x)$ , and there is unique  $x^*$  such that  $R(x^*) = 0$ . Such situations arise, for example, in the solution of stochastic monotone operator problems (cf. Juditsky, Nemirovski, and Tauvel [16] or Bertsekas and Tsitsiklis [5]), including finding equilibria in stochastic convex Nash games (e.g. [16, Sec. 2.1], [5, Ex. 3.5.1(d)]), general saddle-point problems, or multi-user routing problems [5, Ex. 3.5.1(c)]. In this more general setting, we consider the following stochastic and asynchronous iterative process, which extends that for the convex case outlined previously. Each processor performs the following asynchronously and independently:

- (i) Processor reads current problem data  $x$  and counter  $k$
- (ii) Processor receives vector  $g = R(x) + \xi$ , where  $\xi$  is a random (conditionally) mean-zero noise vector, and increments a centralized counter  $k$

- (iii) Processor updates  $x \leftarrow x - \alpha_k g$  via sequential updates  $[x]_j = [x]_j - \alpha_k [g]_j$  for each coordinate  $j \in \{1, \dots, d\}$

As in the convex case, we associate vectors  $x_k$  and  $g_k$  with the update performed using  $\alpha_k$ , and we let  $\xi_k$  denote the noise vector used to construct  $g_k$ . As before, these iterates and assignment of indices again imply that  $x_k$  has the form (2), that is,  $x_k = -\sum_{i=1}^{k-1} \alpha_i E^{ki} g_i$  for diagonal matrices  $E^{ki}$  capturing the updates that have been performed at time  $k$ .

For this iterative process, we define the increasing sequence of  $\sigma$ -fields  $\mathcal{F}_k$  by

$$\mathcal{F}_k = \sigma(\xi_1, \dots, \xi_k, \{E^{ij} : i \leq k+1, j \leq i\}), \quad (6)$$

that is, the noise variables  $\xi_k$  are adapted to the filtration  $\mathcal{F}_k$ , and these  $\sigma$ -fields are the smallest containing both the noise and all index updates that have occurred and that will occur to compute  $x_{k+1}$ . Thus we have  $x_{k+1} \in \mathcal{F}_k$ , and our mean-zero assumption on the noise  $\xi$  is that

$$\mathbb{E}[\xi_k \mid \mathcal{F}_{k-1}] = 0.$$

Our analysis builds off of Polyak and Juditsky's study [26] of averaging in stochastic approximation, and we model our requirements for the convergence of the preceding iteration on those they use for the solution of the nonlinear equality  $R(x^*)$ . First, we assume there is a Lyapunov function  $V$  that functions (essentially) as a squared norm, which satisfies  $V(x) \geq \lambda \|x\|^2$  for all  $x \in \mathbb{R}^d$ ,  $\|\nabla V(x) - \nabla V(y)\| \leq L \|x - y\|$  for all  $x, y$ , that  $\nabla V(0) = 0$ , and  $V(0) = 0$ . Note in particular that this implies

$$\lambda \|x\|^2 \leq V(x) \leq V(0) + \langle \nabla V(0), x - 0 \rangle + \frac{L}{2} \|x\|^2 = \frac{L}{2} \|x\|^2 \quad (7)$$

and that  $\|\nabla V(x)\|^2 \leq L^2 \|x\|^2 \leq (L^2/\lambda)V(x)$ . In addition, we make the following assumptions on the residual function (cf. [26, Assumption 3.2]).

**Assumption D.** *There exists a matrix  $H \in \mathbb{R}^{d \times d}$  with  $H \succ 0$ , a parameter  $0 < \gamma \leq 1$ , constant  $C < \infty$ , and some  $\epsilon > 0$  such that if  $x$  satisfies  $\|x - x^*\| \leq \epsilon$ , then*

$$\|R(x) - H(x - x^*)\| \leq C \|x - x^*\|^{1+\gamma}.$$

Assumption D essentially requires that  $R$  is differentiable at  $x^*$  with derivative matrix  $H \succ 0$ . We also make a few assumptions on the noise process  $\xi$  paralleling Assumption 3.3 of Polyak and Juditsky [26]; specifically, we assume  $\xi$  implicitly depends on  $x \in \mathbb{R}^d$  (so that we may write  $\xi_k = \xi(x_k)$ ), and that the following assumption holds.

**Assumption E.** *The noise vector  $\xi(x)$  decomposes as  $\xi(x) = \xi(0) + \zeta(x)$ , where  $\xi(0)$  is a process satisfying  $\mathbb{E}[\xi_k(0)\xi_k(0)^\top \mid \mathcal{F}_{k-1}] \xrightarrow{P} \Sigma \succ 0$  for some matrix  $\Sigma \in \mathbb{R}^{d \times d}$ , the boundedness condition  $\mathbb{E}[\sup_k \mathbb{E}[\|\xi_k(0)\|^2 \mid \mathcal{F}_{k-1}]] < \infty$ , and*

$$\mathbb{E}[\|\zeta_k(x)\|^2 \mid \mathcal{F}_{k-1}] \leq C \|x - x^*\|^2$$

for some constant  $C < \infty$  and all  $x \in \mathbb{R}^d$ .

As in the convex case, we make one of two additional assumptions, which should be compared with Assumptions C and C'. The first is that  $R$  gives globally strong information about  $x^*$ .

**Assumption F** (Strongly convex residuals). *There exists a constant  $\lambda_0 > 0$  such that for all  $x \in \mathbb{R}^d$ ,  $\langle \nabla V(x - x^*), R(x) \rangle \geq \lambda_0 V(x - x^*)$ .*

Alternatively, we may make an assumption on the boundedness of  $R$ , which we shall see suffices for proving our main results.

**Assumption F'** (Bounded residuals). *There exist  $\lambda_0 > 0$  and  $\epsilon > 0$  such that*

$$\inf_{0 < \|x - x^*\| \leq \epsilon} \frac{\langle \nabla V(x - x^*), R(x) \rangle}{V(x - x^*)} \geq \lambda_0 \quad \text{and} \quad \inf_{\epsilon < \|x - x^*\|} \langle \nabla V(x - x^*), R(x) \rangle > 0.$$

*There also exists some  $C < \infty$  such that  $\|R(x)\| \leq C$  and  $\mathbb{E}[\|\xi_k\|^2 \mid \mathcal{F}_{k-1}] \leq C^2$  for all  $k$  and  $x$ .*

With these assumptions in place, we obtain the following more general version of Theorem 1; indeed, we show in the sequel how Theorem 1 follows from this result.

**Theorem 2.** *Let  $V$  be a function satisfying inequality (7), and let Assumptions D, E, and A hold. Let the stepsizes  $\alpha_k = \alpha k^{-\beta}$ , where  $\frac{1}{\tau-1} + \frac{1}{1+\gamma} < \beta < 1$ . Let one of Assumptions F or F' hold. Then  $x_n \xrightarrow{\text{a.s.}} x^*$  and*

$$\frac{1}{\sqrt{n}} \sum_{k=1}^n (x_k - x^*) \overset{d}{\rightsquigarrow} \mathbf{N}(0, H^{-1} \Sigma H^{-1}).$$

We may compare this result to Polyak and Juditsky's Theorem 2, which gives identical covariance matrix and asymptotic convergence guarantees, but with weaker conditions on the function  $V$  and stepsize sequence  $\alpha_k$ . Our mildly stronger assumptions—namely, Assumptions F and F' are stronger versions of Assumption 3.1 of Polyak and Juditsky [26], which requires only the conditions on  $V$  and  $R$  of Assumption F'—allow our result to apply even in the asynchronous settings considered in this paper.

### 3 Experimental results

We provide empirical results studying the performance of asynchronous stochastic approximation schemes on several simulated and real-world datasets. Our theoretical results suggest that asynchrony should introduce little degradation in solution quality; we also investigate the engineering techniques necessary to truly leverage the power of asynchronous stochastic procedures. In our experiments, we focus on linear and logistic regression, the examples given in Section 2.2; that is, we have data  $(a_i, b_i) \in \mathbb{R}^d \times \mathbb{R}$  (for linear regression) or  $(a_i, b_i) \in \mathbb{R}^d \times \{-1, 1\}$  (for logistic regression), for  $i = 1, \dots, N$ , and objectives

$$f(x) = \frac{1}{2N} \sum_{i=1}^N (\langle a_i, x \rangle - b_i)^2 \quad \text{and} \quad f(x) = \frac{1}{N} \sum_{i=1}^N \log(1 + \exp(-b_i \langle a_i, x \rangle)). \quad (8)$$

We perform each of our experiments using a 48-core Intel Xeon machine with 1 terabyte of RAM, and have put code and binaries to replicate our experiments on CodaLab [11]. The Xeon architecture puts each core onto one of four sockets, where each socket has its own memory. To limit the impact of communication overhead in our experiments, we limit all experiments to at most 12 cores, all on the same socket. Within an experiment—based on the empirical expectations (8)—we

iterate in *epochs*, so that our stochastic gradient procedure loops through random permutations of all examples, touching each example exactly once per epoch in a different random order within each epoch (cf. [27]).<sup>1</sup> We use the following two schemes for the stepsize  $\alpha_k$ .

**Decreasing stepsizes.** We set  $\beta = 0.55$  and let  $\alpha_k = k^{-\beta}$ . The value written on the shared iteration counter  $k$  by one processor may be overwritten by other processors.

**Exponential backoff stepsizes.** We use a fixed stepsize  $\alpha$ , decreasing the stepsize by a factor of 0.95 between each epoch (this matches the experimental protocol of Niu et al. [25] and follows Hazan and Kale [14] and Ghadimi and Lan [13]).

To address issues of hardware resource contention (see Section 3.2 for more on this), in some cases we use a *mini-batching* strategy. Abstractly, in the formulation of the basic problem (1), this means that in each calculation of a stochastic gradient  $g$  we draw  $B \geq 1$  samples  $W_1, \dots, W_B$  i.i.d. according to  $P$ , then set

$$g(x) = \frac{1}{B} \sum_{b=1}^B \nabla F(x; W_b). \quad (9)$$

The mini-batching strategy (9) does not change the (asymptotic) convergence guarantees of asynchronous stochastic gradient descent, as the covariance matrix  $\Sigma = \mathbb{E}[g(x^*)g(x^*)^\top]$  satisfies  $\Sigma = \frac{1}{B} \mathbb{E}[\nabla F(x^*; W) \nabla F(x^*; W)^\top]$ , while the total iteration count is reduced by a factor  $B$ . Lastly, we measure the performance of optimization schemes via *speedup*, defined as

$$\text{speedup} = \frac{\text{average epoch runtime on a single core using stochastic gradient descent}}{\text{average epoch runtime of asynchronous method on } m \text{ cores}}. \quad (10)$$


In our experiments, we see that increasing the number  $m$  of cores does not change the gap in optimality  $f(x_k) - f(x^*)$  after each epoch, so speedup is equivalent to the ratio of the time required to obtain an  $\epsilon$ -accurate solution using a single processor/core to that required to obtain  $\epsilon$ -accurate solution using  $m$  processors/cores.

### 3.1 Efficiency and sparsity


For our first set of experiments, we study the effect that data sparsity has on the convergence behavior of asynchronous methods using the linear regression objective (8). Sparsity has been an essential part of the analysis of many asynchronous and parallel optimization schemes [25, 10, 28], while our theoretical results suggest it should be unimportant, so understanding these effects is important. We generate synthetic linear regression problems with  $N = 10^6$  examples in  $d = 10^3$  dimensions via the following procedure. Let  $p_{\text{nz}} \in (0, 1]$  be the desired fraction of non-zero gradient entries, and let  $\Pi_{p_{\text{nz}}}$  be a random projection operator that zeros out all but a fraction  $p_{\text{nz}}$  of the elements of its argument, meaning that for  $a \in \mathbb{R}^d$ ,  $\Pi_{p_{\text{nz}}}(a)$  uniformly at random chooses  $p_{\text{nz}}d$  elements of  $a$  and leaves them identical, zeroing the remaining elements. We generate data for our linear regression by drawing a random vector  $u^* \sim \mathbf{N}(0, I_{d \times d})$ , then constructing

$$b_i = \langle a_i, u^* \rangle + \varepsilon_i, \quad \text{where } \varepsilon_i \stackrel{\text{i.i.d.}}{\sim} \mathbf{N}(0, 1), \quad \tilde{a}_i \stackrel{\text{i.i.d.}}{\sim} \mathbf{N}(0, I_{d \times d}), \quad \text{and } a_i = \Pi_{p_{\text{nz}}}(\tilde{a}_i) \quad (11)$$

for  $i = 1, \dots, N$ , where  $\Pi_{p_{\text{nz}}}(\tilde{a}_i)$  denotes an independent random sparse projection of  $\tilde{a}_i$ . To measure optimality gap, we directly compute  $x^* = (A^T A)^{-1} A^T b$ , where  $A = [a_1 \ a_2 \ \dots \ a_N]^\top \in \mathbb{R}^{N \times d}$ .



**Figure 1.** Decreasing stepsizes: Optimality gaps for synthetic linear regression experiments showing effects of data sparsity and asynchrony on  $f(x_k) - f(x^*)$  with stepsize  $\alpha_k = k^{-\beta}$ . A fraction  $p_{\text{nz}}$  of each vector  $a_i \in \mathbb{R}^d$  is non-zero.




**Figure 2.** Exponentially decreasing stepsizes: Optimality gaps for synthetic linear regression experiments showing effects of data sparsity and asynchrony on  $f(x_k) - f(x^*)$  with epoch-based stepsizes  $\alpha_{\text{epoch } k} = .95^k$ . A fraction  $p_{\text{nz}}$  of each vector  $a_i \in \mathbb{R}^d$  is non-zero.


In Figures 1 and 2 we plot the results of simulations using densities  $p_{\text{nz}} \in \{.005, .01, .2, 1\}$  and mini-batch size  $B = 10$ , showing the gap  $f(x_k) - f(x^*)$  as a function of the number of epochs for each of the given sparsity levels. Figure 1 gives results for our simulated data (11) using the decreasing stepsize scheme  $\alpha_k = k^{-\beta}$  with  $\beta = .55$ , while Figure 2 gives results using the exponential backoff scheme of [14, 13, 25], where stepsizes are chosen per epoch as  $\alpha_{\text{epoch } k} = .95^k$ . Each plot includes error bars with standard errors over 10 random experiments using different random seeds (the errors are generally too small to see in the plots). We give results using 1, 4, 8, and 10 processor cores (increasing degrees of asynchrony). From the plots, we see that regardless of the number of cores, the convergence behavior is nearly identical, with minor degradations in performance for the sparsest data. (We plot the gaps  $f(x_k) - f(x^*)$  on a logarithmic axis.) Moreover, as the data becomes denser, the more asynchronous methods—larger number of cores—achieve performance essentially identical to the fully synchronous method in terms of convergence versus number of epochs.

In Figures 3 and 4, we plot the speedup achieved for the synthetic regression problem with

<sup>1</sup>Strictly speaking, this violates the stochastic gradient assumption, but it allows direct comparison with the original HOGWILD! code and implementation [25].



**Figure 3.** Decreasing stepsizes: Speedups for synthetic linear regression experiments showing effects of data sparsity on speedup (10) with stepsize  $\alpha_k = k^{-\beta}$ . A fraction  $p_{nz}$  of each vector  $a_i \in \mathbb{R}^d$  is non-zero.



**Figure 4.** Exponentially decreasing stepsizes: Speedups for synthetic linear regression experiments showing effects of data sparsity on speedup (10) with epoch-based stepsizes  $\alpha_{\text{epoch } k} = .95^k$ . A fraction  $p_{nz}$  of each vector  $a_i \in \mathbb{R}^d$  is non-zero.

data (11) using different numbers of cores for the experiments in Figures 1 and 2 (as before, Fig. 3 uses stepsizes  $\alpha_k = k^{-\beta}$  and Fig. 4 uses stepsizes exponentially decreasing between epochs). As a point of comparison, we also implement a synchronized method, which uses multiple cores to compute gradients independently  $g^i$  on each core  $i = 1, \dots, c$ , then computes the average  $\bar{g} = \frac{1}{c} \sum_{i=1}^c g^i$  and uses that to perform a standard stochastic gradient update; this requires explicit synchronization (locking) of the updates, though with sufficiently large batch sizes  $\mathbf{B}$  computation may theoretically overwhelm the communication and locking overhead [7]. For comparison, we use the same batch size  $\mathbf{B} = 10$  for each of the synchronous and asynchronous procedures, and we see that the performance of the naive locking strategy is worse than than the asynchronous gradient method across all data densities  $p_{nz}$ . At higher densities, more computation is necessary within each gradient computation, so that the communication overhead causes less performance degradation for the synchronous method, yet the asynchronous method also benefits and attains better relative performance. We see clearly that data sparsity is not necessary for the asynchronous gradient method to enjoy substantial performance benefits.

No batching ( $B = 1$ )				
Number of cores	1	4	8	10
fraction of L1 misses	$0.0021 \pm 0.0001$	$0.0061 \pm 0.0001$	$0.0096 \pm 0.0001$	$0.0102 \pm 0.0001$
fraction of L2 misses	$0.50 \pm 0.01$	$0.63 \pm 0.01$	$0.76 \pm 0.01$	$0.78 \pm 0.01$
fraction of L3 misses	$0.41 \pm 0.01$	$0.25 \pm 0.01$	$0.24 \pm 0.01$	$0.25 \pm 0.01$
epoch average time (s)	4.55	1.85	1.61	1.47
<b>speedup</b>	<b>1.00</b>	<b><math>2.46 \pm 0.01</math></b>	<b><math>2.83 \pm 0.01</math></b>	<b><math>3.09 \pm 0.01</math></b>

Batch size $B = 10$				
Number of cores	1	4	8	10
fraction of L1 misses	$0.0027 \pm 0.0002$	$0.0033 \pm 0.0001$	$0.0043 \pm 0.0001$	$0.0046 \pm 0.0001$
fraction of L2 misses	$0.44 \pm 0.01$	$0.50 \pm 0.01$	$0.60 \pm 0.01$	$0.63 \pm 0.01$
fraction of L3 misses	$0.35 \pm 0.03$	$0.33 \pm 0.01$	$0.33 \pm 0.01$	$0.33 \pm 0.01$
epoch average time (s)	2.97	0.87	0.58	0.51
<b>speedup</b>	<b>1.00</b>	<b><math>3.42 \pm 0.01</math></b>	<b><math>5.16 \pm 0.02</math></b>	<b><math>5.80 \pm 0.03</math></b>

**Table 1.** Memory traffic for batched updates (9) versus non-batched updates ( $B = 1$ ) for a dense linear regression problem in  $d = 10^3$  dimensions with a sample of size  $N = 10^6$ . Cache misses are substantially higher with  $B = 1$ .


### 3.2 Hardware issues and cache locality

We now detail a set of experiments investigating hardware issues that arise even in the implementation of asynchronous gradient methods. The Intel x86 architecture (as with essentially every processor architecture) organizes memory in a hierarchy, going from level 1 to level 3 (L1 to L3) caches of increasing sizes. An important aspect of the speed of different optimization schemes is the relative fraction of memory *hits*, meaning accesses to memory that is cached locally (in order of decreasing speed, L1, L2, or L3 cache). In Table 1, we show the proportion of cache misses at each level of the memory hierarchy for our synthetic regression experiment with fully dense data ( $p_{\text{nz}} = 1$ ) over the execution of 20 epochs, averaged over 10 different experiments. We compare memory contention when the batch size  $B$  used to compute the local asynchronous gradients (9) is 1 and 10. We see that the proportion of misses for the fastest two levels—1 and 2—of the cache for  $B = 1$  increase significantly with the number of cores, while increasing the batch size to  $B = 10$  substantially mitigates cache incoherence. In particular, we maintain (near) linear increases in iteration speed with little degradation in solution quality (the gap  $f(\hat{x}) - f(x^*)$  output by each of the procedures with and without batching is identical to within  $10^{-3}$ ; cf. Figure 2(d)).

### 3.3 Real datasets

We perform experiments using three different real-world datasets: the Reuters RCV1 corpus [19], the Higgs detection dataset [3], and the Forest Cover dataset [20]. Each represents a binary classification problem, which we formulate using logistic regression (recall Sec. 2.2). We briefly detail relevant statistics of each:

- (1) The Reuters RCV1 dataset consists of  $N \approx 7.81 \cdot 10^5$  data vectors (documents)  $a_i \in \{0, 1\}^d$  with  $d \approx 5 \cdot 10^4$  dimensions; each vector has sparsity approximately  $p_{\text{nz}} = 3 \cdot 10^{-3}$ . Our task is to classify each document as being about corporate industrial topics (CCAT) or not.




**Figure 5.** Decreasing stepsizes: Optimality gaps  $f(x_k) - f(x^*)$  on the (a) RCV1, (b) Higgs, and (c) Forest Cover datasets using stepsize  $\alpha_k = k^{-\beta}$  with  $\beta = .55$ .

- (2) The Higgs detection dataset consists of  $N = 10^6$  data vectors  $\tilde{a}_i \in \mathbb{R}^{d_0}$ , with  $d_0 = 28$ . We quantize each coordinate into 5 bins containing equal fraction of the coordinate values and encode each vector  $\tilde{a}_i$  as a vector  $a_i \in \{0, 1\}^d$  with  $d = 5d_0$  whose non-zero entries correspond to quantiles into which coordinates fall. The task is to detect (simulated) emissions from a linear accelerator.
- (3) The Forest Cover dataset consists of  $N \approx 5.7 \cdot 10^5$  data vectors  $a_i \in \{-1, 1\}^d$  with  $d = 54$ , and the task is to predict forest growth types.

Thus, each dataset gives a different flavor of optimization problem: the first is very sparse and high-dimensional, the second is somewhat sparse and of moderate dimension, while the forest dataset is dense but of relatively small dimension. These allow a broader picture of the performance of the asynchronous gradient method.

We follow the same experimental protocol as in our simulated data experiments. That is, we perform 10 experiments for each dataset using 1, 4, 8, and 10 cores, where each experiment consists of running the asynchronous gradient method for 20 epochs, within each of which examples are accessed according to a new random permutation. We use a batch size  $\mathbf{B} = 10$  for each experiment, and collect standard errors for the (estimated) optimality gaps. In these experiments, as a proxy for the optimal value  $f(x^*)$  we run a synchronous gradient method for 100 epochs, using its best objective value as  $f(x^*)$ . In Figures 5 and 6, we plot the gap  $f(x_k) - f(x^*)$  as a function of epochs, giving standard error intervals, for each of the three datasets. The figures show there is essentially no degradation in objective value when using different numbers of processors, that is, asynchrony appears to have negligible effect for each problem, whether we use stepsizes  $\alpha_k = k^{-\beta}$  (with  $\beta = .55$ ) that we analyze or the epoch-based exponentially decreasing stepsize scheme used by Niu et al. [25], also analyzed by Hazan and Kale [14] and Ghadimi and Lan [13]. In Figures 7 and 8, we plot speedup achieved for these same experiments. The asynchronous gradient method iteration achieves nearly linear speedup of between  $6\times$  and  $8\times$  on each of the datasets using 10 cores.



**Figure 6.** Exponentially decreasing stepsizes: Optimality gaps  $f(x_k) - f(x^*)$  on the (a) RCV1, (b) Higgs, and (c) Forest Cover datasets with epoch-based stepsizes  $\alpha_{\text{epoch } k} = .95^k$ .

## 4 Proofs

In this section, we present proofs of our two main theorems, deferring the proofs of technical lemmas to subsequent appendices.

### 4.1 Proof of Theorem 1

We prove Theorem 1 by a reduction to Theorem 2. For both settings of Theorem 1, we will use  $V(x) = \frac{1}{2} \|x\|^2$  and  $R(x) = \nabla f(x)$ , then apply Theorem 2. With these choices, Assumption D is satisfied with the Hessian  $H = \nabla^2 f(x^*) \succ 0$  and  $\gamma = 1$  by a Taylor expansion of  $f$ , which is assumed continuously twice differentiable near  $x^*$ .

Let us also verify that Assumption E holds for an appropriate noise sequence  $\xi_k$  in the stochastic convex optimization setting. Throughout, we also use the sequence of  $\sigma$ -fields  $\mathcal{F}_k$  defined by expression (6), so that  $x_k \in \mathcal{F}_{k-1}$ , and the counter  $k$  implicitly gives a gradient  $g_k$ , point  $x_k$ , sample  $W_k$ , and noise error

$$\xi_k = g_k - R(x_k) = \nabla F(x_k; W_k) - \nabla f(x_k). \quad (12)$$

That is, we have  $g_k = R(x_k) + \xi_k$  as in the nonlinear setting of Theorem 2. We first verify that  $\xi_k$  is a martingale difference sequence: we have  $\mathbb{E}[\xi_k | \mathcal{F}_{k-1}] = \nabla f(x_k) - \nabla f(x_k) = 0$  because  $W_k$  is independent of  $x_k$ . We also have the decomposition


$$\xi_k = \underbrace{\nabla F(x^*; W_k)}_{\xi_k(0)} + \underbrace{\nabla F(x_k; W_k) - \nabla F(x^*; W_k) - \nabla f(x_k)}_{\zeta_k(x_k)},$$

and that  $\|\nabla f(x_k)\| = \|\nabla f(x_k) - \nabla f(x^*)\| \leq L \|x_k - x^*\|$ , so

$$\mathbb{E}[\|\zeta_k(x_k)\|^2 | \mathcal{F}_{k-1}] = \mathbb{E}[\|\nabla F(x_k; W_k) - \nabla F(x^*; W_k)\|^2 | \mathcal{F}_{k-1}] + \|\nabla f(x_k)\|^2 \leq C \|x_k - x^*\|^2$$

by inequality (3). Moreover, we have

$$\mathbb{E}[\xi_k(0)\xi_k(0)^\top | \mathcal{F}_{k-1}] = \mathbb{E}[\nabla F(x^*; W_k)\nabla F(x^*; W_k)^\top | \mathcal{F}_{k-1}] = \Sigma,$$



**Figure 7.** (Decreasing stepsizes) Logistic regression experiments showing speedup (10) on the (a) RCV1, (b) Higgs, and (c) Forest Cover datasets.

as the random variable  $W_k$  is independent of  $\mathcal{F}_{k-1}$ . That is, Assumption E holds.

Now we show that Assumption F holds whenever Assumption C holds. If  $f$  is strongly convex, then taking  $R(x) = \nabla f(x)$  and  $V(x) = \frac{1}{2} \|x\|^2$  we have for any  $x, y \in \mathbb{R}^d$  that

$$f(y) \geq f(x) + \langle \nabla f(x), y - x \rangle + \frac{\lambda}{2} \|x - y\|^2 \quad \text{and} \quad f(x) \geq f(y) + \langle \nabla f(y), x - y \rangle + \frac{\lambda}{2} \|x - y\|^2.$$

Taking  $y = x^*$  in the preceding expression while noting that  $\nabla f(x^*) = \nabla f(y) = 0$ , we have

$$\langle \nabla V(x - x^*), R(x) \rangle = \langle x - x^*, \nabla f(x) \rangle = \langle \nabla f(x) - \nabla f(x^*), x - x^* \rangle \geq \lambda \|x - x^*\|^2 = 2\lambda V(x - x^*).$$

Clearly,  $V$  has 1-Lipschitz gradient and satisfies  $V(x - x^*) \geq \frac{1}{2} \|x - x^*\|^2$ .

Now we consider conditions on the convex function  $f$  under which Assumption F' is satisfied. In particular, let  $f$  be a differentiable Lipschitz-continuous convex function defined on  $\mathbb{R}^d$ , and assume that  $f$  is locally strongly convex near  $x^*$ , meaning that there exist  $\lambda > 0$  and  $\epsilon > 0$  such that

$$f(y) \geq f(x) + \langle \nabla f(x), y - x \rangle + \frac{\lambda}{2} \|x - y\|^2 \quad \text{for } x, y \text{ s.t. } \|x - x^*\| \leq \epsilon, \|y - x^*\| \leq \epsilon.$$


In particular, as  $\nabla f(x^*) = 0$ , we have  $f(x) \geq f(x^*) + \frac{\lambda}{2} \|x - x^*\|^2$  for all  $x$  such that  $\|x - x^*\| \leq \epsilon$ . We have the following lemma on the growth of such functions.

**Lemma 1.** *Let  $f$  satisfy the conditions in the preceding paragraph. Then*

$$f(x) \geq f(x^*) + \frac{\lambda}{2} \min\{\|x - x^*\|^2, \epsilon \|x - x^*\|\}. \quad (13)$$

Deferring proof of Lemma 1, we show how it implies that the conditions of Assumption F' are satisfied with the Lyapunov function  $V(x) = \frac{1}{2} \|x\|^2$  and residual operator  $R(x) = \nabla f(x)$ . Indeed, by applying the strong convexity inequality with  $y = x^*$ , we have for  $x$  such that  $\|x - x^*\| \leq \epsilon$  that

$$\langle \nabla V(x - x^*), R(x) \rangle = \langle x - x^*, \nabla f(x) \rangle = \langle \nabla f(x) - \nabla f(x^*), x - x^* \rangle \geq \lambda \|x - x^*\|^2.$$



**Figure 8.** Exponentially decreasing stepsizes: Logistic regression experiments showing speedup (10) on the (a) RCV1, (b) Higgs, and (c) Forest Cover datasets with epoch-based stepsize  $\alpha_{\text{epoch } k} = .95^k$ .

Now we claim that

$$\inf_{x: \|x - x^*\| > \epsilon} \langle \nabla V(x - x^*), R(x) \rangle = \inf_{x: \|x - x^*\| > \epsilon} \langle x - x^*, \nabla f(x) \rangle > 0. \quad (14)$$

To see this, note that by claim (13), for  $x$  such that  $\|x - x^*\| \geq \epsilon$ , we have for some constant  $c > 0$  that  $f(x) - f(x^*) \geq c\|x - x^*\|$ , while we have  $f(x^*) \geq f(x) + \langle \nabla f(x), x^* - x \rangle$ , so that

$$\langle \nabla f(x), x - x^* \rangle \geq f(x) - f(x^*) \geq c\|x - x^*\| > c\epsilon \text{ for all } x \text{ s.t. } \|x - x^*\| > \epsilon.$$

Additionally, whenever Assumption C' holds, we have  $\mathbb{E}[\|\nabla F(x; W)\|^2] \leq G^2$ , so that all the conditions of Assumption F' are satisfied. Except for the proof of Lemma 1, this completes the proof of Theorem 1.

**Proof of Lemma 1** Fix  $y \in \mathbb{R}^d$  and let  $h_y(t) = f(x^* + ty/\|y\|) - f(x^*)$ . Notably,  $h_y$  is a one-dimensional convex function, and  $h_y(t) \geq (\lambda/2)t^2$  for  $|t| \leq \epsilon$ . As the slopes of convex functions are non-decreasing (cf. Hiriart-Urruty and Lemaréchal [15, Chapter I]), we have

$$h'_y(\epsilon) = \lim_{\delta \rightarrow 0} \frac{h_y(\epsilon + \delta) - h_y(\epsilon)}{\delta} \geq \frac{h_y(\epsilon) - h_y(0)}{\epsilon} \geq \frac{\lambda\epsilon^2}{2\epsilon} = \frac{\lambda\epsilon}{2}.$$

This inequality implies that for any  $t \geq \epsilon$ , we have

$$\frac{f(x^* + t\frac{y}{\|y\|}) - f(x^*)}{t} = \frac{h_y(t) - h_y(0)}{t} \geq \frac{h_y(\epsilon) - h_y(0)}{\epsilon} \geq \frac{\lambda\epsilon}{2},$$

while for  $0 < t < \epsilon$ , we use that  $t \mapsto (h_y(t) - h_y(0))/t$  is non-decreasing in  $t$  to obtain

$$\frac{f(x^* + t\frac{y}{\|y\|}) - f(x^*)}{t} = \frac{h_y(t) - h_y(0)}{t} \geq \frac{\lambda t^2}{2t} = \frac{\lambda t}{2}.$$

Combining the two preceding displays, we have  $f(x^* + t\frac{y}{\|y\|}) - f(x^*) \geq \frac{\lambda t}{2} \min\{\epsilon, t\}$ , which is equivalent to inequality (13).  $\square$

## 4.2 Proof of Theorem 2

Before beginning the proof of the theorem proper, we state a martingale convergence lemma necessary for our development, then give an outline of the proof to come.

**Lemma 2** (Robbins and Siegmund [30]). *Let  $\mathcal{F}_1 \subset \mathcal{F}_2 \subset \dots$  be a filtration and  $V_n, \beta_n, \kappa_n, \varepsilon_n$  be non-negative  $\mathcal{F}_n$ -measurable random variables such that*

$$\mathbb{E}[V_{n+1} \mid \mathcal{F}_n] \leq (1 + \beta_n)V_n + \kappa_n - \varepsilon_n.$$

*On the event that  $\sum_{n=1}^{\infty} \beta_n < \infty$  and  $\sum_{n=1}^{\infty} \kappa_n < \infty$ , we have  $V_n \xrightarrow{a.s.} V$  for a non-negative random variable  $V$  with  $V < \infty$  almost surely, and  $\sum_{n=1}^{\infty} \varepsilon_n < \infty$  a.s.*

We prove Theorem 2 by relating the sequence  $x_k$  from expression (2) to a sequence whose performance is somewhat easier to analyze, and which has values more closely approximating a “correct” stochastic gradient iteration: we define

$$\tilde{x}_k := - \sum_{i=1}^{k-1} \alpha_i g_i \quad \text{and} \quad \tilde{\Delta}_k := \tilde{x}_k - x^*. \quad (15)$$

With this iteration, we have that  $\tilde{x}_k \in \mathcal{F}_{k-1}$ , where  $\mathcal{F}_k$  is the  $\sigma$ -field defined in expression (6), and (we show) it is close enough to the correct iterates  $x_k$  to give our desired results. The idea of analyzing a corrected sequence for distributed iterations builds out of Bertsekas and Tsitsiklis [5, Chapter 7.8] and has been used [e.g. 9] for distributed and parallel optimization problems.

Because delays may grow unboundedly under Assumption A, we consider the effects of delay increasing polynomially with  $n$ . With this in mind, for the proof and all internal lemmas, we let  $\rho$  be a fixed constant satisfying

$$\frac{1}{\tau - 1} < \rho < \beta - \frac{1}{1 + \gamma}, \quad (16)$$

where the stepsizes  $\alpha_k = \alpha k^{-\beta}$ ,  $\tau$  is the moment in Assumption A, and  $\gamma \in (0, 1]$  is the power in Assumption D. The interval (16) is assumed to be non-empty by the conditions of the theorem. Note that this implies that  $\rho \in (\frac{1}{\tau-1}, \beta - \frac{1}{2})$ . We will show that  $n^\rho$  functions as a bound on the delays in incorporating gradient information.

**Outline of proof** We provide a brief outline before giving the remainder of the proof. First, we show that there is (asymptotically) a finite bound such that the delays  $M_n$  on asynchronous updates are at most of order  $n^\rho$  by Assumption A (Lemmas 3 and 4). Then we show that the “corrected” sequence  $\tilde{x}_k$  (and  $\tilde{\Delta}_k$ ) converges appropriately (Lemma 5), assuming that the true errors  $\Delta_k$  do not diverge, giving almost sure convergence of  $\tilde{\Delta}_k$  using the Robbins-Siegmund martingale convergence theorem (Lemma 2). We use these results to show that  $\Delta_k$ ,  $\tilde{\Delta}_k$ , and  $\tilde{\Delta}'_k$ , where  $\tilde{\Delta}'_k$  is defined by the simpler linear matrix iteration  $v_{k+1} = (I - \alpha_k H)v_k - \alpha_k \xi_k$  with  $\tilde{\Delta}'_k = v_k - x^*$ , are all asymptotically equivalent in probability (Lemmas 6, 7, and 8), as long as the errors  $\Delta_k$  are assumed to stay bounded. In particular, the differences  $\|\Delta_k - \tilde{\Delta}_k\|^2 = O_P(\alpha_k^2 k^{2\rho})$ , that is, they scale quadratically in the stepsize  $\alpha_k$  and with some penalty for delays, and the errors tend to zero as long as  $\rho$  is not too large; asynchrony is dominated by the magnitude of observed gradient noise. Asymptotic normality (Lemma 10) of the equivalent sequences  $\Delta_k, \tilde{\Delta}_k, \tilde{\Delta}'_k$  then follows from results of Polyak and Juditsky [26], which guarantee a central limit theorem for the sequence  $\tilde{\Delta}'_k$ ,

because the error bounds on  $\zeta$  of Assumption E guarantee that  $\xi_k$  eventually behaves like an i.i.d. sequence. Lastly (in Lemma 11), we show that our overarching assumption—that the true errors  $\Delta_k$  did not diverge—in fact holds under the assumptions of the theorem.

We now turn to the proof of Theorem 2 proper.

**Lemma 3.** *Let  $\rho > \frac{1}{\tau-1}$  and  $\mathcal{E}_n$  be the event that  $E^{nk} \neq I_{d \times d}$  for some  $k \leq n - n^\rho$ . Then*

$$\mathbb{P}(\mathcal{E}_n \text{ occurs infinitely often}) = 0.$$

**Proof** We have that  $E^{nk} \neq I$  if and only if  $M_k \geq n - k + 1$ , so that

$$\mathbb{P}(I \neq E^{nk}) \leq \mathbb{P}(M_k \geq n - k + 1) \leq \frac{\mathbb{E}[M_k^\tau]}{(n - k + 1)^\tau} \leq \frac{M^\tau}{(n - k + 1)^\tau}.$$

Letting  $\mathcal{E}_n$  be the event that  $I \neq E^{nk}$  for some  $k \leq n - n^\rho$  as in the statement of the lemma,

$$\begin{aligned} \mathbb{P}(\mathcal{E}_n) &= \mathbb{P}(M_k \geq n - k + 1 \text{ for some } k \leq n - n^\rho) \leq \sum_{k=1}^{n-n^\rho} \frac{M^\tau}{(n - k + 1)^\tau} \\ &= \sum_{k=n^\rho+1}^n \frac{M^\tau}{k^\tau} \lesssim \int_{n^\rho}^n t^{-\tau} dt \lesssim (n^\rho)^{1-\tau} = n^{\rho(1-\tau)}. \end{aligned}$$

Thus we find that

$$\sum_{n=1}^{\infty} \mathbb{P}(\mathcal{E}_n) \lesssim \sum_{n=1}^{\infty} \frac{1}{n^{\rho(\tau-1)}} \stackrel{(i)}{<} \infty,$$

where inequality (i) holds if and only if  $\rho(\tau - 1) > 1$ , or  $\rho > \frac{1}{\tau-1}$ . Applying the Borel-Cantelli lemma gives the result.  $\square$

As an immediate consequence of this lemma, we obtain the following.

**Lemma 4.** *Let  $\alpha_k = \alpha k^{-\beta}$ , where  $\beta \in [0, 1)$ . For any  $\rho \in (\frac{1}{\tau-1}, 1)$ , with probability 1 we have*

$$\limsup_n \sup_{z \in \mathbb{R}_+^n} \frac{\sum_{k=1}^n \|I - E^{nk}\| z_k}{\sum_{k=n-n^\rho}^n z_k} \leq 1 \quad \text{and} \quad \sup_n \sup_{z \in \mathbb{R}_+^n} \frac{\sum_{k=1}^n \alpha_k \|I - E^{nk}\| z_k}{\alpha_n \sum_{k=n-n^\rho}^n z_k} < \infty,$$

where we treat  $0/0 = 1$ .

**Proof** The first statement of the lemma follows from Lemma 3, as with probability 1 over the delays  $M_k$  and delay matrices  $E^{nk}$ , there exists some (random)  $N$  such that  $n \geq N$  implies that  $E^{nk} = I$  for all  $k \leq n - n^\rho$ , so that for  $n \geq N$ , we have for all nonnegative sequences  $z_1, z_2, \dots$  that

$$\frac{\sum_{k=1}^n \|I - E^{nk}\| z_k}{\sum_{k=n-n^\rho}^n z_k} \leq \frac{\sum_{k=n-n^\rho}^n z_k}{\sum_{k=n-n^\rho}^n z_k} = 1.$$

The second follows from the first once we note that for  $k \in [n - n^\rho, n]$ , we have

$$1 \geq \frac{\alpha_n}{\alpha_k} \geq n^{-\beta} (n - n^\rho)^\beta = (1 - n^{\rho-1})^\beta = \exp(-\beta n^{\rho-1})(1 + o(1)) \rightarrow 1$$

as  $n \rightarrow \infty$ . The limit supremum is finite, so the supremum must likewise be finite.  $\square$

In particular, Lemma 4 implies that any sequence  $\alpha_n \sum_{k=1}^n \alpha_k \|I - E^{nk}\| Z_k$  cannot diverge more quickly than  $\alpha_n^2 \sum_{k=n-n^\rho}^n Z_k$ . We will use this fact frequently. For the remainder of the proof of Theorem 2, we define the random variable (implicitly depending on the power  $\rho$  chosen in the interval (16))

$$K_n := \max_{m \leq n} \sup_{z \in \mathbb{R}_+^n} \frac{\sum_{k=1}^m \alpha_k \|I - E^{nk}\| z_k}{\alpha_m \sum_{k=m-m^\rho}^m z_k} \quad \text{and} \quad K_\infty := \limsup_n K_n. \quad (17)$$

As  $K_n$  are non-decreasing, we have  $K_\infty = \lim_n K_n = \sup_n K_n$ , and by Lemma 4, we see that with probability 1 over the delay process, we have  $K_\infty < \infty$ , and moreover, we have  $K_n \in \mathcal{F}_{n-1}$  by definition (6) of the  $\sigma$ -fields  $\mathcal{F}_k$ . For  $t \in \mathbb{R}$  define

$$\mathcal{K}_{n,t} := \{K_n \leq t\} \quad \text{and} \quad \mathcal{K}_{\infty,t} := \bigcap_n \mathcal{K}_{n,t} = \left\{ \sup_n K_n \leq t \right\} \quad (18)$$

be the events that  $K_n$  and  $K_\infty$  are bounded by  $t$ , respectively, noting that  $\mathcal{K}_{n,t} \in \mathcal{F}_{n-1}$  as  $K_n \in \mathcal{F}_{n-1}$  as before. Then Lemma 4 implies

$$\lim_{t \rightarrow \infty} \mathbb{P}(\bigcap_{n \geq 1} \mathcal{K}_{n,t}) = \lim_{t \rightarrow \infty} \mathbb{P}(\mathcal{K}_{\infty,t}) = \lim_{t \rightarrow \infty} \mathbb{P}\left(\sup_n K_n \leq t\right) = 1.$$

Our first lemma, whose proof we provide in Sec. A.1, builds off of the Robbins-Siegmund martingale convergence theorem (Lemma 2) to give an almost sure convergence result for the corrected sequence  $\tilde{\Delta}_n$ .

**Lemma 5.** *Let Assumptions D and A hold and the stepsizes  $\alpha_k = \alpha k^{-\beta}$ .*

(a) *If Assumption F holds, let  $t < \infty$  and assume additionally that  $\sup_n \mathbb{E}[\mathbf{1}\{\mathcal{K}_{n,t}\} \|\Delta_n\|^2] < \infty$ . Then there is a finite random variable  $V_t$  such that  $\mathbf{1}\{\mathcal{K}_{n,t}\} V(\tilde{\Delta}_n) \xrightarrow{a.s.} V_t$  and*

$$\sum_{n=1}^{\infty} \alpha_n \mathbf{1}\{\mathcal{K}_{n,t}\} \left\langle \nabla V(\tilde{\Delta}_n), R(\tilde{x}_n) \right\rangle < \infty.$$

(b) *If Assumption F' holds, there is a finite random variable  $V$  such that  $V(\tilde{\Delta}_n) \xrightarrow{a.s.} V$ , and*

$$\sum_{n=1}^{\infty} \alpha_n \left\langle \nabla V(\tilde{\Delta}_n), R(\tilde{x}_n) \right\rangle < \infty.$$

We can now verify that  $\tilde{\Delta}_n \xrightarrow{a.s.} 0$  under the conditions of Lemma 5. First, let Assumption F' hold, and let  $\epsilon > 0$  be the radius for which  $\langle \nabla V(x - x^*), R(x) \rangle \geq \lambda_0 V(x - x^*)$  for  $\|x - x^*\| \leq \epsilon$ . With  $c_0 := \inf_{x: \|x - x^*\| > \epsilon} \langle \nabla V(x - x^*), R(x) \rangle > 0$ , we have

$$\sum_{n=1}^{\infty} \alpha_n \min\{\lambda_0 V(\tilde{\Delta}_n), c_0\} \leq \sum_{n=1}^{\infty} \alpha_n \left\langle \nabla V(\tilde{\Delta}_n), R(\tilde{x}_n) \right\rangle < \infty,$$

and  $V(\tilde{\Delta}_n) \xrightarrow{a.s.} V$  for some random variable  $V$ . If  $\mathbb{P}(V > 0) > 0$ , there exist realizations of the randomness in the problem such that  $V > 0$ , and for such realizations there must exist  $\epsilon_0 > 0$  such that  $V(\tilde{\Delta}_n) \geq \epsilon_0$  for all sufficiently large  $n$  as  $V(\tilde{\Delta}_n) \xrightarrow{a.s.} V$ ; this contradicts  $\sum_{n=1}^{\infty} \alpha_n = \infty$ , so we must have  $V = 0$  a.s. under Assumption F'.

Under the alternate Assumption F and that  $\sup_n \mathbb{E}[\mathbf{1}\{\mathcal{K}_{n,t}\} \|\Delta_n\|^2] < \infty$  for all  $t$ , we have

$$\sum_{n=1}^{\infty} \mathbf{1}\{\mathcal{K}_{n,t}\} \alpha_n \lambda_0 V(\tilde{\Delta}_n) \leq \sum_{n=1}^{\infty} \mathbf{1}\{\mathcal{K}_{n,t}\} \alpha_n \left\langle \nabla V(\tilde{\Delta}_n), R(\tilde{x}_n) \right\rangle < \infty.$$

As  $\sum_n \alpha_n = \infty$ , we must then have  $V(\tilde{\Delta}_n) \mathbf{1}\{\mathcal{K}_{n,t}\} \xrightarrow{a.s.} 0$ , as we know it converges to something by Lemma 5. Using that  $\lim_{t \rightarrow \infty} \mathbb{P}(\mathcal{K}_{\infty,t}) = 1$  and  $\mathcal{K}_{n,t} \supset \mathcal{K}_{n+1,t}$  for all  $n$ , we find

$$\mathbb{P}(V(\tilde{\Delta}_n) \not\rightarrow 0) = \lim_{t \rightarrow \infty} \mathbb{P}(\mathcal{K}_{\infty,t} \text{ and } V(\tilde{\Delta}_n) \not\rightarrow 0) \leq \limsup_{t \rightarrow \infty} \mathbb{P}\left(\mathbf{1}\{\mathcal{K}_{n,t}\} V(\tilde{\Delta}_n) \not\rightarrow 0\right) = 0$$

by the preceding discussion. In particular, we have

$$V(\tilde{\Delta}_n) \xrightarrow{a.s.} 0 \tag{19}$$

whenever the conditions of Lemma 5 hold.

Now we show that the averages of  $\tilde{\Delta}_n$  and  $\Delta_n$  are asymptotically equivalent in distribution, and we have quantitative control over this equivalence. (See Section A.2 for a proof of this lemma.)

**Lemma 6.** *In addition to the conditions of Lemma 5, assume that  $\beta > \frac{1}{\tau-1} + \frac{1}{2}$  and either (a) Assumption F holds and the sequence  $C_{\Delta,n,t}^2 = \max_{k \leq n} \mathbb{E}[\mathbf{1}\{\mathcal{K}_{k,t}\} \|\Delta_k\|^2]$  satisfies  $\sup_n C_{\Delta,n,t} < \infty$  for all  $t \in \mathbb{R}$  or (b) Assumption F' holds. In case (a), there exists a universal constant  $C$  such that*

$$\mathbb{E}\left[\mathbf{1}\{\mathcal{K}_{n,t}\} \|\tilde{\Delta}_n - \Delta_n\|^2\right] \leq C \alpha_n^2 n^{2\rho} t^2 (C_{\Delta,n-1,t}^2 + 1). \tag{20}$$

Additionally,

$$\sqrt{n} \left( \overline{\tilde{\Delta}_n} - \overline{\Delta_n} \right) = \frac{1}{\sqrt{n}} \sum_{k=1}^n (\tilde{\Delta}_k - \Delta_k) \xrightarrow{p} 0, \quad \frac{1}{\sqrt{n}} \sum_{k=1}^n \|\tilde{\Delta}_k - \Delta_k\| \xrightarrow{p} 0, \tag{21}$$

and

$$V(\Delta_n) \xrightarrow{a.s.} 0. \tag{22}$$

Thus, any distributional convergence results we are able to show on  $n^{-\frac{1}{2}} \sum_{k=1}^n \tilde{\Delta}_k$  will also hold for the uncorrected sequence  $n^{-\frac{1}{2}} \sum_{k=1}^n \Delta_k$  as long as the conditions of Lemma 6 hold. With this in mind, we give an additional equivalence result showing that  $\tilde{\Delta}_k$  is equivalent to an easier to analyze sequence of errors generated from a simpler matrix iteration. Let the noise sequence  $\{\xi_k\}$  be generated as in the iterations (ii)–(iii), and consider the two iterations

$$\tilde{\Delta}_{k+1} = \tilde{\Delta}_k - \alpha_k (R(x_k) + \xi_k) \quad \text{and} \quad \tilde{\Delta}'_{k+1} = (I - \alpha_k H) \tilde{\Delta}'_k - \alpha_k \xi_k. \tag{23}$$

We have the following two results, which show (under slightly different conditions) that the differences between the iterations (23) tends to zero. The proofs of the lemmas are quite similar, so we put material relevant for the proof of both in Section A.3, specializing to each of the lemmas in sections A.3.1 and A.3.2, respectively.

**Lemma 7.** *Let Assumption D hold with some  $\gamma \in (0, 1]$ . Let the stepsizes  $\alpha_k = \alpha k^{-\beta}$ , where  $\frac{1}{1+\gamma} + \frac{1}{\tau-1} < \beta < 1$ . Let Assumption F hold and assume that for each  $t \in \mathbb{R}$  there is a constant  $C_{\Delta,t} < \infty$  such that  $\mathbb{E}[\mathbf{1}\{\mathcal{K}_{n,t}\} \|\Delta_n\|_2^2] \leq C_{\Delta,t}^2$  for all  $n$ . Then*

$$n^{-\frac{1}{2}} \sum_{k=1}^n (\tilde{\Delta}_k - \tilde{\Delta}'_k) \xrightarrow{P} 0.$$

**Lemma 8.** *Let Assumption D hold with some  $\gamma \in (0, 1]$ . Let the stepsizes  $\alpha_k = \alpha k^{-\beta}$ , where  $\frac{1}{1+\gamma} + \frac{1}{\tau-1} < \beta < 1$ . Let Assumption F' hold. Then*

$$n^{-\frac{1}{2}} \sum_{k=1}^n (\tilde{\Delta}_k - \tilde{\Delta}'_k) \xrightarrow{P} 0.$$

With the preceding lemmas in place, we require two additional claims that immediately yield our desired convergence guarantee. The first is the asymptotic normality of the matrix product sequence  $\tilde{\Delta}'_k$ , that is, that  $n^{-\frac{1}{2}} \sum_{k=1}^n \tilde{\Delta}'_k$  is asymptotically normal. The second is that under Assumption F, we have  $\sup_k \mathbb{E}[\|\Delta_k\|^2 \mathbf{1}\{\mathcal{K}_{k,t}\}] < \infty$  for all  $t \in \mathbb{R}$ . We begin with the first result.

**Lemma 9** (Polyak and Juditsky [26], Theorem 1). *Let  $\{\xi_k\}$  be a martingale difference sequence adapted to the filtration  $\mathcal{F}_k$ , so that  $\mathbb{E}[\xi_k | \mathcal{F}_{k-1}] = 0$  and  $\sup_k \mathbb{E}[\|\xi_k\|^2 | \mathcal{F}_{k-1}] < \infty$  with probability 1. Assume additionally that*

$$\lim_{c \rightarrow \infty} \limsup_{k \rightarrow \infty} \mathbb{E}[\|\xi_k\|^2 \mathbf{1}\{\|\xi_k\| > c\} | \mathcal{F}_{k-1}] = 0 \text{ in probability}$$

and that as  $k \rightarrow \infty$  we have

$$\text{Cov}(\xi_k | \mathcal{F}_{k-1}) \xrightarrow{P} \Sigma \succ 0,$$

where  $\text{Cov}(\xi | \mathcal{F}) = \mathbb{E}[\xi \xi^\top | \mathcal{F}]$ . Then if  $H \succ 0$ , the iteration

$$\tilde{\Delta}'_{k+1} = (I - \alpha_k H) \tilde{\Delta}'_k - \alpha_k \xi_k$$

with  $\alpha_k = \alpha k^{-\beta}$  satisfies

$$\frac{1}{\sqrt{n}} \sum_{k=1}^n \tilde{\Delta}'_k \xrightarrow{d} \mathbf{N}(0, H^{-1} \Sigma H^{-1}).$$

We verify that the conditions of Lemma 9 hold in the two settings captured by Theorem 2, that is, under Assumption F and F'. For Assumption F', we immediately have each condition on  $\xi_k$  except that  $\text{Cov}(\xi_k | \mathcal{F}_{k-1}) \xrightarrow{P} \Sigma$  for some positive definite matrix  $\Sigma$ . But we have  $\Delta_k \xrightarrow{a.s.} 0$  as in expression (22), so that by Assumption E

$$\text{Cov}(\xi_k | \mathcal{F}_{k-1}) = \mathbb{E}[\xi_k(0) \xi_k(0)^\top + \zeta(x_k) \zeta(x_k)^\top | \mathcal{F}_{k-1}] = \Sigma + o_P(1) + O_P(\|x_k - x^*\|^2) \xrightarrow{P} \Sigma.$$

Thus, the conditions of Lemma 9 hold under Assumption F'. We now argue that the conditions of the lemma hold under Assumption F and the additional condition that  $\sup_k \mathbb{E}[\|\Delta_k\|^2 \mathbf{1}\{\mathcal{K}_{k,t}\}] < \infty$  for all  $t \in \mathbb{R}$ . In this case, we still know that  $\mathbb{E}[\xi_k | \mathcal{F}_{k-1}] = 0$ , and by Assumption E, we have

$$\begin{aligned} \sup_k \mathbb{E}[\|\xi_k\|^2 | \mathcal{F}_{k-1}] &\leq 2 \sup_k \mathbb{E}[\|\xi_k(0)\|^2 + \|\zeta(x_k)\|^2 | \mathcal{F}_{k-1}] \\ &\leq 2 \sup_k \mathbb{E}[\|\xi_k(0)\|^2 | \mathcal{F}_{k-1}] + C \sup_k \|\Delta_k\|^2 < \infty, \end{aligned}$$

as  $\|\Delta_k\| \xrightarrow{a.s.} 0$  by the result (22). The second condition on the limits of  $\mathbb{E}[\|\xi_k\|^2 \mathbf{1}\{\|\xi_k\| > c\} \mid \mathcal{F}_{k-1}]$  holds similarly, as we have  $\lim_{c \rightarrow \infty} \mathbb{P}(\sup_k \|\Delta_k\| > c) = 0$  because  $\|\Delta_k\| \xrightarrow{a.s.} 0$ , again using the guarantee (22) if  $\sup_k \mathbb{E}[\mathbf{1}\{\mathcal{K}_{k,t}\} \|\Delta_k\|^2] < \infty$ . The covariance condition follows identically to the argument under Assumption F'. Summarizing, we have shown the following result.

**Lemma 10.** *Let either (i) Assumption F hold and assume that  $\sup_k \mathbb{E}[\|\Delta_k\|^2 \mathbf{1}\{\mathcal{K}_{k,t}\}] < \infty$  for each  $t$  or (ii) Assumption F' hold. Then the sequence  $\tilde{\Delta}'_k$  defined by the iteration (23) satisfies*

$$\frac{1}{\sqrt{n}} \sum_{k=1}^n \tilde{\Delta}'_k \overset{d}{\rightsquigarrow} \mathbf{N}(0, H^{-1} \Sigma H^{-1}).$$

By Lemmas 7 and 8 and the probabilistic equivalence (21) of the sequences  $\sqrt{n} \tilde{\Delta}_n$  and  $\sqrt{n} \tilde{\Delta}'_n$ , Lemma 10 implies that (under the conditions of the lemma)

$$\frac{1}{\sqrt{n}} \sum_{k=1}^n \Delta_k \overset{d}{\rightsquigarrow} \mathbf{N}(0, H^{-1} \Sigma H^{-1}).$$

This is the statement of the theorem under Assumption F'; the proof of the theorem will be complete if we can show that under the strong convexity Assumption F we have  $\sup_k \mathbb{E}[\mathbf{1}\{\mathcal{K}_{k,t}\} \|\Delta_k\|^2] < \infty$  for all  $t \in \mathbb{R}$ .

We present a final lemma that gives the boundedness of the sequences  $\mathbb{E}[\mathbf{1}\{\mathcal{K}_{k,t}\} \|\Delta_k\|^2]$ .

**Lemma 11.** *Let Assumption F hold and  $\epsilon > 0$ ,  $t \in \mathbb{R}$ . There exists some  $N = N(\epsilon, t) \in \mathbb{N}$  such that  $n \geq N$  implies*

$$\mathbb{E}[\|\tilde{\Delta}_{n+1}\|^2 \mathbf{1}\{\mathcal{K}_{n+1,t}\}] \leq \epsilon^2 \max_{k \leq n} \max\{1, \mathbb{E}[\mathbf{1}\{\mathcal{K}_{k,t}\} \|\Delta_k\|^2]\}.$$

Now, Lemma 6, inequality (20), and the fact that the stepsize power  $\beta > \rho$  (recall the interval (16)) immediately yields that for any  $\epsilon > 0$  and  $t \in \mathbb{R}$  there is some  $N = N(\epsilon, t) \in \mathbb{N}$  such that  $n \geq N$  implies

$$\mathbb{E}[\mathbf{1}\{\mathcal{K}_{n,t}\} \|\tilde{\Delta}_n - \Delta_n\|^2] \leq \epsilon^2 \max_{k < n} \max\{1, \mathbb{E}[\mathbf{1}\{\mathcal{K}_{k,t}\} \|\Delta_k\|^2]\}.$$

We combine these inequalities and Lemma 11 to argue that under Assumption F, we have  $\sup_k C_{\Delta,k,t}^2 = \sup_k \mathbb{E}[\mathbf{1}\{\mathcal{K}_{k,t}\} \|\Delta_k\|^2] < \infty$ . Indeed, choose  $\epsilon > 0$ ,  $t \in \mathbb{R}$  and  $N = N(\epsilon, t)$  such that  $n \geq N$  implies

$$\mathbb{E}[\|\tilde{\Delta}_n - \Delta_n\|^2 \mathbf{1}\{\mathcal{K}_{n,t}\}] \leq \frac{\epsilon^2}{4} \max_{k < n} \max\{1, \mathbb{E}[\mathbf{1}\{\mathcal{K}_{k,t}\} \|\Delta_k\|^2]\}$$

and

$$\mathbb{E}[\mathbf{1}\{\mathcal{K}_{n,t}\} \|\tilde{\Delta}_n\|^2] \leq \frac{\epsilon^2}{4} \max_{k < n} \max\{1, \mathbb{E}[\mathbf{1}\{\mathcal{K}_{k,t}\} \|\Delta_k\|^2]\}.$$

Then

$$\begin{aligned} \mathbb{E}[\mathbf{1}\{\mathcal{K}_{n,t}\} \|\Delta_n\|^2] &\leq 2\mathbb{E}[\mathbf{1}\{\mathcal{K}_{n,t}\} \|\Delta_n - \tilde{\Delta}_n\|^2] + 2\mathbb{E}[\mathbf{1}\{\mathcal{K}_{n,t}\} \|\tilde{\Delta}_n\|^2] \\ &\leq \epsilon^2 \max_{k < n} \max\{1, \mathbb{E}[\mathbf{1}\{\mathcal{K}_{k,t}\} \|\Delta_k\|^2]\}. \end{aligned}$$

Repeating this chain of inequalities for all  $k$  such that  $N \leq k < n$ , we find that

$$\mathbb{E} \left[ \mathbf{1} \{ \mathcal{K}_{n,t} \} \|\Delta_n\|^2 \right] \leq \epsilon^2 \max_{k \leq N} \max \{ 1, \mathbb{E} [ \mathbf{1} \{ \mathcal{K}_{k,t} \} \|\Delta_k\|^2 ] \}$$

for all  $n > N$ . As  $\max_{k \leq N} \mathbb{E} [\|\Delta_k\|^2] < \infty$  for any finite  $N$ , this shows that there exists a constant  $C_{\Delta,t} < \infty$  such that  $\sup_k \mathbb{E} [\mathbf{1} \{ \mathcal{K}_{k,t} \} \|\Delta_k\|^2] \leq C_{\Delta}^2$  whenever Assumption F holds and the stepsizes  $\alpha_k$  are chosen such that  $\alpha_k = \alpha k^{-\beta}$  for  $\beta \in (\frac{1}{2}, 1)$  and satisfying  $\rho < \beta - \frac{1}{2}$ . This completes the proof of Theorem 2.

## 5 Discussion and conclusions

In this paper, we have analyzed an asynchronous gradient method, based on Niu et al.'s HOGWILD! [25], for the solution of stochastic convex optimization and variational equality problems. Our work shows particularly that asynchrony introduces essentially negligible penalty for stochastic optimization problems under standard optimization assumptions, which can be leveraged in the development of extremely fast optimization procedures. Our experimental results in Section 3 show that there is still work to be done in terms of a deep understanding of implementation of these methods. In particular, even without inherent competition for locks or other synchronization resources in the computer, there can be competition for other resources, such as memory access. As Table 1 demonstrates, even moderately careful control of memory accesses can be extremely beneficial, and without it, asynchronous methods do not enjoy the performance benefits made possible by multi-core and multi-processor systems. It will thus be beneficial, in future work we hope to undertake, to develop an understanding of memory access and use similar to that now well-known in the scientific computing literature (see, for example, Ballard et al. [4]). This understanding will greatly improve the practical effectiveness of stochastic and asynchronous methods.

## A Technical proofs for Theorem 2

In this appendix, we collect the technical proofs required for Theorem 2. We also state a few additional technical lemmas.

**Lemma 12.** *Define  $C_{\Delta,n,t} := \max_{k \leq n} \mathbb{E} [\mathbf{1} \{ \mathcal{K}_{n,t} \} \|\Delta_k\|^2]$ . If Assumption F holds, there is a constant  $c < \infty$  independent of  $n$  and  $C_{\Delta,n,t}$  such that for any  $l \in \mathbb{R}_+$ ,*

$$\mathbb{E} \left[ \mathbf{K}_n^l \mathbf{1} \{ \mathcal{K}_{n,t} \} \sum_{k=n-n^\rho}^{n-1} \|R(x_k) + \xi_k\|^2 \right] \leq c \cdot t^l n^\rho (C_{\Delta,n-1,t}^2 + 1).$$

*If Assumption F' holds, there is a constant  $c < \infty$  independent of  $n$  such that*

$$\mathbb{E} \left[ \sum_{k=n-n^\rho}^{n-1} \|R(x_k) + \xi_k\|^2 \right] \leq cn^\rho.$$

**Proof** We begin with the result under Assumption F, where we use  $\mathbb{E} [\|\Delta_k\|^2 \mathbf{1} \{ \mathcal{K}_{n,t} \}] \leq C_{\Delta,n-1,t}^2$  for  $k \leq n-1$ . We also know that

$$\|R(x_k)\|^2 \lesssim \|\Delta_k\|^2 \quad \text{and} \quad \mathbb{E} [\|\xi_k\|^2 \mid \mathcal{F}_{k-1}] \lesssim \|\Delta_k\|^2 + 1,$$

where we have used Assumption E, and thus we have  $\mathbb{E}[\mathbb{E}[\mathbf{1}\{\mathcal{K}_{k,t}\} \|\xi_k\|^2 \mid \mathcal{F}_{k-1}]] \leq c(C_{\Delta,n-1,t}^2 + 1)$ . In addition, we have  $\|R(x)\| = \|R(x) - R(x^*)\| \leq L \|x - x^*\|$  and  $\mathbf{K}_n^l \mathbf{1}\{\mathcal{K}_{n,t}\} \leq t^l \mathbf{1}\{\mathcal{K}_{n,t}\}$ , so

$$\begin{aligned} \mathbb{E} \left[ \mathbf{K}_n \mathbf{1}\{\mathcal{K}_{n,t}\} \sum_{k=n-n^\rho}^{n-1} \|R(x_k) + \xi_k\|^2 \right] &\leq 2t^l \sum_{k=n-n^\rho}^{n-1} \mathbb{E} \left[ \mathbf{1}\{\mathcal{K}_{n,t}\} \|R(x_k)\|^2 + \mathbf{1}\{\mathcal{K}_{n,t}\} \|\xi_k\|^2 \right] \\ &\leq 2Ct^l n^\rho (LC_{\Delta,n-1,t}^2 + c(C_{\Delta,n-1,t}^2 + 1)), \end{aligned}$$

where we have used that  $\mathbf{1}\{\mathcal{K}_{n,t}\} \leq \mathbf{1}\{\mathcal{K}_{n-1,t}\}$  for all  $n$ . Under Assumption F', the result is simpler: we have simply that  $\mathbb{E}[\|\xi_k\|^2 \mid \mathcal{F}_{k-1}] \leq C$  and  $\|R(x_k)\| \leq C$ , giving the result.  $\square$

We also give two technical results involving integral convergence.

**Lemma 13.** *Let  $c > 0$  and  $\kappa \in (0, 1)$  be constants and  $b \geq a > 0$ . Then*

$$\int_a^b \exp(-c(t^\kappa - a^\kappa)) dt \leq \frac{\max\{2^{\frac{1-\kappa}{\kappa}} - 1, 1\}}{\kappa c} \left[ c^{\frac{\kappa-1}{\kappa}} \Gamma\left(\frac{1}{\kappa}\right) + a^{1-\kappa} \right].$$

See Appendix A.5 for a proof of Lemma 13. The final technical result we use also gives a bound on (essentially) another gamma integral.

**Lemma 14.** *Let  $\beta \in (\frac{1}{2}, 1)$  and  $\rho < \beta - \frac{1}{2}$ . Then*

$$\lim_{n \rightarrow \infty} \sum_{k=1}^n k^{\rho-2\beta} \exp(-c(n^{1-\beta} - k^{1-\beta})) = 0.$$

See Appendix A.6 for a proof of Lemma 14.

## A.1 Proof of Lemma 5

We begin by using the Lipschitz continuity of the gradients of  $V$  to note that

$$\begin{aligned} V(\tilde{\Delta}_{n+1}) &= V(\tilde{\Delta}_n - \alpha_n g_n) \leq V(\tilde{\Delta}_n) - \alpha_n \langle \nabla V(\tilde{\Delta}_n), g_n \rangle + \frac{L\alpha_n^2}{2} \|g_n\|^2 \\ &= V(\tilde{\Delta}_n) - \alpha_n \langle \nabla V(\tilde{\Delta}_n), R(\tilde{x}_n) \rangle - \alpha_n \langle \nabla V(\tilde{\Delta}_n), R(x_n) - R(\tilde{x}_n) \rangle - \alpha_n \langle \nabla V(\tilde{\Delta}_n), \xi_n \rangle + \frac{L\alpha_n^2}{2} \|g_n\|^2. \end{aligned}$$

Taking expectations conditional on  $\mathcal{F}_{n-1}$ , we have  $x_n, \tilde{x}_n \in \mathcal{F}_{n-1}$ , and  $\mathbb{E}[\xi_n \mid \mathcal{F}_{n-1}] = 0$ . Moreover, we have  $g_n = R(x_n) + \xi_n$ , and thus

$$\begin{aligned} \mathbb{E}[V(\tilde{\Delta}_{n+1}) \mid \mathcal{F}_{n-1}] &\leq V(\tilde{\Delta}_n) - \alpha_n \langle \nabla V(\tilde{\Delta}_n), R(\tilde{x}_n) \rangle + \frac{L\alpha_n^2}{2} \mathbb{E}[\|R(x_n) + \xi_n\|^2 \mid \mathcal{F}_{n-1}] \\ &\quad + \alpha_n \|\nabla V(\tilde{\Delta}_n)\| \|R(x_n) - R(\tilde{x}_n)\|. \end{aligned} \tag{24}$$

Using that  $\|R(x_n) + \xi_n\|^2 \leq 2\|R(x_n)\|^2 + 2\|\xi_n\|^2$  and

$$\|R(x_n) - R(\tilde{x}_n)\| \leq L \|x_n - \tilde{x}_n\| = L \left\| \sum_{k=1}^{n-1} \alpha_k (I - E^{nk}) g_k \right\| \leq L \sum_{k=1}^{n-1} \alpha_k \|I - E^{nk}\| \|g_k\|,$$

we find that there is a constant  $C$  such that

$$\begin{aligned} \mathbb{E}[V(\tilde{\Delta}_{n+1}) \mid \mathcal{F}_{n-1}] &\leq V(\tilde{\Delta}_n) - \alpha_n \left\langle \nabla V(\tilde{\Delta}_n), R(\tilde{x}_n) \right\rangle + C\alpha_n^2 \|R(x_n)\|^2 + C\alpha_n^2 \mathbb{E}[\|\xi_n\|^2 \mid \mathcal{F}_{n-1}] \\ &\quad + C\alpha_n \|\nabla V(\tilde{\Delta}_n)\| \sum_{k=1}^{n-1} \alpha_k \|I - E^{nk}\| \|g_k\|. \end{aligned}$$

Recalling the definition (17) of the random variables  $\mathbf{K}_n$  and the selection (16) of the power  $\rho$ , we have  $\sum_{k=1}^{n-1} \alpha_k \|I - E^{nk}\| \|g_k\| \leq \mathbf{K}_n \alpha_n \sum_{k=n-n^\rho}^{n-1} \|g_k\|$ , and using that  $g_k = R(x_k) + \xi_k$ , we obtain

$$\begin{aligned} \mathbb{E}[V(\tilde{\Delta}_{n+1}) \mid \mathcal{F}_{n-1}] &\leq V(\tilde{\Delta}_n) - \alpha_n \left\langle \nabla V(\tilde{\Delta}_n), R(\tilde{x}_n) \right\rangle + C\alpha_n^2 \|R(x_n)\|^2 + C\alpha_n^2 \mathbb{E}[\|\xi_n\|^2 \mid \mathcal{F}_{n-1}] \\ &\quad + C\alpha_n^2 \|\nabla V(\tilde{\Delta}_n)\| \mathbf{K}_n \sum_{k=n-n^\rho}^{n-1} \|R(x_k) + \xi_k\| \end{aligned} \quad (25a)$$

$$\begin{aligned} &\leq V(\tilde{\Delta}_n) - \alpha_n \left\langle \nabla V(\tilde{\Delta}_n), R(\tilde{x}_n) \right\rangle + C\alpha_n^2 \|\Delta_n\|^2 + C\alpha_n^2 \mathbb{E}[\|\xi_n\|^2 \mid \mathcal{F}_{n-1}] \\ &\quad + C\alpha_n^2 n^\rho V(\tilde{\Delta}_n) + C\alpha_n^2 \mathbf{K}_n^2 \sum_{k=n-n^\rho}^{n-1} \|R(x_k) + \xi_k\|^2, \end{aligned} \quad (25b)$$

the final equality following because  $\|R(x)\| = \|R(x) - R(x^*)\| \leq L \|x - x^*\|$ ,  $\|\nabla V(\Delta)\| \leq L \|\Delta\| \leq (L/\sqrt{\lambda})\sqrt{V(\Delta)}$ , and  $ab \leq \frac{1}{2}a^2 + \frac{1}{2}b^2$  for any  $a, b \in \mathbb{R}$ .

We now use the technical Lemma 12, which allows us to control the error terms in inequality (25b). Indeed, by Lemma 12 and our assumption that  $C_{\Delta,t}^2 = \sup_k \mathbb{E}[\mathbf{1}\{\mathcal{K}_{k,t}\} \|\Delta_k\|^2] < \infty$  if Assumption F holds, we obtain

$$\sum_{n=1}^{\infty} \alpha_n^2 \mathbb{E} \left[ \mathbf{K}_n^2 \mathbf{1}\{\mathcal{K}_{n,t}\} \sum_{k=n-n^\rho}^{n-1} \|R(x_k) + \xi_k\|^2 \right] \leq ct^2 (C_{\Delta,t}^2 + 1) \sum_{n=1}^{\infty} \alpha_n^2 n^\rho \lesssim \int_1^{\infty} u^{-2\beta+\rho} du < \infty, \quad (26)$$

the final inequality holding when  $2\beta - \rho > 1$ , or  $\rho < 2\beta - 1$ . In particular, the Robbins-Siegmund convergence theorem (Lemma 2) applies, as we can write (recall inequality (25a) and that  $\mathbf{1}\{\mathcal{K}_{n,t}\} \leq \mathbf{1}\{\mathcal{K}_{n-1,t}\}$ )

$$\mathbb{E}[\mathbf{1}\{\mathcal{K}_{n,t}\} V(\tilde{\Delta}_{n+1}) \mid \mathcal{F}_{n-1}] \leq (1 + \beta_{n-1}) \mathbf{1}\{\mathcal{K}_{n-1,t}\} V(\tilde{\Delta}_n) + \kappa_{n-1} - \varepsilon_{n-1},$$

where  $\beta_{n-1} = C\alpha_n^2 n^\rho$ ,  $\varepsilon_n = \mathbf{1}\{\mathcal{K}_{n,t}\} \alpha_n \left\langle \nabla V(\tilde{\Delta}_n), R(\tilde{x}_n) \right\rangle$ , and

$$\kappa_{n-1} = \mathbf{1}\{\mathcal{K}_{n,t}\} \left[ C\alpha_n^2 \|R(x_n)\|^2 + C\alpha_n^2 \mathbb{E}[\|\xi_n\|^2 \mid \mathcal{F}_{n-1}] + C\alpha_n^2 \mathbf{K}_n^2 \sum_{k=n-n^\rho}^{n-1} \|R(x_k) + \xi_k\|^2 \right]$$

are all  $\mathcal{F}_{n-1}$ -measurable. Moreover,  $\sum_n \beta_n \lesssim \sum_{n=1}^{\infty} n^{\rho-2\beta} < \infty$  because  $\rho < 2\beta - 1$ , and  $\sum_n \mathbb{E}[\kappa_n] < \infty$  by the fact that  $\mathbb{E}[\mathbb{E}[\|\xi_n\|^2 \mid \mathcal{F}_{n-1}]] \lesssim \mathbb{E}[\|\Delta_n\|^2 + 1]$  (this is Assumption E) coupled with Lemma 12 and inequality (26). We thus conclude that

$$\mathbf{1}\{\mathcal{K}_{n,t}\} V(\tilde{\Delta}_n) \xrightarrow{a.s.} V_t \quad \text{and} \quad \sum_{n=1}^{\infty} \alpha_n \mathbf{1}\{\mathcal{K}_{n,t}\} \left\langle \nabla V(\tilde{\Delta}_n), R(\tilde{x}_n) \right\rangle < \infty$$

with probability 1 whenever Assumption F holds in addition to the assumptions of the lemma.

In the somewhat simpler case that Assumption F' holds, we may simply remove all indicator functions  $\mathbf{1}\{\mathcal{K}_{n,t}\}$ , as Lemma 12 shows that we may replace inequality (26) with

$$\sum_{n=1}^{\infty} \alpha_n^2 \sum_{k=n-n^\rho}^{n-1} \mathbb{E}[\|R(x_k) - \xi_k\|^2] \lesssim C \sum_{n=1}^{\infty} \alpha_n^2 n^\rho \lesssim \int_1^{\infty} u^{-2\beta+\rho} du < \infty,$$

while  $\sup_n \mathsf{K}_n < \infty$  with probability 1.

## A.2 Proof of Lemma 6

We may write the difference

$$\tilde{\Delta}_n - \Delta_n = \tilde{x}_n - x_n = \sum_{k=1}^{n-1} \alpha_k (E^{nk} - I) g_k.$$

Recalling the definition (17) of  $\mathsf{K}_n$  and our choice (16) of  $\rho$ , this representation guarantees that

$$\|\tilde{\Delta}_n - \Delta_n\| \leq \mathsf{K}_n \alpha_n \sum_{k=n-n^\rho}^{n-1} \|g_k\|, \quad (27)$$

and so we have

$$\frac{1}{\sqrt{n}} \sum_{k=1}^n \|\tilde{\Delta}_k - \Delta_k\| \leq \frac{\mathsf{K}_n}{\sqrt{n}} \sum_{k=1}^n \alpha_k \left( \sum_{i=k-k^\rho}^{k-1} \|g_i\| \right). \quad (28)$$

Let us first show that the quantity (28) is well-behaved in the simpler case of Assumption F'. Indeed, we have

$$\mathbb{E} \left[ \frac{1}{\sqrt{n}} \sum_{k=1}^n \alpha_k \sum_{i=k-k^\rho}^{k-1} \|g_i\| \right] \lesssim n^{-\frac{1}{2}} \sum_{k=1}^n \alpha_k k^\rho \lesssim n^{-\frac{1}{2}} \int_1^n u^{\rho-\beta} du \asymp n^{\frac{1}{2}+\rho-\beta},$$

which tends to zero if and only if  $\beta > \rho + \frac{1}{2}$ . Then inequality (28) implies  $n^{-\frac{1}{2}} \sum_{k=1}^n \|\tilde{\Delta}_k - \Delta_k\| \leq \mathsf{K}_n Z_n$ , where  $Z_n \xrightarrow{L_1} 0$  and  $\sup_n \mathsf{K}_n < \infty$  with probability 1 by Lemma 4; thus, the convergence (21) holds under Assumption F'.

We turn to the somewhat more challenging case that Assumption F holds and that for our choice of bound  $t$  on  $\mathsf{K}_n$ , there exist constants  $C_{\Delta,n,t}^2 = \max_{k \leq n} \mathbb{E}[\mathbf{1}\{\mathcal{K}_{k,t}\} \|\Delta_k\|^2]$  such that  $C_{\Delta,t} = \sup_n C_{\Delta,n,t} < \infty$ . In this case, inequality (28) and the definition (18) of the event  $\mathcal{K}_{n,t} = \{\mathsf{K}_{n,t} \leq t\}$  imply

$$\mathbb{E} \left[ \mathbf{1}\{\mathcal{K}_{n,t}\} \frac{1}{\sqrt{n}} \sum_{k=1}^n \|\tilde{\Delta}_k - \Delta_k\| \right] \leq \frac{t}{\sqrt{n}} \sum_{k=1}^n \alpha_k \mathbb{E} \left[ \mathbf{1}\{\mathcal{K}_{n,t}\} \sum_{i=k-k^\rho}^{k-1} \|g_i\| \right]$$

Now, we note that for  $k \leq n$ , we have  $\mathcal{K}_{k,t} \supset \mathcal{K}_{n,t}$  so that  $\mathbf{1}\{\mathcal{K}_{n,t}\} \leq \mathbf{1}\{\mathcal{K}_{k,t}\}$ , and

$$\begin{aligned} \mathbb{E}[\|g_k\|^2 \mathbf{1}\{\mathcal{K}_{n,t}\}] &\leq \mathbb{E}[\|g_k\|^2 \mathbf{1}\{\mathcal{K}_{k,t}\}] \leq 2\mathbb{E}[\|R(x_k)\|^2 \mathbf{1}\{\mathcal{K}_{k,t}\}] + 2\mathbb{E}[\|\xi_k\|^2 \mathbf{1}\{\mathcal{K}_{k,t}\}] \\ &\lesssim C_{\Delta,n-1,t}^2 + C_{\Delta,n-1,t}^2 + 1 \end{aligned} \quad (29)$$

by Assumption E and that  $\mathcal{K}_{k,t} \in \mathcal{F}_{k-1}$ . Thus we have  $\mathbb{E}[\|g_k\|^2 \mathbf{1}\{\mathcal{K}_{n,t}\}] \lesssim C_{\Delta,n-1,t} + 1$ , and by Jensen's inequality and inequality (27), we have

$$\mathbb{E} \left[ \mathbf{1}\{\mathcal{K}_{n,t}\} \|\tilde{\Delta}_n - \Delta_n\|^2 \right] \leq t^2 \alpha_n^2 n^\rho \sum_{k=n-n^\rho}^{n-1} \mathbb{E} \left[ \mathbf{1}\{\mathcal{K}_{n,t}\} \|g_k\|^2 \right] \leq C t^2 \alpha_n^2 n^{2\rho} (C_{\Delta,n-1,t}^2 + 1),$$

where  $C$  is some universal constant, by the bound (29). This gives statement (20) of the lemma. To obtain the convergence guarantee (21) in the case of Assumption F and that  $\sup_n C_{\Delta,n,t} < \infty$ , note that

$$\begin{aligned} \sum_{k=1}^n \alpha_k \mathbb{E} \left[ \mathbf{1}\{\mathcal{K}_{n,t}\} \sum_{i=k-k^\rho}^{k-1} \|g_i\| \right] &\lesssim \sum_{k=1}^n \alpha_k k^\rho (C_{\Delta,k-1,t} + 1) \\ &\lesssim (C_{\Delta,n-1,t} + 1) \int_1^n u^{\rho-\beta} du \asymp (C_{\Delta,n-1,t} + 1) n^{1+\rho-\beta}. \end{aligned}$$

In particular, we have

$$\mathbb{E} \left[ \mathbf{1}\{\mathcal{K}_{n,t}\} \frac{1}{\sqrt{n}} \sum_{k=1}^n \|\tilde{\Delta}_k - \Delta_k\| \right] \lesssim (C_{\Delta,n-1,t} + 1) t n^{\frac{1}{2}+\rho-\beta},$$

which tends to 0 if  $\beta > \rho + \frac{1}{2}$ . Thus, we have shown that for any  $\epsilon > 0$ , we have for any  $t > 0$  that

$$\lim_{n \rightarrow \infty} \mathbb{P} \left( \mathcal{K}_{n,t} \text{ and } n^{-\frac{1}{2}} \sum_{k=1}^n \|\tilde{\Delta}_k - \Delta_k\| > \epsilon \right) = 0. \quad (30)$$

We now use expression (30) to get the desired convergence result in the lemma. Let  $D_n = n^{-\frac{1}{2}} \sum_{k=1}^n \|\tilde{\Delta}_k - \Delta_k\|$  be shorthand for our error sum. Fix  $\delta > 0$  and let  $t$  be large enough that  $\mathbb{P}(\mathcal{K}_{\infty,t}) \geq 1 - \delta$ , which we know is possible by Lemma 4. Then as  $\mathcal{K}_{n,t} \subset \mathcal{K}_{\infty,t}$  by definition, we have

$$\mathbb{P}(D_n > \epsilon) \leq \mathbb{P}(\mathcal{K}_{\infty,t} \text{ and } D_n > \epsilon) + \mathbb{P}(\mathcal{K}_{\infty,t}^c) \leq \mathbb{P}(\mathcal{K}_{n,t} \text{ and } D_n > \epsilon) + \delta.$$

Taking the limit as  $n \rightarrow \infty$ , we find from expression (30) that

$$\limsup_{n \rightarrow \infty} \mathbb{P} \left( \frac{1}{\sqrt{n}} \sum_{k=1}^n \|\tilde{\Delta}_k - \Delta_k\| > \epsilon \right) \leq \delta,$$

and as  $\delta > 0$  was arbitrary, we have the desired convergence guarantee (21).

Lastly, we show that expression (22) holds under the conditions of the lemma. We use inequality (27) and the Borel-Cantelli lemma for this. Under Assumption F and the additional condition that  $C_{\Delta,t}^2 = \sup_n \mathbb{E}[\mathbf{1}\{\mathcal{K}_{n,t}\} \|\Delta_n\|^2] < \infty$  for all  $t$  or Assumption F', we have

$$\begin{aligned} \mathbb{P} \left( \mathcal{K}_{n,t}, \|\tilde{\Delta}_n - \Delta_n\| > \epsilon \right) &\leq \epsilon^{-2} \mathbb{E} \left[ \mathbb{K}_n^2 \alpha_n^2 \left( \sum_{k=n-n^\rho}^{n-1} \|g_k\| \right)^2 \mathbf{1}\{\mathcal{K}_{n,t}\} \right] \\ &\stackrel{(i)}{\leq} \frac{t^2 \alpha_n^2 n^\rho \sum_{k=n-n^\rho}^{n-1} \mathbb{E}[\|g_k\|^2 \mathbf{1}\{\mathcal{K}_{n,t}\}]}{\epsilon^2} \stackrel{(ii)}{\lesssim} \frac{t^2 \alpha_n^2 n^{2\rho}}{\epsilon^2}. \end{aligned}$$

Here inequality (i) follows from Jensen's inequality and the fact that  $\mathbf{K}_n \mathbf{1}\{\mathcal{K}_{n,t}\} \leq t$ , and inequality (ii) follows either by the bound (29) (when Assumption F holds) or because  $\mathbb{E}[\|g_k\|^2] \lesssim 1$  for all  $k$  (when Assumption F' holds). In particular, we have

$$\sum_{n=1}^{\infty} \mathbb{P}\left(\mathcal{K}_{n,t}, \|\tilde{\Delta}_n - \Delta_n\| > \epsilon\right) \lesssim \sum_{n=1}^{\infty} n^{2\rho-2\beta} < \infty$$

whenever  $\beta > \rho + \frac{1}{2}$ , which we have already assumed. Thus, we find that under the assumptions of the lemma, we have  $\mathbf{1}\{\mathcal{K}_{n,t}\} \|\tilde{\Delta}_n - \Delta_n\| > \epsilon$  only finitely many times, for any  $t \in \mathbb{R}$ . By Lemma 4, with probability 1 there is some  $t < \infty$  such that  $\mathbf{K}_\infty \leq t$ , that is, as  $\mathcal{K}_{\infty,t} \subset \mathcal{K}_{n,t}$  it must be the case that  $\mathcal{K}_{n,t}$  always holds. So we find that  $\|\tilde{\Delta}_n - \Delta_n\| > \epsilon$  only finitely many times with probability 1. That is,  $\|\tilde{\Delta}_n - \Delta_n\| \xrightarrow{a.s.} 0$ , and the continuity of  $V$  gives the almost sure convergence (22) as desired, as we know that  $\tilde{\Delta}_n \xrightarrow{a.s.} 0$ .

### A.3 Proof of Lemmas 7 and 8

If we define  $B_l^k = \prod_{i=l}^k (I - \alpha_i H)$ , we have that  $\tilde{\Delta}'_{n+1} = B_1^n \Delta_1 - \sum_{k=1}^n \alpha_k B_{k+1}^n \xi^k$ , and additionally we have

$$\sum_{k=1}^n \tilde{\Delta}'_k = \sum_{k=1}^n B_1^{k-1} \Delta_1 - \sum_{k=1}^n H^{-1} \xi^k + \sum_{k=1}^n W_k^n \xi^k,$$

where the matrix  $W_k^n$  is defined by  $W_k^n = \alpha_k \sum_{l=k+1}^{n-1} B_{k+1}^l - H^{-1}$ . This matrix is well-structured, as the following lemma shows.

**Lemma 15** (Polyak and Juditsky [26], Lemma 1). *Let  $\beta \in (0, 1)$ . Then*

$$\sup_{k,n} \|W_k^n\| < \infty \quad \text{and} \quad \lim_n \frac{1}{n} \sum_{k=1}^n \|W_k^n\| = 0.$$

Thus—as we show rigorously shortly—the behavior of  $\sum_{k=1}^n \tilde{\Delta}'_k$  is governed almost completely by  $\sum_{k=1}^n H^{-1} \xi^k$ .

Now, by the iteration (23), we have that

$$\tilde{\Delta}_{k+1} = \tilde{\Delta}_k - \alpha_k \left( R(x_k) + \xi^k \right) = (I - \alpha_k H) \tilde{\Delta}_k + \alpha_k \underbrace{(H \tilde{\Delta}_k - R(x_k))}_{=: Z_k} - \alpha_k \xi^k,$$

so that by analogy with  $\tilde{\Delta}'_k$  we have

$$\sum_{k=1}^n \tilde{\Delta}'_k = \sum_{k=1}^n B_1^{k-1} \Delta_1 - \sum_{k=1}^n H^{-1} \xi^k + \sum_{k=1}^n W_k^n \xi^k - \sum_{k=1}^n H^{-1} Z_k + \sum_{k=1}^n W_k^n Z_k.$$

Using the iteration (23) for  $\tilde{\Delta}'_k$ , we thus have

$$\sum_{k=1}^n \left( \tilde{\Delta}_k - \tilde{\Delta}'_k \right) = \sum_{k=1}^n (H^{-1} - W_k^n) Z_k = \sum_{i=1}^n (H^{-1} - W_k^n) (H \tilde{\Delta}_k - R(x_k)). \quad (31)$$

Thus, to show that  $\sqrt{n}(\widetilde{\Delta}_n - \widetilde{\Delta}'_n) \xrightarrow{P} 0$ , it suffices to show that the rightmost sum in expression (31) is  $o_P(\sqrt{n})$ .

By Lemma 15, we know that  $\sup_{k,n} \|H^{-1} - W_k^n\| < \infty$ , and thus

$$\left\| \sum_{k=1}^n (\widetilde{\Delta}_k - \Delta'_k) \right\| \lesssim \sum_{i=1}^n \left\| H\widetilde{\Delta}_k - R(x_k) \right\| \leq \sum_{k=1}^n \left( \left\| H\widetilde{\Delta}_k - R(\widetilde{x}_k) \right\| + \left\| R(\widetilde{x}_k) - R(x_k) \right\| \right). \quad (32)$$

We consider each of the right-hand terms in inequality (32) in turn, beginning with the second. In this case, Lemma 6 implies that under the conditions of either Lemma 7 or 8, we have

$$\frac{1}{\sqrt{n}} \sum_{k=1}^n \left\| R(\widetilde{x}_k) - R(x_k) \right\| \leq \frac{L}{\sqrt{n}} \sum_{k=1}^n \left\| \widetilde{x}_k - x_k \right\| = \frac{L}{\sqrt{n}} \sum_{k=1}^n \left\| \widetilde{\Delta}_k - \Delta_k \right\| \xrightarrow{P} 0. \quad (33)$$

We now turn to the error part  $H\widetilde{\Delta}_k - R(\widetilde{x}_k)$  of inequality (32). To that end, let  $\epsilon > 0$  be the value in Assumption D such that  $\|H(x - x^*) - R(x)\| \leq C\|x - x^*\|^{1+\gamma}$  for  $x$  such that  $\|x - x^*\| \leq \epsilon$ . Splitting the sum into two parts, Assumption D thus implies

$$\sum_{k=1}^n \left\| H\widetilde{\Delta}_k - R(\widetilde{x}_k) \right\| \leq \sum_{k=1}^n \left\| H\widetilde{\Delta}_k - R(\widetilde{x}_k) \right\| \mathbf{1} \left\{ \left\| \widetilde{\Delta}_k \right\| > \epsilon \right\} + C \sum_{k=1}^n \left\| \widetilde{\Delta}_k \right\|^{1+\gamma}.$$

As we know that  $\widetilde{\Delta}_k \xrightarrow{a.s.} 0$ , with probability 1 over  $\{\xi^1, \xi^2, \dots\} \cup \{E^{ij}\}_{i \geq j}$  (recall the convergence guarantee (19) after Lemma 5) there exists some (random)  $N(\epsilon) < \infty$  such that  $\|\widetilde{\Delta}_k\| \leq \epsilon$  for all  $k \geq N(\epsilon)$ , so that

$$\lim_{n \rightarrow \infty} \sum_{k=1}^n \left\| H\widetilde{\Delta}_k - R(\widetilde{x}_k) \right\| \mathbf{1} \left\{ \left\| \widetilde{\Delta}_k \right\| > \epsilon \right\} < \infty \text{ w.p. } 1,$$

and

$$n^{-\frac{1}{2}} \sum_{k=1}^n \left\| H\widetilde{\Delta}_k - R(\widetilde{x}_k) \right\| \mathbf{1} \left\{ \left\| \widetilde{\Delta}_k \right\| > \epsilon \right\} \xrightarrow{a.s.} 0.$$

It thus remains to argue that  $n^{-\frac{1}{2}} \sum_{k=1}^n \left\| \widetilde{\Delta}_k \right\|^{1+\gamma} \rightarrow 0$  in probability (or otherwise).

Let  $\epsilon > 0$  be such that  $\langle \nabla V(x - x^*), R(x) \rangle \geq \lambda_0 V(x - x^*)$  for all  $x$  such that  $\|x - x^*\| \leq \epsilon$  (such an  $\epsilon$  certainly exists under both Assumptions F and F'). Define the events

$$\mathcal{E}_a^k = \left\{ \left\| \widetilde{\Delta}_i \right\| \leq \epsilon, \text{ all } i \in \{[a], \dots, k\} \right\}.$$

Dividing the sum into two parts, we have

$$\sum_{k=1}^n \left\| \widetilde{\Delta}_k \right\|^{1+\gamma} \leq \sum_{k=1}^n \left\| \widetilde{\Delta}_k \right\|^{1+\gamma} \mathbf{1} \left\{ \mathcal{E}_{k/2}^{k-1} \right\} + \sum_{k=1}^n \left\| \widetilde{\Delta}_k \right\|^{1+\gamma} \left( 1 - \mathbf{1} \left\{ \mathcal{E}_{k/2}^{k-1} \right\} \right).$$

By the fact that  $\widetilde{\Delta}_k \xrightarrow{a.s.} 0$ , we know that there exists some (random but finite)  $N(\epsilon)$  such that  $\|\widetilde{\Delta}_k\| < \epsilon$  for all  $k > N(\epsilon)$ ; the second term in the preceding display is thus finite with probability one and

$$n^{-\frac{1}{2}} \sum_{k=1}^n \left\| \widetilde{\Delta}_k \right\|^{1+\gamma} \left( 1 - \mathbf{1} \left\{ \mathcal{E}_{k/2}^{k-1} \right\} \right) \xrightarrow{a.s.} 0.$$

By combining expressions (31), (32), and (33) with the above display, we see that to prove Lemma 7 or 8, all that remains to show is that

$$n^{-\frac{1}{2}} \sum_{k=1}^n \|\tilde{\Delta}_k\|^{1+\gamma} \mathbf{1} \left\{ \mathcal{E}_{k/2}^{k-1} \right\} \xrightarrow{p} 0. \quad (34)$$

### A.3.1 Proof of Lemma 7

We give a single-step bound on  $V(\tilde{\Delta}_k)$  that we can use to give the desired convergence guarantee under the conditions of Lemma 7. Recall the definitions (17) and (18) of  $\mathbf{K}_n$  and the associated event  $\mathcal{K}_{n,t} = \{\mathbf{K}_n \leq t\}$ , and recall also our assumption (16) that  $\rho \in (\frac{1}{\tau-1}, \beta - \frac{1}{1+\gamma}) \subset (\frac{1}{\tau-1}, \beta - \frac{1}{2})$ , where  $\rho$  is the power used in the definition of  $\mathbf{K}_n$ . We claim that there exist constants  $c > 0$  and  $C < \infty$ , independent of  $C_{\Delta,t}^2 = \sup_k \mathbb{E}[\mathbf{1} \{\mathcal{K}_{k,t}\} \|\Delta_k\|^2]$ , such that for  $l \in \mathbb{R}$  with  $l < k$ , we have

$$\begin{aligned} \mathbb{E} \left[ V(\tilde{\Delta}_{k+1}) \mathbf{1} \left\{ \mathcal{E}_l^k, \mathcal{K}_{k+1,t} \right\} \right] &\leq \mathbb{E} \left[ (1 - c\alpha_k + C\alpha_k^2 k^\rho) V(\tilde{\Delta}_k) \mathbf{1} \left\{ \mathcal{E}_l^{k-1}, \mathcal{K}_{k,t} \right\} \right] \\ &\quad + Ct^2 \alpha_k^2 k^\rho (C_{\Delta,t}^2 + 1). \end{aligned} \quad (35)$$

We temporarily defer proof of this claim and show how to use it to show  $\sum_{k=1}^n \|\tilde{\Delta}_k\|^{1+\gamma} = o_P(\sqrt{n})$ .

Let  $K < \infty$  be large enough that for the constants  $c, C$  in inequality (35), there is a constant  $c'$  such that for  $k \geq K$  and stepsizes  $\alpha_k = \alpha k^{-\beta}$ , we have

$$(1 - c\alpha_i + C\alpha_i^2 i^\rho) \leq \exp(-c'\alpha_i)$$

for  $i \geq k/2$ . This must be possible as we have assumed  $\rho < \beta$ . Then by recursively applying inequality (35), we have for  $k \geq K$ ,

$$\begin{aligned} &\mathbb{E} \left[ V(\tilde{\Delta}_k) \mathbf{1} \left\{ \mathcal{E}_{k/2}^{k-1}, \mathcal{K}_{k,t} \right\} \right] \\ &\leq \exp \left( -c \sum_{i=\lceil k/2 \rceil}^{k-1} \alpha_i \right) \mathbb{E} [V(\tilde{\Delta}_{\lceil k/2 \rceil}) \mathbf{1} \{\mathcal{K}_{\lceil k/2 \rceil, t}\}] + C(C_{\Delta,t}^2 + 1)t^2 \sum_{i=\lceil k/2 \rceil}^{k-1} \alpha_i^2 i^\rho \exp \left( -c \sum_{j=i+1}^{k-1} \alpha_j \right) \\ &\leq \exp \left( -c' k^{1-\beta} \right) \mathbb{E} [V(\tilde{\Delta}_{\lceil k/2 \rceil}) \mathbf{1} \{\mathcal{K}_{\lceil k/2 \rceil, t}\}] + C \sum_{i=\lceil k/2 \rceil}^{k-1} \alpha_i^2 i^\rho \exp \left( -c' \sum_{j=i+1}^{k-1} \alpha_j \right), \end{aligned} \quad (36)$$

where the second inequality follows because  $\sum_{i=l}^k \alpha_i \asymp k^{1-\beta} - l^{1-\beta}$ , and  $k^{1-\beta} - (k/2)^{1-\beta} \geq \frac{1-\beta}{2} k^{1-\beta}$ .

Now, by using the assumption that  $\mathbb{E}[\mathbf{1} \{\mathcal{K}_{k,t}\} V(\tilde{\Delta}_k)] \lesssim \mathbb{E}[\mathbf{1} \{\mathcal{K}_{k,t}\} \|\tilde{\Delta}_k\|^2] \leq C_{\Delta,t}^2$  for all  $k$ , we find that

$$\begin{aligned} \sum_{k=1}^n \mathbb{E} \left[ \|\tilde{\Delta}_k\|^{1+\gamma} \mathbf{1} \left\{ \mathcal{K}_{k,t}, \mathcal{E}_{k/2}^{k-1} \right\} \right] &\leq KC_{\Delta,t}^{1+\gamma} + \sum_{k \geq K}^n \mathbb{E} \left[ \|\tilde{\Delta}_k\|^2 \mathbf{1} \left\{ \mathcal{E}_{k/2}^{k-1}, \mathcal{K}_{k,t} \right\} \right]^{\frac{1+\gamma}{2}} \\ &\stackrel{(i)}{\leq} KC_{\Delta,t}^{1+\gamma} + \sum_{k \geq K}^n \left( \exp \left( -ck^{1-\beta} \right) C_{\Delta,t}^2 + C \sum_{i=k/2}^k \alpha_i^2 i^\rho \exp \left( -c(k^{1-\beta} - i^{1-\beta}) \right) \right)^{\frac{1+\gamma}{2}} \\ &\leq KC_{\Delta,t}^{1+\gamma} + \sum_{k \geq K} \exp \left( -\frac{c(1+\gamma)}{2} k^{1-\beta} \right) + C \sum_{k \geq K} \left( k^{\rho-2\beta} \sum_{i=k/2}^k \exp \left( -c(k^{1-\beta} - i^{1-\beta}) \right) \right)^{\frac{1+\gamma}{2}}, \end{aligned}$$

where step (i) follows from inequality (36). By Lemma 13, we know that

$$\sum_{i=k/2}^k \exp\left(-c(k^{1-\beta} - i^{1-\beta})\right) \lesssim \Gamma\left(\frac{1}{1-\beta}\right) + k^\beta,$$

so that

$$\sum_{k=1}^n \mathbb{E} \left[ \|\tilde{\Delta}_k\|^{1+\gamma} \mathbf{1} \left\{ \mathcal{E}_{k/2}^{k-1}, \mathcal{K}_{k,t} \right\} \right] \leq KC_\Delta^{1+\gamma} + C \sum_{k=K}^n \exp\left(-ck^{1-\beta}\right) + C \sum_{k=K}^n k^{(\rho-\beta)\frac{1+\gamma}{2}}.$$

Noting that  $\sum_{k=1}^n k^{(\rho-\beta)\frac{1+\gamma}{2}} \asymp n^{1-\frac{(\beta-\rho)(1+\gamma)}{2}}$ , we have

$$n^{-\frac{1}{2}} \sum_{k=1}^n \|\tilde{\Delta}_k\|^{1+\gamma} \mathbf{1} \left\{ \mathcal{E}_{k/2}^{k-1}, \mathcal{K}_{k,t} \right\} \xrightarrow{L_1} 0 \quad \text{if } (\beta - \rho)\frac{1+\gamma}{2} > \frac{1}{2}, \quad \text{or } \beta - \rho > \frac{1}{1+\gamma}. \quad (37)$$

Our initial choice of  $\rho$  satisfied this inequality, so we have that for any  $t \in \mathbb{R}$  the preceding convergence guarantee holds.

Now, let  $\delta > 0$  be arbitrary and using Lemma 4, choose  $t$  large enough that  $\mathbb{P}(\mathcal{K}_{\infty,t}) \leq \delta$ . Then

$$\begin{aligned} \mathbb{P} \left( n^{-\frac{1}{2}} \sum_{k=1}^n \|\tilde{\Delta}_k\|^{1+\gamma} \mathbf{1} \left\{ \mathcal{E}_{k/2}^{k-1} \right\} > \epsilon \right) &\leq \mathbb{P} \left( \mathcal{K}_{\infty,t}, n^{-\frac{1}{2}} \sum_{k=1}^n \|\tilde{\Delta}_k\|^{1+\gamma} \mathbf{1} \left\{ \mathcal{E}_{k/2}^{k-1} \right\} > \epsilon \right) + \mathbb{P}(\mathcal{K}_{\infty,t}^c) \\ &\leq \underbrace{\mathbb{P} \left( n^{-\frac{1}{2}} \sum_{k=1}^n \|\tilde{\Delta}_k\|^{1+\gamma} \mathbf{1} \left\{ \mathcal{E}_{k/2}^{k-1} \right\} \mathbf{1} \left\{ \mathcal{K}_{k,t} \right\} > \epsilon \right)}_{\rightarrow 0 \text{ as } n \rightarrow \infty} + \delta, \end{aligned}$$

the convergence to zero a consequence of inequality (37). As  $\delta > 0$  was arbitrary, we see that  $n^{-\frac{1}{2}} \sum_{k=1}^n \|\tilde{\Delta}_k\|^{1+\gamma} \mathbf{1} \left\{ \mathcal{E}_{k/2}^{k-1} \right\} \xrightarrow{P} 0$ , and expression (34) gives the lemma.

**Proof of inequality (35).** Recall the definition of the event  $\mathcal{E}_l^k = \{\|\tilde{\Delta}_i\| \leq \epsilon, \text{ all } i = \lceil l \rceil, \dots, k\}$  and that  $\mathbf{1} \left\{ \mathcal{K}_{k+1,t} \right\} \leq \mathbf{1} \left\{ \mathcal{K}_{k,t} \right\}$ . By inequality (25b) and the fact that  $\mathcal{E}_i^k \in \mathcal{F}_{k-1}$  for any  $i \leq k$  and  $\mathcal{K}_{k,t} \in \mathcal{F}_{k-1}$ , we have

$$\begin{aligned} \mathbb{E} \left[ V(\tilde{\Delta}_{k+1}) \mathbf{1} \left\{ \mathcal{E}_l^k, \mathcal{K}_{k+1,t} \right\} \right] &\leq \mathbb{E} \left[ \mathbb{E} \left[ V(\tilde{\Delta}_{k+1}) \mid \mathcal{F}_{k-1} \right] \mathbf{1} \left\{ \mathcal{E}_l^k, \mathcal{K}_{k,t} \right\} \right] \\ &\leq \mathbb{E} \left[ \left( V(\tilde{\Delta}_k) - \alpha_k \left\langle \nabla V(\tilde{\Delta}_k), R(\tilde{x}_k) \right\rangle + C\alpha_k^2 k^\rho V(\tilde{\Delta}_k) \right) \mathbf{1} \left\{ \mathcal{E}_l^k, \mathcal{K}_{k,t} \right\} \right] \\ &\quad + C\mathbb{E} \left[ \alpha_k^2 \mathbb{K}_k^2 \sum_{i=k-k^\rho}^{k-1} \|R(x_i) + \xi_i\|^2 \mathbf{1} \left\{ \mathcal{E}_l^{k-1}, \mathcal{K}_{k,t} \right\} + \alpha_k^2 (\|\Delta_k\|^2 + 1) \mathbf{1} \left\{ \mathcal{K}_{k,t}, \mathcal{E}_l^{k-1} \right\} \right], \end{aligned}$$

where we have used Assumption E that  $\xi_k = \xi_k(0) + \zeta(x_k)$ , and  $\mathbb{E}[\sup_k \mathbb{E}[\|\xi_k(0)\|^2 \mid \mathcal{F}_{k-1}]] < \infty$ .

We now use Lemma 12 to provide control of the preceding inequality. Under Assumption F coupled with  $\sup_k \mathbb{E}[\mathbf{1} \left\{ \mathcal{K}_{k,t} \right\} \|\Delta_k\|^2] \leq C_{\Delta,t}^2 < \infty$ , Lemma 12 implies that

$$\mathbb{E} \left[ \alpha_k^2 \mathbb{K}_k^2 \sum_{i=k-k^\rho}^{k-1} \|R(x_i) + \xi_i\|^2 \mathbf{1} \left\{ \mathcal{E}_l^{k-1}, \mathcal{K}_{k,t} \right\} \right] \lesssim \alpha_k^2 t^2 k^\rho (C_{\Delta,t}^2 + 1).$$

Noting that  $\|\tilde{\Delta}_k\|^2 \leq CV(\tilde{\Delta}_k)$  and  $\left\langle \nabla V(\tilde{\Delta}_k), R(\tilde{x}_k) \right\rangle \geq \lambda_0 V(\tilde{\Delta}_k)$  by Assumption F, we obtain inequality (35) as desired.

### A.3.2 Proof of Lemma 8

As in the proof of Lemma 7, we show that expression (34) holds. As before, we have  $\rho \in (\frac{1}{\tau-1}, \beta - \frac{1}{1+\gamma})$  as the power used in the definition of  $K_n$ . We begin by considering the progress made by a single step of the iteration with alternate error terms. We first claim that, similar to inequality (35), that there exist constants  $c, C$  such that for any  $l < k$  with  $l \in \mathbb{R}$ , we have

$$\mathbb{E} \left[ V(\tilde{\Delta}_{k+1}) \mathbf{1} \left\{ \mathcal{E}_l^k, \mathcal{K}_{k+1,t} \right\} \right] \leq (1 - c\alpha_k + C\alpha_k^2 k^\rho) \mathbb{E} \left[ V(\tilde{\Delta}_k) \mathbf{1} \left\{ \mathcal{E}_l^{k-1}, \mathcal{K}_{k,t} \right\} \right] + C\alpha_k^2 t^2 k^\rho. \quad (38)$$

Indeed, as in the proof of inequality (35), we use inequality (25a) to obtain

$$\begin{aligned} \mathbb{E} \left[ V(\tilde{\Delta}_{k+1}) \mathbf{1} \left\{ \mathcal{E}_l^k, \mathcal{K}_{k+1,t} \right\} \right] &\leq \mathbb{E} \left[ \mathbb{E} \left[ V(\tilde{\Delta}_{k+1}) \mid \mathcal{F}_{k-1} \right] \mathbf{1} \left\{ \mathcal{E}_l^k, \mathcal{K}_{k,t} \right\} \right] \\ &\leq \mathbb{E} \left[ \left( V(\tilde{\Delta}_k) - \alpha_k \left\langle \nabla V(\tilde{\Delta}_k), R(\tilde{x}_k) \right\rangle + C\alpha_k^2 k^\rho V(\tilde{\Delta}_k) \right) \mathbf{1} \left\{ \mathcal{E}_l^k, \mathcal{K}_{k,t} \right\} \right] \\ &\quad + C\mathbb{E} \left[ \alpha_k^2 K_k^2 \sum_{i=k-k^\rho}^{k-1} \|R(x_i) + \xi_i\|^2 \mathbf{1} \left\{ \mathcal{E}_l^{k-1}, \mathcal{K}_{k,t} \right\} + \alpha_k^2 (\|R(x_k)\|^2 + 1) \mathbf{1} \left\{ \mathcal{K}_{k,t}, \mathcal{E}_l^{k-1} \right\} \right], \end{aligned}$$

where we have used Assumption F' that  $\mathbb{E}[\|\xi_k\|^2 \mid \mathcal{F}_{k-1}] \lesssim 1$  for all  $k$ . Using our assumption that on the event  $\mathcal{E}_l^k$  we have  $\left\langle \nabla V(\tilde{\Delta}_k), R(\tilde{x}_k) \right\rangle \geq \lambda_0 V(\tilde{\Delta}_k)$  and that  $\|R(x)\| \lesssim 1$  for all  $x$  (Assumption F'), we obtain the desired inequality (38).

Again paralleling the proof of Lemma 7, let  $K < \infty$  be large enough that for the constants  $c, C$  in inequality (38), there is a constant  $c'$  such that for  $k \geq K$  and stepsizes  $\alpha_k = \alpha k^{-\beta}$ , we have  $(1 - c\alpha_i + C\alpha_i^2 i^\rho) \leq \exp(-c'\alpha_i)$  for  $i \geq k/2$ . Then by recursively applying inequality (38), we have for  $k \geq K$ ,

$$\begin{aligned} &\mathbb{E} \left[ V(\tilde{\Delta}_k) \mathbf{1} \left\{ \mathcal{E}_{k/2}^{k-1}, \mathcal{K}_{k,t} \right\} \right] \\ &\leq \exp(-c'k^{1-\beta}) \mathbb{E} \left[ V(\tilde{\Delta}_{\lceil k/2 \rceil}) \mathbf{1} \left\{ \mathcal{K}_{\lceil k/2 \rceil, t} \right\} \right] + C \sum_{i=\lceil k/2 \rceil}^{k-1} \alpha_i^2 i^\rho \exp\left(-c' \sum_{j=i+1}^{k-1} \alpha_j\right), \quad (39) \end{aligned}$$

exactly as in the derivation of inequality (36).

The remainder of the proof is completely identical to that of Lemma 7.

### A.4 Proof of Lemma 11

Fix  $n \in \mathbb{N}$  and recall our assumption (16) that  $\rho \in (\frac{1}{\tau-1}, \beta - \frac{1}{2})$ , where  $\rho$  is used in the definition of the ratio  $K_n$  and event  $\mathcal{K}_{n,t}$  (Defs. (17) and (18)), and consider  $\mathbb{E}[V(\tilde{\Delta}_{n+1}) \mathbf{1} \left\{ \mathcal{K}_{n+1,t} \right\}]$ . Combining inequality (25b), the fact that  $\mathbf{1} \left\{ \mathcal{K}_{n,t} \right\}$  is non-increasing (because  $\mathcal{K}_{n+1,t} \subset \mathcal{K}_{n,t}$ ), and  $\mathcal{K}_{n,t} \in \mathcal{F}_{n-1}$ , we see that defining

$$C_{\Delta,n,t}^2 = \max_{k \leq n} \mathbb{E}[\|\Delta_k\|^2 \mathbf{1} \left\{ \mathcal{K}_{k,t} \right\}],$$

we have

$$\begin{aligned} \mathbb{E} \left[ V(\tilde{\Delta}_{n+1}) \mathbf{1} \left\{ \mathcal{K}_{n+1,t} \right\} \right] &\leq \mathbb{E} \left[ V(\tilde{\Delta}_{n+1}) \mathbf{1} \left\{ \mathcal{K}_{n,t} \right\} \right] \\ &\leq \mathbb{E} \left[ \left( V(\tilde{\Delta}_n) - \alpha_n \left\langle \nabla V(\tilde{\Delta}_n), R(\tilde{x}_n) \right\rangle + C\alpha_n^2 n^\rho \|\nabla V(\tilde{\Delta}_n)\|^2 \right) \mathbf{1} \left\{ \mathcal{K}_{n,t} \right\} \right] + Ct^2 \alpha_n^2 n^\rho (C_{\Delta,n,t}^2 + 1). \end{aligned}$$

For the final inequality we have used inequality (29), that is,  $\mathbb{E}[\|g_k\|^2 \mathbf{1}\{\mathcal{K}_{k,t}\}] \leq c(C_{\Delta,n,t}^2 + 1)$ . By using that  $\|\nabla V(x - x^*)\|^2 \leq L^2 \|x - x^*\|^2 \leq \frac{L^2}{\lambda^2} V(x - x^*)$  by our assumptions on  $V$ , we thus obtain

$$\begin{aligned} & \mathbb{E} \left[ V(\tilde{\Delta}_{n+1}) \mathbf{1}\{\mathcal{K}_{n+1,t}\} \right] \\ & \leq \mathbb{E} \left[ \left( V(\tilde{\Delta}_n) - \alpha_n \langle \nabla V(\tilde{\Delta}_n), R(\tilde{x}_n) \rangle + C\alpha_n^2 n^\rho V(\tilde{\Delta}_n) \right) \mathbf{1}\{\mathcal{K}_{n,t}\} \right] + Ct^2 \alpha_n^2 n^\rho (C_{\Delta,n,t}^2 + 1). \end{aligned}$$

Now, noting that  $\langle \nabla V(\tilde{\Delta}_n), R(\tilde{x}_n) \rangle \geq \lambda_0 V(\tilde{\Delta}_n)$  by Assumption F, we have

$$\mathbb{E} \left[ V(\tilde{\Delta}_{n+1}) \mathbf{1}\{\mathcal{K}_{n+1,t}\} \right] \leq (1 - \lambda_0 \alpha_n + C\alpha_n^2 n^\rho) \mathbb{E} \left[ V(\tilde{\Delta}_n) \mathbf{1}\{\mathcal{K}_{n,t}\} \right] + Ct^2 \alpha_n^2 n^\rho (C_{\Delta,n,t}^2 + 1)$$

By recursively applying this inequality, we have

$$\begin{aligned} & \mathbb{E} \left[ V(\tilde{\Delta}_{n+1}) \mathbf{1}\{\mathcal{K}_{n+1,t}\} \right] \\ & \leq \prod_{k=1}^n (1 - \lambda_0 \alpha_k + C\alpha_k^2 k^\rho) \mathbb{E}[V(\Delta_1)] + Ct^2 (C_{\Delta,n,t}^2 + 1) \sum_{k=1}^n \alpha_k^2 k^\rho \prod_{l=k+1}^n (1 - \lambda_0 \alpha_l + C\alpha_l^2 l^\rho). \end{aligned} \quad (40)$$

We state a technical lemma that controls the products above. Let

$$b_l^k := \prod_{i=l}^k (1 - \lambda_0 \alpha_i + C\alpha_i^2 i^\rho).$$

**Lemma 16.** *Let the scalar sequence  $b_l^k$  be defined as above with  $l \leq k$ ,  $\beta > \rho$ , and  $\alpha_i = \alpha i^{-\beta}$  where  $C \geq 4\lambda_0^2$  (so that each term in the product is non-negative). There exist constants  $c_0, c_1, c_2$  (dependent on  $\beta, \alpha, \lambda_0$ , and  $C$ ) such that*

$$b_l^k \leq c_0 \exp \left( -c_1 \sum_{i=l}^k \alpha_i \right) \leq c_0 \exp \left( -c_2 (k^{1-\beta} - l^{1-\beta}) \right). \quad (41)$$

**Proof** This result is similar to a result of Polyak and Juditsky [26, proof of Lemma 1, Part 3], but with some differences for additional powers in the sequence. As  $\alpha_i = \alpha i^{-\beta}$  and we have assumed that  $\beta > \rho$ , there exists some  $K \in \mathbb{N}$  such that for  $k \geq K$  we have  $2C\alpha_k^2 k^\rho \leq \lambda_0 \alpha_k$ , or  $\alpha k^{\rho-\beta} \leq \lambda_0 / (2C)$ . For any  $k \geq K$ , we have  $(1 - \lambda_0 \alpha_k + C\alpha_k^2) \leq 1 - \lambda_0 \alpha_k / 2 \leq \exp(-\frac{\lambda_0}{2} \alpha_k)$ . We find that

$$\begin{aligned} b_l^k &= \prod_{i=l}^k (1 - \lambda_0 \alpha_i + C\alpha_i^2 i^\rho) = \prod_{i \geq l \wedge K}^k (1 - \lambda_0 \alpha_i + C\alpha_i^2 i^\rho) \prod_{i=l}^{(l \wedge K)-1} (1 - \lambda_0 \alpha_i + C\alpha_i^2 i^\rho) \\ &\leq \prod_{i=1}^K \max\{1, 1 - \lambda_0 \alpha_i + C\alpha_i^2 i^\rho\} \prod_{i \geq l \wedge K}^k \exp \left( -\frac{\lambda_0}{2} \alpha_i \right) \\ &\leq \left[ \exp \left( \frac{\lambda_0}{2} \sum_{i=1}^K \alpha_i \right) \prod_{i=1}^K \max\{1, 1 - \lambda_0 \alpha_i + C\alpha_i^2 i^\rho\} \right] \exp \left( -\frac{\lambda_0}{2} \sum_{i=l}^k \alpha_i \right). \end{aligned}$$

The term in the braces  $[\cdot]$  in the preceding product is the constant  $c_0$ , giving the first inequality of expression (41). For the second, note that

$$\sum_{i=l}^k \alpha_i = \alpha \sum_{i=l}^k i^{-\beta} \geq \alpha \int_l^k t^{-\beta} dt = \frac{\alpha}{1-\beta} \left[ k^{1-\beta} - l^{1-\beta} \right],$$

which completes the proof.  $\square$

Applying Lemma 16 in inequality (40), we obtain for constants  $c, C$  independent of  $C_{\Delta, n, t}$  that

$$\mathbb{E} \left[ V(\tilde{\Delta}_{n+1}) \mathbf{1} \{ \mathcal{K}_{n+1, t} \} \right] \leq C \exp(-cn^{1-\beta}) \mathbb{E}[V(\Delta_1)] + Ct^2 (C_{\Delta, n, t}^2 + 1) \sum_{k=1}^n \alpha_k^2 k^\rho \exp \left( -c(n^{1-\beta} - k^{1-\beta}) \right).$$

Now we use the technical Lemma 14, which shows that the final sum tends to zero when  $\alpha_k = \alpha k^{-\beta}$ . In particular, for any  $\epsilon > 0$ , there exists some  $N(\epsilon, t) < \infty$ , independent of  $C_{\Delta, n, t}$ , such that  $n \geq N(\epsilon, t)$  implies

$$C \exp(-cn^{1-\beta}) < \epsilon^2 \quad \text{and} \quad Ct^2 \sum_{k=1}^n \alpha_k^2 k^{\rho-2\beta} \exp \left( -c(n^{1-\beta} - k^{1-\beta}) \right) \leq \epsilon^2.$$

That is, we have

$$\mathbb{E} \left[ V(\tilde{\Delta}_{n+1}) \mathbf{1} \{ \mathcal{K}_{n+1, t} \} \right] \leq \epsilon^2 (\mathbb{E}[V(\Delta_1)] + C_{\Delta, n, t}^2 + 1).$$

As  $\epsilon > 0$  was arbitrary and there are constants  $c, C$  such that  $c \|x - x^*\|^2 \leq V(x - x^*) \leq C \|x - x^*\|^2$  this gives Lemma 11.

## A.5 Proof of Lemma 13

We prove the result via a change of variables. Let  $u = c(t^\kappa - a^\kappa)$ , so that

$$t = (u/c + a^\kappa)^{\frac{1}{\kappa}}, \quad du = \kappa c t^{\kappa-1} dt = \kappa c (u/c + a^\kappa)^{\frac{\kappa-1}{\kappa}} dt, \quad \text{or} \quad dt = (\kappa c)^{-1} (u/c + a^\kappa)^{\frac{1-\kappa}{\kappa}} du.$$

That is, by our change of variables, we have

$$\begin{aligned} \int_a^b \exp(-c(t^\kappa - a^\kappa)) dt &= \frac{1}{\kappa c} \int_0^{c(b^\kappa - a^\kappa)} \left( \frac{u}{c} + a^\kappa \right)^{\frac{1-\kappa}{\kappa}} e^{-u} du \\ &\leq \frac{\max\{2^{\frac{1-\kappa}{\kappa}-1}, 1\}}{\kappa c} \left[ \int_0^{c(b^\kappa - a^\kappa)} \left( \frac{u}{c} \right)^{\frac{1-\kappa}{\kappa}} e^{-u} du + \int_0^{c(b^\kappa - a^\kappa)} a^{1-\kappa} e^{-u} du \right], \end{aligned}$$

where the final inequality follows by convexity of  $t \mapsto t^{\frac{1-\kappa}{\kappa}}$ , for  $\kappa < \frac{1}{2}$  and the fact that  $(t_1 + t_2)^{\frac{1-\kappa}{\kappa}} \leq t_1^{\frac{1-\kappa}{\kappa}} + t_2^{\frac{1-\kappa}{\kappa}}$  for  $\kappa \geq \frac{1}{2}$  (or  $\frac{1-\kappa}{\kappa} \leq 1$ ). Noting that  $\int_0^\infty u^{\frac{1-\kappa}{\kappa}} e^{-u} du = \Gamma(\frac{1}{\kappa})$  and  $\int_0^\infty e^{-u} du = 1$ , we obtain our desired result.

## A.6 Proof of Lemma 14

The quantity in the summation diverges or converges identically to the integral

$$\int_1^n u^{\rho-2\beta} \exp\left(-c(n^{1-\beta} - u^{1-\beta})\right) du \leq \int_1^{an} \exp\left(-c(n^{1-\beta} - u^{1-\beta})\right) dt + \int_{an}^n u^{\rho-2\beta} dt \quad (42)$$

for any  $a \in [0, 1]$ . Now, by concavity of  $u \mapsto u^{1-\beta}$  for  $\beta \in (\frac{1}{2}, 1)$ , we have  $u^{1-\beta} \leq n^{1-\beta} + (1-\beta)n^{-\beta}(u-n)$ , or  $n^{1-\beta} - u^{1-\beta} \geq (1-\beta)(n-u)n^{-\beta}$ . In particular, the first integral on the right side of the display (42) has bound

$$\begin{aligned} \int_1^{an} \exp\left(-c(n^{1-\beta} - u^{1-\beta})\right) du &\leq \int_1^{an} \exp\left(-\frac{c}{n^\beta}(n-u)\right) du = -\int_{\frac{c}{n^\beta}(n-1)}^{\frac{c}{n^\beta}(1-a)n} \frac{n^\beta}{c} \exp(-u) du \\ &\leq \frac{n^\beta}{c} \int_{c(1-a)n^{1-\beta}}^{cn^{1-\beta}} e^{-u} du = \frac{n^\beta}{c} \left[ \exp\left(-c(1-a)n^{1-\beta}\right) - \exp\left(-cn^{1-\beta}\right) \right] \\ &\leq \frac{n^\beta}{c} \exp\left(-c(1-a)n^{1-\beta}\right), \end{aligned} \quad (43)$$

where we made a change of variables. For the second integral, we have

$$\int_{an}^n u^{\rho-2\beta} du = \frac{1}{1-2\beta} (n^{1+\rho-2\beta} - (an)^{1+\rho-2\beta}),$$

and combining this with (43) in the bound (42), we obtain for any  $a \in (0, 1)$  that for constants  $C, c$ ,

$$\sum_{k=1}^n k^{\rho-2\beta} \exp\left(-c(n^{1-\beta} - k^{1-\beta})\right) \leq C \left[ n^\beta \exp\left(-c(1-a)n^{1-\beta}\right) + \frac{n^{1+\rho-2\beta}}{2\beta-1} \left(1 - a^{1+\rho-2\beta}\right) \right].$$

By our assumption that  $\rho < 1$  and  $a \in (0, 1)$ , the first term above converges to zero; our assumption that  $\rho < \beta - \frac{1}{2}$  guarantees that the second does as well.

## References

- [1] A. Agarwal and J. C. Duchi. Distributed delayed stochastic optimization. In *Advances in Neural Information Processing Systems 24*, 2011.
- [2] A. Agarwal, P. L. Bartlett, P. Ravikumar, and M. J. Wainwright. Information-theoretic lower bounds on the oracle complexity of convex optimization. *IEEE Transactions on Information Theory*, 58(5):3235–3249, 2012.
- [3] P. Baldi, P. Sadowski, and D. Whiteson. Searching for exotic particles in high-energy physics with deep learning. *Nature Communications*, 5, July 2014.
- [4] G. Ballard, J. Demmel, O. Holtz, and O. Schwartz. Minimizing communication in numerical linear algebra. *SIAM Journal on Matrix Analysis and Applications*, 32(3):866–901, 2011.
- [5] D. P. Bertsekas and J. N. Tsitsiklis. *Parallel and Distributed Computation: Numerical Methods*. Prentice-Hall, Inc., 1989.

- [6] J. Dean, G. S. Corrado, R. Monga, K. Chen, M. Devin, Q. V. Le, M. Z. Mao, M. Ranzato, A. Senior, P. Tucker, K. Yang, and A. Y. Ng. Large scale distributed deep networks. In *Advances in Neural Information Processing Systems 25*, 2012.
- [7] O. Dekel, R. Gilad-Bachrach, O. Shamir, and L. Xiao. Optimal distributed online prediction using mini-batches. *Journal of Machine Learning Research*, 13:165–202, 2012.
- [8] J. C. Duchi, E. Hazan, and Y. Singer. Adaptive subgradient methods for online learning and stochastic optimization. *Journal of Machine Learning Research*, 12:2121–2159, 2011.
- [9] J. C. Duchi, A. Agarwal, and M. J. Wainwright. Dual averaging for distributed optimization: convergence analysis and network scaling. *IEEE Transactions on Automatic Control*, 57(3): 592–606, 2012.
- [10] J. C. Duchi, M. I. Jordan, and H. B. McMahan. Estimation, optimization, and parallelism when data is sparse. In *Advances in Neural Information Processing Systems 26*, 2013.
- [11] J. C. Duchi, S. Chaturapruek, and C. Ré. Asynchronous stochastic convex optimization, 2015. URL <https://www.codalab.org/worksheets/0x610bcdb722bf48d3b537a65edf0fe72d/>. Code for reproducing experiments.
- [12] Y. M. Ermoliev. On the stochastic quasi-gradient method and stochastic quasi-Feyer sequences. *Kibernetika*, 2:72–83, 1969.
- [13] S. Ghadimi and G. Lan. Optimal stochastic approximation algorithms for strongly convex stochastic composite optimization, I: a generic algorithmic framework. *SIAM Journal on Optimization*, 22(4):1469–1492, 2012.
- [14] E. Hazan and S. Kale. An optimal algorithm for stochastic strongly convex optimization. In *Proceedings of the Twenty Fourth Annual Conference on Computational Learning Theory*, 2011.
- [15] J. Hiriart-Urruty and C. Lemaréchal. *Convex Analysis and Minimization Algorithms I*. Springer, New York, 1993.
- [16] A. Juditsky, A. Nemirovski, and C. Tauvel. Solving variational inequalities with the stochastic mirror-prox algorithm. *Stochastic Systems*, 1(1):17–58, 2011.
- [17] L. Le Cam and G. L. Yang. *Asymptotics in Statistics: Some Basic Concepts*. Springer, 2000.
- [18] E. L. Lehmann and G. Casella. *Theory of Point Estimation, Second Edition*. Springer, 1998.
- [19] D. Lewis, Y. Yang, T. Rose, and F. Li. RCV1: A new benchmark collection for text categorization research. *Journal of Machine Learning Research*, 5:361–397, 2004.
- [20] M. Lichman. UCI machine learning repository, 2013. URL <http://archive.ics.uci.edu/ml>.
- [21] J. Liu, S. J. Wright, C. Ré, V. Bittorf, and S. Sridhar. An asynchronous parallel stochastic coordinate descent algorithm. In *Proceedings of the 31st International Conference on Machine Learning*, 2014.

- [22] A. Nemirovski and D. Yudin. *Problem Complexity and Method Efficiency in Optimization*. Wiley, 1983.
- [23] A. Nemirovski, A. Juditsky, G. Lan, and A. Shapiro. Robust stochastic approximation approach to stochastic programming. *SIAM Journal on Optimization*, 19(4):1574–1609, 2009.
- [24] Y. Nesterov. Primal-dual subgradient methods for convex problems. *Mathematical Programming*, 120(1):261–283, 2009.
- [25] F. Niu, B. Recht, C. Re, and S. Wright. Hogwild: a lock-free approach to parallelizing stochastic gradient descent. In *Advances in Neural Information Processing Systems 24*, 2011.
- [26] B. T. Polyak and A. B. Juditsky. Acceleration of stochastic approximation by averaging. *SIAM Journal on Control and Optimization*, 30(4):838–855, 1992.
- [27] B. Recht and C. Ré. Beneath the valley of the noncommutative arithmetic-geometric mean inequality: conjectures, case-studies, and consequences. In *Proceedings of the Twenty Fifth Annual Conference on Computational Learning Theory*, 2012.
- [28] P. Richtárik and M. Takáč. Parallel coordinate descent methods for big data optimization. *Mathematical Programming*, page Online first, 2015. URL <http://link.springer.com/article/10.1007/s10107-015-0901-6>.
- [29] H. Robbins and S. Monro. A stochastic approximation method. *Annals of Mathematical Statistics*, 22:400–407, 1951.
- [30] H. Robbins and D. Siegmund. A convergence theorem for non-negative almost supermartingales and some applications. In *Optimizing Methods in Statistics*, pages 233–257. Academic Press, New York, 1971.
- [31] A. W. van der Vaart. *Asymptotic Statistics*. Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press, 1998. ISBN 0-521-49603-9.