

Certain Family of Some Beta Distributions Arising from Distribution of Randomly Weighted Average

Rasool Roozegar*

Department of Statistics, Yazd University, P.O. Box 89195-741, Yazd, Iran

Abstract

We give the exact distribution of the average of n independent beta random variables weighted by the selected cuts of $(0, 1)$ by the order statistics of a random sample of size $n - 1$ from the uniform distribution $U(0, 1)$, for each n . A new integral transformation that is similar to generalized Stieltjes transform is given with various properties. The result of Soltani and Roozegar [On distribution of randomly ordered uniform incremental weighted averages: Divided difference approach. *Statist Probab Lett.* 2012;82(5):10121020] with this new transform and also integral representation of the Gauss-hypergeometric function in some parts are employed to achieve the exact distribution. Several examples of the new family are investigated.

Keywords and phrases: Selected order statistics; Beta distribution. Additive Stieltjes transform; Gauss-hypergeometric function; Randomly weighted average, exact distribution.

2010 AMS Subject Classification: 60E05, 46F12, 62E15.

*Corresponding: rroozegar@yazd.ac.ir

1 Introduction

Let $U_{(1)} < U_{(2)} < \dots < U_{(n-1)}$ be order statistics based on a random sample of size $n-1$ from the uniform distribution $U(0, 1)$, $U_{(0)} = 0$ and $U_{(n)} = 1$. Randomly weighted average (RWA) of independent and continuous random variables X_1, \dots, X_n with respective distribution functions F_1, \dots, F_n , is defined by

$$S_n = R_1 X_1 + R_2 X_2 + \dots + R_n X_n, \quad n \geq 2, \quad (1.1)$$

where the proportions $R_i = U_{(i)} - U_{(i-1)}$, $i = 1, \dots, n-1$ and $R_n = 1 - \sum_{i=1}^{n-1} R_i$ are random weights. Suppose that among the above whole order statistics, we select $k-1$ order statistics $U_{(n_1)} < U_{(n_2)} < \dots < U_{(n_{k-1})}$, where $k = k(n)$, $2 \leq k \leq n$ and $n_0 = 0 < n_1 < n_2 < \dots < n_{k-1} < n_k = n$. A general form to S_n will be the RWA of k independent and continuous random variables X_1, \dots, X_k , denoted by $S_{n:n_1, \dots, n_{k-1}}$, which is given by

$$S_{n:n_1, \dots, n_{k-1}} = \sum_{j=1}^k V_j X_j, \quad (1.2)$$

where the random weights V_j are defined by

$$V_j = U_{(n_j)} - U_{(n_{j-1})}, \quad j = 1, 2, \dots, k.$$

In the above we have put the conventions $U_{(n_0)} = 0$, $U_{(n_k)} = 1$ and $r_j = n_j - n_{j-1}$, $j = 1, 2, \dots, k$. Then $U_{(n_j)} = \sum_{i=1}^j V_i$ and $V_j = \sum_{i=n_{j-1}+1}^{n_j} R_i$. The RWA on the form (1.2) was defined and studied by [7]. In fact the random vector $\mathbf{V} = (V_1, V_2, \dots, V_k)$ has the Dirichlet distribution, $Dir(r_1, r_2, \dots, r_k)$, with the probability density function (pdf)

$$f_{\mathbf{V}}(v_1, \dots, v_k) = \frac{\Gamma(n)}{\prod_{j=1}^k \Gamma(r_j)} \prod_{j=1}^{k-1} v_j^{r_j} (1 - \sum_{j=1}^{k-1} v_j)^{r_k},$$

at any point in the canonical simplex $\{(v_1, \dots, v_k) | v_i \geq 0, i = 1, 2, \dots, k, \sum_{j=1}^k v_j = 1\}$ in the $(k-1)$ -dimensional real space \mathbb{R}^{k-1} and zero outside.

To state assertions, we introduce beta distributions, $\tilde{beta}(p, q)$, over $[a, b]$ by the pdf

$$f_{p,q}(x) = \begin{cases} \frac{1}{B(p, q)(b-a)^{p+q-1}} (x-a)^{p-1} (b-x)^{q-1} & \text{if } a < x < b, \\ 0 & \text{otherwise,} \end{cases}$$

for $p, q > 0$ and $B(p, q) = \frac{\Gamma(p)\Gamma(q)}{\Gamma(p+q)}$ is beta function.

Let us denote the usual beta distribution over $[0, 1]$ by $beta(p, q)$ with above pdf when $a = 0$ and $b = 1$. The power semicircle distribution with parameters θ and σ on $(-\sigma, \sigma)$, denoted by $PS(\theta, \sigma)$, is a special case of beta distribution whenever $p = q = \theta + 3/2$, $b = \sigma$ and $a = -\sigma$. We recall that $PS(-1, \sigma)$ is arcsine distribution on $(-\sigma, \sigma)$, for $\theta = -1/2$ the power semicircle density is the uniform on $(-\sigma, \sigma)$ and for $\theta = 0$, it is the semicircle (Wigner) density on $(-\sigma, \sigma)$.

The ordinary Stieltjes transform (ST) and generalized Stieltjes transform (GST) of a distribution H are respectively defined by

$$\mathcal{S}[H](z) = \int_{\mathbb{R}} \frac{1}{z-x} dH(x), \quad z \in \mathbb{C} \cap (\text{supp}H)^c,$$

and

$$\mathcal{S}[H; \rho](z) = \int_{\mathbb{R}} \frac{1}{(z-x)^\rho} dH(x), \quad z \in \mathbb{C} \cap (\text{supp}H)^c, \quad \rho > 0,$$

where \mathbb{C} is the set of complex numbers, $\text{supp}H$ stands for the support of H and ρ is a constant. For more on the ST and GST, see [3].

Soltani and Roozegar [7] effectively apply certain results in divided differences and derive the useful relation between GST of $F_{S_{n:n_1, \dots, n_{k-1}}}$ and GSTs of F_1, \dots, F_n ; more is given there:

$$\mathcal{S}[F_{S_{n:n_1, \dots, n_{k-1}}}; n](z) = \prod_{i=1}^k \mathcal{S}[F_i; r_i](z), \quad z \in \mathbb{C} \bigcap_{i=1}^k (\text{supp} F_i)^c. \quad (1.3)$$

In particular

$$\mathcal{S}[F_{S_n}; n](z) = \prod_{i=1}^n \mathcal{S}[F_i](z), \quad z \in \mathbb{C} \bigcap_{i=1}^n (\text{supp} F_i)^c.$$

The Gauss-hypergeometric function $F_D^{(1)}$, is defined by the series

$$F_D^{(1)}(c, a; b; z) = \sum_{n=0}^{\infty} \frac{(c)_n (a)_n}{(b)_n n!} z^n,$$

where $(a)_0 = 1$ and $(a)_n = a(a+1)(a+2) \cdots (a+n-1)$, $n \geq 1$, denotes the rising factorial.

Gauss-hypergeometric function $F_D^{(1)}$ has the Euler's integral representation of the form

$$F_D^{(1)}(c, a; b; z) = \frac{\Gamma(b)}{\Gamma(a)\Gamma(b-a)} \int_0^1 \frac{t^{a-1}(1-t)^{b-a-1}}{(1-zt)^c} dt. \quad (1.4)$$

For more details on Gauss-hypergeometric function and its properties, see [1] and [2].

There are few examples of known distributions that are distributions of RWA on the form (1.1), S_n , and there is not any examples, to the best of our knowledge, that are distributions of RWA on the form (1.2), $S_{n:n_1, \dots, n_{k-1}}$. Roozegar and Soltani [6] used the investigations of [4] and [5] on the connection between ST and GST of two distributions and introduce new classes of power semicircle laws that are RWA S_n distributions. In this paper we introduce new classes of RWA $S_{n:n_1, \dots, n_{k-1}}$ distributions.

Rest of the paper is organized as follows. In Section 2 we introduce a new transform similar to GST for later analysis. We rewrite the main result of [7] based on this new transform as a Theorem 2.2 in this section. Two new classes of RWA $S_{n:n_1, \dots, n_{k-1}}$ distributions are investigated in Section 3. In Section 4, we present some examples of the new classes.

2 New transform and RWA distribution

First, we define a new univariate characteristic function called an additive Stieltjes transform (AST).

Definition 2.1 *If X is a random variable with distribution H on a subset S of $A = [-a, a]$, $a > 0$, its AST is defined as*

$$\begin{aligned} \mathcal{AS}[H; d](z) &= E_X \left[\frac{1}{(1 - zX)^d} \right] \\ &= \int_S \frac{1}{(1 - zx)^d} dH(x), \quad |z| < \frac{1}{a}, \end{aligned}$$

where d is a positive real number.

The assumptions that d is positive and H has a support in S are needed for the one to one correspondence between AST and H in next theorem.

We have the following relationship between the AST and the GST of a distribution H ,

$$\mathcal{AS}[H; d](z) = \frac{1}{z^d} \mathcal{S}[H; d] \left(\frac{1}{z} \right). \quad (2.1)$$

Theorem 2.1 For distributions H_1 and H_2 in a subset S of $A = [-a, a]$ and any positive real number d , if we have

$$\mathcal{AS}[H_1; d](z) = \mathcal{AS}[H_2; d](z), \quad (2.2)$$

for all $|z| < \frac{1}{a}$, then $H_1 = H_2$.

Proof Since d is a real number and $|zx| < 1$, we have

$$(1 - zx)^{-d} = \sum_{m=0}^{\infty} \frac{\Gamma(d+m)}{\Gamma(d)m!} (zx)^m. \quad (2.3)$$

By Definition 2.1 and equations (2.2) and (2.3), we have

$$\sum_{m=0}^{\infty} \frac{\Gamma(d+m)}{\Gamma(d)m!} z^m \int_S x^m dH_1(x) = \sum_{m=0}^{\infty} \frac{\Gamma(d+m)}{\Gamma(d)m!} z^m \int_S x^m dH_2(x), \quad (2.4)$$

for all $|z| < \frac{1}{a}$. Given an integer k , treating z as variable and equating the corresponding coefficients of z^m ($\frac{\Gamma(d+m)}{\Gamma(d)} \neq 0$, for all m , as d is positive) for each m in the two sums, we obtain

$$\int_S x^m dH_1(x) = \int_S x^m dH_2(x).$$

Hence we see that

$$\int_S P(x) dH_1(x) = \int_S P(x) dH_2(x),$$

where $P(x)$ is any polynomial function of x , and similarly for any continuous function. Thus $H_1 = H_2$. \square

The following theorem enables us to represent a relationship between the AST of the distribution of RWA $S_{n:n_1, \dots, n_{k-1}}$ and to those of X_1, \dots, X_k . This theorem is the main theorem of [7] based on the AST.

Theorem 2.2 Let $S_{n:n_1, \dots, n_{k-1}}$ be the RWA given in (1.2). Assume random variables X_1, \dots, X_k are independent and continuous with distribution functions F_1, \dots, F_k , respectively. Then

$$\mathcal{AS}[F_{S_{n:n_1, \dots, n_{k-1}}}; \sum_{i=1}^k r_i](z) = \prod_{i=1}^k \mathcal{AS}[F_i; r_i](z),$$

for all $|z| < \frac{1}{a}$.

Proof Applying Theorem 3.1 of [7] and then (2.1) to conclude the result. \square

3 Families of RWA on some beta distributions

In this section, we provide some families of RWA $S_{n:n_1, \dots, n_{k-1}}$ distributions.

Theorem 3.1 *Let X_i , $i = 1, 2, \dots, k$ be independent random variables with $\tilde{\text{beta}}(r_i + 1/2, r_i + 1/2)$ distributions. Then the RWA $S_{n:n_1, \dots, n_{k-1}}$ has $\tilde{\text{beta}}(\sum_{i=1}^k r_i + 1/2, \sum_{i=1}^k r_i + 1/2)$ distribution.*

Proof By definition of AST of beta distribution on $[a, b]$, we have

$$\mathcal{AS}[F; r_i](z) = \int_a^b \frac{1}{(1-zx)^{r_i}} \frac{(x-a)^{r_i-1/2}(b-x)^{r_i-1/2}}{B(r_i+1/2, r_i+1/2)(b-a)^{2r_i}} dx.$$

By the change of variable $t = \frac{2x - (b+a)}{b-a}$, we obtain that

$$\mathcal{AS}[F; r_i](z) = \frac{1}{[1 - (a+b)z/2]^{r_i}} \int_{-1}^1 \frac{1}{\left(1 - \frac{z(b-a)}{2 - (a+b)z} t\right)^{r_i}} \frac{(t+1)^{r_i-1/2}(1-t)^{r_i-1/2}}{2^{2r_i} B(r_i+1/2, r_i+1/2)} dt.$$

By Table 3 in [5], it follows that

$$\begin{aligned} \mathcal{AS}[F; r_i](z) &= \frac{1}{[1 - (a+b)z/2]^{r_i}} \left[\frac{2}{1 + \sqrt{1 - \frac{z^2(b-a)^2}{(2 - (a+b)z)^2}}} \right]^{r_i} \\ &= \left[\frac{4}{2 - (a+b)z + 2\sqrt{1 - (a+b)z + abz^2}} \right]^{r_i}. \end{aligned}$$

Using Theorem 2.2 we obtain

$$\mathcal{AS}[F_{S_{n:n_1, \dots, n_{k-1}}}; \sum_{i=1}^k r_i](z) = \left[\frac{4}{2 - (a+b)z + 2\sqrt{1 - (a+b)z + abz^2}} \right]^{\sum_{i=1}^k r_i},$$

which is the AST of $\tilde{\text{beta}}(\sum_{i=1}^k r_i + 1/2, \sum_{i=1}^k r_i + 1/2)$ distribution. Therefore $S_{n:n_1, \dots, n_{k-1}}$ has $\tilde{\text{beta}}(\sum_{i=1}^k r_i + 1/2, \sum_{i=1}^k r_i + 1/2)$ distribution. \square

Another general family of RWA on beta distributions on $[a, b]$ is presented in the following theorem.

Theorem 3.2 Let X_i , $i = 1, 2, \dots, k$ be independent random variables with $\tilde{\text{beta}}(s_i, r_i - s_i)$ distributions. Then the RWA $S_{n:n_1, \dots, n_{k-1}}$ has $\tilde{\text{beta}}(\sum_{i=1}^k s_i, \sum_{i=1}^k r_i - \sum_{i=1}^k s_i)$ distribution.

Proof By using AST of $\tilde{\text{beta}}(s_i, r_i - s_i)$ distribution, we have

$$\mathcal{AS}[F; r_i](z) = \int_a^b \frac{1}{(1-zx)^{r_i}} \frac{(x-a)^{s_i-1} (b-x)^{r_i-s_i-1}}{B(s_i, r_i-s_i)(b-a)^{r_i-1}} dx.$$

Changing of variable $t = \frac{x-a}{b-a}$ implies

$$\mathcal{AS}[F; r_i](z) = \frac{1}{(1-za)^{r_i}} \int_0^1 \frac{1}{(1 - \frac{z(b-a)}{1-za}t)^{r_i}} \frac{t^{s_i-1} (1-t)^{r_i-s_i-1}}{B(s_i, r_i-s_i)} dt.$$

Using (1.4), it follows that

$$\begin{aligned} \mathcal{AS}[F; r_i](z) &= \frac{1}{(1-za)^{r_i}} F_D^{(1)}(r_i, s_i; r_i; \frac{z(b-a)}{1-za}) \\ &= \frac{1}{(1-za)^{r_i}} \frac{1}{(1 - \frac{z(b-a)}{1-za})^{s_i}} \\ &= \frac{1}{(1-za)^{r_i-s_i}} \frac{1}{(1-zb)^{s_i}}. \end{aligned}$$

From Theorem 2.2, we get

$$\mathcal{AS}[F_{S_{n:n_1, \dots, n_{k-1}}}; \sum_{i=1}^k r_i](z) = \frac{1}{(1-za)^{\sum_{i=1}^k (r_i-s_i)}} \frac{1}{(1-zb)^{\sum_{i=1}^k s_i}}.$$

which is the AST of $\tilde{\text{beta}}(\sum_{i=1}^k s_i, \sum_{i=1}^k r_i - \sum_{i=1}^k s_i)$ distribution. Hence $S_{n:n_1, \dots, n_{k-1}}$ has $\tilde{\text{beta}}(\sum_{i=1}^k s_i, \sum_{i=1}^k r_i - \sum_{i=1}^k s_i)$ distribution. \square

The following corollary is an immediate consequence of Theorem 3.2.

Corollary 3.1 For independent random variables X_i , $i = 1, 2, \dots, k$ with $\tilde{\text{beta}}(r_i/2, r_i/2)$ distributions, the RWA $S_{n:n_1, \dots, n_{k-1}}$ has $\tilde{\text{beta}}(\sum_{i=1}^k r_i/2, \sum_{i=1}^k r_i/2)$ distribution.

The following theorem of [6] comes as a corollary to Corollary 3.1.

Corollary 3.2 For every integer $n \geq 2$, a power semicircle distribution with shape parameter $\theta = (n-3)/2$ and any positive range parameter is a RWA S_n distribution.

Proof Consider $k = n$, $r_i = 1$ for $i = 1, 2, \dots, k$, $p = q = 1/2$, $a = -\sigma$ and $b = \sigma$, then use Corollary 3.1 to obtain the result. \square

4 Examples

Theorem 3.1 and Theorem 3.2 in Section 3 give general results on distribution of RWA $S_{n:n_1, \dots, n_{k-1}}$ on some beta distributions on $[a, b]$. In this section we present some special examples of distributions that are randomly weighted average distributions given in (1.2). Other examples of RWA S_n distributions in case of $r_1 = r_2 = \dots = r_k = 1$ and $k = n$ can be found in [6]. For the first and second examples, we use the results of [5].

Example 4.1. Let X_1, X_2 and R be independent random variables with $\tilde{beta}(1/2, 1/2)$, $\tilde{beta}(m-1/2, m-1/2)$ and $beta(1, m-1)$ distributions, respectively. Then randomly weighted average $S = RX_1 + (1 - R)X_2$ has $\tilde{beta}(m - 1/2, m - 1/2)$ distribution. Its AST can be written as

$$\mathcal{AS}[F_S; m](z) = \frac{1}{\sqrt{1 - z(a + b) + abz^2}} \left[\frac{4}{2 - z(a + b) + 2\sqrt{1 - z(a + b) + abz^2}} \right]^{m-1}.$$

Example 4.2. Let X_1, X_2 and R be independent random variables with $\tilde{beta}(3/2, 1/2)$, $\tilde{beta}(m-1/2, m-1/2)$ and $beta(1, m-1)$ distributions, respectively. Then randomly weighted average $S = RX_1 + (1 - R)X_2$ has $\tilde{beta}(m + 1/2, m - 1/2)$ distribution. Its AST is given by

$$\mathcal{AS}[F_S; m](z) = \frac{2}{\sqrt{1 - zab + \sqrt{1 - z(a + b) + abz^2}}} \left[\frac{4}{2 - z(a + b) + 2\sqrt{1 - z(a + b) + abz^2}} \right]^{m-1}.$$

Example 4.3. Let X_1, X_2, \dots, X_k be independent and identically distributed (i.i.d) random variables with common $U(a, b)$ distribution and random vector (V_1, \dots, V_k) has $Dir(2, 2, \dots, 2)$ distribution. Then by Corollary 3.1, the RWA $S_{n:n_1, \dots, n_{k-1}}$ has $\tilde{beta}(k, k)$ distribution.

Example 4.4. Let X_1, X_2, \dots, X_k be i.i.d random variables with common semicircle (Wigner) distribution on $(-\sigma, \sigma)$ and random vector (V_1, \dots, V_k) has $Dir(3, 3, \dots, 3)$ distribution.

Then by Corollary 3.1, the RWA $S_{n:n_1, \dots, n_{k-1}}$ has $\tilde{\text{beta}}(3k/2, 3k/2)$ distribution.

References

- [1] Abramowitz, M., and Stegun, I. A. (Eds.) (2012). *Handbook of mathematical functions: with formulas, graphs, and mathematical tables*. Courier Dover Publications.
- [2] Andrews, G. E., Askey, R., and Roy, R. (1999). *Special functions*. Cambridge University press.
- [3] Debnath, L., and Bhatta, D. (2007). *Integral Transforms and Their Applications*. Chapman and Hall/CRC Press, Boca Raton.
- [4] Demni, N. (2009). On generalized Cauchy-Stieltjes transforms of some beta distributions. *Comm. Stoch. Anal.* 3, 197-210.
- [5] Kubo, I., Kuo, H-H., and Namli, S. (2011). MRM-Applicable measures for the power function of the second order. *Comm. Stoch. Annal.* 5(4), 647-670.
- [6] Roozegar, R. and Soltani, A. R. (2013). Classes of power semicircle laws that are randomly weighted average distributions. *J. Stat. Comput. Simul.* to appear. doi: 10.1080/00949655.2013.806510.
- [7] Soltani, A. R., and Roozegar, R. (2012). On distribution of randomly ordered uniform incremental weighted averages: Divided difference approach. *Statist. Probab. Lett.* 82(5), 1012-1020.