

Kibble–Slepian Formula and Generating Functions for 2D Polynomials

Mourad E. H. Ismail^{a,b,1}, Ruiming Zhang^{c,2,*}

^a*Department of Mathematics, King Saud University, Riyadh, Saudi Arabia*

^b*Department of Mathematics, University of Central Florida, Orlando, Florida 32816, USA*

^c*College of Science, Northwest A&F University, Yangling, Shaanxi 712100, P. R. China*

Abstract

We prove a generalization of the Kibble–Slepian formula (for Hermite polynomials) and its unitary analogue involving the 2D Hermite polynomials recently proved in [17]. We derive integral representations for the 2D Hermite polynomials which are of independent interest. Several new generating functions for 2D q -Hermite polynomials will also be given.

Keywords:

Hermite polynomials, 2D Hermite polynomials, 2D q -Hermite polynomials, Poisson kernels, positivity of kernels, integral operators, multilinear generating functions, Kibble–Slepian formula.

2010 MSC: Primary 33C45, Secondary 42C20.

1. Introduction

The complex Hermite polynomials $\{H_{m,n}(z_1, z_2)\}_{m,n=0}^{\infty}$ may be defined by

$$H_{m,n}(z_1, z_2) = \sum_{k=0}^{m \wedge n} (-1)^k k! \binom{m}{k} \binom{n}{k} z_1^{m-k} z_2^{n-k}. \quad (1.1)$$

*Corresponding author

Email addresses: mourad.eh.ismail@gmail.com (Mourad E. H. Ismail), ruimingzhang@yahoo.com (Ruiming Zhang)

¹Research supported by a grant from the DSFP program at King Saud University and by the National Plan for Science, Technology and Innovation (MAARIFAH), King Abdelaziz City for Science and Technology, Kingdom of Saudi Arabia, Award Number 14-MAT623-02.

²Research partially supported by National Science Foundation of China, Grant No. 11371294.

The polynomials $\{H_{m,n}(z, \bar{z})\}_{m,n=0}^{\infty}$ are orthogonal on \mathbb{R}^2 with respect to $e^{-x^2-y^2}$ and have the generating function

$$\sum_{m,n=0}^{\infty} H_{m,n}(z_1, z_2) \frac{u^m v^n}{m! n!} = e^{uz_1 + vz_2 - uv}. \quad (1.2)$$

They seem to have been considered first by Ito [19] in his study of complex multiple Wiener integrals. Recently they were used in [1] to study Landau levels and were applied in [23] to coherent states, and in [24, 25] to quantum optics and quasiprobabilities, respectively. See also [6, 10, 11]. The reference [14] deals with the spectral properties of the Cauchy transform and the polynomials $\{H_{m,n}(z, \bar{z})\}$ also appear in this context. The polynomials $\{H_{m,n}(z, \bar{z})\}_{m,n=0}^{\infty}$ are essentially the same polynomials as in the monograph [7, (2.6.6)] by Dunkl and Xu.

The Kibble–Slepian formula is Equation (1.4) below. It was first proved by Kibble in 1945 [20] and later by Slepian [22]. Louck [21] gave a proof using Boson operators while Foata [8] gave a purely combinatorial proof. Each proof brings in a new point of view.

Theorem 1.1. *Let $S = (s_{j,k})_{j,k=1}^N$ be an $N \times N$ real symmetric matrix with the Frobenius norm*

$$\|S\|^2 = \sum_{j,k=1}^N |s_{j,k}|^2. \quad (1.3)$$

Assume that $\|S\| < 1$, I_N being an identity matrix of size N , and X being an $N \times 1$ matrix. Then

$$\begin{aligned} & \det(I_N + S)^{-\frac{1}{2}} \exp(X^T S (I_N + S)^{-1} X) \\ &= \sum_K \left(\prod_{1 \leq m \leq n \leq N} \frac{(s_{m,n})^{k_{m,n}}}{2^{k_{m,n}} k_{m,n}!} \right) 2^{-\text{tr}(K)} H_{k_1}(x_1) \cdots H_{k_N}(x_N), \end{aligned} \quad (1.4)$$

where $X = (x_1, x_2, \dots, x_N)^T$, $K = (k_{m,n})_{m,n=1}^N$, $k_{m,n} = k_{n,m}$, $1 \leq m, n \leq N$ and

$$\text{tr}(K) = \sum_{j=1}^N k_{j,j}, \quad k_{\ell} = k_{\ell,\ell} + \sum_{j=1}^N k_{\ell,j}, \quad \ell = 1, \dots, N. \quad (1.5)$$

In (1.4), \sum_K denotes the $\frac{n(n+1)}{2}$ fold sum over all nonnegative integers $k_{m,n} = 0, 1, \dots$ for all positive integers m, n such that $1 \leq m \leq n \leq N$.

It must be noted that the proofs by Louck [21] and Slepian [22] assume S is symmetric and conclude that the expansion in (1.4) holds for $\|S\| < 1$. On the other hand the combinatorial version by Foata [8] makes no assumptions on S but assumes the diagonal elements $h_{j,j}$ vanish, and concludes that the expansion (1.4) holds as a formal power series.

In [17] Ismail proved a similar theorem for the complex Hermite polynomials. His result is essentially the following theorem.

Theorem 1.2. *Let $W = (w_1, w_2, \dots, w_N)^T$, and $H = (h_{m,n})_{m,n=1}^N$ be an $N \times N$ Hermitian matrix with $\|H\| < 1$ in Frobenius norm, and I_N is an $N \times N$ identity matrix. Then*

$$\begin{aligned} & \det(I_N + H)^{-1} \exp(W^* H (I_N + H)^{-1} W) \\ &= \sum_K \prod_{1 \leq m, n \leq N} \frac{(h_{m,n})^{k_{m,n}}}{k_{m,n}!} H_{r_1, c_1}(\overline{w_1}, w_1) \cdots H_{r_N, c_N}(\overline{w_N}, w_N), \end{aligned} \quad (1.6)$$

where $K = (k_{m,n})_{m,n=1}^N$ is a general matrix with nonnegative integer entries, c_n is the sum of the elements of K in column n and r_m is the sum of the elements of K in row m , that is

$$c_n = \sum_{j=1}^N k_{j,n}, \quad r_m = \sum_{\ell=1}^N k_{m,\ell}. \quad (1.7)$$

In this paper we prove the following stronger result without the assumption that $H \in \mathbb{C}^{N \times N}$ is Hermitian.

Theorem 1.3. *Following the notations in Theorem 1.2, we let $W = (w_1, \dots, w_N)^T \in \mathbb{C}^N$, $H \in \mathbb{C}^{N \times N}$, $\|H\|_\infty = \max_{1 \leq j, \ell \leq N} |h_{j,\ell}|$ and $B = \{H : \|H\|_\infty < \frac{1}{N}\}$. Then the series*

$$\sum_K \prod_{1 \leq m, n \leq N} \frac{(h_{m,n})^{k_{m,n}}}{k_{m,n}!} H_{r_1, c_1}(\overline{w_1}, w_1) \cdots H_{r_N, c_N}(\overline{w_N}, w_N)$$

converges absolutely and uniformly for W in any compact subset of \mathbb{C}^N and H in any compact subset of B .

Given $\delta_{j,k} > 0$, $j, k = 1, \dots, N$ and a Hermitian matrix $H_0 = (h_{j,k}^{(0)})_{j,k=1}^N \in B$, let

$$D(H_0, \delta) = \left\{ H : \left| h_{j,j} - h_{j,j}^{(0)} \right| < \delta_{j,j}, \left| u_{\ell,k} - u_{\ell,k}^{(0)} \right| < \delta_{\ell,k}, \left| v_{\ell,k} - v_{\ell,k}^{(0)} \right| < \delta_{\ell,k} \right\}, \quad (1.8)$$

where $1 \leq j, k, \ell \leq N$, $\ell < k$ and

$$u_{\ell,k} = \frac{h_{\ell,k} + h_{k,\ell}}{2}, v_{\ell,k} = \frac{h_{\ell,k} - h_{k,\ell}}{2i}, u_{\ell,k}^{(0)} = \frac{h_{\ell,k}^{(0)} + h_{k,\ell}^{(0)}}{2}, v_{\ell,k}^{(0)} = \frac{h_{\ell,k}^{(0)} - h_{k,\ell}^{(0)}}{2i}. \quad (1.9)$$

If $D(H_0, \delta) \subset B$, then

$$\begin{aligned} & \exp(W^*H(I_N + H)^{-1}W) = \det(I_N + H) \\ & \times \sum_K \prod_{1 \leq m, n \leq N} \frac{(h_{m,n})^{k_{m,n}}}{k_{m,n}!} H_{r_1, c_1}(\bar{w}_1, w_1) \cdots H_{r_N, c_N}(\bar{w}_N, w_N) \end{aligned} \quad (1.10)$$

holds for all $W \in \mathbb{C}^N$ and $H \in D(H_0, \delta)$. In particular, it is not hard to see that $D\left((0)_{j,k=1}^N, \left(\frac{1}{2N}\right)_{j,k=1}^N + \frac{I_N}{2N}\right) \subset B$.

Corollary 1.4. For $N \in \mathbb{N}$, let $W = (\rho_1 e^{i\theta_1}, \dots, \rho_N e^{i\theta_N})^T$ that $\rho_m > 0$, $\theta_m \in \mathbb{R}$ for $m = 1, \dots, N$ in (1.6), H , I_N , K , c_m and r_m are the same as in Theorem 1.2, then

$$\begin{aligned} & \det(I_N + H)^{-1} \exp(W^*H(I_N + H)^{-1}W) \\ &= \sum_K \prod_{m=1}^N \prod_{n=1}^N (-h_{m,n})^{k_{m,n}} \binom{c_m}{k_{1,m}, \dots, k_{N,m}} (\rho_m e^{i\theta_m})^{r_m - c_m} L_{c_m}^{(r_m - c_m)}(\rho_m^2), \end{aligned} \quad (1.11)$$

where $\{L_n^{(\alpha)}(x)\}$ are Laguerre polynomials. In particular, for $x, y > 0$ and $|u|, |v| < \frac{|xy|}{4}$, we have

$$\begin{aligned} & \sum_{0 \leq j < k < \infty} \frac{(u^j v^k + u^k v^j)}{j! k!} C_j(k; x) C_j(k; y) \\ &= \frac{xy}{xy - uv} \exp\left(-\frac{xy(xuv - xy(u+v) + yuv)}{xy - uv}\right) \\ & \quad - \frac{xy}{xy - uv} \exp\left(-\frac{uv(x^2 + y^2)}{xy - uv}\right) I_0\left(2 \frac{\sqrt{uv(xy)^{3/2}}}{xy - uv}\right), \end{aligned} \quad (1.12)$$

where $C_n(x; a)$ is the n -th Charlier polynomial, $I_\alpha(z)$ is the Bessel function of first kind.

Ismail's proof in [17] assumes that H is Hermitian and $\|H\| < 1$ and proves that (1.6) holds as a convergent power series in the variables $h_{j,k}$, $1 \leq j \leq k \leq N$.

Later Ismail and Zeng [18] found a combinatorial proof of Theorem 1.2 where H is not necessary symmetric, but the power series in (1.6) is a formal power series.

The purpose of this paper is to first prove Theorems 1.1 and 1.2 and Corollary 1.4 by using the integral representations

$$H_n(x) e^{-x^2} = \frac{(-2i)^n}{\sqrt{\pi}} \int_{-\infty}^{\infty} t^n e^{-t^2+2ixt} dt, \quad (1.13)$$

and

$$e^{-z\bar{z}} H_{m,n}(z, \bar{z}) = \frac{i^{m+n}}{\pi} \int_{\mathbb{R}^2} w^m \bar{w}^n \exp\left\{-\left(r^2 + s^2\right) - 2i \operatorname{Re}(w\bar{z})\right\} dr ds, \quad (1.14)$$

where $z = x + iy$ and $w = r + is$ such that $r, s, x, y \in \mathbb{R}$. The representation (1.13) is well-known, see for example, formula (4.6.41) in [16], while (1.14) will be proved in §3. Our proof actually proves a stronger version of Theorems 1.1–1.2, where S and H are not assumed to be symmetric and Hermitian, respectively.

It is important to note that the left-hand sides of the multilinear generating functions in Theorems 1.1–1.2 are positive, when S and H are real symmetric and Hermitian, respectively. They contain the Poisson kernels as the special cases when $N = 2$ and the diagonal elements of the matrices involved are zero, [17], [16, §4.7]. Carlitz [5] actually found the Poisson kernel for the 2D Hermite polynomials in 1978, 20 years before [24, 25]. He identified the 2D Hermite polynomials as special cases of a 3D system which he studied in detail but did not derive its orthogonality. Carlitz was not aware that his polynomials are the same as Ito's 2D-Hermite polynomials. Carlitz did not elaborate on the orthogonality of his bivariate or trivariate polynomials.

Section 2 contains the proofs of Theorems 1.1–1.2. Section 3 contains some new formal properties of the 2D Hermite polynomials. In our approach we treat $H_{m,n}(z_1, z_2)$ as a function of two independent complex variables and view the case $z_2 = \bar{z}_1$ as a domain of orthogonality in \mathbb{C}^2 . In Section 4 we derive several multilinear generating functions for the 2D q -Hermite polynomials we introduced in [15]. We do not have a q -analogue of the Kibble–Slepian formula of Theorem 1.1 but the results in §4 would be special cases of such formula. There is no Kibble–Slepian formula known for the one variable q -Hermite polynomials either but their Poisson kernel is known.

2. Proofs

We shall use the multidimensional Taylor series for functions mapping \mathbb{R}^N into \mathbb{R} . For $\alpha = (\alpha_1, \alpha_2, \dots)$ such that $\alpha_1, \alpha_2, \dots$ are nonnegative integers, let $|\alpha| = \alpha_1 + \alpha_2 + \dots$, $\alpha! = \alpha_1! \cdot \alpha_2! \cdots$, and $\mathbf{x}^\alpha = x_1^{\alpha_1} \cdot x_2^{\alpha_2} \cdots$. Additionally, for $n \in \mathbb{N}_0$ and $|\alpha| = n$ we let $\binom{n}{\alpha} = \frac{n!}{\alpha!}$ and $D^\alpha f(\mathbf{x}) = \frac{\partial^{|\alpha|} f(\mathbf{x})}{\partial x_1^{\alpha_1} \cdot \partial x_2^{\alpha_2} \cdots}$.

Theorem 2.1. Assume that f and all its partial derivatives of order $< m$ are differentiable at each point of an open set $S \subset \mathbb{R}^n$. If \mathbf{a} and \mathbf{b} are two points of S such that the line joining \mathbf{a} and \mathbf{b} is contained in S . We further let

$$f^{(k)}(\mathbf{x}; \mathbf{t}) = \sum_{|\alpha|=k} \binom{k}{\alpha} D^\alpha f(\mathbf{x}) \mathbf{t}^\alpha, \quad (2.1)$$

then

$$f(\mathbf{b}) = f(\mathbf{a}) + \sum_{k=1}^{m-1} \frac{1}{k!} f^{(k)}(\mathbf{a}; \mathbf{b} - \mathbf{a}) + \frac{1}{m!} f^{(m)}(\mathbf{z}; \mathbf{b} - \mathbf{a}), \quad (2.2)$$

for some \mathbf{z} on the line segment joining \mathbf{b} and \mathbf{a} .

This is essentially Theorem 12.14 in [3].

Lemma 2.2. Let $S = (s_{j,k})_{j,k=1}^N$ be an $N \times N$ real symmetric matrix and Y an $N \times 1$ complex matrix, then,

$$\exp(-Y^T S Y) = \sum_K \left(\prod_{1 \leq m \leq n \leq N} \frac{(-2s_{m,n})^{k_{m,n}}}{k_{m,n}!} \right) 2^{-\text{tr}(K)} y_1^{k_1} y_2^{k_2} \cdots y_n^{k_n}, \quad (2.3)$$

where $K, k_j, \text{tr}(K)$ are the same as in Theorem 1.1.

Proof. Observe that $\exp(-Y^T S Y)$ is an analytic function in the variables $s_{m,n}$ for $1 \leq m \leq n \leq N$ separately, then,

$$\exp(-Y^T S Y) = \sum_K \left(\prod_{1 \leq m \leq n \leq N} \frac{(s_{m,n})^{k_{m,n}}}{k_{m,n}!} a_{k_{m,n}} \right),$$

then for $1 \leq m = n \leq N$,

$$a_{k_{m,m}} = \frac{\partial^{k_{m,m}} \exp(-Y^T S Y)}{\partial s_{m,m}^{k_{m,m}}} \Big|_{S=0} = (-1)^{k_{m,m}} y_m^{2k_{m,m}},$$

and for $1 \leq m < n \leq N$,

$$a_{k_{m,n}} = \frac{\partial^{k_{m,n}} \exp(-Y^T S Y)}{\partial s_{m,n}^{k_{m,n}}} \Big|_{S=0} = (-2)^{k_{m,n}} y_m^{k_{m,n}} y_n^{k_{m,n}}.$$

We apply Theorem 2.1 and show that the error term $\rightarrow 0$ as $m \rightarrow \infty$ and conclude that

$$\exp(-Y^T S Y) = \sum_K \left(\prod_{1 \leq m \leq n \leq N} \frac{(-2s_{m,n})^{k_{m,n}}}{k_{m,n}!} \right) 2^{-\text{tr}(K)} y_1^{k_1} y_2^{k_2} \cdots y_n^{k_n}.$$

□

Lemma 2.3. Let $H = (h_{j,k})_{j,k=1}^N$ be an $N \times N$ complex matrix and Z an $N \times 1$ complex matrix $Z = (z_1, \dots, z_N)^T$ and $z_j = x_j + iy_j$, $x_j, y_j \in \mathbb{R}$ for $j = 1, \dots, N$, then,

$$\exp(-Z^* H Z) = \sum_K \prod_{j=1}^N (\bar{z}_j)^{r_j} z_j^{c_j} \left(\prod_{1 \leq m, n \leq N} \frac{(-h_{m,n})^{k_{m,n}}}{k_{m,n}!} \right), \quad (2.4)$$

where $K, k_{m,n}, r_m, c_n$ are the same as in Theorem 1.2.

Proof. Observe that $\exp(-Z^* H Z)$ is analytic in the variables $h_{j,k}$, $j, k = 1, \dots, N$ separately, then

$$\exp(-Z^* H Z) = \sum_K \prod_{1 \leq m, n \leq N} \frac{(h_{m,n})^{k_{m,n}}}{k_{m,n}!} a_{k_{m,n}}$$

and

$$a_{k_{m,n}} = \frac{\partial^{k_{m,n}} \exp(-Z^* H Z)}{\partial h_{m,n}^{k_{m,n}}} \Big|_{H=0} = (-\bar{z}_m z_n)^{k_{m,n}}$$

for $k_{m,n} = 0, 1, \dots$ and $m, n = 1, \dots, N$, then,

$$\begin{aligned} \exp(-Z^* H Z) &= \sum_K \prod_{1 \leq m, n \leq N} \frac{(-h_{m,n})^{k_{m,n}} (\bar{z}_m)^{k_{m,n}} (z_n)^{k_{m,n}}}{k_{m,n}!} \\ &= \sum_K \prod_{j=1}^N (\bar{z}_j)^{r_j} z_j^{c_j} \left(\prod_{1 \leq m, n \leq N} \frac{(-h_{m,n})^{k_{m,n}}}{k_{m,n}!} \right). \end{aligned}$$

This completes the proof. □

Remark 1. We have noticed that identities 2.3 and 2.4 can be proved formally without using Theorem 2.1. We observe that for the former we have,

$$\begin{aligned}
\exp(-Y^T S Y) &= \exp\left(-\sum_{i,j=1}^N s_{i,j} y_i y_j\right) = \exp\left(-\sum_{i=1}^N s_{i,i} y_i^2 - 2 \sum_{1 \leq i < j \leq N} s_{i,j} y_i y_j\right) \\
&= \left(\prod_{i=1}^N \sum_{k_{i,i}=0}^{\infty} \frac{(-s_{i,i} y_i^2)^{k_{i,i}}}{(k_{i,i})!} \right) \cdot \left(\prod_{1 \leq m < n \leq N} \sum_{k_{m,n}=0}^{\infty} \frac{(-2s_{m,n} y_m y_n)^{k_{m,n}}}{(k_{m,n})!} \right) \\
&= \sum_{k_{1,1}, k_{2,2}, \dots, k_{N,N}=0}^{\infty} \frac{(-s_{1,1} y_1^2)^{k_{1,1}}}{(k_{1,1})!} \frac{(-s_{2,2} y_2^2)^{k_{2,2}}}{(k_{2,2})!} \cdots \frac{(-s_{N,N} y_N^2)^{k_{N,N}}}{(k_{N,N})!} \\
&\quad \times \sum_{k_{1,2}, k_{1,3}, k_{1,N}, k_{2,3}, \dots, k_{2,N}, \dots, k_{N-1,N}=0}^{\infty} \frac{(-2s_{1,2} y_1 y_2)^{k_{1,2}}}{(k_{1,2})!} \cdots \frac{(-2s_{1,N} y_1 y_N)^{k_{1,N}}}{(k_{1,N})!} \\
&\quad \times \frac{(-2s_{2,3} y_2 y_3)^{k_{2,3}}}{(k_{2,3})!} \cdots \frac{(-2s_{2,N} y_2 y_N)^{k_{2,N}}}{(k_{2,N})!} \cdots \frac{(-2s_{N-1,N} y_{N-1} y_N)^{k_{N-1,N}}}{(k_{N-1,N})!} \\
&= \sum_K \left(\prod_{i=1}^N \left(\frac{y_i}{2}\right)^{k_{i,i}} \right) \prod_{1 \leq m \leq n \leq N} \frac{(-2s_{m,n} y_m y_n)^{k_{m,n}}}{(k_{m,n})!} = \sum_K 2^{-\text{tr}(K)} \prod_{1 \leq m \leq n \leq N} \frac{(-2s_{m,n})^{k_{m,n}}}{(k_{m,n})!} \\
&\quad \times y_1^{2k_{1,1}+k_{1,2}+\dots+k_{1,N}} y_2^{2k_{2,2}+k_{2,3}+\dots+k_{2,N}} \cdots y_{N-1}^{2k_{N-1,N-1}+k_{N-1,N}} y_N^{2k_{N,N}} \\
&= \sum_K \left(\prod_{1 \leq m \leq n \leq N} \frac{(-2s_{m,n})^{k_{m,n}}}{k_{m,n}!} \right) 2^{-\text{tr}(K)} y_1^{k_1} y_2^{k_2} \cdots y_N^{k_N},
\end{aligned}$$

whereas for the latter we have,

$$\begin{aligned}
\exp(-Z^* H Z) &= \exp\left(-\sum_{m,n=1}^N h_{m,n} \overline{z}_m z_n\right) = \prod_{1 \leq m, n \leq N} \exp(-h_{m,n} \overline{z}_m z_n) \\
&= \prod_{1 \leq m, n \leq N} \sum_{k_{m,n}=0}^{\infty} \frac{(-h_{m,n} \overline{z}_m z_n)^{k_{m,n}}}{(k_{m,n})!} = \sum_K \prod_{1 \leq m, n \leq N} \frac{(-h_{m,n})^{k_{m,n}}}{(k_{m,n})!} \\
&\quad \times (\overline{z}_1)^{k_{1,1}+k_{1,2}+\dots+k_{1,N}} \cdots (\overline{z}_N)^{k_{N,1}+k_{N,2}+\dots+k_{N,N}} z_1^{k_{1,1}+k_{2,1}+\dots+k_{N,1}} \cdots \\
&\quad \times z_N^{k_{1,N}+k_{2,N}+\dots+k_{N,N}} = \sum_K \left(\prod_{1 \leq m, n \leq N} \frac{(-h_{m,n})^{k_{m,n}}}{(k_{m,n})!} \right) (\overline{z}_1)^{r_1} \cdots (\overline{z}_N)^{r_N} z_1^{c_1} \cdots z_N^{c_N} \\
&= \sum_K \left(\prod_{j=1}^N \left(\overline{z}_j\right)^{r_j} z_j^{c_j} \right) \left(\prod_{1 \leq m, n \leq N} \frac{(-h_{m,n})^{k_{m,n}}}{(k_{m,n})!} \right).
\end{aligned}$$

Lemma 2.4. For all $m, n \in \mathbb{Z}^+$ and $z \in \mathbb{C}$ we have

$$|H_{m,n}(\bar{z}, z)| \leq e^{|z|^2} \sqrt{m! \cdot n!}. \quad (2.5)$$

Proof. From Equation (1.14) we get

$$\begin{aligned} \pi e^{-|z|^2} H_{m,n}(\bar{z}, z) &\leq \int_{\mathbb{R}^2} |w|^m \cdot |w|^n \exp\{- (r^2 + s^2)\} dr ds \\ &\leq \sqrt{\int_{\mathbb{R}^2} |w|^{2m} \exp\{- (r^2 + s^2)\} dr ds} \sqrt{\int_{\mathbb{R}^2} |w|^{2n} \exp\{- (r^2 + s^2)\} dr ds} \\ &= \sqrt{\int_{\mathbb{R}^2} (r^2 + s^2)^m \exp\{- (r^2 + s^2)\} dr ds} \sqrt{\int_{\mathbb{R}^2} (r^2 + s^2)^n \exp\{- (r^2 + s^2)\} dr ds} \\ &= \pi \sqrt{m! \cdot n!}, \end{aligned}$$

which gives (2.5). \square

We now present our proof of Theorem 1.1.

Proof of Theorem 1.1. First we observe that

$$\|S\|^2 = \sum_{m,n=1}^N |s_{m,n}|^2 = \text{tr}(SS^T) = \sum_{j=1}^N \lambda_j^2,$$

where λ_j , $j = 1, \dots, N$ are the eigenvalues of S , then the matrix $I_N + S$ is positive definite and thus $(I_N + S)^{-1}$ exists and it is positive definite. It is clear that,

$$\begin{aligned} &\det(I_N + S)^{-\frac{1}{2}} \exp(X^T S (I_N + S)^{-1} X) \\ &= \det(I_N + S)^{-\frac{1}{2}} \exp(-X^T (I_N + S)^{-1} X + X^T X), \end{aligned}$$

then (1.4) is equivalent to

$$\begin{aligned} &\det(I_N + S)^{-\frac{1}{2}} \exp(-X^T (I_N + S)^{-1} X) \\ &= \sum_K \left(\prod_{1 \leq m \leq n \leq N} \frac{(s_{m,n})^{k_{m,n}}}{2^{k_{m,n}} k_{m,n}!} \right) 2^{-\text{tr}(K)} \psi_{k_1}(x_1) \cdots \psi_{k_N}(x_N), \end{aligned}$$

where $\psi_n(x) = e^{-x^2} H_n(x)$. Applying the multivariate normal integral [4]

$$\int_{\mathbb{R}^N} \exp(-X^T A X + 2i B^T X) \prod_{j=1}^N dx_j = \sqrt{\frac{\pi^N}{\det A}} e^{-B^T A^{-1} B}, \quad (2.6)$$

where $N \in \mathbb{N}$, A is an $N \times N$ real symmetric positive definite matrix and B, X are $N \times 1$ real matrices, then using Lemma 2.2 and (1.13) we get

$$\begin{aligned} & \det(I_N + S)^{-\frac{1}{2}} \exp(-X^T (I_N + S)^{-1} X) \\ &= \pi^{-\frac{N}{2}} \int_{\mathbb{R}^N} \exp(-Y^T (I_N + S) Y + 2i X^T Y) \prod_{j=1}^N dy_j \\ &= \pi^{-\frac{N}{2}} \int_{\mathbb{R}^N} \exp(-Y^T Y + 2i X^T Y - Y^T S Y) \prod_{n=1}^N dy_n \\ &= \pi^{-\frac{N}{2}} \sum_K \left(\prod_{1 \leq m \leq n \leq N} \frac{(-2s_{m,n})^{k_{m,n}}}{k_{m,n}!} \right) 2^{-\text{tr}(K)} \\ &\quad \times \int_{\mathbb{R}^N} \exp(-Y^T Y + 2i X^T Y) \prod_{n=1}^N y_n^{k_n} dy_n \\ &= \sum_K \left(\prod_{1 \leq m \leq n \leq N} \frac{(-2s_{m,n})^{k_{m,n}}}{k_{m,n}!} \right) 2^{-\text{tr}(K)} \\ &\quad \times \frac{1}{(-2i)^{\sum_{j=1}^N k_j}} \psi_{k_1}(x_1) \cdots \psi_{k_n}(x_n), \end{aligned}$$

where the exchange order of summation and integration is valid because of

$$\int_{\mathbb{R}^N} \exp(-Y^T Y - Y^T S Y) \prod_{n=1}^N dy_n < \infty$$

and an application of Fubini's theorem. Observe that

$$\sum_{j=1}^N k_j = 2 \sum_{1 \leq m \leq n \leq N} k_{m,n},$$

then

$$\begin{aligned} & \det(I_N + S)^{-\frac{1}{2}} \exp(-X^T (I_N + S)^{-1} X) \\ &= \sum_K \left(\prod_{1 \leq m \leq n \leq N} \frac{\left(\frac{s_{m,n}}{2}\right)^{k_{m,n}}}{k_{m,n}!} \right) 2^{-\text{tr}(K)} \psi_{k_1}(x_1) \cdots \psi_{k_n}(x_n), \end{aligned}$$

which proves Theorem 1.1. \square

Proof of Theorem 1.2. Since H is an $N \times N$ Hermitian matrix, then H can be factored into

$$H = U \Lambda U^*$$

where U is an $N \times N$ unitary matrix while Λ is an $N \times N$ real diagonal, say $\Lambda = \text{diag}\{\lambda_1, \dots, \lambda_N\}$, thus,

$$\|H\|^2 = \text{tr}(HH^*) = \sum_{j=1}^N \lambda_j^2 < 1.$$

Hence $I_N + H$ is a positive definite Hermitian matrix and thus it is invertible. It is clear that (1.6) is the same as

$$\begin{aligned} & \det(I_N + H)^{-1} \exp(Z^* (I_N + H)^{-1} Z) \\ &= \sum_K \prod_{1 \leq m, n \leq N} \frac{(h_{m,n})^{k_{m,n}}}{k_{m,n}!} \psi_{r_1, c_1}(\overline{w_1}, w_1) \cdots \psi_{r_N, c_N}(\overline{w_N}, w_N), \end{aligned}$$

where $\psi_{\alpha, \beta}(z_1, z_2) = e^{-z_1 z_2} H_{\alpha, \beta}(z_1, z_2)$. Set $\Gamma = (I_N + H)^{-1}$ to be an $N \times N$ positive definite Hermitian matrix, $\mu = (0, \dots, 0)^T$ an $N \times 1$ complex matrix and $C = (0)_{j,k=1}^N$ in the character function complex normal integral to get [12]

$$\begin{aligned} & \int_{\mathbb{R}^{2N}} \exp(-Z^* (I_N + H) Z + 2i \text{Re}(W^* Z)) \prod_{j=1}^N dx_j dy_j \\ &= \frac{\pi^N \exp(-W^* (I_N + H)^{-1} W)}{\det(I_N + H)} \end{aligned} \tag{2.7}$$

where $W = (w_1, \dots, w_N)^T$ is an $N \times 1$ complex matrix. From Lemma 2.3 to get

$$\begin{aligned}
& \frac{\exp(-W^*(I_N + H)^{-1} W)}{\det(I_N + H)} \\
&= \pi^{-N} \int_{\mathbb{R}^{2N}} \exp(-Z^*Z + 2i \operatorname{Re}(W^*Z) - Z^*HZ) \prod_{j=1}^N dx_j dy_j \\
&= \pi^{-N} \sum_K \left(\prod_{1 \leq m, n \leq N} \frac{(-h_{m,n})^{k_{m,n}}}{k_{m,n}!} \right) \\
&\quad \times \int_{\mathbb{R}^{2N}} \exp(-Z^*Z + 2i \operatorname{Re}(W^*Z)) \prod_{j=1}^N (\bar{z}_j)^{r_j} z_j^{c_j} dx_j dy_j. \\
&= \sum_K \prod_{1 \leq m, n \leq N} \frac{(h_{m,n})^{k_{m,n}}}{k_{m,n}!} \psi_{r_1, c_1}(\bar{w}_1, w_1) \cdots \psi_{r_N, c_N}(\bar{w}_N, w_N),
\end{aligned}$$

and the exchange order of summation and integration can be verified using Fubini's theorem. \square

Proof of Theorem 1.3. By Lemma 2.4 we have

$$|H_{r_1, c_1}(\bar{w}_1, w_1) \cdots H_{r_N, c_N}(\bar{w}_N, w_N)| \leq \exp\left(\sum_{i=1}^N |w_i|^2\right) \sqrt{\prod_{i=1}^N r_i! \cdot \prod_{i=1}^N c_i!}$$

and

$$\begin{aligned}
& \left| \sum_K \prod_{1 \leq j, \ell \leq N} \frac{(h_{j,\ell})^{k_{j,\ell}}}{k_{j,\ell}!} H_{r_1, c_1}(\bar{w}_1, w_1) \cdots H_{r_N, c_N}(\bar{w}_N, w_N) \right| \\
& \leq \exp\left(\sum_{i=1}^N |w_i|^2\right) \sum_K \left\{ \prod_{1 \leq j, \ell \leq N} \frac{(|h_{j,\ell}|)^{k_{j,\ell}} \prod_{i=1}^N r_i!}{k_{j,\ell}!} \right\}^{\frac{1}{2}} \left\{ \prod_{1 \leq j, \ell \leq N} \frac{(|h_{j,\ell}|)^{k_{j,\ell}} \prod_{i=1}^N c_i!}{k_{j,\ell}!} \right\}^{\frac{1}{2}} \\
& \leq \exp\left(\sum_{i=1}^N |w_i|^2\right) \left\{ \sum_K \prod_{1 \leq j, \ell \leq N} \frac{\|H\|_{\infty}^{k_{j,\ell}} \prod_{i=1}^N r_i!}{k_{j,\ell}!} \right\} \left\{ \sum_K \prod_{1 \leq j, \ell \leq N} \frac{\|H\|_{\infty}^{k_{j,\ell}} \prod_{i=1}^N c_i!}{k_{j,\ell}!} \right\}
\end{aligned}$$

by applying the Cauchy–Schwarz inequality. We observe that

$$\begin{aligned} \sum_K \prod_{1 \leq j, \ell \leq N} \frac{\|H\|_\infty^{k_{j,\ell}} \prod_{i=1}^N r_i!}{k_{j,\ell}!} &= \sum_K \prod_{i=1}^N \binom{r_i}{k_{i,1}, \dots, k_{i,N}} \|H\|_\infty^{\sum_{\ell=1}^N k_{i,\ell}} \\ &= \prod_{i=1}^N \sum_{r_i \geq 0} (N\|H\|_\infty)^{r_i} = (1 - N\|H\|_\infty)^{-N} \end{aligned}$$

and

$$\begin{aligned} \sum_K \prod_{1 \leq j, \ell \leq N} \frac{\|H\|_\infty^{k_{j,\ell}} \prod_{i=1}^N c_i!}{k_{j,\ell}!} &= \sum_K \prod_{i=1}^N \binom{c_i}{k_{1,i}, \dots, k_{N,i}} \|H\|_\infty^{\sum_{\ell=1}^N k_{\ell,i}} \\ &= \prod_{i=1}^N \sum_{c_i \geq 0} (N\|H\|_\infty)^{c_i} = (1 - N\|H\|_\infty)^{-N}, \end{aligned}$$

then,

$$\begin{aligned} &\left| \sum_K \prod_{1 \leq j, \ell \leq N} \frac{(h_{j,\ell})^{k_{j,\ell}}}{k_{j,\ell}!} H_{r_1, c_1}(\overline{w_1}, w_1) \cdots H_{r_N, c_N}(\overline{w_N}, w_N) \right| \\ &\leq \exp \left(\sum_{i=1}^N |w_i|^2 \right) (1 - N\|H\|_\infty)^{-2N}. \end{aligned}$$

Hence the series on the right-hand side in (1.10) converges uniformly and absolutely for W in any compact subset of \mathbb{C}^N and H in any compact subset of B .

Clearly, $\det(I_N + H)$ is a polynomial in variables $h_{j,\ell}$. For $H \in \mathbb{C}^N$ we have [13]

$$\|H^*\|_2 = \|H\|_2 \leq \sqrt{N} \|H\|_\infty < \frac{1}{\sqrt{N}} \leq 1.$$

Then

$$H(I_N + H)^{-1} = \sum_{m=1}^{\infty} (-1)^{m-1} H^m$$

converges in norm $\|\cdot\|_2$, and

$$\|H(I_N + H)^{-1}\|_2 \leq \frac{\sqrt{N} \|H\|_\infty}{1 - \sqrt{N} \|H\|_\infty}, \quad |W^* H(I_N + H)^{-1} W| \leq \frac{\sqrt{N} \|H\|_\infty}{1 - \sqrt{N} \|H\|_\infty} \|W\|^2.$$

Consequently, $\exp(W^*H(I_N + H)^{-1}W)$ also converges absolutely and uniformly for W in any compact subset of \mathbb{C}^N and H in any compact subset of B . Let

$$\begin{aligned} F(H, W) &= \exp\left(W^*H(I_N + H)^{-1}W\right) - \det(I_N + H) \\ &\times \sum_K \prod_{1 \leq j, \ell \leq N} \frac{(h_{j,\ell})^{k_{j,\ell}}}{k_{j,\ell}!} H_{r_1, c_1}(\overline{w_1}, w_1) \cdots H_{r_N, c_N}(\overline{w_N}, w_N). \end{aligned}$$

Then for any fixed $W \in \mathbb{C}^N$, $F(H, W)$ is analytic in variables $h_{j,k}$ in B , and $F(H_0, W) = 0$ for any Hermitian matrix $H_0 \in B$ by Theorem 1.2.

Let us introduce a new coordinate system $u_{j,j}$, $u_{\ell,k}$, $v_{\ell,k}$, $1 \leq j, k, \ell \leq N$, $\ell < k$ such that

$$h_{j,j} = u_{j,j}, \quad h_{\ell,k} = u_{\ell,k} + iv_{\ell,k}, \quad \ell < k, \quad h_{k,\ell} = u_{\ell,k} - iv_{\ell,k}, \quad \ell > k.$$

Since this is an invertible linear transformation, any function analytic in $h_{j,k}$, $1 \leq j, k \leq N$ is also analytic in $u_{j,j}$, $u_{\ell,k}$, $v_{\ell,k}$, $1 \leq j, k, \ell \leq N$, $\ell < k$ and vice versa. Furthermore, for any Hermitian matrix $H_0 \in B$ and $\delta_{j,k} > 0$, $1 \leq j, k \leq N$ such that (1.8) and (1.9) and $D(H_0, \delta) \subset B$ are satisfied, then $F(H, W)$ can be expanded into a convergent power series in variables $u_{j,j}$, $u_{\ell,k}$, $v_{\ell,k}$, $1 \leq j, k, \ell \leq N$, $\ell < k$ at $u_{j,j}^{(0)}$, $u_{\ell,k}^{(0)}$, $v_{\ell,k}^{(0)}$, $1 \leq j, k, \ell \leq N$, $\ell < k$ on $D(H_0, \delta)$. Clearly, $D(H_0, \delta)$ contains the following set S :

$$-\delta_{j,j} < u_{j,j} - u_{j,j}^{(0)} < \delta_{j,j}, \quad -\delta_{\ell,k} < u_{\ell,k} - u_{\ell,k}^{(0)} < \delta_{\ell,k}, \quad -\delta_{\ell,k} < v_{\ell,k} - v_{\ell,k}^{(0)} < \delta_{\ell,k},$$

where $1 \leq j, k, \ell \leq N$, $\ell < k$, and H is Hermitian on S . From Theorem 1.2 we know that $F(H, W) = 0$ on S . Hence, all the coefficients in this power series expansion of $F(H, W)$ in variables $u_{j,j}$, $u_{\ell,k}$, $v_{\ell,k}$, $1 \leq j, k, \ell \leq N$, $\ell < k$ at $u_{j,j}^{(0)}$, $u_{\ell,k}^{(0)}$, $v_{\ell,k}^{(0)}$, $1 \leq j, k, \ell \leq N$, $\ell < k$ must be zeros. Thus, $F(H, W) = 0$ holds on $D(H_0, \delta)$, which is the same as (1.10) in $D(H_0, \delta)$. \square

We now come to the proof of Corollary 1.4.

Proof of Corollary 1.4. Let $W = (\rho_1 e^{i\theta_1}, \dots, \rho_N e^{i\theta_N})^T$ such that $\rho_j > 0$, $\theta_j \in \mathbb{R}$ for

$j = 1, \dots, N$ in (1.6) to get

$$\begin{aligned}
& \det(I_N + H)^{-1} \exp(W^* H (I_N + H)^{-1} W) \\
&= \sum_K \prod_{1 \leq m, n \leq N} \frac{(h_{m,n})^{k_{m,n}}}{k_{m,n}!} H_{r_1, c_1}(\overline{w_1}, w_1) \cdots H_{r_N, c_N}(\overline{w_N}, w_N) \\
&= \sum_K \prod_{1 \leq m, n \leq N} \frac{(h_{m,n})^{k_{m,n}}}{k_{m,n}!} H_{r_1, c_1}(\rho_1 e^{i\theta_1}, \rho_1 e^{-i\theta_1}) \cdots H_{r_N, c_N}(\rho_N e^{i\theta_N}, \rho_N e^{-i\theta_N}) \\
&= \sum_K \prod_{m=1}^N \prod_{n=1}^N (-h_{m,n})^{k_{m,n}} \binom{c_m}{k_{1,m}, \dots, k_{N,m}} L_{c_m}^{(r_m - c_m)}(\rho_m^2) (\rho_m e^{i\theta_m})^{r_m - c_m}.
\end{aligned}$$

This establishes 1.11.

To prove 1.12, we let

$$H = \begin{pmatrix} 0 & a \\ b & 0 \end{pmatrix}, \quad W = \begin{pmatrix} x \\ y \end{pmatrix}, \quad |a|, |b| < \frac{1}{4}, \quad a, b, x, y \in \mathbb{R},$$

in (1.11), then

$$\begin{aligned}
& \det(I_N + H)^{-1} \exp(W^* H (I_N + H)^{-1} W) = \frac{\exp\left(-\frac{abx^2 - xy(a+b) + aby^2}{1-ab}\right)}{1-ab} \\
&= \sum_{k_{1,2}, k_{2,1}=0}^{\infty} (-a)^{k_{1,2}} x^{k_{2,1}-k_{1,2}} L_{k_{1,2}}^{(k_{2,1}-k_{1,2})}(x^2) (-b)^{k_{2,1}} y^{k_{1,2}-k_{2,1}} L_{k_{2,1}}^{(k_{1,2}-k_{2,1})}(y^2) \\
&= \sum_{j=0}^{\infty} (ab)^j L_j^{(0)}(x^2) L_j^{(0)}(y^2) + \sum_{0 \leq j < k < \infty} (-a)^j x^{k-j} L_j^{(k-j)}(x^2) (-b)^k y^{j-k} L_k^{(j-k)}(y^2) \\
&\quad + \sum_{0 \leq k < j < \infty} (-a)^j x^{k-j} L_j^{(k-j)}(x^2) (-b)^k y^{j-k} L_k^{(j-k)}(y^2) \\
&= \sum_{j=0}^{\infty} (ab)^j L_j^{(0)}(x^2) L_j^{(0)}(y^2) + \sum_{0 \leq j < k < \infty} \frac{j! a^j b^k}{k!} (xy)^{k-j} L_j^{(k-j)}(x^2) L_j^{(k-j)}(y^2) \\
&\quad + \sum_{0 \leq k < j < \infty} \frac{k! a^j b^k}{j!} (xy)^{j-k} L_k^{(j-k)}(x^2) L_k^{(j-k)}(y^2) \\
&= \sum_{j=0}^{\infty} (ab)^j L_j^{(0)}(x^2) L_j^{(0)}(y^2) + \sum_{0 \leq j < k < \infty} \frac{a^j b^k}{j! k!} (xy)^{k+j} C_j(k; x^2) C_j(k; y^2) \\
&\quad + \sum_{0 \leq k < j < \infty} \frac{a^j b^k}{j! k!} (xy)^{j+k} C_k(j; x^2) C_k(j; y^2),
\end{aligned}$$

where we have applied

$$L_n^{(x-n)}(a) = \frac{(-a)^n}{n!} C_n(x; a).$$

Hence, for $x, y \neq 0$, for u, v sufficiently small, we let

$$a = \frac{u}{xy}, \quad b = \frac{v}{xy}$$

to get

$$\begin{aligned} & \frac{x^2y^2}{x^2y^2 - uv} \exp\left(-\frac{x^2y^2(x^2uv - x^2y^2(u+v) + y^2uv)}{x^2y^2 - uv}\right) \\ &= \sum_{j=0}^{\infty} \left(\frac{uv}{x^2y^2}\right)^j L_j^{(0)}(x^2) L_j^{(0)}(y^2) + \sum_{0 \leq j < k < \infty} \frac{u^j v^k}{j! k!} C_j(k; x^2) C_j(k; y^2) \\ &+ \sum_{0 \leq j < k < \infty} \frac{u^k v^j}{j! k!} C_j(k; x^2) C_j(k; y^2) = \sum_{j=0}^{\infty} \left(\frac{uv}{x^2y^2}\right)^j L_j^{(0)}(x^2) L_j^{(0)}(y^2) \\ &+ \sum_{0 \leq j < k < \infty} \frac{(u^j v^k + u^k v^j)}{j! k!} C_j(k; x^2) C_j(k; y^2). \end{aligned}$$

By

$$\begin{aligned} \sum_{j=0}^{\infty} \left(\frac{uv}{x^2y^2}\right)^j L_j^{(0)}(x^2) L_j^{(0)}(y^2) &= \frac{x^2y^2}{x^2y^2 - uv} \exp\left(-\frac{uv(x^2 + y^2)}{x^2y^2 - uv}\right) {}_0F_1\left(-, 1, \frac{uvx^3y^3}{(x^2y^2 - uv)^2}\right) \\ &= \frac{x^2y^2}{x^2y^2 - uv} \exp\left(-\frac{uv(x^2 + y^2)}{x^2y^2 - uv}\right) I_0\left(2 \frac{\sqrt{uvx^3y^3}}{x^2y^2 - uv}\right) \end{aligned}$$

we have

$$\begin{aligned} & \frac{xy}{xy - uv} \exp\left(-\frac{xy(xuv - xy(u+v) + yuv)}{xy - uv}\right) - \frac{xy}{xy - uv} \exp\left(-\frac{uv(x^2 + y^2)}{xy - uv}\right) I_0\left(2 \frac{\sqrt{uv(xy)^{3/2}}}{xy - uv}\right) \\ &= \sum_{0 \leq j < k < \infty} \frac{(u^j v^k + u^k v^j)}{j! k!} C_j(k; x) C_j(k; y). \end{aligned}$$

□

3. Miscellaneous results

Theorem 3.1. Let $w = r + is$ with $r, s \in \mathbb{R}$ and $z_1, z_2 \in \mathbb{C}$, then we have the moment integral representation

$$e^{-z_1 z_2} H_{m,n}(z_1, z_2) = \frac{1}{\pi i^{m+n}} \int_{\mathbb{R}^2} \bar{w}^m w^n \exp \{-w\bar{w} + iz_1 w + iz_2 \bar{w}\} dr ds. \quad (3.1)$$

In particular we have

$$\begin{aligned} & e^{-r_1 r_2 e^{i(\theta_1 + \theta_2)}} H_{m,n}(r_1 e^{i\theta_1}, r_2 e^{i\theta_2}) \\ &= \frac{i}{2^{m+n+1} \sqrt{\pi}} \int_0^{2\pi} H_{n+m+1} \left(\frac{r_1 e^{i(\theta_1 + \phi)} + r_2 e^{i(\theta_2 - \phi)}}{2} \right) \\ & \times \exp \left\{ -\frac{(r_1 e^{i(\theta_1 + \phi)} + r_2 e^{i(\theta_2 - \phi)})^2}{4} + i(n-m)\phi \right\} d\phi, \end{aligned} \quad (3.2)$$

$$\begin{aligned} & e^{-r^2} H_{m,n}(re^{i\theta}, re^{-i\theta}) = \frac{ie^{i(m-n)\theta}}{2^{m+n+1} \sqrt{\pi}} \\ & \times \int_0^{2\pi} H_{n+m+1}(r \cos \phi) \exp(-r^2 \cos^2 \phi + i(n-m)\phi) d\phi, \end{aligned} \quad (3.3)$$

$$H_{m,n}(w_1, w_2) = \begin{cases} (-1)^n n! w_1^{m-n} L_n^{(m-n)}(w_1 w_2), & m \geq n \\ (-1)^m m! w_2^{n-m} L_m^{(n-m)}(w_1 w_2), & n \geq m \end{cases} \quad (3.4)$$

and

$$\begin{aligned} & \frac{i}{2\sqrt{\pi}} \int_0^{2\pi} H_{n+m+1}(r \cos \phi) \exp(-r^2 \cos^2 \phi + i(n-m)\phi) d\phi \\ &= (-1)^n 2^{m+n} n! r^{m-n} e^{-r^2} L_n^{(m-n)}(r^2). \end{aligned} \quad (3.5)$$

Proof. Let

$$a_{m,n} = \frac{1}{\pi} \int_{\mathbb{R}^2} \bar{w}^m w^n \exp \{-w\bar{w} + iz_1 w + iz_2 \bar{w}\} dr ds$$

then,

$$\sum_{m,n=0}^{\infty} a_{m,n} \frac{u^m}{m!} \frac{v^n}{n!} = \exp(-z_1 z_2) \exp(iz_1 u + iz_2 v + uv).$$

Comparing the above expression with the generating function (1.2) proves Equation (3.1). Let $w = \rho e^{i\phi}$, $z_1 = r_1 e^{i\theta_1}$ and $z_2 = r_2 e^{i\theta_2}$ in (3.1), then

$$\begin{aligned}
& e^{-z_1 z_2} H_{m,n}(z_1, z_2) \\
&= \frac{1}{\pi^{im+n}} \int_{\mathbb{R}^2} \bar{w}^m w^n \exp \{-w\bar{w} + iz_1 w + iz_2 \bar{w}\} dr ds \\
&= \frac{1}{\pi^{im+n}} \int_0^{2\pi} e^{i(n-m)\phi} \left\{ \int_0^\infty \rho^{m+n+1} \exp(-\rho^2 + i\rho(e^{i\phi} z_1 + e^{-i\phi} z_2)) d\rho \right\} d\phi \\
&= \frac{i}{2^{m+n+1} \sqrt{\pi}} \int_0^{2\pi} H_{n+m+1} \left(\frac{r_1 e^{i(\theta_1+\phi)} + r_2 e^{i(\theta_2-\phi)}}{2} \right) \\
&\quad \times \exp \left\{ -\frac{(r_1 e^{i(\theta_1+\phi)} + r_2 e^{i(\theta_2-\phi)})^2}{4} + i(n-m)\phi \right\} d\phi,
\end{aligned}$$

where we used a variant of (1.13) in the last step. This gives (3.2). Let $z_1 = re^{i\theta}$, $z_2 = re^{-i\theta}$ in (3.2) to get,

$$\begin{aligned}
& e^{-r^2} H_{m,n}(re^{i\theta}, re^{-i\theta}) \\
&= \frac{i}{2^{m+n+1} \sqrt{\pi}} \int_0^{2\pi} H_{n+m+1}(r \cos(\theta + \phi)) \exp(-r^2 \cos^2(\theta + \phi) + i(n-m)\phi) d\phi,
\end{aligned}$$

and we establish (3.3). The identification (3.4) is known and follows from (1.1) and the representation of a Laguerre polynomial as a confluent hypergeometric polynomial. Finally (3.5) follows from (3.3) and (3.4). \square

The next result develops mixed relations involving 2D Hermite polynomials and Hermite polynomials.

Theorem 3.2. Let $w_1, w_2, z \in \mathbb{C}$ with $z \neq 0$, $\rho > 0$ and $\theta \in \mathbb{R}$, then we have

$$H_n\left(\frac{w_1 + w_2}{2}\right) = z^n \sum_{j=0}^n \binom{n}{j} H_{j,n-j}\left(zw_1, \frac{w_2}{z}\right) z^{-2j}, \quad (3.6)$$

$$H_n\left(\frac{\rho(z+z^{-1})}{2}\right) = \frac{n!}{(-\rho z)^n} \sum_{j=0}^n \frac{(-\rho^2 z^2)^j}{j!} L_{n-j}^{(2j-n)}(\rho^2), \quad (3.7)$$

$$H_n(\rho \cos \theta) = \frac{n!}{(-\rho e^{i\theta})^n} \sum_{j=0}^n \frac{(-\rho^2 e^{2i\theta})^j}{j!} L_{n-j}^{(2j-n)}(\rho^2), \quad (3.8)$$

$$\int_0^{2\pi} H_n(\rho \cos \theta) e^{-ik\theta} d\theta = \begin{cases} 0 & 2 \nmid (n+k) \\ \frac{2\pi n! (-1)^{(n-k)/2} \rho^k}{(\frac{n+k}{2})!} L_{\frac{n-k}{2}}^{(k)}(\rho^2) & 2 \mid (n+k) \end{cases} \quad (3.9)$$

and

$$\int_0^{2\pi} (H_n(\rho \cos \theta))^2 \frac{d\theta}{2\pi} = \frac{(n!)^2}{\rho^{2n}} \sum_{j=0}^n \frac{\rho^{2j}}{j! j!} (L_{n-j}^{(2j-n)}(\rho^2))^2. \quad (3.10)$$

Proof. Let $u = \alpha^2 t$, $v = \beta^2 t$, $z_1 = 2\beta w_1/\alpha$, $z_2 = 2\alpha w_2/\beta$ in (1.2) to obtain

$$\begin{aligned} & \sum_{m,n=0}^{\infty} H_{m,n}\left(\frac{2\beta w_1}{\alpha}, \frac{2\alpha w_2}{\beta}\right) \frac{\alpha^{2m}}{m!} \frac{\beta^{2n}}{n!} t^{m+n} \\ &= \exp(-(\alpha \beta t)^2 + 2\alpha \beta t (w_1 + w_2)) \\ &= \sum_{n=0}^{\infty} H_n(w_1 + w_2) \frac{(\alpha \beta t)^n}{n!} \end{aligned}$$

and (3.6) follows by equating like coefficients of powers of t . We use the parameter identification $z = \beta/\alpha$, $w_1 = e^{i\theta} \rho \alpha / 2\beta$, $w_2 = e^{-i\theta} \rho \beta / 2\alpha$ in (3.6) and find that

$$\begin{aligned} H_n\left(\frac{\rho \alpha}{2\beta} e^{i\theta} + \frac{\rho \beta}{2\alpha} e^{-i\theta}\right) &= \left(\frac{\beta}{\alpha}\right)^n \sum_{j=0}^n \binom{n}{j} H_{j,n-j}(\rho e^{i\theta}, \rho e^{-i\theta}) \left(\frac{\alpha}{\beta}\right)^{2j} \\ &= \left(\frac{\beta}{\alpha}\right)^n \sum_{j=0}^n \binom{n}{j} (-1)^{n-j} (n-j)! (\rho e^{i\theta})^{2j-n} L_{n-j}^{(2j-n)}(\rho^2) \left(\frac{\alpha}{\beta}\right)^{2j} \\ &= \left(\frac{\beta}{\alpha}\right)^n \frac{n!}{(-\rho e^{i\theta})^n} \sum_{j=0}^n \frac{(-\rho^2 e^{2i\theta})^j}{j!} L_{n-j}^{(2j-n)}(\rho^2) \left(\frac{\alpha}{\beta}\right)^{2j}, \end{aligned}$$

and (3.7) follows. Equations (3.8)–(3.9) follow by taking $z = e^{i\theta}$ in (3.7) and applying the Fourier orthogonality. \square

We list more properties for the 2D Hermite polynomials:

Theorem 3.3. *Let $z_1, z_2, w_1, w_2 \in \mathbb{C}$ and m, n are negative integers, then we have*

$$H_{m,n}(w_1 - iw_2, w_1 + iw_2) = \frac{i^{m-n}}{2^{m+n}} \sum_{j,k=0}^{\min(m,n)} \binom{m}{j} \binom{n}{k} \frac{H_{j+k}(w_1) H_{m+n-j-k}(w_2)}{i^{j-k}}, \quad (3.11)$$

$$\frac{H_{m,n}(z_1 + w_1, z_2 + w_2)}{e^{w_1 w_2 + z_1 w_2 + z_2 w_1}} = \sum_{j,k=0}^{\infty} \frac{(-w_1)^j (-w_2)^k}{j! k!} H_{m+k, n+j}(z_1, z_2), \quad (3.12)$$

$$H_{m,n}(0, 0) = \delta_{m,n} (-1)^n n!, \quad (3.13)$$

Proof. From

$$\begin{aligned} e^{-z_1 z_2} H_{m,n}(z_1, z_2) &= \frac{1}{\pi i^{m+n}} \int_{\mathbb{R}^2} \bar{w}^m w^n \\ &\quad \times \exp \{-w\bar{w} + iz_1 w + iz_2 \bar{w}\} dr ds \\ &= \frac{1}{\pi i^{m+n}} \sum_{j,k=0}^{\infty} \binom{m}{j} \binom{n}{k} i^{m-n-j+k} \int_{\mathbb{R}^2} r^{j+k} s^{m+n-j-k} \\ &\quad \times \exp \{-r^2 - s^2 + ir(z_1 + z_2) + is(iz_1 - iz_2)\} dr ds \\ &= \frac{1}{\pi i^{m+n}} \sum_{j,k=0}^{\infty} \binom{m}{j} \binom{n}{k} i^{m-n-j+k} \int_{\mathbb{R}} r^{j+k} e^{-r^2 + ir(z_1 + z_2)} dr \\ &\quad \times \int_{\mathbb{R}} s^{m+n-j-k} e^{-s^2 + is(iz_1 - iz_2)} dr \\ &= \frac{i^{m-n}}{2^{m+n}} \sum_{j,k=0}^{\infty} \binom{m}{j} \binom{n}{k} i^{-j+k} H_{j+k} \left(\frac{z_1 + z_2}{2} \right) H_{m+n-j-k} \left(\frac{z_1 - z_2}{2} i \right). \end{aligned}$$

or

$$H_{m,n}(z_1, z_2) = \frac{i^{m-n}}{2^{m+n}} \sum_{j,k=0}^{\infty} \binom{m}{j} \binom{n}{k} i^{-j+k} H_{j+k} \left(\frac{z_1 + z_2}{2} \right) H_{m+n-j-k} \left(\frac{z_1 - z_2}{2} i \right).$$

Let $z_1 = w_1 - iw_2$ and $z_2 = w_1 + iw_2$ we get (3.11).

From

$$\begin{aligned}
& e^{-(z_1+w_1)(z_2+w_2)} H_{m,n}(z_1+w_1, z_2+w_2) \\
&= \frac{1}{\pi i^{m+n}} \sum_{j,k=0}^{\infty} \frac{w_1^j w_2^k t^{j+k}}{j! k!} \int_{\mathbb{R}^2} \bar{x}^{m+k} x^{n+j} \exp \{-x\bar{x} + iz_1x + iz_2\bar{x}\} dr ds \\
&= \sum_{j,k=0}^{\infty} \frac{w_1^j w_2^k (-1)^{j+k}}{j! k!} e^{-z_1 z_2} H_{m+k, n+j}(z_1, z_2),
\end{aligned}$$

that is (3.12). Formula (3.13) follows from (1.2). \square

4. q -analogues

We follow the notation for q -shifted factorials and q -series as in [2], [9] and [16]. The 2D q -Hermite polynomials are defined by [15]

$$\frac{H_{m,n}(z_1, z_2 | q)}{(q; q)_m (q; q)_n} = \sum_{k=0}^{m \wedge n} \frac{z_1^{m-k} z_2^{n-k}}{(q; q)_{m-k} (q; q)_{n-k} (q; q)_k}. \quad (4.1)$$

In [15] we also proved the generating function

$$\sum_{m,n=0}^{\infty} H_{m,n}(z_1, z_2 | q) \frac{u^m v^n}{(q; q)_m (q; q)_n} = \frac{(uv; q)_{\infty}}{(uz_1, v z_2; q)_{\infty}}. \quad (4.2)$$

We shall also use the Askey–Wilson integral [2, 9, 16]

$$\int_0^{\pi} \frac{\left(e^{2i\theta}, e^{-2i\theta}; q\right)_{\infty}}{\prod_{j=1}^4 \left(t_j e^{i\theta}, t_j e^{-i\theta}; q\right)_{\infty}} d\theta = \frac{2\pi (t_1 t_2 t_3 t_4; q)_{\infty}}{(q; q)_{\infty} \prod_{1 \leq j < k \leq 4} (t_j t_k; q)_{\infty}}, \quad (4.3)$$

which holds for $\max \{|t_j| : 1 \leq j \leq 4\} < 1$. The trigonometric moments of the q -Hermite weight function are [16]

$$\int_0^{\pi} e^{2ij\theta} \left(e^{2i\theta}, e^{-2i\theta}; q\right)_{\infty} \frac{d\theta}{2\pi} = \frac{(-1)^j}{(q; q)_{\infty}} \left(q^{\binom{j}{2}} + q^{\binom{-j}{2}}\right). \quad (4.4)$$

Theorem 4.1. For $|rz_1| < 1$, $|sz_2| < 1$, we have the generating function

$$\begin{aligned} & \frac{(rs, rs; q)_\infty}{(z_1 z_2 rs; q)_\infty} \\ &= \sum_{m_1, m_2=0}^{\infty} \sum_{n_1, n_2=0}^{\infty} \frac{H_{m_1, n_1}(z_1, z_2 | q) H_{m_2, n_2}(z_1, z_2 | q)}{(q; q)_{m_1} (q; q)_{m_2} (q; q)_{n_1} (q; q)_{n_2}} r^{m_1+m_2} s^{n_1+n_2} \\ & \quad \times \frac{\left(q^{\binom{(m_1+n_2-n_1-m_2)/2}{2}} + q^{\binom{(n_1+m_2-m_1-n_2)/2}{2}} \right)}{(-1)^{(n_1+n_2-m_1-m_2)/2}}, \end{aligned} \quad (4.5)$$

where the summation is over all the nonnegative integers such that $m_1+m_2-n_1-n_2$ is even.

Proof. Multiply the generating functions (4.2) with the z variable being z_1, z_2, z_1, z_2 and set $u_1 = re^{i\theta}, v_1 = se^{-i\theta}, u_2 = re^{-i\theta}, v_2 = se^{i\theta}$. This gives

$$\begin{aligned} & \frac{(rs, rs; q)_\infty}{(rz_1 e^{i\theta}, rz_1 e^{-i\theta}, sz_2 e^{i\theta}, sz_2 e^{-i\theta}; q)_\infty} \\ &= \sum_{m_1, m_2, n_1, n_2=0}^{\infty} \frac{H_{m_1, n_1}(z_1, z_2 | q) H_{m_2, n_2}(z_1, z_2 | q)}{(q; q)_{m_1} (q; q)_{m_2} (q; q)_{n_1} (q; q)_{n_2}} r^{m_1+m_2} s^{n_1+n_2} e^{i\theta(n_1+m_2-m_1-n_2)}. \end{aligned}$$

Multiply the above generating function by $(e^{2i\theta}, e^{-2i\theta}; q)_\infty / (2\pi)$ and integrate over θ in $[-\pi, \pi]$ the apply the case $t_3 = t_4 = 0$ of the Askey–Wilson integral (4.3) we find that

$$\begin{aligned} & 2 \frac{(rs, rs; q)_\infty}{(q; z_1 z_2 rs; q)_\infty} \\ &= \sum_{m_1, m_2=0}^{\infty} \sum_{n_1, n_2=0}^{\infty} \frac{H_{m_1, n_1}(z_1, z_2 | q) H_{m_2, n_2}(z_1, z_2 | q)}{(q; q)_{m_1} (q; q)_{m_2} (q; q)_{n_1} (q; q)_{n_2}} r^{m_1+m_2} s^{n_1+n_2} \\ & \quad \times \int_{-\pi}^{\pi} \frac{(e^{2i\theta}, e^{-2i\theta}; q)_\infty}{e^{i\theta(n_1+m_2-m_1-n_2)}} \frac{d\theta}{2\pi}, \end{aligned}$$

which vanishes unless $m_2 + n_1 - m_1 - n_2$ is even, in which case we get

$$\begin{aligned} & \frac{(rs, rs; q)_\infty}{(q; z_1 z_2 rs; q)_\infty} \\ = & \sum_{m_1, m_2=0}^{\infty} \sum_{n_1, n_2=0}^{\infty} \frac{H_{m_1, n_1}(z_1, z_2 | q) H_{m_2, n_2}(z_1, z_2 | q)}{(q; q)_{m_1} (q; q)_{m_2} (q; q)_{n_1} (q; q)_{n_2}} r^{m_1+m_2} s^{n_1+n_2} \\ & \times \frac{\left(q^{\binom{(m_1+n_2-n_1-m_2)/2}{2}} + q^{\binom{(n_1+m_2-m_1-n_2)/2}{2}} \right)}{(q; q)_\infty (-1)^{(n_1+n_2-m_1-m_2)/2}}, \end{aligned}$$

where the summation is over all the nonnegative integers such that $m_1 + m_2 - n_1 - n_2$ is even. \square

Theorem 4.2. Let $|x_1 z_1|, |x_1 z_2|, |x_1 z_3|, |x_1 z_4| < 1$, then

$$\begin{aligned} & \frac{(r_1 s_1, r_1 s_1, r_2 s_2, r_2 s_2, r_1 s_1, r_2 s_2 z_1 z_2 z_3 z_4; q)_\infty}{(r_1 r_2 z_1 z_2, r_1 r_2 z_1 z_3, r_1 s_2 z_1 z_4, r_2 s_1 z_2 z_3, s_1 s_2 z_2 z_4, r_2 s_2 z_3 z_4, q)_\infty} \quad (4.6) \\ = & \sum_{m_j, n_j \geq 0, 1 \leq j \leq 4}^{\infty} \frac{H_{m_1, n_1}(z_1, z_2 | q) H_{m_2, n_2}(z_1, z_2 | q)}{(q; q)_{m_1} (q; q)_{n_1} (q; q)_{m_2} (q; q)_{n_2}} r_1^{m_1+m_2} r_2^{m_3+m_4} s_1^{n_1+n_2} s_2^{n_3+n_4} \\ & \times \frac{H_{m_3, n_3}(z_3, z_4 | q) H_{m_4, n_4}(z_3, z_4 | q)}{(q; q)_{m_3} (q; q)_{n_3} (q; q)_{m_4} (q; q)_{n_4}} (-1)^M \left[q^{\binom{M}{2}} + q^{\binom{-M}{2}} \right] \quad (4.7) \end{aligned}$$

where the summation is over all nonnegative integers m_j, n_j , $1 \leq j \leq 4$ such that $m_1 + n_2 + m_3 + n_4 - n_1 - m_2 - n_3 - m_4$ is even and $= 2M$.

Proof. Again we start with four cases of the generating function (4.2) with parameters (u_j, v_j) , $1 \leq j \leq 4$ and variables $z_1, z_2, z_1, z_2, z_3, z_4, z_3, z_4$, where $u_1 = r_1 e^{i\theta}$, $v_1 = s_1 e^{-i\theta}$, $u_2 = r_1 e^{-i\theta}$, $v_2 = s_1 e^{i\theta}$, $u_3 = r_2 e^{i\theta}$, $v_3 = s_2 e^{-i\theta}$, $u_4 = r_2 e^{-i\theta}$, $v_4 = s_2 e^{i\theta}$. We multiply the four right-hand sides of (4.2) by $(e^{2i\theta}, e^{-2i\theta}; q)_\infty / (2\pi)$ and integrate over $[-\pi, \pi]$. The use of the Askey–Wilson integral shows that the result is

$$\frac{2 (r_1 s_1, r_1 s_1, r_2 s_2, r_2 s_2, r_1 s_1, r_2 s_2 z_1 z_2 z_3 z_4; q)_\infty}{(q, r_1 r_2 z_1 z_2, r_1 r_2 z_1 z_3, r_1 s_2 z_1 z_4, r_2 s_1 z_2 z_3, s_1 s_2 z_2 z_4, r_2 s_2 z_3 z_4, q)_\infty}.$$

The rest of the proof is similar to the proof of Theorem 4.1 and will be omitted. \square

References

- [1] S. Twareque Ali, F. Bagarello, and G. Honnouvo. Modular structures on trace class operators and applications to Landau levels. *J. Phys. A*, 43(10):105202, 17, 2010.
- [2] George E. Andrews, Richard Askey, and Ranjan Roy. *Special functions*, volume 71 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, UK, 1999.
- [3] Tom M. Apostol. *Mathematical analysis*. Addison-Wesley Publishing Co., Reading, Mass.-London-Don Mills, Ont., second edition, 1974.
- [4] Richard Bellman. *Introduction to matrix analysis*, volume 19 of *Classics in Applied Mathematics*. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 1997. Reprint of the second (1970) edition, With a foreword by Gene Golub.
- [5] L. Carlitz. A set of polynomials in three variables. *Houston J. Math.*, 4(1):11–33, 1978.
- [6] Nicolae Cotfas, Jean Pierre Gazeau, and Katarzyna Górska. Complex and real Hermite polynomials and related quantizations. *J. Phys. A*, 43(30):305304, 14, 2010.
- [7] Charles F. Dunkl and Yuan Xu. *Orthogonal polynomials of several variables*. Encyclopedia of Mathematics and its Applications. Cambridge University Press, Cambridge, UK, second edition, 2014.
- [8] Dominique Foata. Some Hermite polynomial identities and their combinatorics. *Adv. in Appl. Math.*, 2(3):250–259, 1981.
- [9] George Gasper and Mizan Rahman. *Basic hypergeometric series*, volume 96 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, second edition, 2004. With a foreword by Richard Askey.
- [10] Allal Ghanmi. A class of generalized complex Hermite polynomials. *J. Math. Anal. Appl.*, 340(2):1395–1406, 2008.
- [11] Allal Ghanmi. Operational formulae for the complex Hermite polynomials $H_{p,q}(z, \bar{z})$. *Integral Transforms Spec. Funct.*, 24(11):884–895, 2013.

- [12] N. R. Goodman. Statistical analysis based on a certain multivariate complex Gaussian distribution. (An introduction). *Ann. Math. Statist.*, 34:152–177, 1963.
- [13] Roger A. Horn and Charles R. Johnson. *Matrix analysis*. Cambridge University Press, Cambridge, UK, second edition, 2013.
- [14] Abdelkader Intissar and Ahmed Intissar. Spectral properties of the Cauchy transform on $L_2(\mathbb{C}, e^{-|z|^2} \lambda(z))$. *J. Math. Anal. Appl.*, 313(2):400–418, 2006.
- [15] M. E. H. Ismail and R. Zhang. On some 2D orthogonal q -polynomials. *Trans. Amer. Math. Soc.*, 2016. To appear.
- [16] Mourad E. H. Ismail. *Classical and quantum orthogonal polynomials in one variable*, volume 98 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 2009. With two chapters by Walter Van Assche, With a foreword by Richard A. Askey, Reprint of the 2005 original.
- [17] Mourad E. H. Ismail. Analytic properties of complex Hermite polynomials. *Trans. Amer. Math. Soc.*, 368(2):1189–1210, 2016.
- [18] Mourad E. H. Ismail and Jiang Zeng. A combinatorial approach to the 2D-Hermite and 2D-Laguerre polynomials. *Adv. in Appl. Math.*, 64:70–88, 2015.
- [19] Kiyosi Itô. Complex multiple Wiener integral. *Jap. J. Math.*, 22:63–86 (1953), 1952.
- [20] W. F. Kibble. An extension of a theorem of Mehler’s on Hermite polynomials. *Proc. Cambridge Philos. Soc.*, 41:12–15, 1945.
- [21] J. D. Louck. Extension of the Kibble-Slepian formula for Hermite polynomials using boson operator methods. *Adv. in Appl. Math.*, 2(3):239–249, 1981.
- [22] David Slepian. On the symmetrized Kronecker power of a matrix and extensions of Mehler’s formula for Hermite polynomials. *SIAM J. Math. Anal.*, 3: 606–616, 1972.

- [23] K. Thirulogasanthar, G. Honnouvo, and A. Krzyżak. Coherent states and Hermite polynomials on quaternionic Hilbert spaces. *J. Phys. A*, 43(38):385205, 13, 2010.
- [24] Alfred Wünsche. Laguerre 2D-functions and their application in quantum optics. *J. Phys. A*, 31(40):8267–8287, 1998.
- [25] Alfred Wünsche. Transformations of Laguerre 2D polynomials with applications to quasiprobabilities. *J. Phys. A*, 32(17):3179–3199, 1999.