Stephen M. Barnett

School of Physics and Astronomy, University of Glasgow, Glasgow G12 8QQ, UK E-mail: stephen.barnett@glasgow.ac.uk

# James D. Cresser

Department of Physics and Astronomy, Faculty of Science and Engineering, Macquarie University, NSW 2109, Australia

E-mail: james.cresser@mq.edu.au

#### Sarah Croke

School of Physics and Astronomy, University of Glasgow, Glasgow G12 8QQ, UK

E-mail: sarah.croke@glasgow.ac.uk

Abstract. We analyse the properties of a strongly-damped quantum harmonic oscillator by means of an exact diagonalisation of the full Hamiltonian, including both the oscillator and the reservoir degrees of freedom to which it is coupled. Many of the properties of the oscillator, including its steady-state properties and entanglement with the reservoir can be understood and quantified in terms of a simple probability density, which we may associate with the ground-state frequency spectrum of the oscillator.

PACS numbers: 03.65.Yz, 42.50.Lc

## 1. Introduction

Recent technological advances make it possible to realise simple mechanical devices in the microscopic and nanoscopic regimes, the properties of which are determined by quantum effects [\[1\]](#page-16-0). The existence of these represents a remarkable opportunity for fundamental studies of light-matter interactions [\[2\]](#page-16-1) and also the potential for practical application to quantum communications and information processing [\[3\]](#page-16-2). Yet they present also a challenge to existing methods of analysis, many of which were developed to treat more rapidly oscillating systems with weaker couplings.

The quantum theory of machines is built, to a large extent on the theory of oscillators and strongly coupled oscillators, which are coupled to one or more environments each of which is at a characteristic temperature [\[4,](#page-16-3) [5,](#page-16-4) [6\]](#page-16-5). The behaviour of these quantum systems is governed not simply by average properties but also fluctuations, and has been informed by the development of fluctuation theorems for quantum open systems [\[7,](#page-16-6) [8\]](#page-16-7). The coupling to the environment needs to be handled with some care, however, because of the possibility of quantum coherence and the development of entanglement between the system of interest and its environment, with the result that apparently unphysical behaviours may emerge [\[9\]](#page-16-8). The requirement to treat the coupling to the reservoirs with care provides the incentive to return to the problem of a single strongly-damped oscillator and to treat this model exactly.

There exist at present mathematically and physically acceptable methods for treating strongly damped quantum oscillators, but most of these do not retain the simplicity of their classical counterparts. Our aim is to recover at least some of this simplicity. We do not seek to apply familiar techniques developed for weakly damped systems, but proceed instead by means of an exact diagonlisation of the oscillatorreservoir Hamiltonian. This approach is complementary to that adopted by Philbin who has tackled this problem of the oscillator evolution using Green functions [\[10\]](#page-16-9). We hope that the combination of his work and ours will provide a more complete understanding.

Our procedure produces a single positive function of frequency the integral of which is unity. We can interpret this quantity as the frequency spectrum or probability density for the oscillator in its ground state. The steady-state of the oscillator, its entanglement with the environment, its energy and much of its dynamics can be understood in terms of this probability. Before embarking on this programme, let us pause to review some of the background to our study.

## 1.1. Background

The harmonic oscillator has a special place in physics as one of the simplest and most widely employed of physical models. The reasons for its ubiquity, no doubt, are its simplicity and the fact that it is readily analysed. In the quantum domain, the harmonic oscillator is barely more difficult to treat than its classical counterpart and was one of the first dynamical systems to which Schrödinger applied his equation  $[11]$ . Today, both the classical and quantum forms appear in elementary courses on classical and quantum

mechanics.

The damped harmonic oscillator loses energy as a result of coupling to the surrounding environment. In the classical domain it often suffices to describe this in terms of a simple damping coefficient,  $\gamma$ , and an associated stochastic or Langevin force [\[12\]](#page-16-11),  $F(t)$ , which models the effect of environmental fluctuations on the oscillator [\[13,](#page-16-12) [14,](#page-16-13) [15,](#page-16-14) [16,](#page-16-15) [17\]](#page-16-16). The dynamics is described by a simple linear differential equation of the form

<span id="page-2-0"></span>
$$
\ddot{x} + \gamma \dot{x} + \omega_0^2 x = \frac{F(t)}{m},\tag{1}
$$

where  $m$  is the mass of the oscillating particle. There is no requirement for detailed knowledge of the fluctuating force, which may be considered to have a very short correlation time with a magnitude determined by the requirements of thermodynamic equilibrium.

The damped quantum harmonic oscillator requires that explicit account be taken of the quantum nature of the environmental degrees of freedom [\[18\]](#page-17-0), which are most simply described by an ensemble of harmonic oscillators [\[19\]](#page-17-1). If the damping is very weak, so that  $\gamma \ll \omega_0$ , then we can neglect rapidly oscillating terms in the coupling between the oscillator and the environmental oscillators by making the rotating wave approximation, which corresponds to enforcing the conservation of the total number of vibrational quanta, and then the Born and Markov approximations associated with weak coupling and loss of memory in the reservoir [\[20\]](#page-17-2). This leads to the master equations and Heisenberg-Langevin equations that are ubiquitous, most especially, in quantum optics [\[20,](#page-17-2) [21,](#page-17-3) [22,](#page-17-4) [23,](#page-17-5) [24,](#page-17-6) [25,](#page-17-7) [26,](#page-17-8) [27,](#page-17-9) [28,](#page-17-10) [29,](#page-17-11) [30,](#page-17-12) [31\]](#page-17-13).

If the coupling is somewhat stronger then it may not be possible to make the rotating wave approximation and we then need to retain in the Hamiltonian terms that can create or annihilate a pair of quanta, one in the damped oscillator and, at the same time, one in the environment. This leads to the Caldeira-Leggett model [\[10,](#page-16-9) [32,](#page-17-14) [33,](#page-17-15) [34,](#page-17-16) [35,](#page-17-17) [36,](#page-17-18) [37\]](#page-17-19), which we describe in the the following section, and which has been applied to study a wide variety of quantum open systems [\[27,](#page-17-9) [31\]](#page-17-13). A variant on the model has been applied to the quantum theory of light in dielectric [\[38,](#page-17-20) [39\]](#page-17-21) and magneto-dielectric media [\[40,](#page-17-22) [41,](#page-17-23) [42,](#page-17-24) [43\]](#page-17-25). An important complication that seems to be an inevitable consequence working in this strong-coupling regime is the failure of the Markov approximation; attempts to enforce this approximation lead to a master equation that is unphysical in that there exist initial states for which the dynamics leads to negative probabilities [\[27,](#page-17-9) [44,](#page-17-26) [45,](#page-17-27) [46,](#page-17-28) [47\]](#page-17-29). It is possible to derive a master equation but the resulting equation is one that has within it non-trivial time-dependent coefficients [\[37,](#page-17-19) [48\]](#page-17-30). This time-dependence is a clear signature of the non-Markovian nature of the associated evolution. It seems that this non-Markovian character is an inevitable feature of the strongly-damped quantum harmonic oscillator.



<span id="page-3-0"></span>Figure 1. Representation of a single harmonic oscillator (in red) coupled harmonically to a bath of oscillators (in blue).

# 2. Hamiltonian for the strongly-damped harmonic oscillator

Consider a harmonic oscillator of natural angular frequency  $\omega_0$  that is strongly coupled to a large collection of oscillators, with a range of frequencies, through their respective positions as depicted in figure [1.](#page-3-0) We write the Hamiltonian for the combined oscillatorreservoir system in the form [\[35,](#page-17-17) [37\]](#page-17-19)

$$
\hat{H} = \frac{\hat{p}^2}{2m} + \frac{1}{2}m\omega_0^2 \hat{x}^2 + \sum_{\mu} \left( \frac{\hat{p}_{\mu}^2}{2m_{\mu}} + \frac{1}{2}m\omega_{\mu}^2 \hat{x}_{\mu} \right) - \sum_{\mu} m_{\mu} \omega_{\mu}^2 \lambda_{\mu} \hat{x}_{\mu} \hat{x}.
$$
 (2)

If we complete the square we can rewrite this in a minimal-coupling form to arrive at the alternative form:

$$
\hat{H} = \frac{\hat{p}^2}{2m} + \frac{1}{2}m\Omega_0^2 \hat{x}^2 + \sum_{\mu} \left[ \frac{\hat{p}_{\mu}^2}{2m_{\mu}} + \frac{1}{2}m_{\mu}\omega_{\mu}^2 (\hat{x}_{\mu} - \lambda_{\mu}\hat{x})^2 \right],
$$
\n(3)

where

$$
\Omega_0^2 = \omega_0^2 - \sum_{\mu} \frac{m_{\mu}}{m} \omega_{\mu}^2 \lambda_{\mu}^2. \tag{4}
$$

Each term in our Hamiltonian is strictly positive only if this quantity is positive. If it is negative then the second term is also negative and the Hamiltonian is not bounded from below and hence not allowed physically. Hence the positivity of the Hamiltonian places a physical restriction on the strength of the coupling

<span id="page-3-1"></span>
$$
\omega_0^2 > \sum_{\mu} \frac{m_{\mu}}{m} \omega_{\mu}^2 \lambda_{\mu}^2. \tag{5}
$$

It is convenient to rewrite our Hamiltonian in terms of the familiar annihilation and creation operators:

$$
\hat{a} = \sqrt{\frac{m\omega_0}{2\hbar}} \left(\hat{x} + \frac{i\hat{p}}{m\omega_0}\right)
$$

$$
\hat{b}_{\mu} = \sqrt{\frac{m_{\mu}\omega_{\mu}}{2\hbar}} \left(\hat{x}_{\mu} + \frac{i\hat{p}_{\mu}}{m_{\mu}\omega_{\mu}}\right)
$$
(6)

In terms of these operators our Hamiltonian becomes

$$
\hat{H} = \hbar \omega_0 \hat{a}^\dagger \hat{a} + \sum_{\mu} \hbar \omega_{\mu} \hat{b}^\dagger_{\mu} \hat{b}_{\mu} + \sum_{\mu} \frac{\hbar}{2} V_{\mu} \left( \hat{a} + \hat{a}^\dagger \right) \left( \hat{b}_{\mu} + \hat{b}^\dagger_{\mu} \right) \tag{7}
$$

when unimportant constant shifts in the ground-state energies are removed and

$$
V_{\mu} = -\sqrt{\frac{m_{\mu}\omega_{\mu}}{m\omega_{0}}} \omega_{\mu}\lambda_{\mu}.
$$
\n(8)

When written in terms of this quantity, our positivity condition [\(5\)](#page-3-1) becomes

$$
\omega_0 > \sum_{\mu} \frac{V_{\mu}^2}{\omega_{\mu}}.\tag{9}
$$

There is one further refinement that we need to introduce before considering the diagonalisation of the Hamiltonian and this it to introduce a continuum description of the environment. To this end we introduce continuum annihilation and creation operators,  $\hat{b}(\omega)$  and  $\hat{b}^{\dagger}(\omega)$ , satisfying the commutation relations

$$
\left[\hat{b}(\omega), \hat{b}^{\dagger}(\omega')\right] = \delta(\omega - \omega')\tag{10}
$$

and our Hamiltonian becomes [\[39\]](#page-17-21)

<span id="page-4-2"></span>
$$
\hat{H} = \hbar\omega_0 \hat{a}^\dagger \hat{a} + \int_0^\infty d\omega \,\hbar\omega \hat{b}^\dagger(\omega) \hat{b}(\omega) \n+ \int_0^\infty d\omega \,\frac{\hbar}{2} V(\omega) \left(\hat{a} + \hat{a}^\dagger\right) \left[\hat{b}^\dagger(\omega) + \hat{b}(\omega)\right]
$$
\n(11)

and the positivity condition is

<span id="page-4-0"></span>
$$
\omega_0 > \int_0^\infty d\omega \frac{V^2(\omega)}{\omega}.\tag{12}
$$

Our Hamiltonian is quadratic in the annihilation and creation operators for the oscillator and the reservoir and hence leads to linear coupled equations of motion for these operators. We could seek to solve these equations of motion and this would lead to an operator Heisenberg-Langevin equation analogous to [\(1\)](#page-2-0). Here we adopt the different approach of diagonalising the Hamiltonian.

# 2.1. Hamiltonian diagonalisation

We seek to diagonalise the Hamiltonian by finding a complete set of eigenoperators,  $\hat{B}(\omega)$  and their conjugates  $\hat{B}^{\dagger}(\omega)$  that satisfy the operator equations

<span id="page-4-1"></span>
$$
\begin{aligned}\n\left[\hat{B}(\omega), \hat{H}\right] &= \hbar\omega \hat{B}(\omega) \\
\left[\hat{B}^{\dagger}(\omega), \hat{H}\right] &= -\hbar\omega \hat{B}^{\dagger}(\omega)\n\end{aligned} \tag{13}
$$

for all positive frequencies  $\omega$ . These eigenoperators are complete if, in addition to these they also satisfy the condition

$$
\left[\hat{B}(\omega), \hat{B}^{\dagger}(\omega')\right] = \delta(\omega - \omega').\tag{14}
$$

These operator equations are the natural analogues of the more familiar eigenvalue and completeness conditions for the eigenstates of a Hamiltonian. In analogy with the eigenvalue problem, we expand each of the eigenoperators as a superposition of a complete set of operators:

<span id="page-5-2"></span>
$$
\hat{B}(\omega) = \alpha(\omega)\hat{a} + \beta(\omega)\hat{a}^{\dagger} + \int_0^{\infty} d\omega' \left[ \gamma(\omega, \omega')\hat{b}(\omega') + \delta(\omega, \omega')\hat{b}^{\dagger}(\omega') \right], \quad (15)
$$

and then use the eigenoperator equations and completeness condition to determine the coefficients in this expansion. The calculation is a little involved but the main points are summarised in [Appendix A.](#page-14-0)

We can express any of the annihilation and creation operators for the oscillator or the reservoir in terms of our eigenoperators. To do this we write the desired operator as a superposition of all the  $\hat{B}(\omega)$  and  $\hat{B}^{\dagger}(\omega)$  operators and then use the commutation relations to extract the coefficients in this expansion. For the oscillator operators we find

$$
\hat{a} = \int_0^\infty d\omega \left( \alpha^*(\omega) \hat{B}(\omega) - \beta(\omega) \hat{B}^\dagger(\omega) \right) \n\hat{a}^\dagger = \int_0^\infty d\omega \left( \alpha(\omega) \hat{B}^\dagger(\omega) - \beta^*(\omega) \hat{B}(\omega) \right).
$$
\n(16)

The requirement that these operators satisfy the familiar boson commutation relation,  $\left[\hat{a}, \hat{a}^{\dagger}\right] = 1$ , provides a constraint on the functions  $\alpha(\omega)$  and  $\beta(\omega)$  in the form

<span id="page-5-0"></span>
$$
\int_0^\infty d\omega \left[ |\alpha(\omega)|^2 - |\beta(\omega)|^2 \right] = \int_0^\infty d\omega |\alpha(\omega)|^2 \frac{4\omega_0 \omega}{(\omega_0 + \omega)^2} = 1,\tag{17}
$$

where we have used  $(A.10)$ . It is interesting to note that the correctness of this may be verified explicitly by contour integration, but that the proof makes explicit use of the positivity condition [\(12\)](#page-4-0) [\[39\]](#page-17-21).

The integrand in [\(17\)](#page-5-0) is clearly positive (or zero) for all frequencies and hence has the mathematical form of a frequency probability distribution:

<span id="page-5-1"></span>
$$
\pi(\omega) = |\alpha(\omega)|^2 \frac{4\omega_0 \omega}{(\omega_0 + \omega)^2}.
$$
\n(18)

A number of further constraints on this quantity emerge naturally from thinking of it as a probability distribution and from our diagonalisation.

#### 2.2. Physical constraints

We leave until the next section the question of the physical interpretation of the probability density  $\pi(\omega)$  but consider here what can be inferred from the fact that it has the mathematical properties of a probability distribution. To this end let us denote the average value of a function of frequency for this distribution by

<span id="page-5-3"></span>
$$
\langle\langle f(\omega)\rangle\rangle = \int_0^\infty d\omega f(\omega)\pi(\omega). \tag{19}
$$

we note that  $|\alpha(\omega)|^2$  is finite for all  $\omega$  and it follows, therefore, from [\(18\)](#page-5-1) that the average value  $\langle \langle \omega^{-1} \rangle \rangle$  is finite.

It follows from the eigenoperator equations [\(13\)](#page-4-1) that we can write the Hamiltonian in the form

$$
\hat{H} = \int_0^\infty d\omega \,\hbar\omega \,\hat{B}^\dagger(\omega)\hat{B}(\omega) + C,\tag{20}
$$

where C is an unimportant constant. If we substitute into this Hamiltonian our expression for the eigenoperators [\(15\)](#page-5-2) we require that the coefficients of  $\hat{a}^2$  and  $\hat{a}^{\dagger 2}$ should vanish so that

$$
\int_0^\infty d\omega \,\omega |\alpha(\omega)|^2 \left(\frac{\omega - \omega_0}{\omega + \omega_0}\right) = \int_0^\infty d\omega \frac{\pi(\omega)}{4\omega_0} (\omega^2 - \omega_0^2) = 0\tag{21}
$$

which implies that

<span id="page-6-2"></span>
$$
\langle \langle \omega^2 \rangle \rangle = \omega_0^2. \tag{22}
$$

The fact that the square of the mean value cannot exceed the mean value of the square for any probability distribution leads us to deduce that

<span id="page-6-1"></span>
$$
\langle\langle\omega\rangle\rangle<\omega_0.\tag{23}
$$

Finally we can apply the Cauchy-Schwartz inequality to provide a lower bound on the value of  $\langle \langle \omega^{-1} \rangle \rangle$ :

<span id="page-6-0"></span>
$$
\langle \langle \omega^{-1} \rangle \rangle \langle \langle \omega \rangle \rangle \ge 1
$$
  
\n
$$
\Rightarrow \langle \langle \omega^{-1} \rangle \rangle > \frac{1}{\omega_0}.
$$
 (24)

These inequalities are useful in determining the properties of the ground state.

#### 3. Ground-state

It is immediately clear from the form of the Hamiltonian [\(11\)](#page-4-2) that the ground state of the oscillator is not that of the undamped oscillator, that is the state annihilated by  $\hat{a}$ . To see this we need only note that the interaction term  $\hat{a}^\dagger \hat{b}^\dagger(\omega)$  acts on the combined ground state of the non-interacting oscillator and reservoir, adding a quantum to each. This suggests that the true ground state should be a superposition of states with varying numbers of quanta in both the oscillator and the reservoir and hence an entangled state. This situation is reminiscent of the ground state of an atom in quantum electrodynamics, which is dressed by virtual photons [\[49\]](#page-18-0). The dressing of the ground-state atom is responsible for a number of important effects including the Casimir-Polder interaction [\[50,](#page-18-1) [51\]](#page-18-2) and the form of the polarizability of the atom [\[52,](#page-18-3) [53\]](#page-18-4). It is reasonable to expect that it will be similarly significant for our strongly damped oscillator.

The true ground state, which we denote by the ket  $|0\rangle$ , is the zero-eigenvalue eigenstate of all the annihilation operators  $B(\omega)$ :

$$
\hat{B}(\omega)|0\rangle = 0 \qquad \qquad \forall \,\omega. \tag{25}
$$

The ground-state properties of the oscillator in this pure state are described by a mixed

$$
\hat{\rho}_{\rm Osc} = \text{Tr}_{\rm Env} \left( |0\rangle\langle 0| \right). \tag{26}
$$

The most straightforward way to determine the form of this mixed state is to use the characteristic function [\[20\]](#page-17-2):

state density operator obtained by tracing out the environmental degrees of freedom:

$$
\chi(\xi) = \text{Tr} \left[ \hat{\rho} \exp \left( \xi \hat{a}^{\dagger} - \xi^* \hat{a} \right) \right]
$$
  
= \langle 0 | \exp \left( \xi \hat{a}^{\dagger} - \xi^\* \hat{a} \right) | 0 \rangle. (27)

This function provides a complete description of the state and all of its statistical properties. To evaluate it we express  $\hat{a}$  and  $\hat{a}^{\dagger}$  in terms of the eigenoperators<sup>†</sup>

<span id="page-7-2"></span>
$$
\chi(\xi) = \langle 0 | \exp \left\{ \int_0^\infty d\omega \left[ (\xi \alpha + \xi^* \beta) \hat{B}^\dagger(\omega) - (\xi^* \alpha^* + \xi \beta^*) \hat{B}(\omega) \right] \right\} |0\rangle
$$
  
= 
$$
\exp \left( -\frac{1}{2} \int_0^\infty d\omega |\xi \alpha(\omega) + \xi^* \beta(\omega)|^2 \right).
$$
 (28)

This simple form, Gaussian in  $\xi$ , is characteristic of a squeezed thermal state. We can rewrite this characteristic function in terms of our probability density in the form

$$
\chi(\xi) = \exp\left[-\frac{1}{2} \int_0^\infty \pi(\omega) \left(\frac{\omega}{\omega_0} \xi_r^2 + \frac{\omega_0}{\omega} \xi_i^2\right)\right]
$$
  
= 
$$
\exp\left[-\frac{1}{2} \left(\frac{\langle\langle\omega\rangle\rangle}{\omega_0} \xi_r^2 + \langle\langle\omega^{-1}\rangle\rangle\omega_0 \xi_i^2\right)\right],
$$
 (29)

where  $\xi_r$  and  $\xi_i$  are the real and imaginary parts of  $\xi$  respectively. We note, in passing, that the quadrature operators for the oscillator (familiar from quantum optics [\[20\]](#page-17-2)) have unequal uncertainties:

$$
\Delta \left( \frac{\hat{a} + \hat{a}^{\dagger}}{\sqrt{2}} \right) = \sqrt{\frac{\langle \langle \omega^{-1} \rangle \rangle \omega_0}{2}}
$$

$$
\Delta \left( \frac{-i(\hat{a} - \hat{a}^{\dagger})}{\sqrt{2}} \right) = \sqrt{\frac{\langle \langle \omega \rangle \rangle}{2 \omega_0}}.
$$
(30)

The product of these two exceeds  $\frac{1}{2}$ , as it must, by virtue of the Cauchy-Schwartz inequality [\(24\)](#page-6-0).

We have established that in the ground state, the reduced state of the oscillator is Gaussian in position and momentum and, as such, is completely determined by the first and second moments of the position and momentum. These moments are

<span id="page-7-1"></span>
$$
\langle 0|\hat{x}|0\rangle = 0
$$

$$
\langle 0|\hat{p}|0\rangle = 0
$$

$$
\langle 0|\hat{x}^2|0\rangle = \frac{\hbar \langle \langle \omega^{-1} \rangle \rangle}{2m}
$$

<span id="page-7-0"></span>‡ We make use of the operator ordering theorem [\[20\]](#page-17-2)

$$
\exp(\hat{A} + \hat{B}) = \exp(\hat{A}) \exp(\hat{B}) \exp\left(-\frac{1}{2}[\hat{A}, \hat{B}]\right),\,
$$

which holds if the two operators  $\hat{A}$  and  $\hat{B}$  both commute with their commutator  $[\hat{A}, \hat{B}]$ .

$$
\langle 0|\hat{p}^2|0\rangle = \frac{\hbar m \langle\langle \omega \rangle\rangle}{2}
$$
  

$$
\langle 0|\hat{p}\hat{x} + \hat{p}\hat{x}|0\rangle = 0.
$$
 (31)

We note that the frequency of the undamped oscillator,  $\omega_0$ , makes no explicit appearance in these expectation values. The mean energy of the oscillator alone is

$$
\langle 0|\frac{\hat{p}^2}{2m} + \frac{1}{2}m\omega_0^2 \hat{x}^2 |0\rangle = \frac{1}{4}\hbar\omega_0 \left(\frac{\langle\langle \omega \rangle \rangle}{\omega_0} + \langle\langle \omega^{-1} \rangle \rangle \omega_0\right) > \frac{1}{2}\hbar\omega_0.
$$
 (32)

The fact that this exceeds the ground-state energy of the undamped oscillator is a reflection of the fact that there is an energy cost to be paid in order to decouple the oscillator from its environment [\[9\]](#page-16-8). We note, further, that the mean kinetic energy and potential energy for the oscillator alone do not have the same value and that this is in marked contrast to the ground state of the undamped oscillator.

We can diagonalise the density operator for the oscillator alone,  $\rho_{\text{Osc}}$ , by means of a squeezing transformation [\[20\]](#page-17-2) or, equivalently, introducing a new pair of annihilation and creation operators corresponding to a new oscillation frequency,  $\omega_c$ :

$$
\hat{c} = \sqrt{\frac{m\omega_c}{2\hbar}} \left(\hat{x} + i\frac{\hat{p}}{m\omega_c}\right)
$$

$$
\hat{c}^\dagger = \sqrt{\frac{m\omega_c}{2\hbar}} \left(\hat{x} - i\frac{\hat{p}}{m\omega_c}\right).
$$
(33)

To complete the diagonalisation we need only choose  $\omega_c$  such that the expectation values of  $\hat{c}^2$  and  $\hat{c}^{\dagger 2}$  are zero:

$$
\langle 0|\hat{c}^2|0\rangle = \frac{m\omega_c}{2\hbar} \langle 0|\hat{x}^2 - \frac{\hat{p}^2}{m^2\omega_c^2}|0\rangle = 0
$$
  

$$
\Rightarrow \omega_c = \sqrt{\frac{\langle \langle \omega \rangle \rangle}{\langle \langle \omega^{-1} \rangle \rangle}}.
$$
 (34)

This frequency is, by virtue of [\(23\)](#page-6-1) and [\(24\)](#page-6-0), less than  $\omega_0$ , the natural frequency of the undamped oscillator. The appearance of a reduced oscillation frequency is reminiscent of the observed reduction in the oscillation frequency of a damped classical oscillator but unlike in the classical case, the frequency  $\omega_c$  remains real even in the heavily damped regime. We note that Philbin and Horsley arrived at a similar reduced but always real frequency, which they associated with a decrease in the zero point energy [\[54\]](#page-18-5).

The mean number of  $c$ -quanta in the oscillator ground-state is

$$
\bar{n}_c = \langle 0|\hat{c}^\dagger \hat{c}|0\rangle = \frac{1}{2} \left( \sqrt{\langle \langle \omega \rangle \rangle \langle \langle \omega^{-1} \rangle \rangle} - 1 \right),\tag{35}
$$

which exceeds 0, as it should, by virtue of  $(24)$ . When written in terms of the c-quanta, the steady-state density operator takes the form of a thermal Bose-Einstein state, which we can write in the form

<span id="page-8-0"></span>
$$
\hat{\rho}_{\text{Osc}} = \frac{1}{\bar{n}_c + 1} \left( \frac{\bar{n}_c}{\bar{n}_c + 1} \right)^{\hat{c}^\dagger \hat{c}}.
$$
\n(36)

We may interpret this state as a thermal state for the oscillator at the shifted (and reduced) frequency  $\omega_c$  and at an effective "temperature"

$$
T_{\text{eff}} = \frac{\hbar \omega_c}{k_B \ln(1 + \bar{n}_c^{-1})}.
$$
\n(37)

We should note, however, that the true temperature in the ground-state is zero and that this quantity and the frequency  $\omega_c$  are at most only parameters with which to quantify the state of the oscillator and its entanglement with the environment. In particular, the von-Neumann entropy associated with the steady state of the oscillator is

$$
S(\text{Osc}) = (\bar{n}_c + 1)\ln(\bar{n}_c + 1) - \bar{n}_c\ln\bar{n}_c,\tag{38}
$$

and that, by virtue of the Araki-Lieb inequality [\[55,](#page-18-6) [56,](#page-18-7) [57\]](#page-18-8) and the fact that the full state is pure, means that this is also the total entropy of the environment:

$$
S(\text{Env}) = S(\text{Osc})\tag{39}
$$

and that the quantum mutual information [\[57\]](#page-18-8), or index of correlation [\[58\]](#page-18-9), between the oscillator and the environment is

$$
S(\text{Osc}: \text{Env}) = S(\text{Env}) + S(\text{Osc}) - S(\text{Osc}, \text{Env}) = 2S(\text{Osc}). \tag{40}
$$

# 4. Physical meaning of  $\pi(\omega)$

We have seen that the physical properties of the oscillator ground state may readily be expressed in terms of moments of the frequency given the probability distribution  $\pi(\omega)$ . Here we present the case for identifying this probability density with the contribution from the dressed modes, associated with the eigenoperators  $B(\omega)$ , to the the state of the oscillator. It is in, in essence, the spectrum of the true ground-state modes contributing to the oscillator state. We present three arguments to support this interpretation.

Our first justification arises from the form of the expectation values [\(31\)](#page-7-1). We know that the ground state of a harmonic oscillator of frequency  $\omega$  has

$$
\langle \hat{x}^2 \rangle = \frac{\hbar}{2m\omega} \n\langle \hat{p}^2 \rangle = \frac{\hbar m\omega}{2}.
$$
\n(41)

If we treat the state of the oscillator as a mixture of oscillators of different frequencies, each in its ground state, and contribution with weight  $\pi(\omega)$  then the resulting average mean-square values will be

$$
\langle \hat{x}^2 \rangle = \int_0^\infty d\omega \, \pi(\omega) \frac{\hbar}{2m\omega} = \frac{\hbar \langle \langle \omega^{-1} \rangle \rangle}{2m}
$$
\n
$$
\langle \hat{p}^2 \rangle = \int_0^\infty d\omega \, \pi(\omega) \frac{\hbar m\omega}{2} = \frac{\hbar m \langle \langle \omega \rangle \rangle}{2},\tag{42}
$$

which correspond to those obtained above.

Our second point arises from the form of the Hamiltonian for the oscillator

$$
\hat{H}_0 = \frac{\hat{p}^2}{2m} + \frac{1}{2}m\omega_0^2 \hat{x}^2.
$$
\n(43)

We can, by virtue of [\(22\)](#page-6-2), write this as a combination of potentials corresponding to different frequencies but weighted by  $\pi(\omega)$ :

$$
\hat{H}_0 = \frac{\hat{p}^2}{2m} + \frac{1}{2}m \int_0^\infty d\omega \,\pi(\omega)\omega^2 \hat{x}^2
$$

$$
= \frac{\hat{p}^2}{2m} + \frac{1}{2}m \langle \langle \omega^2 \rangle \rangle \hat{x}^2.
$$
\n(44)

Finally we note that the effective mean energy of the oscillator, which is associated with the diagonal form of the density operator  $(36)$  is

$$
\left(\bar{n}_c + \frac{1}{2}\right)\hbar\omega_c = \frac{1}{2}\hbar\int_0^\infty d\omega \,\pi(\omega)\omega = \frac{1}{2}\hbar\langle\langle\omega\rangle\rangle,\tag{45}
$$

which combines the characteristic ground state energies of the dressed oscillators, weighted by the probability distribution  $\pi(\omega)$ . The combination of these three features leads us to interpret  $\pi(\omega)$  as the proportion of the corresponding dressed oscillators contributing to the properties of the damped oscillator. We emphasise that the mathematical results obtained in the preceding section do not require us to adopt this interpretation of  $\pi(\omega)$  but we find it helpful to do so.

## 5. Oscillator dynamics

The exact diagonalisation of the Hamiltonian makes it straightforward to evaluate the time-evolution of any desired property of the oscillator. All that we need do is to express the desired observable in terms of the eigenoperators,  $\hat{B}(\omega)$  and  $\hat{B}^{\dagger}(\omega)$  and then use the time evolution of these operators, the form of which is an elementary consequence of the fact that they are eigenoperators:

$$
\hat{B}(\omega, t) = \hat{B}(\omega, 0)e^{-i\omega t} \n\hat{B}^{\dagger}(\omega, t) = \hat{B}^{\dagger}(\omega, 0)e^{i\omega t}.
$$
\n(46)

In particular, we can determine the time-evolution of the annihilation operator for the oscillator in this way:

<span id="page-10-0"></span>
$$
\hat{a}(t) = \int_0^\infty d\omega \left[ \alpha^*(\omega) \hat{B}(\omega, 0) e^{-i\omega t} - \beta(\omega) \hat{B}^\dagger(\omega, 0) e^{i\omega t} \right]
$$
  
\n
$$
= \int_0^\infty d\omega \left\{ \alpha^*(\omega) e^{-i\omega t} \left[ \alpha(\omega) \hat{a}(0) + \beta(\omega) \hat{a}^\dagger(0) + \int_0^\infty d\omega' \left( \gamma(\omega, \omega') \hat{b}(\omega', 0) + \delta(\omega, \omega') \hat{b}^\dagger(\omega', 0) \right) \right] - \beta(\omega) e^{i\omega t} \left[ \alpha^*(\omega) \hat{a}^\dagger(0) + \beta^*(\omega) \hat{a}(0) + \int_0^\infty d\omega' \left( \gamma^*(\omega, \omega') \hat{b}^\dagger(\omega', 0) + \delta^*(\omega, \omega') \hat{b}(\omega', 0) \right) \right] \right\}.
$$
 (47)

This may be used, together with the initial state of the oscillator and the environment, to evaluate the expectation value of any desired property. We note that if the environment is in a stationary state, so that  $\langle \hat{b}(\omega, 0) \rangle = 0 = \langle \hat{b}^\dagger(\omega, 0) \rangle$ , then the expectation values of the position and momentum operators take a pleasingly simple form:

<span id="page-11-0"></span>
$$
\langle \hat{x}(t) \rangle = \langle \langle \cos(\omega t) \rangle \rangle \langle \hat{x}(0) \rangle + \frac{1}{m} \langle \langle \omega^{-1} \sin(\omega t) \rangle \rangle \langle \hat{p}(0) \rangle
$$
  

$$
\langle \hat{p}(t) \rangle = \langle \langle \cos(\omega t) \rangle \rangle \langle \hat{p}(0) \rangle - m \langle \langle \omega \sin(\omega t) \rangle \rangle \langle \hat{x}(0) \rangle,
$$
 (48)

where the double angle brackets denote, as before, averages over our probability distribution  $\pi(\omega)$  as in [\(19\)](#page-5-3). The form of these equations adds further support to our interpretation of  $\pi(\omega)$  as a frequency probability distribution for the damped oscillator, as they may be viewed as the evolution of an undamped oscillator with a frequency  $\omega$  averaged using this probability distribution. The dissipation arises simply from a dephasing amongst the different frequency components.

The evolution of the mean position and momentum, as given in [\(48\)](#page-11-0), has the necessary short-time form of that for an undamped oscillator

$$
\langle \hat{x}(dt) \rangle = \langle \hat{x}(0) \rangle + \frac{dt}{m} \langle \hat{p}(0) \rangle
$$
  

$$
\langle \hat{p}(dt) \rangle = \langle \hat{p}(0) \rangle - m\omega_0^2 dt \langle \hat{x}(0) \rangle,
$$
 (49)

where we have used the identity  $\langle \langle \omega^2 \rangle \rangle = \omega_0^2$ . The effects of the coupling to the environment enter at order  $dt^3$  and this is an indication of the essentially non-Markovian nature of the strongly-damped oscillator. Our interest is in strongly damped oscillators and so we should note that [\(48\)](#page-11-0) includes the possibilities of both critically-damped and over-damped evolution. The equations contain, moreover, a simple criterion for these, which we may express in terms of our probability distribution. The motion will be oscillatory if  $\langle\langle \cos(\omega t)\rangle\rangle$  has stationary points at times other than at  $t = 0$ . Alternatively, we may state that the motion is under-damped if the derivative of this quantity, that is  $\langle\langle\omega\sin(\omega t)\rangle\rangle$ , is zero for any time other than  $t = 0$ . If it is zero only at  $t = 0$  then the motion is critically-damped or over-damped.

Our expression for the evolved annihilation operator [\(47\)](#page-10-0), together with the corresponding one for the creation operator provide a full description of the oscillator dynamics. This is true for any initial state of the oscillator and, moreover, for any environmental state including, of course, that associated with a finite temperature. As an illustration let us examine the evolution of the characteristic function for an arbitrary initial state of the oscillator coupled to a zero-temperature environment at time  $t = 0$ . With a little effort we find

<span id="page-11-1"></span>
$$
\chi(\xi, t) = \chi[\xi(t), 0] \exp\left[-\frac{1}{2} \int_0^\infty d\omega' \left| \int_0^\infty d\omega \,\mu(\omega, \omega', t) \right|^2\right],\tag{50}
$$

where

$$
\xi(t) = \int_0^\infty d\omega \left[ \eta(\omega, t) \alpha^*(\omega) - \eta^*(\omega, t) \beta(\omega) \right]
$$
  
\n
$$
\mu(\omega, \omega', t) = \eta(\omega, t) \gamma^*(\omega, \omega') - \eta^*(\omega, t) \delta^*(\omega, \omega')
$$
  
\n
$$
\eta(\omega, t) = \left[ \xi \alpha(\omega) + \xi^* \beta(\omega) \right] e^{i\omega t}.
$$
\n(51)

This characteristic function encodes the full dynamics and statistics of the oscillator. As a simple illustration of this we can determine the form of the steady state. To see

this we first note  $\xi(t)$  tends to zero as t tends to infinity and the different frequency components dephase so that

$$
\chi[\xi(\infty), 0] = \chi[0, 0] = 1. \tag{52}
$$

Evaluating the long-time limit of the exponential factor in [\(50\)](#page-11-1) requires some care in the handling of the delta-function and principal part components. We find

$$
\int_0^\infty d\omega' \left| \int_0^\infty d\omega \, \mu(\omega, \omega', \infty) \right|^2 = \int_0^\infty d\omega \left| \xi \alpha(\omega') + \xi^* \beta(\omega') \right|^2, \tag{53}
$$

so the steady-state characteristic function is

$$
\chi(\xi,\infty) = \exp\left(-\frac{1}{2}\int_0^\infty d\omega \left|\xi\alpha(\omega) + \xi^*\beta(\omega)\right|^2\right),\tag{54}
$$

which we recognise as the characteristic function for the oscillator in the global ground state [\(28\)](#page-7-2). This is a most satisfactory and exact result.

#### 6. Weak-coupling limit

The theory developed above was designed to treat the strongly-damped harmonic oscillator, but should also be applicable to the more familiar weakly damped oscillator, for which the oft-employed Born and Markov approximations are applicable and the steady state of the oscillator should be its ground state. We show here that this is indeed the case.

We start by considering the form of the function  $\alpha(\omega)$  in the weak coupling limit. To aid our analysis we rewrite the form given in [\(A.18\)](#page-16-17) as

$$
\alpha(\omega) = \frac{\omega + \omega_0}{\omega_0} \left( \frac{V(\omega)}{|V(\omega)|^2 Y(\omega) - i\pi |V(\omega)|^2} \right).
$$
\n(55)

The weak damping limit corresponds to choosing the coupling to the environment to be small or, more specifically, to  $|V(\omega)|^2 \ll \omega_0$ . It is clear that in this limit,  $|\alpha(\omega)|^2$  will be a sharply peaked function centred around the frequency for which  $Y(\omega) = 0$ . If the integral part in  $Y(\omega)$ , as given in [\(A.15\)](#page-15-1) is small[§](#page-12-0) then this frequency will be close to  $\omega_0$  and we can write

$$
|V(\omega)|^2 Y(\omega) \approx 4(\omega - \omega_0) - 4F(\omega), \tag{56}
$$

where

$$
4F(\omega) = \int_0^\infty d\omega' \left( \frac{\mathbb{P}}{\omega - \omega'} - \frac{1}{\omega + \omega'} \right) |V(\omega')|^2.
$$
 (57)

This leads, in turn, to a corresponding approximate form for  $\alpha(\omega)$ :

$$
\alpha(\omega) \approx \frac{V(\omega)}{2} \frac{1}{\omega - \omega_0 - F(\omega) - i\frac{\pi}{4}|V(\omega)|^2},\tag{58}
$$

where we have set  $\omega = \omega_0$  everywhere except in the denominator. We note that this is of the form that arises from the Fano diagonalisation of our problem if we make the

<span id="page-12-0"></span><sup>§</sup> If it is not then we will need to invoke ideas of renormalisation, a complication we wish to avoid.

rotating wave approximation by omitting from our original Hamiltonian all terms that are products of two creation operators or of two annihilation operarors [\[20,](#page-17-2) [59\]](#page-18-10).

Consistency with the above approximation, which led us to set  $\omega = \omega_0$  leads us to set  $\beta(\omega)$  to zero:

$$
\beta(\omega) \approx 0,\tag{59}
$$

so that the integral over all frequency of  $|\alpha(\omega)|^2$  is unity. Hence in this limit the steadystate characteristic function for our oscillator is

$$
\chi(\xi,\infty) = \exp\left(-\frac{1}{2}|\xi|^2\right),\tag{60}
$$

which we recognise as that for the ground state of the undamped oscillator [\[20\]](#page-17-2). Further, we note that in this limit our probability distribution function,  $\pi(\omega) \approx |\alpha(\omega)|^2$ , approaches a Lorentzian centred on  $\omega_0$ , with some width  $\gamma \|$ . Thus all the complexity of of the original problem is reduced, in the weak-damping limit to just two parameters: a natural oscillation frequency and a damping rate.

It was important to confirm that our more general treatment coincided, in the right limit, with the approximate methods used for weakly damped oscillators. We should note that even if we are working in the weakly damped regime, then our approach offers a systematic way to treat corrections to the results obtained using the Born and Markov approximations, which may play an important role in modelling measurements at the limits of sensitivity.

## 7. Conclusion

We have presented an exact diagonalisation of a simple quantum model of the damped harmonic oscillator, one that is applicable, in particular, to strong damping. We have found that much of the behaviour of the oscillator and may of its properties can be described in terms of a single probability function,  $\pi(\omega)$ , which we may interpret as the contribution of corresponding dressed mode, at frequency  $\omega$  to the oscillator. These properties include the steady state of the oscillator at zero temperature, the entanglement between the oscillator and its environment and also its evolution, both in the familiar under-damped regime but also in the more problematic over-damped regime.

We have applied our diagonalisation to study the properties of the true ground state and have shown that the oscillator part of this pure entangled state coincides with the steady-state of the oscillator in a zero-temperature environment. The diagonalisation is not specific to any particular state of the reservoir, however, and may readily be applied to environments at finite temperature. It may be extended, moreover, to include driving forces, coupled oscillators and multiple reservoirs, with the latter perhaps being at different temperatures [\[60\]](#page-18-11). This may provide some insights into important questions

<span id="page-13-0"></span> $\parallel$  This is not strictly true in the wings of the distribution, of course, as even in this limit we require  $\langle \langle \omega^2 \rangle \rangle = \omega_0^2$ , but the corresponding quantity for a true Lorentzian is divergent.

of principle in the nascent fields of quantum machines and quantum thermodynamics [\[61,](#page-18-12) [62,](#page-18-13) [63,](#page-18-14) [64,](#page-18-15) [65,](#page-18-16) [66\]](#page-18-17). We return to these ideas elsewhere.

#### Acknowledgments

The Hamiltonian diagonalisation upon which much of this work is based was first calculated by Bruno Huttner, in collaboration with SMB, 24 years ago in their work on the quantum electrodynamics of dielectric media. We are most grateful to him and also to Paul Radmore for helpful comments and suggestions.

#### <span id="page-14-0"></span>Appendix A. Fano diagonalisation

We present a brief account of the exact diagonalisation of our Hamiltonian based on methods developed by Fano for the study of configuration interactions [\[67\]](#page-18-18). This idea was applied extended to weakly-coupled oscillators in a quantum study of damped cavity modes [\[20,](#page-17-2) [59\]](#page-18-10). The extension to stronger couplings, with the inclusion of counterrotating couplings, has been given before and applied to the quantum theory of light in dielectric media [\[38,](#page-17-20) [39\]](#page-17-21). We summarise here the analysis presented in [\[39\]](#page-17-21).

Our task is to diagonalise the damped harmonic oscillator Hamiltonian[¶](#page-14-1)

$$
\hat{H} = \hbar\omega_0 \hat{a}^\dagger \hat{a} + \int_0^\infty d\omega \,\hbar\omega \hat{b}^\dagger(\omega) \hat{b}(\omega) \n+ \int_0^\infty d\omega \, \frac{\hbar}{2} \left( \hat{a} + \hat{a}^\dagger \right) \left[ V(\omega) \hat{b}^\dagger(\omega) + V^*(\omega) \hat{b}(\omega) \right],
$$
\n(A.1)

by which we mean rewriting it in the form of a continuum of uncoupled or dressed oscillators:

$$
\hat{H} = \int_0^\infty d\omega \,\hbar\omega \,\hat{B}^\dagger(\omega)\hat{B}(\omega) + C,\tag{A.2}
$$

where  $C$  is an unimportant constant. We proceed by writing the dressed annihilation operators,  $B(\omega)$ , as linear combinations of the bare operators for the oscillator and bath modes:

<span id="page-14-2"></span>
$$
\hat{B}(\omega) = \alpha(\omega)\hat{a} + \beta(\omega)\hat{a}^{\dagger} + \int_0^{\infty} d\omega' \left[ \gamma(\omega, \omega')\hat{b}(\omega') + \delta(\omega, \omega')\hat{b}^{\dagger}(\omega') \right], \quad (A.3)
$$

where  $\alpha\omega$ ,  $\beta(\omega)$ ,  $\gamma(\omega, \omega')$  and  $\delta(\omega, \omega')$  are to be determined.

We require the operator  $B(\omega)$  to be associated with an uncoupled or dressed oscillator of angular frequency  $\omega$ . This requires us to find its form such that the following pair of operator equations are satisfied for every frequency,  $\omega$ :

$$
\left[\hat{B}(\omega),\hat{H}\right] = \hbar\omega\hat{B}(\omega)
$$
\n(A.4)

<span id="page-14-3"></span>
$$
\left[\hat{B}(\omega), \hat{B}^{\dagger}(\omega')\right] = \delta(\omega - \omega'). \tag{A.5}
$$

<span id="page-14-1"></span>**T** For the Hamiltonian of interest in this paper the coupling,  $V(\omega)$ , is real but treating the problem with a more general complex coupling presents no additional difficulties.

Substituting the ansatz  $(A.3)$  into  $(A.4)$  and comparing coefficients of the bare creation and annihilation operators leads to the set of couple equations:

$$
\alpha(\omega)\omega_0 + \frac{1}{2} \int_0^\infty d\omega' \left[ \gamma(\omega, \omega') V(\omega') - \delta(\omega, \omega') V^*(\omega') \right] = \alpha(\omega)\omega \tag{A.6}
$$

<span id="page-15-2"></span>
$$
-\beta(\omega)\omega_0 + \frac{1}{2} \int_0^\infty d\omega' \left[ \gamma(\omega, \omega') V(\omega') - \delta(\omega, \omega') V^*(\omega') \right] = \beta(\omega)\omega \tag{A.7}
$$

$$
\frac{V^*(\omega')}{2} [\alpha(\omega) - \beta(\omega)] + \gamma(\omega, \omega')\omega' = \gamma(\omega, \omega')\omega
$$
\n(A.8)

$$
\frac{V(\omega')}{2} [\alpha(\omega) - \beta(\omega)] - \delta(\omega, \omega')\omega' = \delta(\omega, \omega')\omega.
$$
 (A.9)

Our method of solution is use these to determine the functions  $\beta\omega$ ,  $\gamma(\omega,\omega')$  and  $\delta(\omega, \omega')$  in terms of  $\alpha\omega$  and then to determine this remaining function by enforcing the commutation relation  $(A.5)$ . From  $(A.6)$  and  $(A.7)$  we see that

<span id="page-15-0"></span>
$$
\beta(\omega) = \frac{\omega - \omega_0}{\omega + \omega_0} \alpha(\omega). \tag{A.10}
$$

If we use this to substitute for  $\beta(\omega)$  into the remaining equations then we find

$$
V^*(\omega')\frac{\omega_0}{\omega+\omega_0}\alpha(\omega) = \gamma(\omega,\omega')(\omega-\omega')
$$
\n(A.11)

$$
V(\omega')\frac{\omega_0}{\omega+\omega_0}\alpha(\omega) = \delta(\omega, \omega')(\omega+\omega').
$$
\n(A.12)

Solving the second of these presents no difficulty and we find

$$
\delta(\omega, \omega') = \left(\frac{1}{\omega + \omega'}\right) V(\omega') \frac{\omega_0}{\omega + \omega_0} \alpha(\omega).
$$
\n(A.13)

The first, however, requires careful handling because the behaviour at  $\omega = \omega'$ . Following Fano [\[67\]](#page-18-18), we adopt the method proposed by Dirac [\[68\]](#page-18-19) and write

<span id="page-15-3"></span>
$$
\gamma(\omega,\omega') = \left(\frac{\mathbb{P}}{\omega - \omega'} + Y(\omega)\delta(\omega - \omega')\right)V^*(\omega')\frac{\omega_0}{\omega + \omega_0}\alpha(\omega),\tag{A.14}
$$

where P denotes that the principal part is to be taken on integration and  $Y(\omega)$  is a real function, which we determine by substituting  $(A.14)$  into  $(A.6)$ . We find

<span id="page-15-1"></span>
$$
Y(\omega) = \frac{1}{|V(\omega)|^2} \left[ \frac{2(\omega^2 - \omega_0^2)}{\omega_0} - \int_0^\infty d\omega' \left( \frac{\mathbb{P}}{\omega - \omega'} - \frac{1}{\omega + \omega'} \right) |V(\omega')|^2 \right].
$$
\n(A.15)

If we substitute our operators,  $\hat{B}(\omega)$ , expressed in terms of the function  $\alpha(\omega)$  into the commutation relation [\(A.6\)](#page-15-2) then we find

$$
\left[\hat{B}(\omega), \hat{B}^{\dagger}(\omega')\right] = \alpha(\omega)\alpha^*(\omega')\left\{1 - \left(\frac{\omega - \omega_0}{\omega + \omega_0}\right)\left(\frac{\omega' - \omega_0}{\omega' + \omega_0}\right) \right.\left. + \int_0^\infty d\omega''\left[\left(\frac{P}{\omega - \omega''} + Y(\omega)\delta(\omega - \omega'')\right)\left(\frac{P}{\omega' - \omega''} + Y(\omega)\delta(\omega' - \omega'')\right) \right.\left. - \left(\frac{1}{\omega + \omega''}\right)\left(\frac{1}{\omega' + \omega''}\right)\right]\frac{|V(\omega'')|^2 \omega_0^2}{(\omega + \omega_0)(\omega' + \omega_0)}\right\}.
$$
\n(A.16)

Evaluating the integrals and setting the result equal to  $\delta(\omega - \omega')$  gives<sup>[+](#page-16-18)</sup>

$$
|\alpha(\omega)|^2 = \frac{(\omega + \omega_0)^2}{\omega_0^2 |V(\omega)|^2} \left(\frac{1}{Y^2(\omega) + \pi^2}\right).
$$
\n(A.17)

Note that the diagonalisation does not fix the phase of the complex function  $\alpha(\omega)$  and we are free to choose this as we wish. A convenient choice is to set

<span id="page-16-17"></span>
$$
\alpha(\omega) = \frac{\omega + \omega_0}{\omega_0 V^*(\omega)} \left( \frac{1}{Y(\omega) - i\pi} \right). \tag{A.18}
$$

#### References

- <span id="page-16-0"></span>[1] Chan J, Alegre TPM, Safavi-Naeini AH, Hill JT, Gröblacher S, Aspelmeyer M and Painter O 2011 Laser cooling of a nanomechanical oscillator into its quantum ground state Nature 478 89–92
- <span id="page-16-1"></span>[2] Gigan S, Böhm HR, Paternostro M, Blaser F, Langer G, Hertzberg JB, Schwab KC, Bäuerle D, Aspelmeyer M and Zeilinger A 2006 Self-cooling of a micromirror by radiation pressure Nature 444 67–70
- <span id="page-16-2"></span>[3] Stannigel K, Rabl P, Sørensen AS, Zoller P and Lukin MD 2010 Optomechanical transducers for long-distance quantum communication Phys. Rev. Lett. 105 220501
- <span id="page-16-3"></span>[4] Zhang K, Bariani F and Meystre P 2014 Quantum optomechanical heat engine Phys. Rev. Lett. 112 150602
- <span id="page-16-4"></span>[5] Joshi C, Ohberg P, Cresser JD and Andersson E 2014 Markovian evolution of strongly coupled ¨ harmonic oscillators *Phys. Rev. A* **90** 063815
- <span id="page-16-5"></span>[6] Brunelli M, Xuereb A, Ferraro A, De Chiara G, Kiesel N and Paternostro M 2015 Out-ofequilibrium thermodynamics of quantum optomechanical systems New J. Phys. 17 035016
- <span id="page-16-6"></span>[7] Campisi M, Talkner P and Hänggi P 2009 Fluctuation theorem for arbitrary open quantum systems Phys. Rev. Lett. 102 210401
- <span id="page-16-7"></span>[8] Talkner P, Campisi M and Hänggi P 2009 Fluctuation theorems in driven open quantum systems J. Stat. Mech. P02025
- <span id="page-16-8"></span>[9] Allahverdyan AE and Nieuwenhuizen TM 2000 Extraction of work from a single thermal bath in the quantum regime Phys. Rev. Lett. 85 1799–1802
- <span id="page-16-9"></span>[10] Philbin TG 2012 Quantum dynamics of the damped harmonic oscillator New J. Phys. 14 083043
- <span id="page-16-10"></span>[11] Schrödinger E 1926 Quantisierung als Eigenwertproblem Ann. Phys. 79, 489-527; reprinted in translation as "Quantisation as a problem of proper values (Part II)" in Schrödinger E 1982 Collected Papers on Wave Mechanics 3rd ed. (Providence, Rhode Island: AMS Chelsea Publishing)
- <span id="page-16-11"></span>[12] Langevin P 1908 Sur la théoriedu mouvement brownien Comptes Rend. Acad. Sci. (Paris) 146 530–533; reprinted in a translation by Gythiel A 1997 "On the theory of Brownian motion" Am. J. Phys. 65 1079–1081
- <span id="page-16-12"></span>[13] Uhlenbeck GE and Ornstein LS 1930 On the theory of Brownian motion Phys. Rev. 36, 823–841
- <span id="page-16-13"></span>[14] Wang MC and Uhlenbeck GE 1945 On the theory of Brownian motion II Rev. Mod. Phys. 17 323–342
- <span id="page-16-14"></span>[15] Risken H 1989 The Fokker-Planck equation 2nd ed. (Berlin: Springer)
- <span id="page-16-15"></span>[16] Lemons DS 2002 An Introduction to Stochastic Processes in Physics (Baltimore: Johns Hopkins University Press)
- <span id="page-16-16"></span>[17] Mazo RM 2002 Brownian Motion (Oxford: Oxford University Press)
- <span id="page-16-18"></span><sup>+</sup> This requires the use of the following formula for the product of two principal parts [\[20\]](#page-17-2):

$$
\frac{\mathbb{P}}{\omega - \omega''} \frac{\mathbb{P}}{\omega' - \omega''} = \frac{\mathbb{P}}{\omega' - \omega} \left( \frac{\mathbb{P}}{\omega - \omega''} - \frac{\mathbb{P}}{\omega' - \omega''} \right) + \pi^2 \delta(\omega - \omega'') \delta(\omega' - \omega'').
$$

- <span id="page-17-0"></span>[18] Senitzky IR 1960 Dissipation in quantum mechanics. The harmonic oscillator Phys. Rev. 119 670–679
- <span id="page-17-1"></span>[19] Louisell WH 1964 Radiation and Noise in Quantum Electronics (New York: McGraw-Hill)
- <span id="page-17-2"></span>[20] Barnett SM and Radmore PM 1997 Methods in Theoretical Quantum Optics (Oxford: Oxford University Press)
- <span id="page-17-3"></span>[21] Louisell WH 1973 Quantum Statistical Properties of Radiation (New York: Wiley)
- <span id="page-17-4"></span>[22] Peřina J 1991 Quantum Statistics of Linear and Nonlinear Optical Phenomena 2nd ed. (Dordrecht: Kluwer)
- <span id="page-17-5"></span>[23] Meystre P and Sargent III M 1991 Elements of Quantum Optics 2nd ed. (Berlin: Springer-Verlag)
- <span id="page-17-6"></span>[24] Carmichael H 1993 An Open Systems Approach to Quantum Optics (Berlin: Springer-Verlag)
- <span id="page-17-7"></span>[25] Walls DF and Milburn GJ 1994 Quantum Optics (Berlin: Springer-Verlag)
- <span id="page-17-8"></span>[26] Carmichael HJ 1999 Statistical Methods in Quantum Optics 1 (Berlin: Springer)
- <span id="page-17-9"></span>[27] Breuer H-P and Petruccione F 2002 The Theory of Open Quantum Systems (Oxford; Oxford University Press)
- <span id="page-17-10"></span>[28] Gardiner CW and Zoller P 2004 Quantum Noise 2nd ed. (Berlin: Springer)
- <span id="page-17-11"></span>[29] Ficek Z and Swain S 2004 Quantum Interference and Coherence (New York: Springer)
- <span id="page-17-12"></span>[30] Wiseman HM and Milburn GJ 2010 Quantum Measurement and Control (Cambridge: Cambridge University Press)
- <span id="page-17-13"></span>[31] Weiss U 2012 Quantum Dissipative Systems 4th ed. (New Jersey: World Scientific)
- <span id="page-17-14"></span>[32] Agarwal GS 1971 Brownian motion of a quantum oscillator Phys. Rev. A 4 739–747
- <span id="page-17-15"></span>[33] Ullersma P 1966 An exactly solvable model for Brownian motion I. Derivation of the Langevin equation Physica 32 27-55
- <span id="page-17-16"></span>[34] Ullersma P 1966 An exactly solvable model for Brownian motion II. Derivation of the Fokker-Planck equation and the master equation Physica 32 56–73
- <span id="page-17-17"></span>[35] Caldeira AO and Leggett AJ 1983 Path integral approach to quantum Brownian motion Physica 121 A 587–616
- <span id="page-17-18"></span>[36] Caldeira AO and Leggett AJ 1983 Quantum tunnelling in a dissipative system Ann. Phys. N.Y. 149 374–456
- <span id="page-17-19"></span>[37] Haake F and Reibold R 1985 Strong damping and low-temperature anomalies for the harmonic oscillator Phys. Rev. A 32 2462–2475
- <span id="page-17-20"></span>[38] Huttner B and Barnett SM 1992 Dispersion and loss in a Hopfield dielectric Europhys. Lett. 18 487–492
- <span id="page-17-21"></span>[39] Huttner B and Barnett SM 1992 Quantization of the electromagnetic field in dielectrics Phys. Rev. A 46 4306–4322
- <span id="page-17-22"></span>[40] Kheirandish F and Amooshahi 2006 Electromagnetic field quantisation in a linear polarizable and magnetizable medium Phys. Rev. A 74 042102
- <span id="page-17-23"></span>[41] Kheirandish F and Soltani M 2008 Extension of the Huttner-Barnett model to a magnetodielectric medium Phys. Rev. A 78 012102
- <span id="page-17-24"></span>[42] Kheirandish F, Amooshahi M and Soltani M 2009 A canonical approach to electromagnetic field quantization in a nonhomogeneous and anistropic magnetodielectric medium J. Phys. B: At. Mol. Opt. 42 075504
- <span id="page-17-25"></span>[43] Amooshahi M 2009 Canonical quantisation of the electromagnetic field in an anisotropic polarizable and magnetizable medium J. Math. Phys. 50 062301
- <span id="page-17-26"></span>[44] Munro WJ and Gardiner CW 1996 Non-rotating-wave master equation Phys. Rev. A 53 2633–2640
- <span id="page-17-27"></span>[45] Stenholm S 1996 Simple quantum dynamics in Quantum Dynamics of Simple Systems G. L. Oppo, S. M. Barnett, E. Riis and M. Wilkinson eds. (Bristol: Scottish Universities Summer Schools in Physics and Institute of Physics) pp. 267–313
- <span id="page-17-28"></span>[46] Barnett SM and Cresser JD Quantum theory of friction Phys. Rev. A 72 022107
- <span id="page-17-29"></span>[47] Barnett SM, Jeffers J and Cresser JD 2006 From measurements to quantum friction J. Phys: Condens. Matter 18 S401–S410
- <span id="page-17-30"></span>[48] Hu BL, Paz JP and Zhang Y 1992 Quantum Brownian motion in a general environment: Exact

master equation with nonlocal dissipation and colored noise Phys. Rev. D 45 2843–2861

- <span id="page-18-0"></span>[49] Compagno C, Passante R and Persico F 1995 Atom-Field Interactions and Dressed Atoms (Cambridge: Cambridge University Press)
- <span id="page-18-1"></span>[50] Casimir HBG and Polder D 1948 The influence of retardation on the London-van der Waals forces Phys. Rev. 73 360–372
- <span id="page-18-2"></span>[51] Power EA 1964 Introductory Quantum Electrodynamics (London: Longmans)
- <span id="page-18-3"></span>[52] Loudon R and Barnett SM 2006 Theory of the linear polarizability of a two-level atom J. Phys. B: At. Mol. Opt. Phys. 39 S555–S563
- <span id="page-18-4"></span>[53] Milonni PW, Loudon R, Berman PR and Barnett SM 2008 Linear polarizabilities of two- and three- level atoms Phys. Rev. A 77 043835
- <span id="page-18-5"></span>[54] Philbin TG and Horsley SAR 2013 Damping the zero-point energy of a harmonic oscillator  $arXiv$ : 1304.0977v2 [quant-ph]
- <span id="page-18-6"></span>[55] Araki H and Lieb EH 1970 Entropy inequalities Commun. Math. Phys. 18 160–170
- <span id="page-18-7"></span>[56] Wehrl A 1978 General properties of entropy Rev. Mod. Phys. 50 221–260
- <span id="page-18-8"></span>[57] Barnett SM 2009 Quantum Information (Oxford: Oxford University Press)
- <span id="page-18-9"></span>[58] Barnett SM and Phoenix SJD 1989 Entropy as a measure of quantum optical correlation Phys. Rev. A 40 2404–2409
- <span id="page-18-10"></span>[59] Barnett SM and Radmore PM 1988 Quantum theory of cavity quasimodes Opt. Commun. 68 364–368
- <span id="page-18-11"></span>[60] Martinez EA and Paz JP 2013 Dynamics and thermodynamics of linear quantum open systems Phys. Rev. Lett. 110 130406
- <span id="page-18-12"></span>[61] Gemmer J, Michel M and Mahler G 2004 Quantum Thermodynamics (Berlin: Springer)
- <span id="page-18-13"></span>[62] Vaccaro JA and Barnett SM 2011 Information erasure without an energy cost Proc. R. Soc. A 467 1770–1778
- <span id="page-18-14"></span>[63] Barnett SM and Vaccaro JA 2013 Beyond Landauer erasure Entropy 15 4956–4968
- <span id="page-18-15"></span>[64] Roncaglia AJ, Cerisola F and Paz JP 2014 Work as a generalized measurement Phys. Rev. Lett. 113 0250601
- <span id="page-18-16"></span>[65] Binder F, Vinjanampathy S, Modi K and Goold J 2015 Operational thermodynamics of open quantum systems Phys. Rev. E 91 032119
- <span id="page-18-17"></span>[66] Brandao F, Horodecki M, Ng K, Oppenheim J and Wehner S 2015 The second laws of quantum thermodynamics Proc. Natl. Acad. Sci. USA 112 3275–3279
- <span id="page-18-18"></span>[67] Fano U 1956 Effects of configuration interaction on intensities and phase shifts Phys. Rev. 124 1866–1878
- <span id="page-18-19"></span>[68] Dirac PAM 1927 Uber die Quantenmechanik der Stoßvorgänge Z. Phys. 44 585–595; reprinted in translation as "On the quantum mechanics of collisions" in R. H. Dalitz (ed.) 1995 The Collected Works of P. A. M. Dirac 1924–1948 (Cambridge: Cambridge University Press)