

# On the phase structure of driven quantum systems

Vedika Khemani,<sup>1,2</sup> Achilleas Lazarides,<sup>2</sup> Roderich Moessner,<sup>2</sup> and S. L. Sondhi<sup>1,2</sup>

<sup>1</sup>*Department of Physics, Princeton University, Princeton, NJ 08544*

<sup>2</sup>*Max-Planck-Institut für Physik komplexer Systeme, 01187 Dresden, Germany*

(Dated: May 26, 2022)

Clean and interacting periodically driven quantum systems are believed to exhibit a single, trivial “infinite-temperature” Floquet-ergodic phase. In contrast, here we show that their disordered Floquet many-body localized counterparts can exhibit distinct ordered phases delineated by sharp transitions. Some of these are analogs of equilibrium states with broken symmetries and topological order, while others — genuinely new to the Floquet problem — are characterized by order *and* non-trivial periodic dynamics. We illustrate these ideas in driven spin chains with Ising symmetry.

**Introduction:** Extending ideas from equilibrium statistical mechanics to the non-equilibrium setting is a topic of perennial interest. We consider a question in this vein: Is there a sharp notion of a phase in driven, interacting quantum systems? We find an affirmative answer for Floquet systems<sup>1–3</sup>, i.e. systems whose Hamiltonians depend on time  $t$  periodically,  $H(t+T) = H(t)$ . Unlike in equilibrium statistical mechanics, disorder turns out to be an essential ingredient for stabilizing different phases; moreover, the periodic time evolution allows for the existence (and diagnosis) of phases without any counterparts in equilibrium statistical mechanics.

Naively, Floquet systems hold little promise of a complex phase structure. In systems with periodic Hamiltonians, not even the basic concept of energy survives, being replaced instead with a quasi-energy defined up to arbitrary shifts of  $2\pi/T$ . Indeed, interacting Floquet systems should absorb energy indefinitely from the driving field, as suggested by standard linear response reasoning wherein any nonzero frequency exhibits dissipation. This results in the system heating up to “infinite temperature”, at which point all static and dynamic correlations become trivial and independent of starting state — thus exhibiting a maximally trivial form of ergodicity<sup>4–6</sup>.

To get anything else thus requires a mechanism for localization in the energy domain wherein the absorption from the driving field saturates and the system exhibits a long time state which is sensitive to initial conditions. The current dominant belief is that clean or translationally invariant interacting systems cannot generically exhibit such energy localization<sup>4–6</sup>, although there are computations that suggest otherwise<sup>7–9</sup>. The basic intuition is that modes in translationally invariant systems are spatially extended so that they interact with and transfer energy between each other.

This can be different when disorder spatially localizes the modes, with individual modes exhibiting something like Rabi oscillations while interacting only weakly with distant modes. While the actual situation with interactions is somewhat more involved, several pieces of work<sup>10–12</sup> have made a convincing case for the existence of Floquet energy localization exhibiting a set of properties closely related to those exhibited by time-independent many-body localized<sup>13</sup> (MBL) systems<sup>14</sup>.

In the following we show that such Floquet-MBL systems can exhibit multiple phases. Some of these are driven cousins of MBL phases characterized by broken symmetries and topological order. Remarkably, others are genuinely new to the Floquet setting, characterized by order *and* non-trivial periodic dynamics. Our analysis identifies a key feature of the Floquet problem, the existence of Floquet eigenstates, which permits us to extend the notion of eigenstate order<sup>15–19</sup> to time dependent Hamiltonians. Our work also builds on the discovery of topologically non-trivial Floquet single particle systems and recent advances in their classification<sup>20–28</sup>. As we will explain, non-trivial single particle drives can yet lead to trivial many-body periodic dynamics even without interactions. Thus, the full framework of disorder and interactions is required for the many-body problem.

Below we offer details of this analysis. After a quick review of the Floquet eigenstate formalism, we describe the notion of eigenstate order and generalize it to the Floquet problem. We then illustrate our general ideas for driven Ising spin chains. First we show that there are two Floquet phases, paramagnet (PM) and spin glass (SG), that connect smoothly to phases in the undriven systems. We then identify two *new* phases that do *not*, which we term the Floquet  $0\pi$ -PM and the Floquet  $\pi$ -SG. Along the way we note that, reformulated as fermion problems, these yield instances of Floquet topological order. We close with a summary and outlook.

**Floquet Formalism:** We consider periodic, interacting Hamiltonians which are local in space. The time dependent Schrödinger equation

$$i\frac{d|\psi(t)\rangle}{dt} = H(t)|\psi(t)\rangle \quad (1)$$

has special solutions<sup>1–3</sup>:

$$|\psi_\alpha(t)\rangle = e^{-i\epsilon_\alpha t}|\phi_\alpha(t)\rangle \quad (2)$$

defined by periodic states  $|\phi_\alpha(t)\rangle = |\phi_\alpha(t+T)\rangle$  and quasi-energies  $\epsilon_\alpha$  defined modulo  $2\pi/T$ .

These replace the eigenstates of the time independent problem; in them observables have periodic expectation values, and they form a complete basis. The “Floquet Hamiltonian”  $H_F$  is defined via the time evolution operator over a full period,  $U(T) = e^{-iH_F T}$ . The  $|\phi_\alpha(0)\rangle$

are eigenstates of both  $U(T)$  and  $H_F$ , with eigenvalues  $e^{-i\epsilon_\alpha T}$  and  $\epsilon_\alpha$  respectively.

The question of energy localization relates to the action of  $U(nT) = (U(T))^n$  as  $n \rightarrow \infty$ . An equivalent formulation of Floquet-MBL is that there exists an  $H_F$  which is local and exhibits the generic properties of any fully MBL Hamiltonian, such as a full set of conserved local operators<sup>29–33</sup> ( $l$ -bits), area law eigenstates and failure of the eigenstate thermalization hypothesis (ETH)<sup>34</sup>.

**Eigenstate Order:** Traditionally, phase transitions at non-zero energy densities are considered in the framework of quantum statistical mechanics, signaled by singularities in thermodynamic functions or observables computed in the  $T > 0$  Gibbs state. Work on MBL has led to the realization that this viewpoint is too restrictive<sup>15–19</sup>—instead one can look directly at the many-body eigenstates and eigenvalues and ask if they exhibit singular changes as a parameter is varied. Such transitions have been termed eigenstate phase transitions. This distinction is irrelevant for Hamiltonians obeying ETH, but when ETH fails it becomes important. The passage from ergodicity to localization is an example of an eigenstate transition undetectable by standard ensembles<sup>34–36</sup>—it can take place at  $T = \infty$  and yet the individual many body eigenstates are sharply different in their entanglement properties, with the eigenvalue distributions exhibiting different statistics. Moreover, eigenstate phase transitions can take place between two ETH violating phases<sup>15</sup> and they may even involve a singular rearrangement of the eigenvalues alone.

We now generalize this to the Floquet-MBL regime, leading us to find multiple phases whose existence can be detected in the Floquet many-body eigensystem. In general this will require examination of the full periodic solutions  $|\psi_\alpha(t)\rangle$  for sharply different characteristics. The transition out of the Floquet ergodic phase to a Floquet-MBL phase lacking any order has already been discussed<sup>10,11</sup>. We will now exhibit Floquet-MBL phases with order in the canonically simplest interacting setting: a one dimensional disordered spin chain with the Ising symmetry,

$$H = \sum_i J_i \sigma_i^x \sigma_{i+1}^x + \sum_i h_i \sigma_i^z + J_z \sum_i \sigma_i^z \sigma_{i+1}^z. \quad (3)$$

Carrying out a Jordan-Wigner transformation on only the first two terms gives a p-wave superconducting free-fermion model, whereas the final term is a density-density interaction in the fermion language.

**Floquet Paramagnet and Spin Glass:** We begin with the two phases that *do* exist in undriven systems and demonstrate the stability of these to being (not too strongly) driven. Starting with the non-interacting limit,  $J_z = 0$ , we choose the  $J_i$  and the  $h_i$  to be log-normally distributed with a tunable mean  $\overline{\log(J_i)} \equiv \overline{\log \bar{J}}$ , fixed  $\overline{\log(h_i)} \equiv \overline{\log \bar{h}} = 0$  and two fixed and equal standard deviations  $\delta \log(h_i) = \delta \log(J_i) = 1$ . Work on random, non-interacting transverse field Ising models culminating in Ref. 37 finds a ground state phase diagram which is a




FIG. 1. Level statistics  $\langle r \rangle$  of  $H_F$  for the driven, disordered Ising model (4) averaged over 2000-100,000 samples for the different system sizes.  $\langle r \rangle$  approaches the Poisson limit of .386 with increasing  $L$  deep in the PM and SG phases, showing that these remain well localized. There is a peak in  $\langle r \rangle$  near the non-intersecting critical point at  $\overline{\log J} = \overline{\log h}$  indicating partial delocalization, although the value still remains well below the COE value of .527. (inset): The SG diagnostic  $\chi^{SG}$  defined in (6) which goes to 0 in the paramagnet and approaches a non-zero value in the SG phase.

paramagnet for  $\overline{\log J} < \overline{\log h}$  and a  $Z_2$  breaking ferromagnet for  $\overline{\log J} > \overline{\log h}$ , separated by an infinite disorder fixed point at  $\overline{\log J} = \overline{\log h}$ . The work on eigenstate order has shown that, with disorder and localization, both phases exist at *all* energies with the symmetry-breaking phase exhibiting SG order in individual eigenstates instead of ferromagnetism.

With weak interactions,  $0 < J_z \ll 1$ , the strongly localized PM and SG phases remain (many-body) localized<sup>38,39</sup>. The fate of the SG-PM transition is more sensitive to the inclusion of interactions. It was suggested that it would remain localized<sup>15</sup> and exhibit the same scaling as the non-interacting fixed point<sup>38</sup>; we comment on the analogous question in the Floquet setting below.

We consider a periodic binary drive—computationally much simpler than a monochromatic modulation—switching between two static  $H$ s with  $\overline{\log J}$  differing by 1:

$$H(t) = \sum_i f_s(t) J_i \sigma_i^x \sigma_{i+1}^x + \sum_i h_i \sigma_i^z + J_z \sum_i \sigma_i^z \sigma_{i+1}^z, \quad (4)$$

$$f_s(t) = \begin{cases} 1 & \text{if } 0 \leq t < \frac{T}{4} \text{ or } \frac{3T}{4} < t \leq T \\ e & \text{if } \frac{T}{4} \leq t \leq \frac{3T}{4} \end{cases}.$$

We set  $J_z = 0.1$  in the following. For  $-1 \leq \overline{\log J} \leq 0$  the drive straddles the phase transition, up to small corrections to its location due to the interaction.

Drives consistent with Floquet localization require both small interactions and not too small frequencies. We arrange the latter by defining, for each set of  $(\overline{\log \bar{J}}, \overline{\log \bar{h}})$  parameters, an effective “single-particle bandwidth”,  $W = \max(\sigma_J, \sigma_h)$ , where  $\sigma_h$  and  $\sigma_J$  are the standard deviations of  $h_i$  and  $J_i$  determined from the

parameters of the underlying log-normal distributions. The period is then defined by  $\omega = 2\pi/T = 2W$ . This choice ensures a roughly constant ratio of  $\omega/W$  for different  $\overline{\log J} - \overline{\log \hbar}$  values and thus isolates the effect of tuning the means through the phase diagram.

The lowest frequency in our drives is bigger than the estimated single particle bandwidth but much smaller than the many body bandwidth, so that localization is not a foregone conclusion. In Fig. 1 we characterize the quasi-energy spectrum  $\epsilon_n \in [0, 2\pi)$  using the level statistics of  $H_F$ . We define quasi-energy gaps by  $\delta_n = \epsilon_{n+1} - \epsilon_n$  and the level-statistics ratio  $r = \min(\delta_n, \delta_{n+1})/\max(\delta_n, \delta_{n+1})$ . An average value  $\langle r \rangle$  over all quasienergies and disorder realizations approaching the Poisson limit of .386 signals a lack of level repulsion and hence MBL and ergodicity-breaking. Conversely, delocalization implies the COE value of .527. The trends with increasing system size in Fig. 1 show localization away from the critical region which—given the weak interactions and large frequency of the drive—is close to the undriven, non-interacting transition point  $\overline{\log J} = \overline{\log \hbar}$ . In the interacting critical region we find a peak in  $\langle r \rangle$  which does not grow with system size and is much less than the COE value; we return to this below.

With localization established, we turn to distinguishing the phases. We consider a pair of  $Z_2$  invariant correlators (with  $A = x$  or  $y$ )

$$C_{AA}^\alpha(ij; t) = \langle \phi_\alpha(t) | \sigma_i^A \sigma_j^A | \phi_\alpha(t) \rangle \quad (5)$$

for  $i-j \gg 1$  in any given Floquet eigenstate. We find that for  $\overline{\log J} < \overline{\log \hbar}$  both correlators vanish with increasing system size  $L$  at all  $t$ , signaling a PM, while for  $\overline{\log J} > \overline{\log \hbar}$  are both generically non-zero, though of random sign varying with eigenstate and location, signaling SG order. For the latter, our parameters give  $|C_{xx}^\alpha(ij; t)| \gg |C_{yy}^\alpha(ij; t)|$  at all  $t$  although the more general signature is that  $|C_{xx}^\alpha(ij; t)|$  and  $|C_{yy}^\alpha(ij; t)|$  do not cross for  $0 < t < T$ . For our parameters, it suffices to compute

$$\chi_\alpha^{SG}(t) = \frac{1}{L^2} \sum_{i,j=1}^L (\langle \phi_\alpha(t) | \sigma_i^x \sigma_j^x | \phi_\alpha(t) \rangle)^2 \quad (6)$$

for  $t = 0$  as its variation during the period is qualitatively insignificant. In Fig. 1 (inset) we plot  $\chi_\alpha^{SG}(0)$  averaged over 2000–100,000 disorder realizations for different  $L$ s; the trend indicates that for  $L \rightarrow \infty$ ,  $\chi^{SG}(t) > 0$  in the spin-glass and  $\chi^{SG}(t) \rightarrow 0$  in the paramagnet, thereby sharply distinguishing the two phases.

Three comments are in order. First, recall that it would be sufficient to establish the existence of the Floquet PM—the SG can be obtained by duality<sup>40</sup>. Second, in chains with uniform couplings, both spin and dual spin order vanish in all but one of the Floquet eigenstates (the notion of the “ground-state” is not well defined in a Floquet system) even without interactions—this is the Landau-Peierls prohibition against discrete symmetry breaking in disguise. Localization is essential to avoid this. Third, in the fermionic formulation of the

localized problem, the PM is topologically trivial while the SG is non-trivial. The non-interacting SG phase has zero energy edge Majorana modes in open chains. With interactions, the edge modes remain coherent only in the MBL setting<sup>18,19</sup>. This implies that with localization, the *many-body* Floquet eigenstates on open chains come in conjugate, almost degenerate pairs which can be connected by spectrum generating operators localized near the two edges. In the free fermion problem, such operators (the Dirac fermion creation operators on either edge are examples) toggle the occupation of the bilocal edge Dirac mode and thus connect conjugate many-body states; in spin language, the operators toggle between many-body states with different parity. More concretely, the spectral function of  $\sigma_1^+$ , the spin raising operator on the left edge, in the Floquet eigenbasis

$$\mathcal{A}(\omega) = \frac{1}{2L} \sum_{\alpha\beta} \langle \phi_\alpha(0) | \sigma_1^+ | \phi_\beta(0) \rangle \delta(\omega - (\epsilon_\alpha - \epsilon_\beta)) \quad (7)$$

is a delta function peaked at  $\omega = 0$  (this phase will hence also be labeled the ‘0’ phase below). Finally, we note that the SG displays long-range string order in all eigenstates regardless of boundary conditions. Without disorder, the string order vanishes even in the many body eigenstates of free fermion chains—despite the non-trivial momentum space topology present in their Hamiltonians.

**Paramagnet-Spin Glass Phase transition:** In the non-interacting problem we have strong evidence that the infinite disorder fixed point continues to control the physics near quasienergy zero. We have examined  $H_F$  and we find that all its eigenstates are localized even at the transition, and its structure differs from the canonical strong disorder renormalization group form<sup>37</sup> by short ranged, irrelevant, terms. The ultimate fate of the critical region in the interacting driven problem is an interesting open question<sup>41</sup>, but we note for now that our data on  $\langle r \rangle$  suggests a partially delocalized interacting critical point.  **$\pi$  Spin Glass and  $0\pi$  Paramagnet:** We now present two new Ising phases which exist only in the driven system—the  $\pi$ -SG and the  $0\pi$ -PM. In identifying these new Floquet phases, we have built on existing work on the band topology of translationally invariant free fermion chains and related open spin chains. This work<sup>22</sup> has shown that the Floquet eigenmodes for such chains with *open* boundary conditions can exhibit edge Majorana modes with  $\epsilon_\alpha = \pi/T$  in addition to the better known edge modes with  $\epsilon_\alpha = 0$ . In the MBL setting, this implies that in the ‘ $\pi$ ’ phase the many-body Floquet eigenstates come in pairs separated by quasienergy  $\pi/T$  which can be connected by spectrum generating operators localized near the two edges, *i.e.* the edge spectral function  $\mathcal{A}(\omega)$  (7) shows a delta function peak at  $\omega = \pi/T$ .

We now establish these phases for the binary periodic drive

$$H(t) = \begin{cases} H_z & \text{if } 0 \leq t < T_1 \\ H_x & \text{if } T_1 \leq t < T = T_1 + T_2 \end{cases} \quad (8)$$




FIG. 2. (a) Phase diagram for the binary Ising drive, Eq. (8) without interactions ( $J_z = 0$ ) and disorder. (b) Level Statistics  $\langle r \rangle$  of  $H_F$  in the  $\pi$  phase with parameters defined in the text and disorder averaged over 2000-100,000 realizations for different  $L$ s.  $H_F$  is localized for interaction strengths  $J_z \lesssim 0.1$ . (c) Disorder averaged spectral function  $\mathcal{A}(\omega)$  defined in Eq. (7). Solid (dashed) lines are for the  $\pi$ -SG (0-SG) phase showing a delta function peak at  $\omega = \pi/T$  (0) for small interaction strengths which disappears as the interaction is increased. (d) Time dependence of the  $C_{xx}$  and  $C_{yy}$  correlators defined in Eq. (5) for an eigenstate in the  $\pi$  phase;  $L = 10$  and  $J_z = 0.04$ . The crossings are robust in the  $\pi$  phase.

$$H_z = \sum_{i=1}^L h_i \sigma_i^z + \sum_{i=1}^{L-1} J_z \sigma_i^z \sigma_{i+1}^z,$$

$$H_x = \sum_{i=1}^{L-1} J_i \sigma_i^x \sigma_{i+1}^x + J_z \sigma_i^z \sigma_{i+1}^z.$$

For  $J_z = 0$  and  $h_i = h$ ,  $J_i = J$  these are, essentially, the free fermion drives considered in Refs. 42 and 43. Fig. 2(a) shows the uniform, non-interacting phase diagram with the four possible driven Ising phases. The phases labeled ‘0’ and ‘ $\pi$ ’ have edge Majorana modes at quasienergies 0 and  $\pi/T$  respectively. With disorder and localization, these phases display long-range SG eigenstate order in the correlators (5) for both  $A = x, y$ . Moreover, in the  $\pi$ -SG phase, the time dependence of the  $C_{xx}$  and  $C_{yy}$  correlators over the period is non-trivially correlated: their magnitudes must cross twice during a period. Thus, in this phase, the axis of SG order rotates by an angle  $\pi$  about the  $z$ -axis during the period which can be intuitively understood by thinking semi-classically about the drive (8) at the extremal boundaries of the phase diagram shown in Fig. 2(a). As before, without localization, only one of the Floquet eigenstates will display long-range order in the 0,  $\pi$  phases. The other two phases, labeled PM and  $0\pi$ , have no long-range order and are respectively dual to the 0 and  $\pi$  phases.

We now turn to numerically identifying the localized  $\pi$  phase with disorder and interactions. We pick  $h_i T_1$  uniformly from the interval (1.512, 1.551) and  $J_i T_2$  from (0.393, 1.492), so that all pairs of values  $(h_i T_1, J_j T_2)$  lie in the  $\pi/T$  Majorana region of the free uniform chains. We have confirmed that the free fermion disordered drive exhibits  $\pi/T$  Majoranas for open chains while all other modes are localized in the bulk. In Fig 2b we examine stability to interactions via  $\langle r \rangle$  and clearly observe a transition around  $J_z \approx 0.1$ , with the small  $J_z$  regime being the MBL phase we seek.

Fig. 2c shows the appearance of the  $\pi/T$  Majorana peak in the disorder averaged edge spectral function (7) for system size  $L = 10$  as the delocalization transition is

crossed by decreasing the interaction strength. In contrast, the dashed lines show the spectral function for a similar drive in the 0-SG phase, clearly showing a peak at  $\omega = 0$ . Finally, Fig 2d displays the anticipated time dependence of the SG order in the  $C_{xx}$  and  $C_{yy}$  correlators in a single eigenstate of the interacting system with  $J_z = 0.04$  and  $L = 10$ . The crossings in the correlator within the period are robust in the  $\pi$ -SG phase and topologically distinct from the correlators in the 0-SG phase where there are no crossings. We emphasize that this is a true bulk diagnostic of the phase which, unlike the edge spectral function, is insensitive to boundary conditions. We also note that the non-trivial spin dynamics captured by it *cannot* be obtained without localization.

Finally we turn to the  $0\pi$ -PM. The simplest approach to obtain this phase is to dualize the  $\pi$ -SG. Now the dual SG order is long ranged and rotates by  $\pi$  during the period. With open boundary conditions the duality leads to an additional two states and thus conjugate sets of four MB states—two degenerate pairs of states separated by quasienergy  $\pi/T$ —as required by the fermionic analysis. The edge spectral function shows delta function peaks at both 0 and  $\pi/T$  consistent with the presence of coherent Majorana modes at both 0 and  $\pi/T$  in the non-interacting (and disordered, interacting) problems.

**Summary and open questions:** We have shown that MBL Floquet systems exhibit sharply defined phases bounded by parameter surfaces across which properties of their Floquet eigensystems change in a singular fashion. These phases include the trivial Floquet-ergodic phase and multiple non-trivial non-ergodic phases exhibiting various forms of ordering and dynamics, some of which are entirely new to Floquet systems. While our work builds naturally on very recent progress in the understanding of MBL and its extension to Floquet systems, the net result is still something that is quite surprising given the contentious history of non-equilibrium statistical mechanics. Indeed, it is quite likely that Floquet systems constitute the maximal class for which such a definition of phase structure is possible; with generic time dependences it would not be surprising if heating to in-

finite temperatures is the inevitable result. Going forward we anticipate a more systematic search for Floquet phases and a better understanding of their phase diagrams. In this context we note two studies that include disorder, but not interactions, in Floquet systems<sup>44,45</sup>. One interesting question is whether it is possible for Floquet systems to exhibit coexistence of macroscopically distinct eigenstates at the same quasienergy. It should also be possible to observe the new localized phases by the methods of Ref. 46.

**Acknowledgements:** We are especially grateful to Curt von Keyserlingk and Rahul Roy for many helpful discussions on topology in Floquet systems and to Anushya Chandran, Arnab Das and David Huse for discussions on Floquet and MBL matters. This work was supported by NSF Grant No. 1311781 and the John Templeton Foundation (VK and SLS) and the Alexander von Humboldt Foundation and the German Science Foundation (DFG) via the Gottfried Wilhelm Leibniz Prize Programme at MPI-PKS (SLS) and the SFB 1143.

- 
- <sup>1</sup> J. H. Shirley, *Physical Review* **138**, B979 (1965).  
<sup>2</sup> Y. B. Zel'dovich, *Soviet Physics JETP* (1967).  
<sup>3</sup> H. Sambe, *Phys. Rev. A* **7**, 2203 (1973).  
<sup>4</sup> L. D'Alessio and M. Rigol, *Phys. Rev. X* **4** (2014), 10.1103/physrevx.4.041048.  
<sup>5</sup> A. Lazarides, A. Das, and R. Moessner, *Physical Review E* **90** (2014), 10.1103/physreve.90.012110.  
<sup>6</sup> P. Ponte, A. Chandran, Z. Papić, and D. A. Abanin, *Annals of Physics* **353**, 196 (2015).  
<sup>7</sup> T. Prosen, *Physical Review Letters* **80**, 1808 (1998).  
<sup>8</sup> R. Citro, E. G. D. Torre, L. D'Alessio, A. Polkovnikov, M. Babadi, T. Oka, and E. Demler, *Annals of Physics* **360**, 694 (2015).  
<sup>9</sup> A. Chandran and S. L. Sondhi, arXiv:1506.08836 (2015).  
<sup>10</sup> A. Lazarides, A. Das, and R. Moessner, *Phys. Rev. Lett.* **115** (2015), 10.1103/physrevlett.115.030402.  
<sup>11</sup> P. Ponte, Z. Papić, F. Huvneers, and D. A. Abanin, *Phys. Rev. Lett.* **114** (2015), 10.1103/physrevlett.114.140401.  
<sup>12</sup> D. Abanin, W. De Roeck, and F. Huvneers, arXiv:1412.4752 (2014).  
<sup>13</sup> D. Basko, I. Aleiner, and B. Altshuler, *Annals of Physics* **321**, 1126 (2006).  
<sup>14</sup> For a recent review see<sup>47</sup> and references therein.  
<sup>15</sup> D. A. Huse, R. Nandkishore, V. Oganesyan, A. Pal, and S. Sondhi, *Physical Review B* **88**, 014206 (2013).  
<sup>16</sup> D. Pekker, G. Refael, E. Altman, E. Demler, and V. Oganesyan, *Physical Review X* **4**, 011052 (2014).  
<sup>17</sup> R. Vosk and E. Altman, *Physical Review Letters* **112**, 217204 (2014).  
<sup>18</sup> A. Chandran, V. Khemani, C. Laumann, and S. Sondhi, *Physical Review B* **89**, 144201 (2014).  
<sup>19</sup> Y. Bahri, R. Vosk, E. Altman, and A. Vishwanath, *Nature Communications* **6** (2015).  
<sup>20</sup> D. J. Thouless, *Physical Review B* **27**, 6083 (1983).  
<sup>21</sup> T. Oka and H. Aoki, *Physical Review B* **79**, 081406 (2009).  
<sup>22</sup> L. Jiang, T. Kitagawa, J. Alicea, A. R. Akhmerov, D. Pekker, G. Refael, J. I. Cirac, E. Demler, M. D. Lukin, and P. Zoller, *Physical Review Letters* **106**, 220402 (2011).  
<sup>23</sup> T. Kitagawa, T. Oka, A. Brataas, L. Fu, and E. Demler, *Physical Review B* **84**, 235108 (2011).  
<sup>24</sup> N. H. Lindner, G. Refael, and V. Galitski, *Nature Physics* **7**, 490 (2011).  
<sup>25</sup> T. Kitagawa, E. Berg, M. Rudner, and E. Demler, *Physical Review B* **82**, 235114 (2010).  
<sup>26</sup> M. S. Rudner, N. H. Lindner, E. Berg, and M. Levin, *Physical Review X* **3**, 031005 (2013).  
<sup>27</sup> F. Nathan and M. S. Rudner, (2015), arXiv:1506.07647.  
<sup>28</sup> R. Roy, unpublished (2015).  
<sup>29</sup> M. Serbyn, Z. Papić, and D. A. Abanin, *Physical Review Letters* **111**, 127201 (2013).  
<sup>30</sup> J. Z. Imbrie, (2014), arXiv:1403.7837.  
<sup>31</sup> D. A. Huse, R. Nandkishore, and V. Oganesyan, *Physical Review B* **90**, 174202 (2014).  
<sup>32</sup> V. Ros, M. Müller, and A. Scardicchio, *Nuclear Physics B* **891**, 420 (2015).  
<sup>33</sup> A. Chandran, I. H. Kim, G. Vidal, and D. A. Abanin, *Physical Review B* **91**, 085425 (2015).  
<sup>34</sup> A. Pal and D. A. Huse, *Physical Review B* **82**, 174411 (2010).  
<sup>35</sup> V. Oganesyan and D. A. Huse, *Phys. Rev. B* **75**, 155111 (2007).  
<sup>36</sup> M. Žnidarič, T. Prosen, and P. Prelovšek, *Physical Review B* **77**, 064426 (2008).  
<sup>37</sup> D. S. Fisher, *Physical Review B* **51**, 6411 (1995).  
<sup>38</sup> D. Pekker, G. Refael, E. Altman, E. Demler, and V. Oganesyan, *Physical Review X* **4**, 011052 (2014).  
<sup>39</sup> J. A. Kjäll, J. H. Bardarson, and F. Pollmann, *Physical Review Letters* **113**, 107204 (2014).  
<sup>40</sup> In the tails of the SG phase,  $\log(J) - \overline{\log(h)} \gtrsim 2$ , the values of the couplings  $J$  are much bigger than the fixed interaction strength  $J_z = 0.1$  so the problem effectively looks non-interacting. The PM does not suffer from this problem, and the strongly localized, interacting SG can be simply obtained by dualizing the PM.  
<sup>41</sup> V. Khemani, D. A. Huse, and S. L. Sondhi, *In Progress*.  
<sup>42</sup> V. M. Bastidas, C. Emary, G. Schaller, and T. Brandes, *Phys. Rev. A* **86**, 063627 (2012).  
<sup>43</sup> M. Thakurathi, A. A. Patel, D. Sen, and A. Dutta, *Physical Review B* **88**, 155133 (2013).  
<sup>44</sup> P. Titum, N. H. Lindner, M. C. Rechtsman, and G. Refael, *Physical review letters* **114**, 056801 (2015).  
<sup>45</sup> P. Titum, E. Berg, M. S. Rudner, G. Refael, and N. H. Lindner, arXiv preprint arXiv:1506.00650 (2015).  
<sup>46</sup> M. Schreiber, S. S. Hodgman, P. Bordia, H. P. Lüschen, M. H. Fischer, R. Vosk, E. Altman, U. Schneider, and I. Bloch, arXiv preprint arXiv:1501.05661 (2015).  
<sup>47</sup> R. Nandkishore and D. A. Huse, *Annual Review of Condensed Matter Physics* **6**, 15 (2015).