

Corona graphs as a model of small-world networks

Qian Lv, Yuhao Yi and Zhongzhi Zhang*

Shanghai Key Laboratory of Intelligent Information Processing, School of Computer Science, Fudan University, Shanghai 200433, China

Abstract

We introduce recursive corona graphs as a model of small-world networks. We investigate analytically the critical characteristics of the model, including order and size, degree distribution, average path length, clustering coefficient, and the number of spanning trees, as well as Kirchhoff index. Furthermore, we study the spectra for the adjacency matrix and the Laplacian matrix for the model. We obtain explicit results for all the quantities of the recursive corona graphs, which are similar to those observed in real-life networks.

Keywords: Corona product, Small-world, Spectra, Laplacian matrix

1. Introduction

For decades we want to know what a graph looks like. We want to reveal the principles of the networks' behaviour covered by their complex topology and dynamics. We want to learn about how the network structure evolves over time and how it affects the properties of dynamical processes on it. For the nature of decentrality of real networks, it is hard to observe the networks directly. Instead we observe them by taking snapshots of their network structure and content and keep updating them. Yet this gives little information about the future, since many of them keep growing over time. Thus it is desirable to set up models to fit real networks in both structure and functionality. We can use the models to mimic real-life networks. We also expect that the properties of the models can be proven rigorously, thus we can find the relations between topological and dynamical properties of networks.

*Corresponding author. E-mail address: zhangzz@fudan.edu.cn

Even if it is hard to give closed-formed expressions for some quantities, it would be nice to make them tractable for convenience of estimation.

Among various network models, the ER graph proposed by Erdős and Rényi [1] is the earliest one. It generates a random graph by choosing a constant probability for joining every pair of vertices in the network. The model exhibit interesting statistical properties and has been well studied by many people. However the model lacks some important properties of real-world networks. For example many real-world networks exhibit small-world property [2, 3, 4] with their diameters growing logarithmically in the number of vertices, while maintaining a high clustering coefficient. The Watts-Strogatz (WS) model [3] is a typical graph model with the small-world effect. Nevertheless, because of its randomness, many of its properties cannot be derived precisely, for example eigenvalues of the adjacency and Laplacian matrices. Thus deterministic models are often used to mimic complex networks [5], since their structural [6] and spectral [7] characteristics can be determined analytically. In addition to the small-world effect, another important feature of a network is degree distribution. Many real-world networks exhibit a heavy-tail distribution while some networks have an exponential distribution [8, 9, 10]. The famous preference attachment [11, 12, 13, 14] scheme successfully described the growing process of networks with heavy-tailed degree distribution. This work, like the WS model [3], leads to a network with an exponential degree distribution.

Recently graph (matrix) products have been applied to modelling graphs with the same properties as real-life networked systems, such as the Cartesian product [15], dot product [16] and Kronecker product [17, 18, 19, 20, 21]. A merit of such methods is that the graph/matrix products facilitate estimate of the properties of the generated graphs.

In this paper we introduce a recursive way to generate small-world networks with an exponential degree distribution, based on corona product of graphs. We obtain exact solutions to many structural properties of the networks. Moreover, we derive all the eigenvalues for their adjacency matrix and Laplacian matrix, which are provided in a recursive way. Based on the obtained eigenvalues, we calculate the number of spanning trees, as well as the Kirchhoff index of the networks.

2. Graph Construction

In this paper, we use corona product to generate a small-world network model. Literatures about the corona product and its related graphs are partly established [22, 23]. Let $G = (V(G), E(G))$ be the embedded graph of a network. Suppose the graph is undirected and has vertex set $V(G) = \{1, 2, \dots, N\}$ and edge set $E(G) \subseteq V(G) \times V(G)$. We define the number of vertices N as the order of the graph, and the number of edges as the size of the graph, denoted as $M = |E(G)|$.

Given two graphs G_1 and G_2 , their corona product $G_1 \circ G_2$ is defined as follows.

Definition 2.1. *Let G_1 and G_2 be two graphs with disjoint vertex sets. G_1 has N_1 vertices and G_2 has N_2 vertices. Their corona product $G_1 \circ G_2$ is a new graph which consists of one copy of G_1 and N_1 copies of G_2 . The i -th vertex of G_1 is joint by a new edge with every vertex in the i -th copy of G_2 .*

In this paper we investigate the case where G_2 is the q -complete graph, thus we give the definition of the recursive corona graph.

Definition 2.2. *Let K_q be the q -complete graph ($q \geq 2$), then the g th generation of recursive corona graph (RCG) $C_q(g+1)$ is defined as the corona product of the previous generation of RCG $C_q(g)$ and K_q . More formally, $C_q(g+1)$ is defined as $C_q(g+1) = C_q(g) \circ K_q$, $g \geq 0$, with the initial condition $C_q(0) = K_q$.*

Figure 1 illustrates the construction process for a particular network $C_4(g)$.

3. Structural properties

In this section we derive several important quantities of the RCG, showing that it is an appropriate model for the small-world complex networks. Thanks to the deterministic feature of $C_q(g)$, we can give exact expressions for the properties of the graph. We will give the explicit result for its order, size, degree distribution, degree correlation, average distance, clustering coefficient, number of spanning tree and Kirchhoff index.

We denote by $N(g)$ and $M(g)$ respectively the order and the size of $C_q(g)$. Next we show how to derive these quantities. Assume that the number of vertices and the number of edges that get newly generated at step g is




Figure 1: Construction of the recursive corona graph, showing the first three generations of $C_4(g)$

denoted as $L_V(g)$ and $L_E(g)$. Then it is obvious that we have $L_V(g) = q \times N(g - 1)$ for $g \geq 1$, which leads to the result of $N(g)$ along with the initial condition $N(0) = q$. With respect to the size of the network, we have $L_E(g) = q(q - 1)/2 \cdot N(g - 1) + q \cdot N(g - 1)$, $g \geq 1$, and the initial condition $M(0) = q(q - 1)/2$.

Proposition 3.1. *The order and size of the graph $C_q(g) = (V(g), E(g))$ are, respectively.*

$$N(g) = q(q + 1)^g \quad (1)$$

and

$$M(g) = \frac{1}{2}q((q + 1)^{g+1} - 2) \quad (2)$$

The average degree is $\bar{\delta}(g) = -2(q + 1)^{-g} + q + 1$ which tends to $q + 1$ for large g . Note that many real-life networks are sparse and their average degree tend to a constant value.

3.1. Degree distribution

The degree distribution $\mathcal{P}(\delta)$ for a network is a function indicating the fraction of vertices with degree δ over all vertices. The degree distribution is a very important characteristic of a graph. It is essential to the analysis of many other structural properties.

The cumulative degree distribution [8] is defined as

$$\mathcal{P}_{\text{cum}}(\delta) = \sum_{\delta'=\delta}^{\infty} \mathcal{P}(\delta'),$$

which is often used to analyse the degree distribution of a graph. The quantity gives the fraction of vertices whose degree δ' is greater than or equal to δ . In addition, networks whose degree distributions are exponential: $\mathcal{P}(\delta) \sim e^{-\alpha\delta}$, have also an exponential cumulative distribution with the same exponent:

$$\mathcal{P}_{\text{cum}}(\delta) = \sum_{\delta'=\delta}^{\infty} P(\delta') \approx \sum_{\delta'=\delta}^{\infty} e^{-\alpha\delta'} = \left(\frac{e^\alpha}{e^\alpha - 1} \right) e^{-\alpha\delta}. \quad (3)$$

Next we investigate the degree distribution of $C_q(g)$. We find that at time $g = 0$, the network has q vertices of degree $q - 1$. Now we study the degree of some vertex v at step g . Let the value be $\delta_v(g)$, we look in details about how the quantity evolves. We assume that vertex v is added to the network at step g_v , ($g_v \geq 0$). For any $g_v > 0$, there are $\delta_v(g_v) = q$, where $q - 1$ edges link to other vertices in K_q and the other 1 edge links to $C_q(g - 1)$. At every step every existed vertex increase its degree by q .

Theorem 3.2. *The cumulative degree distribution of the graph $C_q(g)$ follows an exponential distribution: $\mathcal{P}_{\text{cum}}(\delta) \sim (q + 1)^{-\frac{\delta}{q}+1}$.*

Proof. The degree of vertex v at step g , denoted as $\delta_v(g)$, can be written as

$$\delta_v(g + 1) = \delta_v(g) + q. \quad (4)$$

Thus we have

$$\begin{aligned} \delta_v(g) &= q \cdot (g - g_v + 1), (g_v > 0), \text{ and} \\ \delta_v(g) &= q \cdot (g + 1) - 1, (g_v = 0). \end{aligned} \quad (5)$$

This means that the numbers of vertices with the degree equal to $q, 2q, \dots, gq, (g+1)q - 1$ are, respectively, $q^2(q + 1)^{g-1}, q^2(q + 1)^{g-2}, \dots, q^2, q$.

For a certain value of degree δ , we have $\mathcal{P}_{\text{cum}}(\delta) = \left(N(0) + \sum_{g'=1}^{\theta} L_V(g') \right) / N(g)$ where $\theta = \lfloor g - (\delta + q)/q \rfloor$. Therefore we can find

$$\begin{aligned}
\mathcal{P}_{\text{cum}}(\delta) &= \frac{1}{q(q+1)^g} \left(q + \sum_{g'=1}^{\theta} q^2 (q+1)^{g'-1} \right) \\
&= \frac{1}{(q+1)^g} + \frac{q}{(q+1)^g} \sum_{g'=1}^{\theta} (q+1)^{g'-1} \\
&= (q+1)^{\theta-g}.
\end{aligned} \tag{6}$$

For large g we have

$$\mathcal{P}_{\text{cum}}(\delta) = (q+1)^{\theta-g} \sim (q+1)^{-\frac{\delta}{q}+1}.$$

□

3.2. Degree correlation

One important parameter for the degree correlation is the average degree of adjacent vertices of all vertices which is referred to as any vertex with degree δ . We denote the parameter by $k_{\text{nn}}(\delta)$. If $k_{\text{nn}}(\delta)$ increases with δ , this means that the vertices have a tendency to connect to vertices with a similar or larger degree. In this case we claim the graph to be assortative. The considered value of vertices with degree δ which is generated at step g_v can be written as

$$\begin{aligned}
k_{\text{nn}}(\delta) &= \frac{1}{L_V(g_v) \cdot q(g - g_v + 1)} \left(q^2(q(g+1) - 1) + \sum_{g_p=1}^{g_v-1} L_v(g_p) q \cdot q(g - g_p + 1) \right. \\
&\quad \left. + L_v(g_v)(q-1)q(g - g_v + 1) + L_v(g_v) \sum_{i=g_v+1}^g q \cdot q(g - i + 1) \right) \\
&= \frac{1}{2}q(g - g_v + 2) + \frac{1 + q - 2(1+q)^{-g_v+1}}{q(g - g_v + 1)}.
\end{aligned} \tag{7}$$

According to Eq.(5), we can express it as

$$k_{\text{nn}}(\delta) = \frac{1}{2}(q + \delta) + \frac{q + 1 - 2(1+q)^{-g+\frac{\delta}{q}}}{\delta}.$$

Since $q \leq \delta \leq q(g+1) - 1$, therefore we have $0 < (1+q)^{\frac{\delta}{q}-g} < 1$, which means

$$k_{\text{nn}}(\delta) \approx \frac{1}{2}(q + \delta) + \frac{q}{\delta}. \quad (8)$$

As for the initial vertices, everyone of them has the same distribution on the degrees of its neighbors. It leads to

$$\begin{aligned} k_{\text{nn}}(\delta_0) &= \frac{1}{q \cdot (g+1) - 1} \left((q-1)(q(g+1) - 1) + \sum_{i=1}^g q \cdot q(g-i+1) \right) \\ &= \frac{(g+2)q}{2} - \frac{(g+2)q-2}{2(gq+q-1)}, \end{aligned} \quad (9)$$

which yields

$$k_{\text{nn}}(\delta_0) = \frac{1}{2} \left(\delta_0 + q + \frac{1-q}{\delta_0} \right). \quad (10)$$

By checking the results we can see that the considered graph is assortative.

3.3. Average distance

Given a graph $G = (V, E)$, its *average distance* or *mean distance* is defined as: $\mu(G) = \frac{1}{|V(G)|(|V(G)|-1)} \cdot \sum_{u,v \in V(G)} d(u, v)$ where $d(u, v)$ is the distance between the pair of vertices u and v .

Theorem 3.3. *The average distance of graph $C_q(g)$ is*

$$\mu(C_q(g)) = \frac{(q+1)^{-g} (2gq^2(q+1)^{2g-1} + (q+1)^g + (q-2)((q+1)^2)^g)}{q(q+1)^g - 1}. \quad (11)$$

Proof. To begin with, we assume that the summation of distances between all pairs of vertices in $C_q(g)$ is $D(C_q(g))$. The sum of distances between all pairs (u, v) where u belongs to vertex set U and v belongs to a disjoint vertex set V , is denoted as $D(U, V)$.

In order to utilize the recursive construction process to the recursive corona graph we classify the vertex pairs in $C_q(g)$ into 4 different categories W, X, Y and Z . The sum of the distances for the 4 categories is denoted as S_W, S_X, S_Y and S_Z , respectively.

Category W refers to the pairs within the same K_q that we add to the network at step g . Category X refers to the pairs u, v where u is selected from one of the $N(g-1)$ K_q s added at step g and v selected from any other K_q added to the network at the same step. Y refers to the pairs where u is a

new vertex and v is a vertex in $C_q(g-1)$. As for category Z , it indicates the pairs where both u and v are from the previous generation of graph $C_q(g-1)$. Thus we have the following equations:

$$D(C_q(g)) = S_W(g) + S_X(g) + S_Y(g) + S_Z(g), \quad (12)$$

$$S_W(g) = \frac{q(q-1)}{2} \cdot N(g-1), \quad (13)$$

$$\begin{aligned} S_X(g) &= \sum_{\substack{i,j \in V(C_q(g-1)) \\ i < j}} (d_{i,j} + 2) \cdot q^2 \\ &= q^2 S_Z(g) + q^2 N(g-1) \cdot (N(g-1) - 1), \end{aligned} \quad (14)$$

$$\begin{aligned} S_Y(g) &= \sum_{i,j \in V(C_q(g-1))} (d_{i,j} + 1) \cdot q \\ &= 2q S_Z(g) + q N(g-1) \cdot (N(g-1) - 1) + N(g-1) \cdot q, \end{aligned} \quad (15)$$

$$S_Z(g) = D(C_q(g-1)). \quad (16)$$

Combining these recursive expression together we have:

$$D(C_q(g)) = (q+1)^2 D(C_q(g-1)) + \frac{1}{2} q^2 (2q(q+1)^{g-1} - 1) (q+1)^g, \quad (17)$$

with the initial condition $D(C_q(0)) = q(q-1)/2$ we get the result

$$D(C_q(g)) = \frac{1}{2} q (2gq^2(q+1)^{2g-1} + (q+1)^g + (q-2) ((q+1)^2)^g). \quad (18)$$

Dividing $D(C_q(g))$ by $\frac{N(g)(N(g)-1)}{2}$ yields Eq.(11). For large g , we have $\mu(C_q(g)) \sim 2g \sim 2 \log_q N(g)$, which increases logarithmically with the network order. \square

3.4. Clustering coefficient

Clustering coefficient [3] is another crucial quantity used to characterize network structure. Many works about determining clustering coefficient and its related quantities are done on both graph models and graphs in reality [3, 24, 25, 26].

The clustering coefficient of vertex v is defined as the following quantity

$$c(v) = \frac{2\epsilon_v}{\delta_v(\delta_v - 1)}, \quad (19)$$

where ϵ_v is the number of edges between the neighbours of vertex v . The network clustering coefficient $C(G)$ is defined as the average of $c(v)$ among all vertices. That is,

$$c(G) = \frac{1}{|V(G)|} \sum_{v \in V(G)} c(v). \quad (20)$$

Theorem 3.4. *Let $v(\delta)$ be a vertex in $C_q(g)$ whose degree is δ . Except the initial vertices, its clustering coefficient is*

$$c(v(\delta)) = \frac{q-1}{\delta-1} \approx \frac{q-1}{\delta}. \quad (21)$$

Proof. Let us review the intermediate result in calculating the degree distribution that the number of vertices with degree $q, 2q, \dots, gq, (g+1)q-1$ are, respectively, $q^2(q+1)^{g-1}, q^2(q+1)^{g-2}, \dots, q^2, q$. Except the initial vertices, the clustering coefficient of other vertices follow the same rule, that is, a vertex with degree $k \cdot q$ has $k \cdot q$ neighbours, which are evenly distributed in k clusters. Each cluster forms a complete graph K_q . Thus the clustering coefficient of vertex v is derived as

$$c(v) = \frac{kq(q-1)/2}{kq(kq-1)/2} = \frac{q-1}{kq-1}, \quad (g_v > 0), \quad (22)$$

and for initial vertices

$$c(v) = \frac{(q-1)(q-2) + gq(q-1)}{(q(g+1)-1)(q(g+1)-2)}, \quad (g_v = 0). \quad (23)$$

Theorem 3.4 is naturally gained. \square

Theorem 3.5. *The clustering coefficient of RCG network $C_q(g)$ converge to*

$$c(C_q(g)) \sim \frac{q-1}{q+1} \Phi \left(\frac{1}{q+1}, 1, \frac{q-1}{q} \right), \quad (24)$$

when the network order is high enough. In the expression, Φ is the Lerch transcendent function. For large q , the clustering coefficient tends to 1.

Proof. From Eq.(20) and (23) we can obtain the clustering coefficient of $C_q(g)$

$$\begin{aligned}
c(C_q(g)) &= \frac{1}{N(g)} \left[\sum_{k=1}^g \frac{q-1}{kq-1} \cdot q^2(q-1)^{g-k} + \frac{(q-1)(gq+q-2)}{(q(g+1)-1)(q(g+1)-2)} \right] \\
&= \frac{q-1}{q+1} \left(\Phi \left(\frac{1}{q+1}, 1, \frac{q-1}{q} \right) \right. \\
&\quad \left. - \left(\frac{1}{q+1} \right)^g \Phi \left(\frac{1}{q+1}, 1, \frac{gq+q-1}{q} \right) + \frac{(q+1)^{-g+1}}{gq+q-1} \right).
\end{aligned} \tag{25}$$

When g is large enough, we obtain

$$c(C_q(g)) \sim \frac{q-1}{q+1} \Phi \left(\frac{1}{q+1}, 1, \frac{q-1}{q} \right), \tag{26}$$

which is high and tends to 1 when q is also large. Figure 2 gives the clustering coefficient of some networks while they grow. \square




Figure 2: Clustering coefficient of some recursive corona graphs.

3.5. Spanning trees

Next we derive the number of spanning trees in graph $C_q(g)$.

Theorem 3.6. *The number of spanning tree of $C_q(g)$ is*

$$N_{\text{tr}}(C_q(g)) = q^{q-2}(q+1)^{1-q} ((q+1)^{q-1})^{(q+1)^g}. \quad (27)$$

Proof. According to the Cayley's formula [27], the number of spanning trees of a complete graph K_q is equal to q^{q-2} . Since all vertices of a K_q added to the graph are connected to a vertex in the original graph, these $q+1$ vertices consist of a new complete graph K_{q+1} . Therefore we have the following recursive relation of the spanning trees of $C_q(g)$:

$$N_{\text{tr}}(C_q(g)) = ((q+1)^{q-1})^{N(g-1)} N_{\text{tr}}(g-1). \quad (28)$$

Together with the initial condition $N_{\text{tr}} = q^{q-2}$, we can derive the expression for N_{tr} ;

$$N_{\text{tr}}(C_q(g)) = q^{q-2}(q+1)^{1-q} ((q+1)^{q-1})^{(q+1)^g}. \quad (29)$$

□

3.6. Kirchhoff index

Resistance distance is an important character of a graph, which can imply many of its dynamic properties. The Kirchhoff index [28] of a graph refers to the sum of resistance between all vertex pairs in an associated electrical network obtained from the graph by replacing each edge of the graph by a unit resistance. Denote the effective resistance between vertices i and j as $r(i, j)$ or $r_{i,j}$, then the Kirchhoff index R_{Kr} of graph G is defined as

$$R_{\text{Kr}}(G) = \sum_{\substack{i,j \in V(G) \\ i < j}} r_{i,j}. \quad (30)$$

We denote the Kirchhoff index of graph $C_q(g)$ by $R_{\text{Kr}}(C_q(g))$.

Theorem 3.7. *The Kirchhoff index of $C_q(g)$ is*

$$R_{\text{Kr}}(C_q(g)) = (q^3(2g+1) - 2q - 1)(q+1)^{2g-2} + q(q+1)^{g-1}. \quad (31)$$

Proof. We denote by $r(U, V)$ as the sum of all effective resistance between pairs (u, v) in which u and v belong to two disjoint vertex set U and V respectively. Similar to the method we used in calculating the average distance, we classify these pairs into 4 categories W , X , Y and Z , where the definition is exactly the same as used in calculating the average distance. Then the sum of the distances for the four categories is denoted as R_W , R_X , R_Y and R_Z . We have the following equations:

$$R_{\text{Kr}}(C_q(g)) = R_W(g) + R_X(g) + R_Y(g) + R_Z(g), \quad (32)$$

$$R_W(g) = \frac{q(q-1)}{2} \cdot \frac{2}{q+1} \cdot N(g-1), \quad (33)$$

$$\begin{aligned} R_X(g) &= \sum_{\substack{i,j \in V(C_q(g-1)) \\ i < j}} (r_{i,j} + 2 \cdot \frac{2}{q+1}) \cdot q^2 \\ &= q^2 R_Z(g) + \frac{2 \cdot q^2 N(g-1) \cdot (N(g-1) - 1)}{q+1}, \end{aligned} \quad (34)$$

$$\begin{aligned} R_Y(g) &= \sum_{i,j \in V(C_q(g-1))} (r_{i,j} + \frac{2}{q+1}) \cdot q \\ &= 2q R_Z(g) + q \cdot \frac{2}{q+1} N(g-1) \cdot (N(g-1) - 1) \\ &\quad + q \cdot \frac{2}{q+1} \cdot N(g-1), \end{aligned} \quad (35)$$

$$R_Z(g) = R_{\text{Kr}}(C_q(g-1)), \quad (36)$$

which yields

$$R_{\text{Kr}}(C_q(g+1)) = q^2 (2q(q+1)^g - 1) (q+1)^g + (q+1)^2 R_{\text{Kr}}(C_q(g)). \quad (37)$$

Notice that the effective resistance between vertices v and u in a complete graph K_q is $2/q$ since the potential between any other vertices are identical, if we impose potential difference between u and v .

Along with the initial condition $R_k(C_q(0)) = q - 1$ we can deduce

$$\begin{aligned} R_{\text{Kr}}(C_q(g)) &= (q^3 - 2q - 1) (q+1)^{2g-2} + q (2q^2 g (q+1)^g + q + 1) (q+1)^{g-2} \\ &= (q^3 - 2q - 1 + 2gq^3) (q+1)^{2g-2} + (q+1)^{g-1} q \\ &= (q^3(2g+1) - 2q - 1) (q+1)^{2g-2} + q(q+1)^{g-1}. \end{aligned} \quad (38)$$

This completes the proof. \square

4. Spectral analysis

By convention the (unweighted) adjacency matrix $A(G)$ of a graph G is defined as a $N \times N$ matrix with the entry $a_{i,j}$ representing the number of edges incident with endpoints i, j . The degree matrix of G , $D(G)$ is defined as a diagonal matrix with its i th entry on the main diagonal equal to the degree of vertex i . We call $L(G) = D(G) - A(G)$ the Laplacian matrix of graph G . These matrices determine the structure of the graph, and the eigenvalues of $A(G)$ and $L(G)$ are sensitive to many of the structural properties, which have remarkable impact on the dynamic processes superimposed upon the network.

Definition 4.1. *Given $A(C_q(g))$ the adjacency matrix of $C_q(g)$, we define the spectra of $C_q(g)$ as*

$$\sigma(C_q(g)) := \sigma(g) = (\lambda_1(g), \lambda_2(g) \cdots \lambda_n(g)). \quad (39)$$

Similarly, we have:

Definition 4.2. *Given $L(C_q(g))$ as the Laplacian matrix of $C_q(g)$, we define its Laplacian spectra*

$$S(C_q(g)) := S(g) = (\gamma_1(g), \gamma_2(g), \cdots, \gamma_n(g)). \quad (40)$$

4.1. Spectra of Adjacency Matrix

Theorem 4.1. *The relation between $\sigma(g)$ and $\sigma(g-1)$ is*

1. $\frac{\lambda_i(g-1)+q-1 \pm \sqrt{(q-1-\lambda_i(g-1))^2+4q}}{2} \in \sigma(g)$ with multiplicity 1 for $i = 1, \cdots, q(q+1)^{g-1}$ and
2. $-1 \in \sigma(g)$ with multiplicity $(q-1)q(q+1)^{g-1}$.

The result is a corollary of results in [22, 29].

4.2. Spectra of Laplacian Matrix

Theorem 4.2. *The relation between $S(g)$ and $S(g-1)$ is*

1. $\frac{\gamma_i(g-1)+q+1 \pm \sqrt{(\gamma_i(g-1)+q+1)^2-4\gamma_i(g-1)}}{2} \in S(g)$ with multiplicity 1 for $i = 1, \cdots, q(q+1)^{g-1}$ and
2. $q+1 \in S(g)$ with extra multiplicity $(q-1)q(q+1)^{g-1}$.

Note that in the first part $\gamma_i(g-1) = 0$ will generate an eigenvalue equal to $q+1$ with multiplicity 1 in iteration g . So the actual multiplicity of $q+1$ is $(q-1)q(q+1)^{g-1} + 1$ for any $g \geq 1$. The proof of Theorem 4.2 is evident using methods in [22, 23, 29]. For convenience of the following discussion we give a similar proof here:

Proof. The Laplacian matrix of $C_q(g)$ is

$$L(C_q(g)) = \left(\begin{array}{c|ccc} L(C_q(g-1)) + qI_n & -I_n & \cdots & -I_n \\ \hline -I_n & & & \\ \vdots & & & \\ -I_n & & (L(K_q) + I_q) \otimes I_n & \end{array} \right) \quad (41)$$

Let $Y_1, \dots, Y_{N(g-1)}$ be the Laplacian eigenvectors of $C_q(g-1)$ corresponding to the eigenvalues $\gamma_1(g-1), \gamma_2(g-1), \dots, \gamma_{N(g-1)}(g-1)$, respectively. For $i = 1, \dots, N(g-1)$, let

$$\phi_i = \frac{\gamma_i(g-1) + q + 1 + \sqrt{(\gamma_i(g-1) + q + 1)^2 - 4\gamma_i(g-1)}}{2},$$

$$\hat{\phi}_i = \frac{\gamma_i(g-1) + q + 1 - \sqrt{(\gamma_i(g-1) + q + 1)^2 - 4\gamma_i(g-1)}}{2}.$$

Note that $\phi_i, \hat{\phi}_i$ are Laplacian eigenvalues of $C_q(g)$ corresponding to the eigenvectors

$$\begin{pmatrix} Y_i \\ g(\phi_i)Y_i \\ \vdots \\ g(\phi_i)Y_i \end{pmatrix}, \begin{pmatrix} Y_i \\ g(\hat{\phi}_i)Y_i \\ \vdots \\ g(\hat{\phi}_i)Y_i \end{pmatrix},$$

respectively. In fact ϕ_i is obtained by solving

$$\begin{aligned}
\phi_i \begin{pmatrix} Y_i \\ g(\phi_i)Y_i \\ \vdots \\ g(\phi_i)Y_i \end{pmatrix} &= L(C_q(g)) \begin{pmatrix} Y_i \\ g(\phi_i)Y_i \\ \vdots \\ g(\phi_i)Y_i \end{pmatrix} \\
&= \begin{pmatrix} L(C_q(g-1)) + qI_n \\ -I_n \\ \vdots \\ -I_n \end{pmatrix} Y_i + \sum_{j=1}^q \begin{pmatrix} -I_n \\ (L(K_q) + I_n)_{1j}I_n \\ \vdots \\ (L(K_q) + I_n)_{qj}I_n \end{pmatrix} g(\phi_i)Y_i \quad (42) \\
&= \begin{pmatrix} \gamma_i(g-1) + q - q \cdot g(\phi_i) \\ -1 + g(\phi_i) \\ \vdots \\ -1 + g(\phi_i) \end{pmatrix} Y_i.
\end{aligned}$$

Thus we can derive the following equations:

$$\gamma_i(g-1) + q - q \cdot g(\phi_i) = \phi_i, \quad (43)$$

$$-1 + g(\phi_i) = \phi_i \cdot g(\phi_i). \quad (44)$$

From Eq.(44) we can obtain that $g(\phi_i) \neq 0$. Therefore we can substitute Eq.(44) into Eq.(43), we have:

$$(\gamma_i(g-1) + q - \phi_i)(\phi_i - 1) = -q, \quad (45)$$

which leads to the result of the first part of the theorem.

If the Laplacian eigenvalues $\nu_1 = 0, \nu_2 = \nu_3 = \dots = \nu_q = q$ of $L(K_q)$ are correlated with the eigenvectors Z_1, Z_2, \dots, Z_q , respectively, then for $j = 2, \dots, q$, we have

$$L(C_q(g)) \begin{pmatrix} \mathbf{0} \\ Z_j \otimes e_i \end{pmatrix} = (q+1) \begin{pmatrix} \mathbf{0} \\ Z_j \otimes e_i \end{pmatrix}. \quad (46)$$

This completes the proof. \square

Next we use the results of the Laplacian spectra to prove Theorem 3.6 and 3.7. First we give an alternative proof of Theorem 3.6.

Proof. It is known that the number of spanning tree of a graph G has the following form [30, 31]

$$N_{\text{tr}}(G) = \frac{\prod_{i=2}^N \tau_i}{N}, \quad (47)$$

where N is the number of vertices and τ_i refers to N eigenvalues of the graph G . Given the graph is connected, let τ_1 be the unique zero eigenvalue, then τ_i , $i = 2, \dots, N$ are $N - 1$ non-zero eigenvalues of the graph Laplacian.

Theorem 4.2 tells that the Laplacian spectrum of $C_q(g)$ consists of two parts. For the first part, we can derive from Eq.(45) that, in iteration g , γ_i in $C_q(g - 1)$ generates two eigenvalues ϕ_i and $\hat{\phi}_i$ which are subject to the relations $\phi_i \hat{\phi}_i = \gamma_i$ and $\phi_i + \hat{\phi}_i = \gamma_i + q + 1$. In particular the trivial eigenvalue $\gamma_1 = 0$ generates $\phi_i = q + 1$ and $\hat{\phi}_i = 0$. As for the second part, there is an eigenvalue $\gamma = q + 1$ with multiplicity $(q - 1)q(q + 1)^{g-1}$. We denote by $S(g)$ the sum of all non-zero eigenvalues of $L(C_q(g))$ and by $\Upsilon(g)$ the product of all non-zero eigenvalues of $L(C_q(g))$. Then we can obtain

$$\begin{aligned} \Upsilon(g) &= \prod_{i=2}^{N(g)} \gamma_i(g) = (q + 1)^{(q-1)q(q+1)^{g-1}+1} \prod_{i=2}^{N(g-1)} \phi_i \hat{\phi}_i \\ &= (q + 1)^{(q-1)q(q+1)^{g-1}+1} \prod_{i=2}^{N(g-1)} \gamma_i(g - 1) \\ &= \Upsilon(g - 1) \cdot q(q + 1)^{(q-1)q(q+1)^{g-1}+1}. \end{aligned} \quad (48)$$

Eq.(48) and the initial condition $\Upsilon(0) = q^{q-1}$ yield

$$\Upsilon(g) = q^{q-1} (q + 1)^{(q-1)((q+1)^g - 1) + g}. \quad (49)$$

Therefore

$$N_{\text{tr}}(C_q(g)) = q^{q-2} (q + 1)^{(q-1)((q+1)^g - 1)}.$$

□

The result is equivalent to what we derived using combinatorial method.

In the following we give an alternative proof of Theorem 3.7 using the spectra information.

Proof. The Kirchhoff index of a graph G can be expressed as [32, 33]:

$$R_{\text{Kr}}(G) = N \cdot \sum_{i=2}^N \frac{1}{\tau_i}, \quad (50)$$

where N and τ_i are the same as the previous definition. Let

$$S = \sum_{j=2}^N \prod_{\substack{i=2 \\ i \neq j}}^N \gamma_i = \prod_{i=2}^N \gamma_i \cdot \sum_{i=2}^N \frac{1}{\gamma_i}. \quad (51)$$

We can follow the clue of the previous analysis by separating the eigenvalues of its Laplacian matrix into two parts. Recall that the eigenvalues of the Laplacian consist of two parts $\Gamma^{(1)}$ and $\Gamma^{(2)}$ as defined by Theorem 4.2. Assume that $\Gamma^{(1)'} = \Gamma^{(1)} \setminus \{0\}$. For the first part of the eigenvalues of $L(C_q(g))$, we denote them as ϕ_i and $\hat{\phi}_i$, $i = 1, 2, \dots, N(g-1)$. Suppose the original eigenvalue in $L(C_q(g-1))$, which is correlated with ϕ_i and $\hat{\phi}_i$, is γ_i . Then

$$S(g) = S^{(1)}\Upsilon^{(2)} + S^{(2)}\Upsilon^{(1)}, \quad (52)$$

where

$$S^{(1)} = \sum_{\gamma_j \in \Gamma^{(1)'}} \prod_{\substack{\gamma_i \in \Gamma^{(1)' \\ i \neq j}} \gamma_i, \quad (53)$$

$$\gamma^{(2)} = \prod_{\gamma_i \in \Gamma^{(2)}} \gamma_i, \quad (54)$$

$$S^{(2)} = \sum_{\gamma_j \in \Gamma^{(2)}} \prod_{\substack{\gamma_i \in \Gamma^{(2)} \\ i \neq j}} \gamma_i, \quad (55)$$

and

$$\gamma^{(1)} = \prod_{\gamma_i \in \Gamma^{(1)}} \gamma_i. \quad (56)$$

Accordingly we can obtain

$$\begin{aligned}
S^{(1)} &= (q+1) \prod_{i=2}^{N(g-1)} \phi_i \hat{\phi}_i \cdot \left(\sum_{j=2}^{N(g-1)} \left(\frac{1}{\phi_j} + \frac{1}{\hat{\phi}_j} \right) \right) + \prod_{i=2}^{N(g-1)} \phi_i \hat{\phi}_i \\
&= (q+1) \sum_{j=2}^{N(g-1)} \left(\prod_{\substack{i=2 \\ i \neq j}}^{N(g-1)} \phi_i \hat{\phi}_i \cdot (\phi_j + \hat{\phi}_j) \right) + \prod_{i=2}^{N(g-1)} \phi_i \hat{\phi}_i \\
&= \Upsilon(g-1) + (q+1) \sum_{j=2}^{N(g-1)} \prod_{\substack{i=2 \\ i \neq j}}^{N(g-1)} \gamma_i(g-1) \cdot (\gamma_j(g-1) + q+1) \\
&= \Upsilon(g-1) + (q+1) \sum_{j=2}^{N(g-1)} \prod_{\substack{i=2 \\ i \neq j}}^{N(g-1)} \gamma_i(g-1) \cdot \gamma_j(g-1) \\
&\quad + (q+1) \sum_{j=2}^{N(g-1)} \prod_{\substack{i=2 \\ i \neq j}}^{N(g-1)} \gamma_i(g-1) \cdot (q+1) \\
&= (1 + (q+1) \cdot (q(q+1)^{g-1} - 1)) \cdot \Upsilon(g-1) + (q+1)^2 \cdot S(g-1).
\end{aligned} \tag{57}$$

$$\Upsilon^{(2)} = (q+1)^{(q-1)q(q+1)^{g-1}}, \tag{58}$$

$$S^{(2)} = (q-1)q(q+1)^{(q-1)q(q+1)^{g-1}+g-2}, \tag{59}$$

$$\Upsilon^{(1)} = (q+1) \cdot \Upsilon(g-1). \tag{60}$$

Therefore we obtain a recursive relation of $S(g)$. Considering the initial condition $S(0) = (q-1)q^{q-2}$, we derive

$$S(g) = q^{q-2}(q+1)^{(q-1)(q+1)^g+g-q-1} \left(((2g+1)q^3 - 2q - 1)(q+1)^g + q(q+1) \right), \tag{61}$$

thus

$$\begin{aligned}
R_{\text{Kr}}(C_q(g)) &= (q+1)^{g-2} \left(((2g+1)q^3 - 2q - 1)(q+1)^g + q(q+1) \right) \\
&= (q^3(2g+1) - 2q - 1)(q+1)^{2g-2} + q(q+1)^{g-1},
\end{aligned} \tag{62}$$

□

Note that Eq. (62) is consistent with Eq. (27). For large g , the Kirchhoff index displays the following leading behavior:

$$R_{\text{Kr}}(C_q(g)) \sim g \cdot N(g)^2 \sim N(g)^2 \log_{q+1} N(g), \tag{63}$$

5. Conclusion

In this paper, we have introduced a deterministically growing model to generate small-world graph, by using the corona product. The advantage of such a model is that many of its properties can be solved exactly. We have derived explicitly many structural quantities of the small-world model. We have also found the eigenvalues for the adjacency matrix and the Laplacian matrix of the model. In future, the properties of various dynamical processes taking place on the small-world model deserve study.

Acknowledgement

This work was supported by the National Natural Science Foundation of China under Grant No. 11275049.

References

- [1] Erdős P and Rényi A 1960 On the evolution of random graphs *Publ. Math. Inst. Hung. Acad. Sci.* **5** 17–61
- [2] Milgram S 1967 The small world problem *Psychol. Today* **2** 60–67
- [3] Watts D J and Strogatz S H 1998 Collective dynamics of ‘small-world’ networks *Nature* **393** 440–442
- [4] Kleinberg J 2000 The small-world phenomenon: An algorithmic perspective *Proc. 32nd Annual ACM Symp. on Theory of Computing (ACM)* pp 163–170
- [5] Zhang Z and Comellas F 2011 Farey graphs as models for complex networks *Theor. Comput. Sci.* **412** 865 – 875
- [6] Zhang Z and Wu B 2015 Pfaffian orientations and perfect matchings of scale-free networks *Theor. Comput. Sci.* **570** 55–69
- [7] Liu H, Dolgushev M, Qi Y and Zhang Z 2015 Laplacian spectra of a class of small-world networks and their applications *Sci. Rep.* **5** 9024
- [8] Newman M E J 2003 The structure and function of complex networks *SIAM Rev.* **45** 167–256

- [9] Barrat A and Weigt M 2000 On the properties of small-world network models *Eur. Phys. J. B* **13** 547–560
- [10] Ferrer-i-Cancho R, Janssen C and Solé R V 2001 Topology of technology graphs: Small world patterns in electronic circuits *Phys. Rev. E* **64** 046119
- [11] Barabási A L and Albert R 1999 Emergence of scaling in random networks *Science* **286** 509–512
- [12] Albert R and Barabási A L 2002 Statistical mechanics of complex networks *Rev. Mod. Phys.* **74** 47
- [13] Kleinberg J M, Kumar R, Raghavan P, Rajagopalan S and Tomkins A S 1999 The web as a graph: measurements, models, and methods computing and combinatorics *Computing and Combinatorics (Lecture Notes in Computer Science vol 1627)* (Springer) pp 1–17
- [14] Kumar R, Raghavan P, Rajagopalan S and Tomkins A 1999 Extracting large-scale knowledge bases from the web *Proc. 25th Int. Conf. on Very Large Data Bases* (Morgan Kaufmann Publishers Inc.) pp 639–650
- [15] Imrich W and Klavžar S 2000 *Product graphs, structure and recognition (Wiley Series in Discrete Mathematics and Optimization vol 56)* (Wiley-Interscience)
- [16] Young S J and Scheinerman E R 2007 Random dot product graph models for social networks *Proc. 5th Int. Conf. on Algorithms and Models for the Web-graph* pp 138–149
- [17] Weichsel P M 1962 The Kronecker product of graphs *Proc. Amer. Math. Soc.* **13** 47–52
- [18] Leskovec J and Faloutsos C 2007 Scalable modeling of real graphs using Kronecker multiplication *Proc. 24th Int. Conf. on Machine Learning ICML '07* (ACM) pp 497–504
- [19] Mahdian M and Xu Y 2007 Stochastic Kronecker graphs *Proc. 5th Workshop on Algorithms and Models for the Web-Graph* (Springer) pp 179–186

- [20] Leskovec J 2009 Networks, communities and Kronecker products *Proc. 1st ACM Int. Workshop on Complex Networks Meet Information & Knowledge Management* (ACM) pp 1–2
- [21] Leskovec J, Chakrabarti D, Kleinberg J, Faloutsos C and Ghahramani Z 2010 Kronecker Graphs: An Approach to Modeling Networks *J. Mach. Learn. Res.* **11** 985–1042
- [22] Barik S, Pati S and Sarma B K 2007 The spectrum of the corona of two graphs *SIAM J. Discrete Math.* **21** 47–56
- [23] Liu Q 2014 The Laplacian spectrum of corona of two graphs *Kragujevac J. Math.* **38** 163–170
- [24] Holme P and Kim B J 2002 Growing scale-free networks with tunable clustering *Phys. rev. E* **65** 026107
- [25] Tsourakakis C E 2008 Fast counting of triangles in large real networks without counting: Algorithms and laws *Proc. 8th IEEE Int. Conf. on Data Mining* (IEEE Computer Society) pp 608–617
- [26] Tsourakakis C E 2011 Counting triangles in real-world networks using projections *Knowl. Inf. Syst.* **26** 501–520
- [27] Cayley A 1889 A theorem on trees *Quart. J. Math* **23** 376–378
- [28] Klein D J and Randić M 1993 Resistance distance *J. Math. Chem.* **12** 81–95
- [29] Sharma R, Adhikari B and Mishra A 2015 On spectra of corona graphs *Algorithms and Discrete Applied Mathematics (Lecture Notes in Computer Science* vol 8959) (Springer) pp 126–137
- [30] Biggs N 1994 *Algebraic graph theory* 2nd ed (Cambridge university press)
- [31] Tzeng W J and Wu F Y 2000 Spanning trees on hypercubic lattices and nonorientable surfaces *Appl. Math. Lett.* **13** 19–25
- [32] Aldous D and Fill J 2002 Reversible markov chains and random walks on graphs <http://www.stat.berkeley.edu/~aldous/RWG/book.html>
- [33] Ghosh A, Boyd S and Saberi A 2008 Minimizing effective resistance of a graph *SIAM Rev.* **50** 37–66