

# SAMPLING GOLDBACH NUMBERS AT RANDOM

**Ljuben Mutafchiev**

*Department of Mathematics and Science, American University in Bulgaria and  
Institute of Mathematics and Informatics of the Bulgarian Academy of Sciences,  
Blagoevgrad, 2700, Bulgaria  
ljuben@aubg.edu*

## Abstract

Let  $\Sigma_{2n}$  be the set of all partitions of the even integers from the interval  $(4, 2n]$ ,  $n > 2$ , into two odd prime parts. We select a partition from the set  $\Sigma_{2n}$  uniformly at random. Let  $2G_n$  be the number partitioned by this selection.  $2G_n$  is sometimes called a Goldbach number. In [6] we showed that  $G_n/n$  converges weakly to the maximum  $T$  of two random variables which are independent copies of a uniformly distributed random variable in the interval  $(0, 1)$ . In this note we show that the mean and the variance of  $G_n/n$  tend to the mean  $\mu_T = 2/3$  and variance  $\sigma_T^2 = 1/18$  of  $T$ , respectively. Our method of proof is based on generating functions and on a Tauberian theorem due to Hardy-Littlewood-Karamata.

## 1. Introduction and Statement of the Main Result

A Goldbach number is an even positive integer that can be represented as the sum of two primes without regard to order. The representation itself is called a Goldbach partition of the underlying even positive integer. Let  $\mathcal{P} = \{p_1, p_2, \dots\}$  be the sequence of all odd primes arranged in increasing order. For any integer  $k > 2$ , by  $Q_2(2k)$  we denote the number of Goldbach partitions of the number  $2k$ . In 1742 Goldbach conjectured that  $Q_2(2k) \geq 1$ . This problem remains still unsolved (for more details, see e.g. [4], Section 2.8 and p. 594, [8], Section 4.6 and [9], Chapter VI). Let  $\Sigma_{2n}$  be the set of all Goldbach partitions of the even integers from the interval  $(4, 2n]$ ,  $n > 2$ . The cardinality of this set is obviously

$$|\Sigma_{2n}| = \sum_{2 < k \leq n} Q_2(2k). \quad (1)$$

We recently established in [6] that

$$|\Sigma_{2n}| \sim \frac{2n^2}{\log^2 n}, \quad n \rightarrow \infty. \quad (2)$$

Consider now a random experiment. Suppose that we select a partition uniformly at random from the set  $\Sigma_{2n}$ , i.e. we assign the probability  $1/|\Sigma_{2n}|$  to each

Goldbach partition of an even integer from the interval  $(4, 2n]$ . We denote by  $\mathbb{P}$  the uniform probability measure on  $\Sigma_{2n}$ . Let  $2G_n \in (4, 2n]$  be the Goldbach number that is partitioned by this random selection. Using (2), it is not difficult to show that  $G_n/n$  converges weakly, as  $n \rightarrow \infty$ , to a random variable whose cumulative distribution function is

$$F(u) = \begin{cases} 0 & \text{if } u \leq 0, \\ u^2 & \text{if } 0 < u < 1, \\ 1 & \text{if } u \geq 1 \end{cases} \quad (3)$$

(see [6], Theorem 2). It can be easily seen that  $F(u)$  is the distribution function of the random variable  $T = \max\{U_1, U_2\}$ , where  $U_1$  and  $U_2$  are two independent copies of a uniformly distributed random variable in the interval  $(0, 1)$ .

The goal of this present note is to determine asymptotically the first two moments of a random Goldbach number  $2G_n$ . We state our main result in terms of the expectation  $\mathbb{E}$  and variance  $\text{Var}$  of the random variable  $G_n$ , both taken with respect to the probability measure  $\mathbb{P}$ . By  $\mu_T$  and  $\sigma_T^2$  we denote the expected value and the variance of the random variable  $T$ , respectively. (Its cumulative distribution function is given by (3).)

**Theorem 1.** *We have*

$$(i) \quad \lim_{n \rightarrow \infty} \frac{1}{n} \mathbb{E}(G_n) = \mu_T = \frac{2}{3};$$

$$(ii) \quad \lim_{n \rightarrow \infty} \frac{1}{n^2} \text{Var}(G_n) = \sigma_T^2 = \frac{1}{18}.$$

Our method of proof is similar to that in [6]. It is essentially based on a generating function identity and on a classical Tauberian theorem due to Hardy, Littlewood and Karamata (see e.g. [7], Theorem 8.7).

*Remark 1.* Our method of proof and further but similar computations yield also asymptotic expressions for moments of higher order of the variable  $G_n/n$ .

*Remark 2.* The proof of Hardy-Littlewood-Karamata Tauberian theorem may be found e.g. in [3], Chapter 7.

Our paper is organized as follows. Section 2 contains some preliminaries. The proof of Theorem 1 is given in Section 3.

## 2. Preliminary Results

By a prime partition of the positive integer  $n$ , we mean a way of writing it as a sum of primes from the set  $\mathcal{P}$  without regard to order; the summands are called parts. Clearly, Goldbach partitions are prime partitions in which the number of parts is 2.

Consider now the number  $Q_m(n)$  of prime partitions of  $n$  into  $m$  parts ( $1 \leq m \leq n$ ). The bivariate generating function of the numbers  $Q_m(n)$  is of Euler's type, namely,

$$1 + \sum_{n=1}^{\infty} z^n \sum_{m=1}^n Q_m(n) x^m = \prod_{p_k \in \mathcal{P}} (1 - xz^{p_k})^{-1}. \quad (4)$$

(the proof may be found in [1], Section 2.1). It is clear that, for  $n > 4$ ,  $Q_2(n)$  counts the number of Goldbach partitions of  $n$  and that  $Q_2(n) = 0$  if  $n$  is odd.

For any real variable  $z$  with  $|z| < 1$ , we also set

$$f(z) = \sum_{p_k \in \mathcal{P}} z^{p_k}. \quad (5)$$

Differentiating the left- and right-hand sides of (4) twice with respect to  $z$  and setting then  $z = 0$  and  $m = 2$ , we obtain the following identity.

**Lemma 1.** (See [6], Lemma 3.) For  $|z| < 1$ , we have

$$2 \sum_{k>2} Q_2(2k) z^{2k} = f^2(z) + f(z^2), \quad (6)$$

where  $f(z)$  is the function defined by (5).

Further, we will use a Tauberian theorem by Hardy-Littlewood-Karamata [3], Chapter 7. We use it in the form given by Odlyzko [7], Section 8.2.

**Hardy-Littlewood-Karamata Theorem.** (See [7; Theorem 8.7, p. 1225].) Suppose that  $a_k \geq 0$  for all  $k$ , and that

$$g(x) = \sum_{k=0}^{\infty} a_k x^k$$

converges for  $0 \leq x < r$ . If there is a  $\rho > 0$  and a function  $L(t)$  that varies slowly at infinity such that

$$g(x) \sim (r-x)^{-\rho} L\left(\frac{1}{r-x}\right), \quad x \rightarrow r^-, \quad (7)$$

then

$$\sum_{k=0}^n a_k r^k \sim \left(\frac{n}{r}\right)^{\rho} \frac{L(n)}{\Gamma(\rho+1)}, \quad n \rightarrow \infty. \quad (8)$$

*Remark.* A function  $L(t)$  varies slowly at infinity if, for every  $u > 0$ ,  $L(ut) \sim L(t)$  as  $t \rightarrow \infty$ .

### 3. Proof of the Main Result

*Proof of Theorem 1(i).* First, we notice that the definition of the random variable  $G_n$  implies that

$$\frac{1}{n}\mathbb{E}(G_n) = \sum_{2 < k \leq n} \left(\frac{k}{n}\right) \mathbb{P}(G_n = k) = \sum_{2 < k \leq n} \left(\frac{2k}{2n}\right) \frac{Q_2(2k)}{\sum_{2 < j \leq n} Q_2(2j)}. \quad (9)$$

The asymptotic behavior of the denominator in the right-hand side of (9) is completely determined by (1) and (2). Differentiating both sides of (6), we will show next that the series

$$\sum_{k>2} (2k)Q_2(2k)z^{2k-1} = f(z)f'(z) + zf'(z^2) \quad (10)$$

and

$$\sum_{k>2} (2k)(2k-1)Q_2(2k)z^{2k-2} = f'^2(z) + f(z)f''(z) + f(z^2) + 2z^2f'(z^2) \quad (11)$$

satisfy the conditions of Hardy-Littlewood-Karamata theorem. We need the following lemma.

**Lemma 2.** *Let  $f(z)$  be the series defined by (5). Then, as  $z \rightarrow 1^-$ ,*

$$f(z) \sim \frac{1}{(1-z)\log\frac{1}{1-z}}, \quad (12)$$

$$f'(z) \sim \frac{2}{(1-z)^2\log\frac{1}{1-z}}, \quad (13)$$

$$f''(z) \sim \frac{2}{(1-z)^3\log\frac{1}{1-z}}. \quad (14)$$

*Proof.* The proof of (12) is given in [6], Lemma 4. Here we present a complete proof of (13). The proof of (14) is similar.

As usual, by  $\pi(y)$  we denote the number of primes which do not exceed the positive number  $y$ . In (5) we set  $z = e^{-t}$ ,  $t > 0$ , and apply an argument similar to that given by Stong [10] (see also [2]). We have

$$f'(z) \Big|_{z=e^{-t}} = \sum_{p_k \in \mathcal{P}} p_k z^{p_k-1} \Big|_{z=e^{-t}} = \int_0^\infty ye^{-yt} d\pi(y) = I(t) - f(e^{-t}), \quad (15)$$

where

$$I(t) = t \int_0^\infty ye^{yt} \pi(y) dy. \quad (16)$$

In [6] we established that

$$f(e^{-t}) \sim \frac{1}{t \log \frac{1}{t}}, \quad t \rightarrow 0^+. \quad (17)$$

To find the asymptotic of  $I(t)$ , we set in (16)  $y = s/t$  and obtain

$$I(t) = \frac{1}{t} \int_0^\infty s e^{-s} \pi(s/t) ds = \frac{1}{t} (I_1(t) + I_2(t)), \quad (18)$$

where

$$I_1(t) = \int_0^{t^{1/2}} s e^{-s} \pi(s/t) ds,$$

$$I_2(t) = \int_{t^{1/2}}^\infty s e^{-s} \pi(s/t) ds.$$

For  $I_1(t)$  we use the bound  $\pi(s/t) \leq s/t$ . Hence, for enough small  $t > 0$ , we have

$$\begin{aligned} 0 \leq I_1(t) &\leq \frac{1}{t} \int_0^{t^{1/2}} s^2 e^{-s} ds = \frac{1}{t} \left( -s^2 e^{-s} \Big|_0^{t^{1/2}} + 2 \int_0^{t^{1/2}} s e^{-s} ds \right) \\ &= -e^{-t^{1/2}} + O(t^{-1/2}) = O(t^{-1/2}). \end{aligned} \quad (19)$$

The estimate of  $I_2(t)$  follows from the Prime Number Theorem with an error term given by Ingham [5], Theorem 23, p.65. Thus, for  $y > 1$ , we have

$$\pi(y) = \frac{y}{\log y} + O\left(\frac{y}{\log^2 y}\right).$$

Hence, for  $s \geq t^{1/2}$ , we have  $\log s \geq -\frac{1}{2} \log \frac{1}{t}$  and therefore,

$$\begin{aligned} \pi(s/t) &= \left(\frac{s}{t}\right) \frac{1}{\log \frac{1}{t} + \log s} + O\left(\frac{s}{t (\log \frac{1}{t} + \log s)^2}\right) \\ &= \frac{s}{t \log \frac{1}{t}} \left(1 + O\left(\frac{|\log s|}{\log \frac{1}{t}}\right)\right) + O\left(\frac{s}{t \log \frac{1}{t}}\right) \\ &= \frac{s}{t \log \frac{1}{t}} + O\left(\frac{s(1 + |\log s|)}{t \log^2 \frac{1}{t}}\right). \end{aligned}$$

Consequently,

$$\begin{aligned}
I_2(t) &= \frac{1}{t \log \frac{1}{t}} \int_{t^{1/2}}^{\infty} s^2 e^{-s} ds + O\left(\frac{1}{t \log^2 \frac{1}{t}} \int_{t^{1/2}}^{\infty} s^2 (1 + |\log s|) e^{-s} ds\right) \\
&= \frac{1}{t \log \frac{1}{t}} \left( \int_0^{\infty} s^2 e^{-s} ds - \int_0^{t^{1/2}} s^2 e^{-s} ds \right) + O\left(\frac{1}{t \log^2 \frac{1}{t}}\right) \\
&= \frac{1}{t \log \frac{1}{t}} (2 - O(t^{1/2})) + O\left(\frac{1}{t \log^2 \frac{1}{t}}\right) \\
&= \frac{2}{t \log \frac{1}{t}} + O\left(\frac{1}{t^{1/2} \log \frac{1}{t}}\right) + O\left(\frac{1}{t \log^2 \frac{1}{t}}\right) \sim \frac{2}{t \log \frac{1}{t}}, \quad t \rightarrow 0^+. \quad (20)
\end{aligned}$$

Combining (18)-(20), we get

$$I(t) \sim \frac{2}{t^2 \log \frac{1}{t}}, \quad t \rightarrow 0^+.$$

Replacing this asymptotic equivalence and (17) into (15), we conclude that

$$f'(z) |_{z=e^{-t}} = I(t) + O\left(\frac{1}{t \log \frac{1}{t}}\right) \sim \frac{2}{t^2 \log \frac{1}{t}}, \quad t \rightarrow 0^+.$$

Returning to the variable  $z$  by  $t = \log \frac{1}{z}$  ( $z < 1$ ), we obtain

$$f'(z) \sim -\frac{2}{(\log^2 \frac{1}{z}) \log \log \frac{1}{z}}, \quad z \rightarrow 1^-.$$

Now, (13) follows from the obvious observation that

$$\log \frac{1}{z} = -\log z = -\log(1 - (1 - z)) \sim 1 - z, \quad z \rightarrow 1^-.$$

The asymptotic equivalence (14) can be obtained in the same way using the representation

$$f''(z) |_{z=e^{-t}} = \int_0^{\infty} y(y-1) e^{-yt} d\pi(y).$$

One can show that this last integral is  $\sim \frac{2}{t^3 \log \frac{1}{t}}$  as  $t \rightarrow 0^+$  and then substitute again  $t$  by  $-\log z$ . We omit further details. ■

To complete the proof of Theorem 1(i) we recall (10) and observe that (13) implies that

$$f'(z^2) \sim \frac{2}{(1-z^2) \log \frac{1}{1-z^2}} = O\left(\frac{1}{(1-z)^2 \log \frac{1}{1-z}}\right), \quad z \rightarrow 1^-.$$

So, the main contribution to the asymptotic of the right-hand side of (10) is given by the product  $f(z)f'(z)$ . Hence, from (12) and (13) we find that

$$\begin{aligned} \sum_{k>2} (2k)Q_2(2k)z^{2k-1} &= \frac{2}{(1-z)^3 \log^2 \frac{1}{1-z}} + O\left(\frac{1}{(1-z)^2 \log \frac{1}{1-z}}\right) \\ &\sim \frac{2}{(1-z)^3 \log^2 \frac{1}{1-z}}, \quad z \rightarrow 1^-, \end{aligned} \quad (21)$$

which implies that the series  $\sum_{k>2} (2k)Q_2(2k)z^{2k-1}$  satisfies condition (7) of Hardy-Littlewood-Karamata Tauberian theorem with  $r = 1, \rho = 3$  and  $L(t) = \frac{2}{\log^2 t}$ . By (8) we obtain

$$\sum_{2 < k \leq n} (2k)Q_2(2k) \sim \left(\frac{8}{3}\right) \frac{n^3}{\log^2 n}, \quad n \rightarrow \infty. \quad (22)$$

Furthermore, (1) and (2) imply that

$$(2n) \sum_{2 < k \leq n} Q_2(2k) \sim \frac{4n^3}{\log^2 n}, \quad n \rightarrow \infty. \quad (23)$$

Dividing (22) by (23) we see that the limit of the right-hand side of (9) is  $2/3$  as  $n \rightarrow \infty$ . ■

*Remark.* Theorem 1(i) also presents a solution of the following curious problem. Consider a sampling procedure that consists of two steps. In the first step we select, as previously, a Goldbach partition from the set  $\Sigma_{2n}$ , and, in the second step, we select an even number  $2R_n$  from the interval  $(4, 2n]$ . It is then easy to see that the middle part of (9) represents the probability  $Pr(R_n \leq G_n)$  by the total probability formula. Therefore, Theorem 1(i) implies that

$$\lim_{n \rightarrow \infty} Pr(R_n \leq G_n) = \frac{2}{3}.$$

*Proof of Theorem 1(ii).* In the same way, we establish that the leading term in the asymptotic of the right-hand side of (11) is given by the first two terms  $f'^2(z) + f(z)f''(z)$ . By (12)-(14), we have

$$f'^2(z) + f(z)f''(z) \sim \frac{6}{(1-z)^4 \log^2 \frac{1}{1-z}}, \quad z \rightarrow 1^-. \quad (24)$$

Furthermore, by (21),

$$\begin{aligned} &\sum_{k>2} (2k)(2k-1)Q_2(2k)z^{2k-2} \\ &= \sum_{k>2} (2k)^2 Q_2(2k)z^{2k-2} - \sum_{k>2} (2k)Q_2(2k)z^{2k-2} \\ &= \sum_{k>2} (2k)^2 Q_2(2k)z^{2k-2} + O\left(\frac{1}{(1-z)^3 \log^2 \frac{1}{1-z}}\right). \end{aligned} \quad (25)$$

Combining (11), (24) and (21), we obtain

$$\sum_{k>2} (2k)^2 Q_2(2k) z^{2k-2} \sim \frac{6}{(1-z)^4 \log^2 \frac{1}{1-z}}, \quad z \rightarrow 1^-.$$

Applying again Hardy-Littlewood-Karamata Tauberian theorem with  $r = 1, \rho = 4, L(t) = \frac{6}{\log^2 t}$ , we observe that

$$\sum_{2 < k \leq n} (2k)^2 Q_2(2k) \sim \frac{4n^4}{\log^2 n}, \quad n \rightarrow \infty. \quad (26)$$

Similarly to (9), for the second moment of  $G_n$  we have

$$\frac{1}{n^2} \mathbb{E}(G_n^2) = \sum_{2 < k \leq n} \left(\frac{k}{n}\right)^2 \mathbb{P}(G_n = k) = \sum_{2 < k \leq n} \left(\frac{2k}{2n}\right)^2 \frac{Q_2(2k)}{\sum_{2 < k \leq n} Q_2(2k)}. \quad (27)$$

Now, from (26), (27), (1) and (2) it follows that

$$\lim_{n \rightarrow \infty} \frac{1}{n^2} \mathbb{E}(G_n^2) = \frac{1}{2},$$

which, in combination with the result of Theorem 1(i), completes the proof of part (ii). ■

## References

- [1] G.E. Andrews. *The Theory of Partitions*. Encyclopedia Math. Appl., no. 2, Addison-Wesley, Reading, MA, 1976.
- [2] P. Erdős and P. Turán. *On some problems of a statistical group theory, IV*, Acta Math. Acad. Sci. Hungar. **19** (1968), 413-435.
- [3] G.H. Hardy. *Divergent Series*. Oxford Univ. Press, Oxford, 1949.
- [4] G.H. Hardy and E.M. Wright. *An Introduction to the Theory of Numbers*. Oxford Univ. Press, Oxford, 2008.
- [5] A.E. Ingham. *The Distribution of the Prime Numbers*. Cambridge Univ. Press, Cambridge. 1990.
- [6] L. Mutafchiev. *A note on Goldbach partitions of large even integers*. Electron. J. Combin. **22** (2015), # P.1.40.
- [7] A.M. Odlyzko. *Asymptotic Enumeration Methods*. In Handbook of Combinatorics (R. Graham, M. Grötschel and L. Lovász, Eds.), Vol. II, Elsevier, 1995, pp.1063-1229.



- [8] M. Overholt. *A Course in Analytic Number Theory*. Graduate Studies in Mathematics, Vol. 160, American Mathematical Society, Providence, RI, 2014.
- [9] K. Prachar. *Primzahlverteilung*. Springer, Berlin, 1957.
- [10] R. Stong. *The average order of a permutation*. Electron. J. Combin. **5** (1998), #R41.