

Herschel^{*}-HIFI observations of H₂O, NH₃ and N₂H⁺ toward high-mass starless and proto-stellar clumps identified by the Hi-GAL survey

L. Olmi^{1,2}, C.M. Persson³, and C. Codella¹

¹ INAF, Osservatorio Astrofisico di Arcetri, Largo E. Fermi 5, I-50125 Firenze, Italy, e-mail: olmi.luca@gmail.com

² University of Puerto Rico, Rio Piedras Campus, Physics Dept., Box 23343, UPR station, San Juan, Puerto Rico, USA

³ Chalmers University of Technology, Department of Earth and Space Sciences, Onsala Space Observatory, SE-439 92 Onsala, Sweden.

Received Month Date, 2015; accepted Month Date, 2015

ABSTRACT

Context. Our present understanding of high-mass star formation still remains very schematic. In particular, it is not yet clear how much of the difference between low-mass and high-mass star formation occurs during the earliest star formation phases.

Aims. The chemical characteristics of massive cold clumps, and the comparison with those of their low-mass counterparts, could provide crucial clues about the exact role that chemistry plays in differentiating the early phases of low-mass and high-mass star formation. Water, in particular, is a unique probe of physical and chemical conditions in star-forming regions.

Methods. Using the HIFI instrument of *Herschel* we have observed the ortho-NH₃ (1₀ – 0₀) (572 GHz), ortho-H₂O (1₁₀ – 1₀₁) (557 GHz) and N₂H⁺ (6 – 5) (559 GHz) lines toward a sample of high-mass starless and proto-stellar clumps selected from the “*Herschel* Infrared Galactic Plane Survey” (Hi-GAL). We compare our results to previous studies of low-mass and high-mass proto-stellar objects.

Results. At least one of the three molecular lines was detected in 4 (out of 35) and 7 (out of 17) objects in the $\ell = 59^\circ$ and $\ell = 30^\circ$ galactic regions, respectively. All detected sources are *proto*-stellar. The water spectra are complex and consist of several kinematic components, identified through a Gaussian decomposition, and in a few sources inverse and regular P-Cygni profiles have been detected. All water line profiles of the $\ell = 59^\circ$ region are dominated by a broad Gaussian emission feature, indicating that the bulk of the water emission arises in outflows. No such broad emission is detected toward the $\ell = 30^\circ$ objects. The ammonia line in some cases also shows line wings and an inverse P-Cygni profile, thus confirming that NH₃ rotational transitions can be used to probe the dynamics of high-mass star forming regions. Both bolometric and water line luminosity increase with the continuum temperature.

Conclusions. The higher water abundance toward the $\ell = 59^\circ$ sources, characterized by the presence of outflows and shocks, supports a scenario where the abundance of this molecule is linked to the shocked gas. Various indicators suggest that the detected sources toward the $\ell = 30^\circ$ region are in a somewhat later evolutionary phase compared to the $\ell = 59^\circ$ field, although a firm conclusion is limited by the small number of observed sources. We find many similarities with studies carried out toward low-mass proto-stellar objects, but there are indications that the level of infall and turbulence in the high-mass protostars studied here could be significantly higher.

Key words. ISM: molecules – ISM: abundances – Line: formation – Stars: formation

1. Introduction

High-mass stars (OB spectral type, $M \gtrsim 8 M_\odot$), though much rarer than low-mass and intermediate-mass stars, are fundamental in the evolution of galaxies. But, while a reasonably robust evolutionary sequence has been established for the formation and early evolution of individual, our present understanding of high-mass star formation still remains very schematic (Zinnecker & Yorke 2007, Tan et al. 2014). In particular, some of the main issues regarding high-mass star formation that still need to be studied are: (i) what is the physical and dynamical evolutionary path of massive starless¹ clumps² toward the proto-

* *Herschel* is an ESA space observatory with science instruments provided by European-led Principal Investigator consortia and with important participation from NASA.

¹ As identified through the absence of Mid-IR emission and/or the strength of the PACS 70 μ m emission.

² In the literature, the term “core” or “fragment” usually refers to small-scale objects ($\lesssim 0.1$ pc or even $\ll 0.1$ pc), possibly corresponding

stellar and stellar phases? And, (ii) how much of the difference between low-mass and high-mass star formation occurs during the earliest star formation phases?

The high-mass analogues of low-mass Class 0 protostars and starless clumps (e.g., Motte et al. 2007, Bontemps et al. 2010, Olmi et al. 2010, Olmi et al. 2013) constitute a fundamental link between the global processes that regulate star formation on large scales and the properties of newly formed (proto)stars and clusters. Therefore, to better define the evolutionary path from molecular clumps to proto-stars it is crucial to determine the physical and dynamical (and, possibly, chemical) properties of these early phases of high-mass star formation.

The formation of both low-mass and high-mass stars from the initial dense clumps is accompanied by large changes in the

to a later stage of fragmentation, while the term “clump” is generally used for a somewhat larger ($\gtrsim 0.1 - 1$ pc), unresolved object, possibly composed of several cores. The term clump is thus more appropriate to refer to the compact objects described in this work.

Table 1: Observed transitions.

Species	Frequency ^a [GHz]	Transition ^b	E_l [K]	E_u [K]	$\log(A_{ul})$ [s ⁻¹]
ortho-NH ₃	572.4982	$J_K = 1_0 - 0_0$	0	27	-2.80
ortho-H ₂ O	556.9359	$J_{K_a, K_c} = 1_{1,0} - 1_{0,1}$	34	61	-2.46
N ₂ H ⁺	558.9665	$J = 6 - 5$	67	94	-1.96

Notes. ^(a) Frequencies from <http://www.cv.nrao.edu/php/splat/> ^(b) Critical densities are $n_{cr} \gtrsim 10^7 - 10^8 \text{ cm}^{-3}$ for all three lines.

temperature and density structure in their surrounding envelopes. These changes in turn open up different chemical routes. Current observational evidence indicates that significant amounts of volatile molecules, for instance CO, H₂O and NH₃, reside on the surfaces of dust grains in dense clumps. These surface molecules can be formed either in the gas phase via ion-molecule reactions followed by accretion onto the dust particles, or via direct production on the grain surfaces. Different physical conditions, for example during the early formation phases of low-mass and high-mass stars, could thus determine which of these different chemical routes to molecule formation are active and/or dominant.

Table 2: Selected observational and physical properties of the source sample: 1σ noise level, SSB continuum intensity (T_c), beam-filling factor (η_{bf}), evolutionary flag, molecular hydrogen volume density ($n(\text{H}_2)$), and column density ($N(\text{H}_2)$). Luminosity (L), mass (M), dust temperature T_d , source diameter (Full Width at Half Maximum, or FWHM) and distance (d) are all derived from the Hi-GAL observations. Sources with their name in boldface have at least one detection and thus they could be assigned a V_{lsr} .

Source	Gal. Long. [deg]	Gal. Lat [deg]	L [L_\odot]	M [M_\odot]	T_d [K]	Source FWHM ["]	d [kpc]	T_c^a [mK]	$1\sigma^b$ [mK]	V_{lsr}^c [km s^{-1}]	η_{bf}^d	$n(\text{H}_2)^e$ [$\times 10^4 \text{ cm}^{-3}$]	$N(\text{H}_2)^e$ [$\times 10^{22} \text{ cm}^{-2}$]	Evol.Flag ^f
159-251	57.984803	0.0927620	27.2	165.0	10.1	17.3	7.0	0	9	...	0.2	2.7	4.82	1
159-139	58.048693	0.1889100	117.5	2118.0	9.2	27.8	9.1	0	8	...	0.4	3.7	14.0	1
159-376	58.171557	0.7804730	7.7	118.0	8.9	25.3	2.6	0	8	...	0.3	12.0	12.0	1
159-199	58.256292	0.1431680	67.5	323.0	11.5	31.5	7.2	0	8	...	0.4	0.81	2.7	1
159-203	58.464407	0.2996740	11.5	81.0	10.8	20.0	3.5	0	8	...	0.2	6.5	6.9	1
159-129	58.464903	0.4929190	12.6	17.9	...	0	8	...	0.2	1
159-185	58.512225	0.5060260	5.2	132.0	8.3	25.3	2.8	0	8	...	0.3	11.0	11.0	1
159-102	58.517332	0.4718170	8.7	100.0	9.2	25.3	2.8	0	9	...	0.3	1
159-78	58.558761	0.3525380	33.7	3296.0	7.0	25.3	6.2	0	9	...	0.3	25.0	58.0	1
159-314	58.569252	0.0207140	27.6	100.0	11.1	24.8	5.7	0	10	...	0.3	0.53	1.1	1
159-391	58.606871	0.7818820	25.0	109.0	10.7	25.3	4.7	0	9	...	0.4	1
159-80	58.646263	0.5462850	77.4	1027.0	9.1	25.3	6.1	24	10	...	0.3	8.0	19.0	1
159-345	58.693246	0.7685080	25.7	149.0	10.3	25.3	4.9	0	8	...	0.3	4.5	10.0	1
159-136	58.754808	0.6835960	23.1	446.0	8.6	25.3	5.5	0	8	...	0.3	4.8	9.9	1
159-231	58.755031	0.7204550	88.4	137.0	13.8	33.8	5.6	0	9	...	0.5	0.58	1.7	1
159-293	58.810992	0.6553700	14.5	3.0	19.6	17.8	5.2	0	9	...	0.2	0.11	0.15	0
159-667	58.865456	-0.212801	653.6	799.0	12.5	17.0	13.7	0	8	...	0.2	1.8	6.3	2
159-230	59.051623	-0.229814	23.0	79.0	12.1	27.0	2.6	0	8	...	0.4	6.8	7.1	1
159-57	59.151275	-0.268629	185.7	281.0	13.9	24.2	6.2	22	9	...	0.3	2.4	5.4	1
159-191	59.208976	0.1322060	62.8	239.0	11.0	25.3	6.7	0	8	...	0.4	1.1	1.0	1
159-263	59.212680	0.1586920	14.0	170.0	9.2	25.3	4.0	0	9	...	0.5	0.66	2.1	1
159-150	59.300967	0.2429840	47.3	210.0	10.7	25.3	5.9	0	8	...	0.2	13.0	20.0	1
159-104	59.304277	0.2083220	17.0	2.0	23.3	25.3	2.7	0	9	...	0.3	0.18	0.18	1
159-371	59.309467	-0.397915	15.8	227.0	9.6	19.3	6.2	0	8	...	0.2	3.8	6.8	1
159-31	59.541533	-0.101984	20.7	5.0	19.4	25.3	2.3	0	9	...	0.2	60.0	30.0	1
159-441	59.633104	-0.191683	2259.6	294.0	21.6	15.3	2.2	110	8	27.8	0.1	230.0	110.0	2
159-445	59.635570	-0.186242	787.9	82.0	20.9	23.0	2.3	75	8	27.4	0.3	16.0	13.0	2
159-282	59.707751	0.5772220	30.1	112.0	10.9	25.3	4.3	0	8	...	0.1	1
159-409	59.737176	0.5263750	60.7	123.0	13.2	27.8	6.2	0	8	...	0.4	0.7	1.8	1
159-444	59.789451	0.6307310	21.8	9.0	...	37	8	34.3	0.1	2
159-442	59.832526	0.6718730	25.3	13.9	...	28	8	34.2	0.1	2
159-359	59.862726	-0.210802	55.3	429.0	9.8	25.3	6.4	0	11	...	0.5	5.1	18.0	1
159-101	59.880982	-0.102392	85.9	2814.0	8.4	25.9	6.6	0	8	...	0.3	16.0	41.0	1
159-69	59.891888	-0.112074	27.8	92.0	12.2	36.2	1.9	0	8	...	0.5	7.5	8.0	1
159-198	59.946964	0.0610780	5.9	5.0	15.3	13.5	1.9	32	8	...	0.1	8.3	3.2	1

Notes. ^(a) SSB continuum intensity at 557 GHz. ^(b) Rms noise at a resolution of 1 km s^{-1} with subtracted baseline. ^(c) Velocity is specified only for sources with at least a detection. ^(d) Beam-filling factor, see Section 3.4. ^(e) See Section 3.4 for a description of these parameters. ^(f) 0: no detection at $70 \mu\text{m}$, Larson's criterion (Larson 1981) not fulfilled (starless, unbound). 1: no detection at $70 \mu\text{m}$, Larson's criterium fulfilled (pre-stellar). 2: detection at $70 \mu\text{m}$ (proto-stellar).

Table 2: continued

Source	Gal. Long. [deg]	Gal. Lat [deg]	L [L_{\odot}]	M [M_{\odot}]	T_d [K]	Source FWHM [$''$]	d [kpc]	T_c^a [mK]	$1\sigma^b$ [mK]	V_{lsr}^c [km s^{-1}]	η_{br}^d	$n(\text{H}_2)^e$ [$\times 10^4 \text{ cm}^{-3}$]	$N(\text{H}_2)^e$ [$\times 10^{22} \text{ cm}^{-2}$]	Evol.Flag ^f
130-376	29.002507	0.0689530	9328.5	1072.0	19.8	21.8	8.9	23	8	70.5	0.3	4.1	12.0	2
130-512	29.235020	-0.046735	1312.0	1860.0	13.8	12.6	9.9	63	8	61.4	0.1	27.0	51.0	2
130-313	29.862299	-0.043707	53790.2	1866.0	22.9	12.2	8.5	68	8	101.5	0.1	47.0	73.0	2
130-110	29.868288	-0.007673	313.8	1311.0	10.6	14.7	8.5	27	8	...	0.3	10.0	20.0	1
130-327	30.009612	-0.273200	12371.5	964.0	23.5	9.2	6.7	98	8	102.6	0.1	120.0	110	2
130-281	30.121077	-0.548844	435.5	628.0	12.8	25.3	8.1	44	8	...	0.3	2.1	6.4	1
130-375	30.197099	0.3101250	15852.5	2921.0	17.7	12.7	14.2	0	9	...	0.1	14.0	39.0	2
130-432	30.447046	-0.024628	9848.3	271.0	26.7	28.0	11.6	44	8	...	0.2	6.5	22.0	2
130-103	30.575773	0.1188560	186.4	878.0	10.6	25.3	5.7	44	8	55.6	0.3	8.4	18.0	1
130-403	30.627182	-0.063627	1739.5	711.0	15.8	28.0	5.6	38	8	...	0.4	5.3	12.0	2
130-73	30.658809	0.0450200	143.7	962.0	11.1	17.7	5.6	34	8	...	0.2	28.0	42.0	1
130-111	30.661139	0.1419250	392.9	249.0	14.8	25.3	5.6	35	10	...	0.4	7.9	20.0	1
130-132	30.675761	-0.166004	1223.9	1559.0	14.3	27.4	9.0	0	8	...	0.4	3.0	11.0	1
130-42	30.693536	-0.148593	1082.2	723.0	16.7	17.4	5.7	50	9	90.4	0.2	22.0	32.0	2
130-304	30.703241	-0.068204	49112.4	3612.0	22.5	20.4	6.0	350	12	90.7	0.2	57.0	100.0	2
130-43^g	30.822869	0.1343020	2405.6	16437.0	10.2	32.6	8.8	50	8	108.4	0.5	20.0	86.0	2
130-230	30.846821	0.3426510	186.5	425.0	12.0	25.3	6.2	0	8	...	0.4	5.7	21.0	1

Notes. ^(g) Tentative detection (see Section 3.2).

The chemical characteristics of massive cold clumps, and the comparison with those of their low-mass counterparts, are only starting to emerge (Russeil et al. 2010, Marseille et al. 2010). While governed by the same processes as lower mass objects, the much larger luminosity of massive stars during all their evolutionary phases will greatly affect the outcome. Therefore, the comparison of chemical models (e.g., with different weights of grain vs. gas-chemistry) with aimed spectral line observations will provide crucial clues about the exact role that chemistry plays in differentiating the early phases of high-mass star formation from those of low-mass star formation.

The $(1_0 - 0_0)$ ground-state rotational transition of o -NH₃ (e.g., Ho & Townes 1983) has an upper state energy of 27 K (see Table 1), comparable to the rotation-inversion transitions at cm wavelengths. The critical density is, however, four orders of magnitude higher and is thus a better probe of the physical conditions of the dense high-mass starless clumps detected by the Hi-GAL survey. Moreover, its *ortho* symmetry form makes it a complementary tool to the commonly used rotation-inversion transitions of para-NH₃ observable from the ground. Both NH₃ and N₂H⁺ are well known tracers of low-mass pre- and proto-stellar clumps.

Water is another important component in the chemistry of regions of star formation. In cold gas, water is mainly produced on dust grains, and released to the gas phase when the temperature increases ($\gtrsim 100$ K). Water is thus not expected to be found with high abundances in cold, starless clumps (e.g., Bergin & Tafalla 2007, Caselli et al. 2012, Wiström et al. 2014). In fact, recent results from the Herschel Key Program “Water in star-forming regions with *Herschel*” (WISH; van Dishoeck et al. 2011) have shown that during the quiescent phases of star formation, the water abundance is very low. The embedded stage of star formation, however, shows very broad and complex line profiles consisting of many dynamical components (Kristensen & van Dishoeck 2011, Kristensen et al. 2012). The WISH team also found, comparing typical low-, intermediate-, and high-mass young stellar objects, that their spectra were remarkably similar, characterized by broad, complex line profiles and consisting of multiple components. Our observations mainly consists of high-mass clumps in various evolutionary stages, and thus we will also try to identify possible evolutionary differences.

The outline of the paper is as follows. In Section 2 we discuss how our sources were selected and observed with *Herschel*. In Section 3 we describe how the main kinematical and physical parameters of the detected sources were derived. The gas kinematics is further discussed in Section 4 and we draw our conclusions in Section 5.

2. Observations and data reduction

2.1. Source selection

We have selected our targets from the preliminary source catalog obtained toward the two Galactic fields, at $\ell = 30^\circ$ and $\ell = 59^\circ$, observed for the “*Herschel* Infrared Galactic Plane Survey” (Hi-GAL, Molinari et al. 2010) during the Science Demonstration Phase of *Herschel* (Elia et al. 2010). The Hi-GAL (and Hi-GAL360) Project used SPIRE and PACS in parallel mode to carry out an unbiased imaging survey of the Galactic Plane, uniformly sampling a 2° -wide strip, with $-60^\circ < \ell < 60^\circ$, in the 70, 170, 250, 350 and 500 μm photometric bands. The source-identification, flux-extraction and spectral energy density-building in these two Hi-GAL maps were carried out by

adapting to the *Herschel* SPIRE/PACS wavebands the methods and techniques described by Netterfield et al. (2009) and Olmi et al. (2009). From the preliminary catalogs of the $\ell = 30^\circ$ and $\ell = 59^\circ$ fields, we have extracted both high-mass starless clumps and high-mass proto-stellar objects.

Candidate high-mass starless clumps have been selected by imposing that $M_{\text{clump}} > 100 M_\odot$ ($M_{\text{clump}} > 200 M_\odot$) and $L_{\text{fir}} < 100 L_\odot$ ($L_{\text{fir}} < 500 L_\odot$) for the $\ell = 59^\circ$ ($\ell = 30^\circ$) field. Moreover, we have checked that no compact MIPS (“Multiband Imaging Photometer for Spitzer”, Rieke et al. 2004) 24 μm emission were present at the position of the candidate high-mass starless clump. Candidate high-mass proto-stellar objects do have instead a compact MIPS 24 μm counterpart. Finally, the sources have been visually inspected to reject ambiguous detections. This procedure resulted in a total of 81 clumps, including the four candidate high-mass starless clumps already observed with the VLA and the Effelsberg 100-m telescope in the NH₃(1, 1) and (2, 2) inversion lines by Olmi et al. (2010). Out of the initial sample, 52 sources were finally observed with *Herschel*: 17 objects in the $\ell = 30^\circ$ region and 35 in the $\ell = 59^\circ$ region. The observed sources and their basic physical parameters are listed in Table 2. Sources with a detection in at least one of the three observed lines have been highlighted in boldface.

2.2. *Herschel* observations

Using the HIFI instrument (de Graauw et al. 2010) of *Herschel* (Pilbratt et al. 2010) we have observed the o -NH₃($1_0 - 0_0$) line at 572 GHz in the upper side band of band 1b, and simultaneously in the lower side band both the o -H₂O($1_{10} - 1_{01}$) line at 557 GHz, and the N₂H⁺($6 - 5$) line at 559 GHz (see Table 1). We used the wide band spectrometer with a bandwidth of 4 GHz and an effective spectral resolution of 1.1 MHz ($\Delta v = 0.27 \text{ km s}^{-1}$). The half-power beam width and the main-beam efficiency at 572 GHz are 36'' and 0.62, respectively (Mueller et al., 2014³). The total calibration uncertainties are $\approx 9\%$ for band 1⁴ (Roelfsema et al. 2012). We have used the dual beam switching (DBS) observing mode. The reference beams in these observations were located within 3' on either side of the source. The observations were performed during the *Herschel* Open Time (OT2) on April 19 and 20, 2013, and the corresponding observation identifier (OBSID) range is [1342270449, 1342270604].

The data were processed using the standard *Herschel* Interactive Processing Environment⁵, version 11.1, up to level 2, providing fully calibrated double side band spectra in the T_A^* antenna temperature intensity scale where the lines are calibrated on a single side band (SSB) scale. Because HIFI is intrinsically a double side band instrument, the continuum has to be divided by two to be properly scaled. The data quality is excellent with very low intensity ripples in most cases, typically below a few percent of the double side band continuum. However, in some cases the baselines are bad enough that the large rms ($\gtrsim 20 - 30 \text{ mK}$) does not allow a reliable estimate of the continuum temperature to be obtained (see Section 3.3). Therefore, for these sources we set the continuum temperature to 0. The FITS files were exported to the spectral line analysis software packages *xs*⁶ and

³ http://herschel.esac.esa.int/twiki/pub/Public/HiFiCalibrationWeb/HiFiBeamReleaseNote_Sep2014.pdf

⁴ <http://herschel.esac.esa.int/Docs/HIFI/html/ch5.html>

⁵ <http://herschel.esac.esa.int/HIPE/download.shtml>

⁶ <http://www.chalmers.se/rss/oso-en/observations/data-reduction-software>




Fig. 1: *Top panel.* Gaussian decomposition of the o - $\text{H}_2\text{O}(1_{10} - 1_{01})$ spectrum towards the source 159-441. The individual Gaussian components are shown as solid color lines overlaid on the (continuum-subtracted) spectrum. The overall Gaussian fit is shown as a red dashed line, and the residual deviation at a velocity of $\approx 55 \text{ km s}^{-1}$ is visible. The intensity of the component in absorption has been multiplied by a 0.2 factor. *Bottom panel.* Overlay of the o - $\text{H}_2\text{O}(1_{10} - 1_{01})$, o - $\text{NH}_3(1_0 - 0_0)$ and $\text{N}_2\text{H}^+(6-5)$ lines towards the source 159-441.

CLASS⁷. Both polarisations and all three LO-settings were included in the averaged noise-weighted spectra. The resulting averages were convolved to a resolution of 1 km s^{-1} in sources with a low signal-to-noise ratio. All HIFI spectra, in T_A^* units,

⁷ CLASS is part of the GILDAS software package developed by IRAM.

are shown in Appendices A ($\ell = 30^\circ$ region) and B ($\ell = 59^\circ$ region).

3. Results

3.1. Distance determination

Assigning a distance to sources detected with a photometer is a crucial step in giving physical significance to all information extracted from the Hi-GAL data. While reliable distance estimates are available for a limited number of known objects (e.g., HII regions, see Russeil 2003, and masers, see e.g. Green & McClure-Griffiths 2011), this information does not exist for the majority of Hi-GAL sources. We therefore adopted the scheme presented by Russeil et al. (2011) to assign kinematic distances to large lists of sources. A ^{12}CO (or ^{13}CO) spectrum (e.g., from the BUFCRAO Galactic Ring Survey, or GRS, Jackson et al. 2006) is extracted in the line of sight of every Hi-GAL source. Then, the velocity, V_{lsr} , of the brightest spectral component is assigned to that specific source thereby allowing calculation of a kinematical distance. By using extinction maps (derived from the 2MASS point source catalog, see e.g. Schneider et al. 2011) and a catalog of sources with known distances, such as HII regions, masers, and others, the distance ambiguity is resolved and a recommendation given. In this way it is possible to produce a “distance map” having the same pixel size of the CO cube as used to extract the spectra for every target, where the value of the pixel is the assigned distance of the Hi-GAL source(s) falling in that pixel. The typical error on the kinematic distance⁸ (column 8 in Table 2) is estimated to vary between $\sim 0.6 - 0.9 \text{ kpc}$ in the range of longitude $\ell \sim 30^\circ - 60^\circ$.

3.2. Profile decomposition

The observed line profiles, and in particular those of the water spectral lines, show complex shapes, and likely represent contributions from proto-stellar envelopes, molecular outflows, foreground clouds as well as infall motions. These contributions, both in emission and in absorption, are disentangled and parameterized by fitting multiple Gaussians to the line profiles. We show an example in Fig. 1 and list all the results in Table 3. The Gaussian fits have been performed using up to four velocity components and all line parameters were allowed to vary. A similar decomposition of the $\text{H}_2\text{O}(1_{10} - 1_{01})$ (557 GHz) line has also been used by Kristensen et al. (2012) and Herpin et al. (2012).

Table 3 shows that the H_2O line profiles of the $\ell = 59^\circ$ sources are very similar. Almost all water spectra can be described as the sum of a narrow (FWHM $\sim 3 - 5 \text{ km s}^{-1}$), a medium (FWHM $\sim 6 - 10 \text{ km s}^{-1}$) and a broad (up to $\approx 30 \text{ km s}^{-1}$) velocity component. The narrow component is mostly seen in absorption, whereas the medium and broad components have positive intensities. Source 159-444 shows an inverse P-Cygni profile overlaid on a broad component. In source 159-441 and, to a lesser extent, in 159-445 (please note that these two sources are only $21.5''$ apart) if three Gaussian components are used in the fit the red-shifted wing of the spectrum still shows a residual deviation ($\approx 3\sigma$, see Fig. 1) from the overall fit at a velocity of $\approx 55 \text{ km s}^{-1}$. This residual can be eliminated with an additional Gaussian component, which may either have a high velocity ($\approx 50 \text{ km s}^{-1}$), or a velocity comparable to the other components but with a wide line width ($\approx 40 \text{ km s}^{-1}$). In the

⁸ This estimate assumes that the distance ambiguity has been correctly solved.

Table 3: Gaussian fits of the *Herschel*-HIFI observations.

Source	H ₂ O Comp. ^a	E/A ^b	H ₂ O			N ₂ H ⁺			NH ₃		
			V _{lsr} [km s ⁻¹]	T _A [*] [K]	ΔV [km s ⁻¹]	V _{lsr} [km s ⁻¹]	T _A [*] [K]	ΔV [km s ⁻¹]	V _{lsr} [km s ⁻¹]	T _A [*] [K]	ΔV [km s ⁻¹]
159-441	M	E	27.9	0.89	6.3	27.8	0.83	3.1	27.7	0.81	3.6
	B	E	27.0	0.37	26.0	28.2	0.09	10.1
	N	A	26.9	-1.1	3.7	27.7	-0.46	2.0
159-442	M	E	36.7	0.24	6.3	34.2	0.14	3.6	34.8	0.13	6.4
	B	E	36.7	0.12	23.6
	N	A	34.9	-0.30	4.8	35.1	-0.05	2.5
159-444	N	E	32.0	0.21	2.9	34.3	0.20	4.6	34.4	0.21	5.8
	B	E	31.2	0.08	20.7
	N	A	27.7	-0.08	1.0
	N	A	33.9	-0.14	4.8	35.4	-0.18	3.4
159-445	M	E	28.0	0.67	7.2	27.4	0.67	3.1	27.4	0.64	4.2
	B	E	28.4	0.24	28.2
	N	A	27.0	-0.80	3.7	27.7	-0.25	1.7
130-42	90.4	0.03	1.9	91.3	0.08	4.2
130-43 ^d	N	E	108.4	0.07	2.7	...	≲ 0.02 ^c	≲ 0.02 ^c	...
130-304	90.7	0.86	4.1	87.8	0.33	3.1
	M	A	92.9	-0.35	8.5	93.7	-0.31	5.7
130-313	M	E	101.2	0.07	6.5	101.5	0.08	2.7	101.5	0.14	5.8
130-327	102.6	0.07	3.6
	M	A	103.8	-0.11	5.7	104.6	-0.08	4.6
130-376	N	E	70.5	0.07	3.2	...	≲ 0.02 ^c	...	70.5	0.05	2.7
130-512	M	E	63.2	0.11	5.5	61.4	0.06	2.9	61.8	0.15	2.5
	N	A	59.5	-0.06	2.4
	N	A	64.0	-0.06	1.6

Notes. ^(a) Definition of the Gaussian components of the water line: N=narrow, M=medium, B=broad. ^(b) E=emission, A=absorption. ^(c) 3σ limit. ^(d) Tentative detection (see Section 3.2).

first case the additional velocity component could be interpreted as an extremely high velocity (EHV) component. Similar EHV components have previously been reported, e.g., by Kristensen et al. (2012) toward several low-mass protostars and by Leurini et al. (2014) toward the massive protostar IRAS 17233-3606, and are usually associated with shocked gas. However, we do not have further evidence (such as spectra of other water lines and/or spectra at different positions) to confirm that an EHV component indeed exists in source 159-441. In the second scenario, an additional broad velocity component must be invoked, which may be caused by the fact that molecular outflows are not Gaussian shaped. Due to this ambiguity, we choose to show only three Gaussian components for this source. Section 4.2 discusses the water line profile components in detail.

The situation is very different in the $\ell = 30^\circ$ region, with no two profiles being identical. We have no clear detection of a broad component in the water spectra, and the line profile can be fit by fewer Gaussian components. Ammonia also displays complex line shapes with mixed emission and absorption, and only weak signs of outflows. In contrast to water and ammonia, the N₂H⁺ spectra do not show any line asymmetries and we observe only emission. In addition, two sources also show a regular or inverse P-Cygni profile, either in the water line (130-512) or in the NH₃ spectrum (130-327) (see Section 4.2.5). Tables 2, 3 and 4 mark source 130-43 as a tentative detection because recent (June 2015) follow-up observations with the 20-m telescope of

the Onsala Space Observatory have detected the NH₃(1,1) line toward this source at a velocity of $\approx 95.3 \text{ km s}^{-1}$, instead of 108.4 km s^{-1} as shown in Table 3 and Appendix A. Therefore, either the water line in source 130-43 is an artefact of some type, or both lines are actually real and thus the issue remains as to whether or not they belong to the same object. With our current data we cannot answer this question and therefore in this work we will assume that the water line detection is real.

3.3. Correlations between dust- and gas-derived physical parameters

In order to better understand the nature of the detected sources in the $\ell = 30^\circ$ and $\ell = 59^\circ$ regions, as well as to determine the origin of the H₂O emission, we have performed correlations between several parameters. We begin with a comparison which includes both dust-derived physical parameters, as determined from our previous Hi-GAL observations, and HIFI-derived parameters, such as the continuum temperature, T_c . Then, the total H₂O integrated line intensity is compared to different parameters.

In Fig. 2 we plot the clump luminosity vs. mass, as derived from our previous Hi-GAL observations. Since both are distance-dependent quantities, only sources where a distance could be determined are shown. The plot shows that the clumps found in the $\ell = 59^\circ$ region are characterized by lower lumi-




Fig. 2: Clump bolometric luminosity vs. mass, as derived from the Hi-GAL observations for all sources where a distance could be determined. Sources in the $\ell = 30^\circ$ and $\ell = 59^\circ$ regions are represented by the (black) “+” sign and (red) open squares, respectively. Sources with a detection in at least one of the three observed spectral lines are shown with thicker (green) symbols. The dashed lines are loci at constant temperature. Roughly orthogonal to these are loci (dotted lines) of constant $250 \mu\text{m}$ flux density, ranging from 0.1 to 500 Jy, assuming a modified black-body spectral energy distribution with $\beta = 1.5$ and a fixed distance of ~ 6 kpc, equal to the median of the distances of the observed sources. The black dots and the lines joining them represent the evolutionary tracks of clumps with starting envelope mass equal to 120, 470 and $10^3 M_\odot$ (see Section 4.4).

nosity and mass compared to those in the $\ell = 30^\circ$ region (see the detailed statistical analysis by Olmi et al. 2013 and Olmi et al. 2014a). Sources with a detection in at least one of the three observed spectral lines are clearly found at higher luminosity ($L \geq 10^3 L_\odot$), as also confirmed by Table 5. In Fig. 2 we also plot a series of evolutionary tracks which we discuss in Section 4.4.

In Fig. 3 we plot the dust-derived temperatures (T_d , column 6 in Table 2) and luminosities of the clumps in the $\ell = 30^\circ$ and $\ell = 59^\circ$ regions vs. T_c . The continuum flux density associated with T_c is likely to have a contribution from both free-free emission and warm dust thermal emission. We do not have an alternative method to determine the contribution of free-free emission in our sources. Furthermore, because we have only measured T_c at one frequency, we cannot determine how T_c varies with frequency, which could otherwise be used to discriminate between dust and free-free emission. van der Tak et al. (2013), however, found that the free-free emission in their massive proto-stellar clumps is negligible compared to warm dust emission. If this is true also for our sources, then T_c should correlate with the source bolometric luminosity. This is indeed shown in the top panel of Fig. 3, where we also used the Bayesian IDL routine LINMIX_ERR to perform a linear regression to find the slope of the best fit line. The source luminosity is expected to increase




Fig. 3: *Top panel.* Clump bolometric luminosity vs. continuum temperature, T_c . The solid line represents the linear fit to all points (see text); the Spearman rank coefficient is 0.65. *Bottom panel.* Dust-derived temperature, T_d , vs. continuum temperature. Symbols are as in Fig. 2.

with time, as the clump evolves from the starless to the proto-stellar phase (see, e.g., Molinari et al. 2008). Therefore, the top panel of Fig. 3 is also likely to represent an evolutionary sequence. However, the bottom panel of Fig. 3 shows that there is only a marginal correlation between T_d and T_c , which are both distance-independent. The large scatter is certainly due in part to the large uncertainties in T_c , since the baseline rms was in general quite large ($\geq 20 - 30$ mK).

In Fig. 4 we plot various line integrated intensities. The integrated area of the profile has been determined by summing all components in emission. For a Gaussian line profile, the integrated emission is $T_A^{\text{pk}} \Delta V \sqrt{\pi/(4 \ln 2)}$, where T_A^{pk} is the peak antenna temperature and ΔV is the FWHM of the line. Sources with pure absorption spectra in any of the plotted spectral lines




Fig. 4: *Top panel.* Integrated H₂O(1₁₀ – 1₀₁) line intensity plotted vs. the integrated NH₃(1₀ – 0₀) line intensity. Symbols are as in Fig. 2. The solid line represents the linear fit to all points. The Spearman rank coefficient is 0.94. *Middle panel.* Integrated NH₃(1₀ – 0₀) line intensity vs. the integrated N₂H⁺(6 – 5) line intensity. Source 130-304 has not been included in the plot (see text). The Spearman rank coefficient is 0.96. *Bottom panel.* $\int T_A^*(\text{H}_2\text{O}) dv \times d^2$, proportional to the H₂O line luminosity, vs. continuum temperature, T_c . Symbols are as in Fig. 2.

have not been considered. Despite the low number of data points, the top and middle panels show an excellent correlation between the integrated intensities of the H₂O and NH₃ lines, and also between those of the two N-bearing molecules. In the middle panel source 130-304 has not been included in the plot because it has a completely self-absorbed water line profile, and the NH₃ line has an inverse P-Cygni profile. Therefore, the integrated intensity (in emission) of ammonia is very uncertain. The reasonable agreement between the integrated intensities of ammonia and N₂H⁺ suggests that they are emitted from similar volumes of gas, assuming similar excitation conditions.

Finally, in the bottom panel of Fig. 4 we plot the quantity $\int T_A^*(\text{H}_2\text{O}) dv \times d^2$, proportional to the H₂O line luminosity, vs. the continuum temperature, T_c . Although we do not show any linear fit in this case, there seems to be a general trend of increasing line luminosity with T_c . The 130 source with the highest water line luminosity (≥ 60 K km s⁻¹ kpc²) is 130-512 which, however, has a P-Cygni profile (see Section 4.2.5) and thus the actual line luminosity is uncertain. Except for this point, the plot shows that in these objects the “typical” water line luminosity toward the $\ell = 30^\circ$ sources is lower compared to that of the $\ell = 59^\circ$ region.

3.4. Column densities and molecular abundances

3.4.1. RADEX modelling

Since we lack direct information about the excitation temperatures and opacities of the observed transitions, we use the non-equilibrium homogeneous radiative transfer code RADEX⁹ (van der Tak et al. 2007) to correct for possible populations of molecules in unobserved excited levels, in order to relate the observed emission and absorption components to the column densities of H₂O, NH₃ and N₂H⁺.

The principal parameters needed to specify a model are the kinetic temperature, T_k , the average number density of molecular hydrogen n_{H_2} , and the line width (see for example, Persson et al. 2009). The density of each source is computed from the Hi-GAL observed size and mass, assuming a homogeneous, spherical cloud. The kinetic temperature is assumed to be equal to the dust temperature obtained from our Hi-GAL observations. All parameters are listed in Table 2. The background radiation field is an average of the Galactic background radiation in the solar neighbourhood and of the cosmic microwave background radiation.

The column density obtained from RADEX is varied until the model line brightness temperature, T_b , matches the amplitude of the Gaussian fit (Tab. 3). The line brightness temperature is obtained from the antenna temperature after correcting for main-beam efficiency, η_{mb} , and beam-filling factor, η_{bf} , i.e.:

$$T_A^* = T_b \eta_{\text{mb}} \eta_{\text{bf}} = T_b \eta_{\text{mb}} \frac{\theta_{\text{mb}}^2}{\theta_{\text{mb}}^2 + \theta_s^2} \quad (1)$$

with θ_{mb} and θ_s being the FWHM of the main-beam and source, respectively. The values of η_{bf} are also listed in Table 2. For the components in absorption, we match the observed and modelled integrated opacity instead of the brightness temperature (see Section 4.2.3).

The computed RADEX results are listed in Table 4. The results are *not* very sensitive to changes in density because of the high critical densities ($n_{\text{cr}} \gtrsim 10^7 - 10^8$ cm⁻³) of all three

⁹ <http://www.sron.rug.nl/~vdtak/radex/radex.php>

Table 4: Column densities and abundances from RADEX modelling^a.

Source	Comp. ^b	V_{lsr} [km s ⁻¹]	$N(\text{o-H}_2\text{O})^c$ [$\times 10^{14}$ cm ⁻²]	$X(\text{o-H}_2\text{O})^d$ [$\times 10^{-9}$]	V_{lsr} [km s ⁻¹]	$N(\text{N}_2\text{H}^+)^c$ [$\times 10^{14}$ cm ⁻²]	$X(\text{N}_2\text{H}^+)^d$ [$\times 10^{-9}$]	V_{lsr} [km s ⁻¹]	$N(\text{o-NH}_3)^c$ [$\times 10^{14}$ cm ⁻²]	$X(\text{o-NH}_3)^d$ [$\times 10^{-9}$]
159-441	M	27.9	450	41	27.8	0.34	6.2	27.7	37	3.4
	B	27.0	67	28.2	1.3	...
	N	26.9	0.34	0.06	27.7	0.05	0.01
159-442	M	36.7	34.2	34.8
	B	36.7
	N	34.9	35.1
159-444	N	32.0	34.3	34.4
	B	31.2
	N	27.7
	N	33.9	35.4
159-445	M	28.0	350	270	27.4	32	25	27.4	60	46
	B	28.4	300
	N	27.0	0.36	0.6	27.7	0.03	0.04
130-42	90.4	4.3	1.3	91.3	6.0	1.9
130-43 ^g	N	108.4	5	0.58	...	$\lesssim 30$	$\lesssim 4$...	$\lesssim 0.6^e$	$\lesssim 0.06^e$
130-304	90.7	33	3.3	87.8	4.9	0.49
	M	92.9	$\gtrsim 2^f$	$\gtrsim 0.4^f$	93.7	0.52	0.1
130-313	M	101.2	15	2.19	101.5	3.1	0.42	101.5	10	1.4
130-327	102.6	2.4	0.22
	M	103.8	0.73	0.1	104.6	0.21	0.04
130-376	N	70.5	39	33	...	$\lesssim 8$	$\lesssim 7$	70.5	6.1	5.1
130-512	M	63.2	110	22	61.4	26	5.1	61.8	30	5.9
	N	59.5	0.3	0.1
	N	64.0	0.2	0.08

Notes. ^(a) Using our derived $n(\text{H}_2)$ and $T_k = T_{\text{dust}}$ (Table 2). ^(b) N=narrow component, M=medium, B=broad. ^(c) Column density corrected for main beam efficiency and beam-filling factor (Table 2). ^(d) Abundance with respect to molecular hydrogen (Table 2). In the absorption components we have used $N(\text{H}_2)/2$ assuming that the absorption is probing approximately half of the total column density in front of the background continuum. ^(e) 3σ limit. ^(f) Optically thick; τ is assumed to be 5. ^(g) Tentative detection (see Section 3.2).

molecular transitions. However, the column density is sensitive to changes in kinetic temperature if $T_k \lesssim 15$ K. For sources with non-detections in one (or two) of the molecular transitions, the upper limit to the column density has been estimated by using the 3σ limits for T_A^* listed in Table 3.

Abundances with respect to molecular hydrogen, $X(\text{mol}) = N(\text{mol})/N(\text{H}_2)$, are estimated using the derived volume densities and observed source sizes to compute $N(\text{H}_2)$. For the components in absorption we divide the estimated $N(\text{H}_2)$ by a factor of two since the absorption is probing half of the total column density that is in front of the continuum source. For the outflow components we cannot estimate the abundance since $N(\text{H}_2)$ is likely orders of magnitude lower than that of the whole cloud.

Table 4 indicates that the water molecular abundance is possibly higher toward sources in the $\ell = 59^\circ$ region. Our results may be affected by the uncertainties in the derived Hi-GAL physical parameters and other assumptions. However, combined with the result that the typical water line luminosity toward the $\ell = 30^\circ$ sources (where no broad component is detected) is lower compared to that of the $\ell = 59^\circ$ region (see bottom panel of Fig. 4), we conclude that our observations support a scenario where the water abundance is linked to shocked gas.

3.4.2. Column density for the absorption component

We also used an alternative approach to compute the $\text{o-H}_2\text{O}$ column density of the envelope gas responsible for the observed absorption in the water lines, which we also use to double-check the RADEX results. We first derived the optical depth at the dip of the absorption profile as:

$$\tau_{\text{dip}} = -\ln\left(\frac{T_A^{\text{em}} - T_A^{\text{dip}}}{T_A^{\text{em}}}\right) \quad (2)$$

where, in order to include the line contribution to the absorption background, the antenna temperature T_A^{em} represents the sum of all Gaussian components in emission (at the velocity of the dip in the absorption component) and of the continuum temperature T_c . This procedure assumes that the absorber is located in front of both the continuum and the line emitting source. T_A^{dip} represents the peak absolute value of the antenna temperature of the Gaussian component in absorption. The integrated optical depth of the absorption component is then calculated as $\tau_{\text{abs}} = \int \tau dv = 1.06 \tau_{\text{dip}} \Delta V_{\text{abs}}$ where ΔV_{abs} represents the line FWHM of the Gaussian component in absorption. Finally, the column density can be estimated as:

$$N_{\text{tot}} \approx N_l = \frac{8\pi\nu_{\text{ul}}^3 g_l}{c^3 A_{\text{ul}} g_u} \left[1 - \exp\left(-\frac{h\nu_{\text{ul}}}{kT_{\text{ex}}}\right)\right]^{-1} \tau_{\text{abs}} \quad (3)$$

Table 5: Mean values of bolometric luminosity, mass and continuum temperature.

Region	Sources with detection			Sources with no detection ^a		
	Mean L	Mean M	Mean T_c	Mean L	Mean M	Mean T_c
	[L_\odot]	[M_\odot]	[K]	[L_\odot]	[M_\odot]	[K]
$\ell = 30$	16200 ± 17600	3430 ± 3300	93 ± 65	2145 ± 2570	690 ± 305	37 ± 5
$\ell = 59$	1520 ± 735	190 ± 105	63 ± 30	90 ± 64	440 ± 390	26 ± 4

Notes. ^(a) Only sources where all three parameters could be measured were included.

where A_{ul} and ν_{ul} represent the Einstein A coefficient and frequency associated with the transition between the lower and upper level, indicated by the subscripts l and u , respectively, and with degeneracies g_l and g_u . Physical constants are indicated with standard symbols and T_{ex} represents the excitation temperature. Since we do not know T_{ex} we will assume for the moment that $T_{ex} \ll h\nu_{ul}/k = 26.6$ K for the 557 GHz water line. Therefore, the exp factor in Eq.(3) can be neglected.

We thus obtain remarkably similar o -H₂O column densities, varying from about 10^{13} cm⁻² to $\approx 5 \times 10^{13}$ cm⁻² (consistent with the RADEX results), with the highest value obtained toward the source I30-304, where in fact the absorption appears to be almost saturated, since the line profile is nearly down to the zero-temperature level. The column densities of the outer envelope are thus quite similar to those estimated for both low- (Kristensen et al. 2012) and high-mass (van der Tak et al. 2013) star forming regions.

4. Discussion

4.1. Average physical conditions in the $\ell = 30^\circ$ and $\ell = 59^\circ$ regions

As shown by Tables 2 and 3, in the $\ell = 30^\circ$ region about 41% of the sources have a detection in at least one of the observed spectral lines. This percentage drops to about 11% in the $\ell = 59^\circ$ region. We also note that all of the sources with at least one detection are *proto*-stellar in both galactic regions. The significant difference in detections between the two regions may be a consequence of the different fraction of *proto*-stellar sources observed in each region, corresponding to 59% and 14% in $\ell = 30^\circ$ and $\ell = 59^\circ$, respectively. In fact, the fraction of *proto*-stellar sources with at least one detection corresponds to 70% and 80% in the $\ell = 30^\circ$ and $\ell = 59^\circ$ regions, respectively, making the difference in the detection rate less significant.

In Table 5 we list the mean values of the clump mass and bolometric luminosity, as derived from the Hi-GAL observations, for the observed sources in the two galactic regions. We note that the mean values of the luminosity for sources with a detection in one of the observed spectral lines are much higher than the luminosity of sources with no detection at all. The scatter around the mean value is quite high and thus our conclusion is only tentative and will need more data to be confirmed. Furthermore, the mean values in the $\ell = 59^\circ$ region are derived from only two sources, since in sources I59-442 and I59-444 the distance could not be determined reliably.

As with luminosity, Table 5 shows that T_c has a higher mean value (by more than a factor of 2) in sources with at least one detected spectral line, for both observed regions. Given that all of these sources are *proto*-stellar, the mean values of T_c for sources with and without detection are in fact very similar to the mean values of T_c for *proto*-stellar and starless sources, respectively. Compared to luminosity and mass, the mean values of T_c show a

lower scatter, which is still quite high only in the case of sources with detection in the $\ell = 30^\circ$ region. In fact, the scatter for the case of sources with no detection is quite low in both regions, despite the low number of sources. Given the marginal correlation between luminosity and T_c (see Section 3.3) this result is not totally unexpected. However, given that the continuum temperature is a distance-independent quantity this result is more significant in characterizing the sources. Finally, from Table 5 we also note that the mean values of the mass for sources with and without a detection do not show any specific relation. Thus, if we assume tentatively that the difference in the mean value of luminosity (and T_c) between sources with and without detections is indeed real, then this might be a consequence of evolutionary effects rather than an effect associated with more massive clumps (see also Section 4.4).

4.2. Gas kinematics

Our decomposition of the H₂O line profiles (see Section 3.2) is similar to that of Kristensen et al. (2012) toward low-mass protostars, and to the description of the H₂O profiles of van der Tak et al. (2013) toward high-mass protostars. Herpin et al. (2012) found a similar profile decomposition toward the massive protostar W43-MM1. In the following we first describe the general properties of the line shapes in the $\ell = 30^\circ$ and $\ell = 59^\circ$ regions, and then we discuss the possible physical origin of each component.

4.2.1. Line profile components in the $\ell = 59^\circ$ region

Here we summarize the main observational properties of the line profiles of the three transitions, H₂O(1₁₀ - 1₀₁), NH₃(1₀ - 0₀), and N₂H⁺(6 - 5) toward the $\ell = 59^\circ$ region.

H₂O. All detected sources of the $\ell = 59^\circ$ region have a strikingly similar water line-profile, which is always characterized by an absorption dip at the source velocity (as given by N₂H⁺). This absorption dip does not appear to be saturated. As shown by Table 3, the offset between these absorption components and the source velocity does not appear to be significant. The broad and medium Gaussian components have line widths exceeding those of N₂H⁺, which is tracing the more quiescent ambient gas, and are thus likely originating in *proto*-stellar outflows. The water line profile in source I59-444 is somewhat different from that of the other three detected sources in the $\ell = 59^\circ$ region since it is characterized by an inverse P-Cygni profile overlaid on a broad component (see Section 4.2.5).

N₂H⁺ and NH₃. While the N₂H⁺ line-profile looks quite Gaussian, and thus used to set the source velocity, NH₃ shows an asymmetric line-profile which exhibits self-absorption (less evident in I59-442), though not at the exact same velocity as absorption in the water line (see Table 3). Absorption is also present in




Fig. 5: *Top*. Line width of the NH_3 line vs. line width of the N_2H^+ line, when both are detected in emission. The dashed line represents the $\Delta V(\text{NH}_3) = \Delta V(\text{N}_2\text{H}^+)$ locus. *Bottom*. Line width of the broad component of the water line profile vs. the line width of the medium component for the $\ell = 59^\circ$ sources. Symbols are as in Fig. 2.

the NH_3 spectrum of 159-444. Finally, we note that one of the Gaussian components of the NH_3 line toward sources 159-441 and 159-442 has a line width of 10.1 km s^{-1} and 6.4 km s^{-1} (see Table 3). This suggests that the outflows observed in the water line are also partially detected in the ammonia line, at least in source 159-441.

4.2.2. Line profile components in the $\ell = 30^\circ$ region and comparison of line widths

The nature of the observed lines toward this region is significantly different from that of the sources in the $\ell = 59^\circ$ region. The line intensities are in general lower compared to those measured toward the $\ell = 59^\circ$ region, which is likely a consequence of the average larger distance to the $\ell = 30^\circ$ sources.

H_2O . The water line is detected both in emission and absorption, but the line-profiles are much more varied. As already

mentioned, in the sources 130-304 and 130-327 the Gaussian component in absorption appears to be almost saturated. No line wings are detected above the noise level. In several sources (e.g., 130-304 and 130-327) the presence of multiple water absorption lines is evidently tracing the diffuse interstellar gas (which will be the subject of a separate paper, Persson et al., in prep.), that are likely a consequence of the larger amount of diffuse gas encountered along the $\ell = 30^\circ$ direction. In Fig. 4 we have already shown that the water line luminosity toward the $\ell = 30^\circ$ sources is lower compared to that of the $\ell = 59^\circ$ region.

N_2H^+ and NH_3 . As for the $\ell = 59^\circ$ region, the N_2H^+ line-profile looks Gaussian, and (when detected) it is also used to set the source velocity. In most cases the line-profiles of water and ammonia are consistent, i.e., they are either both in emission or absorption. Toward source 130-304 the NH_3 spectrum exhibits an inverse P-Cygni profile and the component in absorption looks almost saturated as in the water line. Toward source 130-327 (and possibly 130-42), ammonia also shows absorption lines along the line of sight, as already observed in the water spectra. The maximum velocity difference between NH_3 and N_2H^+ , when both lines are seen in emission, is $\approx 0.9 \text{ km s}^{-1}$ in source 130-42.

We have compared the line widths of the NH_3 and N_2H^+ lines, as shown in Fig. 5. Except for source 130-304, which shows an inverse P-Cygni profile in the ammonia spectrum, in all other sources we note that $\Delta V(\text{NH}_3) \geq \Delta V(\text{N}_2\text{H}^+)$ including all four detected sources in the $\ell = 59^\circ$ region. We also note that $\Delta V(\text{NH}_3)$ tends to be somewhat higher in the $\ell = 59^\circ$ sources. Since molecular outflows are the most distinctive feature of the detected $\ell = 59^\circ$ sources, this is an indication that the line width of NH_3 is being broadened by these outflows. Likewise, the N_2H^+ emission must then originate in a volume where the molecular gas is not significantly affected by the molecular outflows.

4.2.3. Narrow component

The general appearance of the narrow component, mostly in absorption, indicates a high H_2O column density and a low excitation temperature, suggesting that the absorption is being caused by the outer envelope and ambient cloud. Source 130-376 is the only one showing a clear narrow component in emission, which could be related to temperature, as appearance in emission requires an excitation temperature above the continuum level. We note, in fact, that 130-376 has the *lowest* continuum temperature (23 mK , see Table 2) among sources in the $\ell = 30^\circ$ region with a detectable continuum emission. As mentioned earlier, in the other $\ell = 30^\circ$ sources the water line, either in absorption or emission, is broader than $\approx 5 \text{ km s}^{-1}$ and has thus been classified as a medium component.

As already noted by Kristensen et al. (2012), because the absorption of the narrow component is seen against both the outflow and continuum emission, the absorbing layer must be located in front of both the emitting layers, i.e., the outflows must also be embedded. Figure 6 shows that the narrow component is either centred at the source velocity (offset $\leq 1 \text{ km s}^{-1}$) or blueshifted (130-512) relative to the systemic velocity, as given by N_2H^+ . In most $\ell = 59^\circ$ sources, however, the narrow component is also slightly blueshifted relative to N_2H^+ emission, suggesting that the envelope, where the absorbed narrow component originates, is expanding. In addition, Fig. 6 shows that the narrow component is also blueshifted relative to both the




Fig. 6: Central velocities of the narrow (V_{nar}) and medium (V_{med}) Gaussian components of the observed H_2O lines compared with each other and with the systemic velocity from N_2H^+ . Symbols are as in Fig. 2.

medium and broad (not shown) components. The only source where $V_{\text{nar}}(\text{H}_2\text{O})$ is red-shifted compared to both $V_{\text{med}}(\text{H}_2\text{O})$ and $V_{\text{broad}}(\text{H}_2\text{O})$ is 159-444, which has an inverse P-Cygni profile, which is instead typical of an infalling envelope.

Contrary to van der Tak et al. (2013) we do not find any correlation between the source mass (which is mostly due to the envelope contribution, being derived from the Hi-GAL observations of cold dust emission) and the line width of the narrow component. In Section 3.4 using RADEX we have estimated the $o\text{-H}_2\text{O}$ column density of the envelope gas responsible for the

observed absorption toward all $\ell = 59^\circ$ sources and toward the $\ell = 30^\circ$ sources 130-304, 130-327 and 130-512 (see also Table 3).

4.2.4. Outflows and shocked gas

The H_2O line profiles are characterized by two additional and broader components: a broad emission component, always in emission and visible only towards the $\ell = 59^\circ$ sources, and a medium absorption or emission component seen in almost all sources. The broad and medium line components have widths exceeding the line width typical of the more quiescent envelope (e.g., Jørgensen et al. 2002) and are thus likely related to a proto-stellar outflow.

In the $\ell = 59^\circ$ sources, both Gaussian components are required to fit the line profile. The medium component has already been observed by Kristensen et al. (2012), toward a sample of low-mass proto-stellar objects, by van der Tak et al. (2013) toward a sample of high-mass star forming regions, and also by Herpin et al. (2012) toward the massive protostar W43-MM1. Figure 6 shows that the medium component in most cases is centred at the source velocity (offset $\lesssim 1 \text{ km s}^{-1}$), with $V_{\text{med}}(\text{H}_2\text{O}) - V_{\text{ISR}}(\text{N}_2\text{H}^+)$ having positive values in almost all cases. We already noted in the previous section that the narrow component is blueshifted relative to the medium component. van der Tak et al. (2013) found that their medium component (which they call “narrow outflow component”) usually appears in absorption and is instead blueshifted relative to the narrow component (which they call “envelope component”). However, it should be noted that their decomposition of the line profiles is based on higher-excitation water lines.

Both Kristensen et al. (2012) and van der Tak et al. (2013) associate the origin of the medium component to the region, closer to the central protostar, where the gas is currently being shocked, whereas they associate the broader component with the swept-up gas of the molecular outflow. However, Herpin et al. (2012) report that their medium velocity component is likely due to a combination of turbulence and infall. A more detailed analysis of the different velocity components in shocked molecular gas has been carried out, for example, by Busquet et al. (2014) in the L1157-B1 shock region. They performed a multi-line H_2O survey probing a wide range of excitation conditions. In addition to the component in absorption (probably due to the photodesorption of icy mantles at the edge of the cloud), two gas components, connected with a J-type shock associated with a jet and with different excitation conditions, were found. The first component is a relatively extended ($\sim 10''$ corresponding to $\sim 2 \times 10^4 \text{ AU}$), warm ($\sim 200 \text{ K}$), and dense ($\sim 10^6 \text{ cm}^{-3}$) region, and the second is a more compact (less than $5''$ or $\sim 10^4 \text{ AU}$), hot ($\sim 1000 \text{ K}$) and tenuous ($\sim 10^3 - 10^4 \text{ cm}^{-3}$) component. The water abundance is very high in the hot gas ($\sim 10^{-4}$) and it decreases down to $\sim 10^{-6}$ in the warm component. Both warm and hot components have their peak at low-velocities, with the hot region dominating the total flux at the high end of the velocity range.

In our work, and with only the data presented here, it is not possible to infer the exact physical origin of the medium component of the $\text{H}_2\text{O}(1_{10} - 1_{01})$ transition. However, the correlation between the line widths of the broad and medium components in the $\ell = 59^\circ$ sources, shown in the bottom panel of Fig. 5, suggests an outflow or shock origin for the medium component as well. We also note that Kristensen et al. (2012) speculate that the phenomenon producing the masers should also have a thermal component associated with it, which then produces the

medium component. We analyze the association with masers in Section 4.3.

As far as sources in the $\ell = 30^\circ$ region are concerned, no broad component has been detected. Observationally, this may be a consequence of the larger distances of these sources compared to the $\ell = 59^\circ$ region. Alternatively, the viewing angle of the molecular outflows and/or shocks may also determine whether a source shows a broad, or both a broad and medium component. Because of the small sample size of $\ell = 30^\circ$ sources, such an effect could be statistically significant, although in the $\ell = 59^\circ$ region, with a comparable sample size, *all* sources show a broad component. An alternative, more physical reason for the lack of a broad component toward the $\ell = 30^\circ$ region could be a different evolutionary phase (see Section 4.4).

As far as NH_3 and N_2H^+ are concerned, the line width of the ammonia line is likely affected by the presence of molecular outflows. We note, in particular, that source 159-441 is the only one where the NH_3 profile shows the same three components as the water line. While the water narrow component is generally blueshifted relative to the medium and broad components, all ammonia components have remarkably similar velocities. Three more sources show a NH_3 narrow and medium component (or two narrow components): 159-442, 159-444, and 130-304. The two ammonia components have very similar velocities also in 159-442. In the other two sources, 159-444, and 130-304, however, inverse P-Cygni profiles are observed and will be discussed in the next Section.

4.2.5. Regular and inverse P-Cygni profiles

In a total of three cases, we have observed a regular or inverse P-Cygni profile, which are usually associated to envelope expansion and infall, respectively. An inverse P-Cygni profile is characterised by an absorption feature which is red-shifted relative to an emission feature, and is typically caused by a velocity and excitation gradient in the gas falling in toward a central object. Two sources have inverse P-Cygni profiles, 130-304 and 159-444. In the case of source 159-444, the inverse P-Cygni profile is visible in the spectrum of water as superposed on the broad outflow component. In this source a red-shifted absorption is also visible in the ammonia spectrum. Toward source 130-304 the inverse P-Cygni profile is only detected in the NH_3 spectrum, whereas the water spectrum shows multiple absorption lines from the diffuse interstellar gas, and at the source systemic velocity the water line profile appears almost completely self-absorbed (Section 3.4.2). A recent example of multiple ammonia lines, observed with *Herschel* toward the massive star forming region G34.3+0.15, showing inverse P-Cygni profiles can be found in Hajigholi et al. (submitted). The only regular P-Cygni profile we have detected, toward source 130-512, is also superposed on what looks like an outflow/shock component. These three profiles are thus all different at some level and suggest a great variety of kinematic and excitation conditions.

In order to give the simplest possible description of a contracting or expanding system that can reproduce the observed line profiles, Di Francesco et al. (2001) used a simple model consisting of two slabs moving towards each other with a continuum source between them. This model was a modification of a previous “two-layer” code used by Myers et al. (1996). This model is appealing because of its simplicity, but on the other hand it gives only the crudest picture of the optical depth and excitation temperature of the absorbing and emitting gas. In addition, the model uses a relatively large number of free parameters (eight in total, if none is held fixed), including the excitation temper-




Fig. 7: From top to bottom, inverse P-Cygni profiles detected toward sources 159-444 (H_2O) and 130-304 (NH_3), and regular P-Cygni profile detected toward 130-512 (H_2O). The infall (or expansion, in the case of 130-512) model is shown on top (red, dashed line). The broad outflow component has been subtracted from the spectrum of source 159-444, whereas a narrow absorption component has been added back to the spectrum of 130-512.

ature of the front/rear layers and the continuum source (T_f , T_r , T_c) the filling factor of the absorbing source with respect to the telescope beam (Φ), the peak optical depth of the front and rear layers (τ_o), the turbulent velocity dispersion in each layer (σ), the infall velocity (V_{in}) and the systemic velocity (V_{lsr}).

Table 6: Best-fit parameters of the infall/outflow modelling. See text for a description of the parameters used in the fit.

Source	Species	V_{in} [km s ⁻¹]	σ [km s ⁻¹]	V_{lsr}^a [km s ⁻¹]	τ_o	Φ	T_c^a [K]	T_f^a [K]	T_r [K]
159-444	H ₂ O	0.5	1.5	34.3	1.7	0.14	37.0	4.0	17.2
130-304	NH ₃	2.5	1.7	90.7	3.1	0.01	350.0	4.0	10.5
130-512 ^b	H ₂ O	-2.1	1.6	61.4	1.3	0.01	63.0	4.0	11.0 ^a

Notes. ^(a) Fixed parameter, see text. ^(b) Tentative result.

Therefore, applying the simple two-slab model described above to the two sources with inverse P-Cygni profiles, we obtain the best-fit infall models shown in Fig. 7, with the corresponding best-fit parameters listed in Table 6. In the case of source 159-444, the broad outflow component (and an additional narrow absorption feature, blue-shifted with respect to the inverse P-Cygni profile, see Table 3) have been subtracted to the original spectrum before applying the model. For both sources the value of T_c has been fixed to that shown in Table 2. We have also fixed the value of V_{lsr} to the velocity of N₂H⁺ (see Table 3). In section 4.2.3 we suggested that the gas front layer may have a low excitation temperature and thus we fixed it to the approximate intermediate value ($T_f \simeq 4$ K) shown by Myers et al. (1996) in their Fig.1. As a comparison Hajigholi et al. (submitted) find an excitation temperature of $\simeq 7-8$ K in the absorbing envelope of source G34.3+0.15.

By comparing different infall model profiles to the observations we find that V_{in} and σ can vary by as much as 0.2 – 0.5 km s⁻¹, and are not very sensitive to T_f . However, these tests do not take into account the uncertainty on the beam-filling factor, used to convert the antenna temperature to line brightness temperature (see Section 3.4), which may also affect the best-fit results. Thus, our results show that while the infall velocity toward 159-444 is comparable to the values found by Di Francesco et al. (2001) and Kristensen et al. (2012) (who analyze low-mass protostars), the values of σ (and the value of V_{in} toward 130-304) are somewhat higher, since these authors find $V_{\text{in}}, \sigma \lesssim 1$ km s⁻¹. Our estimated values of σ are consistent with those found by Hajigholi et al. (submitted) toward G34.3+0.15. Some of the high-mass protostars studied by van der Tak et al. (2013) also show regular and inverse P-Cygni profiles, but they do not analyze the line profiles and thus we cannot directly compare our results. Instead, we can compare our results with those of Herpin et al. (2012) toward the massive protostar W43-MM1, where they derive infall velocities as high as 2.9 km s⁻¹ and turbulent velocities ≥ 2 km s⁻¹. These values are much much more similar to our estimates of σ and of the infall velocity toward 130-304.

As far as the only source, 130-512, with a regular P-Cygni profile in our sample, this source shows a narrow absorption component at a velocity of 64 km s⁻¹. Therefore, assuming that this additional absorption feature is not related to the overall P-Cygni profile, and in order to reconstruct the hypothetical P-Cygni profile we added back this component to the water spectrum. The two-layers model described earlier can also be applied to the case of expanding gas, the only change in the model is that the infall velocity becomes negative. We show our best-fit result in the bottom panel of Fig. 7, obtained by fixing the value of T_r , which would otherwise diverge. Clearly, the fit is not very good and it is difficult to determine whether this may be due to contamination of the line profile, up to the level of possibly resulting

in a fake P-Cygni profile. For this reason our results in Table 6 are labelled as tentative.

While the infall and turbulent velocities are distance-independent and, at least in this toy model, do not depend on the mass distribution within the source, determining the mass infall rate would require the knowledge of the characteristic distance of the infall zone or the mass undergoing infall. If the envelopes are in free fall, the velocity at a radius R_{in} in the envelope is given by $V_{\text{in}} = \sqrt{2GM_{\text{enc}}/R_{\text{in}}}$ where M_{enc} is the mass enclosed within the radius R_{in} . Assuming spherical infall symmetry, the mass infall rate can be estimated as (e.g., Kristensen et al. 2012):

$$\dot{M} = 4\pi\mu m_{\text{H}} n_{\text{in}} (2GM_{\text{enc}})^2 V_{\text{in}}^{-3} \quad (4)$$

with n_{in} the density at the infall radius, m_{H} and $\mu = 2.8$ the hydrogen atom mass and mean molecular weight, respectively. Therefore, in order to estimate \dot{M} it is necessary to be able to determine either M_{enc} or the infall radius R_{in} . We do not have a reliable distance determination for source 159-444 (see Table 2), and thus we cannot estimate any mass. In the case of source 130-304 we do have a distance estimate and a mass, but the large mass and relatively large distance (see Table 2) suggest that the estimated mass represents a much larger mass than that enclosed within the infalling region. As an example, if we used a typical infall radius estimated by Herpin et al. (2012) ($R_{\text{in}} \sim 6 \times 10^{17}$ cm) then in source 130-304 we would get a mass infall rate $\dot{M} \sim 5 \times 10^{-2} M_{\odot} \text{ yr}^{-1}$, using as n_{in} the density given in Table 2 and V_{in} from Table 6. Not surprisingly, \dot{M} is comparable to the values estimated by Herpin et al. (2012) toward the massive proto-stellar clump W43-MM1, and is much higher than the infall rates determined by Kristensen et al. (2012) toward low-mass proto-stellar sources.

4.3. Association with maser emission

We searched for masers associated with the detected sources (within a 30'' radius) and found a total of six sources associated with maser emission: all four detected sources in the $\ell = 59^\circ$ region and two more in the $\ell = 30^\circ$ field. Sources 159-441 and 159-445, being very close, are actually associated with the same methanol maser. All sources, except for 159-442, are associated with a methanol 6.7 GHz maser. Source 159-442 is the only one associated with a 22 GHz water maser (B193852.5+235736, Palla et al. 1991), while source 130-304 is also associated with an OH maser.

The methanol masers associated with sources 159-441/159-445 and 159-444 were found by comparing our observations with those carried out by Olmi et al. (2014b) toward the $\ell = 30^\circ$ and $\ell = 59^\circ$ regions in search of 6.7 GHz CH₃OH and 6.0 GHz OH masers. The masers associated with 159-441/159-445 and 159-444 are G59.63–0.19 and G59.78+0.63, respectively, and they have peak flux densities of 0.58 and 0.03 Jy. As discussed by

Olmi et al. (2014b), these two methanol masers belong to a new class of low-brightness masers that could represent an earlier stage of evolution.

The methanol masers have velocities which agree well with the velocities observed in our detected sources (differences are $\lesssim 1-2 \text{ km s}^{-1}$). However, the velocity of the 22 GHz water maser ($V_{\text{lsr}} \approx 50 \text{ km s}^{-1}$) associated with source 159-442 significantly differs from the central velocity of any of the Gaussian components of its H_2O line profile ($V_{\text{lsr}} \approx 35 - 37 \text{ km s}^{-1}$). If we assume that the physical association between these two sources is indeed real, given the small angular separation between them ($\approx 4 \text{ arcsec}$), then one might speculate that the water maser is originated somewhere along the path of the outflow. In fact, the broad Gaussian component in 159-442 has a line width that encompasses the velocity of the water maser.

4.4. Evolutionary trends

The relative intensities of the various components used to decompose the line profiles of the water spectra have already been used in an attempt to identify possible evolutionary sequences. For example, Kristensen et al. (2012) associate the disappearance of the broad component with later evolutionary stages in their low-mass protostars sample. van der Tak et al. (2013) then propose that the appearance of the narrow component (or “envelope” component) in emission is an indicator of a warmer envelope, and thus associated to later evolutionary stages. We note that while in all $\ell = 59^\circ$ sources the narrow component appears in absorption, toward the $\ell = 30^\circ$ region sources 130-43 and 130-376 are characterized by a narrow component in emission (see Table 3; see also Section 3.2 regarding the water detection toward 130-43). In source 130-512 the narrow component is in absorption, but this is a more complex source with a regular P-Cygni profile, as discussed in Section 4.2.5.

An alternative method to analyze the evolutionary stages of our sources makes use of the evolutionary tracks previously shown in Fig. 2. These tracks were determined following a method similar to that discussed by Molinari et al. (2008), who proposed an evolutionary sequence for high-mass protostars in terms of two parameters: the envelope mass and the bolometric luminosity. These authors suggest an evolutionary sequence which is concentrated in two main phases: protostars first accrete mass from their envelopes, and later disperse their envelopes by winds and/or radiation. In Fig. 2 the calculated evolutionary tracks thus rise upward almost vertically in luminosity during the accretion phase, and then proceed horizontally to the left (to lower masses) during the envelope dispersal phase. The positions of our sources in this plot suggest that some (but not all) of the sources with at least one line detection are, for similar ranges of mass, in a more advanced stage of evolution. This is certainly the case for the detected $\ell = 59^\circ$ sources with a well defined distance (159-441 and 159-445), while for sources detected toward the $\ell = 30^\circ$ region a preferential evolutionary phase is not apparent. None of these sources, however, seem to belong to the late phase of envelope dispersal.

There are other indicators that can be used to analyze the relative evolutionary phase of the $\ell = 30^\circ$ and $\ell = 59^\circ$ regions. For example, from the discussion about T_c in Section 4.1 and Table 5, there is a trend¹⁰ of higher T_c toward the detected $\ell = 30^\circ$ sources, suggesting they may be in a more advanced stage of proto-stellar evolution, compared to sources in

¹⁰ We use the word “trend” because of the large scatter shown by the T_c values in Table 5.

the $\ell = 59^\circ$ region. Furthermore, as already noted in Section 3.3, the typical water line luminosity toward the $\ell = 30^\circ$ sources is lower compared to that of the $\ell = 59^\circ$ region. Kristensen et al. (2012) find a similar decrease in water emission between their Class 0 and Class I low-mass protostars. Finally, as stated in Section 4.3 two of the $\ell = 59^\circ$ sources are associated with low-brightness methanol masers that could indicate an earlier phase of evolution. Source 130-304 is the only one associated with an OH maser, which is sometimes described as characterizing later evolutionary stages (Breen et al. 2010) compared to 6.7 GHz methanol masers. Therefore, a firm conclusion about the specific evolutionary phase of the observed sample is limited by the small number of sources. However, various indicators tentatively suggest that the detected sources toward the $\ell = 30^\circ$ region are in a later evolutionary phase compared to those in the $\ell = 59^\circ$ region.

5. Summary and conclusions

Using the HIFI instrument of *Herschel* we have observed the $o\text{-NH}_3(1_0 - 0_0)$ (572 GHz), $o\text{-H}_2\text{O}(1_{10} - 1_{01})$ (557 GHz) and $\text{N}_2\text{H}^+(6 - 5)$ (559 GHz) lines toward a sample of 52 high-mass starless and proto-stellar clumps selected from the Hi-GAL survey. The target sources are concentrated in two galactic fields, at $\ell = 30^\circ$ and $\ell = 59^\circ$, and were observed during the Science Demonstration Phase of *Herschel*. At least one of the three selected molecular lines was detected in 4 out of 35 sources toward the $\ell = 59^\circ$ region, and 7 out of 17 objects toward the $\ell = 30^\circ$ region.

Almost all water spectra can be described as the sum of a narrow ($\text{FWHM} \sim 3 - 5 \text{ km s}^{-1}$), a medium ($\text{FWHM} \sim 6 - 10 \text{ km s}^{-1}$) and a broad (up to $\text{FWHM} \approx 30 \text{ km s}^{-1}$) velocity component. All detected sources of the $\ell = 59^\circ$ region have a strikingly similar water line-profile, characterized by the broad Gaussian component associated with the presence of molecular outflows, which are also observed in the NH_3 line, at least toward one source. Therefore, in these sources there is strong evidence that water line emission is linked to the presence of shocked molecular gas.

The situation is very different in the $\ell = 30^\circ$ region, with no two profiles being identical, and with no clear detection of a broad component in the water spectra. We find that the continuum temperature, T_c , is correlated with the source bolometric luminosity. We also find an indication that the water line luminosity increases with T_c . The detected sources in the $\ell = 59^\circ$ region have both a higher water line luminosity and water abundance, compared to sources detected in the $\ell = 30^\circ$ region. Although we have few sources, this trend constitutes additional support to a scenario where the water abundance is linked to shocks.

In terms of the gas kinematics, the general appearance in absorption of the narrow Gaussian component indicates a high H_2O column density and a low excitation temperature, suggesting that the absorption is being caused by the outer envelope and ambient cloud. The correlation between the line widths of the broad and medium components in the $\ell = 59^\circ$ sources suggests an outflow or shock origin for the medium component. In addition, we also detect in a few sources inverse and regular P-Cygni profiles, though we do not find any link with other source parameters. A firm conclusion about the specific evolutionary phase of the observed sample is limited by the small number of sources. However, various indicators tentatively suggest that the detected sources toward the $\ell = 30^\circ$ region are in a later evolutionary phase compared to those in the $\ell = 59^\circ$ region.

Acknowledgements. HIFI has been designed and built by a consortium of institutes and university departments from across Europe, Canada and the USA under

the leadership of SRON Netherlands Institute for Space Research, Groningen, The Netherlands and with major contributions from Germany, France and the USA. Consortium members are: Canada: CSA, U.Waterloo; France: CESR, LAB, LERMA, IRAM; Germany: KOSMA, MPIfR, MPS; Ireland, NUI Maynooth; Italy: ASI, IAPS-INAf, Osservatorio Astrofisico di Arcetri-INAf; Netherlands: SRON, TUD; Poland: CAMK, CBK; Spain: Observatorio Astronomico Nacional (IGN), Centro de Astrobiología (CSIC-INTA). Sweden: Chalmers University of Technology - MC2, RSS & GARD; Onsala Space Observatory; Swedish National Space Board, Stockholm University - Stockholm Observatory; Switzerland: ETH Zurich, FHNW; USA: Caltech, JPL, NHSC. CMP acknowledges generous support from the Swedish National Space Board.

References

- Bergin, E. A. & Tafalla, M. 2007, *ARA&A*, 45, 339
- Bontemps, S., Motte, F., Csengeri, T., & Schneider, N. 2010, *A&A*, 524, A18
- Breen, S. L., Caswell, J. L., Ellingsen, S. P., & Phillips, C. J. 2010, *MNRAS*, 406, 1487
- Busquet, G., Lefloch, B., Benedettini, M., et al. 2014, *A&A*, 561, A120
- Caselli, P., Keto, E., Bergin, E. A., et al. 2012, *ApJ*, 759, L37
- de Graauw, T., Helmich, F. P., Phillips, T. G., et al. 2010, *A&A*, 518, L6
- Di Francesco, J., Myers, P. C., Wilner, D. J., Ohashi, N., & Mardones, D. 2001, *ApJ*, 562, 770
- Elia, D., Schisano, E., Molinari, S., et al. 2010, *A&A*, 518, L97
- Green, J. A. & McClure-Griffiths, N. M. 2011, *MNRAS*, 417, 2500
- Herpin, F., Chavarría, L., van der Tak, F., et al. 2012, *A&A*, 542, A76
- Ho, P. T. P. & Townes, C. H. 1983, *ARA&A*, 21, 239
- Jackson, J. M., Rathborne, J. M., Shah, R. Y., et al. 2006, *ApJS*, 163, 145
- Jørgensen, J. K., Schöier, F. L., & van Dishoeck, E. F. 2002, *A&A*, 389, 908
- Kristensen, L. E. & van Dishoeck, E. F. 2011, *Astronomische Nachrichten*, 332, 475
- Kristensen, L. E., van Dishoeck, E. F., Bergin, E. A., et al. 2012, *A&A*, 542, A8
- Larson, R. B. 1981, *MNRAS*, 194, 809
- Leurini, S., Gusdorf, A., Wyrowski, F., et al. 2014, *A&A*, 564, L11
- Marseille, M. G., van der Tak, F. F. S., Herpin, F., & Jacq, T. 2010, *A&A*, 522, A40
- Molinari, S., Pezzuto, S., Cesaroni, R., et al. 2008, *A&A*, 481, 345
- Molinari, S., Swinyard, B., Bally, J., et al. 2010, *PASP*, 122, 314
- Motte, F., Bontemps, S., Schilke, P., et al. 2007, *A&A*, 476, 1243
- Myers, P. C., Mardones, D., Tafalla, M., Williams, J. P., & Wilner, D. J. 1996, *ApJ*, 465, L133
- Netterfield, C. B., Ade, P. A. R., Bock, J. J., et al. 2009, *ApJ*, 707, 1824
- Olmi, L., Ade, P. A. R., Anglés-Alcázar, D., et al. 2009, *ApJ*, 707, 1836
- Olmi, L., Anglés-Alcázar, D., Elia, D., et al. 2013, *A&A*, 551, A111
- Olmi, L., Anglés-Alcázar, D., Elia, D., et al. 2014a, *A&A*, 564, A87
- Olmi, L., Araya, E. D., Chapin, E. L., et al. 2010, *ApJ*, 715, 1132
- Olmi, L., Araya, E. D., Hofner, P., et al. 2014b, *A&A*, 566, A18
- Palla, F., Brand, J., Comoretto, G., Felli, M., & Cesaroni, R. 1991, *A&A*, 246, 249
- Persson, C. M., Olberg, M., Hjalmarson, Å., et al. 2009, *A&A*, 494, 637
- Pilbratt, G. L., Riedinger, J. R., Passvogel, T., et al. 2010, *A&A*, 518, L1
- Rieke, G. H., Young, E. T., Engelbracht, C. W., et al. 2004, *ApJS*, 154, 25
- Roelfsema, P. R., Helmich, F. P., Teyssier, D., et al. 2012, *A&A*, 537, A17
- Russeil, D. 2003, *A&A*, 397, 133
- Russeil, D., Pestalozzi, M., Mottram, J. C., et al. 2011, *A&A*, 526, A151
- Russeil, D., Zavagno, A., Motte, F., et al. 2010, *A&A*, 515, A55
- Schneider, N., Bontemps, S., Simon, R., et al. 2011, *A&A*, 529, A1
- Tan, J. C., Beltrán, M. T., Caselli, P., et al. 2014, *Protostars and Planets VI*, 149
- van der Tak, F. F. S., Black, J. H., Schöier, F. L., Jansen, D. J., & van Dishoeck, E. F. 2007, *A&A*, 468, 627
- van der Tak, F. F. S., Chavarría, L., Herpin, F., et al. 2013, *A&A*, 554, A83
- van Dishoeck, E. F., Kristensen, L. E., Benz, A. O., et al. 2011, *PASP*, 123, 138
- Wirström, E. S., Charnley, S. B., Persson, C. M., et al. 2014, *ApJ*, 788, L32
- Zinnecker, H. & Yorke, H. W. 2007, *ARA&A*, 45, 481

Appendix A: Spectra of sources in the $\ell = 30^\circ$ region

All spectra are in T_A^* units and have been smoothed to a spectral resolution of 1 km s^{-1} , unless noted otherwise.




Fig. A.1: Original channel width (0.27 km/s).




Fig. A.3: Original channel width (0.27 km/s).




Fig. A.2: Original channel width (0.27 km/s).




Fig. A.4




Fig. A.5




Fig. A.7




Fig. A.6: Original channel width (0.27 km/s).




Fig. A.8




Fig. A.9




Fig. A.11: Original channel width (0.27 km/s).




Fig. A.10: Original channel width (0.27 km/s).




Fig. A.12: Original channel width (0.27 km/s).




Fig. A.13




Fig. A.15




Fig. A.14: Original channel width (0.27 km/s).




Fig. A.16




Fig. A.17: Original channel width (0.27 km/s).

Appendix B: Spectra of sources in the $\ell = 59^\circ$ region

All spectra are in T_{A}^* units and have been smoothed to a spectral resolution of 1 km s^{-1} , unless noted otherwise.




Fig. B.1




Fig. B.3




Fig. B.2




Fig. B.4




Fig. B.5




Fig. B.7




Fig. B.6




Fig. B.8




Fig. B.9




Fig. B.11




Fig. B.10




Fig. B.12




Fig. B.13




Fig. B.15




Fig. B.14




Fig. B.16




Fig. B.17




Fig. B.19




Fig. B.18




Fig. B.20




Fig. B.21




Fig. B.23




Fig. B.22




Fig. B.24




Fig. B.25




Fig. B.27




Fig. B.26




Fig. B.28




Fig. B.29




Fig. B.31




Fig. B.30




Fig. B.32: Original channel width (0.27 km/s).




Fig. B.33: Original channel width (0.27 km/s).




Fig. B.35: Original channel width (0.27 km/s).




Fig. B.34: Original channel width (0.27 km/s).