Regularity of extremal solutions of semilinear elliptic problems with non-convex nonlinearities on general domains

A. AGHAJANI

School of Mathematics, Iran University of Science and Technology, Narmak, Tehran, Iran. School of Mathematics, Institute for Research in Fundamental Sciences (IPM), P.O.Box: 19395-5746, Tehran, Iran.

Abstract

.

We consider the semilinear elliptic equation $-\Delta u = \lambda f(u)$ in a smooth bounded domain Ω of \mathbb{R}^n with Dirichielt boundary condition, where f is a \mathbb{C}^1 positive and nondeccreasing function in $[0, \infty)$ such that $\frac{f(t)}{t} \to \infty$ as $t \to \infty$. When Ω is an arbitrary domain and f is not necessarily convex, the boundedness of the extremal solution u^* is known only for $n = 2$, established by X. Cabré [\[5\]](#page-9-0). In this paper, we prove this for higher dimensions depending on the nonlinearity f . In particular, we prove that if

$$
\frac{1}{2} < \beta_- := \liminf_{t \to \infty} \frac{f'(t)F(t)}{f(t)^2} \le \beta_+ := \limsup_{t \to \infty} \frac{f'(t)F(t)}{f(t)^2} < \infty
$$

where $F(t) = \int_0^t f(s)ds$, then $u^* \in L^{\infty}(\Omega)$, for $n \leq 6$. Also, if $\beta_- = \beta_+ > \frac{1}{2}$ or $\frac{1}{2} < \beta_- \leq \beta_+ < \frac{7}{10}$, then $u^* \in L^{\infty}(\Omega)$, for $n \leq 9$. Moreover, if $\beta_- > \frac{1}{2}$ then $\tilde{u}^* \in H_0^1(\Omega)$ for $n \geq 2$.

Key words: Extremal solutions; Boundedness; Semilinear elliptic equations MSC(2010). 35K57, 35B65, 35J60

1. Introduction

In this article, we consider the semilinear Dirichlet problem

$$
\begin{cases}\n-\Delta u = \lambda f(u) & x \in \Omega, \\
u > 0 & x \in \Omega, \\
u = 0 & x \in \partial\Omega,\n\end{cases}
$$
\n(1.1)

March 6, 2022

Email address: aghajani@iust.ac.ir (A. AGHAJANI)

where $\Omega \subset \mathbb{R}^n$ is a smooth bounded domain, $n \geq 1$, $\lambda > 0$ is a real parameter, and the nonlinearity $f : [0, \infty] \to R$ satisfies

(*H*) f is C^1 , nondecreasing, $f(0) > 0$ and $\lim_{s \to \infty} \frac{f(s)}{s} = \infty$. By a weak solution solution of (1.1) we mean a nonnegative function $u \in L^1(\Omega)$ so that $f(u) \in L^1_\delta(\Omega) = L^1(\Omega, \delta(x)dx), \ \delta(x) = dist(x, \partial\Omega)$ and

$$
\int_{\Omega} (-\Delta \varphi) u = \int_{\Omega} \lambda f(u) \varphi
$$

holds for any $\varphi \in C^2(\overline{\Omega}), \varphi = 0$ on $\partial \Omega$ (see Brezis et al. [\[3\]](#page-8-0)).

It is well known ([\[3,](#page-8-0) [11,](#page-9-1) [13](#page-9-2)]) that there exists a finite positive extremal parameter λ^* such that for any $0 < \lambda < \lambda^*$, problem (1.1) has a minimal classical solution $u_\lambda \in \mathbb{R}^3$ $C^2(\overline{\Omega})$, while no solution exists, even in the weak sense for $\lambda \geq \lambda^*$. The function $\lambda \to u_\lambda$ is increasing and the increasing pointwise limit $u^*(x) = \lim_{\lambda \uparrow \lambda^*} u_\lambda(x)$ is a weak solution of (1.1) for $\lambda = \lambda^*$ which is called the extremal solution. If $\lambda < \lambda^*$ the solution u_{λ} is obtained by the implicit function theorem and is stable in the sense that the first Dirichlet eigenvalue of the linearized problem at u_λ , $-\Delta - \lambda f'(u_\lambda)$, is positive for all $\lambda \in (0, \lambda_*)$. That is,

$$
\int_{\Omega} |\nabla \varphi|^2 - \int_{\Omega} \lambda f(u)' \varphi^2 \ge 0, \quad \varphi \in H_0^1(\Omega). \tag{1.2}
$$

The regularity and properties of the extremal solutions have been studied extensively in the literature [2-12,15,19] and it is shown that it depends strongly on the dimension n , domain Ω and nonlinearity f .

When f is convex, Nedev in [\[16\]](#page-9-3) proved that $u^* \in L^{\infty}(\Omega)$ for $n = 2, 3$ in any domain Ω. When $2 \le n \le 4$ the best known result was established by Cabré [\[5](#page-9-0)] who showed that $u^* \in L^{\infty}(\Omega)$ for arbitrary nonlinearity f if in addition Ω is convex. Applying the main estimate used in the proof of the results of [\[5](#page-9-0)], Villegas [\[19](#page-9-4)] got the same replacing the condition that Ω is convex with f is convex. Cabré and Capella [\[8](#page-9-5)] proved that $u^* \in L^{\infty}(\Omega)$ if $n \leq 9$ and $\Omega = B_1$. Also, in [\[11\]](#page-9-1), Cabré and Ros-Oton showed that $u^* \in L^{\infty}(\Omega)$ if $n \leq 7$ and Ω is a convex domain of double revolution (see [\[11\]](#page-9-1) for the definition).

By imposing extra assumptions on the convex nonlinearity f satisfies (H) much more is known, see [\[9](#page-9-6)]. Let f is convex and define

$$
\tau_{-} := \liminf_{t \to \infty} \frac{f(t)f''(t)}{f'(t)^2} \le \tau_{+} := \limsup_{t \to \infty} \frac{f(t)f''(t)}{f'(t)^2}.
$$
\n(1.3)

Crandall and Rabinowitz [\[11\]](#page-9-1) proved $u^* \in L^{\infty}(\Omega)$ when $0 < \tau_- \leq \tau_+ < 2+\tau_-+\sqrt{\tau_-}$ and $n < 4 + 2\tau_- + 4\sqrt{\tau_-}$. This result was improved by Ye and Zhou in [\[20](#page-9-7)] and Sanchón in [\[14\]](#page-9-8) establishing that $u^* \in L^{\infty}(\Omega)$ when $\tau > 0$ and $n < 6 + 4\sqrt{\tau-1}$. In [\[14\]](#page-9-8) Sanchón proved that $u \in L^{\infty}$ whenever $\tau_{-} = \tau_{+} \geq 0$ and $n \leq 9$. Recently Cabré, Sanchón and Spruck [\[9](#page-9-6)] proved that if $\tau_+ < 1$ (without assuming $\tau_- > 0$) and $n < 2 + \frac{4}{\tau_+}$ then $u \in L^{\infty}$, and if $\tau_+ = 1$ and $n < 6$ then $u^* \in L^{\infty}$. These results improved by the author in [\[1\]](#page-8-1) as follows

$$
if \ 0 < \tau_{+} < \infty \ \ and \ \ n < \max\{2 + \frac{4}{\tau_{+}} + \frac{4}{\sqrt{\tau_{+}}}, \ 4 + \frac{2}{\tau_{+}} + \frac{4}{\sqrt{\tau_{+}}} \} \ then \ u^{*} \in L^{\infty}(\Omega).
$$

In particular, if $\tau_+ < \frac{2}{9-2\sqrt{14}} \approx 1.318$ and $n < 10$ then $u^* \in L^{\infty}(\Omega)$.

The case when f is not convex and Ω is arbitrary domain, is more challenging and there is nothing much in the literature about the boundedness of the extremal solution. Indeed, in this case, again the best result is due to Cabré $[5]$ who showed that $u^* \in L^{\infty}(\Omega)$ for arbitrary f and Ω in dimension $n = 2$.

In this work we consider problem (1.1) for the case when f is not necessarily convex and Ω is an arbitrary domain and prove the boundedness of the extremal solution in higher dimensions under some extra assumptions on f.

Let f satisfy (H) and define

$$
\beta_{-} := \liminf_{t \to \infty} \frac{f'(t)F(t)}{f(t)^{2}} \le \beta_{+} := \limsup_{t \to \infty} \frac{f'(t)F(t)}{f(t)^{2}},
$$
\n(1.4)

where $F(t) := \int_0^t f(s)ds$, for $t \ge 0$.

The main results of this paper are as follows.

Theorem 1.1. Let f (not necessarily convex) satisfy (H) with $\frac{1}{2} < \beta_- \leq \beta_+ < \infty$ and Ω an arbitrary bounded smooth domain. Let u^* be the extremal solution of problem (1.1) . Then $u^* \in L^{\infty}(\Omega)$ for

$$
n < 4 + 4 \left(\frac{2\beta_+ - 1}{2\beta_+} + \sqrt{\frac{2\beta_- - 1}{\beta_+}} \right). \tag{1.5}
$$

Furthermore, if $\beta_+ < 1$ then $u^* \in L^{\infty}(\Omega)$ for

$$
n < 6 + \frac{4}{2\beta_+ - 1} \left(1 - \beta_+ + \sqrt{\beta_+ (2\beta_- - 1)} \right). \tag{1.6}
$$

As consequences, by the assumption $\frac{1}{2} < \beta_- \leq \beta_+ < \infty$, we have: (a) If $n \leq 6$, then $u^* \in L^{\infty}(\Omega)$. (b) If $\beta_- = \beta_+$ or $\beta_+ < \frac{7}{10}$, then $u^* \in L^{\infty}(\Omega)$ for $n \leq 9$.

It is worth mentioning here that, for a convex nonlinearity f we always have $\beta_+ \geq$ $\beta_- \geq \frac{1}{2}$. Indeed in this case f' is a nondecreasing function, hence we have

$$
f'(t)F(t) = f'(t)\int_0^t f(s)ds \ge \int_0^t f'(s)f(s)ds = \frac{f(t)^2}{2} - \frac{f(0)^2}{2},
$$

now the fact that $f(t) \to \infty$ as $t \to \infty$ gives $\beta_- \geq \frac{1}{2}$. Also for general nonlinearities f (not necessarily convex) satisfy only (H) we always have $\beta_+ \geq \frac{1}{2}$. To see this, 3

by contradiction assume that $0 \leq \beta_+ < \frac{1}{2}$ and take $a \beta \in (\beta_+, \frac{1}{2})$. Then from the definition of β_+ there exists $T > 0$ such that $f'(t)F(t) \leq \beta f(t)^2$ for $t \geq T$, or equivalently, $\frac{d}{dt}(\frac{f(t)}{F(t)}$ $\frac{f(t)}{F(t)^{\beta}}$ \leq 0 for $t \geq T$. Thus, $\frac{f(t)}{F(t)^{\beta}} \leq C := \frac{f(T)}{F(T)^{\beta}}$ $\frac{J(I)}{F(T)^{\beta}}$ for $t \geq T$, and by integration we get $F(t) \leq (C_1 t + C_2)^{\frac{1}{1-\beta}}$ for all $t > T$ and some constants C_1, C_2 . But, from the superlinearity of f we have $\lim_{t\to\infty} \frac{F(t)}{t^2}$ $\frac{v(t)}{t^2} = \infty$, hence we must have $\frac{1}{1-\beta} > 2$ or equivalently $\beta > \frac{1}{2}$ which is a contradiction. Hence, we always have $\beta_+ \geq \frac{1}{2}$.

Example 1.1. Consider problem (1.1) in an arbitrary bounded smooth domain Ω with $f(u) = u^2 + 3u + 3\cos u + 4$. It is easy to see that f satisfies (H), but is not convex (even at infinity). Indeed, we have $f''(u) = 2 - 3 \cos u$, which is negative for all u such that $\cos u > \frac{2}{3}$ (so none of the previous results apply). However, by a simple computation we have $1 > \beta_- = \beta_+ = \frac{2}{3} > \frac{1}{2}$, hence by Theorem 1.1 we get $u^* \in L^{\infty}(\Omega)$ for $n \leq 15$.

Example 1.2. As an another example, consider problem (1.1) with $f(u) = e^u(3 +$ $2\cos u$) and arbitrary bounded smooth domain Ω . Then, f satisfies (H), but is not convex. Indeed, we have $f''(u) = e^u(3 - 4\sin u)$, so $\liminf_{u \to \infty} f''(u) = -\infty$. However, we have, after some simplification,

$$
\frac{f'(t)(F(t) + 4)}{f(t)^2} = \frac{(3 + 2\cos u - \sin u)(3 + \sin u + \cos u)}{(3 + 2\cos u)^2} := \beta(u),
$$

which is a periodic function with period 2π , hence (as computed by Mathematica),

$$
\beta_- = \min_{[0,2\pi]} \beta(u) \approx 0.786244 \quad \text{and} \quad \beta_+ = \max_{[0,2\pi]} \beta(u) \approx 2.08846,
$$

where we used also that $\lim_{t\to\infty} \frac{4f'(t)}{f(t)^2}$ $\frac{f(t)}{f(t)^2} = 0$. Now, using Theorem (1.1) we get $u^* \in L^{\infty}(\Omega)$ for $n \leq 9$.

Now consider the well-known convex nonlinearities $f(t) = e^t$ or $(1+t)^p$, $p > 1$. When $f(t) = e^t$ we have $\beta_- = \beta_+ = 1$ then from Theorem 1.1 we get $u^* \in L^{\infty}(\Omega)$ for $n \leq 9$. Also, for $f(t) = (1+t)^p$, $p > 1$ we have $\beta_- = \beta_+ = \frac{p}{p+1} < 1$, hence from Theorem 1.1 we get

$$
u \in L^{\infty}(\Omega) \quad for \quad n < 2\left(1 + \frac{2p}{p-1} + 2\sqrt{\frac{p}{p-1}}\right).
$$

The above results are well-known in the literature [\[11](#page-9-1), [20](#page-9-7), [14](#page-9-8)].

Also, notice that if f is convex and $\tau_{-} > 0$ then we must have $\beta_{-} > \frac{1}{2}$. Indeed, it is easy to see that the condition $\tau_$ > 0 yields that for every $0 < \tau < \tau_$, there exists $T = T(\tau) > 0$ such that the function $\frac{f'(t)}{f(t) \tau}$ $\frac{f(t)}{f(t)^{\tau}}$ is increasing in $[T, \infty)$, hence

$$
f'(t)F(t) = f(t)^{\tau} \frac{f'(t)}{f(t)^{\tau}} \int_0^t f(s)ds \ge f(t)^{\tau} \int_T^t \frac{f'(s)}{f(s)^{\tau}} f(s)ds = \frac{f(t)^{2-\tau}}{2-\tau} - \frac{f(T)^{2-\tau}}{2-\tau},
$$

now the facts that $f(t) \to \infty$ as $t \to \infty$ and $\tau > \tau_{-}$ was arbitrary give $\beta_{-} \ge \frac{1}{2-\tau_{-}} >$ $\frac{1}{2}$.

To get the regularity of the extremal solution in low dimensions or proving that it is in the energy class (i.e., $u^* \in H_0^1(\Omega)$) we can weaken the assumptions as follows.

Theorem 1.2. Let f (not necessarily convex) satisfy (H) and Ω an arbitrary bounded smooth domain in R^n . Let u^* be the extremal solution of problem (1.1). Then (i) if for some $\epsilon > 0$ there exist $t_0 > 0$ such that we have

$$
\frac{f'(t)F(t)}{f(t)^2} \ge \frac{1}{2} + \frac{\epsilon t}{f(t)}, \quad t > t_0
$$
\n(1.7)

then $u^* \in H_0^1(\Omega)$. In particular this is true if $\beta_- > \frac{1}{2}$. (ii) If for some $0 < \delta \leq 1$, $\lim_{t \to \infty} \frac{f(t)}{t^{2-\delta}}$ $\frac{f(t)}{t^{2-\delta}} = \infty$, and there exist $t_0 > 0$ such that

$$
\frac{f'(t)F(t)}{f(t)^2} \ge \frac{1}{2} + \frac{1}{t^{2-\delta}}, \quad t > t_0
$$
\n(1.8)

then $u^* \in L^{\infty}(\Omega)$ for $n < 5$. In particular this is true if $\beta_{-} > \frac{1}{2}$.

2. Preliminaries and Auxiliary Results

To prove the main results we need the following simple technical lemma based on inequality (1.2), which is used frequently in the literature, for example [\[11,](#page-9-1) [9](#page-9-6), [16](#page-9-3), [20\]](#page-9-7). It is also proved in [\[1\]](#page-8-1) for the general semilinear elliptic equation $-Lu = \lambda f(u)$ with zero Dirichlet boundary condition, but for the convenience of the reader we sketch a proof here for the case $L = \Delta$. Also,

Lemma 2.1. Let u_{λ} be the minimal solution of (1.1) and $g : [0, \infty] \rightarrow [0, \infty]$ be a C^1 function with $g(0) = 0$ and satisfy

$$
H(t) := g(t)^{2} f'(t) - G(t) f(t) \ge 0, \text{ for t sufficiently large}, \qquad (2.1)
$$

where $G(t) := \int_0^t g'(s)^2 ds$. Then $||H(u_\lambda)||_{L^1(\Omega)} \leq C$, where C is a constant independent of λ .

PROOF. Let $u_{\lambda} \in C^2(\overline{\Omega})$ be the minimal classical solution of (1.1) where $0 < \lambda < \lambda^*$, and take $\varphi = g(u_\lambda)$ in the semi-stability condition (1.2). Then we get

$$
\int_{\Omega} g'(u_{\lambda})^2 |\nabla u_{\lambda}|^2 dx - \int_{\Omega} \lambda f'(u_{\lambda}) g(u_{\lambda})^2 dx \ge 0.
$$
 (2.2)

By using the Green's formula one can show that

$$
\int_{\Omega} g'(u_{\lambda})^2 |\nabla u_{\lambda}|^2 dx = \int_{\Omega} \lambda G(u_{\lambda}) f(u_{\lambda}) dx.
$$
\n(2.3)

Using (2.3) in (2.2) we obtain

$$
\int_{\Omega} H(u_{\lambda})dx \le 0.
$$
\n(2.4)

Now from (2.1) there exists $t_0 > 0$ so that $H(t) \geq 0$ for $t \geq t_0$, thus using (2.4) we obtain

$$
\int_{\Omega} |H(u_{\lambda})| dx = \int_{u_{\lambda} \le t_0} |H(u_{\lambda})| dx + \int_{u_{\lambda} \ge t_0} H(u_{\lambda}) dx
$$

$$
\le \int_{u_{\lambda} \le t_0} (|H(u_{\lambda})| - H(u_{\lambda})) dx \le C_0 |\Omega|,
$$

where $|\Omega|$ denotes the Lebesgue measure of Ω and $C_0 := \sup_{t \in [0,t_0]} (|H(t)| - H(t)).$ Now, since C_0 is independent of λ we get the desired result.

The following consequence of the above lemma is essential in the proof of the main results.

Proposition 2.1. Let u_{λ} be the minimal solution of (1.1) and $\xi : [0, \infty] \rightarrow [0, \infty]$ be a C^1 function such that for some $t_0 > 0$ we have $\xi(t) \leq \frac{f'(t)}{f(t)}$ $\frac{f'(t)}{f(t)}, \xi'(t) + \xi(t)^2 \geq 0$ for $t \ge t_0$, and $\frac{E(t)}{f(t)} \to \infty$ as $t \to \infty$ where

$$
E(t) := f(t) \left(\frac{f'(t)}{f(t)} - \xi(t) \right) e^{2 \int_{t_0}^t (\xi(s) + \sqrt{\xi'(s) + \xi(s)^2}) ds}.
$$
 (2.5)

Then $||E(u_\lambda)||_{L^1(\Omega)} \leq C$, where C is a constant independent of λ .

PROOF. Let $g: [0, \infty] \to [0, \infty]$ be a C^1 function with $g(0) = 0$ and

$$
g(t) = e^{\int_{t_0}^t (\xi(s) + \sqrt{\xi'(s) + \xi(s)^2}) ds}
$$
, for $t \ge t_0$.

Also, let $G(t) = \int_0^t g'(s)^2 ds$ as in lemma 2.1. Then using the equality

$$
g'(t) = (\xi(t) + \sqrt{\xi'(t) + \xi(t)^2})g(t)
$$
 for $t \ge t_0$,

we compute

$$
\frac{d}{dt}\left(\xi(t)g(t)^2 - G(t)\right) = \xi'(t)g(t)^2 + 2\xi(t)g(t)g'(t) - g'(t)^2
$$
\n
$$
= g(t)^2\left(\xi'(t) + 2\xi(t)^2 + 2\xi(t)\sqrt{\xi'(t) + \xi(t)^2}\right) - g'(t)^2
$$
\n
$$
= g(t)^2\left(\xi(s) + \sqrt{\xi'(t) + \xi(t)^2}\right)^2 - g'(t)^2
$$
\n
$$
= g'(t)^2 - g'(t)^2 = 0, \text{ for } t \ge t_0,
$$

implies that

$$
G(t) = \xi(t)g(t)^{2} + C_{0}, \text{ where } C_{0} := G(t_{0}) - \xi(t_{0}).
$$
\n(2.6)

Now using (2.6), for $t \ge t_0$ we have

$$
H(t) := g(t)^2 f'(t) - G(t)f(t) = g(t)^2 f'(t) - g(t)2\xi(t)f(t) - C_0f(t) = E(t) - C_0f(t),
$$

which is positive for large $t \geq t_0$ (by the assumption), hence by Lemma 2.1 we get $||H(u_\lambda)||_{L^1(\Omega)} \leq C_1$, where C_1 is a constant independent of λ . However, again by the assumption we have $0 < E(t) < 2H(t)$ for large t that also gives $||E(u_\lambda)||_{L^1(\Omega)} \leq C_2$, where C_2 is a constant independent of λ , which is the desired result. \blacksquare

To prove Theorem 1.2 in the next section, we also need the following rather standard result. For a simple proof see [\[1\]](#page-8-1).

Proposition 2.2. Let f satisfy (H) and u_{λ} be the minimal solution of problem (1.1). If there exists a positive constant C independent of λ such that

$$
||u_{\lambda}||_{L^{1}(\Omega)} \leq C \quad and \quad ||\frac{\tilde{f}(u_{\lambda})^{\alpha}}{u_{\lambda}^{\sigma}}||_{L^{1}(\Omega)} \leq C, \quad for \; some \; 0 \leq \sigma \leq \alpha, \tag{2.7}
$$

where $\tilde{f}(u) = f(u) - f(0)$ and $\alpha > 1$, then

$$
||u_{\lambda}||_{L^{\infty}(\Omega)} \leq \tilde{C} \quad \text{for} \quad n < 2\alpha,
$$
\n^(2.8)

where \tilde{C} is a positive constant independent of λ .

3. Proof of the main results

Proof of Theorem 1.1

By the assumptions we have $\frac{1}{2} < \beta_- \leq \beta_+ < \infty$. Take $\frac{1}{2} < \beta_1 < \beta_2 < \beta_-$ and $\beta_3 \in (\beta_+, \infty)$, then by the definition of β_-, β_+ (see (1.4)) there exists a $t_0 > 0$ such that

$$
\beta_1 < \beta_2 < \frac{f'(t)F(t)}{f(t)^2} < \beta_3
$$
, for $t \ge t_0$. (3.1)

Now let $\xi : [0, \infty] \to [0, \infty]$ be a C^1 function such that $\xi(t) = \beta_1 \frac{f(t)}{F(t)}$ $\frac{f(t)}{F(t)}$ for $t \geq t_0$, where $F(t) := \int_0^t f(s)ds$. Then from (3.1) we have $\xi(t) \leq \frac{f'(t)}{f(t)}$ $\frac{f(t)}{f(t)}$ and

$$
\xi'(t) + \xi(t)^2 = \beta_1 \left(\frac{f'(t)}{F(t)} - (1 - \beta_1) \frac{f(t)^2}{F(t)^2} \right) \ge (2\beta_1 - 1) \frac{f'(t)}{F(t)} \ge \frac{2\beta_1 - 1}{\beta_3} \frac{f'(t)^2}{f(t)^2},
$$
(3.2)

for $t \geq t_0$. Notice that, in (3.2) we used the fact that $\beta_1 < 1$ (because we always have $\beta_- \leq 1$). Also, from (3.1) we have

$$
\frac{f'(t)}{f(t)} - \xi(t) \ge (\beta_2 - \beta_1) \frac{f(t)}{F(t)}, \text{ for } t \ge t_0.
$$

Now let the function $E(t)$ be given as in (2.5) in Proposition 2.1. By the later inequality, (3.1), (3.2) and the fact that $\int_{t_0}^t \xi(s)ds = \beta_1(\ln F(t) - \ln F(t_0))$, we have

$$
E(t) = f(t) \left(\frac{f'(t)}{f(t)} - \xi(t) \right) e^{2 \int_{t_0}^t (\xi(s) + \sqrt{\xi'(s) + \xi(s)^2}) ds} \ge C F(t)^{2\beta_1 - 1} f(t)^{2 + 2\sqrt{\frac{2\beta_1 - 1}{\beta_3}}},
$$
\n(3.3)

where C is a positive constant depends only on f . Now, writing the last inequality in (3.1) as $\frac{f'(t)}{f(t)} < \beta_3 \frac{f(t)}{F(t)}$ $\frac{f(t)}{F(t)}$ for $t_0 > 0$, then integration from t_0 to t gives

$$
F(t) \ge Cf(t)^{\frac{1}{\beta_3}} \quad \text{for} \quad t \ge t_0. \tag{3.4}
$$

Using (3.4) in (3.3) we arrive at

$$
E(t) \ge f(t)^{\gamma}, \text{ where } \gamma := 2 + \frac{2\beta_1 - 1}{\beta_3} + 2\sqrt{\frac{2\beta_1 - 1}{\beta_3}}, \text{ for } t \ge t_0.
$$
 (3.5)

And, since $\frac{E(t)}{f(t)} \to \infty$ t $\to \infty$, from Proposition 2.1 we get $||E(u_\lambda)||_{L^1(\Omega)} \leq C$ and then from (3.5) $|| f(u_\lambda)^\gamma ||_{L^1(\Omega)} \leq C$, where C is a constant independent of λ . Now the standard elliptic regularity theory gives $u^* \in L^{\infty}(\Omega)$ for $n < 2\gamma$, and since β_1 and β_3 were arbitrary in the intervals $(\frac{1}{2}, \beta)$ and (β_+, ∞) , respectively, thus $u^* \in L^{\infty}(\Omega)$ for

$$
n < 4 + 4\left(\frac{2\beta_+ - 1}{2\beta_+} + \sqrt{\frac{2\beta_- - 1}{\beta_+}}\right) := \gamma_1,\tag{3.6}
$$

that proves the first part.

Now assume that $\beta_+ < 1$, then we can also assume that $\beta_3 < 1$. Now, from (3.4) we have $f(t)F(t)^{-\beta_3} \leq C_1$, for $t \geq t_0$, and integration from t_0 to t gives $F(t) \leq C_2 t^{\frac{1}{1-\beta_3}}$, $t \geq t_1$, for some $t_1 \geq t_0$. This together (3.4) implies that $f(t) \leq C_3 t^{\frac{\beta_3}{1-\beta_3}}$, for $t \geq t_1$, that also yields, for some $t_2 \geq t_1$,

$$
f(t)^{\gamma_1} \ge C_4 \frac{f(t)^{\gamma_2}}{t^{\gamma_2}}, \text{ for } t \ge t_2, \gamma_2 := \frac{\beta_3}{2\beta_3 - 1} \gamma_1,
$$

where γ_1 is given in (3.6). Hence, $\left| \frac{\tilde{f}(u_\lambda)^{\gamma_2}}{u_\lambda^{\gamma_2}} \right|$ $\frac{u_{\lambda})^{\gamma_2}}{u_{\lambda}^{\gamma_2}}||_{L^1(\Omega)} \leq C$, where $\tilde{f}(t) = f(t) - f(0)$ and C is a constant independent of λ . Now, from Proposition 2.2 gives $u^* \in L^{\infty}(\Omega)$ for $n < 2\gamma_2$, that proves the second part.

To prove part (a), note that in the case $\beta_+ \geq 1$, it is easy to see that the right hand side of (1.5) is larger than 6 and when β_{+} < 1 we can use (1.6).

To show part (b), note that (from (1.6)) if $6 + \frac{4(1-\beta_+)}{2\beta_+-1} > 9$, which is equivalent to $\beta_+ < \frac{7}{10}$, then $u^* \in L^\infty(\Omega)$ for $n < 9$.

Now assume that $\beta_-=\beta_+$ then we have $\beta_+\leq 1$ (as we always have $\beta_-\leq 1$). Also, from the later part we can consider only the case $\beta_+ \geq \frac{7}{10}$. If $\beta_- = \beta_+ = 1$ then from (1.5) we have $u^* \in L^{\infty}(\Omega)$ for $n < 10$. Also, if $\frac{7}{10} \leq \beta_+ < 1$ we can use (1.6). We need to show that the right hand side of (1.6) is larger than 9. In the case $\frac{7}{10} \leq \beta_+ < 1$ this is equivalent to $68\beta_+^2 + 49 < 124\beta_+$, which obviously holds for $\beta_+ \in [\frac{7}{10}, 1)$.

Proof of Theorem 1.2

Let $\xi : [0, \infty] \to [0, \infty]$ be a C^1 function such that $\xi(t) = \frac{1}{2}$ $f(t)$ $\frac{f(t)}{F(t)}$ for $t \ge t_0$, where t_0 given as in (1.7) . Then from (1.7) we have

$$
\frac{f'(t)}{f(t)} - \xi(t) = \frac{f(t)}{F(t)} \left(\frac{f'(t)F(t)}{f(t)^2} - \frac{1}{2} \right) \ge \epsilon \frac{t}{F(t)}, \text{ for } t \ge t_0,
$$
\n(3.7)

and

$$
\xi'(t) + \xi(t)^2 = \frac{1}{2} \left(\frac{f'(t)F(t)}{f(t)^2} - \frac{1}{2} \right) \frac{f(t)^2}{F(t)^2} > 0 \text{ for } t \ge t_0.
$$
 (3.8)

From (3.7) and (3.8) we get

$$
E(t) = f(t) \Big(\frac{f'(t)}{f(t)} - \xi(t) \Big) e^{2 \int_{t_0}^t (\xi(s) + \sqrt{\xi'(s) + \xi(s)^2}) ds} \ge Ct f(t), \text{ for } t \ge t_0 \tag{3.9}
$$

where C is a positive constant depends only on f . Thus, from Proposition 2.1 we get $u_{\lambda} f(u_{\lambda}) \in L^{1}(\Omega)$ by a constant independent of λ . Multiplying (1.1) by u_{λ} we get

$$
\int_{\Omega} |\nabla u_{\lambda}|^2 dx = \lambda \int_{\Omega} u_{\lambda} f(u_{\lambda}) dx \leq \lambda^* C,
$$

with C independent of λ , which leads to $\int_{\Omega} |\nabla u^*|^2 dx \leq \lambda^* C$. Hence, $u^* \in H_0^1(\Omega)$, that proves part (i).

Let $\xi(t)$ be as defined above. Then, using (1.8) and similar to the proof of part (i) we can show that $E(t) \geq C \frac{f(t)^2}{t^{2-\delta}}$ $\frac{f(t)^2}{t^{2-\delta}}$ for $t \ge t_0$, hence by the assumption $\frac{E(t)}{f(t)} \to \infty$ as $t \to \infty$. Now from Proposition 2.1 we get

$$
\frac{\tilde{f}(u_{\lambda})^2}{u_{\lambda}^{2-\delta}} \in L^{\infty}(\Omega). \tag{3.10}
$$

Now we proceed similar to the proof of Theorem 1.1 in [\[19](#page-9-4)]. From Proposition 2.3 in [\[19](#page-9-4)], there exits a universal constant C_1 independent of f, Ω and λ such that

$$
||u_{\lambda}||_{L^{\infty}(\Omega)} \le C_1 ||\nabla u_{\lambda}||_{L^{4}(\Omega)}.
$$
\n(3.11)

Also, from the continuous inclusion $W^{2,2}(\Omega) \subset W^{1,4}(\Omega)$ and elliptic regularity theory (see [\[2](#page-8-2)]), there exits a constant $C_2 = C_2(\Omega)$ such that

$$
||u_{\lambda}||_{W^{1,4}(\Omega)} \le C_2 ||u_{\lambda}||_{W^{2,2}(\Omega)} \quad \text{and} \quad ||u_{\lambda}||_{W^{2,2}(\Omega)} \le C_2 ||\lambda f(u_{\lambda})||_{L^2(\Omega)}. \tag{3.12}
$$

Now from (3.10) , (3.11) and (3.12) we have (in the following inequalities various constants will be denoted by C)

$$
||u_{\lambda}||_{L^{\infty}(\Omega)} \leq C||\nabla u_{\lambda}||_{L^{4}(\Omega)} \leq C||u_{\lambda}||_{W^{1,4}(\Omega)} \leq C||u_{\lambda}||_{W^{2,2}(\Omega)}
$$

\n
$$
\leq C||\lambda f(u_{\lambda})||_{L^{2}(\Omega)} \leq C\lambda^{*} \Big(\int_{u_{\lambda} \leq 1} f(u_{\lambda})^{2} dx + \int_{u_{\lambda} > 1} f(u_{\lambda})^{2} dx\Big)^{\frac{1}{2}}
$$

\n
$$
\leq C \Big(f(1)^{2}|\Omega| + \int_{u_{\lambda} > 1} \frac{f(u_{\lambda})^{2}}{u_{\lambda}^{2-\delta}} u_{\lambda}^{2-\delta} dx\Big)^{\frac{1}{2}} \leq C \Big(f(1)^{2}|\Omega| + \tilde{C}||u_{\lambda}||_{L^{\infty}(\Omega)}^{2-\delta}\Big)^{\frac{1}{2}},
$$

for every $\lambda \in (0, \lambda_*)$. Hence, we must have

$$
||u_{\lambda}||_{L^{\infty}(\Omega)}^2 \le A + B||u_{\lambda}||_{L^{\infty}(\Omega)}^{2-\delta}, \text{ for every } \lambda \in (0, \lambda_*),
$$

where A and B are positive constants independent of λ . This implies that $||u_\lambda||_{L^\infty(\Omega)} \le$ C with C independent of λ , now letting $\lambda \to \lambda^*$ gives $u^* \in L^{\infty}(\Omega)$.

4. Acknowledgement

This research was in part supported by a grant from IPM (No. 93340123).

References

- [1] A. Aghajani, New a priori estimates for semistable solutions of semilinear elliptic equations, [http://arxiv.org/abs/1508.04723.](http://arxiv.org/abs/1508.04723)
- [2] S. Agmon, A. Douglis, L. Nirenberg, Estimates near the boundary for solutions of elliptic partial differential equations satisfying general boundary conditions, I, Comm. Pure Appl. Math. 12 (1959) 623727.
- [3] H. Brezis, T. Cazenave, Y. Martel and A. Ramiandrisoa, Blow-up for $u_t - \Delta u = g(u)$ revisited, Adv. Differental Equation 1 (1996) 73-90.
- [4] H. Brezis and J. L. Vázquez, Blow-up solutions of some nonlinear elliptic problems, Rev. Mat. Univ. Complut. Madrid 10, 2 (1997) 443-469.
- [5] X. Cabré, Regularity of minimizers of semilinear elliptic problems up to dimension 4, Comm. Pure Appl. Math 63 (2010) 1362-1380.
- [6] X. Cabré, A. Capella, Regularity of radial minimizers and extremal solutions of semi-linear elliptic equations, J. Funct. Anal. 238, 2 (2006) 709-733.
- [7] X. Cabré, A. Capella, and M. Sanchón, Regularity of radial minimizers of reaction equa- tions involving the p-Laplacian, Calc. Var. Partial Differential Equations 34, 4 (2009) 475-494.
- [8] X. Cabré, X. Ros-Oton, Regularity of stable solutions up to dimension 7 in domains of double revolution, Comm. Partial Differential Equations 38, 1 (2013) 135154.
- [9] X. Cabré, M. Sanchón and J. Spruck, A priori estimates for semistable solutions of semilinear elliptic equations, to appear in Discrete Contin. Dyn. Syst., Series A, 36, n. 2. Preprint [http://arxiv.org/abs/1407.0243.](http://arxiv.org/abs/1407.0243)
- [10] A. Capella, J. Dvila, L. Dupaigne, and Y. Sire, Regularity of radial extremal solutions for some nonlocal semilinear elliptic equations, , Comm. PDE 36, 2 (2011) 1353-1384.
- [11] M.G. Crandall and P.H. Rabinowitz, Some continuation and variational methods for positive solutions of nonlinear elliptic eigenvalue problems, Arch. Ration. Mech. Anal. 58 (1975) 207-218.
- [12] J. Dvila, L. Dupaigne, and M. Montenegro, The extremal solution of a boundary reaction problem, Commun. Pure Appl. Anal. 7, 4 (2008) 795-817.
- [13] L. Dupaigne, Stable solutions of elliptic partial differential equations. Chapman & Hall/CRC Monographs and Surveys in Pure and Applied Mathematics, 2011.
- [14] M. Sanchón, Boundedness of the extremal solution of some p-Laplacian problems, Nonlinear Anal. 67 (2007) 281-294.
- [15] M. Sanchón, Regularity of the extremal solution of some nonlinear elliptic problems involving the p-Laplacian, Potential Anal. 27 (2007) 217-224.
- [16] G. Nedev, Regularity of the extremal solution of semilinear elliptic equations, C. R. Acad. Sci. Paris Sér. I Math. 330 (2000) 997-1002
- [17] J. Serrin, Local behavior of solutions of quasi-linear equations, Acta Math. 111 (1964) 247- 302.
- [18] N.S. Trudinger, Linear elliptic operators with measurable coefficients, Ann. Scuola Norm. Sup. Pisa (3) 27 (1973) 265308.
- [19] S. Villegas, Boundedness of extremal solutions in dimension 4, Adv. Math. 235 (2013) 126-133.
- [20] D. Ye, F. Zhou, Boundedness of the extremal solution for semilinear elliptic problems, Commun. Contemp. Math. 4 (2002) 547-558.