

Most Likely Transformations

Torsten Hothorn
Universität Zürich

Lisa Möst
Universität München

Peter Bühlmann
ETH Zürich

Abstract

We propose and study properties of maximum likelihood estimators in the class of conditional transformation models. Based on a suitable explicit parameterisation of the unconditional or conditional transformation function, we establish a cascade of increasingly complex transformation models that can be estimated, compared and analysed in the maximum likelihood framework. Models for the unconditional or conditional distribution function of any univariate response variable can be set-up and estimated in the same theoretical and computational framework simply by choosing an appropriate transformation function and parameterisation thereof. The ability to evaluate the distribution function directly allows us to estimate models based on the exact likelihood, especially in the presence of random censoring or truncation. For discrete and continuous responses, we establish the asymptotic normality of the proposed estimators. A reference software implementation of maximum likelihood-based estimation for conditional transformation models that allows the same flexibility as the theory developed here was employed to illustrate the wide range of possible applications.

Keywords: transformation model, distribution regression, conditional distribution function, conditional quantile function, censoring, truncation.

1. Introduction

In a broad sense, we can understand all statistical models as models of distributions or certain characteristics thereof, especially the mean. All distributions \mathbb{P}_Y for at least ordered responses Y can be characterised by their distribution, quantile, density, odds, hazard or cumulative hazard functions. In a fully parametric setting, all these functions have been specified up to unknown parameters, and the ease of interpretation can guide us in looking at the appropriate function. In the semi- and non-parametric contexts, however, the question arises how we can obtain an estimate of one of these functions without assuming much about their shape. For the direct estimation of distribution functions, we deal with monotonic functions in the unit interval, whereas for densities, we need to make sure that the estimator integrates to one. The hazard function comes with a positivity constraint, and monotonicity is required for the positive cumulative hazard function. These computationally inconvenient restrictions disappear completely only when the log-hazard function is estimated, and this explains the plethora of research papers following this path. However, the lack of any structure in the

A shortened version of this technical report was accepted for publication in the *Scandinavian Journal of Statistics* 2017-06-19.

log-hazard function comes at a price. A too-erratic behaviour of estimates of the log-hazard function has to be prevented by some smoothness constraint; this makes classical likelihood inference impossible. The novel characterisation and subsequent estimation of distributions via their transformation function in a broad class of transformation models that are developed in this paper can be interpreted as a compromise between structure (monotonicity) and ease of parameterisation, estimation and inference. This transformation approach to modelling and estimation allows standard likelihood inference in a large class of models that have so far commonly been dealt with by other inference procedures.

Since the introduction of transformation models based on non-linear transformations of some response variable by [Box and Cox \(1964\)](#), this attractive class of models has received much interest. In regression problems, transformation models can be understood as models for the conditional distribution function and are sometimes referred to as “distribution regression”, in contrast to their “quantile regression” counterpart ([Chernozhukov *et al.* 2013](#)). Traditionally, the models were actively studied and applied in the analysis of ordered categorical or censored responses. Recently, transformation models for the direct estimation of conditional distribution functions for arbitrary responses received interest in the context of counterfactual distributions ([Chernozhukov *et al.* 2013](#)), probabilistic forecasting ([Gneiting and Katzfuss 2014](#)), distribution and quantile regression ([Leorato and Peracchi 2015](#); [Rothe and Wied 2013](#)), probabilistic index models ([Thas *et al.* 2012](#)) and conditional transformation models ([Hothorn *et al.* 2014](#)). The core idea of any transformation model is the application of a strictly monotonic transformation function h for the reformulation of an unknown distribution function $\mathbb{P}(Y \leq y)$ as $\mathbb{P}(h(Y) \leq h(y))$, where the unknown transformation function h is estimated from the data. Transformation models have received attention especially in situations where the likelihood contains terms involving the conditional distribution function $\mathbb{P}(Y \leq y \mid \mathbf{X} = \mathbf{x}) = F_Z(h(y \mid \mathbf{x}))$ with inverse link function F_Z , most importantly for censored, truncated and ordered categorical responses. For partially linear transformation models with transformation function $h(y \mid \mathbf{x}) = h_Y(y) + h_{\mathbf{x}}(\mathbf{x})$, much emphasis has been given to estimation procedures treating the baseline transformation h_Y (*e.g.* the log-cumulative baseline hazard function in the Cox model) as a high-dimensional nuisance parameter. Prominent members of these estimation procedures are the partial likelihood estimator ([Cox 1975](#)) and approaches influenced by the estimation equations introduced by [Cheng *et al.* \(1995\)](#) and [Chen *et al.* \(2002\)](#). Once an estimate for the shift $h_{\mathbf{x}}$ is obtained, the baseline transformation h_Y is then typically estimated by the non-parametric maximum likelihood estimator (see, for example, [Cheng *et al.* 1997](#)). An overview of the extensive literature on the simultaneous non-parametric maximum likelihood estimation of h_Y and $h_{\mathbf{x}}$, *i.e.* estimation procedures not requiring an explicit parameterisation of h_Y , for censored continuous responses is given in [Zeng and Lin \(2007\)](#).

An explicit parameterisation of h_Y is common in models of ordinal responses ([Tutz 2012](#)). For survival times, [Kooperberg *et al.* \(1995\)](#) and [Kooperberg and Clarkson \(1997\)](#) introduced a cubic spline parameterisation of the log-conditional hazard function with the possibility of response-varying effects and estimated the corresponding models by maximum likelihood. [Crowther and Lambert \(2014\)](#) followed up on this suggestion and used restricted cubic splines. Many authors studied penalised likelihood approaches for spline approximations of the baseline hazard function in a Cox model, *e.g.* [Joly *et al.* \(1998\)](#); [Cai and Betensky \(2003\)](#); [Sabanés Bové and Held \(2013\)](#) or [Ma *et al.* \(2014\)](#). Less frequently, the transformation function h_Y was modelled directly. [Mallick and Walker \(2003\)](#); [Chang *et al.* \(2005\)](#)

and McLain and Ghosh (2013) used Bernstein polynomials for h_Y , and Royston and Parmar (2002) proposed a maximum likelihood approach using cubic splines for modelling h_Y and also time-varying effects. The connection between these different transformation models is difficult to see because most authors present their models in the relatively narrow contexts of survival or ordinal data. The lack of a general understanding of transformation models made the development of novel approaches in this model class burdensome. Hothorn *et al.* (2014) decoupled the parameterisation of the conditional transformation function $h(y | \mathbf{x})$ from the estimation procedure, and showed that many interesting and novel models can be understood as transformation models. The boosting-based optimisation of proper scoring rules, however, was only developed for uncensored and right-censored (Möst and Hothorn 2015) observations in the absence of truncation and requires the numerical approximation of the true target function. In a similar spirit, Chernozhukov *et al.* (2013) applied the connection $\mathbb{P}(Y \leq y | \mathbf{X} = \mathbf{x}) = \mathbb{E}(\mathbb{1}(Y \leq y) | \mathbf{X} = \mathbf{x})$ for estimation in the response-varying effects transformation model $\mathbb{P}(Y \leq y | \mathbf{X} = \mathbf{x}) = F_Z(h_Y(y) - \mathbf{x}^\top \boldsymbol{\beta}(y))$; this approach can be traced back to Foresi and Peracchi (1995).

A drawback of all but the simplest transformation models is the lack of a likelihood estimation procedure. Furthermore, although important connections to other models have been known for some time (Doksum and Gasko 1990), it is often not easy to see how broad and powerful the class of transformation models actually is. We address these issues and embed the estimation of unconditional and conditional distribution functions of arbitrary univariate random variables under all forms of random censoring and truncation into a common theoretical and computational likelihood-based framework. In a nutshell, we show in Section 2 that all distributions can be generated by a strictly monotonic transformation of some absolute continuous random variable. The likelihood function of the transformed variable can then be characterised by this transformation function. The parameters of appropriate parameterisations of the transformation function, and thus the parameters of the conditional distribution function in which we are interested, can then be estimated by maximum likelihood under simple linear constraints that allow classical asymptotic likelihood inference, as will be shown in Section 3. Many classical and contemporary models are introduced as special cases of this framework. In particular, all transformation models sketched in this introduction can be understood and estimated in this novel likelihood-based framework. Extensions of classical and contemporary transformation models as well as some novel models are derived from our unified theoretical framework of transformation functions in Section 4, and their empirical performance is illustrated and evaluated in Section 5.

2. The Likelihood of Transformations

Let $(\Omega, \mathfrak{A}, \mathbb{P})$ denote a probability space and (Ξ, \mathfrak{C}) a measurable space with at least ordered sample space Ξ . We are interested in inference about the distribution \mathbb{P}_Y of a random variable Y , *i.e.* the probability space $(\Xi, \mathfrak{C}, \mathbb{P}_Y)$ defined by the $\mathfrak{A} - \mathfrak{C}$ measurable function $Y : \Omega \rightarrow \Xi$. For the sake of notational simplicity, we present our results for the unconditional and ordered case first; regression models and unordered responses are discussed in Section 4.2. The distribution $\mathbb{P}_Y = f_Y \odot \mu$ is dominated by some measure μ and characterised by its density function f_Y , distribution function $F_Y(y) = \mathbb{P}_Y(\{\xi \in \Xi | \xi \leq y\})$, quantile function $F_Y^{-1}(p) = \inf\{y \in \Xi | F_Y(y) \geq p\}$, odds function $O_Y(y) = F_Y(y)/(1 - F_Y(y))$, hazard function $\lambda_Y(y) = f_Y(y)/(1 - F_Y(y))$ or cumulative hazard function $\Lambda_Y(y) = -\log(1 - F_Y(y))$. For

notational convenience, we assume strict monotonicity of F_Y , *i.e.* $F_Y(y_1) < F_Y(y_2) \forall y_1 < y_2 \in \Xi$. Our aim is to obtain an estimate $\hat{F}_{Y,N}$ of the distribution function F_Y from a random sample $Y_1, \dots, Y_N \stackrel{\text{iid}}{\sim} \mathbb{P}_Y$. In the following, we will show that one can always write this potentially complex distribution function F_Y as the composition of a much simpler and *a priori* specified distribution function F_Z and a strictly monotonic transformation function h . The task of estimating F_Y is then reduced to obtaining an estimate \hat{h}_N . The latter exercise, as we will show in this paper, is technically and conceptually attractive.

Let $(\mathbb{R}, \mathfrak{B})$ denote the Euclidian space with Borel σ -algebra and $Z : \Omega \rightarrow \mathbb{R}$ an $\mathfrak{A} - \mathfrak{B}$ measurable function such that the distribution $\mathbb{P}_Z = f_Z \odot \mu_L$ is absolutely continuous (μ_L denotes the Lebesgue measure) in the probability space $(\mathbb{R}, \mathfrak{B}, \mathbb{P}_Z)$. Let F_Z and F_Z^{-1} denote the corresponding distribution and quantile functions. We furthermore assume $0 < f_Z(z) < \infty \forall z \in \mathbb{R}$, $F_Z(-\infty) = 0$ and $F_Z(\infty) = 1$ for a log-concave density f_Z as well as the existence of the first two derivatives of the density $f_Z(z)$ with respect to z ; both derivatives shall be bounded. We do not allow any unknown parameters for this distribution. Possible choices include the standard normal, standard logistic (SL) and minimum extreme value (MEV) distribution with distribution functions $F_Z(z) = \Phi(z)$, $F_Z(z) = F_{\text{SL}}(z) = (1 + \exp(-z))^{-1}$ and $F_Z(z) = F_{\text{MEV}}(z) = 1 - \exp(-\exp(z))$, respectively. In the first step, we will show that there always exists a unique and strictly monotonic transformation g such that the unknown and potentially complex distribution \mathbb{P}_Y that we are interested in can be generated from the simple and known distribution \mathbb{P}_Z via $\mathbb{P}_Y = \mathbb{P}_{g \circ Z}$. More formally, let $g : \mathbb{R} \rightarrow \Xi$ denote a $\mathfrak{B} - \mathfrak{C}$ measurable function. The composition $g \circ Z$ is a random variable on $(\Xi, \mathfrak{C}, \mathbb{P}_{g \circ Z})$. We can now formulate the existence and uniqueness of g as a corollary to the probability integral transform.

Corollary 1. *For all random variables Y and Z , there exists a unique strictly monotonically increasing transformation g , such that $\mathbb{P}_Y = \mathbb{P}_{g \circ Z}$.*

Proof. Let $g = F_Y^{-1} \circ F_Z$ and $Z \sim \mathbb{P}_Z$. Then $U := F_Z(Z) \sim U[0, 1]$ and $Y = F_Y^{-1}(U) \sim \mathbb{P}_Y$ by the probability integral transform. Let $h : \Xi \rightarrow \mathbb{R}$, such that $F_Y(y) = F_Z(h(y))$. From

$$F_Y(y) = (F_Z \circ F_Z^{-1} \circ F_Y)(y) \iff h = F_Z^{-1} \circ F_Y$$

we get the uniqueness of h and therefore g . The quantile function F_Z^{-1} and the distribution function F_Y exist by assumption and are both strictly monotonic and right-continuous. Therefore, h is strictly monotonic and right-continuous and so is g . \square

Corollary 2. *For $\mu = \mu_L$, we have $g = h^{-1}$ and $h'(y) = \frac{\partial h(y)}{\partial y} = f_Z((F_Z^{-1} \circ F_Y)(y))^{-1} f_Y(y)$.*

This result for absolutely continuous random variables Y can be found in many textbooks (*e.g.* Lindsey 1996), Corollary 1 also covers the discrete case.

Corollary 3. *For the counting measure $\mu = \mu_C$, $h = F_Z^{-1} \circ F_Y$ is a right-continuous step function because F_Y is a right-continuous step function with steps at $y \in \Xi$.*

We now characterise the distribution F_Y by the corresponding transformation function h , set up the corresponding likelihood of such a transformation function and estimate the transformation function based on this likelihood. Let $\mathcal{H} = \{h : \Xi \rightarrow \mathbb{R} \mid \mathfrak{C} - \mathfrak{B} \text{ measurable}, h(y_1) < h(y_2) \forall y_1 < y_2 \in \Xi\}$ denote the space of all strictly monotonic transformation functions.

With the transformation function h , we can evaluate F_Y as $F_Y(y | h) = F_Z(h(y)) \forall y \in \Xi$. Therefore, we only need to study the transformation function h ; the inverse transformation $g = h^{-1}$ (used to define a “group family” or “group model” by [Lehmann 1983](#); [Bickel et al. 1993](#)) is not necessary in what follows. The density for absolutely continuous variables Y ($\mu = \mu_L$) is now given by

$$f_Y(y | h) = f_Z(h(y))h'(y).$$

For discrete responses Y ($\mu = \mu_C$) with finite sample space $\Xi = \{y_1, \dots, y_K\}$, the density is

$$f_Y(y_k | h) = \begin{cases} F_Z(h(y_k)) & k = 1 \\ F_Z(h(y_k)) - F_Z(h(y_{k-1})) & k = 2, \dots, K-1 \\ 1 - F_Z(h(y_{k-1})) & k = K \end{cases}$$

and for countably infinite sample spaces $\Xi = \{y_1, y_2, y_3, \dots\}$, we get the density

$$f_Y(y_k | h) = \begin{cases} F_Z(h(y_k)) & k = 1 \\ F_Z(h(y_k)) - F_Z(h(y_{k-1})) & k > 1. \end{cases}$$

With the conventions $F_Z(h(y_0)) := F_Z(h(-\infty)) := 0$ and $F_Z(h(y_K)) := F_Z(h(\infty)) := 1$, we use the more compact notation $f_Y(y_k | h) = F_Z(h(y_k)) - F_Z(h(y_{k-1}))$ in the sequel.

For a given transformation function h , the likelihood contribution of a datum $C = (\underline{y}, \bar{y}] \in \mathfrak{C}$ is defined in terms of the distribution function ([Lindsey 1996](#)):

$$\mathcal{L}(h | Y \in C) := \int_C f_Y(y | h) d\mu(y) = F_Z(h(\bar{y})) - F_Z(h(\underline{y})).$$

This “exact” definition of the likelihood applies to most practical situations of interest and, in particular, allows discrete and (conceptually) continuous as well as censored or truncated observations C . For a discrete response y_k , we have $\bar{y} = y_k$ and $\underline{y} = y_{k-1}$, such that $\mathcal{L}(h | Y = y_k) = f_Y(y_k | h) = F_Z(h(\bar{y})) - F_Z(h(\underline{y}))$. For absolutely continuous random variables Y , we almost always observe an imprecise datum $(\underline{y}, \bar{y}] \subset \mathbb{R}$ and, for short intervals $(\underline{y}, \bar{y}]$, approximate the exact likelihood $\mathcal{L}(h | Y \in (\underline{y}, \bar{y}])$ by the term $(\bar{y} - \underline{y})f_Y(y | h)$ or simply $f_Y(y | h)$ with $y = (\underline{y} + \bar{y})/2$ ([Lindsey 1999](#)). This approximation only works for relatively precise measurements, *i.e.* short intervals. If longer intervals are observed, one speaks of “censoring” and relies on the exact definition of the likelihood contribution instead of using the above approximation ([Klein and Moeschberger 2003](#)). In summary, the likelihood contribution of a conceptually “exact continuous” or left-, right- or interval-censored continuous or discrete observation $(\underline{y}, \bar{y}]$ is given by

$$\mathcal{L}(h | Y \in (\underline{y}, \bar{y}]) \begin{cases} \approx f_Z(h(y))h'(y) & y = (\underline{y} + \bar{y})/2 \in \Xi \quad \text{“exact continuous”} \\ = 1 - F_Z(h(\underline{y})) & y \in (\underline{y}, \infty) \cap \Xi \quad \text{‘right-censored’} \\ = F_Z(h(\bar{y})) & y \in (-\infty, \bar{y}] \cap \Xi \quad \text{‘left-censored’} \\ = F_Z(h(\bar{y})) - F_Z(h(\underline{y})) & y \in (\underline{y}, \bar{y}] \cap \Xi \quad \text{‘interval-censored’}, \end{cases}$$

under the assumption of random censoring. The likelihood is more complex under dependent censoring ([Klein and Moeschberger 2003](#)), but we will not elaborate on this issue. The likelihood contribution $\mathcal{L}(h | Y \in (y_k, y_{k-1}])$ of an ordered factor in category y_k is equivalent to the term $\mathcal{L}(h | Y \in (\underline{y}, \bar{y}])$ contributed by an interval-censored observation $(\underline{y}, \bar{y}]$, when

category y_k is defined by the interval $(\underline{y}, \bar{y}]$. Thus, the expression $F_Z(h(\bar{y})) - F_Z(h(\underline{y}))$ for the likelihood contribution reflects the equivalence of interval-censoring and categorisation at corresponding cut-off points.

For truncated observations in the interval $(y_l, y_r] \subset \Xi$, the above likelihood contribution is defined in terms of the distribution function conditional on the truncation

$$F_Y(y | Y \in (y_l, y_r]) = F_Z(h(y) | Y \in (y_l, y_r]) = \frac{F_Z(h(y))}{F_Z(h(y_r)) - F_Z(h(y_l))} \quad \forall y \in (y_l, y_r]$$

and thus the likelihood contribution changes to (Klein and Moeschberger 2003)

$$\frac{\mathcal{L}(h | Y \in (\underline{y}, \bar{y}])}{F_Z(h(y_r)) - F_Z(h(y_l))} = \frac{\mathcal{L}(h | Y \in (\underline{y}, \bar{y}])}{\mathcal{L}(h | Y \in (y_l, y_r])} \quad \text{when } y_l < \underline{y} < \bar{y} \leq y_r.$$

It is important to note that the likelihood is always *defined* in terms of a distribution function (Lindsey 1999), and it therefore makes sense to directly model the distribution function of interest. The ability to uniquely characterise this distribution function by the transformation function h gives rise to the following definition of an estimator \hat{h}_N .

Definition 1 (Most likely transformation). *Let C_1, \dots, C_N denote an independent sample of possibly randomly censored or truncated observations from \mathbb{P}_Y . The estimator*

$$\hat{h}_N := \arg \max_{h \in \mathcal{H}} \sum_{i=1}^N \log(\mathcal{L}(\tilde{h} | Y \in C_i))$$

is called the most likely transformation (MLT).

Log-concavity of f_Z ensures concavity of the log-likelihood (except when all observations are right-censored) and thus ensures the existence and uniqueness of \hat{h}_N .

Example For an absolutely continuous response Y the likelihood and log-likelihood for h are approximated by the density and log-density evaluated at $y = (\underline{y} + \bar{y})/2$, respectively:

$$\begin{aligned} \mathcal{L}(h | Y \in (\underline{y}, \bar{y}]) &\approx f_Z(h(y))h'(y) \\ \log(\mathcal{L}(h | Y \in (\underline{y}, \bar{y}])) &\approx \log(f_Z(h(y))) + \log(h'(y)). \end{aligned}$$

Strict monotonicity of the transformation function h is required; otherwise the likelihood is not defined. The term $\log(h'(y))$ is not a penalty term, but the likelihood favours transformation functions with a large positive derivative at the observations. If we assume $Y \sim N(\alpha, \sigma^2)$ and for the choice $Z \sim N(0, 1)$ with $F_Z = \Phi$ and $f_Z = \phi$, we can restrict h to linear functions $h(y) = (y - \alpha)\sigma^{-1}$. The likelihood reduces to

$$\mathcal{L}(h | Y \in (\underline{y}, \bar{y}]) \approx f_Z(h(y))h'(y) = \phi((y - \alpha)\sigma^{-1})\sigma^{-1} = \phi_{\alpha, \sigma^2}(y) = f_Y(y | \alpha, \sigma^2).$$

In this simple location-scale family, the most likely transformation is characterised by the parameters of the normal distribution of Y . It is important to note that for other choices of F_Z , the most likely transformation is non-linear; however, the distribution function $F_Y = F_Z(h(y))$ is invariant with respect to F_Z because we can always write h as $F_Z^{-1} \circ F_Y$. In other words,

with $F_Z \neq \Phi$, we can still model normal responses Y ; however, a non-linear transformation function h is required.

Many distributions are defined by a transformation function h , for example, the Box-Cox power exponential family (Rigby and Stasinopoulos 2004), the sinh-arcsinh distributions (Jones and Pewsey 2009), or the T-X family of distributions (Alzaatreh *et al.* 2013). The parameters of these distributions can, for example, be estimated by the GAMLSS approach (Rigby and Stasinopoulos 2005). In what follows, we do not assume any specific form of the transformation function but parameterise h in terms of basis functions. We now introduce such a parameterisation, a corresponding family of distributions, a maximum likelihood estimator and a large class of models for unconditional and conditional distributions.

3. Transformation Analysis

We parameterise the transformation function $h(y)$ as a linear function of its basis-transformed argument y using a basis function $\mathbf{a} : \Xi \rightarrow \mathbb{R}^P$, such that $h(y) = \mathbf{a}(y)^\top \boldsymbol{\vartheta}$, $\boldsymbol{\vartheta} \in \mathbb{R}^P$. The choice of the basis function \mathbf{a} is problem specific and will be discussed in Section 4. The likelihood \mathcal{L} only requires evaluation of h , and only an approximation thereof using the Lebesgue density of “exact continuous” observations makes the evaluation of the first derivative of $h(y)$ with respect to y necessary. In this case, the derivative with respect to y is given by $h'(y) = \mathbf{a}'(y)^\top \boldsymbol{\vartheta}$, and we assume that \mathbf{a}' is available. In the following, we will write $h = \mathbf{a}^\top \boldsymbol{\vartheta}$ and $h' = \mathbf{a}'^\top \boldsymbol{\vartheta}$ for the transformation function and its first derivative, omitting the argument y , and we assume that both functions are bounded away from $-\infty$ and ∞ . For a specific choice of F_Z and \mathbf{a} , the transformation family of distributions consists of all distributions \mathbb{P}_Y whose distribution function F_Y is given as the composition $F_Z \circ \mathbf{a}^\top \boldsymbol{\vartheta}$; this family can be formally defined as follows.

Definition 2 (Transformation family). *The distribution family*

$$\mathbb{P}_{Y,\Theta} = \{F_Z \circ \mathbf{a}^\top \boldsymbol{\vartheta} \mid \boldsymbol{\vartheta} \in \Theta\}$$

with parameter space $\Theta = \{\boldsymbol{\vartheta} \in \mathbb{R}^P \mid \mathbf{a}^\top \boldsymbol{\vartheta} \in \mathcal{H}\}$ is called transformation family of distributions $\mathbb{P}_{Y,\boldsymbol{\vartheta}}$ with transformation functions $\mathbf{a}^\top \boldsymbol{\vartheta} \in \mathcal{H}$, μ -densities $f_Y(y \mid \boldsymbol{\vartheta})$, $y \in \Xi$, and error distribution function F_Z .

The classical definition of a transformation family relies on the idea of invariant distributions, *i.e.* only the parameters of a distribution are changed by a transformation function but the distribution itself is not changed. The normal family characterised by affine transformations is the most well-known example (*e.g.* Fraser 1968; Lindsey 1996). Here, we explicitly allow and encourage transformation functions that change the shape of the distribution. The transformation function $\mathbf{a}^\top \boldsymbol{\vartheta}$ is, at least in principle, flexible enough to generate any distribution function $F_Y = F_Z \circ \mathbf{a}^\top \boldsymbol{\vartheta}$ from the distribution function F_Z . We borrow the term “error distribution” function for F_Z from Fraser (1968), because Z can be understood as an error term in some of the models discussed in Section 4. The problem of estimating the unknown transformation function h , and thus the unknown distribution function F_Y , reduces to the problem of estimating the parameter vector $\boldsymbol{\vartheta}$ through maximisation of the likelihood function. We assume that the basis function \mathbf{a} is such that the parameters $\boldsymbol{\vartheta}$ are identifiable.

Definition 3 (Maximum likelihood estimator).

$$\hat{\boldsymbol{\vartheta}}_N := \arg \max_{\boldsymbol{\vartheta} \in \Theta} \sum_{i=1}^N \log(\mathcal{L}(\mathbf{a}^\top \boldsymbol{\vartheta} \mid Y \in C_i))$$

Based on the maximum likelihood estimator $\hat{\boldsymbol{\vartheta}}_N$, we define plug-in estimators of the most likely transformation function and the corresponding estimator of our target distribution F_Y as $\hat{h}_N := \mathbf{a}^\top \hat{\boldsymbol{\vartheta}}_N$ and $\hat{F}_{Y,N} := F_Z \circ \hat{h}_N$. Because the problem of estimating an unknown distribution function is now embedded in the maximum likelihood framework, the asymptotic analysis benefits from standard results on the asymptotic behaviour of maximum likelihood estimators. We begin with deriving the score function and Fisher information. The score contribution of an “exact continuous” observation $y = (\underline{y} + \bar{y})/2$ from an absolutely continuous distribution is approximated by the gradient of the log-density

$$\begin{aligned} \mathbf{s}(\boldsymbol{\vartheta} \mid Y \in (\underline{y}, \bar{y}]) &\approx \frac{\partial \log(f_Y(y \mid \boldsymbol{\vartheta}))}{\partial \boldsymbol{\vartheta}} = \frac{\partial \log(f_Z(\mathbf{a}(y)^\top \boldsymbol{\vartheta})) + \log(\mathbf{a}'(y)^\top \boldsymbol{\vartheta})}{\partial \boldsymbol{\vartheta}} \\ &= \mathbf{a}(y) \frac{f'_Z(\mathbf{a}(y)^\top \boldsymbol{\vartheta})}{f_Z(\mathbf{a}(y)^\top \boldsymbol{\vartheta})} + \frac{\mathbf{a}'(y)}{\mathbf{a}'(y)^\top \boldsymbol{\vartheta}}. \end{aligned} \quad (1)$$

For an interval-censored or discrete observation y and \bar{y} (the constant terms $F_Z(\mathbf{a}(-\infty)^\top \boldsymbol{\vartheta}) = F_Z(-\infty) = 0$ and $F_Z(\mathbf{a}(\infty)^\top \boldsymbol{\vartheta}) = F_Z(\infty) = 1$ vanish), the score contribution is

$$\begin{aligned} \mathbf{s}(\boldsymbol{\vartheta} \mid Y \in (\underline{y}, \bar{y}]) &= \frac{\partial \log(\mathcal{L}(\mathbf{a}^\top \boldsymbol{\vartheta} \mid Y \in (\underline{y}, \bar{y}]))}{\partial \boldsymbol{\vartheta}} \\ &= \frac{\partial \log(F_Z(\mathbf{a}(\bar{y})^\top \boldsymbol{\vartheta}) - F_Z(\mathbf{a}(\underline{y})^\top \boldsymbol{\vartheta}))}{\partial \boldsymbol{\vartheta}} \\ &= \frac{f_Z(\mathbf{a}(\bar{y})^\top \boldsymbol{\vartheta}) \mathbf{a}(\bar{y}) - f_Z(\mathbf{a}(\underline{y})^\top \boldsymbol{\vartheta}) \mathbf{a}(\underline{y})}{F_Z(\mathbf{a}(\bar{y})^\top \boldsymbol{\vartheta}) - F_Z(\mathbf{a}(\underline{y})^\top \boldsymbol{\vartheta})}. \end{aligned} \quad (2)$$

For a truncated observation, the score function is $\mathbf{s}(\boldsymbol{\vartheta} \mid Y \in (\underline{y}, \bar{y}]) - \mathbf{s}(\boldsymbol{\vartheta} \mid Y \in (y_l, y_r])$.

The contribution of an “exact continuous” observation y from an absolutely continuous distribution to the Fisher information is approximately

$$\begin{aligned} \mathbf{F}(\boldsymbol{\vartheta} \mid Y \in (\underline{y}, \bar{y}]) &\approx -\frac{\partial^2 \log(f_Y(y \mid \boldsymbol{\vartheta}))}{\partial \boldsymbol{\vartheta} \partial \boldsymbol{\vartheta}^\top} \\ &= -\left(\mathbf{a}(y) \mathbf{a}(y)^\top \left\{ \frac{f''_Z(\mathbf{a}(y)^\top \boldsymbol{\vartheta})}{f_Z(\mathbf{a}(y)^\top \boldsymbol{\vartheta})} - \left[\frac{f'_Z(\mathbf{a}(y)^\top \boldsymbol{\vartheta})}{f_Z(\mathbf{a}(y)^\top \boldsymbol{\vartheta})} \right]^2 \right\} \right. \\ &\quad \left. - \frac{\mathbf{a}'(y) \mathbf{a}'(y)^\top}{(\mathbf{a}'(y)^\top \boldsymbol{\vartheta})^2} \right) \end{aligned} \quad (3)$$

(NB: the weight to $\mathbf{a}(y) \mathbf{a}(y)^\top$ is constant one for $F_Z = \Phi$). For a censored or discrete

observation, we have the following contribution to the Fisher information

$$\begin{aligned}
\mathbf{F}(\boldsymbol{\vartheta} \mid Y \in (\underline{y}, \bar{y}]) &= -\frac{\partial^2 \log(\mathcal{L}(\mathbf{a}^\top \boldsymbol{\vartheta} \mid Y \in (\underline{y}, \bar{y}]))}{\partial \boldsymbol{\vartheta} \partial \boldsymbol{\vartheta}^\top} \\
&= -\left\{ \frac{f'_Z(\mathbf{a}(\bar{y})^\top \boldsymbol{\vartheta}) \mathbf{a}(\bar{y}) \mathbf{a}(\bar{y})^\top - f'_Z(\mathbf{a}(\underline{y})^\top \boldsymbol{\vartheta}) \mathbf{a}(\underline{y}) \mathbf{a}(\underline{y})^\top}{F_Z(\mathbf{a}(\bar{y})^\top \boldsymbol{\vartheta}) - F_Z(\mathbf{a}(\underline{y})^\top \boldsymbol{\vartheta})} \right. \\
&\quad \left. - \frac{[f_Z(\mathbf{a}(\bar{y})^\top \boldsymbol{\vartheta}) \mathbf{a}(\bar{y}) - f_Z(\mathbf{a}(\underline{y})^\top \boldsymbol{\vartheta}) \mathbf{a}(\underline{y})]}{[F_Z(\mathbf{a}(\bar{y})^\top \boldsymbol{\vartheta}) - F_Z(\mathbf{a}(\underline{y})^\top \boldsymbol{\vartheta})]^2} \times \right. \\
&\quad \left. [f_Z(\mathbf{a}(\bar{y})^\top \boldsymbol{\vartheta}) \mathbf{a}(\bar{y})^\top - f_Z(\mathbf{a}(\underline{y})^\top \boldsymbol{\vartheta}) \mathbf{a}(\underline{y})^\top] \right\}. \tag{4}
\end{aligned}$$

For a truncated observation, the Fisher information is given by $\mathbf{F}(\boldsymbol{\vartheta} \mid Y \in (\underline{y}, \bar{y}]) - \mathbf{F}(\boldsymbol{\vartheta} \mid Y \in (y_l, y_r])$.

We will first discuss the asymptotic properties of the maximum likelihood estimator $\hat{\boldsymbol{\vartheta}}_N$ in the parametric setting with fixed parameters $\boldsymbol{\vartheta}$ in both the discrete and continuous case. For continuous variables Y and a transformation function parameterised using a Bernstein polynomial, results for sieve maximum likelihood estimation, where the number of parameters increases with N , are then discussed in Subsection 3.2.

3.1. Parametric Inference

Conditions on the densities of the error distribution f_Z and the basis functions \mathbf{a} ensuring consistency and asymptotic normality of the sequence of maximum likelihood estimators $\hat{\boldsymbol{\vartheta}}_N$ and an estimator of their asymptotic covariance matrix are given in the following three theorems. Due to the full parameterisation of the model, the proofs are simple standard results for likelihood asymptotics, and a more complex analysis (as required for estimation equations in the presence of a nuisance parameter h_Y , for example in Cheng *et al.* 1995; Chen *et al.* 2002) is not necessary. We will restrict ourselves to absolutely continuous or discrete random variables Y , where the likelihood is given in terms of the density $f_Y(y \mid \boldsymbol{\vartheta})$. Furthermore, we will only study the case of a correctly specified transformation $h = \mathbf{a}^\top \boldsymbol{\vartheta}$ and refer the reader to Hothorn *et al.* (2014), where consistency results for arbitrary h are given.

Theorem 1. For $Y_1, \dots, Y_N \stackrel{iid}{\sim} \mathbb{P}_{Y, \boldsymbol{\vartheta}_0}$ and under the assumptions (A1) the parameter space Θ is compact and (A2) $\mathbb{E}_{\boldsymbol{\vartheta}_0}[\sup_{\boldsymbol{\vartheta} \in \Theta} |\log(f_Y(Y \mid \boldsymbol{\vartheta}))|] < \infty$ where $\boldsymbol{\vartheta}_0$ is well-separated:

$$\sup_{\boldsymbol{\vartheta}; |\boldsymbol{\vartheta} - \boldsymbol{\vartheta}_0| \geq \epsilon} \mathbb{E}_{\boldsymbol{\vartheta}_0}[\log(f_Y(Y \mid \boldsymbol{\vartheta}))] < \mathbb{E}_{\boldsymbol{\vartheta}_0}[\log(f_Y(Y \mid \boldsymbol{\vartheta}_0))],$$

the sequence of estimators $\hat{\boldsymbol{\vartheta}}_N$ converges to $\boldsymbol{\vartheta}_0$ in probability, $\hat{\boldsymbol{\vartheta}}_N \xrightarrow{\mathbb{P}} \boldsymbol{\vartheta}_0$, as $N \rightarrow \infty$.

Proof. The log-likelihood is continuous in $\boldsymbol{\vartheta}$, and due to (A2), each log-likelihood contribution is dominated by an integrable function. Thus, the result follows from van der Vaart (1998) (Theorem 5.8 with Example 19.7; see note at bottom of page 46). \square

Remark 1. Assumption (A1) is made for convenience, and relaxations of such a condition are given in van de Geer (2000) or van der Vaart (1998). The assumptions in (A2) are rather weak: the first one holds if the functions \mathbf{a} are not arbitrarily ill-posed, and the second one holds if the function $\mathbb{E}_{\boldsymbol{\vartheta}_0}[\log(f_Y(Y \mid \boldsymbol{\vartheta}))]$ is strictly convex in $\boldsymbol{\vartheta}$ (if the assumption would not hold, we would still have convergence to the set $\operatorname{argmax}_{\boldsymbol{\vartheta}} \mathbb{E}_{\boldsymbol{\vartheta}_0}[\log(f_Y(Y \mid \boldsymbol{\vartheta}))]$).

Theorem 2. *Under the assumptions of Theorem 1 and in addition (A3)*

$$\mathbb{E}_{\boldsymbol{\vartheta}_0} \left(\sup_{\boldsymbol{\vartheta}} \left\| \frac{\partial \log f_Y(Y | \boldsymbol{\vartheta})}{\partial \boldsymbol{\vartheta}} \right\|^2 \right) < \infty,$$

(A4) $\mathbb{E}_{\boldsymbol{\vartheta}_0}(\mathbf{a}(Y)\mathbf{a}(Y)^\top)$ and (for the absolutely continuous case $\mu = \mu_L$ only) $\mathbb{E}_{\boldsymbol{\vartheta}_0}(\mathbf{a}'(Y)\mathbf{a}'(Y)^\top)$ are nonsingular, and (A5) $0 < f_Z < \infty$, $\sup |f_Z'| < \infty$ and $\sup |f_Z''| < \infty$, the sequence $\sqrt{N}(\hat{\boldsymbol{\vartheta}}_N - \boldsymbol{\vartheta}_0)$ is asymptotically normal with mean zero and covariance matrix

$$\Sigma_{\boldsymbol{\vartheta}_0} = \left(\mathbb{E}_{\boldsymbol{\vartheta}_0} \left(- \frac{\partial^2 \log f_Y(Y | \boldsymbol{\vartheta})}{\partial \boldsymbol{\vartheta} \partial \boldsymbol{\vartheta}^\top} \right) \right)^{-1},$$

as $N \rightarrow \infty$.

Proof. Because the map $\boldsymbol{\vartheta} \mapsto \sqrt{f_Y(y | \boldsymbol{\vartheta})}$ is continuously differentiable in $\boldsymbol{\vartheta}$ for all y in both the discrete and absolutely continuous case and the matrix

$$\mathbb{E}_{\boldsymbol{\vartheta}_0} \left(\left[\frac{\partial \log f_Y(Y | \boldsymbol{\vartheta})}{\partial \boldsymbol{\vartheta}} \right] \left[\frac{\partial \log f_Y(Y | \boldsymbol{\vartheta})}{\partial \boldsymbol{\vartheta}} \right]^\top \right)$$

is continuous in $\boldsymbol{\vartheta}$ as given in (1) and (2), the transformation family $\mathbb{P}_{Y, \Theta}$ is differentiable in quadratic mean with Lemma 7.6 in van der Vaart (1998). Furthermore, assumptions (A4-5) ensure that the expected Fisher information matrix is nonsingular at $\boldsymbol{\vartheta}_0$. With the consistency and (A3), the result follows from Theorem 5.39 in van der Vaart (1998). \square

Remark 2. *Assumption (A4) is valid for the densities f_Z of the normal, logistic and minimum extreme value distribution. The Fisher information (4) and (4) evaluated at the maximum likelihood estimator $\hat{\boldsymbol{\vartheta}}_N$ can be used to estimate the covariance matrix $\Sigma_{\boldsymbol{\vartheta}_0}$.*

Theorem 3. *Under the assumptions of Theorem 2 and assuming $\mathbb{E}_{\boldsymbol{\vartheta}_0}|\mathbf{F}(\boldsymbol{\vartheta}_0 | Y)| < \infty$, a consistent estimator for $\Sigma_{\boldsymbol{\vartheta}_0}$ is given by*

$$\hat{\Sigma}_{\boldsymbol{\vartheta}_0, N} = \left(N^{-1} \sum_{i=1}^N \mathbf{F}(\hat{\boldsymbol{\vartheta}}_N | Y_i) \right)^{-1}.$$

Proof. With the law of large numbers we have

$$N^{-1} \sum_{i=1}^N \mathbf{F}(\boldsymbol{\vartheta}_0 | Y_i) = N^{-1} \sum_{i=1}^N - \frac{\partial^2 \log f_Y(Y_i | \boldsymbol{\vartheta}_0)}{\partial \boldsymbol{\vartheta} \partial \boldsymbol{\vartheta}^\top} \xrightarrow{\mathbb{P}} \mathbb{E}_{\boldsymbol{\vartheta}_0} \left(- \frac{\partial^2 \log f_Y(Y | \boldsymbol{\vartheta}_0)}{\partial \boldsymbol{\vartheta} \partial \boldsymbol{\vartheta}^\top} \right) = \Sigma_{\boldsymbol{\vartheta}_0}^{-1}.$$

Because the map $\boldsymbol{\vartheta} \mapsto \mathbf{F}(\boldsymbol{\vartheta} | y)$ is continuous for all y (as can be seen from (4) and (4)), the result follows with Theorem 1. \square

Based on Theorems 1-3, we can perform standard likelihood inference on the model parameters $\boldsymbol{\vartheta}$. In particular, we can construct confidence intervals and confidence bands for the conditional distribution function from confidence intervals and bands for the linear functions $\mathbf{a}^\top \boldsymbol{\vartheta}$. We complete this part by formally defining the class of transformation models.

Definition 4 (Transformation model). *The triple $(F_Z, \mathbf{a}, \boldsymbol{\vartheta})$ is called transformation model.*

The transformation model $(F_Z, \mathbf{a}, \boldsymbol{\vartheta})$ fully defines the distribution of Y via $F_Y = F_Z \circ \mathbf{a}^\top \boldsymbol{\vartheta}$ and thus the corresponding likelihood $\mathcal{L}(\mathbf{a}^\top \boldsymbol{\vartheta} \mid Y \in (y, \bar{y}])$. Our definition of transformation models as $(F_Z, \mathbf{a}, \boldsymbol{\vartheta})$ is strongly tied to the idea of structural inference (Fraser 1968) and group families (Lehmann 1983) or group models (Bickel *et al.* 1993). Fraser (1968) described a measurement model \mathbb{P}_Y for Y by an error distribution \mathbb{P}_Z and a structural equation $Y = g \circ Z$, where g is a linear function, thereby extending the location-scale family $Y = \alpha + \sigma Z$ introduced by Fisher (1934) and refined by Pitman (1939). Group models consist of distributions generated by possibly non-linear g . The main difference to these classical approaches is that we parameterise h instead of $g = h^{-1}$. By extending the linear transformation functions g dealt with by Fraser (1968) to non-linear transformations, we approximate the potentially non-linear transformation functions $h = g^{-1} = F_Z^{-1} \circ F_Y$ by $\mathbf{a}^\top \boldsymbol{\vartheta}$, with subsequent estimation of the parameters $\boldsymbol{\vartheta}$. For given parameters $\boldsymbol{\vartheta}$, a sample from \mathbb{P}_Y can be drawn by the probability integral transform, *i.e.* $Z_1, \dots, Z_N \stackrel{\text{iid}}{\sim} \mathbb{P}_Z$ is drawn and then $Y_i = \inf\{y \in \Xi \mid \mathbf{a}(y)^\top \boldsymbol{\vartheta} \geq Z_i\}$. This generalises the method published by Bender *et al.* (2005) from the Cox model to all conditional transformation models.

3.2. Non-parametric Inference

For continuous responses Y , any unknown transformation h can be approximated by Bernstein polynomials of increasing order (Farouki 2012). For uncensored and right-censored responses and under the same conditions for F_Z as stated in Subsection 3.1, McLain and Ghosh (2013) showed that the non-parametric sieve maximum likelihood estimator is consistent with rate of convergence $N^{2/5}$ for h with continuous bounded second derivatives in unconditional and linear transformation models (see Subsection 4.3). In the latter class, the linear shift parameters $\boldsymbol{\beta}$ are asymptotically normal and semi-parametrically efficient. Numerical approximations to the observed Fisher information $\mathbf{F}(\hat{\boldsymbol{\vartheta}}_N \mid Y \in (y, \bar{y}])$ were shown to lead to appropriate standard errors of $\hat{\boldsymbol{\beta}}_N$ by McLain and Ghosh (2013). Hothorn *et al.* (2014) established the consistency of boosted non-parametric conditional transformation models (see Subsection 4.2). For sieve maximum likelihood estimation in the class of conditional transformation models, the techniques employed by McLain and Ghosh (2013) require minor technical extensions, which are omitted here.

In summary, the same limiting distribution arises under both the parametric and the non-parametric paradigm for transformation functions parameterised or approximated using Bernstein polynomials, respectively. In the latter case, the target is then the best approximated transformation function with Bernstein polynomials, say h_N^* (where the index N indicates that we use a more complex approximation when N increases). If the approximation error $h_N^* - h$ is of smaller order than the convergence rate of the estimator, the estimator's target becomes the true underlying transformation function h , and otherwise a bias for estimating h remains.

4. Applications

The definition of transformation models tailored for specific situations “only” requires the definition of a suitable basis function \mathbf{a} and a choice of F_Z . In this section, we will discuss specific transformation models for unconditional and conditional distributions of ordered and unordered categorical, discrete and continuous responses Y . Note that the likelihood function

\mathcal{L} allows all these models to be fitted to arbitrarily censored or truncated responses; for brevity, we will not elaborate on the details.

4.1. Unconditional Transformation Models

Finite Sample Space For ordered categorical responses Y from a finite sample space $\Xi = \{y_1, \dots, y_K\}$, we assign one parameter to each element of the sample space except y_K . This corresponds to the basis function $\mathbf{a}(y_k) = \mathbf{e}_{K-1}(k)$, where $\mathbf{e}_{K-1}(k)$ is the unit vector of length $K-1$, with its k th element being one. The transformation function h is

$$h(y_k) = \mathbf{e}_{K-1}(k)^\top \boldsymbol{\vartheta} = \vartheta_k \in \mathbb{R}, \quad 1 \leq k < K, \quad \text{st} \quad \vartheta_1 < \dots < \vartheta_{K-1},$$

with $h(y_K) = \infty$, and the unconditional distribution function of F_Y is $F_Y(y_k) = F_Z(\vartheta_k)$. This parameterisation underlies the common proportional odds and proportional hazards model for ordered categorical data (Tutz 2012). Note that monotonicity of h is guaranteed by the $K-2$ linear constraints $\vartheta_2 - \vartheta_1 > 0, \dots, \vartheta_{K-1} - \vartheta_{K-2} > 0$ when constrained optimisation is performed. In the absence of censoring or truncation and with $\vartheta_0 = -\infty, \vartheta_K = \infty$, we obtain the maximum likelihood estimator for $\boldsymbol{\vartheta}$ as

$$\begin{aligned} \hat{\boldsymbol{\vartheta}}_N &= \arg \max_{\vartheta_1 < \dots < \vartheta_{K-1}} \sum_{i=1}^N \log(F_Z(\vartheta_{k(i)}) - F_Z(\vartheta_{k(i)-1})) \\ &= \left(F_Z^{-1} \left(N^{-1} \sum_{i=1}^N \mathbf{1}(Y_i \leq y_1) \right), \dots, F_Z^{-1} \left(N^{-1} \sum_{i=1}^N \mathbf{1}(Y_i \leq y_{K-1}) \right) \right)^\top \end{aligned}$$

because $\hat{\pi}_k = N^{-1} \sum_{i=1}^N \mathbf{1}(Y_i = y_k), 1 \leq k < K$ maximises the equivalent multinomial (or empirical) log-likelihood $\sum_{i=1}^N \log(\pi_{k(i)})$, and we can rewrite this estimator as

$$\hat{\pi}_k = N^{-1} \left(\sum_{i=1}^N \mathbf{1}(Y_i \leq y_k) - \mathbf{1}(k > 1) \sum_{i=1}^N \mathbf{1}(Y_i \leq y_{k-1}) \right), \quad 1 \leq k < K.$$

The estimated distribution function $\hat{F}_{Y,N} = F_Z \circ \hat{h}_N$ is invariant with respect to F_Z . Assumption (A4) is valid for these basis functions because we have $\mathbb{E}_{\boldsymbol{\vartheta}_0}(\mathbf{e}_{K-1}(Y)\mathbf{e}_{K-1}(Y)^\top) = \text{diag}(\mathbb{P}(Y = y_k)), 1 \leq k < K$ for $Y \sim \mathbb{P}_{Y,\boldsymbol{\vartheta}_0}$.

If we define the sample space Ξ as the set of unique observed values and the probability measure as the empirical cumulative distribution function (ECDF), putting mass N^{-1} on each observation, we see that this particular parameterisation is equivalent to an empirical likelihood approach and we get $\hat{h}_N = F_Z^{-1} \circ \text{ECDF}$. Note that although the transformation function depends on the choice of F_Z , the estimated distribution function $\hat{F}_{Y,N} = F_Z \circ \hat{h}_N = \text{ECDF}$ does not and is simply the non-parametric empirical maximum likelihood estimator. A smoothed version of this estimator for continuous responses is discussed in the next paragraph.

Infinite Sample Space For continuous responses Y , the parameterisation $h(y) = \mathbf{a}(y)^\top \boldsymbol{\vartheta}$, and thus also $\hat{F}_{Y,N}$, should be smooth in y ; therefore, any polynomial or spline basis is a suitable choice for \mathbf{a} . For the empirical experiments in Section 5, we applied Bernstein

polynomials (for an Overview, see Farouki 2012) of order M ($P = M + 1$) defined on the interval $[\underline{y}, \bar{y}]$ with

$$\begin{aligned} \mathbf{a}_{\text{Bs},M}(y) &= (M+1)^{-1}(f_{\text{Be}(1,M+1)}(\tilde{y}), \dots, f_{\text{Be}(m,M-m+1)}(\tilde{y}), \dots, f_{\text{Be}(M+1,1)}(\tilde{y}))^\top \in \mathbb{R}^{M+1} \\ h(y) &= \mathbf{a}_{\text{Bs},M}(y)^\top \boldsymbol{\vartheta} = \sum_{m=0}^M \vartheta_m f_{\text{Be}(m+1,M-m+1)}(\tilde{y}) / (M+1) \\ h'(y) &= \mathbf{a}'_{\text{Bs},M}(y)^\top \boldsymbol{\vartheta} = \sum_{m=0}^{M-1} (\vartheta_{m+1} - \vartheta_m) f_{\text{Be}(m+1,M-m)}(\tilde{y}) M / ((M+1)(\bar{y} - \underline{y})), \end{aligned}$$

where $\tilde{y} = (y - \underline{y}) / (\bar{y} - \underline{y}) \in [0, 1]$ and $f_{\text{Be}(m,M)}$ is the density of the Beta distribution with parameters m and M . This choice is computationally attractive because strict monotonicity can be formulated as a set of M linear constraints on the parameters $\vartheta_m < \vartheta_{m+1}$ for all $m = 0, \dots, M$ (Curtis and Ghosh 2011). Therefore, application of constrained optimisation guarantees monotonic estimates \hat{h}_N . The basis contains an intercept. We obtain smooth plug-in estimators for the distribution, density, hazard and cumulative hazard functions as $\hat{F}_{Y,N} = F_Z \circ \mathbf{a}_{\text{Bs},M}^\top \hat{\boldsymbol{\vartheta}}_N$, $\hat{f}_{Y,N} = f_Z \circ \mathbf{a}_{\text{Bs},M}^\top \hat{\boldsymbol{\vartheta}}_N \times \mathbf{a}'_{\text{Bs},M}^\top \hat{\boldsymbol{\vartheta}}_N$, $\hat{\lambda}_{Y,N} = \hat{f}_{Y,N} / (1 - \hat{F}_{Y,N})$ and $\hat{\Lambda}_{Y,N} = -\log(1 - \hat{F}_{Y,N})$. The estimator $\hat{F}_{Y,N} = F_Z \circ \mathbf{a}_{\text{Bs},M}^\top \hat{\boldsymbol{\vartheta}}_N$ must not be confused with the estimator $\hat{F}_{Y,N} = \mathbf{a}_{\text{Bs},M}^\top \hat{\mathbf{p}}$ for $Y \in [0, 1]$ obtained from the smoothed empirical distribution function with coefficients $\hat{\mathbf{p}}_{m+1} = \sum_{i=1}^N \mathbb{1}(Y_i \leq m/M) / N$ corresponding to probabilities evaluated at the quantiles m/M for $m = 0, \dots, M$ (Babu *et al.* 2002).

The question arises how the degree of the polynomial affects the estimated distribution function. On the one hand, the model $(\Phi, \mathbf{a}_{\text{Bs},1}, \boldsymbol{\vartheta})$ only allows linear transformation functions of a standard normal and F_Y is restricted to the normal family. On the other hand, $(\Phi, \mathbf{a}_{\text{Bs},N-1}, \boldsymbol{\vartheta})$ has one parameter for each observation and $\hat{F}_{Y,N}$ is the non-parametric maximum likelihood estimator ECDF, which, by the Glivenko-Cantelli lemma, converges to F_Y . In this sense, we cannot choose a “too large” value for M . This is a consequence of the monotonicity constraint on the estimator $\mathbf{a}^\top \hat{\boldsymbol{\vartheta}}_N$, which, in this extreme case, just interpolates the step function $F_Z^{-1} \circ \text{ECDF}$. Empirical evidence for the insensitivity of results when M is large can be found in Hothorn (2017b) and in the discussion.

4.2. Conditional Transformation Models

In the following, we will discuss a cascade of increasingly complex transformation models where the transformation function h may depend on explanatory variables $\mathbf{X} \in \chi$. We are interested in estimating the conditional distribution of Y given $\mathbf{X} = \mathbf{x}$. The corresponding distribution function $F_{Y|\mathbf{X}=\mathbf{x}}$ can be written as $F_{Y|\mathbf{X}=\mathbf{x}}(y) = F_Z(h(y | \mathbf{x}))$. The transformation function $h(\cdot | \mathbf{x}) : \Xi \rightarrow \mathbb{R}$ is said to be conditional on \mathbf{x} . Following the arguments presented in the proof of Corollary 1, it is easy to see that for each \mathbf{x} , there exists a strictly monotonic transformation function $h(\cdot | \mathbf{x}) = F_Z^{-1} \circ F_{Y|\mathbf{X}=\mathbf{x}}$ such that $F_{Y|\mathbf{X}=\mathbf{x}}(y) = F_Z(h(y | \mathbf{x}))$. Because this class of conditional transformation models and suitable parameterisations were introduced by Hothorn *et al.* (2014), we will only sketch the most important aspects here.

Let $\mathbf{b} : \chi \rightarrow \mathbb{R}^Q$ denote a basis transformation of the explanatory variables. The joint basis for both y and \mathbf{x} is called $\mathbf{c} : \Xi \times \chi \rightarrow \mathbb{R}^{d(P,Q)}$; its dimension $d(P,Q)$ depends on the way the two basis functions \mathbf{a} and \mathbf{b} are combined (*e.g.* $\mathbf{c} = (\mathbf{a}^\top, \mathbf{b}^\top)^\top \in \mathbb{R}^{P+Q}$ or $\mathbf{c} = (\mathbf{a}^\top \otimes \mathbf{b}^\top)^\top \in \mathbb{R}^{PQ}$). The conditional transformation function is now parameterised

as $h(y | \mathbf{x}) = \mathbf{c}(y, \mathbf{x})^\top \boldsymbol{\vartheta}$. One important special case is the simple transformation function $h(y | \mathbf{x}) = h_Y(y) + h_{\mathbf{x}}(\mathbf{x})$, where the explanatory variables only contribute a shift $h_{\mathbf{x}}(\mathbf{x})$ to the conditional transformation function. Often this shift is assumed to be linear in \mathbf{x} ; therefore, we use the function $m(\mathbf{x}) = \mathbf{b}(\mathbf{x})^\top \boldsymbol{\beta} = \tilde{\mathbf{x}}^\top \boldsymbol{\beta}$ to denote linear shifts. Here $\mathbf{b}(\mathbf{x}) = \tilde{\mathbf{x}}$ is one row of the design matrix without intercept. These simple models correspond to the joint basis $\mathbf{c}(y, \mathbf{x})^\top \boldsymbol{\vartheta} = \mathbf{a}(y)^\top \boldsymbol{\vartheta}_1 + \mathbf{b}(\mathbf{x})^\top \boldsymbol{\vartheta}_2$, with $h_Y(y) = \mathbf{a}(y)^\top \boldsymbol{\vartheta}_1$ and $h_{\mathbf{x}}(\mathbf{x}) = \mathbf{b}(\mathbf{x})^\top \boldsymbol{\vartheta}_2 = m(\mathbf{x}) = \tilde{\mathbf{x}}^\top \boldsymbol{\beta}$. The results presented in Section 3, including Theorems 1, 2 and 3, carry over in the fixed design case when \mathbf{a} is replaced by \mathbf{c} .

In the rest of this section, we will present classical models that can be embedded in the larger class of conditional transformation models and some novel models that can be implemented in this general framework.

4.3. Classical Transformation Models

Linear Model The normal linear regression model $Y \sim N(\alpha + m(\mathbf{x}), \sigma^2)$ with conditional distribution function

$$F_{Y|\mathbf{X}=\mathbf{x}}(y) = \Phi \left(\frac{y - (\alpha + m(\mathbf{x}))}{\sigma} \right)$$

can be understood as a transformation model with transformation function $h(y | \mathbf{x}) = y/\sigma - \alpha/\sigma - m(\mathbf{x})/\sigma$ parameterised via basis functions $\mathbf{a}(y) = (y, 1)^\top$, $\mathbf{b}(\mathbf{x}) = \tilde{\mathbf{x}}$ and $\mathbf{c} = (\mathbf{a}^\top, \mathbf{b}^\top)^\top$ with parameters $\boldsymbol{\vartheta} = (\sigma^{-1}, -\sigma^{-1}\alpha, -\sigma^{-1}\boldsymbol{\beta}^\top)^\top$ under the constraint $\sigma > 0$, or in more compact notation $(\Phi, (y, 1, \tilde{\mathbf{x}}^\top)^\top, \boldsymbol{\vartheta})$. The parameters of the model are the inverse standard deviation and the inverse negative coefficient of variation instead of the mean and variance of the original normal distribution. For “exact continuous” observations, the likelihood \mathcal{L} is equivalent to least-squares, which can be maximised with respect to α and $\boldsymbol{\beta}$ without taking σ into account. This is not possible for censored or truncated observations, where we need to evaluate the conditional distribution function that depends on all parameters; this model is called Type I Tobit model (Tobin 1958) (although only the likelihood changes under censoring and truncation, but the model does not). Using an alternative basis function \mathbf{c} would allow arbitrary non-normal conditional distributions of Y and the simple shift model $\mathbf{c}(y, \mathbf{x})^\top \boldsymbol{\vartheta} = \mathbf{a}(y)^\top \boldsymbol{\vartheta}_1 + \mathbf{b}(\mathbf{x})^\top \boldsymbol{\vartheta}_2$ is then a generalisation of additive models and leads to the interpretation

$$\mathbb{E}_{Y|\mathbf{X}=\mathbf{x}}(\mathbf{a}(Y)^\top \boldsymbol{\vartheta}_1) = -\mathbf{b}(\mathbf{x})^\top \boldsymbol{\vartheta}_2.$$

The choice $\mathbf{a} = (1, \log)^\top$ implements the log-normal model for $Y > 0$. Implementation of a Bernstein basis $\mathbf{a} = \mathbf{a}_{\text{Bs}, M}$ allows arbitrarily shaped distributions, *i.e.* a transition from the normal family to the transformation family, and thus likelihood inference on $\boldsymbol{\vartheta}_2$ without strict assumptions on the distribution of Y . The transformation $\mathbf{a}_{\text{Bs}, M}(y)^\top \boldsymbol{\vartheta}_1$ must increase monotonically in y . Maximisation of the log-likelihood under the linear inequality constraint $\mathbf{D}_{M+1} \boldsymbol{\vartheta}_1 > 0$, with \mathbf{D}_{M+1} representing first-order differences, implements this requirement.

Continuous “Survival Time” Models For a continuous response $Y > 0$, the model

$$F_{Y|\mathbf{X}=\mathbf{x}}(y) = F_Z \left(\frac{\log(y) - (\alpha + m(\mathbf{x}))}{\sigma} \right)$$

with basis functions $\mathbf{a}(y) = (1, \log(y))$ and $\mathbf{b}(\mathbf{x}) = \tilde{\mathbf{x}}$ and parameters $\boldsymbol{\vartheta} = (-\alpha, \sigma^{-1}, -\boldsymbol{\beta}^\top)^\top$ under the constraint $\sigma > 0$ is called the accelerated failure time (AFT) model. The model $(F_{\text{MEV}}, (1, \log, \tilde{\mathbf{x}}^\top)^\top, (\vartheta_1, 1, -\boldsymbol{\beta}^\top)^\top)$ with $\sigma \equiv 1$ (and thus fixed transformation function \log) is the exponential AFT model because it implies an exponential distribution of Y . When the parameter $\sigma > 0$ is estimated from the data, the model $(F_{\text{MEV}}, (1, \log, \tilde{\mathbf{x}}^\top)^\top, \boldsymbol{\vartheta})$ is called the Weibull model, $(F_{\text{SL}}, (1, \log, \tilde{\mathbf{x}}^\top)^\top, \boldsymbol{\vartheta})$ is the log-logistic AFT model and $(\Phi, (1, \log, \tilde{\mathbf{x}}^\top)^\top, \boldsymbol{\vartheta})$ is the log-normal AFT model. For a continuous (not necessarily positive) response Y , the model $F_{Y|\mathbf{X}=\mathbf{x}}(y) = F_{\text{MEV}}(h_Y(y) - m(\mathbf{x}))$ is called the proportional hazards, relative risk or Cox model. The transformation function h_Y equals the log-cumulative baseline hazard and is treated as a nuisance parameter in the partial likelihood framework, where only the regression coefficients $\boldsymbol{\beta}$ are estimated. Given $\hat{\boldsymbol{\beta}}$, non-parametric maximum likelihood estimators are typically applied to obtain \hat{h}_Y . Here, we parameterise this function as $h_Y(y) = \log(\Lambda_Y(y)) = \mathbf{a}(y)^\top \boldsymbol{\vartheta}_1$ (for example, using $\mathbf{a} = \mathbf{a}_{\text{Bs},M}$) and fit all parameters in the model $(F_{\text{MEV}}, (\mathbf{a}^\top, \tilde{\mathbf{x}}^\top)^\top, (\boldsymbol{\vartheta}_1^\top, -\boldsymbol{\beta}^\top)^\top)$ simultaneously. The model is highly popular because $m(\mathbf{x})$ is the log-hazard ratio to $m(\mathbf{0})$. For the special case of right-censored survival times, this parameterisation of the Cox model was studied theoretically and empirically by [McLain and Ghosh \(2013\)](#). Changing the distribution function in the Cox model from F_{MEV} to F_{SL} results in the proportional odds model $(F_{\text{SL}}, (\mathbf{a}^\top, \tilde{\mathbf{x}}^\top)^\top, (\boldsymbol{\vartheta}_1^\top, -\boldsymbol{\beta}^\top)^\top)$; its name comes from the interpretation of $m(\mathbf{x})$ as the constant log-odds ratio of the odds $O_Y(y | \mathbf{X} = \mathbf{x})$ and $O_Y(y | \mathbf{x} = \mathbf{0})$. An additive hazards model with the conditional hazard function $\lambda_Y(y | \mathbf{X} = \mathbf{x}) = \lambda_Y(y | \mathbf{X} = \mathbf{0}) - \tilde{\mathbf{x}}^\top \boldsymbol{\beta}$ results from the choice $F_Z(z) = F_{\text{Exp}}(z) = 1 - \exp(-z)$ ([Aranda-Ordaz 1983](#)) under the additional constraint $\lambda_Y(y | \mathbf{X} = \mathbf{x}) > 0$. In this case, the function $\mathbf{a}(y)^\top \boldsymbol{\vartheta}_1 > 0$ is the positive baseline cumulative hazard function $\Lambda_Y(y | \mathbf{X} = \mathbf{0})$.

Discrete Models For ordered categorical responses $y_1 < \dots < y_K$, the conditional distribution $F_{Y|\mathbf{X}=\mathbf{x}}(y_k) = F_Z(h_Y(y_k) - m(\mathbf{x}))$ is a transformation model with $\mathbf{a}(y_k) = \mathbf{e}_{K-1}(k)$. The model $(F_{\text{SL}}, (\mathbf{a}^\top, \tilde{\mathbf{x}}^\top)^\top, (\boldsymbol{\vartheta}_1^\top, -\boldsymbol{\beta}^\top)^\top)$ is called the discrete proportional odds model and $(F_{\text{MEV}}, (\mathbf{a}^\top, \tilde{\mathbf{x}}^\top)^\top, (\boldsymbol{\vartheta}_1^\top, -\boldsymbol{\beta}^\top)^\top)$ is the discrete proportional hazards model. Here, $m(\mathbf{x})$ is the log-odds ratio or log-hazard ratio to $m(\mathbf{0})$ independent of k ; details are given in [Tutz \(2012\)](#). For the special case of a binary response ($K = 2$), the transformation model $(F_{\text{SL}}, (\mathbf{1}(k = 1), \tilde{\mathbf{x}}^\top)^\top, (\vartheta_1, -\boldsymbol{\beta}^\top)^\top)$ is the logistic regression model, $(\Phi, (\mathbf{1}(k = 1), \tilde{\mathbf{x}}^\top)^\top, (\vartheta_1, -\boldsymbol{\beta}^\top)^\top)$ is the probit model and $(F_{\text{MEV}}, (\mathbf{1}(k = 1), \tilde{\mathbf{x}}^\top)^\top, (\vartheta_1, -\boldsymbol{\beta}^\top)^\top)$ is called the complementary log-log model. Note that the transformation function h_Y is given by the basis function $\mathbf{a} = \mathbf{1}(k = 1)$, *i.e.* ϑ_1 is just the intercept. The connection between standard binary regression models and transformation models is explained in more detail by [Doksum and Gasko \(1990\)](#).

Linear Transformation Model The transformation model $(F_Z, (\mathbf{a}^\top, \tilde{\mathbf{x}}^\top)^\top, (\boldsymbol{\vartheta}_1^\top, -\boldsymbol{\beta}^\top)^\top)$ for any \mathbf{a} and F_Z is called the linear transformation model and contains all models discussed in this subsection. Note that the transformation of the response $h_Y(y) = \mathbf{a}(y)^\top \boldsymbol{\vartheta}_1$ is non-linear in all models of interest (AFT, Cox, *etc.*), and the term “linear” only refers to a linear shift $m(\mathbf{x})$ of the explanatory variables. Partially linear or additive transformation models allow non-linear shifts as part of a partially smooth basis \mathbf{b} , *i.e.* in the form of an additive model. The number of constraints only depends on the basis \mathbf{a} but not on the explanatory variables.

4.4. Extension of Classical Transformation Models

A common property of all classical transformation models is the additivity of the response transformation and the shift, *i.e.* the decomposition $h(y | \mathbf{x}) = h_Y(y) + h_{\mathbf{x}}(\mathbf{x})$ of the conditional transformation function. This assumption is relaxed by the following extensions of the classical models. Allowing for deviations from this simple model is also the key aspect for the development of novel transformation models in the rest of this section.

Discrete Non-Proportional Odds and Hazards Models For ordered categorical responses, the model $F_{Y|\mathbf{X}=\mathbf{x}}(y_k) = F_Z(h_Y(y_k) - m_k(\mathbf{x}))$ allows a category-specific shift $m_k(\mathbf{x}) = \tilde{\mathbf{x}}^\top \boldsymbol{\beta}_k$; with F_{SL} , this cumulative model is called the non-proportional odds model, and with F_{MEV} , it is the non-proportional hazards model. Both models can be cast into the transformation model framework by defining the joint basis $\mathbf{c}(y_k, \mathbf{x}) = (\mathbf{a}(y_k)^\top, \mathbf{a}(y_k)^\top \otimes \mathbf{b}(\mathbf{x})^\top)^\top$ as the Kronecker product of the two simple basis functions $\mathbf{a}(y_k) = \mathbf{e}_{K-1}(k)$ and $\mathbf{b}(\mathbf{x}) = \tilde{\mathbf{x}}$ (assuming that \mathbf{b} does not contain an intercept term). Note that the conditional transformation function $h(y | \mathbf{x})$ includes an interaction term between y and \mathbf{x} . It is also worth noting that for unordered categorical responses $Y \in \Xi = \{y_1, \dots, y_K\}$, the multinomial model can be estimated by the model $(F_Z, \mathbf{c}, \boldsymbol{\vartheta})$ under any ordering of the response categories because the corresponding conditional density

$$f_{Y|\mathbf{X}=\mathbf{x}}(y_k) = \begin{cases} F_Z(\mathbf{c}(y_k, \mathbf{x})^\top \boldsymbol{\vartheta}) & k = 1 \\ F_Z(\mathbf{c}(y_k, \mathbf{x})^\top \boldsymbol{\vartheta}) - F_Z(\mathbf{c}(y_{k-1}, \mathbf{x})^\top \boldsymbol{\vartheta}) & 1 < k < K \\ 1 - F_Z(\mathbf{c}(y_{k-1}, \mathbf{x})^\top \boldsymbol{\vartheta}) & k = K \end{cases}$$

is invariant with respect to the ordering applied. For F_{SL} , this model is called the cumulative logit or partial proportional odds model (Tutz 2012). The classical multinomial logit (with $F_Z = F_{\text{SL}}$) or probit (with $F_Z = \Phi$) models for an unordered response $Y \in \Xi = \{y_1, \dots, y_K\}$ can be written as models for the density

$$\begin{aligned} \mathbb{P}(Y = y_1 | \mathbf{X} = \mathbf{x}) &= F_Z(\tilde{\mathbf{x}}^\top \boldsymbol{\beta}_1) \\ \mathbb{P}(Y = y_k | \mathbf{X} = \mathbf{x}) &= F_Z(\tilde{\mathbf{x}}^\top \boldsymbol{\beta}_k) - F_Z(\tilde{\mathbf{x}}^\top \boldsymbol{\beta}_{k-1}) \quad 1 < k < K \\ \mathbb{P}(Y = y_K | \mathbf{X} = \mathbf{x}) &= 1 - F_Z(\tilde{\mathbf{x}}^\top \boldsymbol{\beta}_{K1}), \end{aligned}$$

where the parameters $\boldsymbol{\beta}_k$ correspond to parameters in a cumulative model $\mathbb{P}(Y \leq y_k) = F_Z(\tilde{\mathbf{x}}^\top \boldsymbol{\beta}_k)$ for any ordering of the sample space. Of course, the parameter estimates $\hat{\boldsymbol{\beta}}_k$ change when the ordering changes and therefore must not be interpreted directly, but the estimated densities $\hat{\mathbb{P}}(Y = y_k | \mathbf{X} = \mathbf{x})$ are invariant with respect to the ordering applied.

Time-Varying Effects One often studied extension of the Cox model is $F_{Y|\mathbf{X}=\mathbf{x}}(y) = F_Z(h_Y(y) - \tilde{\mathbf{x}}^\top \boldsymbol{\beta}(y))$, where the regression coefficients $\boldsymbol{\beta}(y)$ may change with time y . The Cox model is included with $\boldsymbol{\beta}(y) \equiv \boldsymbol{\beta}$, and the model is often applied to check the proportional hazards assumption. With a smooth parameterisation of time y , for example via $\mathbf{a} = \mathbf{a}_{\text{Bs}, M}$, and linear basis $\mathbf{b}(\mathbf{x}) = \tilde{\mathbf{x}}$, the transformation model $(F_{\text{MEV}}, (\mathbf{a}^\top, \mathbf{a}^\top \otimes \mathbf{b}^\top)^\top, \boldsymbol{\vartheta})$ implements this Cox model with time-varying (linear) effects. This model (with arbitrary F_Z) has also been presented in Foresi and Peracchi (1995) and is called distribution regression in Chernozhukov *et al.* (2013).

4.5. Novel Transformation Models

Due to the broadness of the transformation family, it is straightforward to set up new models for interesting situations by allowing more complex transformation functions $h(y | \mathbf{x})$. We will illustrate this possibility for two simple cases: the independent two-sample situation and regression models for count data. The generic and most complex transformation model is called the conditional transformation model and is explained at the end of this section.

Beyond Shift Effects Assume we observe samples from two groups A and B and want to model the conditional distribution functions $F_{Y|\mathbf{X}=A}(y)$ and $F_{Y|\mathbf{X}=B}(y)$ of the response Y in the two groups. Based on this model, it is often interesting to infer whether the two distributions are equivalent and, if this is not the case, to characterise how they differ. Using an appropriate basis function \mathbf{a} and the basis $\mathbf{b}(\mathbf{x}) = (1, \mathbf{1}(B))^\top$, the model $(F_Z, (\mathbf{a}^\top \otimes \mathbf{b}^\top)^\top, \boldsymbol{\vartheta})$ parameterises the conditional transformation function as $h(y | A) = \mathbf{a}(y)^\top \boldsymbol{\vartheta}_1$ and $h(y | B) = h(y | A) + h_{B-A}(y) = \mathbf{a}(y)^\top \boldsymbol{\vartheta}_1 + \mathbf{1}(B) \mathbf{a}(y)^\top \boldsymbol{\vartheta}_2$. Clearly, the second term is constant zero ($h_{B-A}(y) \equiv 0$) iff the two distributions are equivalent ($F_{Y|\mathbf{X}=A}(y) = F_{Y|\mathbf{X}=B}(y)$ for all y). For the deviation function $h_{B-A}(y) = \mathbf{a}^\top \boldsymbol{\vartheta}_2$, we can apply standard likelihood inference procedures for $\hat{\boldsymbol{\vartheta}}_2$ to construct a confidence band or use a test statistic like $\max(\hat{\boldsymbol{\vartheta}}_2 / \text{se}(\hat{\boldsymbol{\vartheta}}_2))$ to assess deviations from zero. If there is evidence for a group effect, we can use the model to check whether the deviation function is constant, *i.e.* $h_{B-A}(y) \equiv c \neq 0$. In this case, the simpler model $(F_Z, (\mathbf{a}^\top, \mathbf{1}(B))^\top, (\boldsymbol{\vartheta}_1^\top, -\beta)^\top)$ with shift $\beta = -\boldsymbol{\vartheta}_2$ might be easier to interpret. This model actually corresponds to a normal ANOVA model with $F_Z = \Phi$ and $\mathbf{a}(y)^\top = (1, y)^\top$ or the Cox proportional hazards model with $(F_{\text{MEV}}, (\mathbf{a}_{\text{Bs},M}^\top, \mathbf{1}(B))^\top, (\boldsymbol{\vartheta}_1^\top, -\beta)^\top)$.

Count Regression “Without Tears” Simple models for count data $\Xi = \{0, 1, 2, \dots\}$ almost always suffer from over-dispersion or excess zeros. The linear transformation model

$$F_{Y|\mathbf{X}=\mathbf{x}}(y) = F_Z(h_Y(y) - m(\mathbf{x}))$$

can be implemented using the basis function $\mathbf{a}(y) = \mathbf{a}_{\text{Bs},M}(\lfloor y \rfloor)$, and then the parameters of the transformation model $(F_Z, (\mathbf{a}^\top, \tilde{\mathbf{x}}^\top)^\top, \boldsymbol{\vartheta})$ are not affected by over- or under-dispersion because higher moments are handled by h_Y independently of the effects of the explanatory variables $m(\mathbf{x})$. If there are excess zeros, we can set up a joint transformation model

$$F_{Y|\mathbf{X}=\mathbf{x}}(y) = F_Z(h_Y(y) - m(\mathbf{x}) + \mathbf{1}(y=0)(\alpha_0 - m_0(\mathbf{x})))$$

such that we have a two-components mixture model consisting of the count distribution

$$F_{Y|\mathbf{X}=\mathbf{x}}(y) = F_Z(h_Y(y) - m(\mathbf{x})), y \in \Xi$$

and the probability of an excess zero

$$f_{Y|\mathbf{X}=\mathbf{x}}(0) = F_Z(h_Y(0) - m(\mathbf{x}) + (\alpha_0 - m_0(\mathbf{x}))) = F_Z(h_Y(0) + \alpha_0 - \tilde{\mathbf{x}}^\top (\boldsymbol{\beta} + \boldsymbol{\beta}_0))$$

when $m_0(\mathbf{x}) = \tilde{\mathbf{x}}^\top \boldsymbol{\beta}_0$. Hence, the transformation analogue to a hurdle model with hurdle at zero is the transformation model $(F_Z, (\mathbf{a}^\top, \tilde{\mathbf{x}}^\top, \mathbf{1}(y=0), \mathbf{1}(y=0)\tilde{\mathbf{x}}^\top)^\top, (\boldsymbol{\vartheta}_1^\top, \boldsymbol{\beta}^\top, \alpha_0, \boldsymbol{\beta}_0^\top)^\top)$.

Conditional Transformation Models When the conditional transformation function is parameterised by multiple basis functions $\mathbf{a}_j(y), \mathbf{b}_j(\mathbf{x}), j = 1, \dots, J$ via the joint basis

$$\mathbf{c} = (\mathbf{a}_1^\top \otimes \mathbf{b}_1^\top, \dots, \mathbf{a}_J^\top \otimes \mathbf{b}_J^\top)^\top,$$

models of the class $(\cdot, \mathbf{c}, \boldsymbol{\vartheta})$ are called conditional transformation models (CTMs) with J partial transformation functions parameterised as $\mathbf{a}_j^\top \otimes \mathbf{b}_j^\top$ and include all special cases discussed in this section. It is convenient to assume monotonicity for each of the partial transformation functions; thus, the linear constraints for \mathbf{a}_j are repeated for each basis function in \mathbf{b}_j (detailed descriptions of linear constraints for different models in this class are available in [Hothorn 2017b](#)). [Hothorn et al. \(2014\)](#) introduced this general model class and proposed a boosting algorithm for the estimation of transformation functions h for “exact continuous” responses Y . In the likelihood framework presented here, conditional transformation models can be fitted under arbitrary schemes of censoring and truncation and classical likelihood inference for the model parameters $\boldsymbol{\vartheta}$ becomes feasible. Of course, unlike in the boosting context, the number of model terms J and their complexity are limited in the likelihood world because the likelihood does not contain any penalty terms that induce smoothness in the \mathbf{x} -direction. A more detailed overview on the class of conditional transformation models can be found in ([Möst et al. 2016](#)).

A systematic overview of linear transformation models with potentially response-varying effects is given in Table 1. Model nomenclature and interpretation of the corresponding model parameters is mapped to specific transformation functions h and distribution functions F_Z . To the best of our knowledge, models without names have not yet been discussed in the literature, and their specific properties await closer investigation.

5. Empirical Evaluation

We will illustrate the range of possible applications of likelihood-based conditional transformation models. In Subsection 5.2, we will present a small simulation experiment highlighting the possible advantage of indirectly modelling conditional distributions with transformation functions.

5.1. Illustrations

Density Estimation: Old Faithful Geyser The duration of eruptions and the waiting time between eruptions of the Old Faithful geyser in the Yellowstone National Park became a standard benchmark for non-parametric density estimation (the original data were given by [Azzalini and Bowman 1990](#)). The nine parameters of the transformation model $(\Phi, \mathbf{a}_{\text{Bs},8}(\text{waiting}), \boldsymbol{\vartheta})$ were fitted by maximisation of the approximate log-likelihood (treating the waiting times as “exact” observations) under the eight linear constraints $\mathbf{D}_9 \boldsymbol{\vartheta} > 0$. The model depicted in Figure 1A reproduces the classic bimodal unconditional density of waiting time along with a kernel density estimate. It is important to note that the transformation model was fitted likelihood-based, whereas the kernel density estimate relied on a cross-validated bandwidth. An unconditional density estimate for the duration of the eruptions needs to deal with censoring because exact duration times are only available for the day-time measurements. At night, the observations were either left-censored (“short” eruption),




Figure 1: Old Faithful Geyser. Estimated density for waiting times (A) between and duration (B) of eruptions by the most likely transformation model (MLT) and kernel smoothing (function `npudens()` from package `np` in R). Note that the kernel estimator was based on the imputed duration times 2, 3 and 4 for short, medium and long eruptions at night (as are the rugs in B).

interval-censored (“medium” eruption) or right-censored (“long” eruption) as explained by [Azalini and Bowman \(1990\)](#). This censoring was widely ignored in analyses of the Old Faithful data because most non-parametric kernel techniques cannot deal with censoring (see for example [Hyndman and Yao 2002](#)). We applied the transformation model $(\Phi, \mathbf{a}_{Bs,8}(\text{duration}), \boldsymbol{\vartheta})$ based on the exact log-likelihood function under eight linear constraints and obtained the unconditional density depicted in Figure 1B. In [Hothorn \(2017b\)](#), results for $M = 40$ are computed, which led to almost identical estimates of the distribution function.

In addition, we modelled the conditional distribution of the censored duration times given the waiting times using the transformation model $(\Phi, (\mathbf{a}_{Bs,7}(\text{duration})^\top \otimes \mathbf{b}_{Bs,3}(\text{waiting})^\top)^\top, \boldsymbol{\vartheta})$. This conditional transformation model allows a smooth conditional distribution function of duration smoothly varying with waiting time and was fitted by maximisation of the exact log-likelihood under 7×4 linear constraints. The corresponding conditional density in Figure 2 shows that the marginal bimodality can be explained by relatively long durations for short waiting times and two clusters of short and long durations after waiting times longer than 70 minutes. Note that in this model, the choice of F_Z does not influence the estimated conditional distribution or density function if the parameterisation of h is flexible enough to compensate for this change.




Figure 2: Old Faithful Geysers. Conditional distribution of duration given waiting time estimated with a most likely transformation model (MLT) and a conditional kernel density estimator (function `npcdens()` from package `np`). Exact observations are given as dots, censored observations as lines.

Quantile Regression: Head Circumference The Fourth Dutch Growth Study ([Fredriks et al. 2000](#)) is a cross-sectional study on growth and development of the Dutch population younger than 22 years. [Stasinopoulos and Rigby \(2007\)](#) fitted a growth curve to head circumferences (HC) of 7040 boys using a GAMLSS model with a Box-Cox t distribution describing the first four moments of head circumference conditionally on age. The model showed evidence of kurtosis, especially for older boys. We fitted the same growth curves by the conditional transformation model $(\Phi, (\mathbf{a}_{\text{Bs},3}(\text{HC})^\top \otimes \mathbf{b}_{\text{Bs},3}(\text{age}^{1/3})^\top)^\top, \boldsymbol{\vartheta})$ by maximisation of the approximate log-likelihood under 3×4 linear constraints $(\mathbf{D}_4 \otimes \mathbf{I}_4)\boldsymbol{\vartheta} > 0$. Figure 3 shows the data overlaid with quantile curves obtained via inversion of the estimated conditional distributions. The figure very closely reproduces the growth curves presented in Figure 16 of [Stasinopoulos and Rigby \(2007\)](#) and also indicates a certain asymmetry towards older boys.

Survival Analysis: German Breast Cancer Study Group-2 Trial This prospective, controlled clinical trial on the treatment of node-positive breast cancer patients was conducted by the German Breast Cancer Study Group (GBSG-2, [Schumacher et al. 1994](#)). Patients not older than 65 years with positive regional lymph nodes but no distant metastases were included in the study. Out of 686 women, 246 received hormonal therapy, whereas the control group of 440 women did not. Additional variables include age, menopausal status, tumour size, tumour




Figure 3: Head Circumference Growth. Observed head circumference and age of 7040 boys with estimated quantile curves for $p = 0.04, 0.02, 0.1, 0.25, 0.5, 0.75, 0.9, 0.98, 0.996$.

grade, number of positive lymph nodes, progesterone receptor and oestrogen receptor. The right-censored recurrence-free survival time is the response variable of interest.

The Cox model $(F_{\text{MEV}}, (\mathbf{a}_{\text{Bs},10}^\top, \mathbf{1}(\text{hormonal therapy}))^\top, \boldsymbol{\vartheta})$ implements the transformation function $h(y \mid \text{treatment}) = \mathbf{a}_{\text{Bs},10}(y)^\top \boldsymbol{\vartheta}_1 + \mathbf{1}(\text{hormonal therapy})\beta$, where $\mathbf{a}_{\text{Bs},10}^\top \boldsymbol{\vartheta}_1$ is the log-cumulative baseline hazard function parameterised by a Bernstein polynomial and $\beta \in \mathbb{R}$ is the log-hazard ratio of hormonal therapy. This is the classical Cox model with one treatment parameter β but with fully parameterised baseline transformation function, which was fitted by the exact log-likelihood under ten linear constraints. The model assumes proportional hazards, an assumption whose appropriateness we wanted to assess using the non-proportional hazards model $(F_{\text{MEV}}, (\mathbf{a}_{\text{Bs},10}^\top \otimes (1, \mathbf{1}(\text{hormonal therapy})))^\top, \boldsymbol{\vartheta})$ with the transformation function

$$h(y \mid \text{treatment}) = \mathbf{a}_{\text{Bs},10}(y)^\top \boldsymbol{\vartheta}_1 + \mathbf{1}(\text{hormonal therapy})\mathbf{a}_{\text{Bs},10}(y)^\top \boldsymbol{\vartheta}_2.$$

The function $\mathbf{a}_{\text{Bs},10}(y)^\top \boldsymbol{\vartheta}_2$ is the time-varying difference of the log-hazard functions of women without and with hormonal therapy and can be interpreted as the deviation from a constant log-hazard ratio treatment effect of hormonal therapy. Under the null hypothesis of no treatment effect, we would expect $\boldsymbol{\vartheta}_2 \equiv \mathbf{0}$. This monotonic deviation function adds ten linear constraints $\mathbf{D}_{11}\boldsymbol{\vartheta}_1 + \mathbf{D}_{11}\boldsymbol{\vartheta}_2 > \mathbf{0}$, which also ensure monotonicity of the transformation function for treated patients. We first compared the fitted survivor functions obtained from the model including a time-varying treatment effect with the Kaplan-Meier estimators in both




Figure 4: GBSG-2. Estimated survivor functions by the most likely transformation model (MLT) and the Kaplan-Meier (KM) estimator in the two treatment groups (A). Verification of proportional hazards (B): The log-hazard ratio $\hat{\beta}$ (dashed line) with 95% confidence interval (dark grey) is fully covered by a 95% confidence band for the time-varying treatment effect (the time-varying log-hazard ratio is in light grey, the estimate is the solid line) computed from a non-proportional hazards model.

treatment groups. Figure 4A shows a nicely smoothed version of the survivor functions obtained from this transformation model. Figure 4B shows the time-varying treatment effect $\mathbf{a}_{\text{Bs},10}(y)^\top \hat{\boldsymbol{\vartheta}}_2$, together with a 95% confidence band computed from the joint normal distribution of $\hat{\boldsymbol{\vartheta}}_2$ for a grid over time as described by Hothorn *et al.* (2008); the method is much simpler than other methods for inference on time-varying effects (*e.g.* Sun *et al.* 2009). The 95% confidence interval around the log-hazard ratio $\hat{\beta}$ is also plotted, and as the latter is fully covered by the confidence band for the time-varying treatment effect, there is no reason to question the treatment effect computed under the proportional hazards assumption. An alternative method for this type of analysis has been recently suggested by Yang and Prentice (2015).

In the second step, we allowed an age-varying treatment effect to be included in the model $(F_{\text{MEV}}, (\mathbf{a}_{\text{Bs},10}(y)^\top \otimes (\mathbf{1}(\text{hormonal therapy}), 1 - \mathbf{1}(\text{hormonal therapy})) \otimes \mathbf{b}_{\text{Bs},3}(\text{age})^\top)^\top, \boldsymbol{\vartheta})$. For both treatment groups, we estimated a conditional transformation function of survival time y given age parameterised as the tensor basis of two Bernstein bases. Each of the two basis functions comes with 10×3 linear constraints; therefore, the model was fitted under 60 linear constraints. Figure 5 allows an assessment of the prognostic and predictive properties of




Figure 5: GBSG-2. Prognostic and predictive effect of age. The contours depict the conditional survivor functions given treatment and age of the patient.

age. As the survivor functions were clearly larger for all patients treated with hormones, the positive treatment effect applied to all patients. However, the size of the treatment effect varied greatly. The effect was most pronounced for women younger than 30 and levelled off a little for older patients. In general, the survival times were longest for women between 40 and 60 years old. Younger women suffered the highest risk; for women older than 60 years, the risk started to increase again. This effect was shifted towards younger women when hormonal treatment was applied.

Count Regression: Tree Pipit Counts Müller and Hothorn (2004) reported data on the number of tree pipits *Anthus trivialis*, a small passerine bird, counted on 86 forest plots in a light gradient ranging from open and sunny stands (small cover storey) to dense and dark stands (large cover storey). We modelled the conditional distribution of the number of tree pipits on one plot given the cover storey on this plot by the transformation model $(\Phi, (\mathbf{a}^\top \otimes \mathbf{b}_{\text{Bs},4}(\text{cover storey})^\top)^\top, \boldsymbol{\vartheta})$, where $\mathbf{a}(y) = \mathbf{e}_5(y + 1)$, $y = 0, \dots, 4$; the model was fitted under 4×5 linear constraints. In this model of count data, the conditional distribution depends on both the number of counted birds and the cover storey and the effect of cover storey may change with different numbers of birds observed. Figure 6A depicts the observations, and Figure 6B shows the conditional distribution function evaluated for $0, \dots, 5$ observed birds. The conditional distribution function obtained from a generalised additive Poisson (GAM) model with smooth mean effect of cover storey is given in Figure 6C. Despite some




Figure 6: Tree Pipit Counts. Observations (A, the size of the points is proportional to the number of observations) and estimated conditional distribution of number of tree pipits given cover storey by the most likely transformation model (MLT, B) and a generalised additive Poisson model (function `gam()` in package `mgcv`, GAM, C).

overfitting, this model is more restrictive than our transformation model because one mean function determines the whole distribution (the local minima of the conditional distributions as a function of cover storey were constant in Figure 6C, whereas they were shifted towards higher values of cover storey in Figure 6B).

5.2. Simulation Experiment

The transformation family includes linear as well as very flexible models, and we therefore illustrate the potential gain of modelling a transformation function h by comparing a very simple transformation model to a fully parametric approach and to a non-parametric approach using a data-generating process introduced by Hothorn *et al.* (2014).

In the transformation model $(\Phi, ((1, y) \otimes (1, \mathbf{x}^\top))^\top, \boldsymbol{\vartheta})$, two explanatory variables $\mathbf{x} = (x_1, x_2)^\top$ influence both the conditional mean and the conditional variance of a normal response Y . Although the transformation function is linear in y with three linear constraints, the mean and variance of Y given \mathbf{x} depend on \mathbf{x} in a non-linear way. The choices $x_1 \sim U[0, 1]$, $x_2 \sim U[-2, 2]$ with $\boldsymbol{\vartheta} = (0, 0, -1, .5, 1, 0)$ lead to the heteroscedastic varying coefficient model

$$Y = \frac{1}{x_1 + 0.5}x_2 + \frac{1}{x_1 + 0.5}\varepsilon, \quad \varepsilon \sim N(0, 1), \quad (5)$$

where the variance of Y ranges between 0.44 and 4 depending on x_1 . This model can be fitted in the GAMLSS framework under the assumptions that the mean of the normal response depends on a smoothly varying regression coefficient $(x_1 + 0.5)^{-1}$ for x_2 and that the variance is a smooth function of x_1 . This model is therefore fully parametric. As a non-parametric

counterpart, we used a kernel estimator for estimating the conditional distribution function of Y as a function of the two explanatory variables.

From the transformation model, the GAMLSS and kernel estimators, we obtained estimates of $F_{Y|\mathbf{X}=\mathbf{x}}(y)$ over a grid on y, x_1, x_2 and computed the mean absolute deviation (MAD) of the true and estimated probabilities

$$\text{MAD}(x_1, x_2) = \frac{1}{n} \sum_y |F_{Y|\mathbf{X}=\mathbf{x}}(y) - \hat{F}_{Y|\mathbf{X}=\mathbf{x},N}(y)|$$

for each pair of x_1 and x_2 . Then, the minimum, the median and the maximum of the MAD values for all x_1 and x_2 were computed as summary statistics. The most likely transformation approach and its two competitors were estimated and evaluated for 100 random samples of size $N = 200$ drawn from model (5). Cross-validation was used to determine the bandwidths for the kernel-based estimators (function `npcdist()` in package `np`; for details, see Hayfield and Racine 2008). We fitted the GAMLSS models by boosting; the number of boosting iterations was determined via sample splitting (Mayr *et al.* 2012). To investigate the stability of the three procedures under non-informative explanatory variables, we added to the data $p = 1, \dots, 5$ uniformly distributed variables without association to the response and included them as potential explanatory variables in the three models. The case $p = 0$ corresponds to model (5).

Figure 7 shows the empirical distributions of the minimum, median and maximum MAD for the three competitors. Except for the minimum MAD in the absence of any irrelevant explanatory variables ($p = 0$), the conditional distributions fitted by the transformation models were closer to the true conditional distribution function by means of the MAD. This result was obtained because the transformation model only had to estimate a simple transformation function, whereas the other two procedures had a difficult time approximating this simple transformation model on another scale. However, the comparison illustrates the potential improvement one can achieve when fitting simple models for the transformation function instead of more complex models for the mean (GAMLSS) or distribution function (Kernel). The kernel estimator led to the largest median MAD values but seemed more robust than GAMLSS with respect to the maximum MAD. These results were remarkably robust in the presence of up to five non-informative explanatory variables, although of course the MAD increased with the number of non-informative variables p .

6. Discussion

The contribution of a likelihood approach for the general class of conditional transformation models is interesting both from a theoretical and a practical perspective. With the range of simple to very complex transformation functions introduced in Section 4 and illustrated in Section 5, it becomes possible to understand classical parametric, semi-parametric and non-parametric models as special cases of the same model class. Thus, analytic comparisons between models of different complexity become possible. The transformation family $\mathbb{P}_{Y,\Theta}$, the corresponding likelihood function and the most likely transformation estimator are easy to understand. This makes the approach appealing also from a teaching perspective. Connections between standard parametric models (for example, the normal linear model) and potentially complex models for survival or ordinal data can be outlined in very simple notation, placing




Figure 7: Empirical Evaluation. Minimum, median and maximum of the mean absolute deviation (MAD) between true and estimated probabilities for most likely transformation models (MLT), non-parametric kernel distribution function estimation (Kernel) and generalised additive models for location, scale and shape (GAMLSS) for 100 random samples. Values on the ordinate can be interpreted as absolute differences of probabilities. The grey horizontal lines correspond to the median of MLT.

emphasis on the modelling of (conditional) distributions instead of just modelling (conditional) means. Computationally, the log-likelihood $\log \circ \mathcal{L}$ is linear in the number of observations N and, for contributions of “exact continuous” responses, only requires the evaluation of the derivative h' of the transformation function h instead of integrals thereof. Standard optimisers for linearly constrained problems can be applied, and AIC- or BIC-based model selection is possible. Transformation models directly suggest a way to specify conceptual $(F_Z, \mathbf{c}, \boldsymbol{\vartheta})$ and fitted $(F_Z, \mathbf{c}, \hat{\boldsymbol{\vartheta}})$ models in computer languages because “only” the conditional transformation function $h = \mathbf{c}^\top \boldsymbol{\vartheta}$ has to be specified in addition to a “simple” distribution function F_Z (Hothorn 2017a,b).

The results presented in Section 5 are based on only roughly 1000 lines of R code. We tested this reference implementation of most likely transformations against all linear transformation models available in R to date and obtained equivalent regression coefficients $\hat{\boldsymbol{\beta}}$ and their corresponding covariance matrices. Thus, the framework helps to reduce the code base and thereby helps to considerably reduce the number of possible errors in implementations of linear transformation models, while at the same time it allows novel and more complex models to be fitted (Hothorn 2017a,b).

Based on the general understanding of transformation models outlined in this paper, it will be interesting to study these models outside the strict likelihood world. A mixed transformation model for cluster data (Cai *et al.* 2002; Huber-Carol and Vonta 2004; Zeng *et al.* 2005; Choi and Huang 2012) is often based on the transformation function $h(y | \mathbf{x}, i) = h_Y(y) + \delta_i + h_{\mathbf{x}}(\mathbf{x})$ with random intercept (or “frailty” term) δ_i for the i th observational unit. Conceptually, a more complex deviation from the global model could be formulated as $h(y | \mathbf{x}, i) = h_Y(y) + h_Y(y, i) + h_{\mathbf{x}}(\mathbf{x})$, *i.e.* each observational unit is assigned its own “baseline” transformation $h_Y(y) + h_Y(y, i)$, where the second term is an integral zero deviation from h_Y . For longitudinal data with possibly time-varying explanatory variables, the model $h(y | \mathbf{x}(t), t) = h_Y(y, t) + \mathbf{x}(t)\boldsymbol{\beta}(t)$ (Ding *et al.* 2012; Wu and Tian 2013) can also be understood as a mixed version of a conditional transformation model. The penalised log-likelihood $\log(\mathcal{L}(h | y)) - \text{pen}(\boldsymbol{\beta})$ for the linear transformation model $h(y | \mathbf{x}) = h_Y(y) - \tilde{\mathbf{x}}^\top \boldsymbol{\beta}$ leads to Ridge- or Lasso-type regularised models, depending on the form of the penalty term. Priors for all model parameters $\boldsymbol{\vartheta}$ allow a fully Bayesian treatment of transformation models. Instead of the relatively strict model assumption $F_Y = F_Z \circ \mathbf{a}^\top \boldsymbol{\vartheta}$, one could of course allow arbitrary unknown transformation functions $h \in \mathcal{H}$ in the spirit of non-parametric regression and study the quality of the approximation $\mathbf{a}^\top \hat{\boldsymbol{\vartheta}}$ for h for the fixed and random design case.

It is possible to relax the assumption that F_Z is known. The simultaneous estimation of F_Z in the model $\mathbb{P}(Y \leq y | \mathbf{X} = \mathbf{x}) = F_Z(h_Y(y) - \tilde{\mathbf{x}}^\top \boldsymbol{\beta})$ was studied by Horowitz (1996) and later extended by Linton *et al.* (2008) to non-linear functions $h_{\mathbf{x}}$ with parametric baseline transformation h_Y and kernel estimates for F_Z and $h_{\mathbf{x}}$. For AFT models, Zhang and Davidian (2008) applied smooth approximations for the density f_Z in an exact censored likelihood estimation procedure. In a similar setup, Huang (2014) proposed a method to jointly estimate the mean function and the error distribution in a generalised linear model. The estimation of F_Z is noteworthy in additive models of the form $h_Y + h_{\mathbf{x}}$ because these models assume additivity of the contributions of y and \mathbf{x} on the scale of $F_Z^{-1}(\mathbb{P}(Y \leq y | \mathbf{X} = \mathbf{x}))$. If this model assumption seems questionable, one can either allow unknown F_Z or move to a transformation model featuring a more complex transformation function. From this point of view, the distribution function F_Z in flexible transformation models is only a computational device mapping the unbounded transformation function h into the unit interval strictly

monotonically, making the evaluation of the likelihood easy. Then, F_Z has no further meaning or interpretation as error distribution. A compromise could be the family of distributions $F_Z(z | \rho) = 1 - (1 + \rho \exp(z))^{-1/\rho}$ for $\rho > 0$ (suggested by McLain and Ghosh 2013) with simultaneous maximum likelihood estimation of ϑ and ρ for additive transformation functions $h = h_Y + h_{\mathbf{x}}$, as these models are flexible and still relatively easy to interpret.

Another comment concerns the connection between transformation models and quantile regression. For some probability $p \in [0, 1]$, the conditional quantile function is

$$Q_Y(p | \mathbf{X} = \mathbf{x}) := F_Y^{-1}(p | \mathbf{X} = \mathbf{x}) = \inf\{y \in \Xi | F_Z(h(y | \mathbf{x})) \geq p\},$$

and for absolutely continuous Y , we then get $Q_Y(p | \mathbf{X} = \mathbf{x}) = h^{-1}(F_Z^{-1}(p) | \mathbf{X} = \mathbf{x})$. A linear quantile regression model (Koenker 2005) assumes $Q_Y(p | \mathbf{X} = \mathbf{x}) = \alpha(p) + \tilde{\mathbf{x}}^\top \boldsymbol{\beta}(p)$, which cannot be written as a linear transformation model. Thus, non-linear transformation functions are necessary in order to achieve the same flexibility as a linear quantile regression model; however, the full conditional distribution function can be estimated in one step, which avoids computational problems such as quantile crossing. In the most complex case of a transformation model that is invariant with respect to the choice of F_Z , there is a correspondingly complex quantile regression model such that both models are equivalent, as was also noted by Chernozhukov *et al.* (2013). In a certain sense, we can understand transformation models as “inverse quantile regression”. A more detailed analysis of the connection between transformation models (called distribution regression there) and quantile regression can be found in Leorato and Peracchi (2015).

In light of the empirical results discussed in this paper and the theoretical work of McLain and Ghosh (2013) on a Cox model with log-cumulative baseline hazard function parameterised in terms of a Bernstein polynomial with increasing order M , one might ask where the boundaries between parametric, semi-parametric and non-parametric statistics lie. The question how the order M affects results practically has been repeatedly raised; therefore, we will close our discussion by looking at a Cox model with increasing M for the GBSG-2 data. All eight baseline variables were included in the linear predictor, and we fitted the model with orders $M = 1, \dots, 30, 35, 40, 45, 50$ of the Bernstein polynomial parameterising the log-cumulative baseline hazard function. In Figure 8A, the log-cumulative baseline hazard functions start with a linear function ($M = 1$) and quickly approach a function that is essentially a smoothed version of the Nelson-Aalen-Breslow estimator plotted in red. In Figure 8B, the trajectories of the estimated regression coefficients become very similar to the partial likelihood estimates as M increased. For $M \geq 10$, for instance, the results of the “semi-parametric” and the “fully parametric” Cox models are practically equivalent. An extensive collection of such head-to-head comparisons of most likely transformations with their classical counterparts can be found in Hothorn (2017b). Our work for this paper and practical experience with its reference software implementation convinced us that rethinking classical models in terms of fully parametric transformations is intellectually and practically a fruitful exercise.

Acknowledgements

Torsten Hothorn received financial support by Deutsche Forschungsgemeinschaft (DFG) under grant number HO 3242/4-1. We thank Karen A. Brune for improving the language.




Figure 8: GBSG-2. Comparison of exact and partial likelihood for order $M = 1, \dots, 30, 35, 40, 45, 50$ of the Bernstein polynomial approximating the log-cumulative baseline hazard function h_Y . The estimated log-cumulative baseline hazard functions for varying M are shown in grey and the Nelson-Aalen-Breslow estimator is shown in red (A). The right panel (B) shows the trajectories of the regression coefficients $\hat{\beta}$ obtained for varying M , which are represented as dots. The horizontal lines represent the partial likelihood estimates.

References

- Alzaatreh A, Lee C, Famoye F (2013). “A New Method for Generating Families of Continuous Distributions.” *Metron*, **71**(1), 63–79. doi:10.1007/s40300-013-0007-y.
- Aranda-Ordaz FJ (1983). “An Extension of the proportional Hazards Model for Grouped Data.” *Biometrics*, **39**(1), 109–117.
- Azzalini A, Bowman AW (1990). “A Look at Some Data on the Old Faithful Geyser.” *Journal of the Royal Statistical Society: Series C (Applied Statistics)*, **39**(3), 357–365. doi:10.2307/2347385.
- Babu GJ, Canty AJ, Chaubey YP (2002). “Application of Bernstein Polynomials for Smooth Estimation of a Distribution and Density Function.” *Journal of Statistical Planning and Inference*, **105**(2), 377–392. doi:10.1016/S0378-3758(01)00265-8.
- Bender R, Augustin T, Blettner M (2005). “Generating Survival Times to Simu-

- late Cox Proportional Hazards Models.” *Statistics in Medicine*, **24**(11), 1713–1723. doi:10.1002/sim.2059.
- Bickel PJ, Klaassen CAJ, Ritov Y, Wellner JA (1993). *Efficient and Adaptive Estimation for Semiparametric Models*. The Johns Hopkins University Press, Baltimore, U.S.A. and London, U.K.
- Box GEP, Cox DR (1964). “An Analysis of Transformations.” *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, **26**(2), 211–252.
- Cai T, Betensky RA (2003). “Hazard Regression for Interval-Censored Data with Penalized Spline.” *Biometrics*, **59**(3), 570–579. doi:10.1111/1541-0420.00067.
- Cai T, Cheng SC, Wei LJ (2002). “Semiparametric Mixed-Effects Models for Clustered Failure Time Data.” *Journal of the American Statistical Association*, **97**(458), 514–522. doi:10.1198/016214502760047041.
- Chang I, Hsiung CA, Wu YJ, Yany CC (2005). “Bayesian Survival Analysis Using Bernstein Polynomials.” *Scandinavian Journal of Statistics*, **32**(3), 447–466. doi:10.1111/j.1467-9469.2005.00451.x.
- Chen K, Jin Z, Ying Z (2002). “Semiparametric Analysis of Transformation Models with Censored Data.” *Biometrika*, **89**(3), 659–668. doi:10.1093/biomet/89.3.659.
- Cheng SC, Wei LJ, Ying Z (1995). “Analysis of Transformation Models with Censored Data.” *Biometrika*, **82**(4), 835–845. doi:10.1093/biomet/82.4.835.
- Cheng SC, Wei LJ, Ying Z (1997). “Predicting Survival Probabilities with Semiparametric Transformation Models.” *Journal of the American Statistical Association*, **92**(437), 227–235. doi:10.1080/01621459.1997.10473620.
- Chernozhukov V, Fernández-Val I, Melly B (2013). “Inference on Counterfactual Distributions.” *Econometrica*, **81**(6), 2205–2268. doi:10.3982/ECTA10582.
- Choi S, Huang X (2012). “A General Class of Semiparametric Transformation Frailty Models for Nonproportional Hazards Survival Data.” *Biometrics*, **68**(4), 1126–1135. doi:10.1111/j.1541-0420.2012.01784.x.
- Cox DR (1975). “Partial Likelihood.” *Biometrika*, **62**(2), 269–276.
- Crowther MJ, Lambert PC (2014). “A General Framework for Parametric Survival Analysis.” *Statistics in Medicine*, **33**(30), 5280–5297. doi:10.1002/sim.6300.
- Curtis SM, Ghosh SK (2011). “A Variable Selection Approach to Monotonic Regression with Bernstein Polynomials.” *Journal of Applied Statistics*, **38**(5), 961–976. doi:10.1080/02664761003692423.
- Ding AA, Tian S, Yu Y, Guo H (2012). “A Class of Discrete Transformation Survival Models With Application to Default Probability Prediction.” *Journal of the American Statistical Association*, **107**(499), 990–1003. doi:10.1080/01621459.2012.682806.
- Doksum KA, Gasko M (1990). “On a Correspondence Between Models in Binary Regression Analysis and in Survival Analysis.” *International Statistical Review*, **58**(3), 243–252.

- Farouki RT (2012). “The Bernstein Polynomial Basis: A Centennial Retrospective.” *Computer Aided Geometric Design*, **29**(6), 379–419. doi:10.1016/j.cagd.2012.03.001.
- Fisher RA (1934). “Two New Properties of Mathematical Likelihood.” *Proceedings of the Royal Society of London. Series A, Containing Papers of a Mathematical and Physical Character*, **53**, 285–306.
- Foresi S, Peracchi F (1995). “The Conditional Distribution of Excess Returns: An Empirical Analysis.” *Journal of the American Statistical Association*, **90**(430), 451–466. doi:10.1080/01621459.1995.10476537.
- Fraser DAS (1968). *The Structure of Inference*. John Wiley & Sons, New York, U.S.A.
- Fredriks AM, van Buuren S, Burgmeijer RJF, Meulmeester JF, Beuker RJ, Brugman E, Roede MJ, Verloove-Vanhorick SP, Wit J (2000). “Continuing Positive Secular Growth Change in The Netherlands 1955–1997.” *Pediatric Research*, **47**(3), 316–323.
- Gneiting T, Katzfuss M (2014). “Probabilistic Forecasting.” *Annual Review of Statistics and Its Application*, **1**(1), 125–151. doi:10.1146/annurev-statistics-062713-085831.
- Hayfield T, Racine JS (2008). “Nonparametric Econometrics: The np Package.” *Journal of Statistical Software*, **27**(5), 1–32. URL <http://www.jstatsoft.org/v27/i05>.
- Hofner B, Mayr A, Fenske N, Schmid M (2016). *gamboostLSS: Boosting Methods for 'GAMLSS'*. R package version 1.2-2, URL <https://CRAN.R-project.org/package=gamboostLSS>.
- Horowitz JL (1996). “Semiparametric Estimation of a Regression Model with an Unknown Transformation of the Dependent Variable.” *Econometrica*, **64**(1), 103–137.
- Hothorn T (2017a). *mlt: Most Likely Transformations*. R package version 0.1-3, URL <https://CRAN.R-project.org/package=mlt>.
- Hothorn T (2017b). *Most Likely Transformations: The mlt Package*. R package vignette version 0.1-5, URL <https://CRAN.R-project.org/package=mlt.docreg>.
- Hothorn T, Bretz F, Westfall P (2008). “Simultaneous Inference in General Parametric Models.” *Biometrical Journal*, **50**(3), 346–363. doi:10.1002/bimj.200810425.
- Hothorn T, Kneib T, Bühlmann P (2014). “Conditional Transformation Models.” *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, **76**(1), 3–27. doi:10.1111/rssb.12017.
- Huang A (2014). “Joint Estimation of the Mean and Error Distribution in Generalized Linear Models.” *Journal of the American Statistical Association*, **109**(505), 186–196. doi:10.1080/01621459.2013.824892.
- Huber-Carol C, Vonta I (2004). “Frailty Models for Arbitrarily Censored and Truncated Data.” *Lifetime Data Analysis*, **10**(4), 369–388. doi:10.1007/s10985-004-4773-y.
- Hyndman RJ, Yao Q (2002). “Nonparametric Estimation and Symmetry Tests for Conditional Density Functions.” *Journal of Nonparametric Statistics*, **14**(3), 259–278. doi:10.1080/10485250212374.

- Joly P, Commenges D, Letenneur L (1998). “A Penalized Likelihood Approach for Arbitrarily Censored and Truncated Data: Application to Age-Specific Incidence of Dementia.” *Biometrics*, **54**(1), 185–194.
- Jones MC, Pewsey A (2009). “Sinh-arcsinh Distributions.” *Biometrika*, **96**(4), 761–780. doi:10.1093/biomet/asp053.
- Klein JP, Moeschberger MK (2003). *Survival Analysis. Techniques for Censored and Truncated Data*. 2nd edition. Springer, New York, U.S.A.
- Koenker R (2005). *Quantile Regression*. Economic Society Monographs. Cambridge University Press, New York, U.S.A.
- Kooperberg C, Clarkson DB (1997). “Hazard Regression with Interval-Censored Data.” *Biometrics*, **53**(4), 1485–1494.
- Kooperberg C, Stone CJ, Truong YK (1995). “Hazard Regression.” *Journal of the American Statistical Association*, **90**(429), 78–94.
- Lehmann EL (1983). *Theory of Point Estimation*. John Wiley & Sons, New York, U.S.A.
- Leorato S, Peracchi F (2015). “Comparing Distribution and Quantile Regression.” *Technical Report 1511*, Einaudi Institute for Economics and Finance, Rome, Italy. URL <https://ideas.repec.org/p/eie/wpaper/1511.html>.
- Lindsey JK (1996). *Parametric Statistical Inference*. Clarendon Press, Oxford, UK.
- Lindsey JK (1999). “Some Statistical Heresies.” *Journal of the Royal Statistical Society: Series D (The Statistician)*, **48**(1), 1–40.
- Linton O, Sperlich S, van Keilegom I (2008). “Estimation of a Semiparametric Transformation Model.” *The Annals of Statistics*, **36**(2), 686–718. doi:10.1214/009053607000000848.
- Ma J, Heritier S, Lô SN (2014). “On the Maximum Penalized Likelihood Approach for Proportional Hazard Models with Right Censored Survival Data.” *Computational Statistics & Data Analysis*, **74**, 142–156. doi:10.1016/j.csda.2014.01.005.
- Mallick BK, Walker S (2003). “A Bayesian Semiparametric Transformation Model Incorporating Frailties.” *Journal of Statistical Planning and Inference*, **112**(1–2), 159–174. doi:10.1016/S0378-3758(02)00330-0.
- Mayr A, Fenske N, Hofner B, Kneib T, Schmid M (2012). “GAMLSS for High-dimensional Data—A Flexible Approach Based on Boosting.” *Journal of the Royal Statistical Society: Series C (Applied Statistics)*, **61**(3), 403–427. doi:10.1111/j.1467-9876.2011.01033.x.
- McLain AC, Ghosh SK (2013). “Efficient Sieve Maximum Likelihood Estimation of Time-Transformation Models.” *Journal of Statistical Theory and Practice*, **7**(2), 285–303.
- Möst L, Hothorn T (2015). “Conditional Transformation Models for Survivor Function Estimation.” *International Journal of Biostatistics*, **11**(1), 23–50. doi:10.1515/ijb-2014-0006.

- Möst L, Schmid M, Faschingbauer F, Hothorn T (2016). “Predicting Birth Weight with Conditionally Linear Transformation Models.” *Statistical Methods in Medical Research*, **25**(6), 2781–2810. doi:10.1177/0962280214532745.
- Müller J, Hothorn T (2004). “Maximally Selected Two-Sample Statistics as a new Tool for the Identification and Assessment of Habitat Factors with an Application to Breeding Bird Communities in Oak Forests.” *European Journal of Forest Research*, **123**, 218–228. doi:10.1007/s10342-004-0035-5.
- Pitman EJG (1939). “The Estimation of the Location and Scale Parameters of a Continuous Population of Any Given Form.” *Biometrika*, **30**(3/4), 391–421.
- R Core Team (2017). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/>.
- Racine JS, Hayfield T (2014). *np: Nonparametric kernel smoothing methods for mixed data types*. R package version 0.60-2, URL <https://CRAN.R-project.org/package=np>.
- Rigby RA, Stasinopoulos DM (2004). “Smooth Centile Curves for Skew and Kurtotic Data Modelled Using the Box-Cox Power Exponential Distribution.” *Statistics in Medicine*, **23**(19), 3053–3076. doi:10.1002/sim.1861.
- Rigby RA, Stasinopoulos DM (2005). “Generalized Additive Models for Location, Scale and Shape.” *Journal of the Royal Statistical Society: Series C (Applied Statistics)*, **54**(3), 507–554. doi:10.1111/j.1467-9876.2005.00510.x.
- Rothe C, Wied D (2013). “Misspecification Testing in a Class of Conditional Distributional Models.” *Journal of the American Statistical Association*, **108**(501), 314–324. doi:10.1080/01621459.2012.736903.
- Royston P, Parmar MKB (2002). “Flexible Parametric Proportional-hazards and Proportional-odds Models for Censored Survival Data, with Application to Prognostic Modelling and Estimation of Treatment Effects.” *Statistics in Medicine*, **21**(15), 2175–2197. doi:10.1002/sim.1203.
- Sabanés Bové D, Held L (2013). “Comment on Cai and Betensky (2003), On the Poisson Approximation for Hazard Regression.” *Biometrics*, **69**(3), 795–795. doi:10.1111/biom.12088.
- Schumacher M, Basert G, Bojar H, Hübner K, Olschewski M, Sauerbrei W, Schmoor C, Beylerle C, Neumann RLA, Rauschecker H F, for the German Breast Cancer Study Group (1994). “Randomized 2×2 trial evaluating hormonal treatment and the duration of chemotherapy in node-positive breast cancer patients.” *Journal of Clinical Oncology*, **12**, 2086–2093.
- Stasinopoulos DM, Rigby RA (2007). “Generalized Additive Models for Location Scale and Shape (GAMLSS) in R.” *Journal of Statistical Software*, **23**(7), 1–46. URL <http://www.jstatsoft.org/v23/i07>.
- Sun Y, Sundaram R, Zhao Y (2009). “Empirical Likelihood Inference for the Cox Model with Time-dependent Coefficients via Local Partial Likelihood.” *Scandinavian Journal of Statistics*, **36**(3), 444–462. doi:10.1111/j.1467-9469.2008.00634.x.

- Thas O, Neve JD, Clement L, Ottoy JP (2012). “Probabilistic Index Models.” *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, **74**(4), 623–671. doi:10.1111/j.1467-9868.2011.01020.x.
- Tobin J (1958). “Estimation of Relationships for Limited Dependent Variables.” *Econometrica*, **26**(1), 24–36. doi:10.2307/1907382.
- Tutz G (2012). *Regression for Categorical Data*. Cambridge University Press, New York, U.S.A.
- van de Geer S (2000). *Empirical Processes in M-Estimation*. Cambridge University Press, Cambridge, UK.
- van der Vaart AW (1998). *Asymptotic Statistics*. Cambridge University Press, Cambridge, UK.
- Varadhan R (2015). *alabama: Constrained Nonlinear Optimization*. R package version 2015.3-1, URL <https://CRAN.R-project.org/package=alabama>.
- Wood S (2017). *mgcv: Mixed GAM Computation Vehicle with GCV/AIC/REML Smoothness Estimation*. R package version 1.8-17, URL <https://CRAN.R-project.org/package=mgcv>.
- Wu CO, Tian X (2013). “Nonparametric Estimation of Conditional Distributions and Rank-Tracking Probabilities With Time-Varying Transformation Models in Longitudinal Studies.” *Journal of the American Statistical Association*, **108**(503), 971–982.
- Yang S, Prentice RL (2015). “Assessing Potentially Time-dependent Treatment Effect From Clinical Trials and Observational Studies for Survival Data, With Applications to the Women’s Health Initiative Combined Hormone Therapy Trial.” *Statistics in Medicine*, **34**(11), 1801–1817. doi:10.1002/sim.6453.
- Zeng D, Lin DY (2007). “Maximum Likelihood Estimation in Semiparametric Regression Models with Censored Data.” *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, **69**(4), 507–564. doi:10.1111/j.1369-7412.2007.00606.x.
- Zeng D, Lin DY, Yin G (2005). “Maximum Likelihood Estimation for the Proportional Odds Model with Random Effects.” *Journal of the American Statistical Association*, **100**(470), 470–483. doi:10.1198/016214504000001420.
- Zhang M, Davidian M (2008). “Smooth Semiparametric Regression Analysis for Arbitrarily Censored Time-to-Event Data.” *Biometrics*, **64**(2), 567–576. doi:10.1111/j.1541-0420.2007.00928.x.

Appendix

Computational Details

A reference implementation of most likely transformation models is available in the **mlt** package (Hothorn 2017a). All data analyses can be reproduced in the dynamic document Hothorn (2017b). Augmented Lagrangian Minimization implemented in the `auglag()` function of

package **alabama** (Varadhan 2015) was used for optimising the log-likelihood. Package **gamboostLSS** (version 1.2-2, Hofner *et al.* 2016) was used to fit GAMLSS models and kernel density and distribution estimation was performed using package **np** (version 0.60-2, Racine and Hayfield 2014). The additive Poisson model was fitted by **mgev** (version 1.8-17, Wood 2017). All computations were performed using R version 3.4.0 (R Core Team 2017).

Review History: Version 1 by Journal 1 (August 2015–January 2016)

Review for version 1 (<http://arxiv.org/abs/1508.06749v1>). Comments by the authors are printed in *italics*.

Associate Editor

The paper proposes a maximum likelihood approach to a general class of transformation models. The paper is nicely written. The proposed framework can incorporate different types of response variables: continuous, count, censored, truncated. The likelihood framework is attractive as it allows one to make use of many tools developed in the classical maximum likelihood setting.

The following are my main comments on the paper.

1. The abstract says “We propose and study properties of maximum likelihood estimators in the class of conditional transformation models”. However, the paper focuses on the one-sample case, that is, transforming a univariate random variable (without covariates). Hence, one does not see much of the “cascade of increasingly complex transformation models” claimed in the abstract. For this paper to be of more interest to JRSSB readers, the authors should instead focus on the more interesting and practically useful conditional transformations (i.e., the regression case), which were only sketched briefly in Section 4.2?

Regression models are discussed in much detail in Section 4.2 (pages 12-16), in three out of four examples presented in Section 5.1, and in the simulation model described in Section 5.2. The theory was presented in the unconditional case for notational and educational convenience only and carries over to the conditional case as stated in the 2nd paragraph of Section 4.2.

2. Referee 1 is concerned that “once the transformations are parameterized, the asymptotic results in Section 3 are obvious from standard textbooks”. This seems to be a valid concern.

We see this as an advantage, see also the corresponding comment to Referee 1.

I have my own concerns on the theoretical results: the main theorems (Theorem 1-3) are derived under the assumption Y_1, \dots, Y_n are *i.i.d.* from P_{Y, θ_0} , then it’s not much different from a parametric model. To me, it seems that a more reasonable theory should consider: (1) approximate the unknown transformation h using $a^\top \theta$; (2) then apply MLE at the second step. The current theory is only for the second step but ignores the spline approximation error in the first step. For transformation models, the theory would be more relevant to consider approximating h using the possibly misspecified $a^\top \theta$, for which one usually needs to let the number of basis functions goes to infinity at certain speed.

A corresponding consistency result for boosted conditional transformation models treating h as unknown can be found in Hothorn et al. 2014. The more useful asymptotic normality would be much harder to derive in this situation and we therefore decided to stay within the bounds of the modern likelihood world. Note that the comment only applies to continuous models as the transformation function is discrete in discrete models.

There is another related question: h is required to be strictly monotone but the basis function approximation does not automatically guarantee it. This needs to be carefully discussed.

Both parameterisations for discrete and continuous models (Bernstein polynomial) in Section 4.1 allow monotone transformation functions to be estimated under linear constraints. We used a spectral projected gradient method (see Appendix) for optimisation of the likelihood under such linear constraints. The estimated transformation functions are guaranteed to be monotone in y ; the nature and number of constraints are better explained in the revised version.

3. There is the important question that given a real data example, which transformation model should be chosen? When will the new method be more preferable than the alternative methods in the literature, such as Cheng et al. (1995), Chen et al. (2002)?

Classical linear transformation models treat the “baseline” transformation h_Y as a nuisance parameter. Fitting the same models under the full likelihood as suggested here leads to practically the same estimates of the regression coefficients (see Section 5) and a smooth interpolation of the post-hoc NPML estimate of h_Y . It is then relatively simple to construct asymptotic confidence bands for the distribution of the survivor function (see also Section 5). Most importantly, it is hard or even impossible to fit more complex models, for example one with time-varying effects or a conditional transformation model, using the classical approaches but it is straightforward (as we have shown in Section 5) using the most likely transformation approach.

The “Extension of the classical transformation models” appear to have some overlap with the authors’ recent work (Hothorn et al. (2014), Conditional Transformation Models, JRSSB)?

Yes, the class of “conditional transformation models” was introduced in Hothorn et al. 2014, along with a boosting algorithm for the minimisation of proper scoring rules. The 2014 paper is restricted to exact continuous responses. The novel contribution of the present manuscript is a full likelihood framework for arbitrary responses and corresponding likelihood inference procedures.

When will the method proposed in this paper be preferable to those in Hothorn et al. (2014)?

Whenever one is interested in likelihood inference for possibly censored, truncated or discrete responses under weak parametric assumptions in a fairly large model class.

Referee 1

The paper aims to study a unified framework based on transformations to address semi-parametric/nonparametric estimation for discrete, continuous or censored data. First, any

distribution functions are equivalent to some known distributions of transformed random variables. Then the paper proposes a class of parametric models for transformations so estimates the parameters via maximum likelihood estimation. The theoretical results of the parameter estimators are provided. The paper describes a list of examples to illustrate transformations and corresponding results in Section 4. Some numerical evidence is given in Section 5. My serious concern is about the new material presented here. The key motivation for using transformation (the results in Section 2) is available in standard textbooks.

All classical texts study and parameterise transformations $Y = g(Z)$, for example under the name location-scale models, structural inference, group families or group models. We study the case $h(Y) = Z$ and parameterise $h = g^{-1}$. The resulting class of “conditional transformation models” was introduced in Hothorn et al. 2014, along with a boosting algorithm for exact continuous responses. The novel contribution of this manuscript is a full likelihood framework for arbitrary responses and corresponding likelihood inference procedures. This has been clarified in the revision.

Once the transformations are parameterized, the asymptotic results in Section 3 are obvious from standard textbooks.

The simplicity of the approach is an advantage because more complex models can be fitted and analysed in this sound and well-established likelihood theory, whereas much more complex procedures are typically applied in (linear) transformation models. The partial likelihood is the most prominent representative of such “non-standard” estimation approaches for models with high- or even infinite-dimensional nuisance parameters.

Section 4 provides a list of examples to demonstrate how a variety of models can be viewed as from certain transformations, which can be also found in Bickel et al. (1993).

Section 4.2 contains three subsections “Classical Transformation Models”, “Extensions of Classical Transformation Models” and “Novel Transformation Models”. Well-known linear transformation models are shown to be part of the unifying theory presented here in the first part, along with appropriate references. The book by Bickel et al. is an early but not the first reference for a unifying view on linear transformation models (and is restricted to the continuous case). We cited the (as far as we know) first paper in this field by Doksum and Gasko (1990) instead. The models discussed in the remaining two subsections have not been discussed in this context so far or are completely novel.

My specific comments include:

1. in the introduction, it is not clear which transformations were actually used in the literature. Is it $h_Y(y)$ or $h(Y|x)$?

With the few exceptions mentioned in the Introduction, $h_Y(y)$ is typically treated as a nuisance parameter and therefore, it is not parameterised nor estimated directly. The transformation $h(Y|x)$ is a shifted (by a linear predictor) version of $h_Y(y)$ as explained in detail in Section 4.2.

2. the bottom of page 4, what if Y contains both discrete and continuous components?

Y is univariate and either discrete or continuous. For multiple observations $Y_i, i = 1, \dots, N$ of mixed type, the corresponding likelihood contributions (page 5) are simply multiplied (for example in “double censoring”).

3. in Definition 1, does the maximum always exist? If not, what are the conditions for existence?

The distribution of Z needs to be log-concave and the basis a needs to ensure identifiability of the distribution of Y . Both conditions have been added to the revision.

4. in Section 3, originally I thought that the paper would consider nonparametric transformations but it turns out that the actual transformations are parametric in this section. Obviously, a big issue is how to choose basis functions and how much parameterization one needs. The paper has little discussion on this.

We used the term “non-parametric” for indicating the invariance with respect to the distribution of Z which was non-standard and has been revised. The type of the basis functions is problem-specific. In the discrete case, we have $K - 1$ parameters for K levels, in the continuous case we used Bernstein polynomials of order M for modelling smooth transformation functions. For count data, one could use either parameterisation, depending on the number of counts. In the discrete case, the number of basis functions is fix. In the continuous case (as explained in Section 4) the number of basis functions is often limited to two (in all transformation functions allowing linear transformations of Z only) or can be relatively large due to the monotonicity constraints (see Section 4.2). For the Geyser example, we tried up to $M = 70$ basis functions and the resulting distribution function closely approximates the ECDF as one would expect from the comments in subsection 4.1 (results not shown in paper).

5. another concern is that the parameterization should satisfy $h(y_1) < h(y_2)$ if $y_1 < y_2$. This can result in complicated constraints, especially when $h(\cdot)$ depends on covariates x .

The basis functions used in this paper allow monotonicity constraints to be formulated as linear constraints on the parameters (page 11 and 12). Standard algorithms for optimisation under linear constraints can be applied. The necessary constraints are better explained in Sections 4 and 5 in the revision.

6. no computation details are given for the maximization. Indeed, computation is a non-trivial issue in transformation models for censored data.

*Appendix “Computational Details” contains a reference to the optimiser used for the experiments. We have tried other linear constraint optimisers as well. All of them worked reasonably well; the implementation in package **BB** was chosen because of its convenient user interface. One advantage of the framework presented in this paper is the simplicity of the corresponding optimisation problem and thus standard optimisers can be applied (as mentioned in the Discussion). In fact, the evaluation of the likelihood is easier for censored observations as one only needs to compute the distribution function, instead of the density. Note that the core of our reference implementation including all models presented here was written in less than 1000 lines of pure R code, this also highlights the computational advantages of the full likelihood estimation procedure.*

7. in general, when we choose different distributions for Z , it will result in different transformations so parameterization can be different. Would results be sensitive to the choice of Z 's distribution?

It depends on the model: In three out of four examples presented in Section 5.1 the distribution of Y is invariant wrt the distribution of Z . Thus, for flexible enough parameterisations of the transformation function, any choice of Z (subject to the restrictions given in the paper) will practically lead to the same estimated distribution, but of course with different parameters ϑ . In such a situation, we are not interested in the parameter estimates but in the estimated (conditional) distribution, density, hazard or cumulative hazard functions (as in the the density estimation and quantile regression applications). In contrast, the minimum extreme value distribution was used to allow comparisons with a standard Cox model in the survival application. In this model, the choice of Z as the minimum extreme value distribution allows an interpretation of the regression coefficients β as log-hazard ratios; all of this is explained and discussed in Section 4.2.

Referee 2

This paper develops a new theoretical and computational likelihood based framework, which embeds a large class of transformation models (parametric, semi-parametric, and non-parametric) commonly estimated by other procedures. The approach also allows extensions to novel transformation models. Besides the theoretical foundations (complemented by an extensive and exhaustive bibliography), the paper provides several persuasive applications based on real data. It is well written and deserves publication. No revision is necessary.

Review History: Version 2 by Journal 2 (January–April 2016)

Review for a condensed variant of version 2 (v2, <http://arxiv.org/abs/1508.06749v2>). Comments by the authors are printed in *italics*.

Associate Editor

This paper aims to develop a unified transformation based framework for modeling, estimation and inference with a wide range of data types. While the presented formulation of the transformation model seems interesting, as noted by all reviewers, it confines the proposed model to a fully parametric setting through specifying the basis function and F_Z (an analog to "error distribution"). This poses a major limitation of the proposed work. For example, in survival applications, the proposed model may only correspond to a proportional hazards model with parametric baseline hazard function, which is more restrictive than the commonly adopted semi-parametric proportional hazards model.

*This statement is true from a theoretical point of view. Practically, the regression coefficients in a Cox model obtained from the partial likelihood and the full likelihood are numerically equivalent even for rather low-dimensional approximations of h_Y . Empirical evidence was presented in Section 5 of v2 (Figure 6) and we now report on more detailed investigations of this issue in the package vignette for the R add-on package **mlt.docreg** distributed at CRAN. Parameterising the transformation function allows much more flexible models to be formulated and estimated (for example distribution regression models as in the growth curve analysis example) than anything we are aware of in the NPMLE world. As a side-comment, the dominant semiparametric view on survival analysis made estimation of the Cox model for interval censored data rather troublesome for over 40 years (software became available only*

very recently). As a consequence, many analyses of, most prominently, disease-free survival times treating events observed at follow-up examinations as “exact” event times are probably flawed. Convenient estimation of the Cox model under **all** forms of censoring is a by-product of our approach (and is implemented in the **mlt** package).

When positioning the proposed transformation model as a general class of parametric models, there are some important issues left unaddressed:

1. How to decide the form of the basis function in real data analysis?

As explained in sections 4 and 5, there is not much to decide. For discrete observations, each element of the support (except one) gets one parameter assigned and for the Bernstein polynomials used in the empirical parts, M is not a hyper parameter. New empirical evidence can be found in the above mentioned vignette.

2. Similarly, how to choose the distribution function for Z ?

Some conditional transformation models are practically invariant wrt Z (in the Old Faithful and growth curve and partially the GBSG-2 examples), as the fitted distribution functions are numerically the same for different choices of F_Z . For others, Z is part of the model formulation. For example, in the Cox model we need Z to follow a minimum extreme value distribution if we want to interpret the regression coefficients β as log-hazard ratios. As mentioned in the discussion, it is possible to estimate F_Z and McLain and Ghosh (2013) introduce a suitable family of distributions (newly added to discussion) for this exercise.

3. What is the advantage of adopting the transformation model formulation (presented in Definition 4) over direct parametric modeling of the distribution function?

*Section 2 explains that we can generate **all** distributions from a suitable h and not just the ones we can find in textbooks. In this sense, the framework is nonparametric at its heart.*

In addition, I find the presentation of the conditional transformation model quite vague. Is there an implicit assumption that $h(Y|X)$ has a distribution independent of X ?

$F_Z(h(Y|x))$ has a (discrete or continuous) uniform distribution in $[0, 1]$ for every x by the probability integral transform.

Referee 1

This paper provides a mathematical abstraction of the transformation model, and considers inference through fitting the most likely transformation as a maximum likelihood approach. Regularity conditions were provided so that the usual maximum likelihood mathematical results hold for this transformation model as stated in three theorems. Then the authors discussed connections to transformation models in literature, and demonstrated applications on three example data sets. The transformation model has been used extensively in literature, but the mathematical abstraction to summarize the general approach does not seem to appear before. However, I do have concerns about this manuscript.

1. The results in section 3 are rather restatements of standard results. More importantly, they are only stated in the fully parametrized model, which do not include the most

practically useful and academically interesting semi-parametric (non-parametric) transformation model. For fully parametrized model, direct likelihood approach is natural, and it is not clear that this restatement in terms of transformation model brings any extra insight.

See the comment to overall statement by AE.

2. The authors claimed that this likelihood approach clearly illustrate the connections the general transformation models, while the lack of general understanding in previous literature was due to presentation in the relatively narrow contexts of censored or ordinal data (page 4). However, seems to me that a main reason for the authors' simple approach is to ignore the semi-parametric/non-parametric setting that are the main mathematical challenges.

The likelihood does not illustrate the connections between models, the model formulation does. We study and extend transformation models common in the semi- and non-parametric world while making model estimation much simpler (also numerically) by introducing appropriate parameterisations. See also comment to overall statement by AE.

The authors deal with the semi-parametric/nonparametric case by using Bernstein polynomials of fixed degree to approximate the transformation function, then treat this as a parametric fitting. However, for practical purpose, the choice of the degree are critical for data analysis.

*New empirical evidence presented in the package vignette of package **mlt.docreg** suggests that this is actually not the case.*

And for theoretical correctness, the degree should increase with sample size, not fixed.

A new paragraph 3.2. was introduced making the link between the estimation procedure applied here and the asymptotic theory for unconditional and linear transformation models presented by McLain and Ghosh (2013).

The authors simply state that: cannot choose M "too large" (page 18, line 6). This does not really prove any practical guidance nor provide theoretical guarantee for applications. The three examples in section 5, used degrees 8, 3, 5 respectively. There are no discussion on why these degrees are appropriate, and how would the results change with a different choice of these tuning parameters?

See above.

3. Related to the above point, page 20 discussed the connection to the survival time models. But those are rather misleading. For proportion hazards/odds models commonly used in survival analysis, the transformation h_Y is generally assumed unknown (rather than parameterized with Bernstein polynomial here). So this model does not capture the most general PH/PO model, where the nonparametric maximal likelihood estimator was proven to work.

See the comment to overall statement by AE.

Also, for the AFT model, the current model just correspond to some parametric sub-models with both (F, h) assumed known. Those are specific parametric models that do not need this new transformation model abstraction.

*In our opinion it is very interesting, at least from a practical, computational, and educational point of view, to unify all models in this framework. Understanding the connections between models allows one to estimate **all** models under **all** forms of random censoring and truncation using just about 1000 “smart” lines of R code (now available in package **mlt**).*

For the more general AFT model, the F should be assumed unknown, which the transformation model approach in this paper does not cover how to do statistical inference.

See AE point 3.

Referee 2

In this paper, the authors show that one can always write a potentially complex distribution function F_Y as the composition of a priori specified distribution function F_Z and a strictly monotone transformation function h (unknown). The task of estimating F_Y is then reduced to obtaining an estimate of h , which is parameterised in terms of basis functions. The likelihood function of the transformed variable can then be characterised by transformation function and the parameters of interest can be estimated and analysed in the maximum likelihood framework. The authors establish the asymptotic normality of the proposed estimators. A reference software implementation of maximum likelihood-based estimation for conditional transformation models was employed to illustrate the wide range of possible applications.

Major:

1. My major concern is the specification of F_Z , which is assumed to be known. Since Y can be subject to censoring/truncation, how do we choose appropriate F_Z based on censored/truncated data? In particular, for interval-censored or truncated data, how do we decide an appropriate F_Z in the likelihood function? The authors should address this issue.

*F_Z defines the transformation **model** and has nothing to do with its **estimation** under censoring or truncation. Only the likelihood function takes censoring (by integrating over all possible realisations) and truncation (by conditioning) into account. For example, the Cox **model** has nothing to do with censoring, only its **likelihood** (full or partial) allows for this. In the same spirit, we can fit a normal linear model under censoring and truncation without changing anything in the model formulation.*

The Other Comments:

1. page 8, lines -1 through -5: The authors only demonstrate likelihood contribution of censored data. How about truncated data?

The likelihood for truncated observations is given on page 6, line 2, of v2 and page 9, line 12 in your version.

2. page 9, lines -1 through -6: It is not clear to me that $Y \in C_i$, where C_i denotes an independent sample of possibly censored or truncated observations.

C_i are intervals, ie the observations, and Y is the rv.

What is the definition of H ?

The space H is defined on page 4, line -8 in v2 and on page 7, line 14 in your version.

In practice, can we obtain \hat{h}_N ?

Via plug-in, as explained on page 7, line -4 in v2 and page 11, line -8 in your version.

When truncation is present (such as left truncation or double truncation), does the concavity of the log-likelihood hold ? How does this impact the uniqueness of \hat{h}_N ?

The sum of two concave functions is again concave.

3. page 10, line 20: What is the advantage of parameterising the transformation function $h(y)$ in terms of basis functions. Is there any other alternative (e.g. nonparametric estimator)? Please comment on this.

Parameterising h leads to models with well-defined parameters and allows efficient evaluation of the likelihood function.

4. page 11, line 6: When truncation is present, is $\hat{\vartheta}_N$ unique ?

See your point 2.

5. page 13: In Theorems 1 and 2, the authors do not mention any assumption regarding the left and right endpoints of the distribution functions of lifetime, censoring time and truncation time. Are they related to the asymptotic properties of $\hat{\vartheta}_N$?

Sure, but as stated on page 9, line 3 in v2 and page 12, line -3 in your version, the theorems describe the asymptotic distribution for the “exact” case.

6. page 18: The authors should point out how the conditional transformation model relate to semiparametric transformation models (Cheng et al. 1995). How does the proposed method differ from that of Zeng and Lin (2006,2007).

Zeng and Lin (2007) study linear transformation models without explicit parameterisation of the baseline transformation function λ enriched with random effects. The model (without random effects) is a simple linear transformation model known for decades and their estimation procedure is completely different from the one discussed here.

7. page 24: In empirical evaluation, the data illustration can be improved. Please compare the proposed method with the other approaches.

We removed the simulation study presented in v2 from the condensed version, please see Section 5 in all versions of the arXiv report.

Review History: Version 3 by Scandinavian Journal of Statistics (November 2016–March 2017)

Review for a condensed variant of version 3 (v3, <http://arxiv.org/abs/1508.06749v3>). Comments by the authors are printed in *italics*.

Associate Editor

All reviewers think that the paper is interesting and that the unified framework includes a variety of data types and models. The comments are mostly minor. However, all reviewers

and myself think that the paper presentation should be much improved in next version. Please follow the reviewer's specific comments to make revision.

We revised the manuscript taking all suggestions made by the three referees into account. The main changes relate to the following issues

- **Connections between Models.** Two referees suggested to improve the presentation of connections between models. We added a new Table 1 containing a systematic overview on linear and response-varying transformation models. We also stress that models, parameterisation and estimation are not a unity but should be understood as separate issues.
- **Parameter Interpretability.** In some models, transformation functions or shift parameters β or response-varying parameter functions $\beta(y)$ have a clear meaning. We mention interpretability wherever possible and also included this information in our new Table 1.
- **Constraints.** We explicitly present linear constraints for all models discussed in this manuscript.
- **Choice and Meaning of F_Z .** Following a suggestion by referee 3, we added the mean one exponential distribution and a presentation of corresponding additive hazard models. Distributions F_Z with free parameters are discussed in a dedicated paragraph in the last section now.

Referee 1

This study develops a unified framework for studying transformation models and the corresponding (downstream) estimates. What is really nice about this study is that multiple data types, distributions, and models are comprehensively investigated under one framework. Some numerical studies are conducted. I have some minor comments.

1. *The introduction is "too comprehensive" in that a lot of existing studies are mentioned. With so much information, it gets confusing. Please consider better organising discussions on the existing studies. Especially please make it clear which existing studies can be considered under the proposed framework, and which are beyond.*

The common denominator of all studies cited are transformation models. The differences lie in (1) different estimation techniques, such as non-parametric maximum likelihood estimation, estimation equations or optimisation of scoring rules and (2) different parameterisations (or lack thereof). We agree that it is hard to see and understand the connections (we explicitly comment on this problem in the 3rd paragraph) because models, parameterisation and estimation are often treated as a unity. We addressed this issue in two ways. First, the introduction makes it clear that the transformation *models* cited in the introduction can be understood as members of the family of transformation models presented in this manuscript. With the specific parameterisations proposed here, one can also *estimate* this models by maximum likelihood. Second, we added a systematic overview on transformation models and their connections to other models following a suggestion by referee 3 (new Table 1). We hope that this overview will make it easier to understand the connections between different models.

2. *For the density of f_Z , the authors mentioned normal, logistic, and minimum extreme value distributions. And there should be quite a few other choices. Theoretically, computationally, and numerically, does this choice make a difference?*

The main difference is with respect to the interpretation of regression parameters β in linear transformation models. Following a suggestion by Referee 3, we now present a systematic overview on different flavours of transformation models and corresponding parameter interpretations (Table 1). We also added the mean one exponential distribution F_Z because it allows fitting additive hazard models. Some authors studied distribution functions F_Z with free parameters, these approaches are now discussed in a new paragraph in the last section.

3. *The nonparametric inference in Section 3.2 is discussed relatively briefly. For smooth nonparametric estimation, there are multiple candidate approaches. Does that choice make a difference?*

Only from a computational point of view. It is very easy to impose monotonicity on a Bernstein polynomial (make sure the parameters are monotone increasing). This is much harder to achieve for a B-spline, for example. These issues are discussed in Section 4.1.

4. *The illustrations in Section 5.1 do not provide much insight. They can be moved after simulation.*

Feedback from readers showed that many people have difficulties following our path through the transformation model jungle. Presentation of applications usually helps them to get started much easier. We therefore wanted to put strong emphasis on applications. Section 5 illustrates the broadness of transformation models without the need to understand every technical detail. We therefore prefer to keep the current order of presentation.

Referee 2

Authors studied the properties of maximum likelihood estimators in the class of conditional transformation models. The distribution of response variable was written in the priori specified distribution, such as standard normal, standard log-logistic and the minimum extreme value distribution after certain transformation $h(y)$. Then, the same theoretical and computational framework can be applied simply by choosing an appropriate transformation function and parameterizations thereof. For discrete and continuous responses, they established the asymptotic normality of the proposed estimators. The “mlt” package in R can implement the proposed method and empirical evaluation illustrated the wide range of possible applications.

1. *The model was determined by the priori specified distribution and $h_Y(y) + h_x(x)$ and $h_Y(y)$ can be specified or approximated by the spline. The authors have tested the performance of data without censoring and right censoring. Theoretically, this MLE approach can be extended to the interval censored data. However, usually, there are issues in estimation for the interval censored data with spline from the direct MLE, which may still exist in the proposed method. The authors may add discussions on this.*

In fact, linear transformation models of the form $h_Y(y) + h_x(x)$ are the simplest members of the much more general class of conditional transformation models covered in our manuscript. For linear and more complex transformation models, we did not encounter theoretical or computational difficulties when estimating most likely transformations. In her master thesis, Mariia Dobrynina systematically investigated finite sample properties of most likely transformations under interval censoring. She found the R add-on package "mlt" to be correct also for interval censored data. The thesis can be downloaded from <http://user.math.uzh.ch/hothorn/docs/MasterThesisSfS.pdf>.

2. *There is no definition of \tilde{h} in definition 1.*

\tilde{h} is an element of \mathcal{H} (underneath argmax).

Referee 3

In the paper "Most Likely Transformations" the authors propose the use of a flexible class of parametric sieve-type models for conditional and unconditional density estimation. The theory of the models is presented, along with the examples of how the approach would be applied. I found this to be a well written paper on an interesting topic. Please find my main comments on the manuscript below.

1. *I think it's important to give some differences between the proposed approach and the well-known transformation models from Zeng and Lin. It appears that this is all in the introduction, but a sentence focusing and clarifying this aspect would be helpful.*

Zeng and Lin study *nonparametric* maximum likelihood estimator without explicit parameterisations of baseline transformation functions h_Y . We highlighted this fact. The models discussed by Zeng and Lin, however, can be understood as conditional transformation models and thus our likelihood-based estimation procedure can be used for parameterisation and estimation. This is now better explained in the introduction.

2. *Your model also has a clear connection with some grouped survival methods. For example*

- *Aranda-Ordaz FJ. An extension of the proportional- hazards model for grouped data. *Biometrics* 1983; 39: 109–117 and*
- *Tibshirani RJ and Ciampi A. A family of proportional- and additive- hazards models for survival data. *Biometrics* 1983; 39: 141–147.*

The connection to these models with some discussion would be a nice addition.

Thank you very much for this suggestion. We added the corresponding mean one exponential distribution F_Z and discuss the resulting additive hazard models in Section 4 and in Table 1 now. We also added a paragraph discussing the possibility to have free parameters in F_Z in the last section. The piecewise constant parameterisation employed in these two papers is, however, only motivated by the limited computing resources available in the 1980ies. To quote Aranda-Ordaz: "It is often reasonable to consider survival time as essentially a continuous random variable." Thus, with today's computing power, we think it is much more adequate to estimate smooth transformation functions, for example using the method introduced in our paper.

3. *I think you could add a table summarizing the correspondence of the proposed approach to the various popular modeling procedures by (F_Z, a, ϑ) . This is in the paper, but more scattered. A difficult issue with a paper that is as broad as this one, is the presentation of the various of models and methods in a systematic fashion. Section 4 does an honorable job of this, but making the connections is still difficult. A focused table would be a nice start to clarifying the methods, but I recommend that authors look at this issue more generally (see comment 6 as well).*

We thank the referee for this suggestion. We added a table covering linear transformation models as well as response-varying transformation models for binary, polytomous, count and continuous regression, the latter including survival analysis, to Section 4. Indeed, this more systematic overview helps to identify white spots in the model landscape much easier.

4. *The first equation in “Count Regression Without Tears” the first equation has an extra ‘)’.*

Thank you, fixed.

5. *Please explain the issue of the linear constraints of the model in a little more detail. They are mentioned in passing in Section 4, then they appear as a much bigger issue in Section 5. A more focused discussion of the roles that linear constraints can play would be helpful.*

Linear constraints on the model parameters are necessary to ensure monotonicity of the transformation function. For linear transformation models, monotonicity of the “baseline” transformation h_Y is explained in Section 4 and we now added a more detailed discussion to section 4.3 and 4.5 (and added a reference to a technical report explaining the linear constraints for different models). The specific constraints for the empirical studies in Section 5 are now made explicit.

6. *Is there any form that can be given to help the interpretation (even graphically) of covariate parameters in some tangible way? This is attempted in Figure 3, but having “Transformation deviation” as the y-axis makes the plot uninterpretable. Is there an equation that will give some tangible meaning to the coefficients for all possible models (or maybe just all continuous models)? Further, is the sign interpretable. Maybe there is a broad subset of models where a coefficient with a particular sign has a monotonic influence on the outcome (e.g., negative coefficient always extends survival)? Please include some discussion on the interpretation. It does not have to hit these specific points, but a discussion of the specific and general (i.e., sign) interpretations would be beneficial.*

The new table presenting an overview on models also contains interpretations for the baseline transformation function h_Y and the regression coefficients β in linear transformation models of the form $h(y | x) = h_Y(y) - x^\top \beta$. We parameterised a negative shift, because of the interpretation $E(h(Y | x)) = x^\top \beta + c$, ie larger values of the linear predictor correspond to larger conditional expectations (but smaller risks in survival analysis). This relationship is discussed in Section 4.3.

Affiliation:

Torsten Hothorn
Institut für Epidemiologie, Biostatistik und Prävention
Universität Zürich
Hirschengraben 84, CH-8001 Zürich, Switzerland
Torsten.Hothorn@uzh.ch

Lisa Möst
Institut für Statistik
Ludwig-Maximilians-Universität München
Ludwigstraße 33, DE-80539 München, Germany

Peter Bühlmann
Seminar für Statistik
ETH Zürich
Rämistrasse 101, CH-8092 Zürich, Switzerland