An algebraic interpretation of the multivariate *q*-Krawtchouk polynomials

Vincent X. Genest † , Sarah Post $^\ddag$, and Luc Vinet *

†Department of Mathematics, Massachusetts Institute of Technology, 77 Massachusetts Avenue, Cambridge, MA 02139, USA ‡Department of Mathematics, University of Hawai'i at Manoa, 2665 McCarthy Mall, Honolulu, HI 96822, USA *Centre de recherches mathématiques, Université de Montréal, C.P. 6128 succ. Centre-ville, Montréal (Québec) H3C 3J7, Canada

Abstract

The multivariate quantum *q*-Krawtchouk polynomials are shown to arise as matrix elements of "*q*-rotations" acting on the state vectors of many *q*-oscillators. The focus is put on the twovariable case. The algebraic interpretation is used to derive the main properties of the polynomials: orthogonality, duality, structure relations, difference equations and recurrence relations. The extension to an arbitrary number of variables is presented.

Keywords: Multivariate *q*-Krawtchouk polynomials; *q*-oscillator algebra; *q*-rotations **AMS classification numbers:** 33D45, 16T05

1 Introduction

The purpose of this paper is to provide an algebraic model for the multi-variable *q*-Krawtchouk polynomials and to show how this model provides a cogent framework for the characterization of these orthogonal functions. The algebraic interpretation presented here is in terms of matrix elements of unitary "*q*-rotations" acting on *q*-oscillator states; these *q*-rotations are expressed as *q*-exponentials in the generators of several independent q -oscillator algebras. For illustration purposes, the focus will be put on the two-variable case. The algebraic model will lead to a natural and explicit derivation of the main properties satisfied by the bivariate quantum *q*-Krawtchouk polynomials: orthogonality relation, duality property, structure relations, *q*-difference equations and recurrence relations. How this approach generalizes directly to an arbitrary number of variables will also be explained.

The standard (univariate) Krawtchouk polynomials form one of the simplest families of hypergeometric orthogonal polynomials of the Askey scheme [\[19,](#page-21-0) Ch. 9]. These polynomials of degree *n* in *x*, denoted by $\kappa_n(x; p, N)$, have the expression

$$
\kappa_n(x;p,N) = (-1)^n (-N)_n {}_2F_1\left(\begin{array}{c} -n, -x \\ -N \end{array}; \frac{1}{p}\right), \qquad n = 0, 1, \ldots, N,
$$

where $0 < p < 1$ $0 < p < 1$ is a parameter and N is a positive integer¹. In the above, pF_q stands for the generalized hypergeometric series and

$$
(a)_n = a(a+1)\cdots(a+n-1), \qquad (a)_0 := 1,
$$

is the shifted factorial, or Pochhammer symbol [\[1,](#page-20-0) (1.1.2)]. The polynomials $\kappa_n(x; p, N)$ are orthogonal with respect to the binomial distribution, i.e.

$$
\sum_{x=0}^N \binom{N}{x} p^x (1-p)^{N-x} \kappa_n(x;p,N) \kappa_m(x;p,N) = \eta_n \delta_{nm},
$$

where $\eta_n \neq 0$ are normalization coefficients. The Krawtchouk polynomials have a well-known algebraic interpretation as matrix elements of unitary irreducible representations of $\mathfrak{so}(3)$; in point of fact, most of their properties follow from that interpretation (see for example [\[20\]](#page-21-1)[\[28,](#page-22-0) Section 6.8]).

In [\[26,](#page-22-1) [27\]](#page-22-2) Tratnik presented a multi-variable extension of the Askey tableau, giving for each family of the scheme explicit expressions of the polynomials as well as of their orthogonality measures; special cases of these multivariate polynomials occurred earlier, e.g. [\[18\]](#page-21-2). Much later in [\[11\]](#page-21-3), Geronimo and Iliev showed that Tratnik's families of orthogonal polynomials are all bispectral: they exhibited the recurrence relations and the eigenvalue equations that these polynomials satisfy. Despite the fact that the multivariate polynomials defined by Tratnik are expressed in terms of one-variable polynomials, they are non-trivial and arise in a number of different contexts, see for example [\[9,](#page-21-4) [13,](#page-21-5) [18,](#page-21-2) [21,](#page-21-6) [22,](#page-21-7) [24\]](#page-21-8). In two variables, the Krawtchouk polynomials defined by Tratnik have the expression

$$
K_{n_1,n_2}(x_1,x_2;p_1,p_2;N) = \kappa_{n_1}(x_1;p_1,x_1+x_2)\kappa_{n_2}(x_1+x_2-n_1;p_2,N-n_1), \quad n_1,n_2 = 0,1,...N,
$$

with $n_1 + n_2 \leq N$ and are orthogonal with respect to the trinomial distribution

$$
\sum_{\substack{x_1,x_2=0\\x_1+x_2\leq N}}^{N} {N \choose x_1,x_2} \left(\frac{p_1p_2}{1-p_2} \right)^{x_1} \left(\frac{(1-p_1)p_2}{1-p_2} \right)^{x_2} (1-p_2)^N
$$

$$
\times K_{n_1,n_2}(x_1,x_2) K_{m_1,m_2}(x_1,x_2) = \eta_{n_1,n_2} \delta_{n_1m_1} \delta_{n_2m_2},
$$

where $\binom{N}{x_1,x_2}$ are the trinomial coefficients and where $p_1,\,p_2$ are parameters. In [\[10\]](#page-21-9), it was shown that Tratnik's multivariate Krawtchouk polynomials arise in the matrix elements of the reducible unitary rotation group representations on oscillator states 2 2 . It was also shown that Tratnik's Krawtchouk polynomials are in fact a special case of the multivariate Krawtchouk polynomials introduced by Griffiths in [\[12\]](#page-21-10) and by Hoare and Rahman in [\[15\]](#page-21-11). Similarly to the one-variable case, the properties of these multivariate functions follow from their group-theoretical interpretation [\[10\]](#page-21-9); see also [\[17\]](#page-21-12).

Basic analogs of Tratnik's multivariate orthogonal polynomials were introduced by Gasper and Rahman in [\[5\]](#page-21-13). Like Tratnik, Gasper and Rahman explicitly defined their families of multivariate *q*deformed polynomials as convoluted products of one-variable polynomials of the basic Askey scheme and proved their orthogonality relations. In [\[16\]](#page-21-14), Iliev showed that these polynomials, which form a multivariate extension of the basic Askey tableau, are also bispectral. We note here that some of

¹The polynomials $\kappa_n(x; p, N)$ agree with $K_n(x; p, N)$ in [\[19,](#page-21-0) Section 9.11] up to a normalization factor. In this paper, the uppercase K are reserved for the multivariate polynomials.

 2 Let us note that in [\[10\]](#page-21-9) and [\[11\]](#page-21-3), a different definition of Tratnik's two-variable Krawtchouk polynomials was used. The aforementioned algebraic interpretation is valid for both definitions.

these families of orthogonal polynomials, namely the multivariate *q*-Hahn and *q*-Racah polynomials, have already been seen to arise in algebraic and combinatorial contexts [\[23,](#page-21-15) [25\]](#page-21-16). The two-variable *q*-Krawtchouk polynomials introduced in [\[5\]](#page-21-13) have the expression

(1)
$$
\mathbf{K}_{n_1,n_2}(x_1,x_2;\alpha_1,\alpha_2;N) = k_{n_1}(x_1;\alpha_1^{-2},x_1+x_2;q) k_{n_2}(x_1+x_2-n_1;\alpha_2^{-2},N-n_1;q),
$$

with n_1, n_2 are non-negative integers such that $n_1 + n_2 \leq N$. Here $k_n(x, p, N; q)$ stands for the onevariable quantum *q*-Krawtchouk polynomials of degree *n* in *q* [−]*^x* which are defined as [\[19,](#page-21-0) Ch. 14]

(2)
$$
k_n(x; p, N; q) = (-1)^n (q^{-N}; q)_n q^{\binom{n}{2}} 2\phi_1 \left(\frac{q^{-n}, q^{-x}}{q^{-N}}; q, pq^{n+1} \right), \quad n = 0, 1, ..., N,
$$

where $0 < q < 1, p > q^{-N}$ and where N is a positive integer^{[3](#page-2-0)}. In the above

$$
(a;q)_n = (1-a)(1-aq)\cdots(1-aq^{n-1}),
$$
 $(a;q)_0 = 1,$

stands for the *q*-Pochhammer symbol and $r\phi$ _{*s*} is the basic hypergeometric series defined as [\[4,](#page-20-1) (1.2.22)]

$$
{}_r\phi_s\bigg(\n\begin{array}{c}\na_1, a_2, \ldots, a_r \\
b_1, b_2, \ldots, b_s\n\end{array}; q, z\n\bigg) = \sum_{k \geq 0} \frac{(a_1; q)_k (a_2; q)_k \cdots (a_r; q)_k}{(q; q)_k (b_1; q)_k (b_2; q)_k \cdots (b_s; q)_k} \left[(-1)^k q^{\binom{k}{2}}\right]^{1+s-r} z^k.
$$

In this paper, we shall provide an algebraic interpretation of Gasper and Rahman's multivariate (quantum) *q*-Krawtchouk polynomials and show how the properties of these polynomials can be obtained from that interpretation. On the one hand, this work can be considered as a multi-variable generalization of [\[8\]](#page-21-17), where the algebraic interpretation of the one-variable quantum *q*-Krawtchouk in terms of *q*-rotations was investigated. On the other hand, it can be viewed as a *q*-generalization of [\[10\]](#page-21-9) (in the Tratnik special case), where the interpretation of the multivariate Krawtchouk polynomials in terms of rotation group representations on oscillator states was studied. For simplicity the emphasis shall be put on the two-variable case, with the understanding that the approach directly extends to an arbitrary number of variables.

The outline of the paper is as follows. In Section 2, the interpretation of the one-variable quantum *q*-Krawtchouk polynomials in terms of *q*-rotations, investigated in [\[8\]](#page-21-17), is briefly reviewed. In Section 3, the algebraic model for the two-variable quantum *q*-Krawtchouk polynomials is presented. The bivariate quantum *q*-Krawtchouk are seen to arise as matrix elements of unitary three-dimensional *q*-rotations acting on the state space of three mutually commuting copies of the *q*-oscillator algebra. The orthogonality weight and the duality relation for the polynomials are derived. In Section 4, new structure relations for two-variable *q*-Krawtchouk polynomials are found using the algebraic setting. In Section 5, the recurrence relations and the difference equations are obtained and are seen to coincide with those found by Iliev in [\[16\]](#page-21-14). The multivariate case is discussed in the conclusion.

2 Review of the univariate case

In this section, the algebraic interpretation of the one-variable quantum *q*-Krawtchouk polynomials as matrix elements of unitary *q*-rotations acting on the space of two *q*-oscillators is reviewed. Mild generalizations of the *q*-Baker–Campbell–Hausdorff formulas are also presented.

 3 The polynomials $k_n(x; p, N; q)$ agree with $K_n^{qtm}(q^x; p, N, q)$ of [\[19,](#page-21-0) Section 14.14] up to a normalization factor.

2.1 The *q***-oscillator algebra and the Schwinger realization of** $U_q(\mathfrak{sl}_2)$

Let $0 < q < 1$ and consider two mutually commuting copies of the q-oscillator algebra with generators A_{\pm} , A_0 and B_0 , B_{\pm} that satisfy the relations

(3)
$$
[X_0, X_{\pm}] = \pm X_{\pm}, \qquad [X_-, X_+] = q^{X_0}, \qquad X_- X_+ - q X_+ X_- = 1,
$$

with $X \in \{A, B\}$ and where $[X_-, Y_-] = 0$ if $X \neq Y$. The algebra [\(3\)](#page-3-0) has a standard infinite-dimensional representation on the orthonormal basis states

$$
|n_A, n_B\rangle := |n_A\rangle \otimes |n_B\rangle,
$$

where n_A and n_B are non-negative integers. This representation is defined by the following action of the generators on the factors of the direct product states:

$$
(4) \qquad X_{+}|n_{X}\rangle = \sqrt{\frac{1-q^{n_{X}+1}}{1-q}}|n_{X}+1\rangle, \qquad X_{-}|n_{X}\rangle = \sqrt{\frac{1-q^{n_{X}}}{1-q}}|n_{X}-1\rangle, \qquad X_{0}|n_{X}\rangle = n_{X}|n_{X}\rangle,
$$

where $X = A$ or *B*. Note that in this representation, X_+ and X_- are mutual adjoints and X_0 is selfadjoint. It is easily seen that in the $q \uparrow 1$ limit, the representation defined by the actions [\(4\)](#page-3-1) goes to the standard oscillator representation.

The Schwinger realization of the quantum algebra $U_q(\mathfrak{sl}_2)$ is constructed from the two *q*-oscillator algebras [\(3\)](#page-3-0) by taking [\[2\]](#page-20-2)

$$
J_{+} = q^{-\frac{A_0 + B_0 - 1}{4}} A_{+} B_{-},
$$
 $J_{-} = q^{-\frac{A_0 + B_0 - 1}{4}} A_{-} B_{+},$ $J_0 = \frac{A_0 - B_0}{2}.$

Indeed, it is easily verified using the relations [\(3\)](#page-3-0) that the operators J_{\pm} and J_0 satisfy the defining relations of $U_q(\mathfrak{sl}_2)$ which read

$$
[J_0, J_{\pm}] = \pm J_{\pm}, \qquad [J_+, J_-] = \frac{q^{J_0} - q^{-J_0}}{q^{1/2} - q^{-1/2}}.
$$

As can be verified directly using [\(4\)](#page-3-1), the orthonormal *q*-oscillator states defined by

$$
(5) \qquad |n\rangle_N := |n,N-n\rangle, \qquad n=0,1,\ldots,N,
$$

where N is a non-negative integer, support the standard unitary $(N + 1)$ -dimensional irreducible representations of $U_q(\mathfrak{sl}_2)$.

2.2 *q***-analogs of the BCH formulas and** *q***-rotation operators**

Consider the little and big *q*-exponential functions, respectively denoted by $e_q(z)$ and $E_q(z)$, which are defined as follows:

(6)
$$
e_q(z) = \sum_{n=0}^{\infty} \frac{z^n}{(q;q)_n} = \frac{1}{(z;q)_{\infty}}, \quad |z| < 1, \quad \text{and} \quad E_q(z) = \sum_{n=0}^{\infty} \frac{q^{\binom{n}{2}} z^n}{(q;q)_n} = (-z;q)_{\infty}.
$$

One clearly has $e_q(z)E_q(-z) = 1$. In the following, two *q*-analogs of the Baker–Campbell–Hausdorff relation shall be needed. The first relation is of the form

(7)
$$
E_q(\lambda X) Ye_q(-\lambda q^{\alpha} X) = \sum_{n=0}^{\infty} \frac{\lambda^n}{(q;q)_n} [X,Y]_n,
$$

where $[X, Y]_n$ is defined recursively through

$$
[X,Y]_0 = 1, \qquad [X,Y]_{n+1} = q^n X [X,Y]_n - q^{\alpha} [X,Y]_n X, \qquad n = 1,2,...
$$

The second relation reads

(8)
$$
e_q(\lambda X)YE_q(-\lambda q^{\alpha}X) = \sum_{n=0}^{\infty} \frac{\lambda^n}{(q;q)_n} [X,Y]_n',
$$

where $[X, Y]'_n$ is defined recursively by

 $[X, Y]_0' = 1,$ $[X, Y]_{n+1}' = X[X, Y]_n' - q^{n+\alpha} [X, Y]_n' X,$ $n = 0, 1, 2, \ldots$

The identities (7) , (8) can be verified in a straightforward manner by expanding the *q*-exponentials in power series using [\(6\)](#page-3-3). For $\alpha = 0$, these relations are given in [\[3\]](#page-20-3), among others.

Let θ be a real number such that $|\theta| < 1$ and let $U_{AB}(\theta)$ be the operator defined as

(9)
$$
U_{AB}(\theta) =
$$

\n $e_q^{1/2} \left(\theta^2 q^{-A_0} \right) e_q \left(\theta (1-q) q^{-(A_0+B_0)/2} A_+ B_- \right) E_q \left(-\theta (1-q) q^{-(A_0+B_0)/2} A_- B_+ \right) E_q^{1/2} \left(-\theta^2 q^{-B_0} \right).$

This operator will be called a "*q*-rotation" operator. This term, coined by Zhedanov in [\[29\]](#page-22-3), comes from the fact that $U_{AB}(\theta)$ can be viewed as a *q*-analog of an $SU(2)$ element obtained via the exponential map from the algebra to the group; see [\[8,](#page-21-17) [29\]](#page-22-3) for more details. As shown in [\[8\]](#page-21-17), the operator [\(9\)](#page-4-1) is unitary: it satisfies the relations $U^{\dagger}U = 1$ and $UU^{\dagger} = 1$. The unitarity property of $U_{AB}(\theta)$ follows directly from the formulas

$$
e_q(\alpha A - B_+)e_q\left(\frac{\alpha\beta}{(1-q)^2}q^{B_0}\right)e_q(\beta A + B_-) = e_q(\beta A + B_-)e_q\left(\frac{\alpha\beta}{(1-q)^2}q^{A_0}\right)e_q(\alpha A - B_+),
$$

and

$$
E_q(\gamma A_+ B_-) E_q\left(-\frac{\gamma\delta}{(1-q)^2}q^{B_0}\right) E_q(\delta A_- B_+) = E_q(\delta A_- B_+) E_q\left(-\frac{\gamma\delta}{(1-q)^2}q^{A_0}\right) E_q(\gamma A_+ B_-),
$$

where $\alpha, \beta, \gamma, \delta$ are constants or central elements. These formulas can be proved using the *q*-BCH relations [\(7\)](#page-3-2) and [\(8\)](#page-4-0) as well as elementary properties of the *q*-exponential functions; see [\[8\]](#page-21-17).

2.3 Matrix elements and univariate quantum *q***-Krawtchouk polynomials**

Let us now recall how the one-variable quantum *q*-Krawtchouk polynomials arise in this setting. The matrix elements of the unitary *q*-rotation operator [\(9\)](#page-4-1) in the basis [\(5\)](#page-3-4) are defined as

$$
(10) \quad \xi_{n,x}^{(N)}(\theta) = N \langle n | U_{AB}(\theta) | x \rangle_N,
$$

where *n* and *x* take values in $\{0,1,...,N\}$. The matrix elements [\(10\)](#page-4-2) can be written as [\[8\]](#page-21-17)

$$
(11) \quad \xi_{n,x}(\theta) = \omega_x^{(N)}(\theta) \sigma_n^{(N)}(\theta) k_n(x, \theta^{-2}, N; q),
$$

where $k_n(x, p, N; q)$ are the quantum q -Krawtchouk polynomials defined in [\(2\)](#page-2-1). In [\(11\)](#page-4-3), the coefficient $\sigma_n^{(N)}$ has the expression

(12)
$$
\sigma_n^{(N)}(\theta) = \frac{(-1)^n q^{-\binom{n}{2}}}{(q^{-N};q)_n} \sqrt{\binom{N}{n}_{q} \frac{\theta^{2n} q^{-nN}}{(\theta^2 q^{-n};q)_n}},
$$

where

$$
\binom{N}{n}_q = \frac{(q;q)_N}{(q;q)_n(q;q)_{N-n}},
$$

stands for the *q*-binomial coefficient. The expression for $\omega_x^{(N)}(\theta)$, which is defined as the matrix element $_N \langle 0|U_{AB}(\theta)|x\rangle_N$, reads

(13)
$$
\omega_x^{(N)}(\theta) := N \langle 0 | U_{AB}(\theta) | x \rangle_N = (-1)^x q^{\binom{x}{2}} \sqrt{\binom{N}{x}_{q} \frac{(\theta^2 q^{-N}; q)_N}{(\theta^2 q^{-N}; q)_x} \theta^{2x} q^{-xN}}.
$$

From the algebraic interpretation [\(11\)](#page-4-3) of the one-variable quantum *q*-Krawtchouk polynomials, one can derive the main properties of these polynomials [\[8\]](#page-21-17). In what follows, the duality relation

(14)
$$
\xi_{n,x}^{(N)}(\theta) = \xi_{N-x,N-n}^{(N)}(\theta),
$$

which follows from the reality of the matrix elements [\(10\)](#page-4-2), the identity $U_{AB}^{\dagger}(\theta) = U_{AB}^{-1}(\theta)$ and the $\text{observation that } U_{AB}^{-1}(\theta) = U_{BA}(\theta), \text{ shall prove particularly useful.}$

3 A model for the two-variable *q***-Krawtchouk polynomials**

In this section, the algebraic model for the two-variable *q*-Krawtchouk polynomials, which involves a three-dimensional *q*-rotation acting on the states of three *q*-oscillators, is constructed. The orthogonality relation and the duality property of the polynomials are derived from the model.

3.1 The model

Consider the algebra generated by three mutually commuting *q*-oscillators with generators X_0, X_{\pm} , $X \in \{A, B, C\}$, satisfying the commutation relations [\(3\)](#page-3-0) and consider its representation defined by the actions [\(4\)](#page-3-1) on the three-fold tensor product space spanned by the orthonormal basis vectors $|n_A, n_B, n_C\rangle = |n_A\rangle \otimes |n_B\rangle \otimes |n_C\rangle$, where n_A , n_B and n_C are non-negative integers. Let N be a positive integer and introduce the basis states $|n_1,n_2\rangle_N$ defined as

$$
(15) \quad |n_1,n_2\rangle_N := |n_1,n_2,N-n_1-n_2\rangle, \qquad n_1,n_2 \in \{0,1,\ldots,N\}, \qquad n_1+n_2 \le N.
$$

For a given N, the states [\(15\)](#page-5-0) support an $(N+1)(N+2)/2$ -dimensional irreducible representation of the quantum algebra $U_q(\mathfrak{sl}_3)$ realized with three independent *q*-oscillators; see [\[14\]](#page-21-18). Following [\(9\)](#page-4-1), the unitary operator $U_{XY}(\theta)$ effecting the *q*-rotation in the XY "plane" is defined as

(16)
$$
U_{XY}(\theta) =
$$

\n $e_q^{1/2} \left(\theta^2 q^{-X_0} \right) e_q \left(\theta (1-q) q^{-(X_0+Y_0)/2} X_+ Y_- \right) E_q \left(-\theta (1-q) q^{-(X_0+Y_0)/2} X_- Y_+ \right) E_q^{1/2} \left(-\theta^2 q^{-Y_0} \right).$

3.2 Matrix elements and bivariate *q***-Krawtchouk polynomials**

Consider the matrix elements

(17)
$$
\Xi_{n_1,n_2}^{(N)}(x_1,x_2;\theta,\phi) = N \langle n_1,n_2 | U_{BC}(\phi) U_{AB}(\theta) | x_1, x_2 \rangle_N,
$$

of the operator associated to a particular *q*-rotation in three dimensions. To ease the notation, $\Xi_{n_1,n_2}^{(N)}(x_1,x_2;\theta,\phi)$ shall sometimes be written simply as $\Xi_{n_1,n_2}^{(N)}(x_1,x_2)$. An explicit expression for the matrix elements [\(17\)](#page-5-1) can be obtained by inserting a resolution of the identity in between the two *q*-rotation operators $U_{BC}(\phi)$ and $U_{AB}(\theta)$. This leads to

(18)
$$
\Xi_{n_1,n_2}^{(N)}(x_1,x_2;\theta,\phi) = \sum_{\substack{s,t=0 \ s+t \leq N}}^{N} \frac{N}{n_1,n_2} |U_{BC}(\phi)|s,t \rangle_N \frac{N}{s} \langle s,t|U_{AB}(\theta)|x_1,x_2 \rangle_N.
$$

Since the *q*-rotation operator $U_{BC}(\phi)$ leaves invariant the first index when acting on the basis vector $|n_1,n_2\rangle$ _N, it follows from the definition [\(10\)](#page-4-2) and the orthonormality of the basis that

(19)
$$
N \langle n_1, n_2 | U_{BC}(\phi) | s, t \rangle_N = \delta_{n_1, s} \xi_{n_2, t}^{(N-n_1)}(\phi).
$$

Similarly, when acting on the state $|x_1, x_2\rangle_N$, the operator $U_{AB}(\theta)$ leaves invariant the sum of the first two indices $x_1 + x_2$ and one can thus write

(20)
$$
N \langle s, t | U_{AB}(\theta) | x_1, x_2 \rangle_N = \delta_{s+t, x_1+x_2} \xi_{s, x_1}^{(x_1+x_2)}(\theta).
$$

Upon using [\(19\)](#page-6-0) and [\(20\)](#page-6-1) in [\(18\)](#page-6-2), the double summation cancels and one finds

$$
(21) \quad \Xi_{n_1,n_2}^{(N)}(x_1,x_2;\theta,\phi) = \xi_{n_1,x_1}^{(x_1+x_2)}(\theta) \, \xi_{n_2,x_1+x_2-n_1}^{(N-n_1)}(\phi).
$$

With the explicit expression [\(11\)](#page-4-3) of the one-dimensional *q*-rotation matrix elements in terms of the univariate quantum *q*-Krawtchouk polynomials, one has

$$
(22) \quad \Xi_{n_1,n_2}^{(N)}(x_1,x_2;\theta,\phi) = \omega_{x_1}^{(x_1+x_2)}(\theta)\sigma_{n_1}^{(x_1+x_2)}(\theta)k_{n_1}(x_1,\theta^{-2},x_1+x_2;q) \times \omega_{x_1+x_2-n_1}^{(N-n_1)}(\phi)\sigma_{n_2}^{(N-n_1)}(\phi)k_{n_2}(x_1+x_2-n_1,\phi^{-2},N-n_1;q).
$$

Comparing [\(22\)](#page-6-3) with [\(1\)](#page-2-2), it is seen that the matrix elements $\Xi_{n_1,n_2}^{(N)}(x_1,x_2)$ can be expressed in terms of the two-variable q-Krawtchouk polynomials $\mathbf{K}_{n_1,n_2}(x_1,x_2;\alpha_1,\underline{\alpha_2};N)$ with parameters $\alpha_1 = \theta$ and $\alpha_2 = \phi$. One must however verify that the matrix elements $\Xi_{n_1,n_2}^{(N)}(x_1,x_2)$ factorize as a product of the matrix element $_N(0,0|U_{BC}(\phi)U_{AB}(\theta)|x_1,x_2\rangle_N$, a normalization factor depending only on the degree indices n_1 and n_2 , and the polynomials $\mathbf{K}_{n_1,n_2}(x_1,x_2;a_1,a_2;N)$ themselves. A direct, though technical, calculation shows that this is indeed the case and that one can write the matrix elements $\Xi_{n_1,n_2}^{(N)}(x_1,x_2)$ as

$$
(23) \quad \Xi_{n_1,n_2}^{(N)}(x_1,x_2;\theta,\phi) = W_{x_1,x_2}^{(N)}(\theta,\phi) \Sigma_{n_1,n_2}^{(N)}(\theta,\phi) \mathbf{K}_{n_1,n_2}(x_1,x_2;\theta,\phi;N),
$$

where $W^{(N)}_{x_1,x_2}(\theta,\phi)$ is given by

$$
(24) \quad W_{x_1,x_2}^{(N)}(\theta,\phi) = N \langle 0,0 | U_{BC}(\phi) U_{AB}(\theta) | x_1, x_2 \rangle_N
$$

$$
= (-1)^{x_2} \left[(-1)^{N-x_1} (\theta^2 q^{-1})^{x_1+x_2} q^{\binom{x_1}{2}} \binom{N}{x_1,x_2} q^{(\theta^{-2}q)} q^{x_2} q^{x_1} q^{N-x_1-x_2} \phi^{2N} q^{-N(N+1)/2} \right]^{1/2},
$$

where the normalization factor $\Sigma_{n_1,n_2}^{(N)}(\theta,\phi)$ has the expression

$$
(25) \quad \Sigma_{n_1,n_2}^{(N)}(\theta,\phi) = q^{-\binom{n_1}{2} - \binom{n_2}{2}} (-1)^{n_1} \phi^{-n_1} \frac{(q;q)_{N-n_1-n_2}}{(q;q)_N} \times \left[(-1)^{n_1+n_2} \binom{N}{n_1,n_2} q^{\frac{N(n_1+n_2)+2n_1+n_2-n_1n_2 - \binom{n_1}{2} - \binom{n_2}{2}}{q\phi^{-2};q)_{n_1}} \right]^{1/2},
$$

and where $\mathbf{K}_{n_1,n_2}(x_1,x_2;\bm{a}_1,a_2;N)$ are the two-variable quantum q -Krawtchouk polynomials defined in [\(1\)](#page-2-2). In [\(24\)](#page-6-4) and [\(25\)](#page-6-5), $\binom{N}{x,y}_q$ stands for the *q*-trinomial coefficients

$$
\binom{N}{x,y}_q = \frac{(q;q)_N}{(q;q)_x(q;q)_y(q;q)_{N-x-y}}.
$$

Remark. Let us note here that in [\(17\)](#page-5-1), the choice of the ordering of the two *q*-rotations to form a *q*-rotation in three space is important. For example, consider the matrix elements

$$
N \langle n_1, n_2 | U_{AC}(\phi) U_{BC}(\theta) | x_1, x_2 \rangle_N = \xi_{n_1, x_1}^{(N-n_2)}(\theta) \xi_{n_2, x_2}^{(N-x_1)}(\phi),
$$

which can be seen as *q*-analogs of the matrix elements leading to Geronimo and Iliev's two-variable Krawtchouk polynomials [\[11\]](#page-21-3). It can be verified directly that with this choice of ordering one has a factorization the form

$$
N \langle n_1, n_2 | U_{AC}(\phi) U_{BC}(\theta) | x_1, x_2 \rangle_N = N \langle 0, 0 | U_{AC}(\phi) U_{BC}(\theta) | x_1, x_2 \rangle_N P_{n_1, n_2}(x_1, x_2),
$$

where $P_{n_1,n_2}(x_1,x_2)$, with $P_{0,0}(x_1,x_2) = 1$, are not polynomial in the proper variables.

3.3 Orthogonality relations

The orthogonality relation for the two-variable *q*-Krawtchouk polynomials immediately follows from the unitarity of the *q*-rotation $U_{BC}(\phi)U_{AB}(\theta)$, the orthonormality of the basis states and the reality of the matrix elements [\(17\)](#page-5-1). One has

$$
N\langle n_1, n_2 | m_1, m_2 \rangle_N = N\langle n_1, n_2 | U_{BC}(\phi) U_{AB}(\theta) U_{AB}^{\dagger}(\theta) U_{BC}^{\dagger}(\phi) | m_1, m_2 \rangle_N
$$

\n
$$
= \sum_{\substack{x_1, x_2 = 0 \\ x_1 + x_2 \le N}}^N N\langle n_1, n_2 | U_{BC}(\phi) U_{AB}(\theta) | x_1, x_2 \rangle_N N\langle x_1, x_2 | U_{AB}^{\dagger}(\theta) U_{BC}^{\dagger}(\phi) | m_1, m_2 \rangle_N
$$

\n
$$
= \sum_{\substack{x_1, x_2 = 0 \\ x_1 + x_2 \le N}}^N \Xi_{n_1, n_2}^{(N)}(x_1, x_2; \theta, \phi) \Xi_{m_1, m_2}^{(N)}(x_1, x_2; \theta, \phi) = \delta_{n_1 m_1} \delta_{n_2 m_2}.
$$

(26)

N

Upon introducing [\(23\)](#page-6-6) in the above relation, one finds that the two-variable quantum *q*-Krawtchouk polynomials satisfy the orthogonality relation

$$
(27) \sum_{\substack{x_1,x_2=0\\x_1+x_2\leq N}}^{N} \Delta_{x_1,x_2}^{(N)}(\theta,\phi) \mathbf{K}_{n_1,n_2}(x_1,x_2;\theta,\phi;N) \mathbf{K}_{m_1,m_2}(x_1,x_2;\theta,\phi;N) = H_{n_1,n_2}^{(N)}(\theta,\phi) \delta_{n_1m_1} \delta_{n_2m_2},
$$

where $\Delta(x_1, x_2)^{(N)}(\theta, \phi) = [W^{(N)}_{x_1, x_2}(\theta, \phi)]^2$ and $H^{(N)}_{n_1, n_2}(\theta, \phi) = [\Sigma^{(N)}_{n_1, n_2}(\theta, \phi)]^{-2}$. From [\(26\)](#page-7-0), it is easily seen that one can also write a dual orthogonality relation of the form

$$
(28)\quad \sum_{\substack{n_1,n_2=0\\n_1+n_2\leq N}}^{N} \widetilde{\Delta}_{n_1,n_2}^{(N)}(\theta,\phi) \mathbf{K}_{n_1,n_2}(x_1,x_2;\theta,\phi;N) \mathbf{K}_{n_1,n_2}(y_1,y_2;\theta,\phi;N) = \widetilde{H}_{x_1,x_2}^{(N)}(\theta,\phi) \delta_{x_1y_1} \delta_{x_2y_2},
$$

where $\widetilde{\Delta}^{(N)}_{n_1,n_2}(\theta,\phi) = [\Sigma^{(N)}_{n_1,n_2}(\theta,\phi)]^2$ and $\widetilde{H}^{(N)}_{x_1,x_2} = [W^{(N)}_{x_1,x_2}(\theta,\phi)]^{-2}$.

3.4 Duality

The matrix elements [\(17\)](#page-5-1) satisfy a duality relation that allows to interchange the roles of the degree indices n_1 and n_2 with those of the variables indices x_1 and x_2 . This duality relation is obtained by combining the duality relation [\(14\)](#page-5-2) and the expression [\(21\)](#page-6-7) for the matrix elements. One has

$$
\begin{split} \Xi_{n_1,n_2}^{(N)}(x_1,x_2;\theta,\phi) &= \xi_{n_1,x_1}^{(x_1+x_2)}(\theta) \,\xi_{n_2,x_1+x_2-n_1}^{(N-n_1)}(\phi) = \xi_{x_2,x_1+x_2-n_1}^{(x_1+x_2)}(\theta) \xi_{N-x_1-x_2,N-n_1-n_2}^{(N-n_1)}(\phi) \\ &= \xi_{N-x_1-x_2,N-n_1-n_2}^{(N-n_1)}(\phi) \xi_{x_2,x_1+x_2-n_1}^{(x_1+x_2)}(\theta) = \Xi_{N-x_1-x_2,x_2}^{(N)}(N-n_1-n_2,n_2;\phi,\theta), \end{split}
$$

where we have used [\(21\)](#page-6-7) in the first step and [\(14\)](#page-5-2) in the second step; note that the parameters θ and *φ* are interchanged in the duality transformation. In terms of the polynomials $\mathbf{K}_{n_1,n_2}(x_1,x_2;\theta,\phi;N),$ the duality relation [\(29\)](#page-8-0) becomes

$$
(30) \quad (-1)^{x_2+n_2} q^{x_1^2+x_2^2+x_1x_2+x_1-N(x_1+x_2-1)} \left(\frac{(q\theta^{-2};q)_{x_2}(q\phi^{-2};q)_{N-x_1-x_2}}{(q;q)_{x_1}} \right) \mathbf{K}_{n_1,n_2}(x_1,x_2;\theta,\phi;N)
$$

$$
= q^{n_1^2+n_2^2+n_1n_2-n_1-n_2-N(n_1+n_2)} \left(\frac{(q\theta^{-2};q)_{n_1}(q\phi^{-2};q)_{n_2}}{(q;q)_{N-n_1-n_2}} \right) \mathbf{K}_{N-x_1-x_2,x_2}(N-n_1-n_2,n_2;\phi,\theta;N).
$$

4 Structure relations

In this section, the *q*-BCH formulas [\(7\)](#page-3-2), [\(8\)](#page-4-0) and the algebraic model constructed in the previous section are used to obtain structure relations for the two-variable quantum *q*-Krawtchouk polynomials. Raising and lowering relations on the degrees and the variables are found.

4.1 Structure relations with respect to the index *n*¹

We first use the algebraic properties of the three-dimensional *q*-rotation operator $U_{BC}(\phi)U_{AB}(\theta)$ to obtain structures relations on the first degree index *n*1.

4.1.1 Raising relation in n_1

To get a raising relation with respect to the first degree index n_1 , consider the matrix element $N-1\langle n_1, n_2|A-U_{BC}(\phi)U_{AB}(\theta)|x_1, x_2\rangle_N$. On the one hand, using [\(4\)](#page-3-1) and [\(17\)](#page-5-1), one has

$$
(31) \quad N-1 \langle n_1, n_2 | A - U_{BC}(\phi) U_{AB}(\theta) | x_1, x_2 \rangle_N = \sqrt{\frac{1-q^{n_1+1}}{1-q}} \, \Xi^{(N)}_{n_1+1, n_2}(x_1, x_2; \theta, \phi).
$$

On the other hand, one can write

$$
(32) \quad N-1 \langle n_1, n_2 | A - U_{BC}(\phi) U_{AB}(\theta) | x_1, x_2 \rangle_N = N-1 \langle n_1, n_2 | U_{BC}(\phi) U_{AB}(q^{-1/2}\theta) \tilde{A}_- | x_1, x_2 \rangle_N,
$$

where $A_-\$ is of the form

(33)
$$
\tilde{A}_{-} = U_{AB}^{\dagger} (q^{-1/2} \theta) U_{BC}^{\dagger} (\phi) A_{-} U_{BC} (\phi) U_{AB} (\theta) = U_{AB}^{\dagger} (q^{-1/2} \theta) A_{-} U_{AB} (\theta).
$$

A direct calculation using the expression [\(16\)](#page-5-3) and the formulas [\(7\)](#page-3-2) and [\(8\)](#page-4-0) yields

$$
(34) \quad U_{AB}^{\dagger}(q^{-1/2}\theta) A_{-} U_{AB}(\theta) = \sqrt{1 - \theta^2 q^{-1} q^{-B_0}} A_{-} + \theta q^{-1/2} q^{(A_0 - B_0)/2} B_{-}.
$$

To obtain [\(34\)](#page-8-1) from the *q*-Baker–Campbell–Hausdorff relations, one should recall that $U_{XY}^{\dagger} = U_{XY}^{-1}$. Upon substituting [\(34\)](#page-8-1) in [\(32\)](#page-8-2), using the actions [\(4\)](#page-3-1) and comparing with [\(31\)](#page-8-3), one finds

$$
(35) \quad \sqrt{1-q^{n_1+1}} \, \Xi_{n_1+1,n_2}^{(N)}(x_1,x_2;\theta,\phi) = \sqrt{1-\theta^2 q^{-x_2-1}} \sqrt{1-q^{x_1}} \, \Xi_{n_1,n_2}^{(N-1)}(x_1-1,x_2;q^{-1/2}\theta,\phi) + \theta q^{(x_1-x_2)/2} \sqrt{1-q^{x_2}} \, \Xi_{n_1,n_2}^{(N-1)}(x_1,x_2-1;q^{-1/2}\theta,\phi).
$$

Using the expression [\(23\)](#page-6-6) of the matrix elements in terms of the two-variable *q*-Krawtchouk polynomials, the relation [\(35\)](#page-9-0) takes the form

(36)
$$
q^{-n_1}\mathbf{K}_{n_1+1,n_2}(x_1,x_2;q^{1/2}\theta,\phi;N) = q^{-(x_1+x_2)}(1-q^{x_1})(1-\theta^{-2}q^{x_2})\mathbf{K}_{n_1,n_2}(x_1-1,x_2;\theta,\phi;N-1) - (1-q^{-x_2})\mathbf{K}_{n_1,n_2}(x_1,x_2-1;\theta,\phi;N-1).
$$

4.1.2 Lowering relation in *n*¹

A lowering relation on the first degree index n_1 can be obtained by considering instead the matrix ${\rm element} \; _{N+1} \langle n_1,n_2 | A_{+} U_{BC}(\phi) U_{AB} (q^{-1/2}\theta) | x_1,x_2 \rangle_N.$ One has

$$
(37) \quad N+1 \langle n_1, n_2 | A_+ U_{BC}(\phi) U_{AB}(q^{-1/2}\theta) | x_1, x_2 \rangle_N = \sqrt{\frac{1-q^{n_1}}{1-q}} \, \Xi_{n_1-1,n_2}^{(N)}(x_1, x_2; q^{-1/2}\theta, \phi),
$$

and also

$$
(38) \quad N+1 \langle n_1, n_2 | A_+ U_{BC}(\phi) U_{AB}(q^{-1/2}\theta) | x_1, x_2 \rangle_N = N+1 \langle n_1, n_2 | U_{BC}(\phi) U_{AB}(\theta) \tilde{A}_+ | x_1, x_2 \rangle_N,
$$

with

$$
(39) \quad \widetilde{A}_{+} = U_{AB}^{\dagger}(\theta)U_{BC}^{\dagger}(\phi)A_{+}U_{BC}(\phi)U_{AB}(q^{-1/2}\theta).
$$

It is directly seen that [\(39\)](#page-9-1) is the complex conjugate of [\(33\)](#page-8-4). Consequently, we have

(40)
$$
\tilde{A}_+ = A_+ \sqrt{1 - \theta^2 q^{-1} q^{-B_0}} + \theta q^{-1/2} B_+ q^{(A_0 - B_0)/2}.
$$

Upon substituting [\(40\)](#page-9-2) in [\(38\)](#page-9-3), using the actions [\(4\)](#page-3-1) and comparing with [\(37\)](#page-9-4), one finds

$$
(41) \quad \sqrt{1-q^{n_1}}\Xi_{n_1-1,n_2}^{(N)}(x_1,x_2;\theta,\phi) = \sqrt{1-q^{x_1+1}}\sqrt{1-\theta^2q^{-x_2}}\,\Xi_{n_1,n_2}^{(N+1)}(x_1+1,x_2;q^{1/2}\theta,\phi) \n+ \theta q^{(x_1-x_2)/2}\sqrt{1-q^{x_2+1}}\,\Xi_{n_1,n_2}^{(N+1)}(x_1,x_2+1;q^{1/2}\theta,\phi),
$$

where we have taken $\theta \to q^{1/2} \theta$. Using [\(23\)](#page-6-6), this gives

(42)
$$
\frac{q^{n_1}(1-q^{n_1})}{\theta^2} \mathbf{K}_{n_1-1,n_2}(x_1,x_2;\theta,\phi;N) = q^{x_1+2} \mathbf{K}_{n_1,n_2}(x_1,x_2+1;q^{1/2}\theta,\phi;N+1) - q^{x_1+2} \mathbf{K}_{n_1,n_2}(x_1+1,x_2;q^{1/2}\theta,\phi;N+1).
$$

4.2 Structure relations with respect to the index *n*²

We now look for structure relations on the second degree index n_2 .

4.2.1 Raising relation in n_2

To get a raising relation with respect the second degree index n_2 , consider the matrix element $N-1$ $\langle n_1,n_2|q^{A_0/2}B_{-}U_{BC}(\phi)U_{AB}(\theta)|x_1,x_2\rangle_N.$ One has

$$
(43) \quad N-1 \langle n_1, n_2 | q^{A_0/2} B - U_{BC}(\phi) U_{AB}(\theta) | x_1, x_2 \rangle_N = q^{n_1/2} \sqrt{\frac{1 - q^{n_2 + 1}}{1 - q}} \, \Xi_{n_1, n_2 + 1}^{(N)}(x_1, x_2; \theta, \phi).
$$

Proceeding as in the previous subsection, one writes

$$
(44) \quad N-1 \langle n_1, n_2 | q^{A_0/2} B - U_{BC}(\phi) U_{AB}(\theta) | x_1, x_2 \rangle_N = N-1 \langle n_1, n_2 | U_{BC}(q^{-1/2}\phi) U_{AB}(\theta) \tilde{B} - | x_1, x_2 \rangle_N,
$$

where

(45)
$$
\tilde{B}_{-} = U_{AB}^{\dagger}(\theta)U_{BC}^{\dagger}(q^{-1/2}\phi)q^{-A_0/2}B_{-}U_{BC}(\phi)U_{AB}(\theta).
$$

Upon using [\(34\)](#page-8-1) with the *A*s replaced by *B*s and the *B*s replaced by *C*s, one finds

$$
\widetilde{B}_{-}=U_{AB}^{\dagger}(\theta)\left\{q^{(A_0+B_0)/2}\left[\sqrt{1-\phi^2q^{-1}q^{-C_0}}q^{-B_0/2}B_{-}+\phi q^{-1/2}q^{-C_0/2}C_{-}\right]\right\}U_{AB}(\theta).
$$

It is obvious that $q^{A_0+B_0}$ commutes with $U_{AB}(\theta),$ and similarly for all the terms involving the generators C_0 , C_{\pm} . Using the conjugation formula

$$
U_{AB}^{\dagger}(\theta)q^{-B_0/2}B_{-}U_{AB}(\theta)=q^{-B_0/2}B_{-}\sqrt{1-\theta^2q^{-B_0}}-\theta q^{-B_0}A_{-}q^{-A_0/2},
$$

which can be obtained by a straightforward calculation with the help of the *q*-BCH formulas [\(7\)](#page-3-2), [\(8\)](#page-4-0), one obtains

$$
(46)\quad \widetilde{B}_{-}=q^{(A_0+B_0)/2}\Big[\sqrt{1-\phi^2q^{-1}q^{-C_0}}\Big(q^{-B_0/2}B_{-}\sqrt{1-\theta^2q^{-B_0}}-\theta q^{-B_0}A_{-}q^{-A_0/2}\Big)+\phi q^{-1/2}q^{-C_0/2}C_{-}\Big].
$$

Upon inserting [\(46\)](#page-10-0) in [\(44\)](#page-10-1), and comparing with [\(43\)](#page-10-2), one finds

$$
\begin{aligned}\n & (47) \quad q^{n_1/2} \sqrt{1 - q^{n_2 + 1}} \, \Xi_{n_1, n_2 + 1}^{(N)}(x_1, x_2; \theta, \phi) = \phi q^{x_1 + x_2} q^{-N/2} \sqrt{1 - q^{N - x_1 - x_2}} \, \Xi_{n_1, n_2}^{(N-1)}(x_1, x_2; \theta, q^{-1/2} \phi) \\
 &- q^{-1/2} \theta q^{-x_2/2} \sqrt{1 - \phi^2 q^{x_1 + x_2 - N - 1}} \sqrt{1 - q^{x_1}} \, \Xi_{n_1, n_2}^{(N-1)}(x_1 - 1, x_2; \theta, q^{-1/2} \phi) \\
 &+ q^{x_1/2} \sqrt{1 - q^{x_2}} \sqrt{1 - \theta^2 q^{-x_2}} \sqrt{1 - \phi^2 q^{x_1 + x_2 - N - 1}} \, \Xi_{n_1, n_2}^{(N-1)}(x_1, x_2 - 1; \theta, q^{-1/2} \phi).\n \end{aligned}
$$

Using the expression [\(23\)](#page-6-6), one obtains

(48)
$$
q^{N-n_2}\mathbf{K}_{n_1,n_2+1}(x_1,x_2;\theta,q^{1/2}\phi;N)=(1-q^{x_1})(1-\phi^{-2}q^{N-x_1-x_2})\mathbf{K}_{n_1,n_2}(x_1-1,x_2;\theta,\phi;N-1) +q^{x_1}(1-q^{x_2})(1-\phi^{-2}q^{N-x_1-x_2})\mathbf{K}_{n_1,n_2}(x_1,x_2-1;\theta,\phi;N-1) +q^{x_1+x_2}(1-q^{N-x_1-x_2})\mathbf{K}_{n_1,n_2}(x_1,x_2;\theta,\phi;N-1).
$$

The relations [\(36\)](#page-9-5) and [\(48\)](#page-10-3) can be combined to generate $\mathbf{K}_{n_1,n_2}(x_1,x_2; a_1,a_2; N)$ recursively.

4.2.2 Lowering relation in n_2

In order to obtain a lowering relation in the second degree index n_2 , one considers the matrix ele- $\text{ments } {}_{N+1} \langle n_1, n_2 | B_+ q^{A_0/2} U_{BC}(q^{-1/2} \phi) U_{AB}(\theta) | x_1, x_2 \rangle_N.$ One has

$$
(49) \quad N+1 \langle n_1, n_2 | B_+ q^{A_0/2} U_{BC}(q^{-1/2} \phi) U_{AB}(\theta) | x_1, x_2 \rangle_N = q^{n_1/2} \sqrt{\frac{1-q^{n_2}}{1-q}} \, \Xi_{n_1, n_2-1}^{(N)}(x_1, x_2; \theta, q^{-1/2} \phi; N),
$$

and also

$$
(50) \quad N+1 \langle n_1, n_2 | B_+ q^{A_0/2} U_{BC}(q^{-1/2} \phi) U_{AB}(\theta) | x_1, x_2 \rangle_N = N+1 \langle n_1, n_2 | U_{BC}(\phi) U_{AB}(\theta) \tilde{B}_+ | x_1, x_2 \rangle_N,
$$

where \widetilde{B}_+ is given by

(51)
$$
\tilde{B}_{+} = U_{AB}^{\dagger}(\theta)U_{BC}(\phi)B_{+}q^{A_0/2}U_{BC}(q^{-1/2}\phi)U_{AB}(\theta).
$$

It is seen that [\(51\)](#page-11-0) is the complex conjugate of [\(45\)](#page-10-4). Taking the complex conjugate of [\(46\)](#page-10-0), one finds

$$
(52)\quad \widetilde{B}_{+} = \left[\left(\sqrt{1 - \theta^2 q^{-B_0}} B_+ q^{-B_0/2} - \theta q^{-A_0/2} A_+ q^{-B_0} \right) \sqrt{1 - \phi^2 q^{-1} q^{-C_0}} + \phi q^{-1/2} C_+ q^{-C_0/2} \right] q^{(A_0 + B_0)/2}.
$$

Proceeding in the same manner as before leads to the relation

$$
(53) \quad q^{n_1/2}\sqrt{1-q^{n_2}} \Xi_{n_1,n_2-1}^{(N)}(x_1,x_2;\theta,q^{-1/2}\phi) =
$$

\n
$$
-\theta q^{-1/2}q^{-x_2/2}\sqrt{1-\phi^2q^{-1}q^{x_1+x_2-N}}\sqrt{1-q^{x_1+1}} \Xi_{n_1,n_2}^{(N+1)}(x_1+1,x_2;\theta,\phi)
$$

\n
$$
+q^{x_1/2}\sqrt{1-\phi^2q^{-1}q^{x_1+x_2-N}}\sqrt{1-\theta^2q^{-1}q^{-x_2}}\sqrt{1-q^{x_2+1}} \Xi_{n_1,n_2}^{(N+1)}(x_1,x_2+1;\theta,\phi)
$$

\n
$$
+\phi q^{-1/2}q^{x_1+x_2}q^{-N/2}\sqrt{1-q^{N-x_1-x_2+1}} \Xi_{n_1,n_2}^{(N+1)}(x_1,x_2;\theta,\phi).
$$

In terms of the bivariate *q*-Krawtchouk polynomials, this relation amounts to

(54)
$$
\frac{q^{n_1+n_2}(1-q^{n_2})}{\phi^2} \mathbf{K}_{n_1,n_2-1}(x_1,x_2;\theta,q^{-1/2}\phi;N) = -\theta^2 q^{x_1} \mathbf{K}_{n_1,n_2}(x_1+1,x_2;\theta,\phi;N+1) + \theta^2 q^{x_1}(1-\theta^{-2}q^{x_2+1}) \mathbf{K}_{n_1,n_2}(x_1,x_2+1;\theta,\phi;N+1) + q^{x_1+x_2+1} \mathbf{K}_{n_1,n_2}(x_1,x_2;\theta,\phi;N+1).
$$

4.3 Structure relations with respect to the index *x*¹

Proceeding in a "dual" fashion, one can obtain structure relations in the first variable index *x*1.

4.3.1 Lowering relation in *x*¹

In order to find a lowering relation in first variable index x_1 , one considers the matrix element $N^{\langle n}1,n_2|U_{BC}(\phi)U_{AB}(\theta)\,q^{-A_0/2}A_{-}|x_1,x_2\rangle_{N+1}.$ One the one hand, one has

$$
(55) \quad N \langle n_1, n_2 | U_{BC}(\phi) U_{AB}(\theta) q^{-A_0/2} A_- | x_1, x_2 \rangle_{N+1} = q^{-(x_1-1)/2} \sqrt{\frac{1-q^{x_1}}{1-q}} \, \Xi_{n_1, n_2}^{(N)}(x_1-1, x_2; \theta, \phi; N).
$$

On the other hand, one can write

$$
(56)\quad N \langle n_1, n_2 | U_{BC}(\phi) U_{AB}(\theta) q^{-A_0/2} A_- | x_1, x_2 \rangle_{N+1} = N \langle n_1, n_2 | \hat{A}_- U_{BC}(\phi) U_{AB}(\theta) | x_1, x_2 \rangle_{N+1},
$$

where

(57)
$$
\hat{A}_{-} = U_{BC}(\phi)U_{AB}(\theta) q^{-A_0/2} A_{-} U_{AB}^{\dagger}(\theta)U_{BC}^{\dagger}(\phi).
$$

To compute [\(57\)](#page-12-0), one first uses the conjugation formula

$$
(58) \quad U_{AB}(\theta)q^{-A_0/2}A - U_{AB}^{\dagger}(\theta) = q^{-A_0/2}A - \sqrt{1 - \theta^2 q^{-A_0}} - \theta q^{-A_0}B - q^{-B_0/2},
$$

which is obtained straightforwardly from the *q*-BCH formulas [\(7\)](#page-3-2) and [\(8\)](#page-4-0); this lead to

(59)
$$
\widehat{A}_{-} = U_{BC}(\phi) \left[q^{-A_0/2} A_{-} \sqrt{1 - \theta^2 q^{-A_0}} - \theta q^{-1/2} q^{-A_0} q^{-B_0/2} B_{-} \right] U_{BC}^{\dagger}(\phi).
$$

Using [\(58\)](#page-12-1) again with *A* replaced by *B* and *B* replaced by *C*, one finds

$$
(60)\quad \widehat{A}_{-} = \bigg\{ q^{-A_0/2} A_{-} \sqrt{1 - \theta^2 q^{-A_0}} - \theta q^{-1/2} q^{-A_0} \bigg[q^{-B_0/2} B_{-} \sqrt{1 - \phi^2 q^{-B_0}} - \phi q^{-1/2} q^{-B_0} q^{-C_0/2} C_{-} \bigg] \bigg\}.
$$

Upon substituting [\(60\)](#page-12-2) in [\(56\)](#page-11-1), using the actions [\(4\)](#page-3-1) and comparing with [\(55\)](#page-11-2), one finds

(61)
$$
q^{-(x_1-1)/2} \sqrt{1-q^{x_1}} \Xi_{n_1,n_2}^{(N)}(x_1-1,x_2;\theta,\phi) =
$$

\n
$$
q^{-n_1/2} \sqrt{1-q^{n_1+1}} \sqrt{1-\theta^2 q^{-1}q^{-n_1}} \Xi_{n_1+1,n_2}^{(N+1)}(x_1,x_2;\theta,\phi)
$$

\n
$$
-\theta q^{-1/2} q^{-n_1} q^{-n_2/2} \sqrt{1-q^{n_2+1}} \sqrt{1-\phi^2 q^{-1}q^{-n_2}} \Xi_{n_1,n_2+1}^{(N+1)}(x_1,x_2;\theta,\phi)
$$

\n
$$
+\theta \phi q^{-1} q^{-(n_1+n_2)/2} q^{-N/2} \sqrt{1-q^{N-n_1-n_2+1}} \Xi_{n_1,n_2}^{(N+1)}(x_1,x_2;\theta,\phi).
$$

In terms of the polynomials, this amounts to

(62)
$$
\frac{q(1-q^{-x_1})}{\theta^2} \mathbf{K}_{n_1,n_2}(x_1-1,x_2;\theta,\phi;N) = q^{-2n_1} \mathbf{K}_{n_1+1,n_2}(x_1,x_2;\theta,\phi;N+1) + \phi^2 q^{-n_1-2n_2-1} \mathbf{K}_{n_1,n_2+1}(x_1,x_2;\theta,\phi;N+1) - \phi^2 q^{-N-2} (1-q^{N-n_1-n_2+1}) \mathbf{K}_{n_1,n_2}(x_1,x_2;\theta,\phi;N+1).
$$

4.3.2 Raising relation in *x*¹

A raising relation in the first variable index *x*¹ can be obtained by considering the matrix element $N^{\langle n_1,n_2|U_{BC}(\phi)U_{AB}(\theta)A_{+}q^{-A_0/2}|x_1,x_2\rangle_{N-1}.$ One has

$$
(63) \quad N \langle n_1, n_2 | U_{BC}(\phi) U_{AB}(\theta) A_+ q^{-A_0/2} | x_1, x_2 \rangle_{N-1} = q^{-x_1/2} \sqrt{\frac{1 - q^{x_1 + 1}}{1 - q}} \, \Xi_{n_1, n_2}^{(N)}(x_1 + 1, x_2; \theta, \phi),
$$

and also

$$
(64) \quad N \langle n_1, n_2 | U_{BC}(\phi) U_{AB}(\theta) A_+ q^{-A_0/2} | x_1, x_2 \rangle_{N-1} = N \langle n_1, n_2 | \hat{A}_+ U_{BC}(\phi) U_{AB}(\theta) | x_1, x_2 \rangle_{N+1},
$$

where

(65)
$$
\hat{A}_{+} = U_{BC}(\phi)U_{AB}(\theta) A_{+}q^{-A_0/2}U_{AB}^{\dagger}(\theta)U_{BC}^{\dagger}(\phi).
$$

The expression for [\(65\)](#page-12-3) is obtained directly by taking the complex conjugate of [\(60\)](#page-12-2). Substituting the result in [\(64\)](#page-12-4) and comparing with [\(63\)](#page-12-5), one finds

$$
(66) \quad q^{-x_1/2} \sqrt{1 - q^{x_1+1}} \Xi_{n_1,n_2}^{(N)}(x_1+1,x_2;\theta,\phi) = q^{1/2} q^{-n_1/2} \sqrt{1 - \theta^2 q^{-n_1}} \sqrt{1 - q^{n_1}} \Xi_{n_1-1,n_2}^{(N-1)}(x_1,x_2;\theta,\phi)
$$

$$
- \theta q^{-n_1} q^{-n_2/2} \sqrt{1 - q^{n_2}} \sqrt{1 - \phi^2 q^{-n_2}} \Xi_{n_1,n_2-1}^{(N-1)}(x_1,x_2;\theta,\phi)
$$

$$
+ \theta \phi q^{-1/2} q^{-(N+n_1+n_2)/2} \sqrt{1 - q^{N-n_1-n_2}} \Xi_{n_1,n_2}^{(N-1)}(x_1,x_2;\theta,\phi).
$$

Using the expression [\(23\)](#page-6-6) for the matrix elements, this relations translates to

(67)
$$
\mathbf{K}_{n_1,n_2}(x_1+1,x_2;\theta,\phi;N) = q^{-1}(1-q^{n_1})(1-\theta^{-2}q^{n_1})\mathbf{K}_{n_1-1,n_2}(x_1,x_2;\theta,\phi;N-1) +q^{-1-n_1}(1-q^{n_2})(1-\phi^{-2}q^{n_2})\mathbf{K}_{n_1,n_2-1}(x_1,x_2;\theta,\phi;N-1)+q^{-n_1-n_2}\mathbf{K}_{n_1,n_2}(x_1,x_2;\theta,\phi;N-1).
$$

4.4 Structure relations with respect to the index *x*²

We now obtain structure relations with respect to the second variable index x_2 .

4.4.1 Lowering relation in *x*²

In order to obtain a lowering relation with respect to the second variable index x_2 , consider the $\text{matrix element}\ _{N}\langle n_{1},n_{2}|U_{BC}(\phi)U_{AB}(\theta)\,q^{C_{0}/2}B_{-}|x_{1},x_{2}\rangle_{N+1}.$ One has

$$
(68)\quad N\langle n_1,n_2|U_{BC}(\phi)U_{AB}(\theta) q^{C_0/2}B_{-}|x_1,x_2\rangle_{N+1} = q^{(N-x_1-x_2+1)/2}\sqrt{\frac{1-q^{x_2}}{1-q}}\,\Xi_{n_1,n_2}^{(N)}(x_1,x_2-1;\theta,\phi),
$$

and also

$$
(69) \quad N \langle n_1, n_2 | U_{BC}(\phi) U_{AB}(\theta) q^{C_0/2} B_- | x_1, x_2 \rangle_{N+1} = N \langle n_1, n_2 | \hat{B}_- U_{BC}(\phi) U_{AB}(q^{1/2}\theta) | x_1, x_2 \rangle_{N+1},
$$

where

(70)
$$
\hat{B}_- = U_{BC}(\phi)U_{AB}(\theta) q^{C_0/2} B_- U_{AB}^{\dagger} (q^{1/2} \theta) U_{BC}^{\dagger}(\phi)
$$
.

To determine the expression for \widehat{B}_- , one needs the conjugation relation

(71)
$$
U_{AB}(\theta) B - U_{AB}^{\dagger}(q^{1/2}\theta) = \sqrt{1 - \theta^2 q^{-A_0}} B_{-} + \theta q^{(B_0 - A_0)/2} A_{-},
$$

which is again obtained from the *q*-BCH relations. Using this relation, [\(70\)](#page-13-0) becomes

$$
(72)\quad \widehat{B}_{-}=U_{BC}(\phi)\left\{q^{(B_0+C_0)/2}\left[\sqrt{1-\theta^2q^{-A_0}}q^{-B_0/2}B_{-}+\theta q^{-A_0/2}A_{-}\right]\right\}U_{BC}^{\dagger}(\phi).
$$

Upon using [\(58\)](#page-12-1) in the above (with $A \rightarrow B$ and $B \rightarrow C$), one finds

$$
(73) \quad \widehat{B}_{-} = \left\{ q^{(B_0+C_0)/2} \right\} \left\{ \sqrt{1-\theta^2 q^{-A_0}} \left[q^{-B_0/2} B_{-} \sqrt{1-\phi^2 q^{-B_0}} - \phi q^{-B_0} C_{-} q^{-C_0/2} \right] + \theta q^{-A_0/2} A_{-} \right\}.
$$

Substituting the above expression in [\(69\)](#page-13-1) and comparing with [\(68\)](#page-13-2), there comes

$$
(74) \quad q^{(N-x_1-x_2+1)/2}\sqrt{1-q^{x_2}} \equiv_{n_1,n_2}^{(N)}(x_1,x_2-1;\theta,\phi) = \theta q^{N/2} q^{-n_1} \sqrt{1-q^{n_1+1}} \equiv_{n_1+1,n_2}^{(N+1)}(x_1,x_2;q^{1/2}\theta,\phi) +q^{(N-n_1-n_2)/2} \sqrt{1-\theta^2 q^{-n_1}} \sqrt{1-q^{n_2+1}} \sqrt{1-\phi^2 q^{-1} q^{-n_2}} \equiv_{n_1,n_2+1}^{(N+1)}(x_1,x_2;q^{1/2}\theta,\phi) - \phi q^{-1/2} q^{-n_2/2} \sqrt{1-\theta^2 q^{-n_1}} \sqrt{1-q^{N-n_1-n_2+1}} \equiv_{n_1,n_2}^{(N+1)}(x_1,x_2;q^{1/2}\theta,\phi).
$$

In terms of the polynomials $\mathbf{K}_{n_1,n_2}(x_1,x_2;\theta,\phi;N)$ this relation is of the form

(75)
$$
\frac{q^{1-x_1-x_2}}{\theta^2}(1-q^{x_2})\mathbf{K}_{n_1,n_2}(x_1,x_2-1;\theta,\phi;N) = q^{1-2n_1}\mathbf{K}_{n_1+1,n_2}(x_1,x_2;q^{1/2}\theta,\phi;N+1) + \phi^2q^{-n_1-2n_2}(1-\theta^{-2}q^{n_1})\mathbf{K}_{n_1,n_2+1}(x_1,x_2;q^{1/2}\theta,\phi;N+1) -\phi^2q^{-N-1}(1-q^{N-n_1-n_2+1})(1-\theta^{-2}q^{n_1})\mathbf{K}_{n_1,n_2}(x_1,x_2;q^{1/2}\theta,\phi;N+1).
$$

4.4.2 Raising relation in *x*²

To obtain a raising relation in the second variable index x_2 , one considers the matrix element *N* $\langle n_1, n_2 | U_{BC}(\phi) U_{AB}(q^{1/2}\theta) B_{+} q^{C_0/2} | x_1, x_2 \rangle_{N-1}.$ On the one hand, one has

$$
(76)\quad N\langle n_1,n_2|U_{BC}(\phi)U_{AB}(q^{1/2}\theta)B_+q^{C_0/2}|x_1,x_2\rangle_{N-1}=q^{(N-x_1-x_2-1)/2}\sqrt{\frac{1-q^{x_2+1}}{1-q}}\,\Xi_{n_1,n_2}^{(N)}(x_1,x_2+1;\theta,\phi),
$$

and on the other hand one can write

$$
(77) \quad N \langle n_1, n_2 | U_{BC}(\phi) U_{AB}(q^{1/2}\theta) B_+ q^{C_0/2} | x_1, x_2 \rangle_{N-1} = N \langle n_1, n_2 | \hat{B}_+ U_{BC}(\phi) U_{AB}(\theta) | x_1, x_2 \rangle_{N-1},
$$

where

(78)
$$
\hat{B}_+ = U_{BC}(\phi)U_{AB}(q^{1/2}\theta)B_+q^{C_0/2}U_{AB}^{\dagger}(\theta)U_{BC}^{\dagger}(\phi).
$$

Since $\widehat{B}_+ = \widehat{B}^{\dagger}_-$, one directly finds

$$
(79) \quad q^{(N-x_1-x_2-1)/2}\sqrt{1-q^{x_2+1}}\,\Xi_{n_1,n_2}^{(N)}(x_1,x_2+1;q^{1/2}\theta,\phi) = \theta q^{1/2}q^{-n_1}q^{N/2}\sqrt{1-q^{n_1}}\,\Xi_{n_1-1,n_2}^{(N-1)}(x_1,x_2;\theta,\phi) +q^{(N-n_1-n_2)/2}\sqrt{1-\phi^2q^{-n_2}}\sqrt{1-q^{n_2}}\sqrt{1-\theta^2q^{-n_1}}\,\Xi_{n_1,n_2-1}^{(N-1)}(x_1,x_2;\theta,\phi) -\phi q^{-1/2}q^{-n_2/2}\sqrt{1-q^{N-n_1-n_2}}\sqrt{1-\theta^2q^{-n_1}}\,\Xi_{n_1,n_2}^{(N-1)}(x_1,x_2;\theta,\phi).
$$

 $\text{In terms of the polynomials } \mathbf{K}_{n_1,n_2}(x_1,x_2;\theta,\phi;N), \text{this gives}$

(80)
$$
\mathbf{K}_{n_1,n_2}(x_1,x_2+1;q^{1/2}\theta,\phi;N) = q^{-1}(1-q^{n_1})\mathbf{K}_{n_1-1,n_2}(x_1,x_2;\theta,\phi;N-1) +q^{-n_1-1}(1-\phi^{-2}q^{n_2})(1-q^{n_2})\mathbf{K}_{n_1,n_2-1}(x_1,x_2;\theta,\phi;N-1)+q^{-(n_1+n_2)}\mathbf{K}_{n_1,n_2}(x_1,x_2;\theta,\phi;N-1).
$$

5 Bispectrality

In this section, the algebraic model for the two-variable *q*-Krawtchouk polynomials is used to derive their recurrence relations and their eigenvalue equations. These relations coincide with those of [\[16\]](#page-21-14).

5.1 Difference equations

To derive the two eigenvalue equations that determine the polynomials $\mathbf{K}_{n_1,n_2}(x_1,x_2;\alpha_1,\alpha_2;N)$, one could find suitable combinations of the raising and lowering relations on the degree indices n_1 and *n*² obtained in the previous section. However, it will prove more instructive to proceed directly by finding conjugation formulas for the appropriate operators.

5.1.1 First difference equation

Consider the matrix element $_N\langle n_1,n_2|q^{A_0}\,U_{BC}(\phi)U_{AB}(\theta)|x_1,x_2\rangle_N.$ On the one hand, one has

$$
(81) \quad N \langle n_1, n_2 | q^{A_0} U_{BC}(\phi) U_{AB}(\theta) | x_1, x_2 \rangle_N = q^{n_1} \Xi_{n_1, n_2}^{(N)}(x_1, x_2; \theta, \phi).
$$

On the other hand, one can write

$$
(82) \quad N \langle n_1, n_2 | q^{A_0} U_{BC}(\phi) U_{AB}(\theta) | x_1, x_2 \rangle_N = N \langle n_1, n_2 | U_{BC}(\phi) U_{AB}(\theta) \overline{q^{A_0}} | x_1, x_2 \rangle_N,
$$

where

(83)
$$
\widehat{q^{A_0}} = U_{AB}^{\dagger}(\theta)U_{BC}^{\dagger}(\phi) q^{A_0} U_{BC}(\phi)U_{AB}(\theta) = U_{AB}^{\dagger}(\theta) q^{A_0} U_{AB}(\theta).
$$

The conjugation $U_{AB}^{\dagger}(\theta) \, q^{A_0} \, U_{AB}(\theta)$ can be performed with the help of the *q*-BCH formulas. As this computation is not straightforward, we present the details of the calculation. Using the explicit expression [\(16\)](#page-5-3) for the *q*-rotation operator, one writes

$$
\begin{split} \widehat{q^{A_{0}}} &= e_{q}^{1/2} \left(\theta^{2} q^{-B_{0}} \right) e_{q} \left(\theta (1-q) q^{-(A_{0}+B_{0})/2} A_{-} B_{+} \right) E_{q} \left(- \theta (1-q) q^{-(A_{0}+B_{0})/2} A_{+} B_{-} \right) \\ &\times E_{q}^{1/2} \left(- \theta^{2} q^{-A_{0}} \right) \left\{ q^{A_{0}} \right\} e_{q}^{1/2} \left(\theta^{2} q^{-A_{0}} \right) \\ &\times e_{q} \left(\theta (1-q) q^{-(A_{0}+B_{0})/2} A_{+} B_{-} \right) E_{q} \left(- \theta (1-q) q^{-(A_{0}+B_{0})/2} A_{-} B_{+} \right) E_{q}^{1/2} \left(- \theta^{2} q^{-B_{0}} \right). \end{split}
$$

The first conjugation is trivial and one can immediately write

$$
\begin{split} \widehat{q^{A_{0}}} &= e_{q}^{1/2} \left(\theta^{2} q^{-B_{0}} \right) e_{q} \left(\theta (1-q) q^{-(A_{0}+B_{0})/2} A_{-} B_{+} \right) \\ &\times E_{q} \left(- \theta (1-q) q^{-(A_{0}+B_{0})/2} A_{+} B_{-} \right) q^{A_{0}} \, e_{q} \left(\theta (1-q) q^{-(A_{0}+B_{0})/2} A_{+} B_{-} \right) \\ &\times E_{q} \left(- \theta (1-q) q^{-(A_{0}+B_{0})/2} A_{-} B_{+} \right) E_{q}^{1/2} \left(- \theta^{2} q^{-B_{0}} \right). \end{split}
$$

The next step is performed using the identity $q^{A_0}e_q(\lambda A_+)$ = $e_q(q\lambda A_+)q^{A_0}$; one then finds

$$
\widehat{q^{A_0}} = e_q^{1/2} \left(\theta^2 q^{-B_0} \right) J_1 J_2 E_q^{1/2} \left(-\theta^2 q^{-B_0} \right),
$$

where

$$
(84) \quad J_1 = e_q \left(\theta (1-q) q^{-(A_0+B_0)/2} A_- B_+ \right) \left[1 - \theta (1-q) q^{-(A_0+B_0)/2} A_+ B_- \right] E_q \left(-\theta (1-q) q^{-(A_0+B_0)/2} A_- B_+ \right),
$$

and

$$
(85) \quad J_2 = e_q \left(\theta (1-q) q^{-(A_0+B_0)/2} A - B_+ \right) \left[q^{A_0} \right] E_q \left(-\theta (1-q) q^{-(A_0+B_0)/2} A - B_+ \right).
$$

The expression for [\(84\)](#page-15-0) can be obtained from the conjugation formula

$$
e_q(\alpha A - B_+)A_+B_-E_q(-\alpha A - B_+) = A_+B_- + \frac{\alpha}{(1-q)^2}q^{A_0} - \frac{\alpha}{(1-q)^2}\frac{1}{1-\alpha A_-B_+}q^{B_0},
$$

which follows straightforwardly from [\(8\)](#page-4-0). One thus has

$$
J_1 = 1 - \theta (1 - q) q^{-(A_0 + B_0)/2} \left[A_+ B_- + \frac{\theta q^{-(A_0 + B_0)/2}}{(1 - q)} q^{A_0} - \frac{\theta q^{-(A_0 + B_0)/2}}{(1 - q)} \frac{1}{1 - \theta (1 - q) q^{-(A_0 + B_0)/2} A_- B_+} q^{B_0} \right].
$$

One easily finds

$$
J_2 = q^{A_0} \left[1 - \theta (1 - q) q^{-(A_0 + B_0)/2} A_- B_+ \right].
$$

Performing simplifications on the product of J_1 and J_2 using the commutation relations [\(3\)](#page-3-0), one finds

$$
\widehat{q^{A_0}} = e_q^{1/2} \left(\theta^2 q^{-B_0} \right) \left\{ q^{A_0} [1 - \theta^2 q^{-B_0}] - \theta (1 - q) q^{(A_0 - B_0)/2} [1 - \theta^2 q^{-B_0}] A_- B_+ \right.\left. - \theta (1 - q) A_+ B_- q^{(A_0 - B_0)/2} + q^{A_0} \left(\theta^2 - q^{-1} \theta^2 q^{-B_0} + \theta^2 q^{-1} q^{-(A_0 + B_0)} \right) \right\} E_q^{1/2} \left(-\theta^2 q^{-B_0} \right).
$$

The last conjugation is easily carried out; the final result is

$$
(86) \quad \widehat{q^{A_0}} = q^{A_0} \left[1 - (1 + q^{-1}) \theta^2 q^{-B_0} + \theta^2 + q^{-1} \theta^2 q^{-(A_0 + B_0)} \right]
$$

$$
- \theta (1 - q) q^{(A_0 - B_0)/2} \sqrt{1 - \theta^2 q^{-B_0}} A_- B_+ - \theta (1 - q) A_+ B_- \sqrt{1 - \theta^2 q^{-B_0}} q^{(A_0 - B_0)/2}.
$$

Upon using [\(86\)](#page-16-0) in [\(82\)](#page-15-1), using the actions [\(4\)](#page-3-1) and comparing with [\(81\)](#page-15-2), one finds that the matrix $\Xi_{n_1,n_2}^{(N)}(x_1,x_2;\theta,\phi) = \Xi_{n_1,n_2}^{(N)}(x_1,x_2)$ satisfy the eigenvalue equation

$$
(87) \quad q^{n_1} \Xi_{n_1,n_2}^{(N)}(x_1,x_2) = \left[q^{x_1}(1+\theta^2) + q^{-1}\theta^2 q^{-x_2} - (1+q^{-1})\theta^2 q^{x_1-x_2}\right] \Xi_{n_1,n_2}^{(N)}(x_1,x_2)
$$

$$
- \theta q^{-1} q^{(x_1-x_2)/2} \left[(1-q^{x_2+1})(1-q^{x_1})(1-\theta^2 q^{-(x_2+1)}) \right]^{1/2} \Xi_{n_1,n_2}^{(N)}(x_1-1,x_2+1)
$$

$$
- \theta q^{(x_1-x_2)/2} \left[(1-\theta^2 q^{-x_2})(1-q^{x_2})(1-q^{x_1+1}) \right]^{1/2} \Xi_{n_1,n_2}^{(N)}(x_1+1,x_2-1).
$$

Upon using the expression [\(23\)](#page-6-6) for the matrix elements, one finds that the two-variable quantum q -Krawtchouk polynomials $\mathbf{K}_{n_1,n_2}(x_1,x_2;\theta,\phi;N)$ satisfy the 3-point eigenvalue equation

$$
(88) \quad q^{n_1} \mathbf{K}_{n_1,n_2}(x_1,x_2) = \left[q^{x_1}(1+\theta^2) + q^{-1}\theta^2 q^{-x_2} - (1+q^{-1})\theta^2 q^{x_1-x_2}\right] \mathbf{K}_{n_1,n_2}(x_1,x_2) + (1-\theta^2 q^{-x_2-1})(1-q^{x_1}) \mathbf{K}_{n_1,n_2}(x_1-1,x_2+1) + \theta^2 q^{x_1-x_2}(1-q^{x_2}) \mathbf{K}_{n_1,n_2}(x_1+1,x_2-1).
$$

5.1.2 Second difference equation

Another difference equation for the matrix elements $\Xi_{n_1,n_2}^{(N)}(x_1,x_2)$ and the two-variable Krawtchouk polynomials can be obtained from the matrix element $\frac{N}{n_1,n_2|q^{A_0+B_0}}\frac{U_{BC}(\phi)U_{AB}(\theta)|x_1,x_2\rangle_N.$ One can write this matrix elements as

$$
(89) \quad N \langle n_1, n_2 | q^{A_0 + B_0} U_{BC}(\phi) U_{AB}(\theta) | x_1, x_2 \rangle_N = q^{n_1 + n_2} \Xi_{n_1, n_2}^{(N)}(x_1, x_2; \theta, \phi).
$$

but also as

$$
(90) \quad N \langle n_1, n_2 | q^{A_0 + B_0} U_{BC}(\phi) U_{AB}(\theta) | x_1, x_2 \rangle_N = N \langle n_1, n_2 | U_{BC}(\phi) U_{AB}(\theta) \overline{q^{A_0 + B_0}} | x_1, x_2 \rangle_N,
$$

where

$$
(91) \quad \widehat{q^{A_0+B_0}} = U_{AB}^{\dagger}(\theta)U_{BC}^{\dagger}(\phi) \, q^{A_0+B_0} \, U_{BC}(\phi)U_{AB}(\theta).
$$

Using the result [\(86\)](#page-16-0) with the *A*s replaced by *B*s and the *B*s replaced by *C*s, one has

$$
(92) \quad \widehat{q^{A_0+B_0}} = U_{AB}^{\dagger}(\theta) \, q^{A_0+B_0} \left\{ \left[1 - (1+q^{-1})\phi^2 q^{-C_0} + \phi^2 + q^{-1} \phi^2 q^{-(B_0+C_0)} \right] \right. \\ \left. - \phi(1-q)q^{-(B_0+C_0)/2} \sqrt{1-\phi^2 q^{-C_0}} B_- C_+ - \phi q^{-1} (1-q) B_+ C_- \sqrt{1-\phi^2 q^{-C_0}} q^{-(B_0+C_0)/2} \right\} U_{AB}(\theta).
$$

To complete this calculation, one needs to use the conjugation formula

$$
U_{AB}^{\dagger}(\theta)q^{-B_0}U_{AB}(\theta) = q^{-B_0}\Big\{ \Big[1 - (1+q^{-1})\theta^2 q^{-B_0} + \theta^2 + q^{-1}\theta^2 q^{-(A_0+B_0)} \Big] - \theta(1-q)q^{-(A_0+B_0)/2}\sqrt{1-\theta^2 q^{-B_0}}A_-B_+ - \theta q^{-1}(1-q)A_+B_-\sqrt{1-\theta^2 q^{-B_0}}q^{-(A_0+B_0)/2} \Big\},\,
$$

which can be obtained in a fashion similar to the way [\(83\)](#page-15-3) was derived, as well as the formula

$$
U_{AB}^{\dagger}(\theta)q^{-B_0/2}B_{-}U_{AB}(\theta)=q^{-B_0/2}B_{-}\sqrt{1-\theta^2q^{-B_0}}-\theta q^{-B_0}A_{-}q^{-A_0/2},
$$

and its complex conjugate, which are easily obtained from the *q*-BCH relations. The final result is

$$
(93) \quad \widehat{q^{A_0+B_0}} = q^{A_0+B_0} \left\{ 1 - (1+q^{-1})\phi^2 q^{-C_0} + \phi^2 \right\} + q^{-1}\phi^2 q^{A_0-C_0} \left[1 - (1+q^{-1})\theta^2 q^{-B_0} + \theta^2 + q^{-1}\theta^2 q^{-(A_0+B_0)} \right] - q^{-1}\phi^2 \theta (1-q)q^{(A_0-B_0-2C_0)/2} \sqrt{1-\theta^2 q^{-B_0}} A_- B_+ - q^{-1}\phi^2 \theta (1-q)A_+ B_- \sqrt{1-\theta^2 q^{-B_0}} q^{(A_0-B_0-2C_0)/2} - \phi (1-q)q^{(2A_0+B_0)/2} B_- \sqrt{1-\theta^2 q^{-B_0}} q^{-C_0/2} \sqrt{1-\phi^2 q^{-C_0}} C_+ - \phi (1-q) \sqrt{1-\theta^2 q^{-B_0}} B_+ q^{(2A_0+B_0)/2} C_- \sqrt{1-\phi^2 q^{-C_0}} q^{-C_0/2} + \theta \phi q^{-1} (1-q) A_- q^{A_0/2} q^{-C_0/2} \sqrt{1-\phi^2 q^{-C_0}} C_+ + \theta \phi q^{-1} (1-q) q^{A_0/2} A_+ C_- \sqrt{1-\phi^2 q^{-C_0}} q^{-C_0/2} .
$$

As a consequence, we have the following 7-point eigenvalue equation

$$
(94) \quad q^{n_1+n_2} \Xi_{n_1,n_2}^{(N)}(x_1, x_2) = \left[q^{x_1+x_2} \left(1 - (1 + q^{-1}) \phi^2 q^{x_1+x_2-N} + \phi^2 \right) \right.
$$

\n
$$
+ q^{-1} \phi^2 q^{2x_1+x_2-N} \left(1 - (1 + q^{-1}) \theta^2 q^{-x_2} + \theta^2 + q^{-1} \theta^2 q^{-(x_1+x_2)} \right) \right] \Xi_{n_1,n_2}^{(N)}(x_1, x_2)
$$

\n
$$
- q^{-2} \theta \phi^2 q^{(3x_1+x_2-2N)/2} \sqrt{(1 - q^{x_2+1})(1 - q^{x_1})(1 - \theta^2 q^{-(x_2+1)})} \Xi_{n_1,n_2}^{(N)}(x_1 - 1, x_2 + 1)
$$

\n
$$
- q^{-1} \theta \phi^2 q^{(3x_1+x_2-2N)/2} \sqrt{(1 - q^{x_2})(1 - q^{x_1+1})(1 - \theta^2 q^{-x_2})} \Xi_{n_1,n_2}^{(N)}(x_1 + 1, x_2 - 1)
$$

\n
$$
- \phi q^{(3x_1+2x_2-N-2)/2} \sqrt{(1 - q^{x_2})(1 - \theta^2 q^{-x_2})(1 - q^{N-x_1-x_2+1})(1 - \phi^2 q^{x_1+x_2-N-1})} \Xi_{n_1,n_2}^{(N)}(x_1, x_2 - 1)
$$

\n
$$
- \phi q^{(3x_1+2x_2-N)/2} \sqrt{(1 - q^{x_2+1})(1 - \theta^2 q^{-(x_2+1)})(1 - q^{N-x_1-x_2})(1 - \phi^2 q^{x_1+x_2-N})} \Xi_{n_1,n_2}^{(N)}(x_1, x_2 + 1)
$$

\n
$$
+ q^{-1} \theta \phi q^{(2x_1+x_2-N-1)/2} \sqrt{(1 - q^{x_1})(1 - q^{N-x_1-x_2+1})(1 - \phi^2 q^{x_1+x_2-N-1})} \Xi_{n_1,n_2}^{(N)}(x_1 - 1, x_2)
$$

\n
$$
+ q^{-1} \theta \phi q^{(2x
$$

In terms of the polynomials, this relation amounts to

$$
(95) \quad q^{n_1+n_2} \mathbf{K}_{n_1,n_2}(x_1,x_2) = \left[q^{x_1+x_2} \left(1 - (1+q^{-1}) \phi^2 q^{x_1+x_2-N} + \phi^2 \right) \right.
$$

\n
$$
+ q^{-1} \phi^2 q^{2x_1+x_2-N} \left(1 - (1+q^{-1}) \theta^2 q^{-x_2} + \theta^2 + q^{-1} \theta^2 q^{-(x_1+x_2)} \right) \Big| \mathbf{K}_{n_1,n_2}(x_1,x_2)
$$

\n
$$
+ \phi^2 q^{x_1+x_2-N-1} (1-q^{x_1}) (1-\theta^2 q^{-x_2-1}) \mathbf{K}_{n_1,n_2}(x_1-1,x_2+1)
$$

\n
$$
+ \theta^2 \phi^2 q^{2x_1-N-1} (1-q^{x_2}) \mathbf{K}_{n_1,n_2}(x_1+1,x_2-1)
$$

\n
$$
+ q^{x_1} (1-q^{x_2}) (1-\phi^2 q^{x_1+x_2-N-1}) \mathbf{K}_{n_1,n_2}(x_1,x_2-1)
$$

\n
$$
- \phi^2 q^{x_1+x_2} (1-q^{x_1+x_2-N}) (1-\theta^2 q^{-x_2-1}) \mathbf{K}_{n_1,n_2}(x_1,x_2+1)
$$

\n
$$
+ (1-q^{x_1}) (1-\phi^2 q^{x_1+x_2-N-1}) \mathbf{K}_{n_1,n_2}(x_1-1,x_2) - \theta^2 \phi^2 q^{x_1-1} (1-q^{x_1+x_2-N}) \mathbf{K}_{n_1,n_2}(x_1+1,x_2).
$$

The eigenvalue equations [\(88\)](#page-16-1) and [\(95\)](#page-17-0) coincide with those of [\[16\]](#page-21-14).

5.2 Recurrence relations

We shall now derive recurrence relations for the matrix elements $\Xi_{n_1,n_2}^{(N)}(x_1,x_2)$ as well as for the two-variable polynomials $\mathbf{K}_{n_1,n_2}(x_1,x_2)$.

5.2.1 First recurrence relation

To obtain a first recurrence relation for the matrix elements $\Xi_{n_1,n_2}^{(N)}(x_1,x_2)$, one can start from the eigenvalue equation [\(87\)](#page-16-2) and apply the duality relation [\(29\)](#page-8-0) to exchange the roles played by the degree indices n_1 , n_2 and the variable indices x_1 , x_2 . Applying [\(29\)](#page-8-0) on [\(87\)](#page-16-2) and performing the change of variables and parameters

$$
x_2 \leftrightarrow n_2, \qquad N - n_1 - n_2 \leftrightarrow N - x_1 - x_2, \qquad \theta \leftrightarrow \phi,
$$

one finds

$$
q^{-x_1-x_2} \Xi_{n_1,n_2}^{(N)}(x_1,x_2) = -\phi q^{-(n_1+2n_2-N)/2} \sqrt{(1-q^{N-n_1-n_2+1})(1-q^{n_2})(1-\phi^2 q^{-n_2})} \Xi_{n_1,n_2-1}^{(N)}(x_1,x_2)
$$

+
$$
q^{-(n_1+2n_2+1)} \left[q^{n_2+1}(1+\phi^2) + \phi^2 q^{n_1+n_2-N} - (1+q) \phi^2 \right] \Xi_{n_1,n_2}^{(N)}(x_1,x_2)
$$

-
$$
\phi q^{-(n_1+2n_2+N-2)/2} \sqrt{(1-q^{n_2+1})(1-q^{N-n_1-n_2})(1-\phi^2 q^{-n_2-1})} \Xi_{n_1,n_2+1}^{(N)}(x_1,x_2).
$$

In terms of the polynomials, one finds

$$
(96) \quad q^{-x_1-x_2} \mathbf{K}_{n_1,n_2}(x_1,x_2) = q^{-(n_1+2n_1+1)} \Big[q^{n_2+1} (1+\phi^2) + \phi^2 q^{n_1+n_2-N} - (1+q) \phi^2 \Big] \mathbf{K}_{n_1,n_2}(x_1,x_2) + q^{-(n_1+1)} (1-q^{n_2}) (1-q^{n_1+n_2-N-1}) (1-\phi^2 q^{-n_2}) \mathbf{K}_{n_1,n_2-1}(x_1,x_2) - q^{-(3n_2+n_1+1)} \phi^2 \mathbf{K}_{n_1,n_2+1}(x_1,x_2).
$$

5.2.2 Second recurrence relation

To obtain the second recurrence relation, one applies the same procedure as above on the eigenvalue equation [\(94\)](#page-17-1) with the result

$$
(97) \quad q^{-x_1} \Xi_{n_1,n_2}^{(N)}(x_1,x_2) = q^{-(2n_1+n_2+N+1)/2} \theta \phi \sqrt{(1-q^{n_1})(1-q^{N-n_1-n_2+1})(1-\theta^2 q^{-n_1})} \Xi_{n_1-1,n_2}^{(N)}(x_1,x_2)
$$

\n
$$
-q^{-(3n_1+n_2)/2} \theta \sqrt{(1-q^{n_2+1})(1-q^{n_1})(1-\theta^2 q^{-n_1})(1-\phi^2 q^{-n_2-1})} \Xi_{n_1-1,n_2+1}^{(N)}(x_1,x_2)
$$

\n
$$
-q^{-1/2} q^{-(2n_1+n_2)} \theta^2 \phi \sqrt{(1-q^{-n_2})(1-q^{n_1+n_2-N-1})(1-\phi^2 q^{-n_2})} \Xi_{n_1,n_2-1}^{(N)}(x_1,x_2)
$$

\n
$$
+q^{-2n_1} \left[q^{n_1} (1+\theta^2-q^{-1-n_1}(1+q)\theta^2) + q^{-1-n_2}\theta^2 (1+\phi^2+q^{n_1-N-1}\phi^2-q^{-n_2-1}(1+q)\phi^2) \right] \Xi_{n_1,n_2}^{(N)}(x_1,x_2)
$$

\n
$$
-q^{-(3n_1+4n_2+N+4)/2} \theta^2 \phi \sqrt{(1-q^{N-n_1-n_2})(1-q^{n_2+1})(1-\phi^2 q^{-n_2-1})} \Xi_{n_1,n_2+1}^{(N)}(x_1,x_2)
$$

\n
$$
-q^{-(3n_1+n_2+1)/2} \theta \sqrt{(1-q^{n_2})(1-q^{n_1+1})(1-\theta^2 q^{-n_1-1})(1-\phi^2 q^{-n_2})} \Xi_{n_1+1,n_2-1}^{(N)}(x_1,x_2)
$$

\n
$$
+q^{(2n_1+n_2+N+3)/2} \theta \phi \sqrt{(1-q^{n_1+1})(1-q^{N-n_1-n_2})(1-\theta^2 q^{-n_1-1})} \Xi_{n_1+1,n_2}^{(N)}(x_1,x_2).
$$

In terms of the polynomials, one finds

$$
(98) \quad q^{-x_1} \mathbf{K}_{n_1,n_2}(x_1,x_2) = q^{-2n_1} \Big\{ q^{n_1} (1+\theta^2 - q^{-n_1-1}(1+q)\theta^2) + q^{-n_2-1}\theta^2 (1+\phi^2 + q^{n_1-N-1}\phi^2 - q^{-n_2-1}(1+q)\phi^2) \Big\} \mathbf{K}_{n_1,n_2}(x_1,x_2) + q^{-(n_2+2)}\phi^2 (1-q^{n_1})(1-q^{n_1+n_2-N-1})(1-\theta^2 q^{-n_1}) \mathbf{K}_{n_1-1,n_2}(x_1,x_2) + q^{-2(n_2+1)}\phi^2 (1-q^{n_1})(1-\theta^2 q^{-n_1}) \mathbf{K}_{n_1-1,n_2+1}(x_1,x_2) + q^{-(2n_1+2)}\theta^2 (1-q^{n_2})(1-q^{n_1+n_2-N-1})(1-\phi^2 q^{-n_2}) \mathbf{K}_{n_1,n_2-1}(x_1,x_2) - q^{-(3n_2+2n_1+2)}\theta^2 \phi^2 \mathbf{K}_{n_1,n_2+1}(x_1,x_2) - q^{-(3n_1+n_2+1)}\theta^2 \mathbf{K}_{n_1+1,n_2}(x_1,x_2) + \theta^2 \phi^{-2} q^{n_2-3n_1-2} (1-q^{n_2})(1-\phi^2 q^{-n_2}) \mathbf{K}_{n_1+1,n_2-1}(x_1,x_2).
$$

Remark. Let us note that the structure relations with respect to the variable indices x_1, x_2 derived in Subsections (4.3) and (4.4) cannot be obtained by the same duality argument from the structure relations with respect to the degree indices n_1, n_2 given in Subsections (4.1) and (4.2).

6 Conclusion

In this paper, we have constructed an algebraic model for the two-variable quantum *q*-Krawtchouk polynomials. We have shown that these polynomials arise as matrix elements of a three-dimensional *q*-rotation operator expressed in terms of *q*-exponentials that act on the states of three independent *q*-oscillators. We have explained how this model can be used to derive the main properties of the polynomials: orthogonality relation, duality property, structure relations, eigenvalue equations and recurrence relations.

It is apparent that the approach presented here can be extended to any number variables; the only essential difficulty lies in the complicated notation required to describe multivariate polynomials. Nevertheless, let us exhibit how the multivariate quantum *q*-Krawtchouk polynomials of Gasper and Rahman arise in the present framework. Consider $d+1$ independent copies of the *q*-oscillator algebra with generators $A_0^{(i)}$ $_{0}^{\left(i\right) },A_{\pm}^{\left(i\right) }$ and commutation relations

$$
[A_0^{(i)}, A_\pm^{(j)}] = \pm \delta_{ij} A_\pm^{(i)}, \qquad [A_-^{(i)}, A_+^{(i)}] = \delta_{ij} q^{A_0^{(i)}}, \qquad i = 1, 2, \dots, d+1.
$$

Let $\{n_1, n_2, \ldots, n_d\}$ and $\{x_1, x_2, \ldots, x_d\}$ be two sets of non-negative integers such that $\sum_{k=1}^d n_d \le N$ and $\sum_{k=1}^d x_k \leq N$, where N is a positive integer. We define $\mathbf{n} = (n_1, n_2, \ldots, n_{d+1})$ with $n_{d+1} = N - \sum_{k=1}^d n_d$, and similarly for **x**. We shall use the notation:

$$
|\mathbf{y}_k| = y_1 + y_2 + \cdots y_k.
$$

Note that by definition, we have $|\mathbf{y}_{d+1}| = N$ and $|\mathbf{y}_0| = 0$. Consider the representation of the algebra generated by the $d+1$ independent *q*-oscillators on the states

$$
|{\bf n}\rangle_N=\bigotimes_{k=1}^{d+1}|n_k\rangle,
$$

and defined by the following actions of the generators on the factors of the direct product:

$$
(99) \quad A_{+}^{(i)}|n_{i}\rangle = \sqrt{\frac{1-q^{n_{i}+1}}{1-q}}|n_{i}+1\rangle, \qquad A_{-}^{(i)}|n_{i}\rangle = \sqrt{\frac{1-q^{n_{i}}}{1-q}}|n_{i}-1\rangle, \qquad A_{0}^{(i)}|n_{i}\rangle = n_{i}|n_{i}\rangle.
$$

We introduce the *q*-rotation operator in the (i, j) plane which is defined as

$$
U_{i,j}(\theta) = e_q^{1/2} \left(\theta^2 q^{-A_0^{(i)}} \right) e_q \left(\theta (1-q) q^{-(A_0^{(i)} + A_0^{(j)})/2} A_+^{(i)} A_-^{(j)} \right) \\
\times E_q \left(- \theta (1-q) q^{-(A_0^{(i)} + A_0^{(j)})/2} A_-^{(i)} A_+^{(j)} \right) E_q^{1/2} \left(- \theta^2 q^{-A_0^{(j)}} \right).
$$

Let $\Xi_{\mathbf{n}}(\mathbf{x};\theta_1,\dots,\theta_d)$ be the following matrix elements:

$$
\Xi_{\mathbf{n}}(\mathbf{x};\theta_1,\cdots,\theta_d)=N\langle \mathbf{n}|\prod_{k=0}^{d-1}U_{d-k,d-k+1}(\theta_{d-k})\ket{\mathbf{x}}_N.
$$

A direct calculation shows that in terms of the one-variable matrix elements [\(10\)](#page-4-2), the matrix elements $\Xi_{\mathbf{n}}(\mathbf{x};\theta_1,\cdots,\theta_d)$ have the expression

$$
\Xi_{\mathbf{n}}(\mathbf{x};\theta_1,\cdots,\theta_d)=\prod_{k=1}^d\xi_{n_k,|\mathbf{x}_k|-|\mathbf{n}_{k-1}|}^{(|\mathbf{x}_{k+1}|-|\mathbf{n}_{k-1}|)}(\theta_k).
$$

In view of [\(11\)](#page-4-3), this means that these matrix elements are proportional to the multivariate quantum *q*-Krawtchouk polynomials $\mathbf{K}_n(\mathbf{x}; \theta_1, \dots, \theta_d; N)$ which read

$$
\mathbf{K_n}(\mathbf{x};\theta_1,\ldots,\theta_d;N)=\prod_{k=1}^d k_{n_k}\left(|\mathbf{x}_k|-|\mathbf{n}_{k-1}|,\frac{1}{\theta_k^2},|\mathbf{x}_{k+1}|-|\mathbf{n}_{k-1}|;q\right),
$$

where $k_n(x, p, N; q)$ are the one-variable quantum *q*-Krawtchouk polynomials given by [\(2\)](#page-2-1). With the above expression for $\Xi_n(\mathbf{x}; \theta_1, \dots, \theta_d)$, one can proceed with the determination of the properties of the multivariate *q*-Krawtchouk polynomials.

As explained previously, the results presented here can be viewed as a *q*-generalization of [\[10\]](#page-21-9) (in the special Tratnik case), where an algebraic interpretation of Griffiths' multivariate Krawtchouk polynomials was given in terms of the rotation group acting on oscillator states. The approach of [\[10\]](#page-21-9) was generalized in [\[6\]](#page-21-19) and [\[7\]](#page-21-20), where algebraic interpretations of the multivariate Charlier and Meixner polynomials involving the Euclidean and pseudo-rotation groups were found. In light of these results, it would be of interest to generalize the approach developed here to also obtain algebraic models for the multivariate *q*-Charlier and *q*-Meixner polynomials.

Acknowledgments

The authors would like to thank A. Zhedanov for stimulating discussions. VXG is supported by a postdoctoral fellowship from the Natural Sciences and Engineering Research Council of Canada (NSERC). The research of LV is supported in part by NSERC.

References

- [1] G. Andrews, R. Askey, and R. Roy. *Special functions*, volume 71 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, 2001.
- [2] L. C. Biedenharn. The quantum group *SUq*(2) and a *q*-analogue of the boson operator. *J. Phys. A*, 22:L873–L878, 1989.
- [3] R. Floreanini and L. Vinet. Automorphisms of the *q*-oscillator algebra and basic orthogonal polynomials. *Phys. Lett. A*, 180:393–401, 1993.
- [4] G. Gasper and M. Rahman. *Basic Hypergeometric Series*. Second Edition (Corrected and Extended). Cambridge University Press, 2004.
- [5] G. Gasper and M. Rahman. Some systems of multivariable orthogonal *q*-Racah polynomials. *Ramanujan J.*, 13:389–405, 2007.
- [6] V. X. Genest, H. Miki, L. Vinet, and A. Zhedanov. The multivariate Charlier polynomials as matrix elements of the Euclidean group representation on oscillator states. *J. Phys. A*, 47:215204, 2014.
- [7] V. X. Genest, H. Miki, L. Vinet, and A. Zhedanov. The multivariate Meixner polynomials as matrix elements of *SO*(*d*,1) representations on oscillator states . *J. Phys. A*, 47:045207, 2014.
- [8] V. X. Genest, S. Post, L. Vinet, G.-F. Yu, and A. Zhedanov. *q*-Rotations and Krawtchouk polynomials. *Ramanujan J.*, 2015. arXiv: 1408.5292v2.
- [9] V. X. Genest and L. Vinet. The multivariate Hahn polynomials and the singular oscillator. *J. Phys. A*, 47:455201, 2014.
- [10] V. X. Genest, L. Vinet, and A. Zhedanov. The multivariate Krawtchouk polynomials as matrix elements of the rotation group representations on oscillator states. *J. Phys. A*, 46:505203, 2013.
- [11] J. Geronimo and P. Iliev. Bispectrality of multivariable Racah-Wilson polynomials. *Constr. Approx.*, 31:417–457, 2010.
- [12] R. C. Griffiths. Orthogonal Polynomials on the multinomial distribution. *Aust. J. Stat.*, 3:27–35, 1971.
- [13] R. C. Griffiths and D. Spanò. Multivariate Jacobi and Laguerre polynomials, infinite-dimensional extensions, and their probabilistic connections with multivariate Hahn and Meixner polynomials. *Bernoulli*, 17:1095–1125, 2011.
- [14] T. Hayashi. *Q*-analogues of Clifford and Weyl Algebras—Spinor and Oscillator Representations of Quantum Enveloping Algebras. *Commun. Math. Phys.*, 127:129–144, 1990.
- [15] M. Hoare and M. Rahman. A Probablistic Origin for a New Class of Bivariate Polynomials. *SIGMA Symmetry Integrability Geom. Methods Appl.* , 4:89–106, 2008.
- [16] P. Iliev. Bispectral commuting difference operators for multivariable Askey-Wilson polynomials. *Trans. Amer. Math. Soc.*, 363:1577–1598, 2011.
- [17] P. Iliev and P. Terwilliger. The Rahman polynomials and the Lie algebra sl3(C). *Trans. Amer. Math. Soc.*, 364:4225–4238, 2012.
- [18] S. Karlin and J. McGregor. Linear growth models with many types and multidimensional Hahn polynomials. In R. Askey, editor, *Theory and Applications of Special Functions*, pages 261–288. Academic Press, 1975.
- [19] R. Koekoek, P. A. Lesky, and R. F. Swarttouw. *Hypergeometric orthogonal polynomials and their qanalogues*. Springer, 2010.
- [20] T. Koornwinder. Krawtchouk Polynomials, a Unification of Two Different Group Theoretic Interpretations. *SIAM Journal of Mathematical Analysis*, 13(6):1011–1023, November 1982.
- [21] S. Post. Racah Polynomials and Recoupling Schemes of su(1,1). *SIGMA Symmetry Integrability Geom. Methods Appl.* , 11:57–73, 2015.
- [22] H. Rosengren. Multivariable orthogonal polynomials and coupling coefficients for discrete series representations. *SIAM J. Math. Anal.*, 30:233–272, 1998.
- [23] H. Rosengren. Multivariable *q*-Hahn polynomials as coupling coefficients for quantum algebra representations. *Internat. J. Math. and Math. Sci.*, 28:331–358, 2001.
- [24] F. Scarabotti. Multidimensional Hahn polynomials, intertwining functions on the symmetric group and Clebsch-Gordon coefficients. *Methods Appl. Anal.*, 14:355–386, 2007.
- [25] F. Scarabotti. The tree method for multidimensional *q*-Hahn and *q*-Racah polynomials. *Ramanujan J.*, 25:57–91, 2011.
- [26] M. V. Tratnik. Some multivariable orthogonal polynomials of the Askey tableau-Continuous families . *J. Math. Phys.*, 32:2065, 1991.
- [27] M. V. Tratnik. Some multivariable orthogonal polynomials of the Askey tableau-discrete families . *J. Math. Phys.*, 32:1991, 2337.
- [28] N. Ja. Vilenkin and A. U. Klimyk. *Representation of Lie Groups and Special Functions, Vol. 1*, volume 72 of *Mathematics and its Applications (Soviet Series)*. Kluwer, 1991.
- [29] A. Zhedanov. *Q* rotations and other *Q* transformations as unitary nonlinear automorphisms of quantum algebras. *J. Math. Phys.*, 34:2631–2647, 1993.