

On the solvability of confluent Heun equation and associated orthogonal polynomials

Nasser Saad^{1,*}

¹*Department of Mathematics and Statistics, University of Prince Edward Island,
550 University Avenue, Charlottetown, PEI, Canada C1A 4P3.*

Abstract: The present paper analyze the constraints on the confluent Heun type-equation, $(a_{3,1}r^2 + a_{3,2}r)y'' + (a_{2,0}r^2 + a_{2,1}r + a_{2,2})y' - (\tau_{1,0}r + \tau_{1,1})y = 0$, where $|a_{3,1}|^2 + |a_{3,2}|^2 \neq 0$, and $a_{i,j}, i = 3, 2, 1, j = 0, 1, 2$ are real parameters, to admit polynomial solutions. The necessary and sufficient conditions for the existence of these polynomials are given. A three-term recurrence relation is provided to generate the polynomial solutions explicitly. We, then, prove that these polynomial solutions are a source of finite sequences of orthogonal polynomials. Several properties, such as the recurrence relation, Christoffel-Darboux formulas and the moments of the weight function, are discussed. We also show a factorization property of these orthogonal polynomials that allow for the construction of other sequences of orthogonal polynomials. For illustration, we examines the quasi- exactly solvability of the (p, q) -hyperbolic potential $V(r) = -V_0 \sinh^p(r)/\cosh^q(r), V_0 > 0, p \geq 0, q > p$. The associated orthogonal polynomials generated by the solutions of the Schrödinger equation with the $(4, 6)$ -hyperbolic potential are constructed.

PACS numbers: 33C05, 33C47, 47E05, 34B60, 33C15, 34A05, 34B30, 81Q05

Keywords: Confluent Heun equation, polynomial solutions, finite sequences of orthogonal polynomials, Christoffel-Darboux formula, hyperbolic potentials.

I. INTRODUCTION

The second-order differential equation

$$(a_{3,1}r^2 + a_{3,2}r) \frac{d^2y}{dr^2} + (a_{2,0}r^2 + a_{2,1}r + a_{2,2}) \frac{dy}{dr} - (\tau_{1,0}r + \tau_{1,1})y = 0, \quad a_{3,1}^2 + a_{3,2}^2 \neq 0, \quad a_{2,0}^2 + \tau_{1,0}^2 \neq 0, \quad (1)$$

with parameters $a_{i,j} \in \mathbb{R}, i = 3, 2, 1, j = 0, 1, 2$ has two regular singular points $r = 0$ and $r = -a_{3,2}/a_{3,1}$ with exponents $\{0, 1 - a_{2,2}/a_{3,2}\}$ and $\{0, 1 - a_{2,1}/a_{3,1} + a_{2,2}/a_{3,2} + a_{2,0}a_{3,2}/a_{3,1}^2\}$, respectively, in addition to an irregular singular point at infinity unless $a_{2,2} = \tau_{1,1} = 0$.

Although equation (1) generalized the standard confluent Heun equation [1], we shall keep the same name ‘the confluent Heun equation’ for (1) as well. The name confluent driven from the coalescence process of singularities of the general Heun equation [1–3].

Over the past few decades, the confluent Heun equation appeared intensively in mathematical physics applications. Including but not limited to, in studying the spherical Coulomb functions, the spheroidal wave functions, the Mathieu equation and confinement potentials in quantum mechanics. Most recently, it also found some interesting applications in General Relativity and Astrophysics [4–10].

The purpose of this article is twofold. First, we analyze the necessary and sufficient conditions under which the differential equation (1) admit the polynomial solutions $y_n(r) = \sum_{k=0}^n C_k r^k, n = 0, 1, 2, \dots$, Where we set up a scheme for constructing the polynomial coefficients C_k explicitly. Second, we establish a correspondence between each solution $y_n(r)$ and a finite sequence of orthogonal polynomials $\{P_k(\tau_{1,1})\}_{k=0}^n$. We discuss some properties of these orthogonal polynomials such as the recurrence relation, factorization, Christoffel-Darboux formulas, and the moments of the weight function.

As an illustration of these polynomials, we explore the quasi- exact solutions of the one-dimensional Schrödinger

*Electronic address: nsaad@upei.ca

equation with a hyperbolic potential [15]

$$-\frac{\hbar^2}{2m} \frac{d^2}{dx^2} \psi_n(x) - V_0 \frac{\sinh^p(x/d)}{\cosh^q(x/d)} \psi_n(x) = E \psi_n(x), \quad -\infty < x < \infty, \quad (2)$$

with special attention to the case of $p = 4$ and $q = 6$. We show under certain transformations, this equation reduces to a confluent Heun equation of type (1). We discuss the associated sequence of orthogonal polynomials $\{\mathcal{P}_k\}_{k=0}^n$ generated by the solution $\psi_n, n \geq 1$ and investigate some of their properties.

The article organized as follows. In Section II, we gave the necessary and sufficient conditions for the polynomial solvability constraints of equation (1) with a general procedure for formulating the polynomial solution explicitly. In Section III, we establish a correspondence between that the solution y_n and a set of orthogonal polynomials $\{\mathcal{P}_k\}_{k=0}^n$. In that section, we also derive the recurrence relation, the Christoffel-Darboux formula and prove the factorization property of these polynomials. The weight functions and some explicit expressions calculating the moments of the weight functions also discussed. In Section IV, we discuss the quasi- exact solvable quantum model, given by a hyperbolic potential and explicitly construction the finite sequences of orthogonal polynomials associated with the wavefunction solutions.

II. POLYNOMIAL SOLVABILITY CONSTRAINTS

The n -th derivatives of equation (1) yields

$$(a_{3,1}r^2 + a_{3,2}r)y^{(n+2)} + (a_{2,0}r^2 + [a_{2,1} + 2na_{3,1}]r + na_{3,2} + a_{2,2})y^{(n+1)} + ((2na_{2,0} - \tau_{1,0})r + n(n-1)a_{3,1} + na_{2,1} - \tau_{1,1})y^{(n)} + (n(n-1)a_{2,0} - n\tau_{1,0})y^{(n-1)} = 0, \quad n = 0, 1, \dots \quad (3)$$

Thus, the necessary condition for the n -degree polynomial solution is $\tau_{1,0} = na_{2,0}, n = 0, 1, \dots$ while the sufficient condition follow using the recurrence relation

$$(k+1)(ka_{3,2} + a_{2,2})\mathcal{C}_{k+1} + (k(k-1)a_{3,1} + ka_{2,1} - \tau_{1,1})\mathcal{C}_k + ((k-1)a_{2,0} - \tau_{1,0})\mathcal{C}_{k-1} = 0, \quad \mathcal{C}_{-1} = 0, \quad \mathcal{C}_0 = 1, \quad (4)$$

according to the following simple procedure.

For the zero-degree polynomial solution $y_0(r) = 1$: the necessary and sufficient conditions, respectively, reads

$$\tau_{1,0} = 0, \quad \tau_{1,1} = 0. \quad (5)$$

For the first-degree polynomial solution

$$y_1(r) = 1 + \frac{\tau_{1,1}}{a_{2,2}} r, \quad (6)$$

the necessary and sufficient conditions, respectively, reads

$$\tau_{1,0} = a_{2,0}, \quad \begin{vmatrix} -\tau_{1,1} & a_{2,2} \\ -a_{2,0} & a_{2,1} - \tau_{1,1} \end{vmatrix} = 0. \quad (7)$$

For the second-order polynomial solution

$$y_2(r) = 1 + \frac{\tau_{1,1}}{a_{2,2}} r + \frac{2a_{2,0}a_{2,2} - a_{2,1}\tau_{1,1} + \tau_{1,1}^2}{2a_{2,2}(a_{2,2} + a_{3,2})} r^2, \quad (8)$$

the necessary and sufficient conditions, respectively, reads

$$\tau_{1,0} = 2a_{2,0}, \quad \begin{vmatrix} -\tau_{1,1} & a_{2,2} & 0 \\ -2a_{2,0} & a_{2,1} - \tau_{1,1} & 2a_{3,2} + 2a_{2,2} \\ 0 & -a_{2,0} & 2a_{3,1} + 2a_{2,1} - \tau_{1,1} \end{vmatrix} = 0. \quad (9)$$

For the third-order polynomial solution

$$y_3(r) = 1 + \frac{\tau_{1,1}}{a_{2,2}} r + \frac{3a_{2,0}a_{2,2} - a_{2,1}\tau_{1,1} + \tau_{1,1}^2}{2a_{2,2}(a_{2,2} + a_{3,2})} r^2 + \frac{-6a_{2,0}a_{2,2}(a_{2,1} + a_{3,1}) + a_{2,0}(7a_{2,2} + 4a_{3,2})\tau_{1,1} + (a_{2,1} - \tau_{1,1})(2(a_{2,1} + a_{3,1}) - \tau_{1,1})\tau_{1,1}}{6a_{2,2}(a_{2,2} + a_{3,2})(a_{2,2} + 2a_{3,2})} r^3, \quad (10)$$

the necessary and sufficient conditions, respectively, reads

$$\tau_{1,0} = 3a_{2,0}, \quad \begin{vmatrix} -\tau_{1,1} & a_{2,2} & 0 & 0 \\ -3a_{2,0} & a_{2,1} - \tau_{1,1} & 2a_{3,2} + 2a_{2,2} & 0 \\ 0 & -2a_{2,0} & 2a_{3,1} + 2a_{2,1} - \tau_{1,1} & 6a_{3,2} + 3a_{2,2} \\ 0 & 0 & -a_{2,0} & 6a_{3,1} + 3a_{2,1} - \tau_{1,1} \end{vmatrix} = 0. \quad (11)$$

In general, for an n -degree polynomial solution of the confluent Heun equation, we have the following theorem.

Theorem II.1. *The necessary and sufficient conditions for an n -degree polynomial solution of equation (1), respectively, are*

$$\tau_{1,0} = n a_{2,0}, \quad n = 0, 1, \dots, \quad (12)$$

and the vanishes of the $(n+1) \times (n+1)$ determinant

$$\begin{vmatrix} -\tau_{1,1} & a_{2,2} & 0 & 0 & \dots & 0 & 0 \\ -\tau_{1,0} & a_{2,1} - \tau_{1,1} & 2a_{3,2} + 2a_{2,2} & 0 & \dots & 0 & 0 \\ 0 & a_{2,0} - \tau_{1,0} & 2a_{3,1} + 2a_{2,1} - \tau_{1,1} & 6a_{3,2} + 3a_{2,2} & \dots & 0 & 0 \\ 0 & 0 & 2a_{2,0} - \tau_{1,0} & 6a_{3,1} + 3a_{2,1} - \tau_{1,1} & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & \ddots & (n-1)a_{2,0} - \tau_{1,0} & n(n-1)a_{3,1} + na_{2,1} - \tau_{1,1} \end{vmatrix} = 0. \quad (13)$$

Theorem II.1 generalized the two conditions on the standard confluent Heun equation to allow for the n -degree polynomial solutions [9, 10].

Remark II.1: If $a_{3,2} = a_{2,2} = 0$, the differential equation (1) reduces to

$$a_{3,1}r^2 \frac{d^2y}{dr^2} + r(a_{2,0}r + a_{2,1}) \frac{dy}{dr} - (\tau_{1,0}r + \tau_{1,1})y = 0, \quad (14)$$

that have the n -degree polynomial solutions expressed in terms of the confluent hypergeometric function as

$$y_{n;k}(r) = r^k {}_1F_1 \left(k - n; 2k + \frac{a_{2,1}}{a_{3,1}}; -\frac{a_{2,0}}{a_{3,1}} r \right), \quad k = 0, 1, \dots, n, \quad (15)$$

provided that: $\tau_{1,0} = n a_{2,0}$ and $\tau_{1,1} = k(k-1) a_{3,1} + k a_{2,1}$, $k = 0, 1, 2, \dots, n$, $n = 0, 1, 2, \dots$

III. FINITE SEQUENCES OF ORTHOGONAL POLYNOMIALS

Subject to the necessary and sufficient conditions (13), the polynomial solutions (6), (8) and (10) of the differential equation (1) can be generalized as

$$y_n(r) = \sum_{k=0}^n \frac{\mathcal{P}_k^n(\tau_{1,1})}{k! a_{3,2}^k \left(\frac{a_{2,2}}{a_{3,2}} \right)_k} r^k, \quad n = 0, 1, 2, \dots, \quad (16)$$

where $(\alpha)_n$ refer to the pochhammer symbol $(\alpha)_n = \alpha(\alpha+1)\dots(\alpha-n+1)$ satisfies $(-n)_k = 0$ for any positive integers $k \geq n$.

The coefficients $\mathcal{P}_k^n(\zeta)$, $k = 0, \dots, n$ are a sequence of k -degree polynomials in the parameter $\zeta = \tau_{1,1}$. This sequence of polynomials will be our focus in the present work. Indeed, for each n , the solution $y_n(r)$ of the differential equation (1) generate a finite sequence of polynomials $\{\mathcal{P}_k^n(\tau_{1,1})\}_{k=0}^n$ satisfying a number of very interesting properties.

A. Three-term recurrence relation

The first property is that they satisfy a three-term recurrence relation

$$\mathcal{P}_{k+1}^n(\zeta) + (k(k-1)a_{3,1} + ka_{2,1} - \zeta) \mathcal{P}_k^n(\zeta) + k((k-1)a_{3,2} + a_{2,2})((k-1)a_{2,0} - \tau_{1,0}) \mathcal{P}_{k-1}^n(\zeta) = 0, \quad 0 \leq k \leq n-1, \quad (17)$$

initiated with $\mathcal{P}_{-1}^n(\zeta) = 0$, $\mathcal{P}_0^n(\zeta) = 1$. This recurrence relation can be obtained by the fact that $y_n(r)$ satisfies the differential equation (1). Using the necessary condition $\tau_{1,0} = n a_{2,0}$, equation (17) reduce to

$$\mathcal{P}_{k+1}^n(\zeta) = (\zeta - k(k-1)a_{3,1} - k a_{2,1}) \mathcal{P}_k^n(\zeta) - k(k-n-1)a_{2,0}((k-1)a_{3,2} + a_{2,2}) \mathcal{P}_{k-1}^n(\zeta), \quad \mathcal{P}_{-1}^n(\zeta) = 0, \quad \mathcal{P}_0^n(\zeta) = 1, \quad (18)$$

for $k = 0, 1, 2, \dots, n-1$. Here, n is a (fixed) nonnegative integer refer to the degree of the polynomial solution (16).

From the general theory of orthogonal polynomials and according to Favard's Theorem [11–13]), $\{\mathcal{P}_k^n(\zeta)\}_{k=0}^n$ form a sequence of orthogonal polynomials. The first few expressions of these polynomials reads

$$\begin{aligned} \mathcal{P}_1^n(\zeta) &= \zeta, \\ \mathcal{P}_2^n(\zeta) &= \zeta^2 - a_{2,1}\zeta + n a_{2,0} a_{2,2}, \\ \mathcal{P}_3^n(\zeta) &= \zeta^3 - (3a_{2,1} + 2a_{3,1})\zeta^2 + (2a_{2,1}^2 + 2a_{2,1}a_{3,1} + a_{2,0}a_{2,2}n - 2(a_{2,2} + a_{3,2})(a_{2,0} - a_{2,0}n))\zeta - 2na_{2,0}a_{2,2}(a_{2,1} + a_{3,1}) \\ \mathcal{P}_4^n(\zeta) &= \zeta^4 - 2(3a_{2,1} + 4a_{3,1})\zeta^3 + (11a_{2,1}^2 + 26a_{2,1}a_{3,1} + 12a_{3,1}^2 + 2a_{2,0}(a_{2,2}(3n-4) + a_{3,2}(4n-7)))\zeta^2 \\ &\quad - 2(3(a_{2,1} + a_{3,1})(a_{2,1}^2 - 2a_{2,0}a_{2,2} + 2a_{2,1}a_{3,1}) - a_{2,0}a_{3,2}(9a_{2,1} + 6a_{3,1}) + na_{2,0}(a_{2,2}(7a_{2,1} + 10a_{3,1}) + 6(a_{2,1} + a_{3,1})a_{3,2}))\zeta \\ &\quad + 3na_{2,0}a_{2,2}(2(a_{2,1} + a_{3,1})(a_{2,1} + 2a_{3,1}) + a_{2,0}(n-2)(a_{2,2} + 2a_{3,2})). \end{aligned}$$

An important consequence of the three-term recurrence formula (18) is the following Christoffel-Darboux formula.

Theorem III.1. For $\zeta \neq \zeta'$,

$$\sum_{j=0}^k \frac{\mathcal{P}_j^n(\zeta) \mathcal{P}_j^n(\zeta')}{j! (a_{2,0}a_{3,2})^j \left(\frac{a_{2,2}}{a_{3,2}}\right)_j (-n)_j} = \frac{\mathcal{P}_{k+1}^n(\zeta) \mathcal{P}_k^n(\zeta') - \mathcal{P}_k^n(\zeta) \mathcal{P}_{k+1}^n(\zeta')}{k! (a_{2,0}a_{3,2})^k \left(\frac{a_{2,2}}{a_{3,2}}\right)_k (-n)_k (\zeta - \zeta')}, \quad k = 1, \dots, n-1, \quad (19)$$

while for $\zeta' \rightarrow \zeta$

$$\sum_{j=0}^k \frac{(\mathcal{P}_j^n(\zeta))^2}{j! (a_{2,0}a_{3,2})^j \left(\frac{a_{2,2}}{a_{3,2}}\right)_j (-n)_j} = \frac{[\mathcal{P}_{k+1}^n(\zeta)]' \mathcal{P}_k^n(\zeta) - [\mathcal{P}_k^n(\zeta)]' \mathcal{P}_{k+1}^n(\zeta)}{k! (a_{2,0}a_{3,2})^k \left(\frac{a_{2,2}}{a_{3,2}}\right)_k (-n)_k}, \quad k = 1, \dots, n-1. \quad (20)$$

Proof. The recurrence relations (18) for ζ and ζ' reads, respectively,

$$\begin{aligned} \mathcal{P}_{k+1}^n(\zeta) &= \zeta \mathcal{P}_k^n(\zeta) - [k(k-1)a_{3,1} + k a_{2,1}] \mathcal{P}_k^n(\zeta) - k(k-n-1)a_{2,0}((k-1)a_{3,2} + a_{2,2}) \mathcal{P}_{k-1}^n(\zeta), \\ \mathcal{P}_{k+1}^n(\zeta') &= \zeta' \mathcal{P}_k^n(\zeta') - [k(k-1)a_{3,1} + k a_{2,1}] \mathcal{P}_k^n(\zeta') - k(k-n-1)a_{2,0}((k-1)a_{3,2} + a_{2,2}) \mathcal{P}_{k-1}^n(\zeta'). \end{aligned}$$

Multiplying the first by $\mathcal{P}_k^n(\zeta')$ and the second by $\mathcal{P}_k^n(\zeta)$ then subtract, the resulting equation reads

$$(\zeta - \zeta') \mathcal{P}_k^n(\zeta') \mathcal{P}_k^n(\zeta) = Q_{k+1}(\zeta, \zeta') - \lambda_{k+1} Q_k(\zeta, \zeta')$$

where $\lambda_{k+1} = k(k-n-1)a_{2,0}((k-1)a_{3,2} + a_{2,2})$, and $Q_{k+1}(\zeta, \zeta') = \mathcal{P}_{k+1}^n(\zeta) \mathcal{P}_k^n(\zeta') - \mathcal{P}_k^n(\zeta) \mathcal{P}_{k+1}^n(\zeta')$. Thus, recursively over k we have

$$\begin{aligned} (\zeta - \zeta') \mathcal{P}_k^n(\zeta) \mathcal{P}_k^n(\zeta') &= Q_{k+1}(\zeta, \zeta') - \lambda_{k+1} Q_k(\zeta, \zeta') \\ (\zeta - \zeta') \mathcal{P}_{k-1}^n(\zeta) \mathcal{P}_{k-1}^n(\zeta') &= Q_k(\zeta, \zeta') - \lambda_k Q_{k-1}(\zeta, \zeta') \\ &\dots = \dots \\ (\zeta - \zeta') \mathcal{P}_0^n(\zeta) \mathcal{P}_0^n(\zeta') &= Q_1(\zeta, \zeta'). \end{aligned}$$

From which, it is straightforward to obtain

$$\begin{aligned} (\zeta - \zeta') \left[\mathcal{P}_k^n(\zeta) \mathcal{P}_k^n(\zeta') + \lambda_{k+1} \mathcal{P}_{k-1}^n(\zeta) \mathcal{P}_{k-1}^n(\zeta') + \lambda_{k+1} \lambda_k \mathcal{P}_{k-2}^n(\zeta) \mathcal{P}_{k-2}^n(\zeta') + \lambda_{k+1} \lambda_k \lambda_{k-1} \mathcal{P}_{k-3}^n(\zeta) \mathcal{P}_{k-3}^n(\zeta') + \dots \right. \\ \left. + \lambda_{k+1} \lambda_k \lambda_{k-1} \lambda_{k-2} \dots \lambda_2 \mathcal{P}_0^n(\zeta) \mathcal{P}_0^n(\zeta') \right] = Q_{k+1}(\zeta, \zeta') \end{aligned}$$

Divide both sides by $(\zeta - \zeta') \lambda_{k+1} \lambda_k \lambda_{k-1} \lambda_{k-2} \dots \lambda_2$ and sum over k we finally gets

$$\sum_{j=0}^k \frac{\mathcal{P}_j^n(\zeta) \mathcal{P}_j^n(\zeta')}{\lambda_{j+1} \lambda_j \lambda_{j-1} \dots \lambda_2} = (\lambda_{k+1} \lambda_k \lambda_{k-1} \dots \lambda_2)^{-1} \frac{\mathcal{P}_{k+1}^n(\zeta) \mathcal{P}_k^n(\zeta') - \mathcal{P}_k^n(\zeta) \mathcal{P}_{k+1}^n(\zeta')}{\zeta - \zeta'}.$$

Here, the critical polynomial $P_2^1(\zeta)$ factorize every other polynomial $P_{k+2}^n(\zeta)$, $k = 1, 2, \dots$. As another illustration, we note for $n = 2$:

$$\begin{aligned} \mathcal{P}_1^2(\zeta) &= \zeta, \\ \mathcal{P}_2^2(\zeta) &= \zeta^2 - a_{2,1}\zeta + 2a_{2,0}a_{2,2}, \\ \mathcal{P}_3^2(\zeta) &= -4a_{2,0}a_{2,1}a_{2,2} - 4a_{2,0}a_{2,2}a_{3,1} + (2a_{2,1}^2 + 4a_{2,0}a_{2,2} + 2a_{2,1}a_{3,1} + 2a_{2,0}a_{3,2})\zeta - (3a_{2,1} + 2a_{3,1})\zeta^2 + \zeta^3, \\ \mathcal{P}_4^2(\zeta) &= (\zeta - 3a_{2,1} - 6a_{3,1})\mathcal{P}_3^2(\zeta), \\ \mathcal{P}_5^2(\zeta) &= (12a_{2,1}^2 - 4a_{2,0}a_{2,2} + 60a_{2,1}a_{3,1} + 72a_{3,1}^2 - 12a_{2,0}a_{3,2} - (7a_{2,1} + 18a_{3,1})\zeta + \zeta^2)\mathcal{P}_3^2(\zeta). \end{aligned}$$

with $P_3^n(\zeta)$ being a critical polynomial factorize every other polynomial $P_{k+3}^n(\zeta)$, $k = 1, 2, \dots$. Indeed, all the polynomials $\mathcal{P}_{k+n+1}^n(\zeta)$, beyond some critical polynomial $\mathcal{P}_{n+1}^n(\zeta)$, factored into the product

$$\mathcal{P}_{k+n+1}^n(\zeta) = \mathcal{Q}_k^n(\zeta)\mathcal{P}_{n+1}^n(\zeta), \quad k = 0, 1, \dots \quad (25)$$

Interestingly, the quotient polynomials $\{\mathcal{Q}_k^n(\zeta)\}_{k \geq 0}$ form an infinite sequence of orthogonal polynomials. To prove this claim, we substitute (25) into (18) where we re-index the polynomials to eliminate the common factor $\mathcal{P}_{n+1}^n(\zeta)$ from both sides. The recurrence relation (18) then reduces to a three-term recurrence relation for the polynomials $\{\mathcal{Q}_k^n(\zeta)\}_{k \geq 0}$ that reads

$$\mathcal{Q}_k^n(\zeta) = (\zeta - (k+n)((k+n-1)a_{3,1} + a_{2,1}))\mathcal{Q}_{k-1}^n(\zeta) - (k-1)(k+n)a_{2,0}((k+n-1)a_{3,2} + a_{2,2})\mathcal{Q}_{k-2}^n(\zeta), \quad k \geq 1, \quad (26)$$

initiated with $\mathcal{Q}_{-1}^n(\zeta) = 0$, and $\mathcal{Q}_0^n(\zeta) = 1$. Hence, the quotient polynomials $\mathcal{Q}_k^n(\zeta)$ also form a new sequence of orthogonal polynomials for each value of n . The first few polynomials of $\mathcal{Q}_k^n(\zeta)$ can to explicitly evaluate to reads

$$\begin{aligned} \mathcal{Q}_1^n(\zeta) &= \zeta - (n+1)(a_{2,1} + na_{3,1}), \\ \mathcal{Q}_2^n(\zeta) &= \zeta^2 - (2a_{3,1}(1+n)^2 + a_{2,1}(2n+3))\zeta + (n+2)[(1+n)(a_{2,1} + a_{3,1}n)(a_{2,1} + a_{3,1} + a_{3,1}n) - a_{2,0}(a_{2,2} + a_{3,2} + a_{3,2}n)]. \end{aligned}$$

The Christoffel-Darboux formula for this sequence of orthogonal polynomials reads

$$\sum_{j=0}^k \frac{\mathcal{Q}_j^n(\zeta)\mathcal{Q}_j^n(\zeta')}{j!(n+2)_j(a_{2,0}a_{3,2})^j \left(n+1+\frac{a_{2,2}}{a_{3,2}}\right)_j} = \frac{\mathcal{Q}_{k+1}^n(\zeta)\mathcal{Q}_k^n(\zeta') - \mathcal{Q}_k^n(\zeta)\mathcal{Q}_{k+1}^n(\zeta')}{k!(n+2)_k(a_{2,0}a_{3,2})^k \left(n+1+\frac{a_{2,2}}{a_{3,2}}\right)_k (\zeta - \zeta')}, \quad (27)$$

and as $\zeta' \rightarrow \zeta$,

$$\sum_{j=0}^k \frac{(\mathcal{Q}_j^n(\zeta))^2}{j!(n+2)_j(a_{2,0}a_{3,2})^j \left(n+1+\frac{a_{2,2}}{a_{3,2}}\right)_j} = \frac{[\mathcal{Q}_{k+1}^n(\zeta)]'\mathcal{Q}_k^n(\zeta) - [\mathcal{Q}_k^n(\zeta)]'\mathcal{Q}_{k+1}^n(\zeta)}{k!(n+2)_k(a_{2,0}a_{3,2})^k \left(n+1+\frac{a_{2,2}}{a_{3,2}}\right)_k}. \quad (28)$$

C. Moments and weight functions

From the general theory of orthogonal polynomials, there exists a certain weight function $\mathcal{W}(\xi)$ normalized as

$$\int_{\mathcal{S}} \mathcal{W}(\xi) d\xi = 1, \quad (29)$$

for which

$$\int_{\mathcal{S}} \mathcal{P}_k^n(\zeta) \mathcal{P}_{k'}^n(\zeta) \mathcal{W}(\zeta) d\zeta = p_k p_{k'} \delta_{kk'}, \quad k, k' = 0, 1, \dots, n, \quad (30)$$

where \mathcal{S} is the support of the measure $\mathcal{W}(\zeta) d\zeta$ on the real line and $\delta_{kk'}$ is Kronecker's symbol.

Theorem III.3. *For the monic orthogonal polynomials $\{\mathcal{P}_k^n(\zeta)\}_{k=0}^n$*

$$\int_{\mathcal{S}} \zeta^j \mathcal{P}_k^n(\zeta) \mathcal{W}(\zeta) d\zeta = p_k^2 \delta_{kj}, \quad \forall j \leq k. \quad (31)$$

Proof. By writing $p_{k'}^n(\zeta) = \sum_{j=0}^{k'} \alpha_{k';j} \zeta^j$, where $\alpha_{k';k'} = 1$, equation (22) yields

$$\sum_{j=0}^{k'} \alpha_{k';j} \int_{\mathcal{S}} \zeta^j \mathcal{P}_k^n(\zeta) \mathcal{W}(\zeta) d\zeta = p_k p_{k'} \delta_{kk'}, \quad k, k' = 0, 1, \dots, n, \quad \alpha_{k';k'} = 1. \quad (32)$$

For $k' = 0, 1, \dots, n$, equation (32) generate a linear system that reduce to

$$\begin{pmatrix} \int_{\mathcal{S}} \mathcal{P}_k^n(\zeta) \mathcal{W}(\zeta) d\zeta \\ \int_{\mathcal{S}} \zeta \mathcal{P}_k^n(\zeta) \mathcal{W}(\zeta) d\zeta \\ \int_{\mathcal{S}} \zeta^2 \mathcal{P}_k^n(\zeta) \mathcal{W}(\zeta) d\zeta \\ \int_{\mathcal{S}} \zeta^3 \mathcal{P}_k^n(\zeta) \mathcal{W}(\zeta) d\zeta \\ \vdots \\ \int_{\mathcal{S}} \zeta^{k'} \mathcal{P}_k^n(\zeta) \mathcal{W}(\zeta) d\zeta \end{pmatrix} = p^k \begin{pmatrix} 1 & 0 & 0 & 0 & \dots & 0 \\ \alpha_{1;0}^n & 1 & 0 & 0 & \dots & 0 \\ \alpha_{2;0}^n & \alpha_{2;1}^n & 1 & 0 & \dots & 0 \\ \alpha_{3;0}^n & \alpha_{3;1}^n & \alpha_{3;2}^n & 1 & \dots & 0 \\ \alpha_{4;0}^n & \alpha_{4;1}^n & \alpha_{4;2}^n & \alpha_{4;3}^n & \dots & 0 \\ \vdots & \vdots & \vdots & \vdots & \dots & \vdots \\ \alpha_{k';0}^n & \alpha_{k';1}^n & \alpha_{k';2}^n & \alpha_{k';3}^n & \dots & 1 \end{pmatrix}_{(k'+1) \times (k'+1)}^{-1} \times \begin{pmatrix} p_0 \delta_{k0} \\ p_1 \delta_{k1} \\ p_2 \delta_{k2} \\ p_3 \delta_{k3} \\ \vdots \\ p_{k'} \delta_{kk'} \end{pmatrix}_{(k'+1) \times 1}. \quad (33)$$

The inverse of the lower triangular matrix is again lower triangular matrix with ones on the main diagonal. This argument concludes our assertion (27). \square

From Theorem III.3, we also conclude that

$$\int_{\mathcal{S}} \mathcal{W}(\zeta) d\zeta = 1, \quad \int_{\mathcal{S}} \mathcal{P}_k^n(\zeta) \mathcal{W}(\zeta) d\zeta = 0 \quad (k \geq 1), \quad \int_{\mathcal{S}} \mathcal{P}_j^n(\zeta) \mathcal{P}_k^n(\zeta) \mathcal{W}(\zeta) d\zeta = 0, \quad (0 \leq j < k \leq n). \quad (34)$$

Theorem III.4. *The norms of all polynomials $\mathcal{P}_k^n(\xi)$ with $k \geq n + 1$ vanish.*

Proof. The norms p_k can be evaluated using the recurrence relations (18) after multiplying throughout by $\zeta^{k-1} \mathcal{W}(\zeta)$ and taking integral over ζ , we obtain, because of Theorem III.3, a simple two-term recursion relation for the squared norm

$$\mathcal{G}_k = k(k-1-n)a_{2,0}((k-1)a_{3,2} + a_{2,2})\mathcal{G}_{k-1},$$

with a solution given by

$$\mathcal{G}_k = |q_k|^2 = k! (a_{2,0}a_{3,2})^k (-n)_k \left(\frac{a_{2,2}}{a_{3,2}} \right)_k, \quad (35)$$

which is equal to zero for $k = n + 1, n + 2, \dots$ \square

As a result of Theorem III.4, we note

$$\int_{\mathcal{S}} [\mathcal{P}_k^n(\zeta)]^2 \mathcal{W}(\zeta) d\zeta = k! (a_{2,0}a_{3,2})^k (-n)_k \left(\frac{a_{2,2}}{a_{3,2}} \right)_k, \quad 0 \leq k \leq n, \quad (36)$$

$$\int_{\mathcal{S}} \zeta \mathcal{P}_k^n(\zeta) \mathcal{P}_{k-1}^n(\zeta) \mathcal{W}(\zeta) d\zeta = k! (a_{2,0}a_{3,2})^k (-n)_k \left(\frac{a_{2,2}}{a_{3,2}} \right)_k, \quad k = 1, 2, \dots, \quad (37)$$

$$\int_{\mathcal{S}} \zeta [\mathcal{P}_k^n(\zeta)]^2 \mathcal{W}(\zeta) d\zeta = k((k-1)a_{3,1} + a_{2,1})k! (a_{2,0}a_{3,2})^k (-n)_k \left(\frac{a_{2,2}}{a_{3,2}} \right)_k, \quad k = 0, 1, 2, \dots \quad (38)$$

Although we don't have a general formula for evaluating the moments

$$\mu_k = \int_{\mathcal{S}} \zeta^k \mathcal{W}(\zeta) d\zeta, \quad k = 0, 1, \dots, n, \quad (39)$$

it is possible to compute μ_k recursively using Theorem III.3 and the recurrence relation (16). It is not difficult to show that

$$\int_{\mathcal{S}} W(\zeta) d\zeta = 1, \quad \int_{\mathcal{S}} \zeta W(\zeta) d\zeta = 0, \quad \int_{\mathcal{S}} \zeta^2 W(\zeta) d\zeta = p_1^2 = -n a_{2,0} a_{2,2}, \quad a_{2,0} a_{2,2} < 0, \quad (40)$$

and for all $k \geq 2$,

$$\int_{\mathcal{S}} \zeta \mathcal{P}_k^n(\zeta) W(\zeta) d\zeta = 0. \quad (41)$$

For example,

$$\int_{\mathcal{S}} \zeta^3 W(\zeta) d\zeta = a_{2,1} p_1^2, \quad \int_{\mathcal{S}} \zeta^4 W(\zeta) d\zeta = p_2^2 + (a_{2,1} - n a_{2,0} a_{2,2}) p_1^2, \quad \text{and so forth.} \quad (42)$$

Theorem III.5. *The norms of all polynomials $Q_k^n(\xi)$ reads*

$$\mathcal{G}_k^{\mathcal{Q}} = k! (n+2)_k (a_{2,0} a_{3,2})^k \left(n + 1 + \frac{a_{2,2}}{a_{3,2}} \right)_k. \quad (43)$$

Proof. The proof follows by multiplying the recurrence relation (26) by $\zeta^{k-2} W(\zeta)$ and integrate over ζ . This procedure implies the two-term recurrence relation

$$\mathcal{G}_k^{\mathcal{Q}} = k(k+n+1) a_{2,0} ((k+n) a_{3,2} + a_{2,2}) \mathcal{G}_{k-1}^{\mathcal{Q}}, \quad k \geq 1 \quad (44)$$

where $\mathcal{G}_k^{\mathcal{Q}} = \int_{\mathcal{S}} |\mathcal{Q}_k^n(\zeta)|^2 W(\zeta) d\zeta = \int_{\mathcal{S}} \zeta^k Q_k^n(\zeta) d\zeta$. with a solution given (43). \square

IV. EXAMPLE: HYPERBOLIC POTENTIALS

Recently, Downing [15] studied a class of potentials characterized by two physical parameters V_0 and d shaping the potential depth and width respectively

$$V(x) = -V_0 \frac{\sinh^4(x/d)}{\cosh^6(x/d)}, \quad -\infty < x < \infty. \quad (45)$$

In this section, we consider first a more general class of potentials given by [16]

$$V(x) = -V_0 \frac{\sinh^p(x/d)}{\cosh^q(x/d)}, \quad -\infty < x < \infty, \quad V_0 > 0, \quad p \geq 0, \quad q > p. \quad (46)$$

The condition $p \geq 0$ required to avoid a singularity at $x = 0$ and the condition $p < q$ is to ensure the existence of a bound-below potential support the presence of at least one bound state. Indeed, it is well known [19]: *Given a non-positive potential $V \leq 0$ with*

$$\int_{-\infty}^{\infty} V(x) dx < 0, \quad (47)$$

then there exists a bound state (independent of the potential depth and range) with energy $E < 0$ for the Hamiltonian $H = -\frac{\hbar^2}{2m} \frac{d^2}{dx^2} + V(x)$. For the potential (46), it is not difficult to prove that

$$\int_{-\infty}^{\infty} V(x) dx = -\frac{(1 + (-1)^p) V_0 \Gamma\left(\frac{1+p}{2}\right) \Gamma\left(\frac{1}{2}(q-p)\right)}{2\Gamma\left(\frac{1+q}{2}\right)} < 0, \quad p \geq 0, \quad p < q.$$

The stationary one-dimensional Schrödinger equation for a nonrelativistic particle of mass m and energy E in a hyperbolic potential $V(x)$ written as follows

$$-\frac{\hbar^2}{2m} \frac{d^2}{dx^2} \psi(x) - V_0 \frac{\sinh^p(x/d)}{\cosh^q(x/d)} \psi(x) = E \psi(x), \quad V_0 > 0, \quad -\infty < x < \infty. \quad (48)$$

can be written using the substitution $z = x/d$ as

$$-\frac{d^2}{dx^2}\psi(z) - d^2U_0 \frac{\sinh^p(z)}{\cosh^q(z)}\psi(z) = \varepsilon d^2\psi(z), \quad -\infty < z < \infty. \quad (49)$$

where $U_0 = 2mV_0/\hbar^2$ and $\varepsilon = 2mE/\hbar^2$. Upon making the change of variable $\eta = 1/\cosh^2(z)$ such that the domain $-\infty < z < \infty$ maps to $0 < \eta < 1$, equation (49) yields

$$4\eta^2(1-\eta)\frac{d^2\psi(\eta)}{d\eta^2} + [4\eta - 6\eta^2]\frac{d\psi(\eta)}{d\eta} + (\varepsilon d^2 + d^2U_0\eta^{(q-p)/2}(1-\eta)^{p/2})\psi(\eta) = 0, \quad 0 < \eta < 1, \quad p \geq 0, \quad q > p. \quad (50)$$

It is straightforward to verify that $\eta = 0$ is a regular singular point with exponents $\{\pm d\sqrt{-\varepsilon}/2\}$ where $\varepsilon < 0$ and the singular point $\eta = 1$ is also a regular singular point with exponents $s = \{0, 1/2\}$ utilizing the symmetric and anti-symmetric solutions [15], respectively. From the classical theory of ordinary differential equation, we know that if $\eta = \infty$ is a regular singular point as well, then the differential equation (50) is a hypergeometric-type equation with exact solutions expressed in terms of the Gauss hypergeometric function ${}_2F_1$. This claim can be easily verified in the case of $p = 0$ and $q = 2$ where the differential equation (50) turn to be Fuchsian with the three regular singular points at $\eta = 0, 1$, and ∞ .

The point $\eta = \infty$ is an irregular singular point only if $p > 2$ and $q > p$. A reason for $V(x) = -V_0 \sinh^4(\eta)/\cosh^6(\eta)$ to be a quasi-exactly solvable potential that will be our focus because of the recent interest. In this case, equation (50) reads

$$\eta^2(1-\eta)\psi''(\eta) + \eta\left[1 - \frac{3}{2}\eta\right]\psi'(\eta) + \frac{1}{4}(\varepsilon d^2 + U_0 d^2\eta(1-\eta)^2)\psi(\eta) = 0. \quad (51)$$

Because of the asymptotic behavior at the singular points, the exact solutions takes the form

$$\psi(\eta) = \eta^{\beta/2}(1-\eta)^s e^{-\alpha\eta/2} f(\eta), \quad \beta, \alpha > 0, \quad s = 0, 1/2, \quad 0 < \eta < 1, \quad (52)$$

we obtain, for the function $f(\eta)$, the following differential equation

$$f''(\eta) + \left[\frac{1+4s}{2(\eta-1)} + \frac{1+\beta}{\eta} - \alpha\right]f'(\eta) + \left[\frac{-\alpha + \beta + \beta^2 + 4s - 4\alpha s + 4\beta s}{4(\eta-1)} + \frac{-2\alpha - \beta - 2\alpha\beta - \beta^2 - 4s - 4\beta s + \alpha^2}{4\eta}\right]f(\eta) = 0 \quad (53)$$

where $s = 0, 1/2$, $\beta = d\sqrt{-\varepsilon}$ and $\alpha = d\sqrt{U_0}$. Equation (53) can be written as

$$4\eta(\eta-1)f''(\eta) + [-4\alpha\eta^2 + (6+4\alpha+4\beta+8s)\eta - 4(1+\beta)]f'(\eta) + [(\alpha(\alpha-2\beta-4s-3)\eta - \alpha^2 + 2\alpha(1+\beta) + (1+\beta)(\beta+4s))]f(\eta) = 0 \quad (54)$$

Using Theorem II.1, the necessary condition for the differential equation (54) to admit the polynomial solutions is

$$\alpha - 2\beta = 4n + 4s + 3$$

that implies the following eigenvalue spectra:

$$\varepsilon_n = -\frac{(4n+4s+3-d\sqrt{U_0})^2}{4d^2}, \quad d^2U_0 > (4n+4s+3)^2. \quad (55)$$

Again using Theorem II.1, the sufficient condition follows by the determinant

$$\begin{vmatrix} \mu_0 & \delta_1 & 0 & 0 & 0 & \dots & 0 & 0 & 0 \\ \gamma_1 & \mu_1 & \delta_2 & 0 & 0 & \dots & 0 & 0 & 0 \\ 0 & \gamma_2 & \mu_2 & \delta_3 & 0 & \dots & 0 & 0 & 0 \\ 0 & 0 & \gamma_3 & \mu_3 & \delta_4 & \dots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \ddots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & 0 & \dots & \gamma_{n-1} & \mu_{n-1} & \delta_n \\ 0 & 0 & 0 & 0 & 0 & \dots & 0 & \gamma_n & \mu_n \end{vmatrix} = 0, \quad (56)$$

where

$$\mu_j = -\alpha^2 + 2\alpha(2j + \beta + 1) + (2j + \beta + 1)(2j + \beta + 4s), \quad \gamma_j = 4\alpha(1 - j + n), \quad \delta_j = -4j(\beta + j).$$

The exact solutions are given explicitly by the equation

$$\psi_n(\eta) = \eta^{d\sqrt{-\varepsilon_n}} (1 - \eta)^s e^{-d\sqrt{U_0}\eta/2} \sum_{k=0}^n \frac{\mathcal{P}_k^n(\zeta)}{k! (1 + \beta)_k} \left(-\frac{\eta}{4}\right)^k, \quad n = 0, 1, 2, \dots, \quad (57)$$

where the coefficients $\mathcal{P}_k^n(\zeta)$, with $\zeta \equiv \tau_{1,1} = (1 + d\sqrt{-\varepsilon_n})(d\sqrt{-\varepsilon_n} + 4s) + 2d(1 + d\sqrt{-\varepsilon_n})\sqrt{U_0} - d^2U_0$ are evaluated using the following three-term recurrence relation

$$\mathcal{P}_{k+1}^n(\zeta) = (\zeta - 2k(1 + 2\alpha + 2\beta + 2k + 4s))\mathcal{P}_k^n(\zeta) - 16k(k - n - 1)\alpha(\beta + k)\mathcal{P}_{k-1}^n(\zeta), \quad \mathcal{P}_{-1}^n(\zeta) = 0, \quad \mathcal{P}_0^n(\zeta) = 1, \quad (58)$$

with the square norm given by

$$\int_S [\mathcal{P}_k^n(\zeta)]^2 \mathcal{W}(\zeta) d\zeta = 2^{4k} k! \alpha^k (-n)_k (1 + \beta)_k, \quad 0 \leq k \leq n. \quad (59)$$

The first few solutions are:

- For $n = 0$, we have

$$\varepsilon_0 = -\frac{(4s + 3 - d\sqrt{U_0})^2}{4d^2}, \quad f_0(\eta) = 1, \quad (60)$$

subject to the root finding of the equation

$$7d^2U_0 - 12d\sqrt{U_0} - 3(4s + 1)(4s + 5) = 0. \quad \text{where} \quad U_0 > (4s + 3)^2/d^2. \quad (61)$$

The critical polynomial reads

$$\mathcal{P}_1^0(\alpha, \beta; s) = -\alpha^2 + 2\alpha(1 + \beta) + (1 + \beta)(\beta + 4s).$$

- For $n = 1$, we have

$$\varepsilon_1 = -\frac{(4s + 7 - d\sqrt{U_0})^2}{4d^2}, \quad f_1(\eta) = 1 - \left(\frac{\alpha^2 - 2\alpha(1 + \beta) - (1 + \beta)(\beta + 4s)}{4(1 + \beta)} \right) \eta. \quad (62)$$

subject to the root finding of the equation

$$\alpha^4 - 4(2 + \beta)\alpha^3 + 2(3 + 5\beta + \beta^2 - 4(2 + \beta)s)\alpha^2 + 4(1 + \beta)(3 + 4\beta + \beta^2 + 4n + 4(3 + \beta)s)\alpha + (1 + \beta)(3 + \beta)(\beta + 4s)(2 + \beta + 4s) = 0 \quad (63)$$

where $\alpha = d\sqrt{U_0}$, $\beta = (7 + 4s - d\sqrt{U_0})/2$ and $d^2U_0 > (7 + 4s)^2$. The critical polynomial in this case reads

$$\mathcal{P}_2^1(\alpha, \beta; s) = 16\alpha(1 + \beta) + (-\alpha^2 + 2\alpha(1 + \beta) + (1 + \beta)(\beta + 4s))(-\alpha^2 + 2\alpha(1 + \beta) + (1 + \beta)(\beta + 4s) - 2(3 + 2\alpha + 2\beta + 4s)).$$

- For $n = 2$, we have

$$\begin{aligned} \varepsilon_2 = & -\frac{(4s + 11 - d\sqrt{U_0})^2}{4d^2}, \\ f_2(\eta) = & 1 - \frac{\alpha^2 - 2\alpha(1 + \beta) - (1 + \beta)(\beta + 4s)}{4(1 + \beta)} \eta \\ & + \frac{32\alpha(1 + \beta) - (6 + \alpha^2 + 2\alpha(1 - \beta) + 4s + \beta(3 - \beta - 4s))(-\alpha^2 + 2\alpha(1 + \beta) + (1 + \beta)(\beta + 4s))}{32(1 + \beta)(2 + \beta)} \eta^2, \end{aligned} \quad (64)$$

subject to the the root finding of the equation

$$\begin{aligned} & 32\alpha(2 + \beta)(-\alpha^2 + 2\alpha(1 + \beta) + (1 + \beta)(\beta + 4s)) \\ & + (-\alpha^2 + 2\alpha(5 + \beta) + (5 + \beta)(4 + \beta + 4s))(32\alpha(1 + \beta) + (-\alpha^2 + 2\alpha(1 + \beta) + (1 + \beta)(\beta + 4s))) \\ & \times (-\alpha^2 + 2\alpha(3 + \beta) + (3 + \beta)(2 + \beta + 4s)) = 0 \end{aligned} \quad (65)$$

where $\alpha = d\sqrt{U_0}$, $\beta = (11 + 4s - d\sqrt{U_0})/2$ and $d^2U_0 > (11 + 4s)^2$. The critical polynomial reads in this case as

$$\begin{aligned} \mathcal{P}_3^2(\alpha, \beta; s) = & 32\alpha(1 + \beta)(-\alpha^2 + 2\alpha(1 + \beta) + (1 + \beta)(\beta + 4s) - 4(5 + 2\alpha + 2\beta + 4s)) \\ & + (2\alpha(1 + \beta) - \alpha^2 + (1 + \beta)(\beta + 4s))(32\alpha(2 + \beta) + (-\alpha^2 + 2\alpha(1 + \beta) + (1 + \beta)(\beta + 4s) - 2(3 + 2\alpha + 2\beta + 4s))) \\ & \times (-\alpha^2 + 2\alpha(1 + \beta) + (1 + \beta)(\beta + 4s) - 4(5 + 2\alpha + 2\beta + 4s)) \end{aligned}$$

V. CONCLUSIONS

We have given the sufficient and necessary conditions that ensure the existence of the polynomial solutions of a general confluent Heun equation. It is indeed exciting to know that these polynomial solutions are the source of finite sequences of orthogonal polynomials. The number of the orthogonal polynomials in each sequence equal to the degree of the polynomial solution of the confluent Heun equation. Although we discussed several properties of these orthogonal polynomials, a general theory that explain the zeros and evaluation of the discrete weight function will be of great interest. We hope the present study will stimulate further research on the theory of finite orthogonal polynomials.

VI. ACKNOWLEDGMENTS

Partial financial support of this work under Grant No. GP249507 from the Natural Sciences and Engineering Research Council of Canada is gratefully acknowledged.

References

-
- [1] A. Ronveaux (Ed.) *Heun's Differential Equations*, The Clarendon Press Oxford University Press, New York (1995).
 - [2] A. Decarreau, M. C. Dumont-Lepage, P. Maroni, A. Robert, and A. Ronveaux, *Formes canoniques des équations confluentes de l'équation de Heun*, Ann. Soc. Sci. Bruxelles. **T92(I-II)** (1978) 53 - 78.
 - [3] A. Decarreau, P. Maroni, and A. Robert, *Sur les équations confluentes de l'équation de Heun*, Ann.Soc. Sci. Bruxelles. **T92(III)** (1978) 151 - 189.
 - [4] R. Boyack and J. Lekner, *Confluent Heun functions and separation of variables in spheroidal coordinates* J. Math. Phys. **52** (2011) 073517.
 - [5] M. S. Cunha and H. R. Christiansen, *Confluent Heun functions in gauge theories on thick braneworlds*, Phys. Rev.D **84** (2011) 085002.
 - [6] E.S. Cheb-Terrab, *Solutions for the General, Confluent and Biconfluent Heun equations and their connection with Abel equations*, J. Phys. A: Math. Gen. **37** (2004) 9923.
 - [7] L.J. El-Jaick, D. B. Bartolomeu D. B. Figueiredo, *On certain solutions for Confluent and Double-Confluent Heun equations*, J. Math. Phys. **49** (2008) 083508.
 - [8] R. L. Hall, N. Saad, and K. D. Sen, *Soft-core Coulomb potentials and Heun's differential equation*, J. Math. Phys. J. Math. Phys. **51** (2010) 022107.
 - [9] P. P. Fiziev, *Novel relations and new properties of confluent Heun's functions and their derivatives of arbitrary order*, J. Phys. A: Math. Theor. **43** (2010) 035203.
 - [10] P. P. Fiziev and D. Staicova, *Application of the confluent Heun functions for finding the quasinormal modes of nonrotating black holes*, Phys. Rev. D **84** (2011) 127502.
 - [11] J. Favard, *Sur les polynomes de Tchebicheff*, C. R. Acad. Sci. Paris **200** (1935) 2052 - 2053.
 - [12] T. S. Chihara, *An introduction to orthogonal polynomials*, Gordon and Breach, 1978; reprinted, Dover, 2011.
 - [13] M. E. H. Ismail, *Classical and quantum orthogonal polynomials in one variable*, Cambridge University Press, 2005; corrected reprint, 2009.
 - [14] C. M. Bender, G. V. Dunne, *Quasi-exactly solvable systems and orthogonal polynomials*, J. Math. Phys. **37** (1996) 6 - 11.
 - [15] C. A. Downing, *On a solution of the Schrödinger equation with a hyperbolic double-well potential*, J. Math. Phys. **54** (2013) 072101.
 - [16] W. Fa-Kai, T]Y. Zhan-Ying, L. Chong, Y. Wen-Li and Z. Yao-Zhong, *Exact polynomial solutions of Schrödinger equation with various hyperbolic potentials*, Commun. Theor. Phys. **61** (2014) 153 - 159.
 - [17] A. Krajewska, A. Ushveridze, and Z. Walczak, *BenderDunne orthogonal polynomials general theory*, Mod. Phys. Lett. A **12** (1997) 1131
 - [18] N. Saad, R. L. Hall, and H. Ciffci, *Sextic anharmonic oscillators and orthogonal polynomials*, J. Phys. A: Math. Gen. **39** (2006) 8477.
 - [19] K. Chadán, N.N. Khuri, A. Martín and T.T. Wu, *Bound States in one and two spatial dimensions*, J. Math.Phys. **44** (2003) 406, arXiv:math-ph/0208011.