PROOF OF CONVERGENCE FOR THE LATTICE MONOMER-DIMER CLUSTER EXPANSION I, A SIMPLIFIED MODEL

P. FEDERBUSH DEPARTMENT OF MATHEMATICS UNIVERSITY OF MICHIGAN ANN ARBOR, MI 48109-1043, USA

ABSTRACT. We present some promising ideas to treat the problem of making completely rigorous the development of our expression for $\lambda_d(p)$ of the monomer-dimer problem on a *d*-dimensional hypercubic lattice

$$\lambda_d(p) = \frac{1}{2} \left(p \ln(2d) - p \ln(p) - 2(1-p) \ln(1-p) - p \right) + \sum_{k=2}^{\infty} a_k(d) p^k \quad (A)$$

where $a_k(d)$ is a sum of powers $(1/d)^r$ for

$$k - 1 \le r \le k/2 \tag{B}$$

In fact as we will point out one has allready rigorously established the convergence of the sum in A for small p. It is the d dependence of $a_k(d)$ that has yet to be rigorously shown. We do not now know how to complete the proof.

1. The Model Problem

We start by presenting the model problem. It is interesting in its own right, and is after all the study of this paper. r > 0 is given and real J_i are given, $i \ge 2$ satisfying $|J_i| \leq r^i$

$$Z = Z(N, p) = \sum_{\alpha_i} \prod_i \left((J_i p^i N)^{\alpha_i} \cdot \frac{1}{\alpha_i!} \right)$$
(2)

The $\alpha_i, i \geq 2$ are non-negative integers, and the sum over the a_i in (2) is over all values of the α_i , restricted by

$$\sum_{i=2}^{\infty} i\alpha_i \le \frac{pN}{2} \tag{3}$$

(1)

We will prove

We define

Theorem. There is a $p_0 > 0$ such that for $0 \le p \le p_0$

$$\lim_{N \to \infty} \frac{\ln Z}{N} = \sum_{2}^{\infty} p^{i} J_{i} \tag{4}$$

The easiest way to prove this theorem (probably) must be to estimate the corresponding sum to that in (2) with the α_i satisfying the complementary inequality

$$\sum_{i=2}^{\infty} i\alpha_i > \frac{pN}{2} \tag{5}$$

P. FEDERBUSH

If this part is "small enough" then Z becomes approximately equal

$$Z \cong \prod_{i} \left(\sum_{\alpha_i} (J_i p^i N)^{\alpha_i} \cdot \frac{1}{\alpha_i!} \right)$$
(6)

and the theorem is easily proved

We follow a much more devious route to the proof, in this paper. A route whose steps can all be paralleled in the actual problem.

2. INTRODUCTION

The cluster expansion approach to λ_d of the dimer problem on a hypercubic lattice was presented in [1]. A formal argument was given for the expansion

$$\lambda_d \sim \frac{1}{2} \ln(2d) - \frac{1}{2} + \sum_i C_i \frac{1}{d^i}$$
 (7)

At present the status of this putative asymptotic expansion is still not clear. In [4], working with Friedland, a simple extension of the expansion formalism of [1] was made to treat the monomer-dimer problem. This yielded the expression

$$\lambda_d(p) = \frac{1}{2} \left(p \ln(2d) - p \ln p - 2(1-p) \ln(1-p) - p \right) + \sum_{k=2} a_k(d) p^k$$
(8)

where $a_k(d)$ is a sum of powers of $(1/d), (1/d)^r$, with

$$k-1 \ge r \ge k/2. \tag{9}$$

The situation is a little complicated. The cluster expansion formalism yields an expression for $a_k(d)$ as a function of the Mayer Series coefficients of the dimer gas. In [4], repeated in eq (12)–(20) of [3], another route was obtained to derive the $a_k(d)$ from the mayer series coefficients. Although the two routes certainly give the same answer, this has not been rigorously established. The inscrutable [WORDS] identity that must be proved in detail in [5]. It is child's play to see that the second route leads rigorously to an expression (8) for $\lambda_d(p)$ where the sum converges. But it is the expression in (8) derived by the first route (still not proved rigorously) for which the $a_k(d)$ have the expression in powers of 1/d described before (9). Proof of the convergence of the cluster expansion, which we enter into in this paper, will show that the two expressions for all $a_k(d)$ are equal, that the sum in (8) converges for small p, and that the $a_k(d)$ have the indicated dependence on d.

The limit that must be evaluated to rigorously establish the cluster expansion development is presented in Section 3. The relation of the limit of the model problem, eq (4), to this limit will be clear. In succeeding sections the theorem of eq (4) is proven, taking care continuously to carry out steps as they can easily be applied to the general limit of Section 3. The basic strategy is to arrange $\ln Z$ as the sum of terms, chunks. Within the chunks some sums are replaced by contour integrals (in many complex variables). A single stationary point of the integrand [WORDS] in the limit of those integrals.

3. The Object of Study

We must analyze Z^*

$$Z^* = \sum_{\alpha_i} \beta(N, \sum i\alpha_i) \prod \bar{J}_i^{\alpha_i} \frac{N^{\sum \alpha_i}}{\prod (a_i!)}$$
(10)

This is eq (5.24) of [2]. Here the $\alpha_i, i \leq 2$, are non-negative integers and are restricted by eq (3)

$$\beta(N, jN) = e^{NH(p,j)} \tag{11}$$

with

$$\sum i\alpha_i = jN \tag{12}$$

and

$$H(p,j) = j \ln p + (1-2j) \ln(1-2j) + j - \frac{p}{2} \left(1 - \frac{2j}{p}\right) \cdot \ln\left(1 - \frac{2j}{p}\right)$$
(13)

$$\equiv j\ln p + \tilde{H}(p,j) \tag{14}$$

Eq (11) is eq (5.16) of [2], eq (13) is eq (5.17) of [2]. Eq (14) defines \tilde{H} . We also define $\tilde{\beta}$

$$\beta(N, \sum i\alpha_i) \equiv p^{\sum i\alpha_i} \tilde{\beta}(N, \sum i\alpha_i) \equiv p^{\sum i\alpha_i} e^{N\tilde{H}(p,j)}$$
(15)

 Z^* becomes

$$Z^* = \sum_{\alpha_i} \tilde{\beta}(N, \sum i\alpha_i) \prod (\bar{J}_i p^i N)^{\alpha_i} \frac{1}{\prod (\alpha_i !)}$$
(16)

Sums are again restricted by (3).

One wants to study

$$\lim_{N \to \infty} \frac{\ln Z^*}{N} \tag{17}$$

The similarity between eq (16) and eq (17) and the pair of equations, eq (2) and eq (4), is obvious. The \bar{J}_i have a weak dependence on N. (They are asymptotically constant.)

The desired limit of (17) we do not detail now. This limit might be found in (5.31) and (5.32) of [2], or in an entirely different form in the discussion surrounding (24)–(28) in [3], where another reference is given.

4. FROM SUMS TO CONTOUR INTEGRALS

In the next section Z or Z^* from (2) or (16) will be arranged into a sum of terms called chunks. In some of these chunks there will be a designated set of indices, S such that a portion of the chunk is of the form

$$\prod_{i \in \mathcal{S}} \left(\sum_{i=0}^{m_i} \frac{(J_i p^i N)^{\alpha_i}}{\alpha_i!} \right) \beta \tag{18}$$

Here if it is Z we are working with $\beta = 1$, it is Z^* , J_i becomes \bar{J}_i ; from now on such trivial differences will not be commented on. In (18) the other α_i are not summed, having been assigned certain values and β may depend on α_i . The J_i for i in Swill be negative, it will be important to control cancellations between positive and negative terms in evaluating (18) accurately enough. This motivated the use of contour integrals. In fact dealing with Z a simpler treatment is possible as will be pointed out later, but we want to use a method that applies to both Z and Z^* .

We set $-a \equiv J_i N p^i$ and note

$$\sum_{\alpha=0}^{n} \frac{(-a)^{\alpha}}{\alpha!} f(\alpha) = \frac{1}{2\pi i} \oint_{C} dz \frac{\pi}{\sin \pi z} \frac{a^{z}}{z!} f(z)$$
(19)

where the contour C is counterclockwise and contains $\{0, 1, ..., n\}$ and no other singularities of the integrand. Employing the identity (19) in (18) for all the α_i with *i* in S we have converted all the sums in our chunk to a single multivariable contour integral. Analytic properties of $\Gamma(z)$ and β will be dealt with later.

The division of Z and Z^* into chunks to convert sums from having limits as given by (3) to limits as in (19)

$$0 \le \alpha_i \le m_i \tag{20}$$

In the space of allowed α_i we are fitting hyper rectangles. This messy procedure is not illucidated. It is we feel the central idea of the proof.

5. Disection into Chunks

We write this section in the language of Z^* of (16), changes for Z of (2) trivial. We first introduce a number of parameters. There is \tilde{M}

$$\tilde{M} = \frac{pN}{4} \tag{21}$$

and m_i , for $i \geq 2$,

$$m_i = \frac{1}{i^{2+\epsilon i}} \tag{22}$$

Each chunk is assigned a "level", a non-negative integer, and is either "free" or "boxed". It requires patience to develop the construction of these chunks.

We let \mathcal{P} be the subset of indices for which $\overline{J}_i \geq 0$ if $i \in \mathcal{P}$ and \mathcal{N} be the subset for which $\overline{J}_i < 0$. Each chunk has a unique α_i assigned to the $i \in \mathcal{P}$, say $\alpha_i = t_i$. Thus in a chunk some of the α_i may be summed over, but not the α_i with $i \in \mathcal{P}$. For any chunk we define

$$R_0 = \sum_{i \in \mathcal{P}} i t_i \tag{23}$$

Level-zero free chunks. A level zero chunk is uniquely specified by the set of t_i , $i \in \mathcal{P}$. If

$$R_0 \ge \tilde{M} \tag{24}$$

then it is a free level-zero chunk. Its precise definition is

$$\prod_{i \in \mathcal{P}} (\operatorname{subs}(\alpha_i = t_i)) \sum_{\substack{\alpha_i, i \in \mathcal{N} \\ \sum_{i \in \mathcal{N}} i\alpha_i \le \frac{pN}{2} - R_0}} \tilde{\beta}(N, \sum_{i \alpha_i}) \left(\prod_i (\bar{J}_i p^i N)^{\alpha_i} \frac{1}{\prod(\alpha_i!)} \right)$$
(25)

We are using a Mapple-like notation. In (25) the α_i for $i \in \mathcal{P}$ are the set equal to t_i , and the remaining α_i are summed over subject to the restriction from (3). As with all the chunks, this chunk is some subsum of the terms in (16). Different chunks are disjoint, the union of all the chunks giving all terms in (16).

Level-zero boxed chunks. Here

$$R_0 < M \tag{26}$$

We define C_0 by

$$C_0 = \sup_{x} \left\{ x \mid \sum_{i \in \mathcal{N}} i \cdot \text{floor}(xU_i) \le \frac{pN}{2} - R_0 \right\}$$
(27)

where floor(α) is the largest integer $\leq \alpha$. We the set

$$m_0(i) = \text{floor}(C_0 U_i) \tag{28}$$

The level-zero boxed chunk defined by the $t_i, i \in \mathcal{P}$ is then

$$\prod_{i \in \mathcal{P}} (\operatorname{subs}(\alpha_i = t_i)) \sum_{\alpha_{a_1} = 0}^{m_0(a_i)} \cdots \sum_{\alpha_{a_s} = 0}^{m_0(a_s)} M$$
(29)

Here M indicates everything after the sums in (16). a_1, \ldots, a_s are the indices labelling elements of \mathcal{N} . By using (27) we have found the biggest box, of a certain shape, we can insert in the sum. That is we are picking the upper limits in (29) as large as possible, with a certain fixed ratio between them

The $P_0 \geq \tilde{M}$ in the free chunk will lead to enough smallness in estimates later that one will not have to study cancellations between signed terms via the contour integrals of (19). For the boxed chunks one will need to do so.

Level-one chunks. The level-zero chunks fail to exhaust all terms In (16) because of terms containing some α_i for some $i \in \mathcal{N}$ exceeding the upper limits in (29). We give a subset \mathcal{B}_1 of \mathcal{N} and to each i in \mathcal{B}_1 we associate a t_i with

$$t_i > m_0(i) \tag{30}$$

We call the augmented set \mathcal{B}_1 of indices and associated t_i , $\overline{\mathcal{B}}_1$. We set

$$R_1 = \sum_{i \in \mathcal{B}_1} it_i \tag{31}$$

If $R_0 + R_1 \ge \tilde{M}$ we have the level-one free chunk given as

$$\prod_{i \in \mathcal{P} \cup \mathcal{B}_1} (\operatorname{subs}(\alpha_i = t_i)) \sum_{\substack{\alpha_i, i \in \mathcal{N}_1 \\ \sum_{i \in \mathcal{N}_1} i \alpha_i \leq \frac{pN}{2} - R_0 - R_1}} M$$
(32)

We have defined $\mathcal{N}_1 = \mathcal{N} - \mathcal{B}_1$. To define the boxed chunk we define

$$C_1 = \sup_{x} \left\{ x \mid \sum_{i \in \mathcal{N}_1} i \operatorname{floor}(xU_i) \le \frac{pN}{2} - R_0 - R_1 \right\}$$
(33)

and

$$m_1(i) = \text{floor}(C_1 U_i) \tag{34}$$

Then the level-one boxed chunk is given as

$$\prod_{\in \mathcal{P} \cup \mathcal{B}_1} (\operatorname{subs}(\alpha_i = t_i)) \sum_{\alpha_{a_1} = 0}^{m_1(a_1)} \cdots \sum_{\alpha_{a_s} = 0}^{m_1(a_s)} M$$
(35)

 $i \in \mathcal{P} \cup \mathcal{B}_1$ α_a and here the a_1, \ldots, a_s label elements of \mathcal{N}_1 .

The set of t_i associated to the i in $\mathcal{P} \cap \mathcal{B}_1$ uniquely label the level-one chunks.

General level chunks. We assume we have defined chunks of level-zero through level-n, and we will derive expressions for the level-(n + 1) chunks. Thus we have

$$\mathcal{N} = \mathcal{N}_0 \supset \mathcal{N}_1 \supset \mathcal{N}_2 \cdots \supset N_n \tag{36}$$

$$\mathcal{B}_i \subset \mathcal{N}_{i-1}, \qquad i = 1, 2, \dots, n \tag{37}$$

$$\mathcal{N}_i = \mathcal{N}_i - \mathcal{B}_i, \qquad i = 1, 2, \dots, n \tag{38}$$

$$\bar{\mathcal{B}}_i, \qquad i = 1, 2, \dots, n \tag{39}$$

That is, we have an assignment $\alpha_i = t_i$ for *i* in each \mathcal{B}_i . We set

$$R_k = \sum_{i \in \mathcal{B}_k} it_i, \qquad k = 1, 2, \dots, n \tag{40}$$

and have C_0, C_1, \ldots, C_n with

$$C_k = \sup_{x} \left\{ x \mid \sum_{i \in \mathcal{N}_k} i \operatorname{floor}(xU_i) \le \frac{pN}{2} - \sum_{0}^k Rk \right\}, \qquad k = 0, 1, \dots, n$$
(41)

We set

$$m_k(i) = \text{floor}(C_k U_i), \qquad k = 0, 1, \dots, n \tag{42}$$

and for i in \mathcal{B}_k , $k = 2, \ldots, n$ one has

$$m_{k-1}(i) < t_i \le m_{k-2}(i)$$
 (43)

For k = 1 the analogous condition is given by (30); there is in this case no upper bound here imposed. It is not difficult to see the C_i from a decreasing sequence.

To go to the next level, select the set $\mathcal{B}_{n+1} \subset \mathcal{N}_n$ and for $i \in \mathcal{B}_{n+1}$ require

$$m_n(i) < t_0 \le m_{n-1}(i)$$
 (44)

The rest follows immediately.

Let us specify free and boxed chunks of level-(n + 1). The free level n + 1 chunk arises if

$$\sum_{i=0}^{n+1} R_i \ge \tilde{M} \tag{45}$$

and then is given as

$$\prod_{i \in \mathcal{P} \cup (\mathcal{N} - \mathcal{N}_{n+1})} (\operatorname{subs}(\alpha_i = t_i)) \sum_{\substack{\alpha_i, i \in \mathcal{N}_{n+1} \\ \sum_{i \in \mathcal{N}_{n+1}} i\alpha_i \leq \frac{pN}{2} - \sum_{i=0}^{n+1} R_i}} M_i$$
(46)

If (45) is not satisfied the boxed chunk is given as

$$\prod_{i \in \mathcal{P} \cup (\mathcal{N} - \mathcal{N}_{n+1})} (\operatorname{subs}(\alpha_i = t_i)) \sum_{\alpha_{a_1} = 0}^{m_{n+1}(a_1)} \cdots \sum_{\alpha_{a_s} = 0}^{m_{n+1}(a_s)} M$$
(47)

where a_1, \ldots, a_s are the indices of \mathcal{N}_{n+1} .

6. Smallness from high occupation

In this section we find an upper bound on

$$\prod_{i} \left(\frac{(|\bar{J}_i| p^i N)^{\alpha_i}}{\alpha_i!} \right) \tag{48}$$

subject to the restriction

$$\sum_{i} i\alpha_i \ge \tilde{M} \tag{49}$$

We use (3)

$$|\bar{J}_i| \le r^i \tag{50}$$

and define

$$Q = \prod_{i} \left(\frac{((pr)^{i}N)^{\alpha_{i}}}{\alpha_{i}!} \right)$$
(51)

One then has

$$\ln Q \le F = N \sum_{i} [ix_i \ln(pr) - x_i \ln x_i + x_i]$$
(52)

using the well known inequality $n \ln(n/e) + 1 \leq \ln n!$, where we have set

$$\alpha_i = x_i N \tag{53}$$

$$\tilde{M} = \tilde{m}N$$
 (54)

Then we have

$$f = \frac{F}{N} = \sum_{i} [ix_{i} \ln(pr) - x_{i} \ln x_{i} + x_{i}]$$
(55)

We ignore the restriction that α_i must be an integer. We note that our upper bound will apply if we restrict the set of indices in the products of (48) or (51), or require the α_i to be integers, each of these would lead to a lower upper bound.

Now we want to maximize f subject to

$$h = \sum_{i} x_i = \tilde{m} \tag{56}$$

We apply Lagrange multipliers

$$\frac{d}{dx_i}(f - \lambda h) = 0 \tag{57}$$

getting

$$x_i = (pre^{-\lambda})^i \tag{58}$$

and then, from (56)

$$\sum_{i=2}^{\infty} (pre^{-\lambda})^i = \tilde{m}$$
(59)

one gets for $\tilde{m} = cp$ as p goes to zero

$$F \cong -N\tilde{m}|\ln(p/\sqrt{\tilde{m}})| \tag{60}$$

7. THE CONTOUR INTEGRALS, DISTORTING THE CONTOURS

We have displayed the expressions for boxed chunks of level-zero, level-one, and general level-(n+1) in eqs. (29), (35), (47). We isolate the portion of the expression shown in eq. (18), and relabelling indices look at our present object or study

$$A = \sum_{\alpha_1=0}^{m_1} \cdots \sum_{\alpha_s=0}^{m_s} \prod_{i=1}^s \left(\frac{(J_{d_i} p^{d_i} N)^{\alpha_i}}{\alpha_i!} \right) \beta \tag{61}$$

Here all the J_i are negative. We set

$$a_i = -J_{d_i} p^{d_i} N \tag{62}$$

and rewrite A as

$$A = \frac{1}{(2\pi i)^s} \oint_{C_1} dz_1 \cdots \oint_{C_s} dz_s \prod_{i=1}^s \left(\frac{\pi}{\sin \pi z_i} \frac{a_i^{z_i}}{\Gamma(z_i+1)} \right) \beta \tag{63}$$

where we take C_i as hugging the interval $[-1/2, m_i + 1/2]$, encircling it counterclockwise.

We let g be the integrand of (63) so that

$$A = \frac{1}{(2\pi i)^s} \oint_{C_1} dz_1 \cdots \oint_{C_s} dz_s g \tag{64}$$

and seek a stationary point of g, so we solve together

$$\frac{\partial}{\partial z_i}g = 0, \qquad i = 1, \dots, s \tag{65}$$

We now use

$$\Gamma(z)\Gamma(1-z) = \frac{\pi}{\sin \pi z} \tag{66}$$

to write g as

$$g = \prod_{i=1}^{s} \left(-a_i^{z_i} \Gamma(-z_i) \right) \beta \tag{67}$$

We look at the equations (65) for a stationary point of Z, where then $\beta = 1$, getting

$$\frac{\partial}{\partial z_i}(-a_i^{z_i}\Gamma(-z_i)) = 0 \tag{68}$$

or taking exponentials and setting $w_i = -z_i$

$$\frac{d}{dw_i}[-w_i\ln a + \ln\Gamma(w)] = 0 \tag{69}$$

We now use the standard large w approximation

$$\ln n! \cong n \ln n - n \tag{70}$$

to get

$$w_i \cong a_i + 1 \tag{71}$$

or

$$z_i = z_{i0} \cong -a_i - 1 \tag{72}$$

at the stationary point.

In general let the coordinates of the stationary point be $z_i = z_{i0}$. We then first shift the contour C_i to C'_i where C'_i hugs the interval $[-1/2 + \text{floor}(z_{i0}), m_i + 1/2]$ counterclockwise. One hits no singularities of g to distort the contour so. Next we stretch the contour to infinity, plus and minus, in the imaginary directions, leaving contours C''_i . C''_i consists of two line segments parallel to the imaginary axis

$$C_{i}'' = [-\infty i - 1/2 + \text{floor}(z_{i}), \infty i - 1/2 + \text{floor}(z_{i})]$$
union
$$[-\infty i + m_{i} + 1/2, \infty i + m_{i} + 1/2]$$
(73)

Again it will not be hard to show that no singularities of the integrand are crossed in the distortion, and the segments at ∞ to close the C_i'' contours contribute nothing.

In treating Z^* and studying

$$A = \frac{1}{(2\pi i)^s} \oint_{C_i''} dz_1 \cdots \oint_{C_s''} dz_s g \tag{74}$$

one will have to show the limit in eq (17) picks out just the value of the integrand at the center of the first line segment in (73). In studying Z we will use a simpler route to study (61). (For one thing we do not want to worry about situations where a_i is not large, that is not guaranteed by (1)... perhaps there is no problem here at all.)

8. Completion of the Proof for Z

In this section we carry through the proof of the theorem, see eq. (4), for Z, much in the way we plan to complete the proof for Z^* , see eq. (17). An important difference is that we do not need use of contour integration. We divide Z into a sum of three terms

$$Z = Z(N, p) = T_1 + T_2 + T_3$$
(75)

 $T_1(N,p)$ is the sum of level-zero boxed chunks, $T_2(N,p)$ is the sum of the rest of the boxed chunks, and $T_3(N,p)$ is the sum of the free chunks. We will arrive at the theorem by proving, for small enough p, that

$$\lim_{N \to \infty} \frac{\ln T_1}{N} = \sum_2 p^i J_i \tag{76}$$

$$\lim_{N \to \infty} \frac{\ln T_2}{N} < \sum_2 p^i J_i \tag{77}$$

$$\lim_{N \to \infty} \frac{\ln T_3}{N} < \sum_2 p^i J_1 \tag{78}$$

Study of T_3 . We may overestimate T_3 as follows

$$T_3 \le \sum_{\alpha_i} Q \tag{79}$$

where Q is from (51) and the α_i are restricted by

$$\sum i\alpha_i \ge \frac{pN}{4} \tag{80}$$

Then we write

$$\sum_{\alpha_{i}} Q = \sum_{\alpha_{i}} \prod_{i} \left(\frac{((pr)^{i}N)^{\alpha_{i}}}{\alpha_{i}!} \right)$$

$$\leq \sum_{\alpha_{i}} \prod_{i} \left(\frac{((pr)^{i}N)^{\alpha_{i}/2}}{(\alpha_{i}/2)!} \right) \prod_{i} \left(\frac{((pr)^{i}N)^{\alpha_{i}/2}}{(\alpha_{i}/2)!} \right)$$
(81)

all subject to the restriction (80). This yields

$$\sum_{\alpha_i} Q \le AB \tag{82}$$

with

$$A = \sup \prod_{i} \left(\frac{((pr)^{i}N)^{\alpha_{i}/2}}{(\alpha_{i}/2)!} \right)$$
(83)

and

$$B = \sum_{i} \prod_{i} \left(\frac{((pr)^{i}N)^{\alpha_{i}/2}}{(\alpha_{i}/2)!} \right)$$
(84)

In (82) and (83) restriction (80) is enforced, but getting a weaker inequality do not impose it in (84). For A from (60) we get

$$A \cong e^{-N[WORDS]} \tag{85}$$

Looking at B we note

$$\sum_{i} \frac{a^{i/2}}{(i/2)!} = \sum_{i} \frac{a^{i}}{i!} + \sum_{i} \frac{a^{1/2+i}}{(1/2+i)!}$$
$$= \sum_{i} \frac{a^{i}}{i!} + a^{1/2} \sum_{i} \frac{a^{i}}{(1/2+i)!}$$
$$\leq (1 + ca^{1/2}) \sum_{i} \frac{a^{i}}{i!} \leq (1 + ca^{1/2})e^{a}$$
(86)

Equation (78) follows for p small enough with little work.

9. Dealing with Z_1 and Z_2

In this section we study the methods used to treat T_1 and T_2 to derive (76) and (77). We discuss the proof of (76) alone, since (77) requires no new ideas. We write T_1 , the sum of the level zero boxed chunks as

$$T_1 = \sum_{\beta} A^{\beta} B^{\beta} \tag{87}$$

with

$$A^{\beta} = \prod_{i \in \mathcal{P}} \frac{(J_i p^i N)^{t_i}}{t_i!}$$
(88)

where $t_i = t_i(\beta)$ satisfy

$$R_0 = R_0(\beta) = \sum it_i < \frac{pN}{4} \tag{89}$$

 $C_0(\beta)$ is given by (27), and $m_0(i)$ by (28), using which we define

$$B^{\beta} = \prod_{i \in \mathcal{N}} \left(\sum_{\alpha_i=0}^{m_0(i)} \frac{(J_i p^i N)^{\alpha_i}}{\alpha_i!} \right)$$
(90)

We now set

$$B_0 = \prod_{i \in \mathcal{N}} (e^{J_i p^i N}) \tag{91}$$

We write

$$A^{\beta}B^{\beta} = A^{\beta}B_0 + A^{\beta}E^{\beta} \tag{92}$$

and

$$\sum A^{\beta}B^{\beta} = B_0 \sum A^{\beta} + \sum A^{\beta}E^{\beta}$$
(93)

We study the second term in the right side of (93) first

2

$$\left|\sum A^{\beta} E^{\beta}\right| \le \left(\sum A^{\beta}\right) \sup_{\beta} |E^{\beta}| \tag{94}$$

If we were studying Z^* instead of Z we would have used contour integral techniques, but in the present case we use simpler methods

$$\sum A^{\beta} \le e^{\sum_{i \in \mathcal{P}} J_i p^i N} \tag{95}$$

To study $|E^{\beta}|$ we first introduce

$$g(a,n) = \sum_{i=n+1}^{\infty} \left(\frac{|a|^i}{i!}\right) e^{+|a|}$$
(96)

10

and

$$g_i = g(J_i p_i N, m_0(i)) \tag{97}$$

In terms of the quantities we have

$$|E^{\beta}| \le e^{\sum_{i} J_{i} p^{i} N} \cdot \left(\prod_{i \in \mathcal{N}} (1+g_{i}) - 1\right)$$
(98)

The dependence on β comes from the dependence of g_i on the $m_0(i)$ which depend on β .

If $x_i \ge 0$ and $\sum_i \le 1$ one has the inequality

$$\prod (1+x_i) - 1 \le e \sum x_i \tag{99}$$

One we know $\sum g_i \leq 1$ we now have

$$\left|\sum A^{\beta} E^{\beta}\right| \le e^{\sum J_i p^i N} \cdot e \cdot \sup_{\beta} \left(\sum_{i \in \mathcal{N}} g_i\right)$$
(100)

Limit (76) follows from

$$\lim_{N \to \infty} \frac{1}{N} \ln \left(B_0 \sum A^\beta \right) = \sum J_i p^i \tag{101}$$

and

$$\lim_{N \to \infty} \frac{\left|\sum A^{\beta} E^{\beta}\right|}{B_0 \sum A^{\beta}} = 0$$
(102)

We choose to see (101) by approximating $\sum A^{\beta}$ by its largest term, a justifiable procedure in this case. Note that all terms in sum are positive This is the method used in [1] to treat (27) there, and again in [2] to treat (5.24) therein. In both these cases the method is used "formally" where it does not really apply since there are positive and negative terms. It is the work of the present paper and its sequel, part 2, to show the results in [1] and [2] are none the less correct.

We turn to understanding limit (102). From (100) we have

$$\frac{\left|\sum A^{\beta} E^{\beta}\right|}{B_0 \sum A^{\beta}} \le \frac{e^{\sum J_i p^i N}}{B_0 \sum A^{\beta}} \cdot e \cdot \sup_{\beta} \left(\sum_{i \in \mathcal{N}} g_i\right)$$
(103)

From (101), now presumed true this becomes

$$\frac{\left|\sum A^{\beta} E^{\beta}\right|}{B_0 \sum A^{\beta}} \le e^{N\varepsilon(N)} \cdot e \cdot \sup_{\beta} \left(\sum_{i \in \mathcal{N}} g_i\right)$$
(104)

for some $\varepsilon(N)$ that goes to zero with N. We may deduce an upper bound on $\sum_{i \in \mathcal{N}} g_i$ [WORDS] $\sum_{i \in \mathcal{N}} \bar{g}_i$ with \bar{g}_i, g_i computed with $R_0 = \frac{pN}{4}$. We get an accurate upper bound for the $\sum_{i \in \mathcal{N}} \bar{g}_i$ by looking at

$$h(r^2 p^2 N, \gamma p N) \tag{105}$$

with

$$h(a,n) = \frac{a^n/n!}{e^{-a}}$$
(106)

$$\ln h(r^2 p^2 N, \gamma p N) \le -(\gamma p N) \ln \left(\frac{\gamma p N}{r^2 p^2 N}\right) + (\gamma p N) - r^2 p^2 N \tag{107}$$

implying

$$h \le e^{-\tilde{\gamma}N} \tag{108}$$

with $\tilde{\gamma} > 0$. This and (104) yield (102) and finally (76).

References

- [1] P. Federbrush, Computation of Terms in the Asymptotic Expansion of Dimer λ_d for High Dimension Physics Letters A, 374 (2009) 131-133.
- [2] P. Federbush and S. Friedland, An Asymptotic Expansion and Recursive Inequalities for the Monomer-Dimer Problem, J. Stat. Phys. 143, 306 (2011).
- [3] P. Butera, P. Federbush, and M. Peruici, Higher-order Expansion for the Entropy of a Dimer or a Monomer-Dimer System on d-dimensional Lattices, Phys. Rev. E 87, 062113 (2013).
- [4] P. Federbush, The Dimer Gas Mayer Series, the Monomer-Dimer $\lambda_d(p)$, the Federbush Relation, arXiv:1207.1252.
- [5] P. Federbush, For the Monomer-Dimer $\lambda_d(p)$, the Master Algebraic Conjecture. arXiv:1209.0987.