Two dimensional heteroclinic attractor in the generalized Lotka-Volterra system.

Valentin S. Afraimovich^{*}, Gregory Moses[†], Todd Young[‡]

August 21, 2018

Abstract

We study a simple dynamical model exhibiting sequential dynamics. We show that in this model there exist sets of parameter values for which a cyclic chain of saddle equilibria, O_k , $k = 1, \ldots, p$, have two dimensional unstable manifolds that contain orbits connecting each O_k to the next two equilibrium points O_{k+1} and O_{k+2} in the chain $(O_{p+1} = O_1)$. We show that the union of these equilibria and their unstable manifolds form a 2-dimensional surface with boundary that is homeomorphic to a cylinder if p is even and a Möbius strip if p is odd. If, further, each equilibrium in the chain satisfies a condition called "dissipativity," then this surface is asymptotically stable.

1 Background

In the last decade it became clear that typical processes in many neural and cognitive networks are realized in the form of sequential dynamics (see [\[23\]](#page-25-0), [\[14\]](#page-24-0), [\[24\]](#page-25-1), [\[26\]](#page-25-2), [\[25\]](#page-25-3), [\[2\]](#page-24-1) and references therein). The dynamics are exhibited as sequential switching among metastable states, each of which represents a collection of simultaneously activated nodes in the network, so that at most instants of time a single state is activated. Such dynamics are consistent with the winner-less competition principle [\[26,](#page-25-2) [2\]](#page-24-1). In the phase space of a mathematical model of such a system each state corresponds to an invariant saddle set and the switchings are determined by trajectories joining these invariant sets. In the simplest case these invariant sets may be saddle equilibrium points coupled by heteroclinic trajectories, and they form a heteroclinic sequence (HS). This sequence can be stable if all saddle equilibria have one-dimensional unstable manifolds [\[2\]](#page-24-1), in the sense that there is an open set of initial points such that trajectories going through them follow the heteroclinic ones in the HS, or unstable if some of the unstable manifolds are two-or-more dimensional. In the latter case properties of trajectories in a neighborhood of the HS were studied in [\[2\]](#page-24-1) and [\[3\]](#page-24-2). General results on the stability of heteroclinic sequences have been obtained

[∗]Universidad Autonoma de San Luis Potosi, IICO

[†]Corresponding author, Department of Mathematics, Ohio University, gm192206@ohio.edu

[‡]Ohio University, Department of Mathematics

by Krupa and Melbourne [\[17,](#page-25-4) [18\]](#page-25-5) in the form of necessary conditions that may also be sufficient in the presence of certain algebraic conditions.

An instability of a HS may be caused if some initial conditions follow trajectories on the unstable manifold different from the heteroclinic ones. M. Rabinovich suggested [\[22\]](#page-25-6) considering the case when all trajetories on the unstable manifold of any saddle in a HS are heteroclinic to saddles in the HS, which implies an assumption that the HS is, in fact, a heteroclinic cycle. In this case one can expect some kind of stability, not of the HS, of course, but of the object in the phase space formed by all heteroclinic trajectories of the saddles in the HS. We study in our paper the Rabinovich problem. We deal here with the generalized Lotka-Volterra model [\[2\]](#page-24-1) that is a basic model of sequential dynamics for which unstable sets are realized as saddle equilibrium points. All variables and parameters may take only nonnegative values, so we work in the positive orthant of the phase space \mathbb{R}^n . We impose some restrictions on parameters under which all unstable manifolds of the saddle point are two-dimensional and all trajectories on them (in the positive orthant) are heteroclinic in some specific way (see below). We prove that they form a piece-wise smooth manifold homeomorphic to the cylinder if the number of the saddle points in the HS is even or to the Möbius band if it is odd. We prove also that under the additional assumption that each equilibrium is dissipative (see below), then this manifold is the maximal attractor for some absorbing region (in the positive orthant). Trajectories in this region may follow different heteroclinic trajectories and may manifest some kind of weak complex behavior.

Although our motivations are neurological, we observe that heteroclinic networks (and thus, potentially, high-dimensional counterparts of the same) are ubiquitous, appearing in applications that range from celestial dynamics [\[16\]](#page-24-3) to evolutionary game theory [\[11\]](#page-24-4).

2 Notation and Results

We begin our consideration of two-dimensional heteroclinic channels with the study of a series of Lotka-Volterra equations. Lotka-Volterra models are widely used in the context of heteroclinic sequences where all the unstable manifolds are one dimensional, so that each equilibrium is connected to exactly one subsequent equilibrium (e.g. $[13, 28, 10, 9]$ $[13, 28, 10, 9]$ $[13, 28, 10, 9]$ $[13, 28, 10, 9]$). It has been recently shown that they ask provide a general method for embedding directed graphs into a system of ordinary differential equations [\[6\]](#page-24-8). We remark that although [\[6\]](#page-24-8) allows graphs of high valency to be modeled by heteroclinic networks, and some work has been done on the stability of such systems (e.g. [\[15\]](#page-24-9), which considers the competing dynamics of the "overlapping" channels $\lambda_1 \to \lambda_2 \to \lambda_3 \to \lambda_1$ and $\lambda_1 \to \lambda_2 \to \lambda_4 \to \lambda_1$), study of heteroclinic networks has usually only considered one-dimensional unstable manifolds. In keeping with the discussion of the introduction, we consider the dynamics of a system such that initial conditions arbitrarily close to any of $p \leq n$ saddle nodes may be mapped into neighborhoods of either one of two other saddle nodes. In particular, a simple general model for heteroclinic sequential dynamics was given in [\[4\]](#page-24-10) by

$$
\dot{x}_i = F_i(x) = x_i(\sigma_i - \sum_{j=1}^n \rho_{ij} x_j) \quad \text{for } i = 1, ..., n,
$$
\n(1)

Figure 1: A rough representation of the dynamics of the system defined by Equation [1,](#page-1-0) with $p = 5$.

where all of the parameters are assumed to be positive. The constants ρ_{ij} have biological meaning, representing inhibition of mode i by mode j . For the sake of simplicity, we assume that $\rho_{ii} = 1$ for all i. Further, since the variable x is assumed to encode biological information that is necessarily non-negative, e.g. activation levels or chemical concentrations, we restrict the system to the first closed orthant, $\overline{\mathbb{R}^n_+} = \{x \in \mathbb{R}^n : x_i \geq 0, 1 \leq i \leq n\}$. The system is so constructed that it contains saddle points lying on the axes, with σ_i being the *i*-th coordinate of the *i*-th saddle along the *i*-th axis $(\sigma_i > 0)$, i.e., the system [\(1\)](#page-1-0) has n equilibrium points of the form $O_k = (0, ..., 0, \sigma_k, 0, ...)$, for $k = 1, ..., n$. There may be other equilibria as well, but they are not relevant for our purposes; we only study transitions between the n equilibria just defined.

In the present work we suppose that the first p equilibria points are sequentially connected by a set of 2-dimensional unstable manifolds. For each $k, 1 \leq k \leq p$, there will be a heteroclinic orbit connecting O_k to O_{k+1} and a heteroclinic orbit connecting O_k to O_{k+2} . Furthermore, the system is closed in the sense that $O_{p+i} = O_i$, i.e. p is the modulus of the subscript. In the following, we will consider the restrictions necessary to enforce such dynamics.

To ensure that there are heteroclinic trajectories between O_k and both O_{k+1} and O_{k+2} , we apply eigenvalue conditions to the system. Consider first O_1 . The linearization of the vector field $F(x)$ [\(1\)](#page-1-0) at O_1 is given by the upper triangular matrix:

$$
DF(O_1) = \begin{pmatrix} -\sigma_1 & -\sigma_1 \rho_{12} & -\sigma_1 \rho_{13} & -\sigma_1 \rho_{14} & \cdots & -\sigma_1 \rho_{1n} \\ 0 & \sigma_2 - \rho_{21} \sigma_1 & 0 & 0 & \cdots & 0 \\ 0 & 0 & \sigma_3 - \rho_{31} \sigma_1 & 0 & \cdots & 0 \\ 0 & 0 & 0 & \sigma_4 - \rho_{41} \sigma_1 & \cdots & \vdots \\ \vdots & \vdots & \vdots & 0 & \ddots & \vdots \\ 0 & 0 & 0 & 0 & \cdots & \sigma_n - \rho_{n1} \sigma_1 \end{pmatrix}, \quad (2)
$$

and so the eigenvalues appear on the diagonal. The matrices $DF(O_k)$ have a similar simple structure, zeros everywhere except on the diagonal and on the k -th row. It is then easy to see that the eigenvalues of DF at O_k are

$$
\lambda_k^k = -\sigma_k \quad \text{and} \quad \lambda_j^k = \sigma_j - \rho_{jk}\sigma_k, \text{ for } j \neq k.
$$

In particular, the eigenvalues are all real. One can also see from the structure of [\(2\)](#page-2-0) that $DF(O_k)$ has a full set of eigenvectors, even if some eigenvalues are repeated.

Note that because of the particular form of the equations, all coordinate axes, planes and hyperplanes are invariant. Thus, in order that trajectories can travel from O_k to O_{k+1} or O_{k+2} , it is sufficient to put the restriction $0 < \lambda_{k+2}^k, \lambda_{k+1}^k$, and $\lambda_j^k < 0$ otherwise, to ensure that they can only go in those directions. In particular, for each $k, 1 \leq k \leq p$, we require that

$$
0 < \min_{i=1,2} \{ \sigma_{k+i} - \rho_{k+i,k} \sigma_k \} \text{ (indices mod } p), \tag{3}
$$

$$
\lambda_j = \sigma_j - \rho_{jk}\sigma_k < 0, \quad \text{for } j \neq k, k+1, k+2 \mod p. \tag{4}
$$

Note also that $-\sigma_k < 0$. These inequalities guarantee that each equilibrium is a hyperbolic saddle with 2 unstable directions and $n-2$ stable directions. Let $\widetilde{W}_k^u = W^u(O_k) \cap \overline{\mathbb{R}^n_+}$ be the unstable manifold of O_k restricted to the positive orthant. We show below that

$$
\Gamma \equiv \bigcup_{k=1}^p (\widetilde{W}_k^u \cup O_k)
$$

forms a piecewise smooth surface that we will classify topologically as follows.

Theorem 2.1. Suppose that inequalities [\(3\)](#page-3-0) and [\(4\)](#page-3-1) hold for each k, $1 \leq k \leq p$ and that each unstable manifold $\overline{W}^u(O_k)$ is contained in a compact forward invariant set as specified in Lemma [3.10](#page-8-0) (see also Remark 3.11). When p is even, the union of unstable manifolds Γ is homeomorphic to a cylinder. When p is odd, Γ is homeomorphic to a Möbius strip

The proof of this theorem, which is slightly involved, is left for the appendix.

Consider the following definition.

Definition 2.2. Let Σ^u and Σ^s be the set of stable and unstable eigenvalues, respectively, of the linearization of a vector field at a saddle equilibrium, i.e., $\max Re(\Sigma^s) < 0$ and $\min Re(\Sigma^u) > 0$. We say that the saddle is **dissipative** if

$$
\max Re(\Sigma^u) < -\max Re(\Sigma^s).
$$

In other words, the weakest stable eigenvalue is stronger than the strongest unstable eigen-value. (See [\[4\]](#page-24-10).)

In terms of the specific vector field under study all the eigenvalues in question are real and for each k we have:

$$
\max_{i=1,2} \{ \sigma_{k+i} - \rho_{k+i,k} \sigma_k \} < \min_{j \neq k, k+1, k+2} \{ |\sigma_j - \rho_{jk} \sigma_k|, \sigma_k \} \quad \text{(indices mod } p). \tag{5}
$$

The main goal of this manuscript is to show that, under the condition that each saddle equilibrium is dissipative, then Γ is asymptotically stable. Specifically, our main theorem is:

Theorem 2.3. Suppose that inequalities [\(3\)](#page-3-0), [\(4\)](#page-3-1) and [\(5\)](#page-3-2) hold for each k, $1 \leq k \leq p$ and that each unstable manifold $\widetilde{W}^u(O_k)$ is contained in a compact forward invariant set as specified in Section [3.3.](#page-8-1) Then Γ is asymptotically stable.

The proof of Theorem [2.3](#page-3-3) breaks roughly into two independent pieces. We start by considering the trajectory of a representative point that is ϵ -close to Γ, but is distant from each of the fixed points O_i . In such a case, the dynamics of the system are controlled largely by three consecutive saddles O_i , O_{i+1} , and O_{i+2} . In Section [3,](#page-4-0) we consider the restriction of [\(1\)](#page-1-0) to three consecutive dimensions; we will gain information on the full-dimensional system by viewing it as a perturbation of this restriction. The second part of the proof is to consider the dynamics as a trajectory passes near a fixed point; we consider this in Section [4.](#page-10-0)

In Section [4.2](#page-12-0) we prove the main theorem. In Section [4.3](#page-15-0) we show that there is a non-empty parameter set for which the conditions of the theorem are satisfied.

3 Three Dimensions

We begin our proof of Theorems 2.1 and 2.3 with a study of the restriction of the system to three dimensional sub-spaces corresponding to three consecutive coordinate directions. Through a series of geometric lemmas, we prove the main result of the section, Theorem [3.12,](#page-9-0) which provides information on the behavior of trajectories inside this invariant subspace. This will be used in later sections to complete the proofs of the main results by providing information on those parts of the full phase space where all but three coordinates are small.

3.1 Set-up

Let O_i , O_{i+1} , and O_{i+2} be any three consecutive equilibria and restrict the system [\(1\)](#page-1-0) to the three dimensions spanned by consecutive coordinates x_i , x_{i+1} , and x_{i+2} . For convenience, we will refer to these three variables as x_1, x_2 , and x_3 . Our goal in three dimensions is to show the existence of a compact, forward invariant set containing O_1 , O_2 , and O_3 such that any trajectory with an initial value in the interior of that set converges to O_3 .

Restricted to three dimensions, the equations [\(1\)](#page-1-0) are reduced to

$$
\begin{aligned}\n\dot{x}_1 &= x_1(\sigma_1 - x_1 - \rho_{12}x_2 - \rho_{13}x_3), \\
\dot{x}_2 &= x_2(\sigma_2 - x_2 - \rho_{21}x_1 - \rho_{23}x_3), \\
\dot{x}_3 &= x_3(\sigma_3 - x_3 - \rho_{31}x_1 - \rho_{32}x_2).\n\end{aligned} \tag{6}
$$

The restriction of the dimension of the unstable manifolds via the eigenvalue conditions, and the positivity conditions on σ_1, σ_2 , and σ_3 yield the following inequalities:

$$
-\sigma_1 < 0 < \sigma_j - \rho_{j1}\sigma_1, \ \ j = 2, 3,\tag{7}
$$

$$
-\sigma_2 < 0 < \sigma_3 - \rho_{32}\sigma_2,\tag{8}
$$

$$
\sigma_1 - \rho_{12}\sigma_2 < 0,\tag{9}
$$

$$
\sigma_1 - \rho_{13}\sigma_3 < 0,\tag{10}
$$

$$
\sigma_2 - \rho_{23}\sigma_3 < 0. \tag{11}
$$

Of those inequalities, [\(7\)](#page-4-1) controls the behavior of the system at O_1 , [\(8\)](#page-4-2) - [\(9\)](#page-4-3) control the behavior of the system at O_2 , and (10) - (11) at O_3 . The point $(\sigma_1, 0, 0)$ is a saddle with a two-dimensional unstable manifold, $(0, \sigma_2, 0)$ is a saddle with a one-dimensional unstable manifold, and $(0, 0, \sigma_3)$ is a sink for the system [\(6\)](#page-4-6). We remark that although (6) has the form of the May-Leonard model, the particular parameter restrictions under consideration yield simple dynamics (see Theorem [3.12\)](#page-9-0), and prevent the more complex behavior usually studied in that context. In particular, they are inconsistent with the symmetric May-Leonard model as it was introduced in [\[19\]](#page-25-8).

Figure 2: Illustration of dynamics of the system projected onto the first three coordinates. All consecutive triplets (x_i, x_{i+1}, x_{i+2}) (where $x_{p+i} = x_i$) of coordinates possess the same qualitative dynamics.

For each i, we will be interested in the points where $\dot{x}_i = 0$; each such set is the union of the plane $x_i = 0$ and some "nontrivial" plane. We designate those planes:

$$
P_1 := \{ \sigma_1 - \rho_{12} x_2 - \rho_{13} x_3 - x_1 = 0 \},\tag{12}
$$

$$
P_2 := \{ \sigma_2 - \rho_{21} x_1 - \rho_{23} x_3 - x_2 = 0 \},\tag{13}
$$

$$
P_3 := \{\sigma_3 - \rho_{31}x_1 - \rho_{32}x_2 - x_3 = 0\},\tag{14}
$$

where $\dot{x}_i = 0$ on P_i .

We will also refer to the plane passing through the points $(\sigma_1, 0, 0)$, $(0, \sigma_2, 0)$, and $(0, 0, \sigma_3)$, which we denote by Σ . Observe that Σ is given by the equation

$$
\Sigma := \frac{x_1}{\sigma_1} + \frac{x_2}{\sigma_2} + \frac{x_3}{\sigma_3} = 1.
$$

For ease of discussion, we will also name the coordinate planes:

$$
P_{12} := \{(x_1, x_2, x_3) : x_3 = 0\},
$$

\n
$$
P_{23} := \{(x_1, x_2, x_3) : x_1 = 0\},
$$

\n
$$
P_{13} := \{(x_1, x_2, x_3) : x_2 = 0\}.
$$

Because of the positivity conditions on the parameters and variables, we may use P_1 , P_2 , P_3 , and Σ as shorthand for the intersection of those planes with the first octant without the risk of confusion. We observe that the intersection of Σ and of each P_i with $\overline{\mathbb{R}^n_+}$ is a compact triangle, a fact we will use repeatedly in the following section.

In order to describe the dynamics of orbits, we are interested in when one plane lies "above" another in the first octant. Consider the following definition:

Definition 3.1. Observe that each plane P_i can be written as the graph of a function $z^{i}(x_1, x_2)$. A plane P_i dominates a plane P_j if $(x_1, x_2, z^{i}(x_1, x_2)) \in P_i$ and $(x_1, x_2, z^{j}(x_1, x_2)) \in P_i$ P_j implies that $z^j(x_1, x_2) < z^i(x_1, x_2)$ for all $x_1, x_2 > 0$. A plane P_i is dominated by a plane P_j if P_j dominates P_i .

3.2 Geometrical Lemmas

Each plane P_i divides $\overline{\mathbb{R}^3_+}$ into two regions, one where \dot{x}_i is positive and another where it is negative. This allows information about \dot{x}_i to be gained from purely geometric information. For example, \dot{x}_1 is positive below P_1 , and negative above it. Since P_3 dominates P_1 (i.e. is always above it), we instantly see that $\dot{x}_1|_{P_3} < 0$ (see Corollary [3.5,](#page-7-0) given below). We introduce geometric lemmas giving the information we can gain in this way; the proofs of all of them are parallel to one another, and can be summarized as follows: since the system is restricted to the first octant, we compare two planes (triangles) by seeing where they intersect the x_1, x_2 , and x_3 axes. Denote the compact triangle thus formed by a plane S as T_S and the non-zero component of its vertex on the *i*-th axis as S^i . Then a plane S dominates a plane R if $R^i \leq S^i$ for $i = 1, 2, 3$, with at least one of those a strict equality, i.e. if its vertices are farther from the origin.

Lemma 3.2. The plane P_1 is dominated by the plane Σ .

Proof. We consider where each plane intersects each axis:

- The planes P_1 and Σ both intersect the x_1 −axis at the point O_1 .
- The plane P_1 intersects the x_2 axis at $(0, \frac{\sigma_1}{\sigma_1})$ $\frac{\sigma_1}{\rho_{12}}$, 0), while Σ intersects the axis at O_2 . We know from [\(9\)](#page-4-3) that $\frac{\sigma_1}{\rho_{12}} < \sigma_2$.
- The plane P_1 intersects the x_3 -axis at $(0,0,\frac{\sigma_1}{\sigma_1})$ $\frac{\sigma_1}{\rho_{13}}$, while Σ intersects the axis at O_3 . We know from [\(10\)](#page-4-4), that $\frac{\sigma_1}{\rho_{13}} < \sigma_3$.

 \Box

Since $P_1^i \leq \Sigma^i$ for all *i*, P_1 is dominated by Σ .

Lemma 3.3. The plane Σ is dominated by P_3 .

Proof. We consider where each plane intersects each axis:

- The plane P_3 intersects the x_1 -axis at the point $(\frac{\sigma_3}{\rho_{31}}, 0, 0)$, while Σ intersects the axis at O_1 . We know from [\(7\)](#page-4-1) that $\sigma_1 < \frac{\sigma_3}{\sigma_1}$ $\frac{88}{\rho_{31}}$.
- The plane P_3 intersects the x_2 axis at $(0, \frac{\sigma_3}{\sigma_3})$ $\frac{\sigma_3}{\rho_{32}}$, 0), while Σ intersects the axis at O_2 . We know from [\(8\)](#page-4-2) that $\sigma_2 < \frac{\sigma_3}{\sigma_3}$ ρ_{32} .
- The planes P_3 and Σ both intersect the x_3 axis at O_3 .

Since $\Sigma^i \leq P_3^i$ for all *i*, P_3 dominates Σ .

The property "is dominated by" is clearly transitive, so the following corollary holds.

Corollary 3.4. P_1 is dominated by P_3 .

Since $\dot{x}_1 < 0$ above the P_1 plane, the following corollary follows immediately.

Corollary 3.5. On the plane P_3 , $\dot{x}_1 < 0$.

If we further had that Σ dominates P_2 , then the eigenvalue conditions introduced in [\[4\]](#page-24-10) and summarized as (3) - (8) would be sufficient to ensure the existence of a positively invariant region. It happens, however, that this is not the case.

Lemma 3.6. The plane Σ neither dominates nor is dominated by P_2 .

Proof. We consider where each plane intersects each axis:

- P_2 intersects the x_1 -axis at the point $(\frac{\sigma_2}{\rho_{21}}, 0, 0)$, while Σ intersects the axis at O_1 . We know from [\(7\)](#page-4-1) that $\sigma_1 < \frac{\sigma_2}{\sigma_2}$ $\frac{\sigma_2}{\rho_{21}}$, and therefore Σ does not dominate P_2 .
- P_2 intersects the x_3 -axis at $(0,0,\frac{\sigma_2}{\sigma_2})$ $\frac{\sigma_2}{\rho_{23}}$, while Σ intersects the axis at O_3 . We know from [\(11\)](#page-4-5) that $\frac{\sigma_2}{\rho_{23}} < \sigma_3$, and therefore P_2 does not dominate Σ .

Thus, neither plane dominates the other.

$$
\Box
$$

The situation is somewhat salvaged by the following.

Lemma 3.7. If $\frac{\sigma_2}{\rho_{21}} \leq \frac{\sigma_3}{\rho_{31}}$ $\frac{\sigma_3}{\rho_{31}}$, then P_2 is dominated by P_3 .

Proof. In the proof of Lemma [3.3](#page-6-0) (second bullet point), we established that $P_2^2 \n\t\le P_3^2$. In the proof of Lemma [3.6](#page-7-1) (second bullet point), we established that $P_2^3 \n\t\le P_3^3$. All that remains for P_3 to dominate P_2 is for $P_2^1 \le P_3^1$, which occurs if and only if $\frac{\sigma_2}{\rho_{21}} \le \frac{\sigma_3}{\rho_{31}}$ $\frac{\sigma_3}{\rho_{31}}.$

The parameter restriction in [\(3.7\)](#page-7-2) written in terms of the general systems gives that for each $k, 1 \leq k \leq p$,

$$
\frac{\sigma_{k+1}}{\rho_{k+1,k}} \le \frac{\sigma_{k+2}}{\rho_{k+2,k}} \quad \text{(indices mod } p). \tag{15}
$$

Similarly to Corollary [3.5,](#page-7-0) we have the following:

 \Box

Corollary 3.8. In the region of parameter space where the hypotheses of Lemma [3.7](#page-7-2) are satisfied, $\dot{x}_2|_{P_3} < 0$.

We note here one additional observation.

Lemma 3.9. The rectangular box:

$$
B = \{x : 0 \le x_i \le \sigma_i, i = 1, \dots, n\}
$$

is forward invariant with respect to the system [\(1\)](#page-1-0). Further, the unstable manifolds \widetilde{W}_{k}^{u} are all contained in B.

Forward invariance follows immediately from the differential equations [\(1\)](#page-1-0) and the assumption that $\rho_{ii} = 1$. The conclusion that the unstable manifold at O_1 is inside B follows easily by noting that the unstable eigenspace at O_1 (restricted to the first orthant) is strictly inside B.

Figure 3: The plane P_1 (dotted) is dominated by Σ (solid) which is dominated by P_3 (dashes). If the hypotheses of Lemma [3.7](#page-7-2) are satisfied, then the plane P_2 (dashes and dots) is dominated by P_3 .

Rather than requiring that P_3 dominates P_2 , we could require only that P_3 dominates P_2 inside the forward invariant box B. Since P_3^1 is strictly outside of B by Lemma [3.3](#page-6-0) (see Figure [3.2\)](#page-8-2), this produces a strictly greater set of allowable parameter values.

3.3 Existence of a Positively Invariant Region

We remarked that our goal in three dimensions was to show the existence of a compact, forward-invariant set whose trajectories converge to O_3 . We now carry this out.

Lemma 3.10. In the region of parameter space where the inequalities (7) - (11) and the hypotheses of Lemma [3.7](#page-7-2) are satisfied, the planes P_3 , P_{23} , P_{13} , and P_{12} enclose a positively invariant region, i.e. no trajectory leaves in positive time.

Proof. No trajectory can leave through any of the $x_i = 0$ planes, since $\dot{x}_i |_{x_i \text{-plane}} = 0$. The outward normal vector to P_3 is $N = \langle \rho_{31}, \rho_{32}, 1 \rangle$, and its scalar product with the vector field [\(1\)](#page-1-0) is given by $\bar{n} \cdot F = \rho_{31} \dot{x}_1 + \rho_{32} \dot{x}_2$ when $\dot{x}_3 = 0$. This is negative by Corollary [3.5](#page-7-0) and Corollary [3.8.](#page-8-3) \Box

Remark 3.11. Since the pair of inequalities $\rho_{31} \dot{x}_1 < 0$ and $\rho_{32} \dot{x}_2 < 0$ form a sufficient, but not necessary, condition for the inequality $\rho_{31}\dot{x}_1 + \rho_{32}\dot{x}_2 < 0$ to be satisfied, a positively invariant region might exist even if the hypothesis of Lemma [3.7](#page-7-2) is not satisfied. For instance, since the box B (Lemma [3.9\)](#page-8-2) is forward invariant in may be that $\dot{x}_2|_{P_3\cap B}$ is negative even when the hypotheses of Lemma [3.7](#page-7-2) fail. Lemma [3.10](#page-8-0) can therefore be extended to a region of parameter space that includes the region defined by Lemma [3.7](#page-7-2) as a proper subset.

Theorem 3.12. Suppose that P_3 , $x_1 = 0$, $x_2 = 0$, and $x_3 = 0$ enclose a positively invariant region. Any trajectory $\phi^t(x)$ in the above-described region that is not contained in P_{12} (the $x_3 = 0$ plane) goes to $(0, 0, \sigma_3)$ as $t \to +\infty$.

Proof. Any trajectory in a compact positively invariant region has a non-empty ω -limit set. Fix a trajectory with initial condition x in the interior of the region, and let q be an arbitrary point in the ω -limit set Ω_x . The proof breaks into three parts: we prove that q lies on P_3 , that it lies on P_{23} , and that it lies on P_{13} ; the intersection of these planes is $(0, 0, \sigma_3)$. The proofs of the second and third statements are essentially the same as the proof of the first statement, which we treat in detail.

By way of contradiction, suppose that q does not lie on P_3 . First of all, note that q does not lie on P_{12} , because by assumption, the $x \notin P_{12}$, and since the trajectory increases in the x_3 variable, $\phi^t(x)$ cannot approach $P_{12} = \{x_3 = 0\}$

Since \dot{x}_3 is continuous, and q lies on neither P_3 nor P_{12} , the regions where $\dot{x}_3 = 0$, we can find a spherical neighborhood $O_r(q)$ centered at q with radius r such that $\dot{x}_3|_{O_r(q)} > \epsilon > 0$ for some ϵ . It follows that q is not a fixed point, and Ω_x is not the singleton $\{q\}$, since ω –limit sets are forward invariant [\[21\]](#page-25-9).

The x_3 component of $\phi^t(x)$ is non-decreasing in the positively invariant set and it is bounded above, since it is bounded above by σ_3 . Thus it has a limit x_3^* . It follows by continuity that the x_3 component of any point in Ω_x is also x_3^* . We have supposed that $q \in \Omega_x$ but $q \notin P_3$; now consider $\phi^t(q)$. Since $\dot{x}_3 > 0$ at q, and thus in a neighborhood of q, the x_3 component of $\phi^t(q)$ must be strictly increasing as a function of time along the forward solution near q. This contradicts that the x_3 component is x_3^* everwhere on Ω_x .

We repeat the argument twice. First, q must lie on the plane P_{23} ; otherwise $\dot{x}_1(q) < 0$, and the same contradiction will be obtained. Then, using the same argument, we see that it must lie on P_{13} . We observe here, with reference to Lemma [3.7,](#page-7-2) that even if P_2 is not dominated by P_3 , it is dominated by it on the restriction to P_{23} , and the argument therefore goes through. Thus q must lie on the single point, $(0, 0, \sigma_3)$, and our proof is complete. \Box

4 The Dynamics in the Full Phase Space

4.1 Local Dynamics Near an Equilibrium O_k

We assumed in [\(5\)](#page-3-2) that the saddle points O_k are dissipative. Denote by ν the ratio:

$$
\nu \equiv \frac{\min |\{Re(\Sigma^s)\}|}{\max \{Re(\Sigma^u)\}}.
$$
\n(16)

We call ν the *minimal saddle value* of the equilibrium (see [\[27\]](#page-25-10)). The equilibrium is dissipative if $\nu > 1$.

The main implication of this assumption is that a trajectory starting at an initial value at a distance ϵ from the stable manifold of the saddle comes to a point of distance on the order of ϵ^{ν} from the unstable manifold after going through a neighborhood of the saddle. Formulating this result more strictly, we label the variables in a neighborhood of O_k into the 2-dimensional unstable subspace $\eta = (x_{k+1}, x_{k+2})$ and the $(n-2)$ -dimensional stable subspace $\xi = (x_j), j \neq k+1, k+2$. Let $|\cdot|$ denote the sup norm in these local coordinates. By the Stable Manifold Theorem, for each k there exists $\delta_k > 0$ such that in a δ_k -neighborhood of O_k , the unstable manifold $W^u(O_k)$ is the graph of a (smooth) function $\xi = h_k^u(\eta)$. Let δ be the minimum of these δ_k and consider the δ -neighborhood V_k of each O_k , $k = 1, \ldots, p$

For fixed k, and $0 < \epsilon_k$ sufficiently small, define a pair of sections,

$$
S_0 = \{ (\xi, \eta) : |\xi| = \delta, |\eta| \le \epsilon_k \} \quad \text{and} \quad S_1 = \{ (\xi, \eta) : |\xi| \le \delta, |\eta| = \delta \}.
$$

By the classical Shil'nikov variables technique (see [\[27\]](#page-25-10)), there exists ϵ_k sufficiently small so that every forward solution starting at S_0 will intersect S_1 before leaving V_k . Let ϵ be the minimum of the ϵ_k 's needed in the neighborhood of each O_k . Let T denote the time at which such a solution intersects S_1 .

Theorem 4.1. If δ and ϵ are sufficiently small and if $(\xi(0), \eta(0)) \in S_0$ and $(\xi(T), \eta(T)) \in$ S_1 then $|\xi(T)| \le C |\eta(0)|^{\nu-e}$, where $C > 0$ is independent of the initial point and $\nu - e > 1$.

The sections S_0 and S_1 , together with Theorem [4.1,](#page-10-1) are illustrated in two dimensions in Figure [4.1](#page-10-1)

Proof. For ease of notation, consider O_1 . Note from [\(2\)](#page-2-0) that the eigenvector corresponding to the first diagonal element, $-\sigma_1$, can have only one non-zero component and that is in the x_1 direction. The eigenvector corresponding to the j-th diagonal element, $j \neq 1$, has non-zero components in the x_1 and x_j directions only.

Now take into account that the hyperplane $\{(x_2 = x_3 = 0)\}\$ is invariant under the flow of the equations and tangent to the stable eigenspace of $DF(O_1)$. It thus coincides locally with and contains $W^s(O_1)$.

Next note that the unstable eigenspace has no non-zero components in the coordinate directions x_4, x_5, \ldots, x_n and that the hyperplane where they are zero is invariant under

Figure 4: A schematic diagram of Theorem [4.1.](#page-10-1) Initial conditions on a section intersecting a stable manifold are mapped in finite time to a section intersecting an unstable manifold, resulting in a contraction.

the flow. It thus follows that the x_4, x_5, \ldots, x_n coordinates of the unstable manifold are all zero. Thus the unstable manifold $W^u(O_1)$ is given locally in the δ -neighborhood of O_1 as the graph $x_1 = \sigma_1 + h_1^u(x_2, x_3)$ where $h(0,0) = 0$ and h is smooth. Consider the local change of variables that "straightens out" the unstable manifold:

$$
X_1 = x_1 - \sigma_1 - h(x_2, x_3). \tag{17}
$$

Under this smooth change of coordinates [\(1\)](#page-1-0) becomes:

$$
\begin{aligned}\n\dot{X}_1 &= -\sigma_1 X_1 - f(X_1, x_2, \dots, x_n) \\
\dot{x}_2 &= x_2 (\sigma_2 - x_2 - \rho_{21} (X_1 + \sigma_1 + h(x_2, x_3)) - \rho_{23} x_3 - \rho_{24} x_4 - \dots - \rho_{2n} x_n) \\
\dot{x}_3 &= x_3 (\sigma_3 - x_3 - \rho_{31} (X_1 + \sigma_1 + h(x_2, x_3)) - \rho_{32} x_2 - \rho_{34} x_4 - \dots - \rho_{3n} x_n) \\
\dot{x}_4 &= x_4 (\sigma_4 - x_4 - \rho_{41} (X_1 + \sigma_1 + h(x_2, x_3)) - \rho_{42} x_2 - \rho_{43} x_3 - \dots - \rho_{4n} x_n) \\
&\vdots \\
\dot{x}_n &= x_n (\sigma_n - x_n - \rho_{n1} (X_1 + \sigma_1 + h(x_2, x_3)) - \rho_{n2} x_2 - \rho_{n3} x_3 - \dots - \rho_{n,n-1} x_{n-1}).\n\end{aligned} \tag{18}
$$

Since $W^s(O₁)$ is contained in the coordinate hyperplane, we do not need a change of variables corresponding to W^s . Noting the $f(0, x_2, \ldots, x_n) \equiv 0$, we use the MVT to define a new function $\frac{f(X_1,x_2,...,x_n)-0}{X_1-0} = f_1(X_1,x_2,...,x_n)$, and rewrite the equations to obtain:

$$
\begin{aligned}\n\dot{X}_1 &= -\sigma_1 X_1 + f_1(X_1, x_2, \dots, x_n) X_1 \\
\dot{x}_2 &= x_2 \left(\sigma_2 - \rho_{21} \sigma_1 - x_2 - \rho_{21} (X_1 + h(x_2, x_3)) - \rho_{23} x_3 - \rho_{24} x_4 - \dots - \rho_{2n} x_n \right) \\
\dot{x}_3 &= x_3 \left(\sigma_3 - \rho_{31} \sigma_1 - x_3 - \rho_{31} (X_1 + h(x_2, x_3)) - \rho_{32} x_2 - \rho_{34} x_4 - \dots - \rho_{3n} x_n \right) \\
\dot{x}_4 &= x_4 \left(\sigma_4 - \rho_{41} \sigma_1 - x_4 - \rho_{41} (X_1 + h(x_2, x_3)) - \rho_{42} x_2 - \rho_{43} x_3 - \dots - \rho_{4n} x_n \right) \\
&\vdots = \vdots\n\end{aligned}\n\tag{19}
$$

Now note that distances from $W^s(O_1)$ and $W^u(O_1)$ are not effected by the coordinate change [\(17\)](#page-11-0). If we now denote $\xi = (X_1, x_4, \ldots, x_n)$ then $|\xi|$ is the distance to the unstable manifold and $|\eta|$ is the distance to the stable manifold (in the sup norm).

It now follows immediately from [\(19\)](#page-11-1) that inside the δ neighborhood of O_1 the unstable directions satisfy the estimates:

$$
\dot{x}_2 \le x_2(\sigma_2 - \rho_{21}\sigma_1 - e^u)
$$
 and $\dot{x}_3 \le x_3(\sigma_3 - \rho_{31}\sigma_1 - e^u)$,

where we can take $e^u > 0$ arbitrarily small by starting with δ and ϵ small. Thus, by a simple application of Gronwall's inequality, solutions starting on S_0 and remaining in our δ neighborhood of O_1 must satisfy:

$$
|\eta(t)| \le |\eta(0)| e^{(\lambda^{uu} - e^u)t},
$$

where λ^{uu} is the maximal unstable eigenvalue, i.e.,

$$
\lambda^{uu} = \max_{j=2,3} \{ \sigma_j - \rho_{j1} \sigma_1 \}.
$$

Thus if T is defined by $|\eta(T)| = \delta$, then

$$
T \ge \frac{1}{\lambda^{uu} - e^u} \ln(\frac{\delta}{\eta(0)}).
$$

Similarly, using Gronwall's inequality once again we obtain:

$$
|\xi(t)| \le |\xi(0)|e^{(\lambda^{ls} + e^s)t},
$$

where λ^{ls} is the "leading" stable eigenvalue, i.e. since the eigenvalues are real, the negative eigenvalue with the smallest absolute value. In terms of our parameters:

$$
\lambda^{ls} = \max\{-\sigma_1, \{\sigma_j - \rho_{j1}\sigma_1, j = 4,\ldots, n\}\}.
$$

Thus

$$
|\xi(T)| \le C(\delta) |\eta(0)|^{(\nu - e)}
$$

where e can be taken arbitrarily small. By the assumption that O_1 is dissipative, $\nu > 1$ and so we can make $\nu - e > 1$.

 \Box

4.2 Stable Heteroclinic Surface

We are now in a position to prove that the union of the unstable manifolds of our system, restricted to the positive orthant, form an asymptotically stable forward invariant set under appropriate parameter restrictions.

Theorem (Theorem [2.3,](#page-3-3) restated). Suppose that inequalities (3) , (4) and (5) hold for each $k, 1 \leq k \leq p$ and that each unstable manifold $\widetilde{W}^u(Q_k)$ is contained in a compact forward invariant set as in Section [3.3.](#page-8-1) Then $\Gamma \equiv \bigcup_{k=1}^p (\widetilde{W}_k^u \cup O_k)$ is asymptotically stable.

We consider the role each inequality of the hypothesis plays in the theorem. The inequalities [\(3\)](#page-3-0) and [\(4\)](#page-3-1) state that

$$
0 < \min_{i=1,2} \{ \sigma_{k+i} - \rho_{k+i,k} \sigma_k \} \text{ (indices mod } p \text{), for each } 1 \le k \le p, \text{ and}
$$
\n
$$
\lambda_j = \sigma_j - \rho_{jk} \sigma_k < 0, \quad \text{ for } j \neq k, k+1, k+2 \mod p, \text{ for each } 1 \le k \le p.
$$

These inequalities are fundamental to the problem we are considering. They ensure that there is a heteroclinic channel from each equilibrium O_k to the equilibria O_{k+1} and O_{k+2} (the first inequality), and that there are no heteroclinic channels to the other equilibria (the second inequality).

The inequality (5) states that for each k,

$$
\max_{i=1,2} \{\sigma_{k+i} - \rho_{k+i,k}\sigma_k\} < \min_{j \neq k,k+1,k+2} \{|\sigma_j - \rho_{jk}\sigma_k|, \sigma_k\} \quad \text{(indices mod } p\text{)}.
$$

This is the dissipativity condition. Informally, it may be taken to say that when a trajectory that is close to one of the unstable manifolds whose union is Γ passes near one of the saddle fixed points, the distance of the trajectory to Γ contracts exponentially.

It will become clear, from the proof of Theorem [2.3,](#page-3-3) that [\(5\)](#page-3-2) is a much stronger condition than is necessary. It ensures not only asymptotic stability, but a sort of monotonic asymptotic stability, such that whenever the trajectory passes near some O_k , its distance to Γ contracts exponentially. If, on the other hand, [\(5\)](#page-3-2) held for some, but not all, values of k, then the trajectory would at times contract exponentially towards Γ , and at other times drift away from Γ , and stability would depend on how these attractive and repulsive forces average over time. Attempting to formulate a replacement condition for [\(5\)](#page-3-2) that is necessary as well as sufficient is extremely nontrivial.

The final hypothesis, that each unstable manifold is contained in a compact forward invariant set, is necessary. We have seen one set of inequalities that ensure that such sets exist,

$$
\frac{\sigma_{k+1}}{\rho_{k+1,k}} \le \frac{\sigma_{k+2}}{\rho_{k+2,k}} \quad \text{(indices mod } p),
$$

which are sufficient but not necessary. In Section [4.3,](#page-15-0) we see that there is a nonempty open region of parameter space where the hypotheses of Theorem [2.3](#page-3-3) hold. From our current discussion, we see that the theorem applies to a larger region.

We recall that Γ is asymptotically stable if given any neighborhood U of Γ (restricted to $\overline{\mathbb{R}^n_+}$) there exists an ϵ -neighborhood of Γ , say $V_{\epsilon}(\Gamma) \subset \overline{\mathbb{R}^n_+}$, such that if $x_0 \in V_{\epsilon}(\Gamma)$ then $x(t, x_0) \in U$ for $t > 0$ and $\lim_{t \to \infty} \text{dist}(x(t, x_0), \Gamma) = 0$, where $x(t, x_0)$ is the solution of the initial value problem [\(1\)](#page-1-0) with $x(0, x_0) = x_0$. When we speak of an open ϵ -neighborhood of Γ , we are speaking of a set that is open in the subspace topology. That is, an ϵ -neighborhood in $\overline{\mathbb{R}^n_+}$ is an ϵ -neighborhood in \mathbb{R}^n intersected with $\overline{\mathbb{R}^n_+}$.

Proof of Theorem [2.3.](#page-3-3) For each $k = 1, ..., p$, let $V(O_k)$ be a sufficiently small δ neighborhood of O_k , such that Theorem [4.1](#page-10-1) can be applied within each $V(O_k)$ and δ does not depend on k .

Let z_0 be a representative point at an initial condition ϵ -close to Γ. Choose ϵ such that $\epsilon < \delta$. Then we can classify the dynamics as either local if $z_0 \in V(O_k)$ for some k, and global otherwise.

We observe that if $z_0 = O_k$ for any k, its behavior is trivial, and likewise, if z_0 lies on a coordinate plane, it remains on that plane while converging exponentially to some O_k . We therefore assume without loss of generality that neither of these cases hold.

Suppose that $z_0 \notin V(O_k)$ for any k. The point is ϵ -close to Γ, and since Γ is a finite union, we can say that z_0 is ϵ -close to \widetilde{W}^u_α , where α is fixed and depends on z_0 . Consider the projection of the system onto the three-dimensional subspace spanned by the axes x_{α} , $x_{\alpha+1}$, and $x_{\alpha+2}$. In three dimensions, the specific route a solution takes has not been important; a trajectory in the invariant set may go straight to a neighborhood of $O_{\alpha+2}$, or it may detour to $O_{\alpha+1}$, but the net result is the same (Theorem [3.12\)](#page-9-0). We now formally differentiate between these two cases.

We consider two cases: either the positive semitrajectory of z_0 intersects $V(O_{\alpha+2})$ without first intersecting $V(O_{\alpha+1})$ (case (i)), or the positive semitrajectory intersects $V(O_{\alpha+1})$, then intersects $V(O_{\alpha+2})$ (case (ii)).

Before proceeding, we recall the definitions of S_0 and S_1 given in Theorem [4.1,](#page-10-1) and similarly define such sections S_0^q g_0^q , S_1^q for $1 \le q \le p$. Without loss of generality, we assume that $z_0 \in S_1^1$.

Suppose that case (i) occurs. For each $k = 1, ..., p$, let $\hat{V}(O_k)$ be the projection of $V(O_k)$ into the three dimensions spanned by x_k, x_{k+1} , and x_{k+2} ; note that x_k is negligable for $k \neq \alpha, \alpha+1$, and $\alpha+2$. Then the projection of the orbit onto \mathbb{R}^3 intersects $\hat{V}(O_{\alpha+2})$ before it can intersect $\hat{V}(O_{\alpha+1})$. In \mathbb{R}^3 , we know that all trajectories inside of \widetilde{W}^u_α that do not intersect $\hat{V}(P_{\alpha+1})$ come to a neighborhood of $O_{\alpha+2}$ in bounded time, where the bound does not depend on the initial condition. We may consider the non-projected, full-dimensional space as a "perturbation" of the projected space, and cite smooth dependence of initial conditions; the trajectory going through a slightly perturbed initial point corresponding to such a case must enter $V(O_{\alpha+2})$ in a well-behaved way. In particular if z_0 belongs to S_0^{α} then a mapping from a neighborhood of z_0 on S_0^{α} to $S_0^{\alpha+2}$ is well defined and Lipschitzcontinuous.

Suppose that case (ii) occurs. Then once an orbit of z_0 enters $V(O_{\alpha+1})$, it starts to manifest the dissipative behavior. In particular, if $d(x, W^u_\alpha) < \epsilon$, then after passing through $V(O_{\alpha+1}),$ $d(\widetilde{x}, W_{\alpha+1}^u) < C\epsilon^{\nu}$, by Theorem [4.1.](#page-10-1) Once the representative point leaves $V(O_{\alpha+1})$, we may apply (i), viewing its position after leaving the neighborhood as an initial condition that does not re-enter the neighborhood $V(O_{\alpha+1})$. Thus when the representative point finally enters $V(O_{\alpha+2})$, its distance to Γ has been contracted by an order of $Ce^{\nu_{\alpha+1}-e_{\alpha+1}}$, where $1 < \nu_{\alpha+1} - e_{\alpha+1}$ and C absorbs both the constant C from Theorem [4.1](#page-10-1) and a Lipschitz constant.

Suppose now that $z_0 \in V(O_k)$, where k is now fixed. Then the trajectory leaves $V(O_k)$ without increasing its distance from the unstable manifold, and passes into $V(O_{k+1})$ as just described. We may then apply Theorem [4.1.](#page-10-1) As the trajectory passes through $V(O_k)$, its distance from the unstable manifold is contracted on an order of $\epsilon^{\nu_k-e_k}$.

Since the mapping contracts in the global dynamics and is Lipschitz (or contracting) in the local dynamics, simple inductions yields that as a representative point moves through the system, its distance from the manifold changes from ϵ to $c_1 \epsilon^{\nu_1 - \epsilon_1}$ to $c_2 \epsilon^{(\nu_1 - \epsilon_1)(\nu_2 - \epsilon_2)}$, and so on.

For a fixed *i*, the value ν_i , representing a ratio of eigenvalues, is likewise fixed. The value e_i is not; it depends on the distance between the representative point and the stable manifold as the trajectory enters $V(O_i)$, which changes from one instance to the next. For a given i, however, there is some maximal value that e_i can take, since the system is constantly contracting towards the manifold and e_i goes to 0 along with that distance. Thus there exists a global value, $1 < \nu < \nu_i - e_i$ for all i and all e_i , such that passing from the first to the *p*-th unstable manifold is a contraction of order ce^{v^p} . □

4.3 Existence of Parameter Sets

Throughout the paper, we have put a number of restrictions on the parameters of the system. One must ask whether the specified inequalities may be satisfied.

Lemma 4.2. There are sets of positive parameters values $\{\sigma_i\}$, $i = 1, \ldots, n$, and $\{\rho_{ik}\}$, $j, k = 1, \ldots, n$, with non-empty interior such that the inequalities [\(3\)](#page-3-0), [\(4\)](#page-3-1), [\(5\)](#page-3-2) and [\(15\)](#page-7-3) hold.

Proof. First note that given $\{\sigma_i\}$ the inequalities [\(3\)](#page-3-0) and [\(4\)](#page-3-1) are completely uncoupled and all trivially have positive solutions $\{\rho_{jk}\}\$. For each k and $i = 1, 2$ these are:

$$
\rho_{jk} < \frac{\sigma_j}{\sigma_k} \tag{20}
$$

We note that with these inequalities the stable eigenvalues may be freely chosen to take any negative value and the unstable eigenvalues any positive values. Since the restrictions [\(5\)](#page-3-2) concern only relative orderings of those eigenvalues at each O_k (k fixed), it is clear that [\(5\)](#page-3-2) is satisfied for open subsets of the previously chosen sets.

The final remaining inequality [\(15\)](#page-7-3) concerns on $\rho_{k+1,k}$ and $\rho_{k+2,k}$ and so to finish the proof we need only to consider whether this restriction on those values is consistent with the previous restrictions on those parameters, namely [\(3\)](#page-3-0) and [\(5\)](#page-3-2), but not [\(4\)](#page-3-1). Note that the inequalities in [\(3\)](#page-3-0) can be written as:

$$
\frac{\sigma_{k+1}}{\rho_{k+1,k}} > \sigma_k \quad \text{ and } \quad \frac{\sigma_{k+2}}{\rho_{k+2,k}} > \sigma_k.
$$

The constraint [\(15\)](#page-7-3) only requires that

$$
\frac{\sigma_{k+1}}{\rho_{k+1,k}} \le \frac{\sigma_{k+2}}{\rho_{k+2,k}}.
$$

There is clearly no inconsistency in these inequalities. The final inequalities [\(5\)](#page-3-2) involve each $\rho_{k+1,k}$ and $\rho_{k+2,k}$ independently of the others. First they require that for each k and $i = 1, 2$

$$
\sigma_k - \rho_{k+1,k} \sigma_k < \sigma_k.
$$

This is equivalent to

$$
\frac{\sigma_{k+i}}{\sigma_k} - 1 < \rho_{k+1,k}
$$

which can clearly be satisfied along with [\(20\)](#page-15-1). If $\{\rho_{k+i,k}\}\$, $i=1,2$ have already been chosen to satisfy [\(3\)](#page-3-0) and [\(15\)](#page-7-3), then values for ρ_{jk} can be chosen to satisfy [\(5\)](#page-3-2) by simply choosing them sufficiently large, i.e., for each $j, k, j \neq k, k + 1, k + 2, \rho_{jk}$ must satisfy:

$$
\sigma_{k+1} - \rho_{k+i,k} \sigma_k < \rho_{jk} \sigma_k - \sigma_j, \quad i = 1, 2.
$$

This can be rewritten as:

$$
\rho_{jk} > \frac{\sigma_{k+i} + \sigma_j}{\sigma_k} - \rho_{k+i,k}, \quad i = 1, 2.
$$

Thus ρ_{ik} can take any value greater than the maximum of these two values. Previously we had only required (in (4)) that these parameters satisfy:

$$
\rho_{jk} > \frac{\sigma_j}{\sigma_k}.
$$

Thus, with all other choices of consistent parameters, any large enough ρ_{jk} , will also be consistent. □

For example if $\sigma_i = \sigma$ are all the same, then [\(3\)](#page-3-0), [\(4\)](#page-3-1) and [\(15\)](#page-7-3) are satisfied if simple :

$$
0 < \rho_{k+2,k} \le \rho_{k+1,k} < 1
$$

and

$$
\rho_{jk} > 1, \quad \text{for } j \neq k, k+1, k+2.
$$

The dissipative requirement [\(5\)](#page-3-2) will be satisfied if further:

$$
2\sigma - \rho_{k+i,k} < \rho_{jk}
$$
, for $i = 1, 2$ and $j \neq k, k + 1, k + 2$.

For instance, the parameter values: $\sigma_i = 1$, $\rho_{k+1,k} = .9$, $\rho_{k+2,k} = .8$ and $\rho_{jk} = 1.3$ for $j \neq k, k + 1, k + 2$, (indices mod p) strictly satisfy all of the inequalities.

5 Conclusion

We proved in the paper that under some conditions the generalized Lotka-Volterra system admits a two-dimensional attractor that consists of saddles and the unstable manifolds joining them into a heteroclinic system. Thus, trajectories inside the attractor manifest completely regular features. However, behavior of wandering trajectories in the basin of the attractor could be treated as weakly chaotic. Indeed, one can introduce an oriented graph with vertices identified with the saddle equilibrium points O_k and edges identified with heteroclinic trajectories joining O_k and O_{k+1} (belonging to the coordinate plane $P_{k,k+1}$) or O_k and O_{k+2} (belonging to the plane $P_{k,k+2}$). It is possible to show that for each finite path through this graph there exists an open set of initial points in the basin such that the trajectory going through any of these points follows the corresponding heteroclinic trajectories. The number of paths grows exponentially with the length, so the number of pieces of trajectories with different behavior (in fact all of them are (ϵ, T) separated for some values of ϵ and T) grows as $T \to \infty$, the metric complexity function grows with time. A similar effect has been observed in [\[1\]](#page-24-11), where it was called weak transient chaos. We intend to describe its properties in another publication.

In the context of the motivating application, functional sequential dynamics in neural networks, the two-dimensional heteroclinic attractor Γ in the phase space of dynamical system [\(1\)](#page-1-0) may be thought of as a mathematical image of diverse sequential dynamics based on the parallel performance of not one but two different modalities. Many interesting applications of this may be found in cognitive science. For example, the learning and performing of sensory-motor human behaviors in many situations require the integration or binding of the sequential stimuli of one modality with the sequential stimuli of another. This seems to be the case in one of the most important cognitive functions: sequential working memory. In performance of a cross-modal working memory task, there may be two sequentially discrete neural processes (different chains of metastable states) that represent simultaneous neural activities corresponding to cross-modal transfer of information in the working memory (see e.g. $|20|$).

ACKNOWLEDGEMENT. V.A. was partially supported by the grant RNF 14-41-00044 of the Russian Science Foundation during his stay at Nizhny Novgorod University, and by a Glidden Professorship Award during his stay at Ohio University. The authors thank M. I. Rabinovich for bringing to their attention the problems considered in this manuscript. We thank the referees for their comments and insight.

A The Topological Form of Γ

Theorem [2.1](#page-3-4) states that Γ depends on the parity of p; this is because the number of connected components in the boundary of Γ depends on the parity of p.

Proposition A.1. If p is even, then the boundary $\partial \Gamma$ has two connected components. If p is odd, then $\partial \Gamma$ has one component.

Proof. If p is even, then one connected component of the boundary will include the trajectories connecting the saddles O_n where n is even, and another will include the trajectories connecting the saddles O_m , where m is odd.

If p is odd, then as in the previous case, the saddles O_n where n is even are contained in the same component, and the saddles O_n where n is odd are likewise contained in a single component. Furthermore, O_{p-1} and O_1 are contained in the same component, where $p-1$ is even, since $O_1 = O_{p+1}$. Thus all saddles O_n are contained in the same component. \Box

Figure [5](#page-18-0) will help visualize the situation.

Figure 5: A diagram representing $p = 6$ (left) and $p = 5$ (right). We clearly observe two distinct boundaries for $p = 6$, and one for $p = 5$.

It appears from the diagram that when p is even, Γ is a cylinder, and when p is odd, Γ is a Möbius strip. We will formalize that intuition. We first observe that the components whose union is the unstable manifold Γ are not literally triangles, as they are depicted in Figure [5,](#page-18-0) but rather curved surfaces. We cite a definition from algebraic topology (e.g. [\[8\]](#page-24-12)).

Definition A.2. Suppose X is a compact Hausdorff space. A curved triangle in X is a subspace A of X and a homeomorphism $H : T \rightarrow A$ where T is a closed triangular region in the plane. A triangulation of X is a collection of curved triangles $A_1, ..., A_n$ in X whose union is X and such that for $i \neq j$, the intersection $A_i \cap A_j$ is either empty, or a vertex of both A_i and A_j , or an edge of both. We also require, if h_i is the homeomorphism associated with A_i , that when $A_i \cap A_j = e$ is an edge of both, then the map $h_j^{-1}h_i$ is a linear homeomorphism of the edge $h_i^{-1}(e)$ of T_i with the edge $h_j^{-1}(e)$ of T_j .

All compact surfaces have a triangulation, but we will prove in particular that the decomposition of Γ into equilibria and the closure of their unstable manifolds, each restricted to the first orthant, forms a triangulation of Γ.

Definition A.3. Consider the system of differential equations [\(1\)](#page-1-0). For each α , we call the closure of $\widetilde{W}^u_\alpha \cup O_\alpha$ a heteroclinic triangle, T_α .

Note that:

$$
T_{\alpha} = \widetilde{W}_{\alpha}^{u} \cup O_{\alpha} \cup O_{\alpha+1} \cup O_{\alpha+2} \cup \Gamma_{\alpha+1,\alpha+2}
$$

where $\Gamma_{\alpha+1,\alpha+2}$ is the heteroclinic orbit from $O_{\alpha+1}$ to $O_{\alpha+2}$. This follows from the invariance of the $(\alpha, \alpha + 1, \alpha + 2)$ -plane and Theorem [3.12.](#page-9-0)

Theorem A.4. A heteroclinic triangle T_{α} is homeomorphic to a closed triangle in the plane.

Certainly the boundary of a heteroclinic triangle T_{α} , that is O_{α} , $O_{\alpha+1}$, and $O_{\alpha+2}$, and the smooth paths connecting them, is homeomorphic to the boundary of a closed triangle in the plane. Further, the interior of T_{α} is $\overline{W}_{\alpha}^{u}$, by the unstable manifold theorem and

Theorem [3.12,](#page-9-0) homeomorphic to an open disk in the plane (and so also to the interior of a triangular region). The only complication may arise from the behavior of the unstable manifold near the edges or vertices. For instance, the \widetilde{W}^u_α could be "folded" as it approaches the edge connecting $O_{\alpha+1}$ and $O_{\alpha+2}$ so that a neighborhood of a point on the edge is not locally homeomorphic to a point on an edge of a closed triangle in the plane.

Consider the dynamics projected onto three consecutive dimensions; for simplicity of notation, we will use the standard x, y, z coordinates, and label the saddle points on each axis O_x , O_y , and O_z , where there are heteroclinic connections $O_x \rightarrow O_y \rightarrow O_z$ and $O_x \rightarrow O_z$, which we denote as Γ_{xy} , Γ_{xz} , and Γ_{yz} . We denote the corresponding heteroclinic triangle by T_x .

Proposition A.5. Each orbit in $\widetilde{W}^u(O_x)$ is uniquely identified with an angle $0 \le \phi \le \pi$.

Proof. We define the usual δ -neighborhoods $N_{\delta}(O_x)$, $N_{\delta}(O_y)$, and $N_{\delta}(O_z)$. By the Unstable Manifold Theorem, we may choose δ small enough that $W^u(O_x)$ is the graph of a function of (y, z) in $N_{\delta}(O_k)$. Using the monotonicity of the z coordinate inside the invariant region, we may also choose δ small enough such that once a trajectory leaves $N_{\delta}(O_x)$ it cannot return to it. By the previous observations, the other coordinates (other than x, y , and z) of $\widetilde{W}^u(O_x)$ are all zero.

Now consider that each orbit in $\widetilde{W}^u(Q_x)$ has a unique intersection point w with the boundary of $N_{\delta}(O_x)$. Since $\widetilde{W}^u(O_x)$ is a graph over (y, z) consider the projection \bar{w} of w onto (y, z) -plane. Let ϕ denote the angle of the ray through the origin and \bar{w} with the z-axis. Note that ϕ lies between 0 and $\pi/2$. The extreme angles $\phi = 0$ and $\phi = \pi/2$ correspond to the heteroclinic orbits Γ_{xz} and Γ_{xy} respectively. \Box

Definition A.6. For a point $p \in \widetilde{W}^u(O_x) \cup O_x$ define $d(p)$ to be the distance from O_x to p along the orbit containing p. For a point $p \in \widetilde{W}^s(O_z) \cup O_z$ define $e(p)$ to be the distance from p to O_z along the orbit containing p.

Proposition A.7. The arc lengths d and e are finite and continuous where defined.

Proof. By basic existence theory the orbits are smooth and therefore arc length is locally well-defined on them.

Denote the solution with initial value p as $p(t)$. Recall from the Unstable Manifold Theorem that $p(t) \to O_x$ as $t \to -\infty$ and in fact

$$
|O_x - p(t)| \le Ce^{(\lambda^{lu} - e^u)t}, \quad \text{for } t \le 0,
$$
\n
$$
(21)
$$

where λ^{lu} is the leading unstable eigenvalue, i.e. in this case the minimum of

 $\sigma_2 - \rho_{21}\sigma_1$ and $\sigma_3 - \rho_{31}\sigma_1$,

and e^u may be chosen to be arbitrarily small.

Now consider the length of the orbit:

$$
d(p) = \int_{-\infty}^{0} \sqrt{\dot{x}^2 + \dot{y}^2 + \dot{z}^2} dt
$$

=
$$
\int_{\tau}^{0} \sqrt{\dot{x}^2 + \dot{y}^2 + \dot{z}^2} dt + \int_{-\infty}^{\tau} \sqrt{\dot{x}^2 + \dot{y}^2 + \dot{z}^2} dt
$$
 (22)

If we substitute the equations [\(6\)](#page-4-6) into the integrals, then substitute [\(21\)](#page-19-0) into the second integral one easily sees that this integral goes to zero as $\tau \to -\infty$. Thus $d(p)$ is finite.

Now consider the orbits starting at points q close to p. Fix $\epsilon > 0$ and chose τ so that the remainder integral above is less than $\epsilon/3$. Since [\(21\)](#page-19-0) implies that $|q(t) - p(t)|$ goes to zero exponentially as $t \to -\infty$, we can make the difference in the remainders less than $2\epsilon/3$ if $q(\tau)$ is sufficiently close to $p(\tau)$. This we can accomplish by requiring $q(0)$ and $p(0)$ sufficiently close (by continuous dependence on initial conditions). We can also make the integrals from τ to 0 less than $\epsilon/3$ by choosing $q(0)$ close to $p(0)$. For such q , $|d(q)-d(p)| < \epsilon$.

The arc length $e(p)$ is finite for any p not in $O_x \cup \Gamma_{xy} \cup O_y$ since O_z is a stable node in the xyz-subspace and all orbits approach O_z exponentially in time and so the same type estimates as above hold here. Continuity of this distance also follows by a similar proof as for d . \Box

Corollary A.8. All orbits on T_x have finite length and this length is a continuous function of an initial point on $\widetilde{W}^u(Q_x) \setminus (\Gamma_{xy} \cup Q_y \cup \Gamma_{yz})$

This follows from the continuity of d and e where they are defined.

Lemma A.9. The arc length d can be extended continuously to the edge Γ_{yz} . The arc length is finite and uniformly bounded for all orbits that make up T_x .

Proof. Denote by D_{ϕ} the length of the orbit in $\widetilde{W}^u(O_x)$ identified by the angle ϕ for $0 \leq \phi < \pi/2$.

Note that distance is well defined along the edges Γ_{xy} and Γ_{yz} . In fact since $\Gamma_{xy} \subset W^u(O_x) \cap$ $\widetilde{W}^s(O_y)$, and $\Gamma_{yz} \subset \widetilde{W}^u(O_y) \cap \widetilde{W}^s(O_z)$ the arguments above show that the lengths of these two solutions arcs are finite. Let D_{xy} denote the length of Γ_{xy} and D_{yz} the length of Γ_{yz} . Set

$$
D_{\pi/2} = D_{xy} + D_{yz},
$$

i.e. the combined length of Γ_{xy} and Γ_{yz} .

For $p \in \Gamma_{yz}$ set:

$$
d(p) = D_{\pi/2} - e(p),
$$

i.e. the length to p along Γ_{xy} and Γ_{yz} . We claim that d thus defined is continuous at $p \in \Gamma_{yz}.$

Fix $\epsilon > 0$. Consider a δ (sup norm) neighborhood N of O_y . For any ϕ sufficiently close to $\pi/2$, the orbit on T_x indexed by ϕ passes through N. Denote by P^- the point where Γ_{xy}

intersects N and by P^+ the point where Γ_{yz} intersects N. By continuity of d and e we may chose such a δ sufficiently small that:

$$
D_{xy} - d(P^-) < \epsilon/6 \quad \text{ and } \quad D_{yz} - e(P^+) < \epsilon/6.
$$

Further, require that δ be sufficiently small so that: $\delta \left(\frac{2}{\lambda^{ls} + e^s} + \frac{1}{\lambda^{u} + e^u} \right) < \epsilon/6$.

Given ϕ sufficiently close to $\pi/2$, denote by Q^- and Q^+ the points where the orbit with angle ϕ intersects N. Again by continuity of d and e for all ϕ sufficiently close to $\pi/2$ we have:

$$
|d(Q^-) - d(P^-)| < \epsilon/6
$$
 and $e(Q^+) - e(P^+) < \epsilon/6$.

Let q be a point sufficiently close to p so that the above conditions on ϕ are satisfied and such that $|e(q) - e(p)| < \epsilon/6$.

Now $d(q)$ will be equal $d(Q^-)$ plus the arc length from Q^- to Q^+ plus the arc length from Q^+ to q. Denote these arc lengths by $\ell(Q^-, Q^+)$ and $\ell(Q^+, q)$ respectively.

Within the neighborhood N we can transform coordinates to straighten the unstable manifold Γ_{yz} . In this subspace this unstable manifold is the graph of a function of z only and has that has the form:

$$
y = \sigma_y + h(z).
$$

We thus straighten $\widetilde{W}^u(O_y)$ by the coordinate change $Y = y - \sigma_y + h(z)$.

By the Shilnikov variables technique [\[27\]](#page-25-10), we have that the stable variables satisfy:

$$
|(x(t), Y(t))| < \delta e^{(\lambda^{ls} + e^s)t}, \quad 0 \le t \le \tau,
$$

while the unstable variable z satisfies:

$$
|z(t)| < \delta e^{(\lambda^u - e^u)(t-\tau)}, \quad 0 \le t \le \tau,
$$

where τ is the time of passage through N. We then obtain:

$$
\ell(Q^{-}, Q^{+}) = \int_{0}^{\tau} \sqrt{\dot{x}^{2} + \dot{y}^{2} + \dot{z}^{2}} dt
$$

$$
< \int_{0}^{\tau} |\dot{x}| + |\dot{y}| + |\dot{z}| dt
$$

$$
< \delta \int_{0}^{\tau} 2e^{(\lambda^{ls} + e^{s})t} + e^{(\lambda^{u} + e^{u})(t-\tau)}
$$

$$
< \delta \frac{2}{\lambda^{ls} + e^{s}} + \frac{1}{\lambda^{u} + e^{u}}
$$

$$
< \frac{\epsilon}{6}.
$$
 (23)

We have that

$$
d(q) - d(p) = d(Q^-) + \ell(Q^-, Q^+) + \ell(Q^+, q) - (D_{xy} + D_{yz} - e(p))
$$

= $d(Q^-) + \ell(Q^-, Q^+) + e(Q^+) - e(q)$
 $- D_{xy} + d(P^-) - d(P^-) - D_{yz} + e(P^+) - e(P^+) + e(p)$ (24)
= $d(Q^-) - d(P^-) + \ell(Q^-, Q^+) + e(Q^+) - e(P^+) + d(P^-) - D_{xy} + e(p) - e(q).$

Combining the above estimates we have: $|d(q) - d(p)| < \epsilon$.

Now let D_{ϕ} , $0 \leq \phi \leq \pi/2$ be the arc length of the orbit determined by the angle ϕ . By the continuity of d on all of T_x , except O_x and the continuity of e in a neighborhood of O_z , D_ϕ depends continuously on ϕ . It is thus uniformly bounded. \Box

Proof of Theorem [A.4.](#page-18-1) For each coordinate pair $p = (\phi(p), d(p))$, we normalize the arclength d by leting $u = d/D_{\phi}$ Note that in this normalized distance $u(O_z) = 1$ and so u is continuous on T_x .

We will show that the heteroclinic triangle T_x with corners O_x , O_y , and O_z is homeomorphic to the triangle ABC in the plane with corners at $A = (0,0), B = (b,1/2)$ and $C = (1,0)$.

Now for a given point w on T identified by (ϕ, u) , consider the map:

$$
H: w \mapsto (u, v) = (u, h(u, \phi)).
$$

where $v = h(u, \phi)$ is given by the continuous map:

$$
v = \begin{cases} \frac{u}{2b} \tan(\phi/2) & \text{if } 0 \le u \le b\\ \frac{1-u}{2(1-b)} \tan(\phi/2) & \text{if } b < u \le 1. \end{cases}
$$

The map H is a homeomorphism. It is one-to-one at all points. It maps O_x , O_y and O_z onto A, B and C respectively. At interior points it is a local homeomorphism because it is a composition of local homeomorphisms. At the corners O_x and O_z , considering the mappings as polar coordinates make it clear that H is a local homeomorphism there. It is \Box also clear that H is one-to-one along the edges Γ_{xz} , Γ_{xy} and Γ_{yz} .

Thus Figure [5](#page-18-0) represents not merely an easy-to-understand representation of Γ, but a triangulation. We now investigate the orientibility of Γ. The most common definition of orientibility, in terms of normal vectors, is not useful in this situation, but having triangulated Γ, we may use instead a less common, but still standard definition (see e.g. [\[7\]](#page-24-13)).

Definition A.10. Consider some arbitrary triangle in the triangulation of a manifold. Assign to each triangle in the triangulation a value of clockwise. This assigns corresponding directions to each side of each of the triangles. Now let Δ_i and Δ_j be triangles sharing a side; observe that the common side has been assigned two different directions, i.e. two adjacent triangles with the same orientation conflict on their shared side. If this can be done without contradiction, then the surface is orientable. If a contradiction is reached, the surface is non-orientable.

It is a standard result of algebraic topology that orientibility is independent of the specific triangulation used.

Using this definition and the specific triangulation we have defined, the following lemma can be easily proven.

Lemma A.11. If p is odd, Γ is non-orientable. If p is even, Γ is orientable.

Proof. Let p be odd. We consider the triangulation by heteroclinic triangles. Start with Δ_1 , the triangle defined by O_1 , O_2 , and O_3 and without loss of generality assign it the clockwise orientation. Thus the direction on the $O_1 - O_2$ side of the triangle is given the direction $O_2 \to O_1$. Likewise, the triangle Δ_i defined by O_p , O_1 , and O_2 is assigned the clockwise orientation, and the $O_1 - O_2$ side of the triangle is given the direction $O_2 \rightarrow O_1$. But Δ_i and Δ_i are adjacent triangles, and the fact that they give the same direction to the $O_1 - O_2$ side means that the surface is non-orientable.

Let p be even. We consider the triangulation by heteroclinic triangles. Giving the triangle defined by O_1 , O_2 , and O_3 the clockwise orientation, and thus giving the $O_1 - O_2$ edge the $O_2 \rightarrow O_1$ direction, we clearly observe that making the O_p , O_1 , O_2 triangle clockwise gives the $O_1 - O_2$ boundary, the only place where orientibility could be broken, the conflicting $O_1 \rightarrow O_2$ direction. □

We cite one more result, a classification theorem [\[8\]](#page-24-12).

Theorem A.12. Given a compact connected triangulable 2-manifold Y with boundary such that ∂Y has k components, Y is homeomorphic to X-with-k-holes, where X is S^2 or the n-fold torus T_n or the m-fold projective plane P_m .

A "hole" in the sense of the theorem is a set homeomorphic to an small ϵ open ball.

Our extensive build-up makes the proof of the theorem almost trivial.

Theorem A.13. If p is odd, then Γ is homeomorphic to a Möbius strip. If p is even, Γ is homeomorphic to a cylinder.

Proof. Let p be odd. Of the possible homeomorphic images named in Theorem [A.12,](#page-23-0) only the projective plane is non-orientable; since we know from Lemma [A.1](#page-17-0) that Γ has only one boundary component, Γ is homeomorphic to the projective plane with a ball removed, which is homeomorphic to the Möbius strip.

Let p be even. Then Γ is a rectangle with two of its edges identified. Since the "right-hand" point of the bottom edge is identified with the "right-hand" point of the top edge, it is a cylinder. \Box

References

- [1] Afraimovich V S, Cuevas D and Young T 2013 Sequential dynamics of master-slave systems Dynamical Systems: an International Journal 28(2) 154 - 172
- [2] Afraimovich V S, Tristan I, Huerta R and Rabinovich M 2008 Winnerless competition principle and prediction of the transient dynamics in a Lotka-Volterra model CHAOS 18 43103
- [3] Afraimovich V S, Tristan I, Varona P and Rabinovich M 2013 Transient dynamics in complex systems: heteroclinic sequences with multidimensional unstable manifolds Discontinuity, Nonlinearity and Complexity $2(1)$ 21 - 41
- [4] Afraimovich V S, Zhigulin V and Rabinovich M 2014 On the origin of the reproducible activity in neural circuits CHAOS 14 1123-1129
- [5] Ashwin P, Karabacak O and Nowotny T 2011 Criteria for robustness of heteroclinic cycles in neural microcircuits The Journal of Mathematical Neuroscience 1(13)
- [6] Ashwin P and Postlethwaite C M 2013 On designing heteroclinic networks from graphs Physica D 265 26-39.
- [7] Conlon L 2008 Differentiable Manifolds (Birkhauser: New York)
- [8] George T 2011 The classification of surfaces with boundary available at: http://www.math.uchicago.edu/∼[may/VIGRE/VIGRE2011/REUPapers/George.pdf](http://www.math.uchicago.edu/~may/VIGRE/VIGRE2011/REUPapers/George.pdf)
- [9] Guckenheimer J and Holmes P 1988 Structurally stable heteroclinic cycles Math. Proc. Camb. Phil. Soc.103 189-192
- [10] Luis A. Gonzalez-Diaz L, Gutiérrez-Sénchez E, Varona P and Cabrera J 2013 Winnerless competition in coupled Lotka-Volterra maps Physical Review E88
- [11] Hofbauer J and Sigmund K 1998 Evolutionary Games and Population Dynamics (Cambridge University Press) 220-232
- [12] Homburg A J and Sandstede B 2010 Homoclinic and heteroclinic bifurcations in vector fields Handbook of Dynamical Systems Volume 3 Broer H, Hasselblatt B and Takens F ed. 379-524
- [13] Horchler A, Daltorio K, Chiel H and Quinn R 2015 Designing responsive pattern generators: stable heteroclinic channel cycles for modeling and control Bioinspiration \mathscr *Biomimetics* 10(2)
- [14] Jones L M, Fontanini A, Sadacca B F, Miller P and Katz P B 2007 Natural stimuli evoke dynamic sequences of states in sensory cortical ensembles *Proc. Natl. Acad. Sci.* USA 104 18772-7
- [15] Kirk V and Silber M 1994 A Competition Between Heteroclinic Cycles Nonlinearity 7 1605-1621
- [16] Koon W, Lo M, Marsden J and Ross S 2000 Heteroclinic connections between periodic orbits and resonance transitions in celestial mechanics $Chaos \, 10(2) \, 427-469$
- [17] Krupa M and Melbourne I, Asymptotic stability of heteroclinic cycles in systems with symmetry Ergodic Theory and Dynamical Systems 15 121-147
- [18] Krupa M and Melbourne I, Asymptotic stability of heteroclinic cycles in systems with symmetry, II Proceedings of the Royal Society of Edinburgh: Section A Mathematics 134(6) 1177-1197
- [19] May R and Leonard W 1975 Nonlinear aspects of competition between three species SIAM Journal of Applied Mathematics 29(2) 243-253
- [20] Ohara S, Lenz F and Zhou Y D 2006 Sequential neural processes of tactile-visual crossmodal working memory Neuroscience 139(1) 299-309
- [21] Perko L 1991 Differential Equations and Dynamical Systems (Springer: New York) 194
- [22] Rabinovich M 2013 private communication
- [23] Rabinovich M, Huerta R and Laurent G 2008 Transient dynamics for neural processing Science 321 48-50
- [24] Rabinovich M, Varona P, Tristan I and Afraimovich V S 2014 Chunking dynamics: heteroclinics in mind Front. Comput. Neuroscience 8
- [25] Rabinovich M I, Varona P, Selverston A I and Abarbanel H D I 2006 Dynamical principles in neuroscience Rev. Mod. Physics 78 1213-65
- [26] Rabinovich M, Volkovski A, Lecandra P, Huerta R, Abarbanel H D I and Laurent G 2011 Dynamical encoding by networks of competing neuron groups: winnerless competition Phys. Rev. Letters 87
- [27] Shilnikov L P, Shilnikov A L, Turaev D and Chua L 1998 Methods of Qualitative Theory In Nonlinear Dynamics Part 1 (World Scientific) 82-83
- [28] Schwappach C, Hutt A and Graben P 2015 Metastable dynamics in heterogeneous neural fields Front Syst Neurosci.9(97)
- [29] Wolkowicz G 2006 Interpretation of the generalized asymmetric May-Leonard model of three species competition as a food web in a chemostat Fields Institute Communications 48 279-289.