Two-sex mosquito model for the persistence of Wolbachia

Ling Xue ^{a*} and Carrie A. Manore^a and Panpim Thongsripong ^b and James M. Hyman^a

 ^a Department of Mathematics, Center for Computational Science, Tulane University, New Orleans, LA 70118;
 ^bSchool of Public Health and Tropical Medicine, Tulane University, New Orleans, LA 70112

We develop and analyze an ordinary differential equation model to investigate the transmission dynamics of releasing *Wolbachia*-infected mosquitoes to establish an endemic infection in a population of wild uninfected mosquitoes. *Wolbachia* is a genus of endosymbiotic bacteria that can infect mosquitoes and reduce their ability to transmit dengue virus. Although the bacterium is transmitted vertically from infected mothers to their offspring, it can be difficult to establish an endemic infection in a wild mosquito population. Our transmission model for the adult and aquatic-stage mosquitoes takes into account *Wolbachia*-induced fitness change and cytoplasmic incompatibility. We show that, for a wide range of realistic parameter values, the basic reproduction number, R_0 , is less than one. Hence, the epidemic will die out if only a few *Wolbachia*-infected mosquitoes are introduced into the wild population. Even though the basic reproduction number is less than one, an endemic *Wolbachia* infection unstable endemic equilibrium, and a high-infection stable endemic equilibrium, an intermediate-infection unstable endemic equilibrium, and a high-infection stable endemic equilibrium. We analyze the impact of reducing the wild mosquito population before introducing the infected mosquitoes and observed that the most effective approach to establish the infection in the wild is based on reducing mosquitoes in both the adult and aquatic stages.

Keywords: dengue; vertical transmission; backward bifurcation; cytoplasmic incompatibility; basic reproduction number

1. Introduction

We use a disease transmission model to investigate the conditions for releasing *Wolbachia*-infected mosquitoes that will establish an endemic infection in a population of wild uninfected mosquitoes. The *Wolbachia* infected mosquitoes are less able to transmit dengue virus, and the goal of the modeling effort is to better understand how this bacterium can be used to control vector-borne diseases. We observed that the basic reproductive number for the *Wolbachia* model is less than one for typical model parameters, and so small *Wolbachia* infestations will die out. However, the model predicts that there is a critical threshold, and if a sufficient number of infected mosquitoes are released, then an endemic *Wolbachia*-infected population of mosquitoes can be established. We also investigated how this critical threshold can be reduced by first decreasing the population of wild mosquitoes.

Dengue is the world's most significant and widespread arthropod-borne viral disease [1]. Each year, 400

^{*}Corresponding author. Email: lxue2@tulane.edu

million people are infected with dengue virus in more than 100 countries [2], while the other one-third of the world's population is at risk. To date, there are no vaccines or specific therapy available. Traditional control strategies focusing on reducing population of *Aedes* mosquito vectors have failed to slow the current dengue pandemic, especially in tropical communities [3]. The main vector, *Aedes aegypti*, rebounds in many areas and the secondary vector, *Aedes albopictus* keeps expanding its geographic distribution, leading to 30 fold increase in cases over the past 50 years [4], which necessitates effective novel alternatives to break dengue transmission cycles [3] targeting *Aedes aegypti* and *Aedes albopictus*.

Increasing attention has been paid to controlling the spread of dengue by targeting mosquito longevity by introducing genetically modified mosquitoes or introducing endosymbiotic *Wolbachia* bacteria to shorten the mosquito lifespan [3, 5, 6]. That is, *Wolbachia*-infected mosquitoes are released to create a sustained infection in the wild (uninfected) population. If the infection is sustained, then the wild infected mosquitoes will be less effective in transmitting dengue fever. We create and analyze a mathematical model to help understand the underlying dynamics of *Wolbachia*-infected mosquitoes that are needed to create a sustained endemic *Wolbachia* infection. Once the *Wolbachia* disease transmission model is well understood, one of our future goals will be to couple this model with a mosquito-human model for the spread of dengue.

1.1. Wolbachia Bacteria

The wMel strain of *Wolbachia pipientis* bacteria is a maternally inherited endosymbiont infecting more than 60% of all insect species. This strain has the ability to alternate host reproduction through parthenogenesis, which results in the development of unfertilized eggs, male killing, feminization, and cytoplasmic incompatibility (CI) [7, 8] that prevents the eggs from forming viable offspring. The latter includes strategies for both the suppression and replacement of medically important mosquito populations. Cytoplasmic incompatibility is an incompatibility between the sperms and eggs induced by *Wolbachia* infection [9] and has received considerable attention as a method to control vector-borne diseases [10]. *Wolbachia* even induces resistance to dengue virus in *Aedes aegypti* [11] and limits transmission of dengue virus in *Aedes albopictus* [12].

Uninfected females only mate successfully with uninfected males, while infected females can mate successfully with both uninfected and infected males [13]. If a male fertilizes a female harboring the same type of infection, the offspring still can survive [6]. When *Wolbachia*-infected males mate with uninfected females, or females infected with a different *Wolbachia* strain, then the CI often results in killing the embryos. [9]. Therefore, infected females have a reproduction advantage over uninfected females due to protection from CI [14].

To successfully transmit dengue virus, a vector must imbibe virus particles during blood-feeding and survive to the point that the pathogen can be biologically transmitted to the next vertebrate host [15]. This time period, called extrinsic incubation period (EIP), varies with ambient temperature, many other climatic factors, and characteristics of the vector-parasite system [16]. Vectors that survive long enough to transmit the pathogen are called effective vectors [15]. Typically dengue virus has an incubation period as long as two weeks to transmit through *Aedes aegypti* populations [1]. A life-shortening strain of *Wolbachia* may halve the life span of *Aedes aegypti* [17]. *Wolbachia* infection may reduce the rate of disease transmission due to the reduction on the lifespan of infected mosquitoes or the interference with mosquito susceptibility to dengue virus.

1.2. Existing Mosquito-Wolbachia Models

Ordinary differential equation (ODE) models have been developed to explore key factors that determine the success of applying *Wolbachia* to dengue control. A single-sex model for *Wolbachia* infection with both age-structured and unstructured models were presented to study the stability and equilibrium based on the assumption that *Wolbachia* infection leads to increased mortality or reduced birth rate [18]. A model assuming a fixed ratio of females and males addressed how pathogen protection affects *Wolbachia* invasion [19]. Age-structured and unstructured models combining males and females were found to be different in terms of existence and stability of equilibrium solutions [18]. A stochastic model for female mosquitoes was developed to investigate the impact of introduction frequency on establishment of *Wolbachia* [20].

Discrete-time models explored the impact of the type of immigration and the temporal dynamics of the host population on the spread of *Wolbachia*, assuming equal sex ratio between males and females [21]. Discrete generation models for female mosquitoes were built to understand unstable equilibrium produced by reduced lifespan or lengthened development [22]. Reaction-diffusion and integro-difference equation model has been used to analyze the impact of insect dispersal and infection spread on invasion of *Wolbachia* [23].

An ordinary differential equation (ODE) model was developed to evaluate the desirable properties of the *Wolbachia* strain to be introduced to female mosquitoes, assuming that *Wolbachia*-infected mosquitoes have reduced lifespan and reduced capability to transmit dengue, and equal fraction of male and female mosquitoes [24]. A continuous time non-spatial model and a reaction-diffusion model incorporating lifespan shortening and CI were developed to study factors that determine the spatial spread of *Wolbachia* through a population of female *Aedes aegypti* mosquitoes assuming constant population size and perfect maternal transmission of *Wolbachia* [25]. A two-sex deterministic model with deterministic immature life stages and stochastic female adult life stage was developed to understand *Wolbachia* invasion into uninfected host population [26].

A single strain model, two strain model, and spatial model were developed to study whether multi-stain of *Wolbachia* can coexist in a spatial context [27]. A two-sex ODE model taking into account different death rates, but the same egg laying rates of *Wolbachia*-infected and *Wolbachia* uninfected mosquitoes [28] showed the basic reproduction number is always less than one, and the complete infection equilibrium is locally asymptotically stable (LAS) due to positive determinant of the Jacobian matrix for the system. Simulations showed that dengue epidemics will not occur when *Wolbachia* infection is sufficiently prevalent [28]. A two-sex mosquito model assuming complete vertical transmission and equal death rates for male and female mosquitoes was developed and four steady states were found [29].

Most of these models consider either a single-sex model for adult mosquitoes, or assume a fixed ratio between the number of male and female mosquitoes. Also, most of the models assume homogeneous death rates and egg laying rates for *Wolbachia*-free and *Wolbachia*-infected mosquitoes. Our model addresses both of these issues. Our emphasis is to understand how *Wolbachia* infection can be established in a wild population of mosquitoes.

1.3. Results

We proposed a compartmental two-sex model to investigate the underlying mechanisms that may contribute to invasion and sustainable establishment of *Wolbachia* in mosquito populations. We assigned female and male mosquitoes to different classes to understand corresponding roles that they are playing in the spread of *Wolbachia* in mosquito populations.

We showed that, for a wide range of realistic parameter values, the basic reproduction number, R_0 , for this model is less than one. Hence, the epidemic will die out if only a few *Wolbachia*-infected mosquitoes are introduced into the wild population. Even though the basic reproduction number is less than one, an endemic *Wolbachia* infection can be established if a sufficient number of infected mosquitoes are released. This threshold effect can be explained as a backward bifurcation with three coexisting equilibria: a stable zero-infection equilibrium, an intermediate-infection unstable endemic equilibrium, and a high-infection stable endemic equilibrium.

If the number of infected individuals is below the unstable endemic equilibrium, then the infection decays to the zero-infection equilibrium. Conversely, if the number of infected mosquitoes is greater than unstable endemic equilibrium, then the solution tends to the stable high-infection equilibrium. We identified the relationships between dimensionless combinations of model parameters and the initial conditions for *Wolbachia* to be attracted to the high-infection state. As expected, the number of infected female mosquitoes needed to be released to establish the infection in a wild *Wolbachia*-free population decreases as R_0 increases to one. We analyzed the impact of reducing the wild mosquito population before introducing the infected mosquitoes. We found that the most effective approach of reducing the number of infected mosquito populations in both the adult and aquatic stages before the release. This could be accomplished by recursive spraying, or a combination of spraying and larvae control.

Our main findings are:

- (1) Three equilibria, disease free equilibrium (DFE), endemic equilibrium (EE), and complete infection equilibrium (CIE) coexist when $R_0 < 1$. Disease free equilibrium is a steady state when all individuals are *Wolbachia*-free. Endemic equilibrium is a steady state when some individuals are *Wolbachia*-free, the rest are infected with *Wolbachia*. Complete infection equilibrium is a steady state when all individuals are infected with *Wolbachia*.
- (2) The backward bifurcation analysis of our *Wolbachia* transmission model predicts that if $R_0 < 1$, then there is a critical threshold for the number of infected mosquitoes released in the wild before the infection can be established. If we release too few *Wolbachia* infected mosquitoes, then *Wolbachia* infection will die out.
- (3) Killing both aquatic state (eggs and larvae) and adult mosquitoes before releasing the infected mosquitoes greatly increases the chance that the infection will be established. Our model quantifies what fraction of wild mosquitoes must be killed before releasing *Wolbachia* infected mosquitoes.

After introducing the mathematical model, we summarize the key results from the analysis and numerical simulations. We conclude with a discussion of the relevance, importance, and future directions for this work.

2. Description of Model Framework

We developed an ODE model incorporating adult females (F), adult males (M), and an aggregated aquatic (A) stage that includes the egg, larvae, and pupae stages. The population dynamics of mosquitoes without taking into account *Wolbachia* is in the Appendix, Equation 9. The vertical transmission of *Wolbachia* from infected females to their offspring is a key factor in establishing an endemic infected population.

Mosquitoes are grouped into six compartments: susceptible aquatic stage, A_u , infected aquatic stage, A_w , susceptible female mosquitoes, F_u , infected female mosquitoes, F_w , susceptible male mosquitoes, M_u , and infected male mosquitoes, M_w . The eclosion rates of susceptible female and male mosquitoes hatching from eggs are $\psi \theta A_u$ and $\psi(1-\theta)A_u$, respectively. Similarly, the birth rates of infected female and male mosquitoes are $\psi \theta A_w$ and $\psi(1-\theta)A_w$. Death rates of uninfected male mosquitoes and infected male mosquitoes are $\mu_a A_u$ and $\mu_a A_w$. Death rates of uninfected female mosquitoes and infected female mosquitoes are $\mu_{fu}F_u$ and $\mu_{fw}F_w$. Death rates of uninfected male mosquitoes and infected male mosquitoes are $\mu_{mu}M_u$ and $\mu_{mw}M_w$. Development rates of uninfected aquatic stage and infected aquatic stage of mosquitoes are ψA_u and ψA_w . The model parameters are described in Table 1.

The model (Fig. 1) describing population dynamics of aquatic stage, adult male, and adult female mosquitoes is given by:

$$\frac{\mathrm{d}A_u}{\mathrm{d}t} = B_{uu} + v_u(B_{wu} + B_{ww}) - \mu_a A_u - \psi A_u \tag{1a}$$

$$\frac{\mathrm{d}A_w}{\mathrm{d}t} = v_w (B_{wu} + B_{ww}) - \mu_a A_w - \psi A_w \tag{1b}$$

$$\frac{\mathrm{d}F_u}{\mathrm{d}t} = b_f \psi A_u - \mu_{fu} F_u \tag{1c}$$

$$\frac{\mathrm{d}F_w}{\mathrm{d}t} = b_f \psi A_w - \mu_{fw} F_w \tag{1d}$$

$$\frac{\mathrm{d}M_u}{\mathrm{d}t} = b_m \psi A_u - \mu_{mu} M_u \tag{1e}$$

$$\frac{\mathrm{d}M_w}{\mathrm{d}t} = b_m \psi A_w - \mu_{mw} M_w \quad . \tag{1f}$$

Because the vertical transmission and birth rates, B_{**} , depend on the sex of the infected or uninfected parents, the model included the four egg laying situations

$$B_{uu} = \phi_u F_u m_u \left(1 - \frac{N_A}{K_a} \right) \tag{2a}$$


$$B_{uw} = 0 \tag{2b}$$

$$B_{wu} = \phi_w F_w m_u \left(1 - \frac{N_A}{K_a} \right) \tag{2c}$$

$$B_{ww} = \phi_w F_w m_w \left(1 - \frac{N_A}{K_a} \right) \quad . \tag{2d}$$

Here $m_u = \frac{M_u}{M_w + M_u}$ and $m_w = 1 - m_u$ are the fractions of uninfected and infected male mosquitoes. B_{uu} is the egg laying rate of uninfected females mating with uninfected males. B_{uw} is the egg laying rate of uninfected males. M_A is the total number of aquatic stage of mosquitoes, and K_a is carrying capacity of aquatic stage of mosquitoes. These equations reflect the observations in Table 2 that

- Mating of uninfected males with uninfected females produce uninfected offspring.
- Mating of infected males with uninfected females leads to death of embryos before hatching due to cytoplasmic incompatibility.
- Uninfected males and infected females produce a fraction, denoted by v_w , of infected offspring by vertical transmission.



(3)

Figure 1. The birthing rates (2) capture that when the uninfected males mate with uninfected females, they produce uninfected offspring. When infected males mate with uninfected females, then CI causes the embryos to die before hatching. Uninfected males mating with infected females produce a fraction, denoted by v_w , of infected offspring by vertical transmission. Cross of infected males with infected females produces a fraction of infected offspring.

• Cross of infected males with infected females produces a fraction of infected offspring.

2.1. Model Analysis

We compute the basic reproduction number and the equilibria, and analyze the stability of the equilibrium points.

We use the next generation matrix approach to compute the basic reproduction number [30]. Only infected compartments are considered for ease of computation:

$$\frac{d}{dt} \begin{bmatrix} A_w & F_w & M_w \end{bmatrix}^T = \mathscr{F} - \mathscr{V} = \begin{bmatrix} (v_w F_w \phi_w m_u + v_w F_w \phi_w m_w) \left(1 - \frac{N_A}{K_a}\right) \\ b_f \psi A_w \\ b_m \psi A_w \end{bmatrix} - \begin{bmatrix} \mu_a A_w + \psi A_w \\ \mu_{fw} F_w \\ \mu_{mw} M_w \end{bmatrix},$$

where $\mathscr{F} = (\mathscr{F}_i)$ is a vector for new infected, and $\mathscr{V} = (\mathscr{V}_i)$ is a vector for transfer between compartments. Jacobian matrices for transmission, F, and transition, V, [30] are defined as:

$$F = \left[\frac{\partial \mathscr{F}_i(x^0)}{\partial x_j}\right], \quad V = \left[\frac{\partial \mathscr{V}_i(x^0)}{\partial x_j}\right],\tag{4}$$

where x^0 represents the disease free equilibrium, and x_j is the number or proportion of infected individuals in compartment $j, j = 1, 2, \dots, m$.

- M_u : Number of susceptible male mosquitoes
- M_w : Number of male mosquitoes infected with Wolbachia
- F_u : Number of susceptible female mosquitoes
- F_w : Number of female mosquitoes infected with Wolbachia
- A_u : Number of susceptible aquatic stage of mosquitoes
- A_w : Number of infected aquatic stage of mosquitoes
- N_A : Total number of aquatic stage of mosquitoes
- K_a : Carrying capacity of aquatic stage of mosquitoes
- b_f : Fraction of births that are female mosquitoes.
- b_m : Fraction of births that are male mosquitoes $= 1 b_f$.
- m_w : Fraction of the male mosquitoes that are infected = $M_w/(M_w + M_u)$.
- m_u : Fraction of the male mosquitoes that are uninfected = $1 m_w$.
- v_w : Fraction of infected mosquito eggs produced by infected female mosquitoes.
- v_u : Fraction of uninfected mosquito eggs produced by infected female mosquitoes = $1 v_w$.
- ϕ_u : Per capita egg laying rate by *Wolbachia*-free mosquito eggs. Number of eggs/time
- ϕ_w : Per capita egg laying rate by *Wolbachia*-infected mosquito eggs. Number of eggs/time
- ψ : Per capita development rate of mosquito eggs. Time⁻¹
- μ_a : Per capita death rate of aquatic stage of mosquitoes. Time⁻¹
- μ_{fu} : Per capita death rate of uninfected female mosquitoes. Time⁻¹
- μ_{fw} : Per capita death rate of infected female mosquitoes. Time⁻¹
- μ_{mu} : Per capita death rate of uninfected male mosquitoes. Time⁻¹
- μ_{mw} : Per capita death rate of infected male mosquitoes. Time⁻¹

Table 1. State variables and parameters for the model (1)

Male	Female	Offspring
Uninfected	Uninfected	Uninfected
Uninfected	Infected	Some offspring are infected
Infected	Uninfected	Unviable
Infected	Infected	Some offspring are infected

Table 2. The model allows for *Wolbachia* to be transmitted vertically from infected parents to their offspring. The offspring of male and female uninfected mosquitoes are infected. Some of the offspring of male and female infected mosquitoes are infected, as are the offspring of an uninfected male and an infected female mosquitoes. The offspring of an infected male and uninfected female mosquito are unviable.

The unique disease free equilibrium is

$$A_u^0 = K_a \left(1 - \frac{1}{R_{0u}} \right)$$
$$F_u^0 = b_f \frac{\psi}{\mu_{fu}} A_u^0$$
$$M_u^0 = b_m \frac{\psi}{\mu_{mu}} A_u^0$$
$$A_w^0 = F_w^0 = M_w^0 = 0$$

where $R_{0u} = \frac{b_f \phi_u \psi}{(\mu_a + \psi)\mu_{fu}}$ is the threshold for *Wolbachia*-free offspring, $\frac{b_f \phi_u}{\mu_{fu}}$ is the total number of eggs laid by uninfected female mosquitoes, $\frac{\psi}{\mu_a + \psi}$ is the probability that aquatic stage of mosquitoes survive to the point when they develop into adult mosquitoes. R_{0u} is the number of female eggs that develop into adult mosquitoes. When $R_{0u} > 1$, then the mosquito population may grow; otherwise, the population will decrease.

The Jacobian matrix of \mathscr{F} evaluated at DFE is:

$$F = \begin{bmatrix} 0 & v_w m_u^0 \phi_w \left(1 - \frac{N_A^0}{K_a} \right) & 0\\ b_f \psi & 0 & 0\\ b_m \psi & 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & \frac{v_w (\mu_a + \psi) \mu_{fu} \phi_w}{b_f \psi \phi_u} & 0\\ b_f \psi & 0 & 0\\ b_m \psi & 0 & 0 \end{bmatrix},$$

where $m_u^0 = \frac{M_u^0}{M_w^0 + M_u^0}$. The Jacobian matrix of $\mathscr V$ evaluated at DFE is:

$$V = \begin{bmatrix} \mu_a + \psi & 0 & 0 \\ 0 & \mu_{fw} & 0 \\ 0 & 0 & \mu_{mw} \end{bmatrix}$$

The next generation matrix for infected compartments at disease free equilibrium is:

$$FV^{-1} = \begin{bmatrix} 0 & \frac{v_w(\mu_a + \psi)\mu_{fu}\phi_w}{b_f\psi\mu_{fw}\phi_u} & 0\\ \frac{b_f\psi}{\mu_a + \psi} & 0 & 0\\ \frac{b_m\psi}{\mu_a + \psi} & 0 & 0 \end{bmatrix}.$$

The basic reproduction number is the largest abosulute eigenvalue of FV^{-1} , denoted by $\rho(FV^{-1})$:

$$R_0 = \rho(FV^{-1}) = \sqrt{v_w \frac{\phi_w}{\mu_{fw}} \left(\frac{\phi_u}{\mu_{fu}}\right)^{-1}} \quad . \tag{5}$$

Wolbachia-infected and *Wolbachia*-free and females produce $\frac{\phi_w}{\mu_{fw}}$ and $\frac{\phi_u}{\mu_{fu}}$ eggs during their lifetime, respectively. Hence, $\frac{\phi_w}{\mu_{fw}} \left(\frac{\phi_u}{\mu_{fu}}\right)^{-1}$ is the ratio of the number of eggs produced by *Wolbachia*-infected females to the number of eggs produced by Wolbachia-free females during their lifetime. R_0 is geometric mean of vertical transmission rate and the ratio of the total number of eggs produced by Wolbachia-infected females to the total number of eggs produced by Wolbachia-free females.

When the variables are ordered as: $[A_u \ F_u \ M_u \ A_w \ F_w \ M_w]^T$, Jacobian matrix at equilibrium $[A_u^* F_u^* M_u^* A_w^* F_w^* M_w^*]^T$ for system of equations (1) is:

$$J = \begin{bmatrix} c & a & b & | & c_2 & a_2 & b_2 \\ b_f \psi & -\mu_{fu} & 0 & | & 0 & 0 & 0 \\ b_m \psi & 0 & -\mu_{mu} & | & 0 & 0 & 0 \\ -- & -- & -- & -- & -- \\ c_3 & 0 & b_3 & | & c_1 & a_1 & b_1 \\ 0 & 0 & 0 & | & b_f \psi & -\mu_{fw} & 0 \\ 0 & 0 & 0 & | & b_m \psi & 0 & -\mu_{mw} \end{bmatrix} = \begin{bmatrix} A & C \\ D & B \end{bmatrix},$$

where

$$\begin{aligned} a &= \phi_u m_u^* \left(1 - \frac{A_u^* + A_w^*}{K_a} \right) \\ b &= \phi_u m_w^* \frac{F_u^*}{M_w^* + M_u^*} \left(1 - \frac{A_u^* + A_w^*}{K_a} \right) \\ c &= -(\mu_a + \psi) - \phi_u m_u^* \frac{F_u^*}{K_a} - \phi_w v_u \frac{F_w^*}{K_a} \\ a_1 &= v_w \phi_w \left(1 - \frac{A_u^* + A_w^*}{K_a} \right) \\ b_1 &= 0 \\ c_1 &= -(\mu_a + \psi) - \phi_w v_w \frac{F_w^*}{K_a} \\ a_2 &= \phi_w v_u \left(1 - \frac{A_u^* + A_w^*}{K_a} \right) \\ b_2 &= -\phi_u m_u^* \frac{F_u^*}{M_w^* + M_u^*} \left(1 - \frac{A_u^* + A_w^*}{K_a} \right) \\ c_2 &= -\phi_u m_u^* \frac{F_u^*}{K_a} - \phi_w v_u \frac{F_w^*}{K_a} \\ b_3 &= 0 \\ c_3 &= -\phi_w v_w \frac{F_w^*}{K_a} \end{aligned}$$

At disease free equilibrium, D = 0, then the eigenvalues of J are eigenvalues of A and B. Matrices A and B are:

$$A = \begin{bmatrix} -(\mu_a + \psi) - \phi_u \frac{F_u^0}{K_a} & \frac{\mu_{fu}(\mu_a + \psi)}{b_f \psi} & 0\\ b_f \psi & -\mu_{fu} & 0\\ b_m \psi & 0 & -\mu_{mu} \end{bmatrix}, \quad B = \begin{bmatrix} -(\mu_a + \psi) & \frac{v_w(\mu_a + \psi)\mu_{fu}\phi_w}{b_f \psi\phi_u} & 0\\ b_f \psi & -\mu_{fw} & 0\\ b_m \psi & 0 & -\mu_{mw} \end{bmatrix}.$$

If $R_{0u} > 1$, then all eigenvalues of A are negative. If $R_0 < 1$, then all eigenvalues of B are negative. Therefore, the system (1) at disease free equilibrium is LAS whenever $R_{0u} > 1$ and $R_0 < 1$.

2.1.1. Complete Vertical Transmission

If $R_0 < 1$ and $R_{0w} > 1$ and the vertical transmission is 100%, $(v_w = 1)$, then the ratio of the infected to the uninfected aquatic stage mosquitoes, k, is

$$k = \frac{A_w^*}{A_u^*} = \frac{\mu_{mw}}{\mu_{mu}} (R_0^{-2} - 1) \quad ,$$

and there is a unique endemic equilibrium:

$$A_u^* = \frac{K_a}{1+k} \left(1 - R_{0w}^{-1} \right) \tag{6a}$$

$$A_w^* = k A_u^* \tag{6b}$$

$$F_u^* = b_f \frac{\psi}{\mu_{fu}} A_u^* \tag{6c}$$

$$F_w^* = k F_u^* \tag{6d}$$

$$M_u^* = b_m \frac{\psi}{\mu_{mu}} A_u^* \tag{6e}$$

$$M_{w^*} = k M_u^*, \tag{6f}$$

where $R_{0w} = \frac{v_w b_f \psi \phi_w}{(\mu_a + \psi) \mu_{fw}}$. If we further assume that $\mu_{mu} = \mu_{mw}$, then $k = R_0^{-2} - 1$ and the unique complete infection equilibrium (CIE) is:

$$A_u = 0$$

$$A_w = K_a \left(1 - R_{0w}^{-1}\right)$$

$$F_u = 0$$

$$F_w = b_f \frac{\psi}{\mu_{fw}} A_w$$

$$M_u = 0$$

$$M_w = b_m \frac{\psi}{\mu_{mw}} A_w.$$

Jacobian matrix of system of equations (1) at complete Wolbachia infection equilibrium for $v_w = 1$ is:

Eigenvalues of J_{cw} are composed of three eigenvalues of A and three eigenvalues of B. The eigenvalues of A are all negative. Characteristic polynomial of B is:

$$\lambda^2 + \left(\mu_{fw} + \frac{\phi_w b_f \psi}{\mu_{fw}}\right)\lambda + \phi_w b_f \psi - (\mu_a + \psi)\mu_{fw} = 0 \quad . \tag{7}$$

If $R_{0w} > 1$, all eigenvalues of B are negative. Therefore, the complete infection equilibrium is LAS whenever $R_{0w} > 1$, as shown in Figure 2(a).

Parameter	Baseline Value	Ranges	References
b_f	0.5	0.34 - 0.6	[31 - 33]
ϕ_u	50/day	0 - 75	[32]
ϕ_w	51/day	0 - 75	[32]
ψ	$0.01/\mathrm{day}$		Assume
μ_a	0.02/day		Assume
v_w	0.9	0 - 1	Assume
μ_{fu}	$0.061/\mathrm{day}$	1/55 - 1/11	[32, 34, 35]
μ_{fw}	$0.068/\mathrm{day}$	1/55 - 1/11	[32, 34, 35]
μ_{mu}	$0.068/\mathrm{day}$	1/31 - /7	[32, 34, 35]
μ_{mw}	$0.068/\mathrm{day}$	1/31 - /7	[32, 34, 35]
K_a	100,000		Assume

Table 3. Baseline values for parameters in Model (1).

Vertical	Disease Free	Endemic	Complete Infection
Transmission	Equilibrium (DFE)	Equilibrium (EE)	Equilibrium (CIE)
Complete	$R_0 < 1$ and $R_{0u} > 1$,	$R_0 < 1$ and $R_{0w} > 1$, unstable	$R_{0w} > 1$, LAS
	LAS		
Incomplete	$R_0 < 1$ and $R_{0u} > 1$,	$0.5 < v_w < 1$ and $4v_w(v_w - 1) <$	does not exist
	LAS	$R_0 < 1$, when $k = k_1$, LAS, when	
		$k = k_2$, unstable	

Table 4. Threshold condition for existence of disease free equilibrium, endemic equilibrium, and complete infection equilibrium and their stability. R_{0u} is the threshold for *Wolbachia*-free mosquito population, and R_{0w} is the threshold for *Wolbachia*-infected mosquito population. Only when $R_{0u} > 1$, *Wolbachia*-free may grow, and only when $R_{0w} > 1$, *Wolbachia*-infected population may grow.

2.1.2. Incomplete Vertical Transmission

When vertical transmission is incomplete, i.e., $0 < v_w < 1$, then at endemic equilibrium,

$$\frac{A_w^*}{A_u^*} = \frac{v_w - v_u \frac{\mu_{mw}}{\mu_{mu}} \pm \sqrt{\left(v_u \frac{\mu_{mw}}{\mu_{mu}} - v_w\right)^2 - 4v_u v_w (R_0^{-2} - 1) \frac{\mu_{mw}}{\mu_{mu}}}}{2v_u} \quad . \tag{8}$$

We assume $\mu_{mu} = \mu_{mw}$, and let $k = \frac{A_w^*}{A_u^*}$. When $\left(v_u \frac{\mu_{mw}}{\mu_{mu}} - v_w\right)^2 > 4v_u v_w (R_0^{-2} - 1) \frac{\mu_{mw}}{\mu_{mu}}$, that is, $R_0 > 2\sqrt{v_w v_u}$. Equation 7 has two roots: $k_1 = \frac{2v_w - 1 + \sqrt{1 - 4v_w v_u R_0^{-2}}}{2v_u}$ and $k_2 = \frac{2v_w - 1 - \sqrt{1 - 4v_w v_u R_0^{-2}}}{2v_u}$.

If $0.5 < v_w < 1$ and $4v_w(v_w - 1) < R_0 < 1$, two endemic equilibria exist as shown in Figure 2(b), 2(c), and 2(d). When $k = k_1$, the endemic equilibrium is LAS. When $k = k_2$, the equilibrium is not LAS and backward bifurcation occurs. If $v_w \leq 0.5$ and $R_0 \leq 1$, then endemic equilibrium does not exist, only DFE exists. When $k_1 = k_2$, then $R_0 = 2\sqrt{v_w v_u}$. Let $R_0^* = 2\sqrt{v_w v_u}$, which is the intersection of unstable and stable endemic equilibrium. When $R_0 > 1$, a unique endemic equilibrium point exists with $k = k_1$. It is proven to be LAS by numerical simulations.




Figure 2. Bifurcation diagrams for Wolbachia vertical transmission. ϕ_u and μ_{fu} are varying, other parameter values are the same as those baseline values in Table 3. Denote the intersection of two endemic equilibrium, that is, the intersection of the black dashed line and x-axis, as $R_0^* = \sqrt{4v_w v_u}$. When $R_0 < 1$ and $v_w < 0.5$, no endemic equilibria exist. When $R_0 < 1$ and $v_w > 0.5$, as the vertical transmission rate increases so does R_0^* and the LAS equilibrium approaches a constant. If we increase the number of infected females, then the endemic equilibrium may become stable endemic or complete infection equilibrium. If we decrease the number of infected females at endemic equilibrium, then the endemic equilibrium may become disease free equilibrium. WIF denotes Wolbachia-infected female mosquitoes.

3. Results

We observed that Wolbachia can persist when $R_0^* < R_0$, where R_0^* is the turning point of the backward bifurcation. Three equilibria, namely, disease free equilibrium, endemic equilibrium, and complete infection equilibrium coexist when $R_0 < 1$, $R_{0w} > 1$, $v_w = 1$, and $R_{0u} > 1$ as shown in Figure 2(a). The disease free equilibrium is LAS whenever $R_0 < 1$ and $R_{0u} > 1$, and complete infection equilibrium exists and is LAS as long as $R_{0w} > 1$. The unique endemic equilibrium is not LAS, which can become disease free equilibrium by decreasing the number of infected individuals or become complete infection equilibrium by increasing the number of infected individuals. Figures 2(b), 2(c), and 2(d) showed that two endemic equilibrium points exist when $\sqrt{4v_w v_u} < R_0 < 1$ and $v_w > 0.5$, and only one of them is LAS proven by numerical simulations. When v_w is larger, R^* is closer to one. The condition for the existence of the equilibria and their stability are summarized in Table 4.




Figure 3. Thresholds for fraction of infected individuals vary with reproduction number. $A_{u0} + A_{w0} = A_u^0$, $F_{u0} + F_{w0} = F_u^0$, $M_{u0} + M_{w0} = M_u^0$. When $R_0 < 1$, the smaller R_0 is, the larger number of infected female mosquitoes are needed to be released for *Wolbachia* to be endemic. The *Wolbachia* infection is only sustained if the fraction of WIF mosquitoes is above the red dotted line.

Initial condition thresholds for an epidemic to occur vary with the vertical transmission rate as shown in Table 5. When $v_w = 0.9$, the epidemic will spread if at least 40% of the population are initially infected. When $v_w = 0.95$, the epidemic will spread if at least 34% of the population are initially infected. When $v_w = 1$, the epidemic will spread if at least 28% of the population are initially infected. When the vertical transmission rate is high, the threshold for the number of initially infected individuals to start *Wolbachia* epidemic is low.

The initial condition threshold for an epidemic to occur varies with the ratio of death rates of Wolbachia-infected male mosquitoes to death rates of Wolbachia-free male mosquitoes, $\frac{\mu_{mw}}{\mu_{mu}}$, while fixing other parameters. The larger $\frac{\mu_{mw}}{\mu_{mu}}$ is, the larger the initial number of infected individuals required to start a Wolbachia epidemic. With higher vertical transmission rate, a smaller percentage of Wolbachia carriers can invade, as shown in Table 5. Similarly, if we increase ϕ_w or μ_{fu} , or decrease ϕ_u or μ_{fw} , then the threshold for initial infection is smaller. Increasing $\frac{\psi}{\mu_a+\psi}$ will increase the epidemic threshold for initial infection. The thresholds for fraction of initial infections decreasing with the basic reproduction number is shown in Figure 3.

The reproduction number is very sensitive to the vertical transmission rate, egg laying rates of *Wolbachia*-infected infected mosquitoes, egg laying rates of *Wolbachia*-free mosquitoes, and death rates of *Wolbachia*-infected female mosquitoes and *Wolbachia*-free female mosquitoes as shown in Equation (5). The reproduction number varies directly with either the vertical transmission rate, the egg laying rates of *Wolbachia*-infected mosquitoes, or the death rates of *Wolbachia*-free female mosquitoes. The reproduction number varies inversely with egg laying rates of *Wolbachia*-free mosquitoes, or the death rates of *Wolbachia*-free female mosquitoes, or the death rates of *Wolbachia*-free female mosquitoes, or the death rates of *Wolbachia*-free mosquitoes, or the death rates of *Wolbachia*-infected female mosquitoes, but towards opposite direction.

We compared five strategies before the release of Wolbachia-infected female mosquitoes:

- DFE: releasing Wolbachia-infected female mosquitoes at the disease free equilibrium,
- KHA: first killing half of the aquatic stage of mosquitoes,
- KHM: first killing half of the wild adult mosquitoes,
- KHM2: first killing half of the wild adult mosquitoes, and then killing half of the adult mosquitoes again after two weeks,
- KHMA: first killing half of the wild mosquitoes and half aquatic stage of mosquitoes.

Initial condition	$v_w = 0.9$	$v_w = 0.95$	$v_w = 1$
A_{u0}	$\leq 60\% A_u^0$	$\leq 66\% A_u^0$	$\leq 72\% A_u^0$
A_{w0}	$\geq 40\% A_u^0$	$\geq 34\% A_u^0$	$\geq 28\% A_u^0$
F_{u0}	$\leq 60\% F_u^0$	$\leq 66\% F_u^0$	$\leq 72\% F_u^0$
F_{w0}	$\geq 40\% F_u^0$	$\geq 34\% F_u^0$	$\geq 28\% F_u^0$
M_{u0}	$\leq 60\% M_u^0$	$\leq 66\% M_u^0$	$\leq 72\% M_u^0$
M_{w0}	$\geq 40\% M_u^0$	$\geq 34\% M_u^0$	$\geq 28\% M_u^0$

Table 5. Initial condition thresholds for epidemic to occur with different vertical transmission rates. When the number is *Wolbachia*-carrying mosquitoes is above the threshold, *Wolbachia*-infected mosquitoes can petsit, otherwise, they are wiped out by *Wolbachia*-free mosquitoes.

Scenario	minimum release ra-	minimum release ra-	minimum release ra-
	tio when $R_0 = 0.85$	tio when $R_0 = 0.9$	tio when $R_0 = 0.95$
DFE	1.007	0.571	0.259
KHA	0.549	0.317	0.151
KHM	0.478	0.244	0.107
KHM2	0.329	0.165	0.072
KHMA	0.210	0.112	0.051

Table 6. Different population suppression strategies applied before release of *Wolbachia* infected female mosquitoes can reduce the minimum number of *Wolbachia*-infected mosquitoes that can lead to persistence of *Wolbachia*. The top row indicates that you have to release 1.007 times as many infected female mosquitoes as there are wild infect mosquitoes to establish an infection in the wild that starts at the DFE when $R_0 = 0.85$. The number of infected mosquitoes that need to be released to establish an infection decreases with the strategies: KHA denotes killing half of the aquatic stage mosquitoes, KHM denotes killing half of adult wild mosquitoes once, then kill half adult wild mosquitoes again after two weeks, and finally KHMA denotes killing half adult wild mosquitoes and half aquatic stage of mosquitoes.

The ratios of the minimum number of *Wolbachia*-infected female mosquitoes that can lead to persistence of *Wolbachia* to the number of female mosquitoes at disease free equilibrium are listed in Table 6 in decreasing order. Notice that killing the adult mosquitoes only once is not an effective strategy because the aquatic stage mosquitoes hatch and quickly replace the wild uninfected population. Killing both the adult and aquatic (larvae) stage mosquitoes before releasing the infected mosquitoes is the most effective strategy.

4. Discussion

We developed a model considering two sex of aquatic stage and adult mosquitoes, diversity in the death rates of *Wolbachia*-infected mosquitoes and *Wolbachia*-free mosquitoes, and egg laying rates of *Wolbachia*-infected female mosquitoes and *Wolbachia*-free female mosquitoes. The general model is not constrained to particular weather condition, specific *Wolbachia* strains, or specific mosquito species, and it can be easily adapted to *Aedes aegypti* or *Aedes albopictus* at any location with parameters calibrated using realistic environmental factors, such as temperature and rainfall etc.

We found conditions for the existence of multiple equilibria and backward bifurcation. If vertical transmission is complete, then a unique endemic equilibrium exists and is not LAS when $R_0 < 1$ and $R_{0u} > 1$. Backward bifurcation occurs when endemic equilibrium changes into disease free equilibrium if we decrease the initial number of infected individuals, or it reaches another LAS equilibrium if we increase the number of initially infected individuals. When vertical transmission is incomplete but more than 50%, and $R_0 < 1$, two endemic equilibria coexist, but only one endemic equilibrium is LAS such that it becomes disease free equilibrium or endemic equilibrium by perturbation. Since *Wolbachia*-infected mosquitoes are less capable of transmitting dengue virus, complete *Wolbachia*-infection is the ideal case for dengue control.

When $R_0 > 1$, Wolbachia can spread with a small number of initially infected mosquitoes, although very slowly. Population replacement may occur. When $R_0 < 1$, Wolbachia can spread if initial number of infected individuals exceeds a threshold, which depends on vertical transmission rate, ratio of egg laying rates of Wolbachia-infected females to egg laying rates of Wolbachia-free female mosquitoes, and ratio of death rates of infected female mosquitoes to death rates of uninfected female mosquitoes. A smaller number of Wolbachia-infected female mosquitoes is needed to be released for persistence of Wolbachia if a population suppression strategy is implemented before the release.

The reproduction number is the product of the vertical transmission rate, ratio of the egg laying rates of *Wolbachia*-infected mosquitoes to egg laying rates of *Wolbachia*-free mosquitoes, and the ratio of death rates of *Wolbachia*-free mosquitoes to death rates of *Wolbachia* infected mosquitoes. If the total number of eggs produced by *Wolbachia*-infected mosquitoes through vertical transmission is more than the total number of eggs produced by *Wolbachia*-free mosquitoes, then $R_0 > 1$, such that complete infection or endemic equilibrium is LAS.

The number of Wolbachia-infected mosquitoes required for sustained Wolbachia infection depends on vertical transmission rate, v_w , the ratio of the number of eggs laid by Wolbachia-infected mosquitoes to the number of eggs laid by Wolbachia-free mosquitoes, $\frac{\phi_w}{\mu_f w}$, the ratio of the death rates of Wolbachia-free females to death rates of Wolbachia-infected females, $\frac{\mu_{fu}}{\mu_{fw}}$, and the ratio of the death rates of Wolbachia-free males to death rates of Wolbachia-infected males $\frac{\mu_{mu}}{\mu_{mw}}$. These parameters depend on pariculiar Wolbachia strain. If the life span of a mosquito is shorter, then the mosquitoes will lay fewer eggs. Once we know the specific Wolbachia strain that a specific mosquito species, such as Aedes aegypti or Aedes albopictus is carrying, we can estimate the number of infected individuals needed to be released for sustainable Wolbachia establishment using this model.

We find that reducing both the aquatic stage and adult uninfected mosquitoes before releasing *Wolbachia*-infected infected female mosquitoes is the most effective strategy to reduce the number of *Wolbachia*-infected female mosquitoes needed for *Wolbachia* persistence (Table 6). The second most effective strategy was to repeatedly kill the wild uninfected mosquitoes (to reduce both the adult and the aquatic stage mosquitoes) before releasing the infected mosquitoes.

The model and analysis can help in understanding how *Wolbachia* can invade and persist in mosquito populations. In future research, we will couple our model to a dengue fever transmission model and analyze the impact that a bacteria infected mosquito population has on the spread of dengue.

Acknowledgments

This work was supported by the endowment for the Evelyn and John G. Phillips Distinguished Chair in Mathematics at Tulane University, the National Science Foundation MPS award DMS-1122666, and the NIH/NIGMS program for Models of Infectious Disease Agent Study (MIDAS) award U01GM097661. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Science Foundation or the National Institutes of Health. The authors thank Michael Robert

for his many helpful comments.

Appendix

The system of equations for mosquito population dynamics is:

$$\frac{\mathrm{d}A}{\mathrm{d}t} = \phi F\left(1 - \frac{A}{K_a}\right) - \mu_a A - \psi A \tag{9a}$$

$$\frac{\mathrm{d}F}{\mathrm{d}t} = \theta\psi A - \mu_{fu}F\tag{9b}$$

$$\frac{\mathrm{d}M}{\mathrm{d}t} = (1-\theta)\psi A - \mu_{mu}M\tag{9c}$$

The parameters for this model are described in Table 1.

THEOREM 4.1 The zero equilibrium for mosquito population dynamics is LAS when $R_{0u} < 1$, while the steady state is LAS when $R_{0u} > 1$, where $R_{0u} = \frac{\phi \theta \psi}{\mu_{fu}(\mu_a + \psi)}$.

Proof. Jacobian matrix of system (9) is:

$$J = \begin{bmatrix} -\mu_a - \psi - \phi \frac{F}{K_a} & \phi \left(1 - \frac{A}{K_a} \right) & 0\\ \theta \psi & -\mu_{fu} & 0\\ (1 - \theta)\psi & 0 & -\mu_{mu} \end{bmatrix}.$$

The Jacobian matrix for no-infection equilibrium is:

$$J_0 = \begin{bmatrix} -\mu_a - \psi & \phi & 0\\ \theta \psi & -\mu_{fu} & 0\\ (1 - \theta)\psi & 0 & -\mu_{mu} \end{bmatrix}.$$

The characteristic polynomial of J_0 is:

$$(\lambda + \mu_{mu})[(\lambda + \mu_a + \psi)(\lambda + \mu_{fu}) - \phi\theta\psi] = 0$$
(10)

If $R_{0u} < 1$, then all eigenvalues are negative, the zero equilibrium is LAS. The Jacobian matrix at steady state is:

$$J_{ss} = \begin{bmatrix} -\mu_a - \psi - \phi \frac{F^*}{K_a} & \phi \left(1 - \frac{A^*}{K_a} \right) & 0\\ \theta \psi & -\mu_{fu} & 0\\ (1 - \theta)\psi & 0 & -\mu_{mu} \end{bmatrix}.$$

Characteristic polynomial of J_{ss} is:

$$(\lambda + \mu_{mu}) \left[\lambda + \mu_a + \psi + \frac{\phi \theta \psi}{\mu_{fu}} \left(1 - \frac{(\mu_a + \psi)\mu_{fu}}{\phi \theta \psi} \right) \right] (\lambda + \mu_{fu}) - \phi \theta \psi = 0$$
(11)

If $R_{0u} > 1$, then all eigenvalues of J_{ss} are negative, and the non-zero steady state is LAS.

THEOREM 4.2 The zero equilibrium: (0,0,0) is globally asymptotically stable (GAS) when $R_{0u} < 1$. The steady state: $(K_a(1-\frac{1}{R_{0u}}), \frac{K_a\theta\psi}{\mu_{fu}}(1-\frac{1}{R_{0u}}), \frac{K_a(1-\theta)\psi}{\mu_{mu}}(1-\frac{1}{R_{0u}}))$ is GAS when $R_{0u} > 1$.

Proof. We follow the approach in [36]. First we consider the subsystem:

$$\frac{\mathrm{d}A}{\mathrm{d}t} = \phi F\left(1 - \frac{A}{K_a}\right) - \mu_a A - \psi A \tag{12a}$$

$$\frac{\mathrm{d}F}{\mathrm{d}t} = \theta\psi A - \mu_{fu}F\tag{12b}$$

where f = (A, F) is C^1 on an open set $D \subset \mathbb{R}^2$.

Let $\delta_1 = \delta_2 = 1$, $\delta_1 \frac{df_2}{dx_1} = \theta \psi > 0$, $\delta_1 \frac{df_1}{dx_2} = \phi \ge 0$, and $\delta_2 \frac{df_2}{dx_2} = \mu_{fu} > 0$. According to the definition of tridiagonal feedback [36], system 12 is a monotone tridiagonal feedback system with Poincaré-Bendixson property [37].

Recall Theorem 2 in [38]. If the systems of ODEs $dx/dt = f(x), x \subset D$ satisfies:

- (1) The system exists on a compact absorbing set $K \subset D$.
- (2) A unique equilibrium point E exists and is LAS.
- (3) The system has Poincaré-Bendixson property.
- (4) Each periodic orbit of the system is asymptotically stable.

Then E is globally asymptotically stable in D.

To prove that each periodic orbit $\Omega = p(t) : 0 \le t \le w$ of system (12) is asymptotically stable, we follow [39] and Theorem 3 in [36]. We need to prove that the linear system $\frac{dz(t)}{dt} = J_F^{[2]}(p(t))z(t)$ is asymptotically stable, where $J_F^{[2]}$ is the second additive compound matrix of the Jacobian matrix J_F associated with system (12). For system (12), $J_F^{[2]} = -\left(\psi + \mu_a + \frac{\phi}{K_a}F + \mu_{fu}\right)$. We build the following linear system with one equation and the right hand side is the compound matrix of the Jacobian matrix J_F .

$$\frac{\mathrm{d}X}{\mathrm{d}t} = -\left(\psi + \mu_a + \frac{\phi}{K_a}F + \mu_{fu}\right)X.$$

Let Lyapunov function V(X, A, Y) = |X|.

The right derivative of V along the solution paths (X) and (A, F) is: $D_+(V(t) = -(\psi + \mu_a + \frac{\phi}{K_a}F + \mu_{fu})|X|$, which implies that $V(t) \to 0$, and $X(t) \to 0$ as $t \to \infty$. Therefore, the linear system 12 is asymptotically stable, and the solution (A, F) is asymptotically orbitally stable with asymptotic phase.

By the same argument in [36], the system 12 is uniformly persistent in $D \subset \mathbb{R}^2$. The zero equilibrium (0,0,0) is isolated and the largest compact invariant outside D is (A^*, F^*) , which is absorbing and the system 12 is uniformly persistent [40]. The conditions for Theorem 2 in [38] are all satisfied. Therefore, (0,0) is GAS whenever $R_{0u} < 1$, and (A^*, F^*) exists and is GAS when $R_{0u} > 1$.

Following [41] and Theorem 4 in [36], (0,0,0) is GAS whenever $R_{0u} < 1$ and (A^*, F^*, M^*) is GAS whenever $R_{0u} > 1$.

Theorem 4.3

$$\mathcal{D} = \left\{ \begin{pmatrix} A \\ F \\ M \end{pmatrix} \in \mathbb{R}^3 \middle| \begin{array}{c} 0 \le A \le K_a, \\ 0 \le F \le \frac{\psi \theta K_a}{\mu_{fu}}, \\ 0 \le M \le \frac{\psi (1-\theta) K_a}{\mu_{mu}} \end{array} \right\}.$$
(13)

is an invariant region under the flow induced by (9).

Proof. The proof directly follows the proofs for Lemma 4.2 and 4.3 in [42].

References

- R. V. Gibbons, D. W. Vaughn, Dengue: an escalating problem, BMJ (Clinical research ed.) 324 (2002) 1563– 1566.
- [2] J. L. Kyle, E. Harris, Global spread and persistence of dengue, Annual Review of Microbiology 62 (2008) 71–92.
- [3] T. Walker, P. H. Johnson, L. A. Moreira, I. Iturbe-Ormaetxe, F. D. Frentiu, C. J. McMeniman, Y. S. Leong, Y. Dong, J. Axford, P. Kriesner, A. L. Lloyd, S. A. Ritchie, S. L. O'Neill, A. A. Hoffmann, The wMel Wolbachia strain blocks dengue and invades caged Aedes aegypti populations, Nature 476 (2011) 450–453.
- [4] Z. Lam, D. Burke, M. R. Capeding, C. K. Chong, Preparing for introduction of a dengue vaccine: Recommendations from the 1st dengue v2v asia-pacific meeting, Vaccine 29 (2011) 9417–9422.
- [5] M. S. Blagrove, C. Arias-Goeta, A. B. Failloux, S. P. Sinkins, Wolbachia strain wmel induces cytoplasmic incompatibility and blocks dengue transmission in Aedes albopictus, Proceedings of the National Academy of Sciences of the United States of America 109 (2012) 255–260.
- [6] C. J. McMeniman, R. V. Lane, B. N. Cass, A. W. Fong, M. Sidhu, Y. Wang, S. L. O'Neill, Stable introduction of a life-shortening Wolbachia infection into the mosquito Aedes aegypti, Science 323 (2009) 141–144.
- [7] K. Bourtzis, S. L. O'Neill, Wolbachia infections and arthropod reproduction, Bioscience 480 (1998) 287–293.
- [8] S. L. O'Neill, A. Hoffman, J. H. Werren, et al., Influential passengers: inherited microorganisms and arthropod reproduction., Oxford University Press, 1997.
- [9] A. A. Hoffmann, M. Turelli, Cytoplasmic incompatibility in insects, in: S. L. O'Neill, A. A. Hoffmann, J. H. Werren (Eds.), Influential passengers: Inherited microorganisms and arthropod reproduction, Oxford University Press, Oxford, 1997, pp. 42–80.
- [10] E. Suh, S. L. Dobson, Reduced competitiveness of Wolbachia infected Aedes aegypti larvae in intra- and inter-specific immature interactions, Journal of invertebrate pathology 114 (2013) 173–177.
- [11] G. Bian, Y. Xu, P. Lu, Y. Xie, Z. Xi, The endosymbiotic bacterium Wolbachia induces resistance to dengue virus in Aedes aegypti, PloS Neglected Tropical Diseases 6 (2010).
- [12] L. Mousson, K. Zouache, C. Arias-Goeta, V. Raquin, P. Mavingui, A. B. Failloux, The native Wolbachia symbionts limit transmission of dengue virus in Aedes albopictus, PLoS neglected tropical diseases 6 (2012) e1989.
- [13] S. L. Dobson, E. J. Marsland, W. Rattanadechakul, Mutualistic Wolbachia infection in Aedes albopictus: accelerating cytoplasmic drive, Genetics 160 (2002) 1087–1094.
- [14] A. R. Weeks, M. Turelli, W. R. Harcombe, K. T. Reynolds, A. A. Hoffmann, From parasite to mutualist: rapid evolution of Wolbachia in natural populations of drosophila, PLoS biology 5 (2007) e114.
- [15] J. L. Rasgon, L. M. Styer, T. W. Scott, Wolbachia-induced mortality as a mechanism to modulate pathogen transmission by vector arthropods, Journal of medical entomology 40 (2003) 125–132.
- [16] D. M. Watts, D. S. Burke, B. A. Harrison, R. E. Whitmire, A. Nisalak, Effect of temperature on the vector efficiency of Aedes aegypti for dengue 2 virus, The American Journal of Tropical Medicine and Hygiene 36 (1987) 143–152.
- [17] L. A. Moreira, I. Iturbe-Ormaetxe, J. A. Jeffery, G. Lu, A. T. Pyke, L. M. Hedges, B. C. Rocha, S. Hall-Mendelin, A. Day, M. Riegler, L. E. Hugo, K. N. Johnson, B. H. Kay, E. A. McGraw, A. F. van den Hurk, P. A. Ryan, S. L. O'Neill, A Wolbachia symbiont in Aedes aegypti limits infection with dengue, Chikungunya, and Plasmodium, Cell 139 (2009) 1268–1278.
- [18] J. Z. Farkas, P. Hinow, Structured and unstructured continuous models for Wolbachia infections, Bulletin of

mathematical biology 72 (2010) 2067–2088.

- [19] C. Souto-Maior, J. S. Lopes, E. Gjini, C. J. Struchiner, L. Teixeira, M. G. M. Gomes, Heterogeneity in symbiotic effects facilitates Wolbachia establishment in insect populations, Theor Ecol (2014).
- [20] V. A. Jansen, M. Turelli, H. C. Godfray, Stochastic spread of Wolbachia, Proceedings.Biological sciences / The Royal Society 275 (2008) 2769–2776.
- [21] P. A. Hancock, S. P. Sinkins, H. C. Godfray, Population dynamic models of the spread of Wolbachia, The American Naturalist 177 (2011) 323–333.
- [22] M. Turelli, Cytoplasmic incompatibility in populations with overlapping generations, Evolution; international journal of organic evolution 64 (2010) 232–241.
- [23] P. Schofield, Spatially explicit models of turelli-hoffmann Wolbachia invasive wave fronts, Journal of theoretical biology 215 (2002) 121–131.
- [24] H. Hughes, N. F. Britton, Modelling the use of Wolbachia to control dengue fever transmission, Bulletin of mathematical biology 75 (2013) 796–818.
- [25] J. G. Schraiber, A. N. Kaczmarczyk, R. Kwok, M. Park, R. Silverstein, F. U. Rutaganira, T. Aggarwal, M. A. Schwemmer, C. L. Hom, R. K. Grosberg, S. J. Schreiber, Constraints on the use of lifespan-shortening Wolbachia to control dengue fever, Journal of theoretical biology 297 (2012) 26–32.
- [26] P. R. Crain, J. W. Mains, E. Suh, Y. Huang, P. H. Crowley, S. L. Dobson, Wolbachia infections that reduce immature insect survival: predicted impacts on population replacement, BMC evolutionary biology 11 (2011) 290-2148-11-290.
- [27] M. J. Keeling, F. M. Jiggins, J. M. Read, The invasion and coexistence of competing Wolbachia strains, Heredity 91 (2003) 382–388.
- [28] J. Koiller, M. A. da Silva, M. O. Souza, C. Codeco, A. Iggidr, G. Sallet, Aedes, Wolbachia and Dengue, Technical Report, Institute for Research in Computer Science and Automation, 2014.
- [29] M. Z. Ndii, R. I. Hickson, G. N. Mercer, Modelling the introduction of Wolbachia into Aedes aegypti mosquitoes to reduce dengue transmission, ANZIAM J 53 (2012) 213–227.
- [30] P. van den Driessche, J. Watmough, Reproduction numbers and sub-threshold endemic equilibria for compartmental models of disease transmission, Mathematical biosciences 180 (2002) 29–48.
- [31] L. P. Lounibos, R. L. Escher, Sex ratios of mosquitoes from long-term censuses of florida tree holes, Journal of the American Mosquito Control Association 24 (2008) 11–15.
- [32] H. Delatte, G. Gimonneau, A. Triboire, D. Fontenille, Influence of temperature on immature development, survival, longevity, fecundity, and gonotrophic cycles of Aedes albopictus, vector of chikungunya and dengue in the Indian ocean, Journal of medical entomology 46 (2009) 33–41.
- [33] L. C. Monteiro, J. R. de Souza, C. M. de Albuquerque, Eclosion rate, development and survivorship of Aedes albopictus (Skuse)(Diptera: Culicidae) under different water temperatures, Neotropical entomology 36 (2007) 966–971.
- [34] L. M. Styer, S. L. Minnick, A. K. Sun, T. W. Scott, Mortality and reproductive dynamics of Aedes aegypti (Diptera: Culicidae) fed human blood, Vector borne and zoonotic diseases (Larchmont, N.Y.) 7 (2007) 86–98.
- [35] W. C. Marquardt, B. C. Kondratieff, Biology of Disease Vectors, Elsevier Academic Press, Burlington, MA, 2005.
- [36] Y. Dumont, J. M. Tchuenche, Mathematical studies on the sterile insect technique for the Chikungunya disease and Aedes albopictus, Journal of mathematical biology 65 (2012) 809–854.
- [37] J. Mallet-Paret, G. Sell, The poincarébendixson theorem for monotone cyclic feedback systems with delay, J Differ Equ 125 (1996) 441–489.
- [38] M. Li, L. Wang, Global stability in some seir epidemic models, IMA 126 (2002) 295311.
- [39] J. M. Muldowney, Compound matrices and ordinary differential equations, Rocky Mountain J. Math. 20 (1990) 857–872.

- [40] J. Hofbauer, J. W. H. So, Uniform persistence and repellers for maps, Proc AMS 107 (1980) 1137–1142.
- [41] M. Vidyasagar, Decomposing techniques for large-scale systems, IEEE Transactions on automic control AC-25 (1980) 892–898.
- [42] D. Moulay, M. A. Aziz-Alaoui, M. Cadivel, The chikungunya disease: modeling, vector and transmission global dynamics, Mathematical biosciences 229 (2011) 50–63.