

ZERO-ONE LAW OF HAUSDORFF DIMENSIONS OF THE RECURRENT SETS

DONG HAN KIM AND BING LI*

ABSTRACT. Let (Σ, σ) be the one-sided shift space with m symbols and $R_n(x)$ be the first return time of $x \in \Sigma$ to the n -th cylinder containing x . Denote

$$E_{\alpha, \beta}^{\varphi} = \left\{ x \in \Sigma : \liminf_{n \rightarrow \infty} \frac{\log R_n(x)}{\varphi(n)} = \alpha, \limsup_{n \rightarrow \infty} \frac{\log R_n(x)}{\varphi(n)} = \beta \right\},$$

where $\varphi : \mathbb{N} \rightarrow \mathbb{R}^+$ is a monotonically increasing function and $0 \leq \alpha \leq \beta \leq +\infty$. We show that the Hausdorff dimension of the set $E_{\alpha, \beta}^{\varphi}$ admits a dichotomy: it is either zero or one depending on φ, α and β .

Key Words: First return time, Hausdorff dimension.

AMS Subject Classification (2010): 28A80

1. INTRODUCTION

Let $m \geq 2$ be an integer and (Σ, σ) be one-sided shift space with m symbols, more precisely, $\Sigma = \{0, 1, \dots, m-1\}^{\mathbb{N}}$ and $\sigma(x) = (x_{i+1})_{i=1}^{\infty}$ for $x = (x_i)_{i=1}^{\infty} \in \Sigma$. A usual metric d on Σ is given as

$$d(x, y) = m^{-\inf\{k \geq 0 : x_{k+1} \neq y_{k+1}\}}$$

for $x = (x_i)_{i=1}^{\infty}, y = (y_i)_{i=1}^{\infty} \in \Sigma$. For $n \geq 1$ and $x \in \Sigma$, define the first return time of x to the initial word of length n as

$$R_n(x) = \inf\{j \geq 1 : x_{j+1}x_{j+2} \cdots x_{j+n} = x_1x_2 \cdots x_n\},$$

$$R'_n(x) = \inf\{j \geq n : x_{j+1}x_{j+2} \cdots x_{j+n} = x_1x_2 \cdots x_n\}.$$

* Corresponding author.

This work was supported by the National Research Foundation of Korea(NRF) (2012R1A1A2004473), and by NSFC (no. 11201155 and 11371148).

That is,

$$R_n(x) = \inf\{j \geq 1 : \sigma^j(x)|_n = x|_n\} = \inf\{j \geq 1 : d(\sigma^j(x), x) \leq m^{-n}\},$$

where $x|_n$ is the prefix of x with the length n . Both R_n and R'_n are the same except for the case of very short return time. However, R'_n is slightly easier to investigate.

Let ν be any σ -invariant ergodic Borel probability measure on Σ . Ornstein and Weiss [16] proved that for ν -almost all $x \in \Sigma$,

$$\lim_{n \rightarrow \infty} \frac{\log R'_n(x)}{n} = h_\nu(\sigma),$$

where $h_\nu(\sigma)$ denotes the measure-theoretic entropy of ν with respect to σ (see also [9]).

The topic of the first return time of a point in a dynamical system is originated in the famous Poincaré recurrence theorem, (see [7, p. 61]) which states that μ -almost all $x \in X$ is recurrent in the sense

$$\liminf_{n \rightarrow \infty} d(T^n x, x) = 0, \quad (1.1)$$

where $(X, \mathcal{B}, \mu, T, d)$ is a metric measure-preserving dynamical system, by which we mean that (X, d) is a metric space and has a countable base, \mathcal{B} is a sigma-field containing the Borel sigma-field of X and (X, \mathcal{B}, μ, T) is a measure-preserving dynamical system. Boshernitzan [3] improved (1.1) by a quantitative version

$$\liminf_{n \rightarrow \infty} n^{1/\alpha} d(T^n x, x) < \infty$$

for μ -almost every (a.e.) $x \in X$, where α is the dimension of the space in some sense. Let

$$\tau_r(x) = \inf\{n \geq 1 : d(T^n x, x) < r\}$$

and define the lower and upper recurrence rates of $x \in X$ as

$$\underline{R}(x) = \liminf_{r \rightarrow 0} R_r(x), \quad \overline{R}(x) = \limsup_{r \rightarrow 0} R_r(x),$$

where $R_r(x) = \frac{\log \tau_r(x)}{-\log r}$. Barreira and Saussol [2] proved that

$$\underline{R}(x) = \underline{d}_\mu(x), \quad \overline{R}(x) = \overline{d}_\mu(x), \quad \mu - \text{a.e.} \quad (1.2)$$

with the condition that μ has a so-called long return time (see [2]) and $\underline{d}_\mu(x) > 0$ for μ -a.e. x , where $\underline{d}_\mu(x)$, $\overline{d}_\mu(x)$ are the lower and upper pointwise dimensions of μ at $x \in X$ respectively. As a consequence of (1.2), if $\alpha > \underline{d}_\mu(x)$, then

$$\liminf_{n \rightarrow \infty} n^{1/\alpha} d(T^n x, x) = 0,$$

which is a reformulation of Boshernitzan's result. Some other sufficient conditions of (1.2) were given in the literatures (for instance, [19, 20, 21]). The distribution of the first hitting time of the dynamical system is also considered (see [8, 10]).

It is shown by Tan and Wang [23, Theorem 1.3] (see also [11]) that for a positive function ψ defined on \mathbb{N} we have

$$\dim_{\text{H}} \{x \in \Sigma : d(\sigma^n(x), x) < \psi(n) \text{ infinitely often}\} = \frac{1}{1+b}, \quad (1.3)$$

where

$$b = \liminf_{n \rightarrow \infty} \frac{-\log_m \min\{\psi(n), 1\}}{n}.$$

In 2001, Feng and Wu [6] studied the exceptional sets

$$\left\{ x \in \Sigma : \liminf_{n \rightarrow \infty} \frac{\log R'_n(x)}{n} = \alpha, \limsup_{n \rightarrow \infty} \frac{\log R'_n(x)}{n} = \beta \right\}$$

with $0 \leq \alpha \leq \beta \leq \infty$ and proved that those sets are always of Hausdorff dimension one no matter what α and β are (see also [4, 18, 22]). Lau and Shu [12] extended this result to the dynamical systems with specification property by considering the topological entropy instead of Hausdorff dimension. Olsen [14] studied the set of the points for which the set of accumulation points of $\frac{\log R'_n(x)}{n}$ is a given interval for the self-conformal system satisfying a certain separation condition (see also [15]). He proved such set is of full Hausdorff dimension which can be applied to the case of the N -adic transformation with $N \in \mathbb{N}$. Ban and Li [1] generalised this result to β -transformation for any $\beta > 1$ including the cases of full shifts, subshift of finite type, specification, synchronizing etc.

In [6] and [16], the authors considered the recurrence time $R'_n(x)$ with the exponential rate. What will happen if we replace the exponential rate with the polynomial rate? Denote

$$E'_{\alpha,\beta} = \left\{ x \in \Sigma : \liminf_{n \rightarrow \infty} \frac{\log R'_n(x)}{\log n} = \alpha, \limsup_{n \rightarrow \infty} \frac{\log R'_n(x)}{\log n} = \beta \right\}.$$

Peng [17] proved that $\dim_{\text{H}} E'_{\alpha,\beta} = 1$ for any $1 \leq \alpha \leq \beta \leq +\infty$, where \dim_{H} means the Hausdorff dimension of some set.

However, for the short return time case it is worth to distinguish R_n and R'_n . Let $0 \leq \alpha \leq \beta \leq \infty$ and denote

$$E_{\alpha,\beta} = \left\{ x \in \Sigma : \liminf_{n \rightarrow \infty} \frac{\log R_n(x)}{\log n} = \alpha, \limsup_{n \rightarrow \infty} \frac{\log R_n(x)}{\log n} = \beta \right\}.$$

Once the condition $\alpha \geq 1$ fails for R_n , we have $\dim_{\text{H}} E_{\alpha,\beta} = 0$ (see Corollary 1.2), which is the complement of the result in [17]. This result indicates that the Hausdorff dimensions of the recurrence sets with the polynomial rates may drop from the full dimension which is different with that in [6]. Furthermore, such dimensions are either zero or one.

Now we consider the case of general rate. More precisely, let $\varphi : \mathbb{N} \rightarrow \mathbb{R}^+$ be a monotonically increasing function. Denote

$$E_{\alpha,\beta}^{\varphi} = \left\{ x \in \Sigma : \liminf_{n \rightarrow \infty} \frac{\log R_n(x)}{\varphi(n)} = \alpha, \limsup_{n \rightarrow \infty} \frac{\log R_n(x)}{\varphi(n)} = \beta \right\}.$$

We completely calculate the Hausdorff dimensions of the sets $E_{\alpha,\beta}^{\varphi}$ by the following theorem, which is a generalization and complement of [6] and [17].

Write

$$\liminf_{n \rightarrow \infty} \frac{\varphi(n)}{\log n} = \delta \quad \text{and} \quad \limsup_{n \rightarrow \infty} \frac{\varphi(n)}{\log n} = \gamma.$$

Theorem 1.1. *For any $0 \leq \alpha \leq \beta \leq \infty$, we have*

$$\dim_{\text{H}}(E_{\alpha,\beta}^{\varphi}) = \begin{cases} 1, & \text{if } \alpha \geq \frac{1}{\gamma} \text{ and } \beta \geq \frac{1}{\delta}, \\ 0, & \text{otherwise,} \end{cases}$$

with the convention $\frac{1}{0} = \infty$ and $\frac{1}{\infty} = 0$.

Application of Theorem 1.1 to $\varphi(n) = \log n$ implies the following.

Corollary 1.2. *Let $0 \leq \alpha \leq \beta \leq \infty$. Then*

$$\dim_{\text{H}} E_{\alpha, \beta} = \begin{cases} 1 & \text{if } \alpha \geq 1, \\ 0 & \text{if } \alpha < 1. \end{cases}$$

From the results in [6, 17] and Theorem 1.2, when $\varphi(n) = n$ or $\varphi(n) = \log n$, the values of $\dim_{\text{H}}(E_{\alpha, \beta}^{\varphi})$ just depend on α ($\beta \geq \alpha$ is a natural condition since otherwise, the set $(E_{\alpha, \beta}^{\varphi})$ will be empty). The new phenomenon arises for general φ from Theorem 1.1, that is, the values of $\dim_{\text{H}}(E_{\alpha, \beta}^{\varphi})$ may depend on β as well.

2. PROOF OF THE MAIN RESULT

Firstly we prove the zero-dimensional part of Theorem 1.1 by the following proposition, which is a consequence of (1.3).

Proposition 2.1. *Let $\alpha < 1$. Then*

$$\dim_{\text{H}} \left\{ x \in \Sigma : \liminf_{n \rightarrow \infty} \frac{\log R_n(x)}{\log n} \leq \alpha \right\} = 0. \quad (2.1)$$

Proof. For any $0 < \varepsilon < 1 - \alpha$, the condition

$$\liminf_{n \rightarrow \infty} \frac{\log R_n(x)}{\log n} \leq \alpha$$

implies there exist infinitely many n 's such that $R_n(x) < n^{\alpha + \varepsilon}$, that is, $n > R_n(x)^{1/(\alpha + \varepsilon)}$. By the definition of $R_n(x)$, we obtain $d(\sigma^{R_n(x)}(x), x) < m^{-n} < m^{-R_n(x)^{1/(\alpha + \varepsilon)}}$. Therefore the desired sets in (2.1) are the subsets of the set $\left\{ x \in \Sigma : d(\sigma^n(x), x) < m^{-n^{1/(\alpha + \varepsilon)}}$ infinitely often $\right\}$. We complete the proof by the application of (1.3). \square

Proof of the zero-dimensional part of Theorem 1.1. Suppose that $\alpha\gamma < 1$ with $\alpha < \infty$ and $\gamma < \infty$ or $\beta\delta < 1$ with $\beta < \infty$ and $\delta < \infty$. Since

$$\begin{aligned} \liminf_{n \rightarrow \infty} \frac{\log R_n(x)}{\log n} &\leq \left(\liminf_{n \rightarrow \infty} \frac{\log R_n(x)}{\varphi(n)} \right) \cdot \left(\limsup_{n \rightarrow \infty} \frac{\varphi(n)}{\log n} \right) = \alpha\gamma, \\ \liminf_{n \rightarrow \infty} \frac{\log R_n(x)}{\log n} &\leq \left(\limsup_{n \rightarrow \infty} \frac{\log R_n(x)}{\varphi(n)} \right) \cdot \left(\liminf_{n \rightarrow \infty} \frac{\varphi(n)}{\log n} \right) = \beta\delta, \end{aligned}$$

we have

$$\liminf_{n \rightarrow \infty} \frac{\log R_n(x)}{\log n} < 1.$$

By Proposition 2.1, such point set is zero dimensional, which implies that

$$\dim_{\text{H}}(E_{\alpha, \beta}^{\varphi}) = 0. \quad \square$$

Now we concentrate on the proof of the one-dimensional part of Theorem 1.1. The idea is to construct a subset of $E_{\alpha, \beta}^{\varphi}$ with full Hausdorff dimension for different cases of φ, α, β . The following technical lemma provides such subsets with dimension one, which is a generalization of Lemma 1 in [6] and the ideas of proofs are same.

Lemma 2.2. *Let $\{n_i\}_{i \geq 1}$ and $\{\ell_i\}_{i \geq 1}$ be two strictly increasing sequences of natural numbers satisfying the following conditions:*

(i) $\ell_{i+1} \geq \ell_i + n_i + 3$;

(ii) $\lim_{i \rightarrow \infty} \frac{i(n_i+3)}{\ell_i} = 0$.

Then the set

$$A(\{n_i\}, \{\ell_i\}) := \{x \in \Sigma : R_n(x) = \ell_{i+1} \text{ for all large } n \text{ with } n_i < n \leq n_{i+1}\}$$

is of full Hausdorff dimension.

Proof. Let $p > 2$ be a natural number. Define

$$F_p = \{x \in \Sigma : x_j = 0 \text{ for } 1 \leq j \leq p, x_{pk+1} = x_{pk+p} = 1 \text{ for } k \geq 1\}.$$

By the dimensional formula of the self-similar set (see [5, p.130], see also [6]), we know that

$$\dim_{\text{H}} F_p = \frac{p-2}{p}.$$

We will construct a map $g : F_p \rightarrow A(\{n_i\}, \{\ell_i\})$ in the following and show that g^{-1} is nearly Lipschitz on $g(F_p)$, that is, for any $\varepsilon > 0$, there exists $M > 0$ such that $d(g(x), g(y)) < 2^{-k}$ implies $d(x, y) < 2^{-(1-\varepsilon)k}$ for all $k \geq M$.

Let $x \in F_p$ and $x^{(0)} = x$. Find the minimal k_0 such that $\ell_{k_0} - 1 > p$ and let $x^{(1)} = x^{(2)} = \dots = x^{(k_0-1)} = x^{(0)}$. Assume that the sequence

$x^{(k-1)} = (x_i^{(k-1)})$ has been defined ($k \geq k_0$) and now we define $x^{(k)}$. Put the word $w_k(x) = 1(x^{(k-1)}|_{n_k})\overline{x_{n_k+1}^{(k-1)}}1$, where $x^{(k-1)}|_{n_k}$ means the prefix word of $x^{(k-1)}$ with length n_k and $\overline{x_{n_k+1}^{(k-1)}} = x_{n_k+1}^{(k-1)} + 1 \pmod{m}$. Then $x^{(k)}$ is given by inserting $w_k(x)$ in $x^{(k-1)}$ at the place ℓ_k , i.e.,

$$x^{(k)} = x_1^{(k-1)} x_2^{(k-1)} \cdots x_{\ell_k-1}^{(k-1)} w_k(x) x_{\ell_k}^{(k-1)} x_{\ell_k+1}^{(k-1)} \cdots .$$

Since $\{\ell_k\}$ is increasing to infinity and $x^{(k)}|_{\ell_k-1} = x^{(k-1)}|_{\ell_k-1}$, the limit of $\{x^{(k)}\}$ exists, denoted by x^* . Indeed, x^* is obtained by inserting the sequence of words $\{w_k(x)\}$ in $x = x_1^{(0)} x_2^{(0)} \cdots x_n^{(0)} \cdots$.

Now we prove $x^* \in A(\{n_i\}, \{\ell_i\})$. In fact, let $n_i < n \leq n_{i+1}$ for some $i \geq 1$ with $n_i > p$. Since $x^*|_{\ell_{i+1}+n_{i+1}+3} = x^{(i+1)}|_{\ell_{i+1}+n_{i+1}+3}$ (by condition (i)), by the construction of $x^{(i+1)}$, we know that $R_n(x^*) = R_n(x^{(i+1)}) \leq \ell_{i+1}$. It remains to show that $R_n(x^{(i+1)}) < \ell_{i+1}$ can not happen, that is, $x^{(i+1)}|_n$ cannot reappear in $x^{(i+1)}|_{\ell_{i+1}+n-1} = (x^{(i)}|_{\ell_{i+1}-1})1(x^{(i)}|_{n-1})$. In fact, since $x^{(i+1)}|_n$ begins with p consecutive zeros which does not appear in x except at the beginning by the structure of $x \in F_p$, the only possible places where $x^{(i+1)}|_n$ may appear are $w_j(x)$ for some $1 \leq j \leq i$. However, $x^{(i+1)}|_n$ cannot appear in $w_j(x)$ for all $1 \leq j \leq i$ because the maximal length of common prefixes between $x^{(i+1)}|_n$ and $w_j(x)$ is n_j due to the existence of $\overline{x_{n_j+1}^{(j-1)}}$ in $w_j(x)$ and $n > n_j$. Therefore, $R_n(x^*) = \ell_{i+1}$, that is, $x^* \in A(\{n_i\}, \{\ell_i\})$. So we have the map $g : F_p \rightarrow A(\{n_i\}, \{\ell_i\})$ defined as $g(x) = x^*$.

In the following, we show that g^{-1} is nearly Lipschitz on $g(F_p)$. Indeed, suppose $d(x^*, y^*) < m^{-k}$ for some $x, y \in F_p$ and $\ell_i \leq k < \ell_{i+1}$, then $x_1^* x_2^* \cdots x_k^* = y_1^* y_2^* \cdots y_k^*$. Since $x = g^{-1}(x^*)$ is obtained by removing the parts of $w_j(x)$ from x^* and similar for y , it turns out that $x_1 x_2 \cdots x_{k'} = y_1 y_2 \cdots y_{k'}$, where

$$k = k - \sum_{j=1}^i (n_j + 3).$$

Since $\sum_{j=1}^i (n_j + 3) \leq i(n_i + 3)$, by condition (ii) and $\ell_i \leq k$, for any $\varepsilon > 0$, there exists $M > 0$ such that for all $k > M$, we have

$$\sum_{j=1}^i (n_j + 3) \leq i(n_i + 3) < \ell_i \varepsilon < \varepsilon k.$$

So $d(x, y) \leq m^{-k'} < m^{-(1-\varepsilon)k}$. Thus

$$d(g^{-1}(x^*), g^{-1}(y^*)) \leq d(x^*, y^*)^{1-\varepsilon}.$$

Therefore, $\dim_{\mathbb{H}} g(F_p) \geq (1 - \varepsilon) \dim_{\mathbb{H}} F_p$ (see [5, Proposition 2.3]). So $\dim_{\mathbb{H}} g(F_p) \geq \dim_{\mathbb{H}} F_p$ by letting $\varepsilon \rightarrow 0$. Notice that $A(\{n_i\}, \{\ell_i\}) \supset g(F_p)$ and $\dim_{\mathbb{H}} F_p = \frac{p-2}{p}$, we have

$$\dim_{\mathbb{H}} A(\{n_i\}, \{\ell_i\}) \geq \frac{p-2}{p}.$$

We conclude the assertion of the lemma by letting $p \rightarrow \infty$. \square

The proof of the one-dimensional part relies on the applications of Lemma 2.2 by constructing proper sequences $\{n_i\}$ and $\{\ell_i\}$, and verifying the corresponding $A(\{n_i\}, \{\ell_i\})$ is a subset of $E_{\alpha, \beta}^{\varphi}$. For any $x \in A(\{n_i\}, \{\ell_i\})$, since $R_n(x) = \ell_i$ for all $n_{i-1} + 1 \leq n \leq n_i$, we know that

$$\begin{aligned} \liminf_{n \rightarrow \infty} \frac{\log R_n(x)}{\varphi(n)} &= \liminf_{i \rightarrow \infty} \frac{\log \ell_i}{\varphi(n_i)}, \\ \limsup_{n \rightarrow \infty} \frac{\log R_n(x)}{\varphi(n)} &= \limsup_{i \rightarrow \infty} \frac{\log \ell_i}{\varphi(n_{i-1} + 1)}. \end{aligned}$$

So the subsequences $\{n_i\}$ and $\{n_i + 1\}$ are essential for the lower and upper recurrence rates. The following two technical lemmas are needed for the construction of the sequences $\{n_i\}$ and $\{\ell_i\}$.

Lemma 2.3. *For a positive function $\varphi(n)$ and a constant $C \geq 1$ we can choose a sequence of positive integers $\{n_i\}_{i=1}^{\infty}$ satisfying that*

$$\limsup_{i \rightarrow \infty} \frac{\varphi(n_i)}{\log n_i} = \gamma, \quad \liminf_{i \rightarrow \infty} \frac{\varphi(n_i + 1)}{\log(n_i + 1)} = \delta$$

and

$$\lim_{i \rightarrow \infty} \frac{\log n_{i+1}}{\log n_i} = C, \quad \lim_{i \rightarrow \infty} \frac{\log n_i}{i} = \infty.$$

Proof. First, suppose that $C > 1$. Choose $n_1 > e$. For each n_k choose an integer m_k such that

$$\log(\log m_k) - \log(\log n_k) \geq k \log C$$

and

$$\left| \frac{\varphi(m_k)}{\log m_k} - \gamma \right| < \frac{1}{k} \text{ for } 0 \leq \gamma < \infty, \quad \frac{\varphi(m_k)}{\log m_k} > k \text{ for } \gamma = \infty.$$

Denote

$$d_k = \left\lfloor \frac{\log(\log m_k) - \log(\log n_k)}{\log C} \right\rfloor \geq k \quad (2.2)$$

where $\lfloor \cdot \rfloor$ is the floor function. For $0 \leq j \leq d_k$ define r_{k+j} by letting

$$\log(\log r_{k+j}) = \log(\log n_k) + \frac{j}{d_k} \cdot (\log(\log m_k) - \log(\log n_k)). \quad (2.3)$$

Then for $1 \leq j \leq d_k$

$$\log \left(\frac{\log r_{k+j}}{\log r_{k+j-1}} \right) = \frac{\log(\log m_k) - \log(\log n_k)}{d_k}.$$

Since

$$\log C \leq \frac{\log(\log m_k) - \log(\log n_k)}{d_k} < \frac{(d_k + 1) \log C}{d_k} \leq \left(1 + \frac{1}{k}\right) \log C,$$

we have for $1 \leq j \leq d_k$

$$C \leq \frac{\log r_{k+j}}{\log r_{k+j-1}} < C^{1+1/k}. \quad (2.4)$$

Let $n_{k+j} = \lceil r_{k+j} \rceil$ for $1 \leq j \leq d_k$, where $\lceil t \rceil$ is the ceiling function.

Note that $n_{k+d_k} = m_k$.

Next, we choose m'_k such that

$$\log(\log m'_k) - \log(\log n_{k+d_k}) \geq (k + d_k) \log C$$

and

$$\left| \frac{\varphi(m'_k + 1)}{\log(m'_k + 1)} - \delta \right| < \frac{1}{k + d_k}, \quad \text{for } 0 \leq \delta < \infty,$$

$$\frac{\varphi(m'_k + 1)}{\log(m'_k + 1)} > k + d_k, \quad \text{for } \delta = \infty.$$

Then, denote d'_k the same way as in (2.2) (replacing m_k, n_k by m'_k, n_{k+d_k} respectively) and define r_{k+d_k+j} as (2.3) for $1 \leq j \leq d'_k$ and

$n_{k+d_k+j} = \lceil r_{k+d_k+j} \rceil$ for $1 \leq j \leq d'_k$. Note that $n_{k+d_k+d'_k} = m'_k$. We defined $n_{k+1}, n_{k+1}, \dots, n_{k+d_k+d'_k}$ from n_k . Starting from $n_{k+d_k+d'_k}$ we repeat this procedure again. Inductively, we obtain a sequence $\{n_k\}$ by this procedure satisfying

$$\limsup_k \frac{\varphi(n_k)}{\log n_k} = \gamma, \quad \liminf_k \frac{\varphi(n_k + 1)}{\log(n_k + 1)} = \delta.$$

On the other hand, we deduce from (2.4) by letting k to infinity that

$$\lim_{i \rightarrow \infty} \frac{\log r_{i+1}}{\log r_i} = C.$$

Since $r_i \leq n_i < r_i + 1$ and r_i goes to infinity, we get

$$\lim_{i \rightarrow \infty} \frac{\log n_{i+1}}{\log n_i} = C.$$

Also by (2.4) we have

$$\log n_i \geq \log r_i \geq C^{i-1} \log r_1 = C^{i-1} \log n_1.$$

It follows by the assumption of $C > 1$ that

$$\lim_{i \rightarrow \infty} \frac{\log n_i}{i} = \infty.$$

Now assume that $C = 1$. Let $n_1 = 3$. Then $1^2 \leq \log n_1 < 2^2$. For each integer n_k with $k^2 \leq \log n_k < (k+1)^2$, choose an integer m_k such that

$$\log m_k \geq (k+1)^2$$

and

$$\left| \frac{\varphi(m_k)}{\log m_k} - \gamma \right| < \frac{1}{k} \text{ for } 0 \leq \gamma < \infty, \quad \frac{\varphi(m_k)}{\log m_k} > k, \text{ for } \gamma = \infty.$$

Denote

$$d_k = \left\lfloor \sqrt{\log m_k} - k \right\rfloor. \quad (2.5)$$

For each $1 \leq j < d_k$ put

$$n_{k+j} = \left\lceil \exp((k+j)^2) \right\rceil \text{ and } n_{k+d_k} = m_k.$$

Then for all $1 \leq j \leq d_k$

$$(k+j)^2 \leq \log n_{k+j} < (k+j+1)^2. \quad (2.6)$$

Next, we choose m'_k such that

$$\log m'_k \geq (k + d_k + 1)^2$$

and

$$\left| \frac{\varphi(m'_k + 1)}{\log(m'_k + 1)} - \delta \right| < \frac{1}{k + d_k}, \quad \text{for } 0 \leq \delta < \infty,$$

$$\frac{\varphi(m'_k + 1)}{\log(m'_k + 1)} > k + d_k, \quad \text{for } \delta = \infty.$$

Then, denote $d'_k = \lfloor \sqrt{\log m'_k} - (k + d_k) \rfloor$ as before and define $n_{k+d_k+j} = \lfloor \exp((k + d_k + j)^2) \rfloor$ ($1 \leq j < d'_k$) and $n_{k+d_k+d'_k} = m'_k$. Repeating this procedure, we obtain a sequence $\{n_k\}$ satisfying

$$\limsup_k \frac{\varphi(n_k)}{\log n_k} = \gamma, \quad \liminf_k \frac{\varphi(n_k + 1)}{\log(n_k + 1)} = \delta$$

and

$$k^2 \leq \log n_k < (k + 1)^2,$$

which implies that

$$\lim_{i \rightarrow \infty} \frac{\log n_{i+1}}{\log n_i} = 1, \quad \lim_{i \rightarrow \infty} \frac{\log n_i}{i} = \infty. \quad \square$$

Lemma 2.4. *Let $\varphi(n)$ be a positive monotone increasing function which tends to infinity as $n \rightarrow \infty$.*

(i) *We can choose a sequence of positive integers $\{m_i\}_{i=1}^{\infty}$ satisfying that for each $i \geq 1$*

$$\varphi(m_{i+1}) - \varphi(m_i + 1) \leq 3, \quad \varphi(m_{i+1}) - \varphi(m_i) > 1.$$

(ii) *We can choose a sequence of positive integers $\{m_i\}_{i=1}^{\infty}$ satisfying that*

$$\frac{\varphi(m_{i+1})}{\varphi(m_i + 1)} \rightarrow 1 \quad \text{and} \quad \frac{\log m_{i+1}}{\log m_i} \rightarrow 1$$

and for each i , either $m_{i+1} \geq m_i \log m_i$ or $\varphi(m_{i+1}) - \varphi(m_i) > 1$.

Proof. (i) Choose $n_1 \geq 3$. For each n_i we choose n_{i+1} as

$$n_{i+1} = \min\{n : \varphi(n) > \varphi(n_i) + 1\}.$$

and define a sequence $\{m_i\}$ as

$$m_i = \begin{cases} n_i, & \text{if } \varphi(n_{i+1}) - \varphi(n_i) \leq 2, \\ n_{i+1} - 1, & \text{if } \varphi(n_{i+1}) - \varphi(n_i) > 2. \end{cases}$$

Then, we have

$$n_i \leq m_i \leq n_{i+1} - 1 \quad (2.7)$$

and

$$\varphi(n_{i+1} - 1) - \varphi(n_i) \leq 1. \quad (2.8)$$

If $\varphi(n_{i+1}) - \varphi(n_i) \leq 2$, then we have from (2.7) and (2.8)

$$\begin{aligned} \varphi(m_{i+1}) - \varphi(m_i + 1) &\leq \varphi(n_{i+2} - 1) - \varphi(n_i) \\ &= \varphi(n_{i+2} - 1) - \varphi(n_{i+1}) + \varphi(n_{i+1}) - \varphi(n_i) \\ &\leq 1 + 2 = 3 \end{aligned}$$

and

$$\varphi(m_{i+1}) - \varphi(m_i) \geq \varphi(n_{i+1}) - \varphi(n_i) > 1.$$

If $\varphi(n_{i+1}) - \varphi(n_i) > 2$, we have $m_i = n_{i+1} - 1$ so that by (2.8)

$$\varphi(m_{i+1}) - \varphi(m_i + 1) \leq \varphi(n_{i+2} - 1) - \varphi(n_{i+1}) \leq 1$$

and

$$\begin{aligned} \varphi(m_{i+1}) - \varphi(m_i) &\geq \varphi(n_{i+1}) - \varphi(n_{i+1} - 1) \\ &= \varphi(n_{i+1}) - \varphi(n_i) + \varphi(n_i) - \varphi(n_{i+1} - 1) > 1. \end{aligned}$$

(ii) Choose $n_1 \geq 3$. For each n_i we choose n_{i+1} as

$$n_{i+1} = \begin{cases} \min\{n : \varphi(n) > \varphi(n_i) + 1\}, & \text{if } \varphi(\lceil n_i \log n_i \rceil) > \varphi(n_i) + 1, \\ \lceil n_i \log n_i \rceil, & \text{otherwise} \end{cases}$$

and define a sequence $\{m_i\}$ as

$$m_i = \begin{cases} n_i, & \text{if } \varphi(n_{i+1}) - \varphi(n_i) \leq 2, \\ n_{i+1} - 1, & \text{if } \varphi(n_{i+1}) - \varphi(n_i) > 2. \end{cases}$$

Then, we have

$$n_{i+1} < n_i \log n_i + 1, \quad n_i \leq m_i \leq n_{i+1} - 1 \quad (2.9)$$

and

$$\varphi(n_{i+1} - 1) - \varphi(n_i) \leq 1. \quad (2.10)$$

If $\varphi(n_{i+1}) - \varphi(n_i) \leq 2$, then we have from (2.9) and (2.10)

$$\begin{aligned} \varphi(m_{i+1}) - \varphi(m_i + 1) &\leq \varphi(n_{i+2} - 1) - \varphi(n_i) \\ &= \varphi(n_{i+2} - 1) - \varphi(n_{i+1}) + \varphi(n_{i+1}) - \varphi(n_i) \\ &\leq 1 + 2 = 3 \end{aligned}$$

and if $\varphi(n_{i+1}) - \varphi(n_i) > 2$, we have $m_i = n_{i+1} - 1$ so that by (2.10)

$$\varphi(m_{i+1}) - \varphi(m_i + 1) \leq \varphi(n_{i+2} - 1) - \varphi(n_{i+1}) \leq 1.$$

Thus, we have

$$\lim_{i \rightarrow \infty} \frac{\varphi(m_{i+1})}{\varphi(m_i + 1)} = 1.$$

It follows from (2.9) that

$$\begin{aligned} m_{i+1} &\leq n_{i+2} - 1 < n_{i+1} \log n_{i+1} < (n_i \log n_i + 1) \log(n_i \log n_i + 1) \\ &\leq (m_i \log m_i + 1) \log(m_i \log m_i + 1), \end{aligned}$$

thus

$$\lim_{i \rightarrow \infty} \frac{\log m_{i+1}}{\log m_i} = 1.$$

If $\varphi(n_{i+1}) - \varphi(n_i) \leq 2$ and $\varphi(\lceil n_i \log n_i \rceil) > \varphi(n_i) + 1$, then

$$\varphi(m_{i+1}) - \varphi(m_i) \geq \varphi(n_{i+1}) - \varphi(n_i) > 1.$$

If $\varphi(n_{i+1}) - \varphi(n_i) \leq 2$ and $\varphi(\lceil n_i \log n_i \rceil) \leq \varphi(n_i) + 1$, then

$$m_{i+1} \geq n_{i+1} = \lceil n_i \log n_i \rceil = \lceil m_i \log m_i \rceil \geq m_i \log m_i.$$

If $\varphi(n_{i+1}) - \varphi(n_i) > 2$, then, by (2.10), we have

$$\begin{aligned} \varphi(m_{i+1}) - \varphi(m_i) &\geq \varphi(n_{i+1}) - \varphi(n_{i+1} - 1) \\ &= \varphi(n_{i+1}) - \varphi(n_i) + \varphi(n_i) - \varphi(n_{i+1} - 1) > 1. \quad \square \end{aligned}$$

Now we are ready to show the left part of Theorem 1.1.

Proof of the one-dimensional part of Theorem 1.1. Recall

$$\limsup_{n \rightarrow \infty} \frac{\varphi(n)}{\log n} = \gamma, \quad \liminf_{n \rightarrow \infty} \frac{\varphi(n)}{\log n} = \delta.$$

(i) Suppose that $\alpha = \beta = \infty$.

Choose $n_i = i$ and $\ell_i = \max\{e^{i\varphi(i)}, \ell_{i-1} + n_{i-1} + 3, i^2(i+3)\}$, $\ell_0 = 1$. Then $\{n_i\}$ and $\{\ell_i\}$ satisfy the conditions of Lemma 2.2, thus for any $x \in A(\{n_i\}, \{\ell_i\})$, note that $R_n(x) = \ell_i$ for all $n_{i-1} + 1 \leq n \leq n_i$, we get

$$\lim_{n \rightarrow \infty} \frac{\log R_n(x)}{\varphi(n)} \geq \lim_{i \rightarrow \infty} \frac{\log \ell_i}{\varphi(n_i)} \geq \lim_{i \rightarrow \infty} \frac{i\varphi(i)}{\varphi(i)} = \infty.$$

Therefore, $A(\{n_i\}, \{\ell_i\}) \subset E_{\alpha, \beta}^\varphi$.

(ii) Suppose that $\beta = \infty$, $0 \leq \alpha < \infty$ and $\gamma \in [1/\alpha, \infty]$ for $\alpha > 0$, $\gamma = \infty$ for $\alpha = 0$.

We can choose an increasing sequence of positive integers $\{n_i\}$ such that

$$\lim_{i \rightarrow \infty} \frac{\varphi(n_i)}{\log n_i} = \gamma,$$

$$\varphi(n_i) > i\varphi(n_{i-1} + 1), \quad \log n_i > i\varphi(n_{i-1} + 1), \quad \log n_i > \log n_{i-1} + i^2 + 2$$

and if $\gamma = \infty$,

$$\frac{\varphi(n_{i+1})}{\log n_{i+1}} > \frac{\varphi(n_i)}{\log n_i}.$$

Let

$$\ell_i := \begin{cases} \lceil e^{\alpha\varphi(n_i)} \rceil, & \text{if } 0 < \alpha < \infty, \gamma = \infty, \\ \lceil (n_i)^{\alpha\gamma} \log n_i \rceil, & \text{if } 0 < \alpha < \infty, \gamma < \infty \\ \lceil n_i \log n_i \rceil, & \text{if } \alpha = 0, \gamma = \infty. \end{cases}$$

If $0 < \alpha < \infty, \gamma = \infty$, then for i large enough to make $\frac{\varphi(n_{i+1})}{\log n_{i+1}} > \frac{\varphi(n_i)}{\log n_i} > \frac{2}{\alpha}$ and $n_i \geq 3$,

$$\begin{aligned}
\ell_{i+1} - \ell_i &\geq e^{\alpha\varphi(n_{i+1})} - e^{\alpha\varphi(n_i)} - 1 = e^{\alpha\varphi(n_i)} (e^{\alpha(\varphi(n_{i+1})-\varphi(n_i))} - 1) - 1 \\
&> e^{2\log n_i} \left(e^{\alpha \left(\frac{\log n_{i+1}}{\log n_i} \varphi(n_i) - \varphi(n_i) \right)} - 1 \right) - 1 \\
&= (n_i)^2 \left(e^{\frac{\alpha\varphi(n_i)}{\log n_i} (\log n_{i+1} - \log n_i)} - 1 \right) - 1 \\
&> (n_i)^2 (e^{2(\log n_{i+1} - \log n_i)} - 1) - 1 \\
&= (n_{i+1})^2 - (n_i)^2 - 1 = (n_{i+1} - n_i)(n_{i+1} + n_i) - 1 \geq n_i + 3
\end{aligned}$$

and

$$\begin{aligned}
\lim_{i \rightarrow \infty} \frac{i(n_i + 3)}{\ell_i} &\leq \lim_{i \rightarrow \infty} \frac{i(n_i + 3)}{n_i^2} \\
&= \left(\lim_{i \rightarrow \infty} \frac{n_i + 3}{n_i} \right) \cdot \left(\lim_{i \rightarrow \infty} \frac{i}{n_i} \right) = 1 \cdot 0 = 0.
\end{aligned}$$

For other cases of $0 < \alpha < \infty, \gamma < \infty$ and $\alpha = 0, \gamma = \infty$, for large i with $n_i \geq 4$

$$\ell_{i+1} - \ell_i \geq n_i(\log n_{i+1} - \log n_i) - 1 > 2n_i - 1 \geq n_i + 3$$

and

$$\begin{aligned}
\lim_{i \rightarrow \infty} \frac{i(n_i + 3)}{\ell_i} &\leq \lim_{i \rightarrow \infty} \frac{i(n_i + 3)}{n_i \log n_i} \\
&= \left(\lim_{i \rightarrow \infty} \frac{n_i + 3}{n_i} \right) \cdot \left(\lim_{i \rightarrow \infty} \frac{i}{\log n_i} \right) = 1 \cdot 0 = 0.
\end{aligned}$$

Therefore, $\{n_i\}$ and $\{\ell_i\}$ satisfy the conditions of Lemma 2.2. Since

$$\lim_{i \rightarrow \infty} \frac{\log \ell_i}{\varphi(n_i)} = \begin{cases} \alpha, & \text{if } 0 < \alpha < \infty, \gamma = \infty, \\ \lim \frac{\log \ell_i}{\log n_i} \lim \frac{\log n_i}{\varphi(n_i)} = \alpha\gamma \cdot \frac{1}{\gamma} = \alpha, & \text{if } 0 < \alpha < \infty, \gamma < \infty, \\ \lim \frac{\log \ell_i}{\log n_i} \lim \frac{\log n_i}{\varphi(n_i)} = 1 \cdot \frac{1}{\gamma} = 0, & \text{if } \alpha = 0, \gamma = \infty, \end{cases}$$

for any $x \in A(\{n_i\}, \{\ell_i\})$ we get

$$\liminf_{n \rightarrow \infty} \frac{\log R_n(x)}{\varphi(n)} = \liminf_{i \rightarrow \infty} \frac{\log \ell_i}{\varphi(n_i)} = \alpha$$

and

$$\begin{aligned} \limsup_{n \rightarrow \infty} \frac{\log R_n(x)}{\varphi(n)} &= \limsup_{i \rightarrow \infty} \frac{\log \ell_i}{\varphi(n_{i-1} + 1)} \\ &= \begin{cases} \lim \frac{\log \ell_i}{\varphi(n_i)} \lim \frac{\varphi(n_i)}{\varphi(n_{i-1} + 1)} = \alpha \cdot \infty = \infty, & \text{if } 0 < \alpha < \infty, \gamma = \infty, \\ \lim \frac{\log \ell_i}{\log n_i} \lim \frac{\log n_i}{\varphi(n_{i-1} + 1)} = \alpha \gamma \cdot \frac{1}{\gamma} = \alpha, & \text{if } 0 < \alpha < \infty, \gamma < \infty, \\ \lim \frac{\log \ell_i}{\log n_i} \lim \frac{\log n_i}{\varphi(n_{i-1} + 1)} = 1 \cdot \frac{1}{\gamma} = 0, & \text{if } \alpha = 0, \gamma = \infty. \end{cases} \end{aligned}$$

Thus $A(\{n_i\}, \{\ell_i\}) \subset E_{\alpha, \beta}^\varphi$.

In the rest of the proof we only consider the case of finite α, β . Define

$$A := \begin{cases} \alpha\gamma, & \text{if } 0 < \alpha, \gamma < \infty, \\ 1, & \text{if } \alpha = 0, \gamma = \infty, \\ \infty, & \text{if } \alpha > 0, \gamma = \infty, \end{cases} \quad B := \begin{cases} \beta\delta, & \text{if } 0 < \beta, \delta < \infty, \\ 1, & \text{if } \beta = 0, \delta = \infty, \\ \infty, & \text{if } \beta > 0, \delta = \infty. \end{cases}$$

(iii) Suppose that $0 \leq \alpha \leq \beta < \infty$ and $A = B = \infty$, i.e., $0 < \alpha \leq \beta < \infty$ and $\delta = \gamma = \infty$.

Let $\{m_i\}$ be the sequence given by Lemma 2.4 (i), thus $\varphi(m_{i+1}) \leq \varphi(m_i + 1) + 3$ and $\varphi(m_{i+1}) > \varphi(m_i) + 1$.

Let $\tilde{\ell}_1 = e^{\beta\varphi(m_1)}$. For each k with $\tilde{\ell}_k = e^{\beta\varphi(m_k)}$, we will choose k' and define $\tilde{\ell}_{k+1}, \tilde{\ell}_{k+2}, \dots, \tilde{\ell}_{k'}$, where $\tilde{\ell}_{k'} = e^{\beta\varphi(m_{k'})}$. Then by repeating the procedure inductively we define $\{\tilde{\ell}_i\}$. Let $\tilde{\ell}_k = e^{\beta\varphi(m_k)}$, then there exists a minimal integer $d \geq 1$ such that

$$\varphi(m_{k+d}) \geq \left(\frac{2\beta}{\alpha} - 1 \right) \varphi(m_k).$$

Define for $1 \leq j < d$

$$\tilde{\ell}_{k+j} = e^{\frac{2\beta-\alpha}{2}\varphi(m_k) + \frac{\alpha}{2}\varphi(m_{k+j})} > e^{\alpha\varphi(m_{k+j})},$$

and

$$\tilde{\ell}_{k+d} = e^{\alpha\varphi(m_{k+d})}, \quad \tilde{\ell}_{k+d+1} = e^{\beta\varphi(m_{k+d+1})}.$$

Then continue this procedure for $k' = k + d + 1$.

For $0 \leq j < d - 1$

$$\begin{aligned} \tilde{\ell}_{k+j+1} - \tilde{\ell}_{k+j} &= \tilde{\ell}_{k+j} \left(e^{\frac{\alpha}{2}(\varphi(m_{k+j+1}) - \varphi(m_{k+j}))} - 1 \right) \\ &> \tilde{\ell}_{k+j} \frac{\alpha(\varphi(m_{k+j+1}) - \varphi(m_{k+j}))}{2} > \frac{\alpha}{2} \tilde{\ell}_{k+j} \end{aligned}$$

and

$$\begin{aligned} \tilde{\ell}_{k+d} - \tilde{\ell}_{k+d-1} &= \tilde{\ell}_{k+d-1} \left(e^{\alpha\varphi(m_{k+d}) - \frac{2\beta-\alpha}{2}\varphi(m_k) - \frac{\alpha}{2}\varphi(m_{k+d-1})} - 1 \right) \\ &> \tilde{\ell}_{k+d-1} \left(\alpha\varphi(m_{k+d}) - \frac{2\beta-\alpha}{2}\varphi(m_k) - \frac{\alpha}{2}\varphi(m_{k+d-1}) \right) \\ &\geq \tilde{\ell}_{k+d-1} \left(\frac{\alpha}{2}\varphi(m_{k+d}) - \frac{\alpha}{2}\varphi(m_{k+d-1}) \right) > \frac{\alpha}{2} \tilde{\ell}_{k+d-1}, \end{aligned}$$

$$\begin{aligned} \tilde{\ell}_{k+d+1} - \tilde{\ell}_{k+d} &= \tilde{\ell}_{k+d} \left(e^{\beta\varphi(m_{k+d+1}) - \alpha\varphi(m_{k+d})} - 1 \right) \\ &> \tilde{\ell}_{k+d} (\beta\varphi(m_{k+d+1}) - \alpha\varphi(m_{k+d})) > \beta \tilde{\ell}_{k+d} > \frac{\alpha}{2} \tilde{\ell}_{k+d}. \end{aligned}$$

Since $\delta = \infty$, for large i_0 so as to $\varphi(m_i) > \frac{3}{\alpha} \log m_i$ for $i \geq i_0$ we have

$$\tilde{\ell}_{i+1} - \tilde{\ell}_i > \frac{\alpha}{2} \tilde{\ell}_i \geq \frac{\alpha}{2} e^{\alpha\varphi(m_i)} > \frac{\alpha}{2} (m_i)^3,$$

thus for $i \geq i_0$

$$\tilde{\ell}_i > \frac{\alpha}{2} (m_i)^3 \geq \frac{\alpha}{2} i^2 m_i.$$

Thence, $\{m_i\}$ and $\{\ell_i = \lceil \tilde{\ell}_i \rceil\}$ satisfy the conditions of Lemma 2.2.

Thus, for any $x \in A(\{m_i\}, \{\ell_i\})$ we get

$$\liminf_{n \rightarrow \infty} \frac{\log R_n(x)}{\varphi(n)} = \liminf_{i \rightarrow \infty} \frac{\log \ell_i}{\varphi(m_i)} = \alpha$$

and

$$\begin{aligned} \limsup_{n \rightarrow \infty} \frac{\log R_n(x)}{\varphi(n)} &= \limsup_{i \rightarrow \infty} \frac{\log \ell_i}{\varphi(m_{i-1} + 1)} = \limsup_{i \rightarrow \infty} \left(\frac{\log \ell_i}{\varphi(m_i)} \frac{\varphi(m_i)}{\varphi(m_{i-1} + 1)} \right) \\ &= \left(\limsup_{i \rightarrow \infty} \frac{\log \ell_i}{\varphi(m_i)} \right) \left(\lim_{i \rightarrow \infty} \frac{\varphi(m_i)}{\varphi(m_{i-1} + 1)} \right) = \beta \cdot 1 = \beta. \end{aligned}$$

That is, $A(\{m_i\}, \{\ell_i\}) \subset E_{\alpha, \beta}^{\varphi}$.

(iv) Suppose that $0 \leq \alpha \leq \beta < \infty$ and $1 \leq A < B = \infty$.

Since $\delta = \infty$, we deduce that $\gamma = \infty$ and $A = 1$. We can choose a sequence $\{n_i\}$ satisfying that

$$\lim_{i \rightarrow \infty} \frac{\log n_i}{\varphi(n_{i-1} + 1)} = \beta > 0.$$

Note that combined with the assumption $\lim_{i \rightarrow \infty} \frac{\varphi(n_i)}{\log n_i} = \infty$, we get $\frac{\log n_{i+1}}{\log n_i}$ tends to infinity as $i \rightarrow \infty$, which implies that $\frac{i}{\log n_i}$ converges to 0. Let

$$\ell_i = \lceil n_i \log n_i \rceil.$$

Then for large i such that $\log \frac{n_{i+1}}{n_i} \geq 2$ and $n_i \geq 4$

$$\ell_{i+1} - \ell_i \geq n_i(\log n_{i+1} - \log n_i) - 1 > 2n_i - 1 \geq n_i + 3$$

and

$$\begin{aligned} \lim_{i \rightarrow \infty} \frac{i(n_i + 3)}{\ell_i} &\leq \lim_{i \rightarrow \infty} \frac{i(n_i + 3)}{n_i \log n_i} \\ &= \left(\lim_{i \rightarrow \infty} \frac{n_i + 3}{n_i} \right) \cdot \left(\lim_{i \rightarrow \infty} \frac{i}{\log n_i} \right) = 1 \cdot 0 = 0. \end{aligned}$$

Therefore, $\{n_i\}$ and $\{\ell_i\}$ satisfy the conditions of Lemma 2.2. Thus, for any $x \in A(\{n_i\}, \{\ell_i\})$ we get

$$\liminf_{n \rightarrow \infty} \frac{\log R_n(x)}{\varphi(n)} = \liminf_{i \rightarrow \infty} \frac{\log \ell_i}{\varphi(n_i)} = \left(\lim_{i \rightarrow \infty} \frac{\log \ell_i}{\log n_i} \right) \left(\lim_{i \rightarrow \infty} \frac{\log n_i}{\varphi(n_i)} \right) = 1 \cdot 0 = 0$$

and

$$\begin{aligned} \limsup_{n \rightarrow \infty} \frac{\log R_n(x)}{\varphi(n)} &= \limsup_{i \rightarrow \infty} \frac{\log \ell_i}{\varphi(n_{i-1} + 1)} = \limsup_{i \rightarrow \infty} \left(\frac{\log \ell_i}{\log n_i} \frac{\log n_i}{\varphi(n_{i-1} + 1)} \right) \\ &= \left(\lim_{i \rightarrow \infty} \frac{\log \ell_i}{\log n_i} \right) \left(\lim_{i \rightarrow \infty} \frac{\log n_i}{\varphi(n_{i-1} + 1)} \right) = 1 \cdot \beta = \beta. \end{aligned}$$

That is, $A(\{n_i\}, \{\ell_i\}) \subset E_{\alpha, \beta}^{\varphi}$.

(v) Suppose that $0 \leq \alpha \leq \beta < \infty$ and $1 \leq A \leq B < \infty$. By Lemma 2.3 there exists a sequence $\{n_i\}$ satisfying that

$$\limsup_{i \rightarrow \infty} \frac{\varphi(n_i)}{\log n_i} = \gamma, \quad \liminf_{i \rightarrow \infty} \frac{\varphi(n_i + 1)}{\log(n_i + 1)} = \delta, \quad \lim_{i \rightarrow \infty} \frac{\log n_{i+1}}{\log n_i} = \frac{B}{A}.$$

Let

$$\ell_i = \lceil (n_i)^A \log n_i \rceil.$$

Then $\{n_i\}$ and $\{\ell_i\}$ satisfies the conditions of Lemma 2.2.

If $x \in A(\{n_i\}, \{\ell_i\})$, then $R_n(x) = \ell_i$ for each $n_{i-1} < n \leq n_i$. Thus for $x \in A(\{n_i\}, \{\ell_i\})$ we have for $n_{i-1} + 1 \leq n \leq n_i$

$$\frac{\log \ell_i}{\varphi(n_i)} \leq \frac{\log R_n(x)}{\varphi(n)} \leq \frac{\log \ell_i}{\varphi(n_{i-1} + 1)},$$

where the equalities holds for $n = n_i$ and $n = n_{i-1} + 1$ respectively. Therefore, we have

$$\begin{aligned} \limsup_{n \rightarrow \infty} \frac{\log R_n(x)}{\varphi(n)} &= \limsup_{i \rightarrow \infty} \frac{\log \ell_i}{\varphi(n_{i-1} + 1)} \\ &= \limsup_{i \rightarrow \infty} \left(\frac{\log \ell_i}{\log n_i} \cdot \frac{\log n_i}{\log(n_{i-1} + 1)} \cdot \frac{\log(n_{i-1} + 1)}{\varphi(n_{i-1} + 1)} \right) \\ &= \lim_{i \rightarrow \infty} \frac{\log \ell_i}{\log n_i} \cdot \lim_{i \rightarrow \infty} \frac{\log n_i}{\log(n_{i-1} + 1)} \cdot \lim_{i \rightarrow \infty} \frac{\log(n_{i-1} + 1)}{\varphi(n_{i-1} + 1)} \\ &= A \cdot \frac{B}{A} \cdot \frac{1}{\delta} = \frac{B}{\delta} = \beta \end{aligned}$$

and

$$\begin{aligned} \liminf_{n \rightarrow \infty} \frac{\log R_n(x)}{\varphi(n)} &= \liminf_{i \rightarrow \infty} \frac{\log \ell_i}{\varphi(n_i)} = \liminf_{i \rightarrow \infty} \left(\frac{\log \ell_i}{\log n_i} \cdot \frac{\log n_i}{\varphi(n_i)} \right) \\ &= \lim_{i \rightarrow \infty} \frac{\log \ell_i}{\log n_i} \cdot \liminf_{i \rightarrow \infty} \frac{\log n_i}{\varphi(n_i)} = A \cdot \frac{1}{\gamma} = \alpha. \end{aligned}$$

Therefore, $x \in E_{\alpha, \beta}^\varphi$. It follows that $A(\{n_i\}, \{\ell_i\}) \subset E_{\alpha, \beta}^\varphi$.

(vi) Suppose that $0 \leq \alpha \leq \beta < \infty$ and $1 \leq B < A \leq \infty$, i.e., $\alpha > 0$ and $0 < \delta < \infty$.

We may assume that for all n

$$\delta \leq \frac{\varphi(n)}{\log n} \leq \gamma.$$

Since $\beta\delta < \alpha\gamma$ and $\alpha \leq \beta$, we get $\gamma > \delta$. Put

$$C := \begin{cases} \frac{\alpha\gamma - \beta\delta}{\gamma - \delta}, & \text{if } \gamma < \infty, \\ \alpha & \text{if } \gamma = \infty, \end{cases} \quad D := \begin{cases} \frac{(\beta - \alpha)\gamma\delta}{\gamma - \delta}, & \text{if } \gamma < \infty, \\ (\beta - \alpha)\delta & \text{if } \gamma = \infty. \end{cases}$$

and let

$$\rho(x) := Cx + D.$$

Note that, $C > 0$ and $D \geq 0$. Thus $1 \leq \beta\delta \leq \rho(x) \leq \alpha\gamma$ for all $\delta \leq x \leq \gamma$.

Let $\{m_i\}$ be the sequence given by Lemma 2.4 (ii), thus we have

$$\begin{aligned} \rho\left(\frac{\varphi(m_{i+1})}{\log m_{i+1}}\right) \log m_{i+1} - \rho\left(\frac{\varphi(m_i)}{\log m_i}\right) \log m_i \\ \geq C(\varphi(m_{i+1}) - \varphi(m_i)) := \Delta_i. \end{aligned} \quad (2.11)$$

Choose

$$\tilde{\ell}_i = m_i^{\rho(\varphi(m_i)/\log m_i)} \varphi(m_i) \log m_i \quad \text{and} \quad \ell_i = \lfloor \tilde{\ell}_i \rfloor.$$

Then we get

$$\begin{aligned} \tilde{\ell}_{i+1} - \tilde{\ell}_i &\geq m_i^{\rho(\varphi(m_i)/\log m_i)} (e^{\Delta_i} \varphi(m_{i+1}) \log m_{i+1} - \varphi(m_i) \log m_i) \\ &\geq m_i \varphi(m_i) (e^{\Delta_i} \log m_{i+1} - \log m_i) \\ &\geq m_i \varphi(m_i) ((1 + \Delta_i) \log m_{i+1} - \log m_i) \\ &= m_i \varphi(m_i) (\Delta_i \log m_{i+1} + (\log m_{i+1} - \log m_i)), \end{aligned}$$

where the first and the second inequalities are respectively from (2.11) and $\rho(\varphi(m_i)/\log m_i) \geq 1$, and the third holds since $e^x \geq 1 + x$. By Lemma 2.4 (ii),

$$\tilde{\ell}_{i+1} - \tilde{\ell}_i \geq m_i \varphi(m_i) \min(C \log m_{i+1}, \log \log m_i).$$

Thus, for i large enough that $m_i \geq 4$, $\varphi(m_i) \geq 2$, $C \log m_{i+1} \geq 1$ and $\log \log m_i \geq 1$ we get

$$\ell_{i+1} - \ell_i > \tilde{\ell}_{i+1} - \tilde{\ell}_i - 1 \geq 2m_i - 1 \geq m_i + 3.$$

It follows from Lemma 2.4 (ii) that

$$\varphi(m_{i+1}) \log m_{i+1} - \varphi(m_i) \log m_i \geq \min(\varphi(m_i) \log \log m_i, \log m_i),$$

thus for large $i \geq i_0$ with $\log m_{i_0} > \varphi(m_1) \log \log m_{i_0}$ we have

$$\begin{aligned} \varphi(m_i) \log m_i &\geq \varphi(m_{i_0}) \log m_{i_0} + \sum_{j=i_0}^{i-1} \min(\varphi(m_1) \log \log m_j, \log m_j) \\ &\geq \varphi(m_{i_0}) \log m_{i_0} + \varphi(m_1) \sum_{j=i_0}^{i-1} \log \log m_j, \end{aligned}$$

which implies that

$$\begin{aligned} \frac{\tilde{\ell}_i}{im_i} &= \frac{m_i^{\rho(\varphi(m_i)/\log m_i)} \varphi(m_i) \log m_i}{im_i} \geq \frac{\varphi(m_i) \log m_i}{i} \\ &> \frac{\varphi(m_1)}{i} \sum_{j=i_0}^{i-1} \log \log m_j \geq \frac{\varphi(m_1)}{i} \sum_{j=i_0}^{i-1} \log \log j \\ &\geq \frac{\varphi(m_1)}{i} \int_{i_0}^{i-1} \log \log x \, dx \rightarrow \infty \end{aligned}$$

as $i \rightarrow \infty$. Hence, $\{m_i\}$ and $\{\ell_i\}$ satisfy the condition of Lemma 2.2.

By the choice of ℓ_i we have

$$\rho \left(\frac{\varphi(m_i)}{\log m_i} \right) \frac{\log m_i}{\varphi(m_i)} \leq \frac{\log \ell_i}{\varphi(m_i)} \leq \rho \left(\frac{\varphi(m_i)}{\log m_i} \right) \frac{\log m_i}{\varphi(m_i)} + \frac{\log(\varphi(m_i) \log m_i)}{\varphi(m_i)}$$

Using the fact that

$$\frac{\rho(x)}{x} = C + \frac{D}{x}$$

is monotone decreasing for $\delta \leq x \leq \gamma$, we deduce that

$$\alpha = \frac{\rho(\gamma)}{\gamma} \leq \rho \left(\frac{\varphi(m_i)}{\log m_i} \right) \frac{\log m_i}{\varphi(m_i)} \leq \frac{\rho(\delta)}{\delta} = \beta.$$

It follows that

$$\alpha \leq \frac{\log \ell_i}{\varphi(m_i)} \leq \beta + \frac{\log \varphi(m_i) + \log \log m_i}{\varphi(m_i)}$$

If $x \in A(\{m_i\}, \{\ell_i\})$, then for large n satisfying $m_{i-1} < n \leq m_i$, $R_n(x) = \ell_i$, thus

$$\frac{\log \ell_i}{\varphi(m_i)} \leq \frac{\log R_n(x)}{\varphi(n)} \leq \frac{\log \ell_i}{\varphi(m_{i-1} + 1)} = \frac{\log \ell_i}{\varphi(m_i)} \frac{\varphi(m_i)}{\varphi(m_{i-1} + 1)},$$

which implies that

$$\alpha \leq \frac{\log R_n(x)}{\varphi(n)} \leq \left(\beta + \frac{\log \varphi(m_i) + \log \log m_i}{\varphi(m_i)} \right) \frac{\varphi(m_i)}{\varphi(m_{i-1} + 1)}.$$

It follows from the fact that $\varphi(n) \geq (\delta/2) \log n$ for sufficiently large n and $\log x/x \rightarrow 0$ as $x \rightarrow \infty$,

$$\lim_{i \rightarrow \infty} \frac{\log \varphi(m_i)}{\varphi(m_i)} = 0, \quad \lim_{i \rightarrow \infty} \frac{\log \log m_i}{\varphi(m_i)} = 0.$$

Therefore, using Lemma 2.4 (ii) we get

$$\limsup_{n \rightarrow \infty} \frac{\log R_n(x)}{\varphi(n)} \leq \beta \cdot \lim_{i \rightarrow \infty} \frac{\varphi(m_i)}{\varphi(m_{i-1} + 1)} = \beta.$$

Since the sequence $\{m_i\}$ satisfies Lemma 2.4 (ii) and φ is increasing,

$$\begin{aligned} \limsup_{n \rightarrow \infty} \frac{\varphi(n)}{\log n} &= \limsup_{i \rightarrow \infty} \left(\max_{m_{i-1}+1 \leq n \leq m_i} \frac{\varphi(n)}{\log n} \right) \leq \limsup_{i \rightarrow \infty} \frac{\varphi(m_i)}{\log m_{i-1}} \\ &= \limsup_{i \rightarrow \infty} \frac{\varphi(m_i)}{\log m_i} \frac{\log m_i}{\log m_{i-1}} = \limsup_{i \rightarrow \infty} \frac{\varphi(m_i)}{\log m_i} \cdot \lim_{i \rightarrow \infty} \frac{\log m_i}{\log m_{i-1}} \\ &= \limsup_{i \rightarrow \infty} \frac{\varphi(m_i)}{\log m_i} \leq \limsup_{n \rightarrow \infty} \frac{\varphi(n)}{\log n}, \end{aligned}$$

which implies that

$$\limsup_{i \rightarrow \infty} \frac{\varphi(m_i)}{\log m_i} = \limsup_{n \rightarrow \infty} \frac{\varphi(n)}{\log n} = \gamma.$$

Similarly, we get

$$\liminf_{i \rightarrow \infty} \frac{\varphi(m_i)}{\log m_i} = \liminf_{n \rightarrow \infty} \frac{\varphi(n)}{\log n} = \delta.$$

Choose a subsequence $\{m_{i_k}\}$ of $\{m_i\}$ such that $\lim_{k \rightarrow \infty} \frac{\varphi(m_{i_k})}{\log m_{i_k}} = \gamma$. By the continuity of ρ , then

$$\lim_{k \rightarrow \infty} \frac{\log R_{m_{i_k}}(x)}{\varphi(m_{i_k})} = \lim_{k \rightarrow \infty} \frac{\log \ell_{i_k} \log m_{i_k}}{\log m_{i_k} \varphi(m_{i_k})} = \frac{1}{\gamma} \lim_{k \rightarrow \infty} \rho \left(\frac{\varphi(m_{i_k})}{\log m_{i_k}} \right) = \alpha.$$

Hence, we get $\liminf_{n \rightarrow \infty} \frac{\log R_n(x)}{\varphi(n)} = \alpha$ and $\limsup_{n \rightarrow \infty} \frac{\log R_n(x)}{\varphi(n)} = \beta$ in a similar way. Thus $x \in E_{\alpha, \beta}^\varphi$. We have established that $A(\{m_i\}, \{\ell_i\}) \subset E_{\alpha, \beta}^\varphi$.

Therefore, we deduce that

$$A(\{n_i\}, \{\ell_i\}) \subset E_{\alpha, \beta}^\varphi$$

for all cases, which implies that

$$\dim_{\text{H}} E_{\alpha, \beta}^\varphi = 1.$$

by Lemma 2.2. □

REFERENCES

- [1] J.-C. Ban and B. Li, *The multifractal spectra for the recurrence rates of beta-transformations*, J. Math. Anal. Appl. 420 (2014), 1662–1679.
- [2] L. Barreira and B. Saussol, *Hausdorff dimension of measures via Poincare recurrence*, Comm. Math. Phys. 219 (2001), 443–463.
- [3] M. Boshernitzan, *Quantitative recurrence results*, Invent. Math. 113 (1993), 617–631.
- [4] H.-B. Chen, Z.-X. Wen and M. Yu, *The multifractal spectra of certain planar recurrence sets in the continued fraction dynamical system*, J. Math. Anal. Appl. 422 (2015), no. 2, 1264–1276.
- [5] K. J. Falconer, *Fractal geometry: Mathematical foundations and applications*, John Wiley & Sons, Ltd., Chichester, 1990.
- [6] D. J. Feng and J. Wu, *The Hausdorff dimension of recurrent sets in symbolic spaces*, Nonlinearity. 14 (2001), 81–85.
- [7] H. Furstenberg, *Recurrence in Ergodic Theory and Combinatorial Number Theory*, Princeton University Press, Princeton, New Jersey, 1981.
- [8] S. Galatolo, *Dimension and hitting time in rapidly mixing systems*, Math. Res. Lett. 14 (2007), no. 5, 797–805.
- [9] S. Galatolo, D. H. Kim and K. K. Park, *The recurrence time for ergodic systems with infinite invariant measures*, Nonlinearity 19 (2006), no. 11, 2567–2580.
- [10] S. Galatolo and M. J. Pacifico, *Lorenz-like flows: exponential decay of correlations for the Poincaré map, logarithm law, quantitative recurrence*, Ergodic Theory Dynam. Systems 30 (2010), no. 6, 1703–1737.
- [11] R. Hill and S. Velani, *The ergodic theory of shrinking targets*, Invent. Math. 119 (1995) 175–198.
- [12] K. S. Lau and L. Shu, *The spectrum of Poincaré recurrence*, Ergod. Th. Dynam. Sys. 28 (2008), 1917–1943.
- [13] P. Mattila, *Geometry of sets and measures in Euclidean spaces, Fractals and rectifiability*, Cambridge Studies in Advanced Mathematics, 44. Cambridge University Press, Cambridge, 1995.
- [14] L. Olsen, *First return times: multifractal spectra and divergence points*, Discrete Contin. Dyn. Syst. 10 (2004), no. 3, 635–656.
- [15] L. Olsen, *Applications of multifractal divergence points to sets of numbers defined by their N -adic expansion*, Math. Proc. Cambridge Philos. Soc. 136 (2004), no. 1, 139–165.
- [16] D. Ornstein and B. Weiss, *Entropy and data compression schemes*, IEEE Trans. Inform. Theory 39 (1993), 78–83.

- [17] L. Peng, *Dimension of sets of sequences defined in terms of recurrence of their prefixes*, C. R. Math. Acad. Sci. Paris 343 (2006), no. 2, 129–133.
- [18] L. Peng, B. Tan and B.-W. Wang, *Quantitative Poincaré recurrence in continued fraction dynamical system*, Sci. China Math. 55 (2012), no. 1, 131–140.
- [19] J. Rousseau, *Recurrence rates for observations of flows*, Ergodic Theory Dynam. Systems 32 (2012), no. 5, 1727–1751.
- [20] J. Rousseau and B. Saussol, *Poincaré recurrence for observations*, Trans. Amer. Math. Soc. 362 (2010), no. 11, 5845–5859.
- [21] B. Saussol, *Recurrence rate in rapidly mixing dynamical system*, Discrete. Contin. Dyn. Sys. ser.A 15 (2006), 259–267.
- [22] B. Saussol and J. Wu, *Recurrence spectrum in smooth dynamical system*, Nonlinearity. 16 (2003), 1991–2001.
- [23] B. Tan and B.-W. Wang, *Quantitative recurrence properties for beta-dynamical system*, Adv. Math. 228 (2011), no. 4, 2071–2097.

¹DEPARTMENT OF MATHEMATICS EDUCATION, DONGGUK UNIVERSITY-SEOUL, SEOUL, 100-715 KOREA

E-mail address: kim2010@dongguk.edu

²DEPARTMENT OF MATHEMATICS, SOUTH CHINA UNIVERSITY OF TECHNOLOGY, GUANGZHOU, 510641, P.R. CHINA

E-mail address: scbingli@scut.edu.cn