

Lagrangian tetragons and instabilities in Hamiltonian dynamics

Michael Entov^a, Leonid Polterovich^b

January 12, 2021

Abstract

We present a new existence mechanism, based on symplectic topology, for orbits of Hamiltonian flows connecting a pair of disjoint subsets in the phase space. The method involves function theory on symplectic manifolds combined with rigidity of Lagrangian submanifolds. Applications include superconductivity channels in nearly integrable systems and dynamics near a perturbed unstable equilibrium.

Contents

1	Introduction and main results	2
1.1	Interlinking: a chord existence mechanism	2
1.2	Introducing Lagrangian tetragons	4
1.3	Superconductivity channels	7
1.4	Unstable equilibrium	8
1.5	A Poisson bracket invariant	9
1.6	Interlinking and exact Lagrangians	10
2	Preliminaries	13
3	Connecting trajectories of smooth vector fields	14

¹Partially supported by the Israel Science Foundation grant # 1096/14 and by M. & M. Bank Mathematics Research Fund.

²Partially supported by the Israel Science Foundation grant 178/13 and the European Research Council Advanced grant 338809.

4	Poisson bracket invariants	15
5	Lagrangian tetragons	19
5.1	Construction and interlinking of Lagrangian tetragons	20
5.2	Non-exactness of Lagrangians in Liouville manifolds	26

1 Introduction and main results

Given a Hamiltonian flow and a pair of disjoint subsets in the phase space, does there exist an orbit connecting these subsets? Various instances of this question arise in the study of instabilities in Hamiltonian dynamics. In this paper we combine two seemingly remote approaches to detecting such orbits:

- the one based on the Poisson bracket invariants coming from function theory on symplectic manifolds [4];
- a geometric construction due to Mohnke [20], yielding what we call below a *Lagrangian tetragon*, which enabled him at the time to confirm a version of Arnold’s chord conjecture in Reeb dynamics.

The obtained package turns out to be efficient in a number of specific models, including superconductivity channels in nearly integrable systems and dynamics near a perturbed unstable equilibrium that will be discussed below. The proposed existence mechanism for connecting orbits is robust with respect to perturbations of the Hamiltonian in the uniform norm.

1.1 Interlinking: a chord existence mechanism

SETTING THE STAGE. Given an arbitrary smooth (possibly time-dependent) vector field v on a smooth manifold, the piece of its integral trajectory defined over a time interval $[t_0, t_1]$, $t_0 < t_1$, is called a *chord of v* , or a *chord of the flow of v* , of *time-length* $t_1 - t_0$. If such a chord passes at the time t_0 through a set X_0 and at the time t_1 through a set X_1 we say that it is a *chord from X_0 to X_1* .

If the flow of a vector field is defined everywhere for all times, we say that the vector field is *complete*.

Let (M, ω) be a connected symplectic manifold. Consider a smooth time-periodic Hamiltonian $G : M \times \mathbb{S}^1 \rightarrow \mathbb{R}$, where $\mathbb{S}^1 = \mathbb{R}/\mathbb{Z}$. Given a pair of

disjoint compact subsets $X_0, X_1 \subset M$, a (*Hamiltonian*) *chord* of G from X_0 to X_1 is a chord from X_0 to X_1 of the Hamiltonian vector field defined by G . If this vector field is complete, we say that G is *complete*.

SEPARATION. A function $G \in C^\infty(M \times \mathbb{S}^1)$ Δ -*separates* two disjoint compact sets $Y_0, Y_1 \subset M$ if

$$\Delta = \Delta(G; Y_0, Y_1) := \min_{Y_1 \times \mathbb{S}^1} G - \max_{Y_0 \times \mathbb{S}^1} G > 0 .$$

Note that in this definition the order of sets Y_0 and Y_1 is important: G is larger on Y_1 than on Y_0 .

INTERLINKING.

Definition 1.1. Let $(X_0, X_1), (Y_0, Y_1)$ be two pairs of disjoint sets: $X_0 \cap X_1 = Y_0 \cap Y_1 = \emptyset$.

We say that the pair (Y_0, Y_1) κ -*interlinks* the pair (X_0, X_1) , $\kappa > 0$, if every complete (time-dependent) Hamiltonian which Δ -separates Y_0 and Y_1 admits a chord from X_0 to X_1 of time-length $\leq \kappa/\Delta$.

If this property is known to hold only for complete *autonomous* Hamiltonians on M , we say that the pair (Y_0, Y_1) *autonomously* κ -*interlinks* the pair (X_0, X_1) .

Interlinking is the central phenomenon discussed in the present paper. The existence of interlinking pairs is non-obvious and will be discussed below. The interlinking of (X_0, X_1) and (Y_0, Y_1) can be formulated in terms of a Poisson bracket invariant of the quadruple (X_0, X_1, Y_0, Y_1) – see Section 1.5.

We will also discuss a stable version of the interlinking phenomenon. Namely, given a closed connected manifold K , define the K -*stabilization* of a subset $X \subset M$ as is its product $X \times K \subset M \times T^*K$ with the zero-section K of T^*K (the trivial case $K = T^*K = \{\text{a point}\}$ is also allowed; thus, a set can be viewed as a trivial stabilization of itself). We say that a pair (Y_0, Y_1) *stably* κ -*interlinks* the pair (X_0, X_1) if for all $m \in \mathbb{Z}_{\geq 0}$ the pair $(Y_0 \times \mathbb{T}^m, Y_1 \times \mathbb{T}^m)$ κ -interlinks the pair $(X_0 \times \mathbb{T}^m, X_1 \times \mathbb{T}^m)$, where \mathbb{T}^m is the m -dimensional torus.

ROBUSTNESS. Observe that the existence of a Hamiltonian chord of G provided by the interlinking is robust with respect to perturbations of G for which the perturbed Hamiltonian $G + F$ is complete and the perturbation term F is sufficiently small on $Y_0 \cup Y_1$. Indeed,

$$\Delta(G + F; Y_0, Y_1) \geq \Delta(G; Y_0, Y_1) - |\Delta(F; Y_0, Y_1)| . \tag{1}$$

Therefore if (Y_0, Y_1) κ -interlinks (X_0, X_1) and G γ -separates X_0 and X_1 , the perturbed Hamiltonian $G + F$ has a chord from X_0 to X_1 of time-length $\leq \kappa/(\gamma - \delta)$, where $\delta := |\Delta(F; Y_0, Y_1)|$ is assumed to be less than γ .

Let us emphasize that the perturbation F can be arbitrarily large away from $Y_0 \cup Y_1$ and can have large derivatives everywhere – thus the dynamics generated by $G + F$ might be completely different from the dynamics generated by G and the chord of $G + F$ does not have to be close in any sense to the chord of G .

1.2 Introducing Lagrangian tetragons

Next, we illustrate the notion of interlinking for a special class of examples which plays a key role in dynamical applications discussed further in the paper. The following construction originates in the work of Mohnke [20] (for another application of Mohnke’s construction to Hamiltonian dynamics see [19]).

Let (Σ^{2k-1}, ξ) , $k \geq 1$, be a (not necessarily closed) contact manifold with a co-orientable contact structure ξ and let L be a closed connected Legendrian submanifold L of Σ . (If $k = 1$, the contact structure ξ is formed by the zero subspaces of the tangent spaces of Σ and the Legendrian submanifold L is just a point).

Let us make the following additional choices:

(C1) Pick a contact 1-form λ_0 on Σ , $\xi = \ker \lambda_0$ (if $k = 1$ we let λ_0 be any non-vanishing 1-form on Σ). Denote by $\psi_t : \Sigma \rightarrow \Sigma$ the Reeb flow of λ_0 . (Recall that a contact form λ_0 defines a vector field, called Reeb vector field v , by two conditions: $i_v d\lambda_0 = 0$ and $\lambda_0(v) = 1$. The flow of v is called the Reeb flow of λ_0).

(C2) Pick any $T > 0$ such that $\psi_t(L)$ is well-defined and disjoint from L for all $t \in (0, T]$ – such a T exists since L closed and tangent to the contact structure ξ while the Reeb vector field v generating the flow $\{\psi_t\}$ is nowhere tangent to ξ .

(C3) Let $0 < R_0 < R_1$.

(C4) Let K be a closed connected manifold identified with the zero-section of T^*K .

We will view the symplectization of (Σ, ξ) as the symplectic manifold $(\Sigma \times \mathbb{R}_+, d(s\lambda_0))$, where s is the coordinate on the \mathbb{R}_+ -factor of $\Sigma \times \mathbb{R}_+$. Consider the following quadruple of Lagrangian submanifolds (with boundary) $\mathcal{F}, \mathcal{C}, \mathcal{L}, \mathcal{H} \subset \Sigma \times \mathbb{R}_+ \times T^*K$:

$$\begin{aligned} \mathcal{F} &:= \bigcup_{0 \leq t \leq T} \psi_t(L) \times R_0 \times K, & \mathcal{C} &:= \bigcup_{0 \leq t \leq T} \psi_t(L) \times R_1 \times K, \\ \mathcal{L} &:= \psi_T(L) \times [R_0, R_1] \times K, & \mathcal{H} &:= L \times [R_0, R_1] \times K. \end{aligned}$$

Definition 1.2. The union $\Lambda := \mathcal{F} \cup \mathcal{C} \cup \mathcal{L} \cup \mathcal{H}$ is called a *Lagrangian tetragon*. The sets \mathcal{F} and \mathcal{C} are called its *floor* and *ceiling* while \mathcal{L} and \mathcal{H} the *low* and the *high* walls, respectively.

We shall need the following glossary.

Assume that $U \subset \Sigma$ is a domain, $\mathcal{I} \subset \mathbb{R}_+$ is an open interval containing $[R_0, R_1]$ and W is an open tubular neighborhood of K in T^*K , so that $U \times \mathcal{I} \times W$ contains the Lagrangian tetragon Λ . The image of Λ in an arbitrary symplectic manifold (M, ω) under a symplectic embedding $U \times \mathcal{I} \times W \rightarrow M$ will be called a *Lagrangian tetragon in M* . We will view it as a “transplant” of the *original* Lagrangian tetragon $\Lambda \subset \Sigma \times \mathbb{R}_+ \times T^*K$ to (M, ω) . For the remainder of this section we will use the same notation for an original Lagrangian tetragon and for its “transplant”.

A Lagrangian tetragon $\Lambda = \mathcal{F} \cup \mathcal{C} \cup \mathcal{L} \cup \mathcal{H} \subset (M, \omega)$ is called (*stably/autonomously*) κ -*interlinked* (*in (M, ω)*) if the pair $(\mathcal{L}, \mathcal{H})$ (*stably/autonomously*) κ -interlinks the pair $(\mathcal{F}, \mathcal{C})$ in (M, ω) . Roughly speaking, every complete (time-periodic) Hamiltonian on (M, ω) which is larger on the high wall than on the low wall admits a chord from the floor to the ceiling.

Example 1.3 (Prototype example). Take Σ to be the circle $\mathbb{S}^1 = \mathbb{R}/\mathbb{Z}$ equipped with the trivial 0-dimensional contact structure determined by the contact form $\lambda_0 = du$, where u is the coordinate on \mathbb{R} . Fix a point $v \in \mathbb{S}^1$ considered as a Legendrian submanifold of Σ , along with some numbers $0 < R_0 < R_1$ and $0 < T < 1$. Let $K = \{\text{a point}\}$. The corresponding Lagrangian tetragon Λ in the symplectization of Σ is a quadrilateral of area $a = T(R_1 - R_0)$ in the cylinder $\mathbb{S}^1(u) \times \mathbb{R}_+(s)$, with the area form $ds \wedge du$. It was shown in [4] that Λ is stably a -interlinked. Moreover, this result is sharp: Λ is not a' -interlinked with any $a' < a$ (see Remark 5.11).

This example can be extended to higher dimensions in two different ways.

Example 1.4. Let $T^*\mathbb{T}^k = \mathbb{R}^k \times \mathbb{T}^k$, $k > 1$, be the cotangent bundle of the torus with the canonical coordinates $p \in \mathbb{R}^k$ and $q \in \mathbb{T}^k = \mathbb{R}^k/\mathbb{Z}^k$ so that the symplectic structure is given by $dp \wedge dq$. Let $|\cdot|$ denote the Euclidean norm. Consider the contact manifold $\Sigma = \{|p| = 1\} \subset T^*\mathbb{T}^k$ (called the space of co-oriented contact elements to the torus), equipped with the contact form $\lambda_0 = pdq$. The corresponding Reeb flow is simply the Euclidean geodesic flow $(p, q) \rightarrow (p, q + pt)$. The symplectization $\Sigma \times \mathbb{R}_+$ is identified symplectically with $T^*\mathbb{T}^k \setminus \mathbb{T}^k$ by $(p, q, s) \mapsto (sp, q)$. Consider the Lagrangian tetragon Λ in $T^*\mathbb{T}^k$ associated to the Legendrian fiber $\{|p| = 1, q = 0\}$ of Σ , some $0 < R_0 < R_1$, $T > 0$ and $K = \{\text{a point}\}$. It is well-defined provided $0 < T < 1/2$. Observe that its floor, ceiling and the low wall lie in the boundary of the domain

$$E := \{R_0 < |p| < R_1\} \times \{|q| < T\},$$

while the high wall is the central fiber of the fibration $E \rightarrow \{|q| < T\}$.

Theorem 1.5. Λ is stably κ -interlinked with $\kappa = T(R_1 - R_0)$.

For the proof see Section 5.2.

Example 1.6. Consider the unit sphere \mathbb{S}^{2k-1} in the standard symplectic space $\mathbb{R}^{2k} = \mathbb{C}^k$ equipped with the complex coordinates $z = p + iq$ (in the vector notation) and the symplectic form $dp \wedge dq$. The sphere carries a contact structure ξ_{st} given by its field of complex tangencies which is defined by the contact form $\lambda_0 = \frac{1}{2}(pdq - qdp)$. The corresponding Reeb flow is given by $z \mapsto e^{2it}z$. The symplectization $\Sigma \times \mathbb{R}_+$ is identified symplectically with $\mathbb{C}^k \setminus 0$ by $(z, s) \mapsto \sqrt{s}z$. Consider the Lagrangian tetragon Λ in \mathbb{C}^k associated to the Legendrian sphere $\{|p| = 1, q = 0\}$, some numbers $0 < R_0 < R_1$, $T = \pi/4$, and $K = \{\text{a point}\}$. Observe that the high and the low walls of Λ lie in spherical shells $\{R_0^{1/2} \leq |p| \leq R_1^{1/2}, q = 0\}$ and $\{p = 0, R_0^{1/2} \leq |q| \leq R_1^{1/2}\}$ and in the p - and in the q -space, respectively. Its floor and ceiling lie in the $(2k - 1)$ -dimensional spheres in \mathbb{C}^k of radii $R_0^{1/2}$ and $R_1^{1/2}$, respectively.

Theorem 1.7. Λ is stably κ -interlinked with $\kappa = \pi(R_1 - R_0)/4$.

For the proof see Section 5.2.

1.3 Superconductivity channels

Let us now present applications of the results above to specific Hamiltonian systems.

The first application is related to an example that appears in the famous work of Nekhoroshev [23] on the long-term stability of nearly integrable systems¹.

Namely, let ϕ, I denote the standard action-angle coordinates on the cotangent bundle $T^*\mathbb{T}^k = \mathbb{T}^k \times \mathbb{R}^k$ of a torus \mathbb{T}^k (the ϕ coordinates are defined mod 1 and the standard symplectic form on $T^*\mathbb{T}^k$ is written as $dI \wedge d\phi$). Nekhoroshev considered (analytic) Hamiltonians of the form

$$H(\phi, I) = h(I) + \epsilon f(I, \phi), \quad (2)$$

where ϵ is a small parameter, and studied the long-term behavior $I(t)$ of the action variables I along trajectories of the flow generated by H . His main discovery was that if h is a so-called steep function (that is, its restriction to any affine subspace of \mathbb{R}^k has only isolated critical points), then, as long as ϵ is sufficiently small, $I(t)$ stays close to $I(0)$ for exponentially long times. At the same time Nekhoroshev gave an explicit example of a Hamiltonian system of type (2) with two degrees of freedom showing that if h is not steep, then even for small ϵ the actions $I(t)$ may grow linearly fast with t (the so-called “fast diffusion” phenomenon) along certain chords of H projecting to straight intervals in a level set of h in the I -coordinate space (the so-called “superconductivity channels”).

Here we present a multi-dimensional Nekhoroshev-type example where the linear growth phenomenon is robust with respect to perturbations of the Hamiltonian that are C^0 -small on a certain “thin” subset of the phase space (but may be C^1 -large everywhere).

Namely, consider the time-independent Hamiltonian

$$H(p, q, p', q') = \sum_{i=1}^k p_i h_i(p') + U(q) \quad (3)$$

on the symplectic manifold

$$M := T^*\mathbb{T}^k(p, q) \times T^*\mathbb{T}^m(p', q').$$

¹Our interest to this model was triggered by a recent paper [6] by Bounemoura and Kaloshin.

Here the p, p' and q, q' -coordinates correspond, respectively, to the I and ϕ -coordinates above, h_i are arbitrary smooth functions with $h_i(0) = 0$, and U is a non-constant potential. One easily checks that H is complete. If, for instance, all h_i 's are linear, H is quadratic, albeit non-convex, in momenta.

Looking at the Hamiltonian flow of H one readily sees that if $p'(0) = 0$, then $q(t) = q(0)$ for all t and $p(t) = p(0) - \nabla U(q(0)) \cdot t$. Thus choosing $q(0)$ to be a non-critical point of the potential, we see that the increment of the momentum $|p(t) - p(0)|$ grows with linear speed along a straight line (a *superconductivity channel*).

Assume now that the potential U attains a local maximum $\beta := U(0)$ at the point $0 \in \mathbb{T}^k$. Take $r < 1/2$ and assume that $\alpha := \max_{\{|q|=r\}} U < \beta$.

Take the Lagrangian tetragon in $T^*\mathbb{T}^k$ as in Example 1.3, if $k = 1$, and as in Example 1.4, if $k > 1$. Stabilizing it by the zero-section $\mathbb{T}^m \subset T^*\mathbb{T}^m$, we get a Lagrangian tetragon in $M = T^*\mathbb{T}^k \times T^*\mathbb{T}^m$. Denote by $\mathcal{F}, \mathcal{C}, \mathcal{L}, \mathcal{H}$ its floor, ceiling and walls. Observe that H γ -separates the walls \mathcal{L} and \mathcal{H} with $\gamma := \beta - \alpha$. As an immediate consequence of Theorem 1.5 (if $k > 1$) and of the statement at the end of Example 1.3 (if $k = 1$) we get the following result.

Corollary 1.8. *For every complete Hamiltonian $G = H + F$, where $F \in C^\infty(M \times \mathbb{S}^1)$ and $\delta := |\Delta(F; \mathcal{L}, \mathcal{H})| < \gamma$, there exists a Hamiltonian chord of G from \mathcal{F} to \mathcal{C} of time-length $\leq (R_1 - R_0)r/(\gamma - \delta)$. The increment of $|p|$ along this chord equals $R_1 - R_0$.*

1.4 Unstable equilibrium

The simplest model of an unstable equilibrium is provided by the quadratic Hamiltonian $H = \frac{1}{2}(|p|^2 - |q|^2)$ on the standard symplectic vector space $(\mathbb{R}^{2k}, dp \wedge dq)$ (as before, $|\cdot|$ denotes the Euclidean norm). Consider the Lagrangian tetragon $\Lambda \subset \mathbb{R}^{2k}$ described in Example 1.6. Denote by $\mathcal{F}, \mathcal{C}, \mathcal{L}, \mathcal{H}$ its floor, ceiling and walls. Take any point of the form $z = u$, $u \in \mathbb{R}^k$, lying in the sphere $\{|p| = R_0^{1/2}, q = 0\} \subset \mathcal{H}$. By definition, the point $z' = e^{i\pi/4}z$ lies on the floor \mathcal{F} . At the same time one readily sees that z' belongs to the unstable manifold $\{p = q\}$ of the fixed point 0 of the Hamiltonian flow of H . The trajectory of z' has the form $e^t z'$, and hence it eventually hits the ceiling \mathcal{C} . Thus we have found a chord of H from \mathcal{F} to \mathcal{C} .

Observe also that H γ -separates \mathcal{L} and \mathcal{H} with $\gamma = R_0$. Therefore, by Theorem 1.7, a chord connecting \mathcal{F} and \mathcal{C} persists under sufficiently C^0 -small

perturbations of H on \mathcal{L} and \mathcal{H} , yielding the following corollary.

Corollary 1.9. *For every complete Hamiltonian $G = H + F$, where $F \in C^\infty(M \times \mathbb{S}^1)$ and $\delta := |\Delta(F; \mathcal{L}, \mathcal{H})| < R_0$, there is a Hamiltonian chord of G from \mathcal{F} to \mathcal{C} of time-length $\leq \pi(R_1 - R_0)/(4(R_0 - \delta))$. The increment of $|(p, q)|$ along this chord equals $R_1^{1/2} - R_0^{1/2}$.*

Here is another similar setting where Theorem 1.7 can be applied. Let $G : \mathbb{R}^{2k}(p, q) \times \mathbb{S}^1(t) \rightarrow \mathbb{R}$ be a complete Hamiltonian of the form

$$G(p, q, t) = |p|^2/2 + U(q, t).$$

Assume that for some $0 < R_0 < R_1$

$$\max_{\{R_0^{1/2} \leq |q| \leq R_1^{1/2}\} \times \mathbb{S}^1} U =: -\beta \leq 0$$

and that $U(0, t) = 0$ for all $t \in \mathbb{S}^1$. Then G γ -separates \mathcal{L} and \mathcal{H} with $\gamma = R_0/2 + \beta$. Thus, Theorem 1.7 yields the following corollary.

Corollary 1.10. *There exists a Hamiltonian chord of G from \mathcal{F} to \mathcal{C} of time-length bounded from above by $\pi(R_1 - R_0)/(2R_0 + 4\beta)$.*

This corollary can be viewed as a generalization of the known result in the theory of the inverse Lagrange-Dirichlet problem about the instability of the equilibrium point of a mechanical system corresponding to a (non-strict) local maximum of the potential (see [15, 24]) – such an instability follows immediately if the potential U above is taken to be time-independent and with a local maximum $U(0) = 0$ at $q = 0$.

1.5 A Poisson bracket invariant

It has been shown in [4] that the existence of connecting trajectories of Hamiltonian flows can be proved by means of a certain invariant involving the Poisson bracket. In the present paper we refine this techniques in order to establish interlinking for Lagrangian tetragons.

Given a symplectic manifold (M, ω) and a quadruple of compact sets $X_0, X_1, Y_0, Y_1 \subset M$ with $X_0 \cap X_1 = Y_0 \cap Y_1 = \emptyset$, define a number

$$pb_4^+(X_0, X_1, Y_0, Y_1) := \inf \max_M \{F, G\},$$

where the infimum is taken over all compactly supported functions $F, G : M \rightarrow \mathbb{R}$ such that $F|_{X_0} \leq 0$, $F|_{X_1} \geq 1$, $G|_{Y_0} \leq 0$, $G|_{Y_1} \geq 1$. Let us explain the notation: pb stands for the ‘‘Poisson bracket’’, 4 for the number of sets and $+$ for the fact that we are considering the maximum of the Poisson bracket, instead of the uniform norm as it was done in [4].

Write $\widehat{X}_i, \widehat{Y}_i$ for the \mathbb{S}^1 -stabilizations of X_i and Y_i in $M \times T^*\mathbb{S}^1$.

Consider the set Υ (resp., Υ_{aut}) of all $\kappa > 0$ such that the pair (Y_0, Y_1) κ -interlinks (resp., autonomously κ -interlinks) the pair (X_0, X_1) . Let

$$\bar{\kappa} := \inf \Upsilon, \quad \bar{\kappa}_{aut} := \inf \Upsilon_{aut}.$$

If $\Upsilon = \emptyset$ (resp., $\Upsilon_{aut} = \emptyset$), set $\bar{\kappa} := +\infty$ (resp., $\bar{\kappa}_{aut} := +\infty$). One easily checks that $\bar{\kappa}_{aut} \leq \bar{\kappa}$ and $\Upsilon = [\bar{\kappa}, +\infty) \subset \Upsilon_{aut} = [\bar{\kappa}_{aut}, +\infty)$.

Theorem 1.11.

$$1/pb_4^+(X_0, X_1, Y_0, Y_1) = \bar{\kappa}_{aut} \leq \bar{\kappa} \leq 1/pb_4^+(\widehat{X}_0, \widehat{X}_1, \widehat{Y}_0, \widehat{Y}_1).$$

In particular, if $pb_4^+(\widehat{X}_0, \widehat{X}_1, \widehat{Y}_0, \widehat{Y}_1) =: 1/\kappa > 0$, then the pair (Y_0, Y_1) κ -interlinks the pair (X_0, X_1) , and if $pb_4^+(X_0, X_1, Y_0, Y_1) =: 1/\kappa > 0$, then the pair (Y_0, Y_1) autonomously κ -interlinks the pair (X_0, X_1) .

The proof follows the lines of [4] with the following amendments.

1. We adjust the argument of [4] to complete but not necessarily compactly supported Hamiltonians appearing in the definition of interlinking.
2. The results of [4] do not say whether the Hamiltonian chord connecting X_0 and X_1 goes from X_0 to X_1 or from X_1 to X_0 . This is why we introduce a refined version of the Poisson bracket invariant and use a recent theorem of A.Fathi (see Theorem 3.1) to detect Hamiltonian chords going between two sets in a given direction.

1.6 Interlinking and exact Lagrangians

Let us discuss our method of proof of the interlinking of a Lagrangian tetragon. For the sake of transparency, let us focus on the simplest case of autonomous interlinking of a Lagrangian tetragon $\Lambda = \mathcal{F} \cup \mathcal{C} \cup \mathcal{L} \cup \mathcal{H}$ in the symplectization $\Sigma \times \mathbb{R}_+$ of a contact manifold (Σ, ξ) . By Theorem 1.11,

in order to establish the interlinking (respectively, the autonomous interlinking) for a Lagrangian tetragon Λ , it suffices to show the positivity of $\widehat{pb}_4^+(\mathcal{F}, \mathcal{C}, \mathcal{L}, \mathcal{H})$ (respectively, of $pb_4^+(\mathcal{F}, \mathcal{C}, \mathcal{L}, \mathcal{H})$). We will prove this positivity as follows.

Assume that Λ was constructed using a Legendrian submanifold $L \subset (\Sigma, \xi)$, a contact form λ_0 on Σ and real parameters $0 < R_0 < R_1$ and $T > 0$, see (C1)-(C3) in Section 1.2 above².

The Lagrangian tetragon Λ is a singular Lagrangian submanifold with corners. One can smoothen its corners and get a smooth Lagrangian submanifold Λ_ε in $(\Sigma \times \mathbb{R}_+, d(s\lambda_0))$ diffeomorphic to $L \times \mathbb{S}^1$. One easily checks that the Lagrangian isotopy class of Λ_ε in $(\Sigma \times \mathbb{R}_+, d(s\lambda_0))$ depends only on the pair (Σ, L) and not on $\lambda_0, R_0, R_1, T > 0$ and ε .

We will say that the pair (Σ, L) is *exact*, if the above-mentioned Lagrangian isotopy class of Λ_ε in $(\Sigma \times \mathbb{R}_+, d(s\lambda_0))$ contains an exact Lagrangian submanifold – that is, a Lagrangian submanifold Λ' such that $(s\lambda_0)|_{\Lambda'}$ is an exact form. Otherwise (Σ, L) will be called *non-exact*.

Similarly, we will say that the pair (Σ, L) is *stably non-exact*, if for any $m \in \mathbb{Z}_{\geq 0}$ the Lagrangian isotopy class of $\Lambda_\varepsilon \times \mathbb{T}^m$ in the exact symplectic manifold $(\Sigma \times \mathbb{R}_+ \times T^*\mathbb{T}^m, d(s\lambda_0 + pdq))$ does not contain an exact Lagrangian submanifold. (Here pdq is the canonical 1-form on $T^*\mathbb{T}^m$). Clearly, stable non-exactness implies non-exactness.

Theorem 1.12. *Let (Σ, L) be a pair as above and let $\Lambda = \mathcal{F} \cup \mathcal{C} \cup \mathcal{L} \cup \mathcal{H} \subset \Sigma \times \mathbb{R}_+$ be a Lagrangian tetragon constructed using L , a contact form on (Σ, ξ) and some parameters R_0, R_1, T .*

A. If the pair (Σ, L) is non-exact, then

$$pb_4^+(\mathcal{F}, \mathcal{C}, \mathcal{L}, \mathcal{H}) = \left((R_1 - R_0)T \right)^{-1},$$

and thus, by Theorem 1.11, Λ is autonomously $(R_1 - R_0)T$ -interlinked in $(\Sigma \times \mathbb{R}_+, d(s\lambda_0))$.

B. If the pair (Σ, L) is stably non-exact, then

$$\widehat{pb}_4^+(\mathcal{F}, \mathcal{C}, \mathcal{L}, \mathcal{H}) = \left((R_1 - R_0)T \right)^{-1},$$

and thus, by Theorem 1.11, Λ is $(R_1 - R_0)T$ -interlinked in $(\Sigma \times \mathbb{R}_+, d(s\lambda_0))$.

²The manifold K in (C4) is assumed to be the point.

We refer to Section 5.1 for the proof and a more general version of this result. Theorem 1.12 gives rise to the following question.

Question 1.13. *Do exact pairs exist?*

We shall show, by using methods of symplectic topology, that the answer is negative provided (Σ, ξ) is the ideal contact boundary of a Liouville manifold (see Section 5.2 below for the statement of the result and Remarks 5.6 and 5.10 for its generalization to certain non-compact contact manifolds). Note that there do exist contact manifolds whose symplectizations contain *some* exact closed Lagrangian submanifolds (see [22]), but we do not know whether such a Lagrangian submanifold can be constructed as a smoothed Lagrangian tetragon.

Remark 1.14. Incidentally, Theorem 1.12 yields the following result in Reeb dynamics. As above, let $\psi_t : \Sigma \rightarrow \Sigma$ be the Reeb flow of the contact form λ_0 on Σ . Let (Σ, L) be a non-exact pair and assume that, as in the assumption (C2) in the construction of a Lagrangian tetragon, $\psi_t(L)$ is well-defined and disjoint from L for all $t \in (0, T]$ for some $T > 0$. Put $L' := \psi_T(L)$. Let λ be an arbitrary contact form on Σ that defines the same co-orientation of ξ as λ_0 and has a complete Reeb flow.

We claim that the Reeb flow of λ has a chord from L to L' . Moreover, the time-length of this chord does not exceed³ $T \max_Y(\lambda/\lambda_0)$ with $Y := \bigcup_{t \in [0, T]} \psi_t(L)$.

Indeed, look at the Lagrangian tetragon $\Lambda = \mathcal{F} \cup \mathcal{C} \cup \mathcal{L} \cup \mathcal{H}$ associated to L , λ_0 , T and parameters $R =: R_1 > 1 =: R_0 > 0$. Combining the anti-symmetry of the Poisson bracket invariant (see (6) below) with Theorem 1.12 we get that

$$pb_4^+(\mathcal{H}, \mathcal{L}, \mathcal{F}, \mathcal{C}) = ((R - 1)T)^{-1},$$

and hence the pair $(\mathcal{F}, \mathcal{C})$ autonomously κ -interlinks the pair of the walls $(\mathcal{H}, \mathcal{L})$ in $(\Sigma \times \mathbb{R}_+, d(s\lambda_0))$ with $\kappa := (R - 1)T$. The Reeb flow of λ on Σ lifts to a Hamiltonian flow on $(\Sigma \times \mathbb{R}_+, d(s\lambda_0))$ equivariant with respect to the multiplicative \mathbb{R}_+ -action on $\Sigma \times \mathbb{R}_+$ and generated by the Hamiltonian $H := s(\lambda_0/\lambda)$. Choose R large enough and note that H Δ -separates \mathcal{F} and \mathcal{C} with $\Delta := R \min_Y(\lambda_0/\lambda) - \max_Y(\lambda_0/\lambda)$. Hence, for such R , the Reeb flow

³The published version of the paper gives a wrong upper bound on the time-length. We correct this mistake here.

of λ admits a chord of time-length $\leq \kappa/\Delta$ from L to L' . Letting $R \rightarrow +\infty$, we get that the Reeb flow of λ has a chord from L to L' of time-length $\leq T/\min_Y(\lambda_0/\lambda) = T \max_Y(\lambda/\lambda_0)$, which proves the claim. Let us mention that Legendrian contact homology should be a more adequate and powerful technique for detecting Reeb chords connecting L and L' , see e.g. [8, 1].

ORGANIZATION OF THE PAPER: In Section 2 we recall the necessary preliminaries. In Section 3 we formulate a recent result of A.Fathi on the existence of chords for smooth vector fields which is crucial for our studies. In Section 4 we discuss the Poisson bracket invariants and their relation to the existence of Hamiltonian chords. Section 5 is the central section of the paper – in this section we prove the interlinking of Lagrangian tetragons.

2 Preliminaries

Let (M^{2n}, ω) be a connected (not necessarily closed) symplectic manifold.

Given an open set $U \subset M$, set $C_c^\infty(U)$ to be the space of compactly supported smooth functions on M .

Further on we always identify $\mathbb{R}/\mathbb{Z} = \mathbb{S}^1$. Given a Hamiltonian $G : M \times \mathbb{S}^1 = M \times \mathbb{R}/\mathbb{Z} \rightarrow \mathbb{R}$, we denote $G_t := G(\cdot, t)$ and say that G is *compactly supported*, if $\text{supp } G := \bigcup_{t \in \mathbb{S}^1} \text{supp } G_t$ is a compact subset of M . For a bounded $G \in C^\infty(M)$ denote

$$\|G\| := \sup_M |G|.$$

For $G \in C^\infty(M)$ define a vector field $\text{sgrad } G$ by $i_{\text{sgrad } G} \omega = -dG$. Given $F, G \in C^\infty(M)$, define the Poisson bracket $\{F, G\}$ by

$$\begin{aligned} \{F, G\} &:= \omega(\text{sgrad } G, \text{sgrad } F) = dF(\text{sgrad } G) = -dG(\text{sgrad } F) = \\ &= L_{\text{sgrad } G} F = -L_{\text{sgrad } F} G. \end{aligned}$$

The *Hamiltonian flow*, or just a *flow*, of $G : M \times \mathbb{S}^1 \rightarrow \mathbb{R}$ is, by definition, the flow of the (time-dependent) vector field $\text{sgrad } G_t$.

A subset of (M, ω) is called *displaceable*, if it can be completely displaced from its closure by the flow of a compactly supported (time-dependent) Hamiltonian.

We say that a (possibly open) symplectic manifold is of *bounded geometry at infinity*, if it is geometrically bounded in the sense of [3] or convex at

infinity in the sense of [9]. Let us note that for any smooth manifold N the cotangent bundle T^*N , equipped with the standard symplectic structure, is of bounded geometry at infinity.

A symplectic manifold (M, ω) is called *exact*, if ω is exact. A Lagrangian submanifold L of an exact symplectic manifold $(M, d\lambda)$ is called *exact* if the cohomology class $[\lambda|_L] \in H^1(L; \mathbb{R})$ is zero.

3 Connecting trajectories of smooth vector fields

In this section let M be any smooth manifold, v a complete smooth time-independent vector field on M and $X_0, X_1 \subset M$ disjoint compact subsets of M . Denote by $T(X_0, X_1; v)$ the infimum of the time-lengths of the chords of v from X_0 to X_1 (if there is no such chord, set $T(X_0, X_1; v) := +\infty$). Define

$$L_{max}(X_0, X_1; v) := \inf_F \max_M L_v F,$$

where the infimum is taken over all smooth compactly supported functions F on M such that $F|_{X_0} \leq 0$, $F|_{X_1} \geq 1$.

A basic calculus argument shows that

$$T(X_0, X_1; v) \geq 1/L_{max}(X_0, X_1; v).$$

The following theorem has been proved by A.Fathi.

Theorem 3.1 (A.Fathi, [12]). $T(X_0, X_1; v) = 1/L_{max}(X_0, X_1; v)$.

In particular, together with the compactness of X_0 this implies that if $L_{max}(X_0, X_1; v) > 0$ there exists a chord of v from X_0 to X_1 of time-length $1/L_{max}(X_0, X_1; v)$.

Remark 3.2. Replace the maximum of $L_v F$ by $\|L_v F\|$ in the definition of $L_{max}(X_0, X_1; v)$ and call the resulting quantity $L_0(X_0, X_1; v)$. Then, as it was shown in [4, Section 4.1] by a rather basic averaging argument,

$$\min\{T(X_0, X_1; v), T(X_1, X_0; v)\} = 1/L_0(X_0, X_1; v). \quad (4)$$

In other words, if $L_0(X_0, X_1; v) > 0$, then there exists *either* a chord of v from X_0 to X_1 *or* a chord of v from X_1 to X_0 . (Note that none of the results in [4] about the existence of chords says anything about the direction of chords!)

Theorem 3.1 is much more difficult than (4) – in fact, Fathi’s proof of Theorem 3.1 uses, along with the averaging argument similar to the one in [4, Section 4.1], some ingenious arguments from the general theory of metric spaces.

4 Poisson bracket invariants

Let (M, ω) be a connected symplectic manifold.

We say that sets $X_0, X_1, Y_0, Y_1 \subset M$ form an *admissible quadruple*, if they are compact and $X_0 \cap X_1 = Y_0 \cap Y_1 = \emptyset$.

Assume U is an open subset of M and $X_0, X_1, Y_0, Y_1 \subset U$ is an admissible quadruple. Define

$$\begin{aligned} pb_4^U(X_0, X_1, Y_0, Y_1) &:= \inf_{F, G} \|\{F, G\}\|, \\ pb_4^{U,+}(X_0, X_1, Y_0, Y_1) &:= \inf_{F, G} \max_M \{F, G\}, \\ pb_4^{U,-}(X_0, X_1, Y_0, Y_1) &:= \inf_{F, G} (-\min_M \{F, G\}), \end{aligned}$$

where the infimum in all the cases is taken over all $F, G \in C_c^\infty(U)$ such that

$$F|_{X_0} \leq 0, F|_{X_1} \geq 1, G|_{Y_0} \leq 0, G|_{Y_1} \geq 1. \quad (5)$$

One can prove similarly to [4] that this class of pairs (F, G) can be replaced, without changing the infimums, by a smaller class where the inequalities for F, G are replaced by the equalities on some open neighborhoods of the sets X_0, X_1, Y_0, Y_1 .

Let us now define a stabilized version of $pb_4^{U,\pm}$. Identify the cotangent bundle $T^*\mathbb{S}^1$ with the cylinder $\mathbb{R} \times \mathbb{S}^1$ equipped with the coordinates r and $\theta \pmod{1}$ and the standard symplectic structure $dr \wedge d\theta$. As above, write $\widehat{X}_i, \widehat{Y}_i$ for the \mathbb{S}^1 -stabilizations of X_i and Y_i in $M \times T^*\mathbb{S}^1$. Given an open set $U \subset M$ and an admissible quadruple $X_0, X_1, Y_0, Y_1 \subset U$, set

$$\begin{aligned} \widehat{pb}_4^{U,+}(X_0, X_1, Y_0, Y_1) &:= pb_4^{U \times T^*\mathbb{S}^1,+}(\widehat{X}_0, \widehat{X}_1, \widehat{Y}_0, \widehat{Y}_1), \\ \widehat{pb}_4^{U,-}(X_0, X_1, Y_0, Y_1) &:= pb_4^{U \times T^*\mathbb{S}^1,-}(\widehat{X}_0, \widehat{X}_1, \widehat{Y}_0, \widehat{Y}_1). \end{aligned}$$

If $U = M$, or it is clear from the context what X_0, X_1, Y_0, Y_1 are meant, we will omit the corresponding indices and sets in the notation for pb_4, pb_4^\pm and $\widehat{pb}_4, \widehat{pb}_4^\pm$.

The following properties of $pb_4^\pm, \widehat{pb}_4^\pm$ follow easily from the definitions, similarly to the corresponding properties for pb_4 (cf. [4]):

ANTI-SYMMETRY:

$$pb_4^{U,+}(X_0, X_1, Y_0, Y_1) = pb_4^{U,-}(X_1, X_0, Y_0, Y_1) = pb_4^{U,-}(X_0, X_1, Y_1, Y_0) = \quad (6)$$

$$= pb_4^{U,-}(Y_0, Y_1, X_0, X_1) = pb_4^{U,+}(Y_1, Y_0, X_0, X_1),$$

$$pb_4 \geq \max\{pb_4^+, pb_4^-\}. \quad (7)$$

Similar claims hold also for $\widehat{pb}_4^{U,\pm}$.

BEHAVIOR UNDER PRODUCTS:

Suppose that M and N are connected symplectic manifolds. Equip $M \times N$ with the product symplectic form. Let $K \subset N$ be a compact subset. Then for every collection X_0, X_1, Y_0, Y_1 of compact subsets of M

$$pb_4^{M \times N, \pm}(X_0 \times K, X_1 \times K, Y_0 \times K, Y_1 \times K) \leq pb_4^{M, \pm}(X_0, X_1, Y_0, Y_1) \quad (8)$$

and a similar claim holds also for \widehat{pb}_4^\pm .

In particular, for any $U \subset M, X_0, X_1, Y_0, Y_1 \subset U$,

$$\widehat{pb}_4^{U, \pm}(X_0, X_1, Y_0, Y_1) \leq pb_4^{U, \pm}(X_0, X_1, Y_0, Y_1).$$

MONOTONICITY: Assume $U \subset W$ are opens sets in M and $X'_0, X'_1, Y'_0, Y'_1 \subset U \subset W$ is an admissible quadruple. Let X_0, X_1, Y_0, Y_1 be another admissible quadruple such that $X_0 \subset X'_0, X_1 \subset X'_1, Y_0 \subset Y'_0, Y_1 \subset Y'_1$. Then

$$pb_4^{W, \pm}(X_0, X_1, Y_0, Y_1) \leq pb_4^{U, \pm}(X'_0, X'_1, Y'_0, Y'_1). \quad (9)$$

A similar inequality holds also for \widehat{pb}_4^\pm .

SEMI-CONTINUITY:

Suppose that a sequence $X_0^{(j)}, X_1^{(j)}, Y_0^{(j)}, Y_1^{(j)}, j \in \mathbb{N}$, of ordered collections converges (in the sense of the Hausdorff distance between sets) to a collection X_0, X_1, Y_0, Y_1 . Then

$$\limsup_{j \rightarrow +\infty} pb_4^\pm(X_0^{(j)}, X_1^{(j)}, Y_0^{(j)}, Y_1^{(j)}) \leq pb_4^\pm(X_0, X_1, Y_0, Y_1). \quad (10)$$

A similar inequality holds also for \widehat{pb}_4^\pm .

Proof of Theorem 1.11. Let us prove that

$$1/pb_4^+(X_0, X_1, Y_0, Y_1) = \bar{\kappa}_{aut}. \quad (11)$$

To prove that

$$1/pb_4^+(X_0, X_1, Y_0, Y_1) \leq \bar{\kappa}_{aut}. \quad (12)$$

we need to prove that

$$pb_4^+(X_0, X_1, Y_0, Y_1) \geq 1/\kappa \quad (13)$$

for any $\kappa \in \Upsilon_{aut}$. Pick such a $\kappa \in \Upsilon_{aut}$ and any $F, G \in C_c^\infty(M)$ satisfying (5). Then the pair (Y_0, Y_1) autonomously κ -interlinks the pair (X_0, X_1) and, since G 1-separates Y_0 and Y_1 , this means that there exists a chord of G from X_0 to X_1 of time-length $\leq \kappa$. Restricting F to the chord and applying the mean value theorem from the basic calculus one readily obtains that the function $L_{\text{sgrad } G} F = \{F, G\}$ takes a value greater or equal to $1/\kappa$ at some point of the chord and thus $\max_M \{F, G\} \geq 1/\kappa$. Since this holds for any $F, G \in C_c^\infty(M)$ satisfying (5), we obtain (13) and hence (12).

Let us prove that

$$k_{aut} := 1/pb_4^+(X_0, X_1, Y_0, Y_1) \leq \bar{\kappa}_{aut}. \quad (14)$$

Equivalently, this means to prove the following: Let $G : M \rightarrow \mathbb{R}$ be a complete Hamiltonian that Δ -separates Y_0 and Y_1 . We need to show that there exists a chord of G from X_0 to X_1 of time-length $\leq \kappa_{aut}/\Delta$.

We may assume without loss of generality that $\max_{Y_0} G = 0$, $\min_{Y_1} G = 1$, $\Delta = 1$ (the general case is reduced to this one if one replaces G with $u \circ G$ for an appropriate function $u : \mathbb{R} \rightarrow \mathbb{R}$). Let g_t be the flow of G . Since the flow g_t is defined for all $t \in \mathbb{R}$, the following subset of M is well-defined and compact:

$$\Theta_{aut} := \bigcup_{0 \leq t \leq \kappa_{aut}} g_t(X_0) \cup Y_0 \cup Y_1.$$

Let $G' : M \rightarrow \mathbb{R}$ be a compactly supported Hamiltonian which coincides with G on an open neighborhood of Θ_{aut} . Since G coincides with G' on Y_0 and Y_1 , we get that

$$\max_{Y_0} G = \max_{Y_0} G' = 0, \quad \min_{Y_1} G = \min_{Y_1} G' = 1.$$

Since

$$\kappa_{aut} = 1/pb_4^+(X_0, X_1, Y_0, Y_1) > 0,$$

we get that

$$\begin{aligned} \max_M \{F', G'\} &= \max_M L_{\text{sgrad } G'} F' \geq \\ &\geq L_{\text{max}}(X_0, X_1; \text{sgrad } G') \geq pb_4^+(X_0, X_1, Y_0, Y_1) = 1/\kappa_{aut} \end{aligned}$$

for any compactly supported $F' : M \rightarrow \mathbb{R}$ such that $F'|_{X_0} \leq 0$, $F'|_{Y_1} \geq 1$. Therefore, by Theorem 3.1, there exists a chord of G' from X_0 to X_1 of time-length $\leq \kappa_{aut}$. This chord of G' is also a chord of G – indeed, G' coincides with G on a neighborhood of Θ_{aut} and therefore for any $t \in [0, \kappa_{aut}]$ the time-[0, t] flows of G and G' coincide on X_0 .

Thus we have proved the existence of a chord of G from X_0 to X_1 of time-length $\leq \kappa_{aut}$. This finishes the proof of (14). Together (12) and (14) imply (11).

Let us now prove that

$$\bar{\kappa} \leq \kappa := 1/pb_4^+(\widehat{X}_0, \widehat{X}_1, \widehat{Y}_0, \widehat{Y}_1). \quad (15)$$

Equivalently, this means to prove the following: Let $G : M \times \mathbb{S}^1 \rightarrow \mathbb{R}$ be a complete Hamiltonian that Δ -separates Y_0 and Y_1 . We need to show that there exists a chord of G from X_0 to X_1 of time-length $\leq \kappa/\Delta$.

The argument is very similar to the argument used above in the autonomous case. Namely, consider an *autonomous* Hamiltonian

$$H : M \times T^*\mathbb{S}^1 \rightarrow \mathbb{R}, \quad (x, r, \theta) \rightarrow G(x, \theta) + r,$$

generating a Hamiltonian flow $h_t : M \times T^*\mathbb{S}^1 \rightarrow M \times T^*\mathbb{S}^1$. Note that the projection $M \times T^*\mathbb{S}^1 \rightarrow M$ maps each trajectory of h_t to a trajectory of g_t of the same time-length. Thus, it suffices to prove the existence of a chord of H from \widehat{X}_0 to \widehat{X}_1 of time-length $\leq \kappa/\Delta$.

The Hamiltonian flow h_t of H is defined for all times, since so is the Hamiltonian flow g_t of G . Note that $r = 0$ on $\widehat{Y}_0, \widehat{Y}_1$. Therefore

$$\max_{\widehat{Y}_0} H = \max_{Y_0 \times \mathbb{S}^1} G, \quad \min_{\widehat{Y}_1} H = \min_{Y_1 \times \mathbb{S}^1} G,$$

meaning that H Δ -separates \widehat{Y}_0 and \widehat{Y}_1 . Thus, H is a complete autonomous Hamiltonian on $M \times T^*\mathbb{S}^1$ that Δ -separates \widehat{Y}_0 and \widehat{Y}_1 . By the result in the autonomous case, H has a chord from \widehat{X}_0 to \widehat{X}_1 of time-length $\leq \kappa/\Delta$, as required. This finishes the proof of (15) and of the theorem. \square

5 Lagrangian tetragons

The proof of the positivity of \widehat{pb}_4 for Lagrangian tetragons relies on the following proposition which is based on a method from [4] relating deformations of symplectic forms and Poisson brackets.

Proposition 5.1. *Let $(M, \omega = d\lambda)$ be a connected (possibly open) exact symplectic manifold with a fixed primitive λ and let $\Lambda \subset M$ be a closed Lagrangian submanifold. Let $F, G \in C_c^\infty(M)$. Assume*

1. *(M, ω) does not admit exact closed Lagrangian submanifolds Lagrangian isotopic to Λ .*
2. *The form FdG is closed on Λ .*
3. *There exists a number $c > 0$ so that the cohomology class of FdG on Λ satisfies $[FdG|_\Lambda] = -c[\lambda|_\Lambda]$.*

Then $\max_M\{F, G\} \geq c$.

Proof of Proposition 5.1. Following [4], consider the deformation

$$\omega_\tau := \omega + \tau dF \wedge dG, \quad \tau \in \mathbb{R}.$$

A direct calculation shows that

$$dF \wedge dG \wedge \omega^{n-1} = -\frac{1}{n}\{F, G\}\omega^n.$$

and thus

$$\omega_\tau^n = (1 - \tau\{F, G\})\omega^n.$$

Thus ω_τ is symplectic for any $\tau \in I := [0, 1/\max_M\{F, G\})$.

For any $\tau \in I$ the form ω can be mapped (using Moser's method [21]) to ω_τ by a compactly supported isotopy $\vartheta_\tau : (M, \omega_\tau) \rightarrow (M, \omega)$. Note that, by condition 2, Λ is Lagrangian with respect to ω_τ . Therefore the manifold $\Lambda_\tau := \vartheta_\tau(\Lambda)$ is a Lagrangian submanifold of (M, ω) Lagrangian isotopic to Λ .

Assume, by contradiction, that $\max_M\{F, G\} < c$, i.e. $a := 1/c \in I$. As we noted above Λ_a is a Lagrangian submanifold with respect to ω Lagrangian isotopic to L . We claim that Λ_a , and hence Λ , is Lagrangian isotopic to an exact Lagrangian submanifold of (M, ω) – this would yield a contradiction with condition 1 and thus prove that, in fact, $\max_M\{F, G\} \geq c$.

Indeed, by condition 3, Λ is exact with respect to the primitive $\lambda_a := \lambda + aFdG$ of ω_a . Therefore $\Lambda_a = \vartheta_a(\Lambda)$ is exact with respect to the primitive form $\alpha := (\vartheta_a^*)^{-1}\lambda_a$ of $d\alpha = \omega$. Since the isotopy ϑ_τ and the functions F, G are compactly supported, we get that $\beta := \alpha - \lambda$ is a closed compactly supported 1-form on M .

Let $f_t : M \rightarrow M$ be the locally Hamiltonian flow of β generated by the vector field v with $i_v\omega = \beta$. Since β is compactly supported, f_t is defined for all t . Also, since f_t is a locally Hamiltonian flow, each f_t is a symplectomorphism of (M, ω) . Observe that for all t

$$\frac{d}{dt}f_t^*\lambda = f_t^*L_v\lambda = d(f_t^*(i_v\lambda)) + f_t^*\beta$$

and

$$f_t^*\beta = \beta.$$

Therefore for all t

$$\frac{d}{dt}[f_t^*\lambda] = [\beta],$$

which yields

$$[f_1^*\lambda|_{\Lambda_a}] = [\lambda|_{\Lambda_a}] + [\beta|_{\Lambda_a}] = [\alpha|_{\Lambda_a}] = 0.$$

Thus $f_1(\Lambda_a)$ is an exact Lagrangian submanifold of (M, ω) with respect to λ , and $f_t(\Lambda_a)$, $t \in [0, 1]$, is the desired Lagrangian isotopy connecting $\Lambda_a = f_0(\Lambda_a)$ to $f_1(\Lambda_a)$. This finishes the proof of the claim and of the proposition. \square

5.1 Construction and interlinking of Lagrangian tetragons

Let (Σ^{2k-1}, ξ) , $k \geq 1$, be a connected (not necessarily closed) contact manifold with a co-orientable contact structure ξ (if $k = 1$, the contact structure ξ is formed by the zero subspaces of the tangent spaces). Let L be a closed connected Legendrian submanifold of Σ (in the case $k = 1$ the submanifold L is just a point).

Pick a contact form λ_0 on Σ : $\xi = \{\lambda_0 = 0\}$. (In the case $k = 1$ let λ_0 be any non-vanishing form on Σ). Denote by $\psi_t : \Sigma \rightarrow \Sigma$ the Reeb flow of λ_0 . (In the case $k = 1$ the Reeb vector field v is defined just by the condition

$\lambda_0(v) \equiv 1$). Pick $T > 0$ so that $\psi_t(L)$ is well-defined and disjoint from L for all $t \in (0, T]$. Let $0 < R_0 < R_1$.

Let $(\Sigma \times \mathbb{R}_+, d(s\lambda_0))$ be the symplectization of (Σ, ξ) (here s is the coordinate on the \mathbb{R}_+ -factor of $\Sigma \times \mathbb{R}_+$).

Define a *Lagrangian tetragon* $\Lambda'' \subset \Sigma \times \mathbb{R}_+$:

$$\begin{aligned} \mathcal{F}'' &:= \bigcup_{0 \leq t \leq T} \psi_t(L) \times R_0, & \mathcal{C}'' &:= \bigcup_{0 \leq t \leq T} \psi_t(L) \times R_1, \\ \mathcal{L}'' &:= \psi_T(L) \times [R_0, R_1], & \mathcal{H}'' &:= L \times [R_0, R_1], \\ \Lambda'' &:= \mathcal{F}'' \cup \mathcal{C}'' \cup \mathcal{L}'' \cup \mathcal{H}''. \end{aligned}$$

This is a singular Lagrangian submanifold. We will define its smoothening as follows.

Consider an embedding $\Phi : L \times (\mathbb{R}_+ \times [0, T]) \rightarrow \Sigma \times \mathbb{R}_+$ given by

$$\Phi(x \times (s, t)) := (\psi_t(x), s).$$

Then for any smooth embedded loop $\gamma \subset [R_0, R_1] \times [0, T]$ the restriction of Φ to $L \times \gamma$ is a Lagrangian embedding. Choose a family γ_ε of smooth embedded loops in $[R_0, R_1] \times [0, T]$ C^0 -converging to the boundary of the rectangle $[R_0, R_1] \times [0, T]$ as $\varepsilon \rightarrow 0$ and denote by

$$\Lambda''_\varepsilon := \Phi(L \times \gamma_\varepsilon) \subset \Sigma \times \mathbb{R}_+$$

the resulting smooth Lagrangian submanifold, called a *smoothened Lagrangian tetragon* in $\Sigma \times \mathbb{R}_+$.

Let K be a closed connected manifold. Then

$$\Lambda' := \Lambda'' \times K$$

is a *Lagrangian tetragon* in $\Sigma \times \mathbb{R}_+ \times T^*K$ and $\Lambda'_\varepsilon := \Lambda''_\varepsilon \times K$ is its smoothening.

Assume now that U is a domain in Σ , $\mathcal{I} \subset \mathbb{R}_+$ is an open interval containing $[R_0, R_1]$ and W is an open tubular neighborhood of K in T^*K so that Λ' is contained in $U \times \mathcal{I} \times W$. Denote by η the standard 1-form on T^*K . Let $(M, \omega = d\lambda)$ be an exact connected symplectic manifold and let $\nu : U \times \mathcal{I} \times W \rightarrow M$ be a symplectic embedding such that

$$\nu^*\lambda = s\lambda_0 \oplus \eta. \tag{16}$$

Denote by $\Lambda := \nu(\Lambda')$ the resulting *Lagrangian tetragon* in M and let

$$\mathcal{F} := \nu(\mathcal{F}'' \times K), \mathcal{C} := \nu(\mathcal{C}'' \times K), \mathcal{L} := \nu(\mathcal{L}'' \times K), \mathcal{H} := \nu(\mathcal{H}'' \times K),$$

be, respectively, its floor, ceiling, low wall and high wall. Let $\Lambda_\varepsilon := \nu(\Lambda'_\varepsilon)$ be its smoothening.

Note that for sufficiently small ε all Lagrangian submanifolds Λ_ε are Lagrangian isotopic in (M, ω) . From this point on we consider only such ε .

Theorem 5.2. *Assume that M does not admit closed exact Lagrangian submanifolds Lagrangian isotopic to Λ_ε . Then*

$$pb_4^{M,+}(\mathcal{F}, \mathcal{C}, \mathcal{L}, \mathcal{H}) = \left((R_1 - R_0)T \right)^{-1} > 0,$$

and thus, Λ is autonomously $(R_1 - R_0)T$ -interlinked.

The autonomous interlinking of Λ stated in the theorem follows, by Theorem 1.11, from the lower bound on $pb_4^{M,+}(\mathcal{F}, \mathcal{C}, \mathcal{L}, \mathcal{H})$. An immediate consequence of Theorem 5.2 is as follows:

Corollary 5.3. *Assume that for every $m \in \mathbb{N}$, $M \times T^*\mathbb{T}^m$ does not admit closed exact Lagrangian submanifolds Lagrangian isotopic to $\Lambda_\varepsilon \times \mathbb{T}^m$. Then Λ is stably $(R_1 - R_0)T$ -interlinked.*

Proof of Theorem 1.12: Part A immediately follows from Theorem 5.2 with $K = \{\text{point}\}$, $M = \Sigma \times \mathbb{R}_+$ and $\nu = Id$, and, similarly, part B follows from Corollary 5.3. \square

Proof of Theorem 5.2. 1) First, let us prove

$$pb_4^{M,+}(\mathcal{F}, \mathcal{C}, \mathcal{L}, \mathcal{H}) \leq \left((R_1 - R_0)T \right)^{-1}. \quad (17)$$

By the symplectic invariance, the monotonicity (see (9)) and the product (see (8)) properties of pb_4^+ , it suffices to prove that

$$pb_4^{\Sigma \times \mathbb{R}_+,+}(\mathcal{F}'', \mathcal{C}'', \mathcal{L}'', \mathcal{H}'') \leq \left((R_1 - R_0)T \right)^{-1}. \quad (18)$$

Let us consider a Hamiltonian of the form $u(s)$ on $(\Sigma \times \mathbb{R}_+, d(s\lambda_0))$ such that:

- $u(s) = 0$ outside $(R_0 + \delta_1, R_1 + \delta_1)$,
- $u(R_1) = 1$,
- u is non-decreasing on $[R_0, R_1]$ and its derivative there satisfies $u'(s) \leq (R_1 - R_0)^{-1} + \delta_2$,

Here δ_1, δ_2 are small positive constants. The obvious relation between the Hamiltonian flow of u and the Reeb flow of λ_0 easily implies that the flow of u does not have chords from \mathcal{H}'' to \mathcal{L}'' of time-length $\leq (R_1 - R_0)^{-1}T - \epsilon$, where $\epsilon \rightarrow 0$ as $\delta_1, \delta_2 \rightarrow 0$. Note also that u 1-separates \mathcal{F}'' and \mathcal{C}'' . Therefore Theorem 1.11 implies that

$$pb_4^{M,+}(\mathcal{H}'', \mathcal{L}'', \mathcal{F}'', \mathcal{C}'') \leq \left((R_1 - R_0)T \right)^{-1},$$

which, by (6), yields (18) and hence (17). Let us emphasize that the above argument involves Hamiltonian chords connecting the walls of the Lagrangian tetragon, even though the actual applications deal with the chords connecting its floor and ceiling.

2) Let us now prove

$$pb_4^{M,+}(\mathcal{F}, \mathcal{C}, \mathcal{L}, \mathcal{H}) \geq \left((R_1 - R_0)T \right)^{-1}. \quad (19)$$

The sets Λ_ε C^0 -converge to Λ as $\varepsilon \rightarrow 0$. Moreover, we can subdivide Λ_ε into four sets $\mathcal{F}_\varepsilon, \mathcal{C}_\varepsilon, \mathcal{L}_\varepsilon, \mathcal{H}_\varepsilon$ that C^0 -converge, respectively, to $\mathcal{F}, \mathcal{C}, \mathcal{L}, \mathcal{H}$ as $\varepsilon \rightarrow 0$. In view of (10) we need to prove that

$$pb_4^{M,+}(\mathcal{F}_\varepsilon, \mathcal{C}_\varepsilon, \mathcal{L}_\varepsilon, \mathcal{H}_\varepsilon) \geq \left((R_1 - R_0)T - \delta(\varepsilon) \right)^{-1}$$

for some $\delta(\varepsilon) \rightarrow 0$ as $\varepsilon \rightarrow 0$.

Denote by D_{γ_ε} the disk bounded by γ_ε in $[R_0, R_1] \times [0, T]$. Set

$$\Gamma_\varepsilon := \Phi(\{\text{point}\} \times \gamma_\varepsilon) \subset \Lambda_\varepsilon''$$

and let

$$D_{\Gamma_\varepsilon} := \Phi(\{\text{point}\} \times D_{\gamma_\varepsilon})$$

be the disk bounded by Γ_ε .

Pick arbitrary smooth functions $F, G \in C_c^\infty(M)$ satisfying the following conditions:

$$F|_{Op(\mathcal{F}_\varepsilon)} = G|_{Op(\mathcal{L}_\varepsilon)} = 0, \quad F|_{Op(\mathcal{C}_\varepsilon)} = G|_{Op(\mathcal{H}_\varepsilon)} = 1. \quad (20)$$

(Here Op stands for *some* open neighborhood of a set). Note that, by (20),

$$dF \wedge dG|_{Op(\Lambda_\varepsilon)} \equiv 0. \quad (21)$$

and thus $FdG|_{\Lambda_\varepsilon}$ is a closed form, which verifies condition 2 of Proposition 5.1.

We claim that $[FdG|_{\Lambda_\varepsilon}] = -c[\lambda|_{\Lambda_\varepsilon}]$ for some $c > 0$. In order to check it, we need to show that the integrals of λ and FdG over any $a \in H_1(\Lambda_\varepsilon; \mathbb{R})$ are proportional with the proportionality coefficient $-c$ for some $c > 0$ independent of a . Note that

$$H_1(\Lambda_\varepsilon; \mathbb{R}) \cong H_1(L; \mathbb{R}) \oplus \mathbb{R} \oplus H_1(K; \mathbb{R}),$$

where the isomorphism is given by ν_* and the middle summand \mathbb{R} is generated by the homology class of the curve $\nu(\Gamma_\varepsilon)$. Thus, it is enough to check the proportionality for $a = \nu_*b$, where $b \in H_1(L; \mathbb{R}) \oplus \mathbb{R} \oplus H_1(K; \mathbb{R})$ is one of the following types:

1. $b \in H_1(L; \mathbb{R})$.
2. $b = [\Gamma_\varepsilon]$.
3. $b \in H_1(K; \mathbb{R})$.

Let us consider the first case. Note that the form η vanishes on the zero section $K \subset T^*K$. Thus, by (16),

$$(\nu^*\lambda)|_{\Lambda_\varepsilon} = (s\lambda_0)|_{\Lambda_\varepsilon}, \quad (22)$$

and hence

$$\lambda(\nu_*b) = (s\lambda_0)(b) = 0,$$

since $b \in H_1(L; \mathbb{R})$ and the contact form λ_0 vanishes on the Legendrian submanifold L . Since FdG is zero on a neighborhood of \mathcal{H} and b can be represented by a chain in \mathcal{H} , we also get $[FdG](b) = 0$.

Let us move to the second case: $b = [\Gamma_\varepsilon]$. A direct computation using (22) and the Stokes theorem shows that

$$\lambda(\nu_*[\Gamma_\varepsilon]) = (s\lambda_0)(b) = \int_{D_{\Gamma_\varepsilon}} d(s\lambda_0) = (R_1 - R_0)T - \delta(\varepsilon) =: c > 0,$$

where the latter number c is the Euclidean area of D_{γ_ε} . At the same time another direct calculation using (20) yields

$$\int_{\Gamma_\varepsilon} \nu^*(FdG) = -1.$$

Thus, in this case

$$FdG(\nu_*b) = -\lambda(\nu_*b)/c.$$

Finally, consider the third case: $b \in H_1(K; \mathbb{R})$. By (22) and since $s\lambda_0$ vanishes on K ,

$$\lambda(\nu_*b) = (s\lambda_0)(b) = 0.$$

Moreover, ν_*b can be represented by a curve that lies in \mathcal{H} . Since, by (20), FdG is zero on a neighborhood of \mathcal{H} , we get that

$$FdG(\nu_*b) = 0.$$

Thus, $[FdG|_{\Lambda_\varepsilon}] = -c[\lambda|_{\Lambda_\varepsilon}]$ for $c > 0$. This verifies the condition 3 of Proposition 5.1. Condition 1 is a part of the hypothesis of the theorem that we are proving. Therefore Proposition 5.1 can be applied and we get

$$\max_M \{F, G\} \geq 1/c = \left((R_1 - R_0)T - \delta(\varepsilon) \right)^{-1}.$$

Since this is true for any $F, G \in C_c^\infty(M)$ satisfying (20), we get

$$pb_4^{M,+}(\mathcal{F}_\varepsilon, \mathcal{C}_\varepsilon, \mathcal{L}_\varepsilon, \mathcal{H}_\varepsilon) \geq \left((R_1 - R_0)T - \delta(\varepsilon) \right)^{-1},$$

where $\delta(\varepsilon) \rightarrow 0$ as $\varepsilon \rightarrow 0$, as required.

This proves (19). Together with (17) proved above this implies the proposition. \square

Remark 5.4. Recall that a symplectic manifold (M, ω) is called *symplectically aspherical* if $[\omega]$ vanishes on all homology classes in $H_2(M, \mathbb{R})$ represented by spheres, and a Lagrangian submanifold $Z \subset (M, \omega)$ is called *weakly exact* if $[\omega]$ vanishes on the subspace $H_2^D(M, Z; \mathbb{R}) \subset H_2(M, Z; \mathbb{R})$ generated by discs whose boundary lies on Z .

Theorem 5.2 admits a rather straightforward generalization to the case when (M, ω) is a symplectically aspherical symplectic manifold and the Lagrangian isotopy class of Λ_ε does not contain weakly exact Lagrangian submanifolds.

Namely, for such (M, ω) and Λ assume in addition that

$$H_2^D(M, \Lambda; \mathbb{R}) = \text{Im } \nu_{\sharp} + Q,$$

where

$$\nu_{\sharp} : H_2^D(U \times \mathcal{I} \times W, \Lambda'; \mathbb{R}) \rightarrow H_2^D(M, \Lambda; \mathbb{R})$$

is induced by ν and $Q \subset H_2^D(M, \Lambda; \mathbb{R})$ is a subgroup such that $\omega(Q) = 0$ and $\partial(Q)$ lies in the image of the natural inclusion $H_1(L; \mathbb{R}) \rightarrow H_1(\Lambda; \mathbb{R})$. Under these assumptions one can deduce the same conclusion as in Theorem 5.2.

5.2 Non-exactness of Lagrangians in Liouville manifolds

In this section we describe classes of Lagrangian tetragons in exact symplectic manifolds $(M, d\lambda)$ whose Lagrangian isotopy classes contain no exact Lagrangian submanifolds. Applying Corollary 5.3 to such tetragons, we shall prove our main results on the interlinking stated in the introduction.

Recall that an exact symplectic manifold $(M, \omega = d\lambda)$, with a fixed primitive λ of the symplectic form ω , is called *Liouville* if the following two conditions hold:

- the flow θ_t of the Liouville vector field v given by $i_v \omega = \lambda$ is complete;
- there exists a closed connected hypersurface Σ in M transversal to v bounding a domain $U \subset M$ with compact closure such that $M = U \sqcup \bigcup_{t \geq 0} \theta_t(\Sigma)$.

In this situation $\lambda_0 := \lambda|_{\Sigma}$ is a contact 1-form on Σ defining a contact structure $\xi = \ker \lambda$. The set $\bigcup_{t \in \mathbb{R}} \theta_t(\Sigma) \subset M$ can be canonically symplectically identified with the symplectization $\Sigma \times \mathbb{R}_+$ of Σ , so that λ is identified with the form $s\lambda_0$ on $\Sigma \times \mathbb{R}_+$. We call the contact manifold (Σ, ξ) *the ideal contact boundary* of the Liouville manifold M . One readily checks that the ideal contact boundary is uniquely defined up to a contactomorphism.

Examples of Liouville manifolds include, for instance, the cotangent bundle T^*V of a closed manifold V equipped with the canonical Liouville form

and the linear vector space $\mathbb{R}^{2n}(p, q)$ with the Liouville form $(pdq - qdp)/2$. Here the ideal contact boundaries are the unit sphere bundle of V with respect to any Riemannian metric, and the standard contact sphere, respectively. The product of two Liouville manifolds is again Liouville.

In what follows we focus on the case when (Σ, ξ) is the ideal contact boundary of a Liouville manifold M , and fix the notation θ_t for the Liouville flow on M .

Theorem 5.5. *The exact symplectic manifold $(\Sigma \times \mathbb{R}_+ \times T^*\mathbb{T}^n, d(s\lambda_0 + pdq))$ admits no closed exact Lagrangian submanifolds. In particular, the pair (Σ, L) is stably non-exact for every choice of a closed Legendrian submanifold $L \subset \Sigma$.*

Proof. Assume on the contrary that there exists a closed exact Lagrangian submanifold Z of $(\Sigma \times \mathbb{R}_+ \times T^*\mathbb{T}^n, d(s\lambda_0 + pdq))$. Let Z_t be the image of Z under the Liouville flow of $M \times T^*\mathbb{T}^n$ given by $(x, p, q) \mapsto (\theta_t(x), tp, q)$. Note that Lagrangian submanifolds Z_t are exact for all t , and Z_τ is disjoint from Z for sufficiently large τ . This contradicts to a theorem by Gromov stating that a symplectically aspherical and geometrically bounded at infinity symplectic manifold – and, in particular, any Liouville manifold – does not admit a displaceable closed exact Lagrangian submanifold [14], [3]. \square

Remark 5.6. By a theorem of Gromov [14] (cf. [3]), any closed weakly exact Lagrangian submanifold of a geometrically bounded symplectically aspherical symplectic manifold (M, ω) is non-displaceable. The geometric boundedness assumption, which holds, for instance for Liouville manifolds and cotangent bundles, enables one to control the behavior of pseudo-holomorphic curves in M “at infinity”. A minor modification of the above arguments yields the following claim:

Let $(\Sigma, \ker \lambda_0)$ be an arbitrary (not necessarily closed) contact manifold. Assume that for some $a > 0$ the domain $\Sigma \times (a, +\infty) \subset (\Sigma \times \mathbb{R}_+, d(s\lambda_0))$ admits a symplectic embedding ν to a geometrically bounded symplectic manifold (M, ω) . Let $L \subset \Sigma$ be a closed connected Legendrian submanifold so that the morphism $H_2^D(\Sigma, L) \rightarrow H_2^D(M, L)/\ker \omega$ induced by ν is onto. (Here $\ker \omega$ is the kernel of the symplectic area map $D \mapsto \int_D \omega$ on $H_2^D(M, L)$.) Then the pair (Σ, L) is stably non-exact.

If ω is exact, i.e. $\omega = d\lambda$, the assumption on the morphism of the relative homology groups can be removed, provided $\nu^*\lambda = s\lambda_0$.

It would be interesting to find meaningful dynamical applications of this result.

We have just proved that any Lagrangian tetragon in the symplectization $(\Sigma \times \mathbb{R}_+, d(s\lambda_0))$ of Σ is not Lagrangian isotopic to an exact Lagrangian submanifold in $(\Sigma \times \mathbb{R}_+, d(s\lambda_0))$. *Can it be Lagrangian isotopic to an exact Lagrangian submanifold in M ?* This question is still open. Below we answer it in negative in two particular cases.

First, we consider the class of subcritical Weinstein manifolds. Recall that a Liouville manifold is *Weinstein* if the Liouville vector field v is gradient-like for a proper bounded from below Morse function f with finite number of critical points. It is known that the indices of all critical points of f are $\leq n$, where $\dim M = 2n$. A Weinstein manifold M is called *subcritical* if for some choice of f all the indices are $< n$, and *critical* otherwise.

Example 5.7. The standard symplectic vector space $\mathbb{R}^{2n}(p, q)$ equipped with the Liouville form $\lambda = (pdq - qdp)/2$ is Weinstein subcritical. Indeed, the Liouville vector field is $v = (p\partial/\partial p + q\partial/\partial q)/2$. It is gradient like for the proper non-negative function $f(p, q) = p^2 + q^2$ whose only critical point is non-degenerate minimum at 0. The ideal contact boundary of \mathbb{R}^{2n} is the unit sphere $\mathbb{S}^{2n-1} \subset \mathbb{R}^{2n}$ equipped with standard contact structure ξ defined by the restriction of λ .

More generally, the product $M \times \mathbb{R}^2$ with any Liouville M is subcritical. In contrast to that, the cotangent bundle T^*V of a closed manifold V is critical.

Theorem 5.8. *Assume that $(M, \omega = d\lambda)$ is subcritical. Then the exact symplectic manifold $(M \times T^*\mathbb{T}^m, d(\lambda + pdq))$ admits no exact Lagrangian submanifolds.*

Proof. Biran and Cieliebak showed [5] that every compact subset of M (and a fortiori of $M \times T^*\mathbb{T}^m$) is displaceable. The existence of closed exact Lagrangian submanifolds would again violate the theorem of Gromov cited in the previous proof, yielding a contradiction. \square

Proof of Theorem 1.7. We work in the situation of Example 5.7. By Theorem 5.8 and Corollary 5.3, any Lagrangian tetragon $\Lambda \subset (\mathbb{R}^{2n}, dp \wedge dq)$, built using any closed connected Legendrian submanifold of (\mathbb{S}^{2n-1}, ξ) and some R_0, R_1, T , is stably $(R_1 - R_0)T$ -interlinked in $(\mathbb{R}^{2n}, dp \wedge dq)$. \square

Next, let V be a closed manifold, and $\Sigma \subset T^*V$ be the unit sphere bundle with respect to any Riemannian metric on V . Even though T^*V is a critical Weinstein manifold, we have the following result.

Theorem 5.9. *The symplectic manifold $T^*V \times T^*\mathbb{T}^m$ admits no exact Lagrangian submanifolds Lagrangian isotopic to Λ_ε .*

Proof. Assume, by contradiction, that $Z \subset T^*V \times T^*\mathbb{T}^m$ is an exact Lagrangian submanifold Lagrangian isotopic to Λ_ε . Then the \mathbb{Z}_2 -degree of the projections $\Lambda_\varepsilon \rightarrow V \times \mathbb{T}^m$ and $Z \rightarrow V \times \mathbb{T}^m$ are equal. By Kragh's theorem [17, Cor. 1.1], the latter degree is non-zero. (Recall that Kragh's theorem says that the \mathbb{Z}_2 -degree of the projection $X \rightarrow Y$ of *any* closed exact Lagrangian submanifold X of the cotangent bundle of *any* manifold Y is non-zero). Thus, the \mathbb{Z}_2 -degree of the projection $\Lambda_\varepsilon \rightarrow V \times \mathbb{T}^m$ is non-zero.

On the other hand, note that the Lagrangian isotopy type of Λ_ε in $T^*V \times T^*\mathbb{T}^m$ is independent of the parameters R_0, R_1, T (where ε is always chosen sufficiently small depending on R_0, R_1, T). Thus, without loss of generality, we may assume that T is sufficiently small (depending on R_0 and R_1) so that $\pi(\Lambda''_\varepsilon)$ lies in a small neighborhood of $\pi(L)$ in V . Note that $\pi(L) \subsetneq V$, since $\dim L = \dim V - 1$. Therefore the projection $\pi : \Lambda''_\varepsilon \rightarrow V$ is not surjective. Hence the projection $\pi \times Id : \Lambda_\varepsilon \rightarrow V \times \mathbb{T}^m$ also is not surjective and therefore the \mathbb{Z}_2 -degree of the projection $\Lambda_\varepsilon \rightarrow V \times \mathbb{T}^m$ is zero. We have got a contradiction. This finishes the proof. \square

Remark 5.10. In fact, the same argument works verbatim for tetragons in T^*V with non-compact V .

As we noted above, Theorem 5.5 implies that if Σ is the unit cotangent bundle (with respect to an arbitrary Riemannian metric) of a *closed* manifold V , then for any closed connected Legendrian submanifold L of Σ , the pair (Σ, L) is non-exact. The version of Theorem 5.9 for an arbitrary V immediately yields the non-exactness of (Σ, L) for any V .

Proof of Theorem 1.5. The Lagrangian tetragon Λ in $T^*\mathbb{T}^k$ appearing in the statement of the theorem is exactly the one constructed as above in the case where Σ is the unit sphere bundle with respect to the Euclidean metric on $V := \mathbb{T}^k$, L is a Legendrian fiber of Σ , $K = \{\text{a point}\}$, and $0 < R_0 < R_1$, $0 < T < 1/2$ are some numbers. Thus, Theorem 5.9 and Corollary 5.3 can be applied, yielding that Λ is stably $(R_1 - R_0)T$ -interlinked. \square

Remark 5.11. In view of Theorem 1.11 this means that $(R_1 - R_0)T$ is the smallest κ for which the Lagrangian tetragon Λ appearing in Theorem 1.5 is κ -interlinked.

Acknowledgments. We thank Vadim Kaloshin for very useful discussions and stimulating questions on Hamiltonian instabilities and Albert Fathi for communicating to us Theorem 3.1. We are grateful to Paul Biran and Tobias Ekholm for useful comments on Reeb dynamics and to an anonymous referee for corrections. L.P. thanks the University of Chicago, where parts of this paper were written, for an excellent research atmosphere.

References

- [1] Alves, M.R.R., *Legendrian contact homology and topological entropy*, preprint arXiv:1410.3381, 2014.
- [2] Arnold, V., *Mathematical methods of classical mechanics*. Springer-Verlag, New York, 1989.
- [3] Audin, M., Lalonde, F., Polterovich, L., *Symplectic rigidity: Lagrangian submanifolds*, in *Holomorphic curves in symplectic geometry*, 271-321, Progr. Math., **117**, Birkhäuser, Basel, 1994.
- [4] Buhovsky, L., Entov, M., Polterovich, L., *Poisson brackets and symplectic invariants*, *Selecta Math.* **18** (2012), 89-157.
- [5] Biran, P., Cieliebak, K., *Lagrangian embeddings into subcritical Stein manifolds*, *Israel J. Math.* **127** (2002), 221-244.
- [6] Bounemoura, A., Kaloshin, V., *Generic diffusion for a class of non-convex Hamiltonians with two degrees of freedom*, *Moscow Math. J.*, **14** (2014), 181-203.
- [7] Cieliebak, K., Eliashberg, Y., *From Stein to Weinstein and back. Symplectic geometry of affine complex manifolds*. AMS, Providence, RI, 2012.
- [8] Ekholm, T., *Rational SFT, linearized Legendrian contact homology, and Lagrangian Floer cohomology*, in “*Perspectives in analysis, geometry, and topology*”, 109-145, Birkhäuser, Boston, 2012.

- [9] Eliashberg, Y., Gromov, M., *Convex symplectic manifolds*, in *Several complex variables and complex geometry, Part 2 (Santa Cruz, CA, 1989)*, 135-162, Proc. Sympos. Pure Math., **52**, Part 2, AMS, Providence, RI, 1991.
- [10] Eliashberg, Y., *A few remarks about symplectic filling*, Geom. Topol. **8** (2004), 277-293.
- [11] Etnyre, J., *On symplectic fillings*, Algebr. Geom. Topol. **4** (2004), 73–80.
- [12] Fathi, A., *An Urysohn-type theorem under a dynamical constraint*, J. of Modern Dynamics **10** (2015), 331-338.
- [13] Floer A., *Morse theory for Lagrangian intersections*, Journal of Differential Geometry **28** (1988), 513-547.
- [14] Gromov, M., *Pseudoholomorphic curves in symplectic manifolds*, Invent. Math. **82** (1985), 307-347.
- [15] Hagedorn, P., *Die Umkehrung der Stabilitätssätze von Lagrange-Dirichlet und Routh*, Arch. Rational Mech. Anal. **42** (1971), 281-316.
- [16] Hofer, H., *Lagrangian embeddings and critical point theory*, Ann. Inst. H. Poincaré Anal. Non Linéaire **2** (1985), 407–462.
- [17] Kragh, T., *Parametrized ring-spectra and the nearby Lagrangian conjecture (with an appendix by Mohammed Abouzaid)*, Geom. Topol. **17** (2013), 639-731.
- [18] Laudенbach, F., Sikorav, J.-C., *Persistence d'intersection avec la section nulle au cours d'une isotopie hamiltonienne dans un fibré cotangent*, Invent. Math. **82** (1985), 349–357.
- [19] Lisi, S., *Homoclinic orbits and Lagrangian embeddings*, Int. Math. Res. Not. IMRN (2008), no. 5, Art. ID rnm 151, 16 pp.
- [20] Mohnke, K., *Holomorphic disks and the chord conjecture*, Ann. of Math. **154** (2001), 219-222.
- [21] Moser, J., *On the volume elements on a manifold*, Trans. AMS **120** (1965), 288-294.

- [22] Murphy, E., *Closed exact Lagrangians in the symplectization of contact manifolds*, preprint, arXiv:1304.6620, 2013.
- [23] Nehorošhev, N.N., *An exponential estimate of the time of stability of nearly integrable Hamiltonian systems*, Russian Math. Surveys **32** (1977), 1-65.
- [24] Taliaferro, S., *An inversion of the Lagrange-Dirichlet stability theorem*, Arch. Rational Mech. Anal. **73** (1980), 183-190.