## $AdS_2/CFT_1$ , Whittaker vector and Wheeler-DeWitt equation

Tadashi Okazaki<sup>1</sup>

Department of Physics and Center for Theoretical Sciences, National Taiwan University, Taipei 10617, Taiwan

## Abstract

We study the energy representation of conformal quantum mechanics as the Whittaker vector without specifying classical Lagrangian. We show that a generating function of expectation values among two excited states of the dilatation operator in conformal quantum mechanics is a solution to the Wheeler-DeWitt equation and it corresponds to the  $AdS_2$  partition function evaluated as the minisuperspace wave function in Liouville field theory. We also show that the dilatation expectation values in conformal quantum mechanics lead to the asymptotic smoothed counting function of the Riemann zeros.

<sup>&</sup>lt;sup>1</sup>tadashiokazaki@phys.ntu.edu.tw

The holographic principle [1, 2] states that quantum gravity on (d+1)-dimensional manifold can be described by a theory on its *d*-dimensional boundary. The  $AdS_{d+1}/CFT_d$ correspondence [3] which is one of the greatest productions of string theory provides the most successful realization as the relationship between effective gauge theories of the brane dynamics and string theory on the near horizon AdS geometry. The  $AdS_2/CFT_1$  could conceivably be the most significant case in that all the extremal black holes contain an AdS<sub>2</sub> factor in their near horizon geometry [4, 5]. In spite of a lot of interesting works [6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30] it would be fair to say that we have not yet gained a fully satisfactory understanding of the correspondence due to the peculiarities of the  $AdS_2/CFT_1$ , including the fact that only the AdS<sub>2</sub> has two disconnected boundaries and it is a long-standing question whether the dual  $CFT_1$  description is a single CFT or two systems living on the two boundaries.

In this letter we study conformal quantum mechanics (CQM) without specifying classical Lagrangian description. One important consequence is a generic evidence of the  $AdS_2/CFT_1$  as the relationship between a generating function of the dilatation expectation values in the boundary CQM and a partition function of the bulk  $AdS_2$ . We show that the expectation values are not associated with the ground state but with two excited states in the correspondence. Our result is in favor of the statement [31, 25, 32, 33, 34] that the dual  $CFT_1$ 's on two boundaries of  $AdS_2$  space-time are excited and entangled. We also claim that the energies on the boundary CQM would be responsible for the  $AdS_2$  radius so that the ground state would correspond to a flat space in the bulk with an infinitely large  $AdS_2$  radius.

Another intriguing thing is the speculative implication of the resulting expectation values of the dilatation operator in CQM. When we consider the DFF-model [35] as CQM, the dilatation operator takes the form of xp. Such operator has been proposed as a strong candidate for the realization of the Hilbert-Pólya conjecture that the imaginary part of the non-trivial Riemann zeros are eigenvalues of a self-adjoint operator (see [36, 37, 38, 39, 40, 41] and references therein). Unlike a lot of efforts undertaken thus far we here propose a novel approach to obtain the counts of the Riemann zeros from CQM point of view. The fact that the operator xp is the dilatation operator in CQM rather than the Hamiltonian enables us to jump into a fairly general setting beyond the operator with the form of xp. We show that the expectation values of the dilatation operator in CQM naturally lead to the asymptotic form of the smoothed counting function of the Riemann zeros.

We shall begin by considering CQM that is invariant under the conformal symmet-

ric transformation of the time coordinate t [35]

$$\delta t = \epsilon_1 + \epsilon_2 t + \epsilon_3 t^2 \tag{1}$$

where  $\epsilon_1$ ,  $\epsilon_2$  and  $\epsilon_3$  are identified with the infinitesimal parameters of the translation, the dilatation and the conformal boost transformation respectively. The corresponding generators, the Hamiltonian H = id/dt, the dilatation operator D = it (d/dt), and the conformal boost operator  $K = it^2 (d/dt)$  obey the commutation relations

$$[H,D] = iH \tag{2}$$

$$[K,D] = -iK \tag{3}$$

$$[H,K] = 2iD\tag{4}$$

of  $SL(2,\mathbb{R})$ , which we call the one-dimensional conformal group. Hence the Hilbert space of conformal quantum mechanical system exhibits the  $SL(2,\mathbb{R})$  conformal symmetry and the physical states would be classified by its irreducible representation. Since we wish to make all integrals convergent, we require the unitarity of the representation. The classification of the irreducible unitary representations of  $SL(2,\mathbb{R})$ was studied in [42] and the irreducible unitary representation of  $SL(2,\mathbb{R})$  conformal group as a function of the continuous time coordinate t can be generalized by taking the principal spherical series of the representation which is induced by the one-dimensional representation of the Borel subgroup. Let  $V_{\lambda}$  be a set of the irreducible unitary representations of  $SL(2,\mathbb{R})$  with weight  $\lambda$ . Then the Hamiltonian H, the dilatation operator D and the conformal boost operator K can be expressed as

$$H = i\frac{d}{dt} \tag{5}$$

$$D = it\frac{d}{dt} + \frac{\lambda}{2i} \tag{6}$$

$$K = it^2 \frac{d}{dt} + \frac{1}{i}\lambda t,\tag{7}$$

satisfying the commutation relations (2), (3) and (4). The unitarity implies that  $\frac{1}{2}(\lambda + 1)$  is pure imaginary. The finite conformal transformation is

$$t^n \to \frac{(\alpha t + \beta)^n}{(\gamma t + \delta)^{\lambda + n}} \tag{8}$$

where the parameters  $\alpha$ ,  $\beta$ ,  $\gamma$  and  $\delta$  are the elements of real two by two matrices with determinant one

$$A = \begin{pmatrix} \alpha & \gamma \\ \beta & \delta \end{pmatrix}, \qquad \qquad \alpha \delta - \beta \gamma = 1 \tag{9}$$

which forms the one-dimensional conformal group  $SL(2,\mathbb{R})$ . The quadratic Casimir operator is given by

$$C_2 = HK - iD - D^2$$
$$= \frac{\lambda^2}{4} + \frac{\lambda}{2}.$$
 (10)

Let us define the ground state  $|0\rangle_{\Delta}$  by

$$H|0\rangle_{\Delta} = 0 \tag{11}$$

$$D|0\rangle_{\Delta} = \Delta|0\rangle_{\Delta}.\tag{12}$$

From the equations (10)-(12) we see that the eigenvalue  $\Delta$  of the ground state  $|0\rangle_{\Delta}$  is  $\Delta = \frac{\lambda}{2i}$  and we thus write the ground state as  $|0\rangle_{\lambda}$ . Now consider the energy eigenstate obeying

$$H|E\rangle = E|E\rangle. \tag{13}$$

This eigenvector  $|E\rangle$  is known as the Whittaker vector in the representation theory of  $SL(2,\mathbb{R})$  [43, 44, 45]. Given eigenvalue E and irreducible representation with weight  $\lambda$ , there is a unique Whittaker vector and we can write

$$|E\rangle_{\lambda} = -\sum_{k} \sum_{n} C_{k} \frac{(-EK)^{n}}{n!\lambda(\lambda-1)\cdots(\lambda-n+1)} |0\rangle_{\lambda,k}$$
(14)

where k parametrizes the degenerate ground states which might arise from some symmetries in the theories and  $C_k$  is the weighted coefficient for k-th ground state. For simplicity let us assume the vanishing of tunneling amplitudes;  $\lambda_{,l}\langle 0|0\rangle_{\lambda,k} = \delta_{l,k}$  and the normalization of the ground states;  $\sum_k |C_k|^2 = 1$  so that we shall omit the indices k. Correspondingly we can take the dual energy eigenstate vector  $\lambda \langle E|$  as the dual Whittaker vector which satisfies the relation <sup>2</sup>

$$_{\lambda}\langle E|K = E_{\lambda}\langle E| \tag{15}$$

and it can be represented by

$$_{\lambda}\langle E| := -_{\lambda}\langle 0| \sum_{k,n} C_k^* \frac{(-EH)^n}{n!\lambda(\lambda-1)\cdots(\lambda-n+1)}.$$
(16)

With the Whittaker vector  $|E\rangle_{\lambda}$  and its dual vector  $_{\lambda}\langle E|$  as the energy eigenstate vector and its dual vector in conformal quantum mechanics, we will consider the

 $<sup>^2 {\</sup>rm The}$  choice of pair of the Whittaker vector and the dual Whittaker vector results in the consistent Hamiltonian reduction.

situation where the theory is coupled to two-dimensional bulk theory to investigate the  $AdS_2/CFT_1$  correspondence. Let us consider the function defined by

$$W_{\lambda, E_L, E_R}(\phi) := {}_{\lambda} \left\langle E_L | e^{-2i\phi D} | E_R \right\rangle_{\lambda}$$
(17)

where  $E_L$ ,  $E_R$  are the eigenvalues of the Whittaker vector (13) and the dual Whittaker vector (15) respectively. Here  $\phi$  is regarded as the restriction of some bulk field in AdS<sub>2</sub> space-time on the boundary that is coupled to the dilatation operator D in conformal quantum mechanics on the boundaries. By acting the quadratic Casimir operator (10) on the function (17), one obtains the differential equation

$$\left[\frac{1}{2}\frac{\partial^2}{\partial\phi^2} - \frac{\partial}{\partial\phi} - 2E_L E_R e^{2\phi}\right] W_{\lambda, E_L, E_R}(\phi) = \left(\frac{\lambda^2}{2} + \lambda\right) W_{\lambda, E_L, E_R}(\phi).$$
(18)

The function  $W_{\lambda,E_L,E_R}(\phi)$  is known as the  $SL(2,\mathbb{R})$  Whittaker function [43, 44, 45, 46]. At first sight one might expect that the Whittaker function  $W_{\lambda,E_L,E_R}$  plays a role of the generating function of the expectation values of the operator D. However, the unitarity asserts that the eigenvalue  $\Delta$  of the operator D associated with the ground state  $|0\rangle_{\lambda}$  are not real-valued observables. Alternatively if we consider a shifted operator  $(D-\frac{i}{2})$ , the corresponding eigenvalues would provide real-valued physical quantities. Furthermore in a more precise treatment one can express the bulk filed  $\phi$  as  $\beta\phi_0$  where  $\phi_0$  is the time-dependent part while  $\beta$  is the time-independent part. Instead of the Whittaker function (17) let us consider the function

$$\Psi_{\lambda,\beta,E_L,E_R}(\phi_0) := \lambda \langle E_L | e^{-2i\beta\phi_0(D-\frac{i}{2})} | E_R \rangle_{\lambda}.$$
(19)

Applying the quadratic Casimir operator (10) on the function (19), we get

$$\left[\frac{1}{2}\frac{\partial^2}{\partial\phi_0^2} - 2\beta^2 E_L E_R e^{2\beta\phi_0}\right]\Psi_{\lambda,\beta,E_L,E_R}(\phi_0) = \frac{1}{2}\beta^2 \left(\lambda + 1\right)^2 \Psi_{\lambda,\beta,E_L,E_R}(\phi_0).$$
(20)

We see that the equation (20) is the Wheeler-DeWitt equation that is encountered in the minisuperspace approximation of Liouville field theory (LFT).

Local properties of LFT can be described by the Lagrangian

$$\mathcal{L} = \frac{1}{4\pi} \partial^{\mu} \phi \partial_{\mu} \phi + \mu e^{2b\phi}$$
(21)

where b is the dimensionless coupling constant and  $\mu$  is the cosmological coupling constant. The equation of motion is

$$\Delta \phi = 4\pi b \mu e^{2b\phi}.\tag{22}$$

In two dimensions it is always possible to make any metric  $g_{\mu\nu}$  conformally flat by coordinate redefinition  $g_{\mu\nu} = e^{2b\phi}\eta_{\mu\nu}$ . Furthermore, in two dimensions, the curvature

can be determined by the scalar curvature. Equation (22) asserts that the curvature is a negative constant  $-8\pi b^2 \mu$  and  $g_{\mu\nu}$  describes a two-dimensional surface with constant negative curvature, thus the corresponding Lorentzian surface can be identified with AdS<sub>2</sub> space-time locally.

In order to quantize LFT via canonical quantization, we shall employ the Fourier decomposition of the Liouville field  $\phi$  and its canonical momentum  $\Pi$  on the cylinder

$$\phi(t,\sigma) = \phi_0(t) + \sum_{n \neq 0} \frac{i}{n} \left[ a_n(t)e^{-in\sigma} + b_n(t)e^{in\sigma} \right]$$
  
$$\Pi(t,\sigma) = p_0(t) + \sum_{n \neq 0} \left[ a_n(t)e^{-in\sigma} + b_n(t)e^{in\sigma} \right]$$
(23)

with  $a_n^{\dagger} = a_{-n}, \ b_n^{\dagger} = b_{-n}$ . The canonical relation

$$[\phi(t,\sigma),\Pi(t,\sigma')] = i\delta(\sigma - \sigma') \tag{24}$$

leads to the commutation relations

$$[\phi_0, p_0] = i, \qquad [a_n, a_m] = \frac{n}{2} \delta_{n, -m}, \qquad [b_n, b_m] = \frac{n}{2} \delta_{n, -m}, \qquad (25)$$

which imply that  $a_n, b_n$  are creation operators while  $a_{-n}, b_{-n}$  are annihilation operators. The spectrum of LFT has been discussed in the minisuperspace approximation [47]. The minisuperspace approximation was originally proposed in quantum gravity [48, 49] where the problem is simplified by only treating the zero mode and truncating the higher excited modes. Whether physics in minisuperspace quantization gives a faithful properties of quantum gravity remains an open question, however, it has been discussed that the minisuperspace approximation would be exact in pure two-dimensional gravity [50, 51]. Replacing the canonical momentum  $p_0 = \frac{\dot{\phi}_0}{2\pi}$  with differential operator  $-i(\partial/\partial\phi_0)$ , we obtain the minisuperspace Schrödinger equation

$$\left[-\frac{1}{2}\frac{\partial^2}{\partial\phi_0^2} + 2\pi\mu e^{2b\phi_0}\right]\Psi_P(\phi_0) = 2P^2\Psi_P(\phi_0)$$
(26)

where P is the Liouville momentum, the eigenvalue of the Hamiltonian and  $\Psi_P(\phi_0)$ is the minisuperspace wave function. For  $\phi_0 \to -\infty$  the interaction is small and the wave function is a linear combination of  $e^{\pm iP\phi_0}$ . Because of the complete reflection potential at  $\phi_0 \to 0$ , P is restricted to be positive so that the incoming wave uniquely determines the reflected wave.

The function  $\Psi_{\lambda,\beta,E_L,E_R}(\phi_0)$  in CQM plays a role of the generating function of the expectation values evaluated among two excited states. On the other hand, the Liouville wave function  $\Psi_P(\phi_0)$  would be regarded as the partition function of AdS<sub>2</sub>, which is a function of the boundary values in the sense that a partition function generically transforms as a wave function under a change of polarization on field space specified at a boundary [52, 53, 54, 55] and LFT describes  $AdS_2$  space-time in the classical solution. Therefore we come to the interesting conclusion that  $AdS_2$  bulk mode  $\phi$  that behaves as the zero-mode  $\phi_0$  near the boundary is dual to a source term  $\phi_0(D-\frac{i}{2})$  in the CQM on the boundary of  $AdS_2$  via  $AdS_2/CFT_1$  correspondence

$$\Psi_{\lambda,\beta,E_L,E_R}(\phi_0) = \left\langle E_L \left| e^{-2i\beta\phi_0 \left( D - \frac{i}{2} \right)} \right| E_R \right\rangle_{\text{CQM}}$$
$$= Z_{\text{AdS}_2}(\phi|_{\text{bdy}} = \phi_0) = \Psi_P(\phi_0). \tag{27}$$

We observe that the expectation values in the CQM are evaluated between two excited states defined by the Whittaker vector (13) and the dual Whittaker vector (15) in the correspondence (27). The two distinct states in the correspondence (27) would would enable us to have two independent dynamical systems. This is consistent to the statement [20, 25, 34, 28] that unlike higher dimensional cases  $AdS_2$  in the global coordinate has two boundaries and the dual conformal field theory of asymptotically  $AdS_2$  is realized as two systems or two copies of  $CFT_1$  on the two boundaries although we cannot exclude the possibility of a single  $CFT_1$  as the dual description. The appearance of the excited states could also be in favor of the statement that in the Lorentzian AdS/CFT operator expectation values in excited CFT states differ from vacuum expectation values due to the existence of normalizable propagating states in the bulk [32, 33].

The comparison of the two Wheeler-DeWitt equations (20) and (26) identifies the coupling constant b in LFT with the constant parameter  $\beta$  in CQM and establishes the dictionary of parameters between the bulk LFT of AdS<sub>2</sub> and the boundary CQM as

$$\frac{\pi\mu}{b^2} = E_L E_R \tag{28}$$

$$\frac{P^2}{b^2} = -\frac{1}{4} \left(\lambda + 1\right)^2 = \left(\Delta - \frac{i}{2}\right)^2.$$
(29)

Therefore the relations (28) and (29) indicate that the quantum gravity on  $AdS_2$  can be described by two conformal quantum mechanical systems on the boundary with energies  $E_L$  and  $E_R$  as a holographic principle [1, 2].

The equation (28) says that the excited states with non-vanishing energy eigenvalues  $E_L$ ,  $E_R$  in the boundary CQM are needed to realize negative constant curvature of AdS<sub>2</sub> space-time which is generated by the non-trivial interaction term with finite and non-vanishing parameters b,  $\mu$  in two-dimensional gravity theory. It is illustrative to compare our result with the analogous statement in the AdS<sub>3</sub>/CFT<sub>2</sub> that the  $AdS_3$  radius  $l_3$  is represented by the central charge c of the CFT<sub>2</sub> through the Brown-Henneaux relation  $c = \frac{3l_3}{2G_3}$  [56] where  $G_3$  is the three-dimensional Newton constant. Instead of the central charge the energies of the states play an important role in the  $AdS_2/CFT_1$ , however, the relation between the  $AdS_2$  radius and the energies is even more attractive in that one of the other parameters in LFT necessarily comes about. In terms of the coupling constant b controlling the quantum effect in LFT we can write the  $AdS_2$  radius  $l_2$  as

$$\frac{1}{\sqrt{E_L E_R}} = 2b^2 l_2. \tag{30}$$

The semiclassical analysis in LFT is valid for small b, and accordingly our formula (30) would reflect the fact that the ground states with vanishing energies in the boundary CQM force the AdS<sub>2</sub> radius to go to infinity and the classical AdS<sub>2</sub> geometry then becomes flat space-time.

We see from the equation (29) that the Liouville momentum P in two-dimensional gravity theory corresponds to the conformal dimension of the ground state in the dual CQM. As we have discussed, the unitarity condition in CQM requires that  $\frac{1}{2}(\lambda + 1)$  is pure imaginary. This is consistent to the fact that the Liouville momentum P is real in the dual two-dimensional gravity theory.

We would like to emphasize that the correspondence (27) and the dictionaries (28), (29), (30) are quite universal since we have not specified the conformal quantum mechanical systems so far. However, if we contain more specific information characterizing dynamical properties and symmetries, there would be more fruitful statements (the GKP-Witten relation [57, 58]) in the  $AdS_2/CFT_1$  as the extension of the relation (27)

$$\left\langle e^{h_0 \mathcal{O}} \right\rangle_{\text{CQM}} = Z_{\text{AdS}_2}(h|_{\text{bdy}} = h_0)$$
 (31)

where  $h_0$  is some function of the boundary values for the bulk field h while  $\mathcal{O}$  is the dual operator in CQM. For a non-flat space the left values in the correspondence (31) are generally presumed to be computed between two excited states from the relation (30). It would be interesting that there has been proposals for such relation associated with the DFF-model in [28] and with the counting of microstates of BPS extremal black holes in [26, 25].

We now consider the resulting expectation values

$${}_{\lambda}\langle E_L | \left( D - \frac{i}{2} \right) | E_R \rangle_{\lambda} = \frac{i}{2} \frac{\delta}{\delta \phi} \Psi_{\lambda, E_L, E_R}(\phi) \Big|_{\phi=0}$$
(32)

and its possible application to one of the deepest mathematical problem, the Riemann hypothesis. The equation (20) has two linearly independent solutions, which are known

to be cylindric functions [59]. By requiring the unitarity we can write the solutions as

$$\Psi_{\lambda,E_L,E_R}(\phi) = \frac{1}{i} K_{\lambda+1} \left( 2\sqrt{E_L E_R} e^{\phi} \right)$$
(33)

where  $K_{\nu}(z)$  is the Macdonald function. We should note that the Macdonald functions of purely imaginary order  $\lambda + 1$  with positive argument are real. The prefactor  $\frac{1}{i}$  in the generating function (33) guarantees the reality condition of the expectation values in CQM. Making use of the recurrence relation

$$K_{\nu-1}(z) + K_{\nu+1}(z) = -2\frac{d}{dz}K_{\nu}(z)$$
(34)

and the formula (32) we can write the expectation values between the excited states as

$$_{\lambda}\langle E_{L}|\left(D-\frac{i}{2}\right)|E_{R}\rangle_{\lambda} = -\frac{z}{4}\left(K_{\lambda}(z)+K_{\lambda+2}(z)\right)$$
(35)

where  $z = 2\sqrt{E_L E_R}$ . Let us now consider a one particle conformal quantum mechanical model known as the DFF-model [35]

$$S = \frac{1}{2} \int dt \left( \dot{x}^2 - \frac{g}{x^2} \right) \tag{36}$$

with g being a dimensionless coupling constant. For the DFF-model the dilatation operator can be expressed as  $D = -\frac{1}{2}xp + \frac{i}{4}$  where p is the canonical momentum and the equation (35) becomes

$${}_{\lambda}\langle E_L | \left( xp + \frac{i}{2} \right) | E_R \rangle_{\lambda} = \frac{z}{2} \left( K_{\lambda}(z) + K_{\lambda+2}(z) \right).$$
(37)

It is speculated that the Riemann zeros would be realized as eigenvalues of the operator which takes the form of xp [36, 37, 38] as a promising candidate of the Riemann operator in the Hilbert-Pólya conjecture whose eigenvalues are the imaginary part of the non-trivial Riemann zeros. Berry and Keating [37, 38] identified the operator xpwith the Hamiltonian and imposed the conditions  $|x| \ge l_x$ ,  $|p| \ge l_p$  so that  $l_x l_p = 2\pi\hbar$ in the phase space. Then they found that the semiclassical number N(E) of states with the energy between 0 and E is given by the area in the phase space divided by the Planck cell  $h = 2\pi$ 

$$N(E) = \frac{E}{2\pi} \left( \log \frac{E}{2\pi} - 1 \right) + \mathcal{O}(1) \tag{38}$$

and observed that (38) precisely coincides with the asymptotics of the smoothed counting function of the number of Riemann zeros [60]. Connes [36] introduced the constraints  $|x| \leq \Lambda$ ,  $|p| \leq \Lambda$  where  $\Lambda$  is a common cutoff and counted the number of such semiclassical states as

$$N(E) = \frac{E}{\pi} \log \Lambda - \frac{E}{2\pi} \left( \log \frac{E}{2\pi} - 1 \right).$$
(39)

He interpreted the counting formula (39) as missing spectral lines associated to the smooth Riemann zeros which arise in the limit  $\Lambda \to \infty$ , however, it was reinterpreted as a finite size correction from a physical system later in [39]. According to these semiclassical proposals of the Hilbert-Pólya conjecture it has been desirable to replace these artificially imposed semiclassical regularizations of the operator xp with the proper quantum treatment which naturally generates a discrete spectrum. In order to obtain the discrete spectrum via quantization, there has been proposed various attempts including the modifications of the xp operator (see, e.g., [40, 41]) and the adoption of the regularization methods (see, e.g., [39]). Nevertheless, these attempts seem to be quite artificial and difficult fo find the quantum mechanical explanation to follow these ideas.

Here we wish to propose a novel perspective to acquire the distribution of the spectrum of the Riemann operator from conformal quantum mechanics point of view. Consider now the eigenfunction  $\Phi_{\rho}(x)$  of the operator  $\left(xp + \frac{i}{2}\right)$  in the equation (37) satisfying

$$\left[\frac{1}{i}x\frac{d}{dx} + \frac{i}{2}\right]\Phi_{\rho}(x) = \rho\Phi_{\rho}(x).$$
(40)

They take the form

$$\Phi_{\rho}(x) = Cx^{\frac{1}{2}+i\rho} \tag{41}$$

with C a constant of integration. The non-trivial zeros of the Riemann zeta function  $\zeta(s)$  which is conjectured to be  $s = \frac{1}{2} + i\rho$ ,  $\rho \in \mathbb{R}$  in the Riemann hypothesis appears in the power  $x^s$  of the eigenfunction  $\Phi_{\rho}(x)$ . The eigenvalues  $\rho$  of the operator  $\left(xp + \frac{i}{2}\right)$ which would be the candidates of the Riemann zeros can be continuous in the position eigenfunction. This is the same situation as has been already discussed in many literatures.

However, in CQM the operator  $\left(xp + \frac{i}{2}\right)$  should not be recognized as the Hamiltonian but rather as the dilatation operator whose expectation values can be measured by the energy eigenstates as

$$D(z;\rho) = \frac{z}{2} \left( K_{1-i\rho}(z) + K_{1+i\rho}(z) \right).$$
(42)

Note that the expression (42) can evidently be lifted to arbitrary conformal quantum mechanical systems by qualifying  $\rho$  as the eigenvalue of the operator  $-2\left(D-\frac{i}{2}\right)$ .

Although almost all quantum approaches so far have tried to identify the operator xp with the Hamiltonian and simultaneously diagonalize it with the position x or the momentum p, CQM would provide an alternative avenue to the Riemann hypothesis as the diagonalization of the dilatation operator. We observe that the ground state  $|0\rangle_{\lambda}$  is the eigenfunction of both of the Hamiltonian and the dilatation operator. Since the excited states are not eigenvectors of the dilatation operator, the limit  $E_L$ ,  $E_R \to 0$  of the expectation values (42) would naturally give rise to the eigenfunction of the operator  $-2(D - \frac{i}{2})$  multiplied by its eigenvalue  $\rho$ . In other words, the limit in which  $z = 2\sqrt{E_L E_R}$  goes to zero yields the definite eigenvalue  $\rho$  and the distribution function  $D(\rho)$  of the ground state. Hence the expectation values (42) are in some sense the regularized functions which produce the distribution of the eigenvalues  $\rho$  as

$$\rho D(\rho) = \lim_{z \to 0} \frac{z}{2} \left( K_{1-i\rho}(z) + K_{1+i\rho}(z) \right)$$
(43)

where we have used the relation  $\lambda = -(1 + i\rho)$  and the formula  $K_{\nu}(z) = K_{-\nu}(z)$ . The asymptotic behavior of the Macdonald function

$$K_{1+i\rho}(z) \sim \sqrt{\frac{\pi}{z}} e^{-\frac{\pi}{2}\rho} \left(\frac{2\rho}{ze}\right)^{i\rho} \tag{44}$$

for large  $\rho$  allows us to write (43) as

$$\rho D(\rho) = \lim_{z \to 0} \sqrt{\pi z} e^{-\frac{\pi}{2}\rho} \cos\left[\rho \ln\left(\frac{2\rho}{ze}\right)\right].$$
(45)

The semiclassical distribution of (45) for large  $\rho$  is realized when the cosine function is at its maximum

$$\cos\left[\rho\ln\left(\frac{2\rho}{ze}\right)\right] = 1\tag{46}$$

so that

$$\frac{\rho}{\pi} \left[ \ln \left( \frac{\rho}{E_L E_R} \right) - 1 \right] = 2n, \quad \forall n \in \mathbb{Z}.$$
(47)

Since the expression (47) diverges when  $E_L, E_R \to 0$ , a low energy cutoff is required to make sense of the expression (47). Let us introduce the cutoff  $\Lambda$  such that  $E_L E_R = 2\pi/\Lambda$ . Then we obtain the behavior of the large eigenvalues  $\rho$  as

$$N(\rho) = \frac{\rho}{2\pi} \ln \Lambda + \frac{\rho}{2\pi} \left( \ln \frac{\rho}{2\pi} - 1 \right).$$
(48)

Remarkably the first term is a continuum in the limit  $\Lambda \to \infty$  while the second term leads to the asymptotics of the counting function of the Riemann zeros as in (38) and (39). It would be interesting to note that the equation (48) also counts the large conformal dimensions for the ground state in CQM. Combining the semiclassical realization (48) of the counting Riemann zeros with our proposed holographic correspondence (29) would indicate underlying profound relation among essential ingredients in number theory, in quantum mechanics and in gravity.

## Acknowledgments

The author would like to thank Pei-Ming Ho, Kazuo Hosomichi, Takeo Inami and Dharmesh Jain for communications and discussions and Yu Nakayama for enlightening comments and remarks about Liouville field theory and Michael Berry, Jon Keating, Paul Townsend and especially Germán Sierra for helpful comments and explanations of their works on Riemann zeros. This work was supported by National Taiwan University and the National Center for Theoretical Science (NCTS).

## References

- G. 't Hooft, "Dimensional reduction in quantum gravity," in *Salamfest* 1993:0284-296, pp. 0284–296, 1993. gr-qc/9310026.
- [2] L. Susskind, "The World as a hologram," J. Math. Phys. 36 (1995) 6377-6396, hep-th/9409089.
- J. M. Maldacena, "The Large N limit of superconformal field theories and supergravity," Adv. Theor. Math. Phys. 2 (1998) 231-252, hep-th/9711200.
- [4] H. K. Kunduri, J. Lucietti, and H. S. Reall, "Near-horizon symmetries of extremal black holes," *Class. Quant. Grav.* 24 (2007) 4169–4190, 0705.4214.
- [5] P. Figueras, H. K. Kunduri, J. Lucietti, and M. Rangamani, "Extremal vacuum black holes in higher dimensions," *Phys. Rev.* D78 (2008) 044042, 0803.2998.
- [6] A. Strominger, "AdS(2) quantum gravity and string theory," *JHEP* 9901 (1999) 007, hep-th/9809027.
- [7] J. M. Maldacena, J. Michelson, and A. Strominger, "Anti-de Sitter fragmentation," JHEP 9902 (1999) 011, hep-th/9812073.
- [8] T. Nakatsu and N. Yokoi, "Comments on Hamiltonian formalism of AdS / CFT correspondence," Mod. Phys. Lett. A14 (1999) 147–160, hep-th/9812047.

- [9] P. Townsend, "The M(atrix) model / AdS(2) correspondence," hep-th/9903043.
- M. Spradlin and A. Strominger, "Vacuum states for AdS(2) black holes," JHEP 9911 (1999) 021, hep-th/9904143.
- [11] M. Cadoni and S. Mignemi, "Asymptotic symmetries of AdS(2) and conformal group in d = 1," Nucl. Phys. B557 (1999) 165–180, hep-th/9902040.
- [12] J. D. Blum, "Supersymmetric quantum mechanical description of four-dimensional black holes," JHEP 0001 (2000) 006, hep-th/9907101.
- [13] J. Navarro-Salas and P. Navarro, "AdS(2) / CFT(1) correspondence and near extremal black hole entropy," Nucl. Phys. B579 (2000) 250–266, hep-th/9910076.
- M. Caldarelli, G. Catelani, and L. Vanzo, "Action, Hamiltonian and CFT for 2-D black holes," JHEP 10 (2000) 005, hep-th/0008058.
- [15] M. Cadoni, P. Carta, D. Klemm, and S. Mignemi, "AdS(2) gravity as conformally invariant mechanical system," *Phys. Rev.* D63 (2001) 125021, hep-th/0009185.
- [16] S. Bellucci, A. Galajinsky, E. Ivanov, and S. Krivonos, "AdS(2) / CFT(1), canonical transformations and superconformal mechanics," *Phys.Lett.* B555 (2003) 99–106, hep-th/0212204.
- [17] A. Strominger, "A Matrix model for AdS(2)," JHEP 0403 (2004) 066, hep-th/0312194.
- [18] C. Leiva and M. S. Plyushchay, "Conformal symmetry of relativistic and nonrelativistic systems and Ads / CFT correspondence," Annals Phys. 307 (2003) 372–391, hep-th/0301244.
- [19] A. Giveon and A. Sever, "Strings in a 2-d extremal black hole," JHEP 0502 (2005) 065, hep-th/0412294.
- [20] T. Azeyanagi, T. Nishioka, and T. Takayanagi, "Near Extremal Black Hole Entropy as Entanglement Entropy via AdS(2)/CFT(1)," *Phys.Rev.* D77 (2008) 064005, 0710.2956.
- [21] M. Alishahiha and F. Ardalan, "Central Charge for 2D Gravity on AdS(2) and AdS(2)/CFT(1) Correspondence," JHEP 08 (2008) 079, 0805.1861.

- [22] R. K. Gupta and A. Sen, "Ads(3)/CFT(2) to Ads(2)/CFT(1)," JHEP 0904 (2009) 034, 0806.0053.
- [23] T. Hartman and A. Strominger, "Central Charge for AdS(2) Quantum Gravity," JHEP 04 (2009) 026, 0803.3621.
- [24] A. Galajinsky, "Particle dynamics on AdS(2) x S\*\*2 background with two-form flux," *Phys. Rev.* D78 (2008) 044014, 0806.1629.
- [25] A. Sen, "Quantum Entropy Function from AdS(2)/CFT(1) Correspondence," Int. J. Mod. Phys. A24 (2009) 4225–4244, 0809.3304.
- [26] A. Sen, "Entropy Function and AdS(2) / CFT(1) Correspondence," JHEP 0811 (2008) 075, 0805.0095.
- [27] A. Sen, "State Operator Correspondence and Entanglement in  $AdS_2/CFT_1$ ," Entropy 13 (2011) 1305–1323, 1101.4254.
- [28] C. Chamon, R. Jackiw, S.-Y. Pi, and L. Santos, "Conformal quantum mechanics as the CFT<sub>1</sub> dual to AdS<sub>2</sub>," *Phys.Lett.* B701 (2011) 503–507, 1106.0726.
- [29] J. Molina-Vilaplana and G. Sierra, "An xp model on AdS<sub>2</sub> spacetime," Nucl. Phys. B877 (2013) 107–123, 1212.2436.
- [30] R. Jackiw and S.-Y. Pi, "Conformal Blocks for the 4-Point Function in Conformal Quantum Mechanics," *Phys. Rev.* D86 (2012) 045017, 1205.0443.
- [31] A. B. Zamolodchikov and A. B. Zamolodchikov, "Liouville field theory on a pseudosphere," hep-th/0101152.
- [32] V. Balasubramanian, P. Kraus, and A. E. Lawrence, "Bulk versus boundary dynamics in anti-de Sitter space-time," *Phys. Rev.* D59 (1999) 046003, hep-th/9805171.
- [33] V. Balasubramanian, P. Kraus, A. E. Lawrence, and S. P. Trivedi, "Holographic probes of anti-de Sitter space-times," *Phys. Rev.* D59 (1999) 104021, hep-th/9808017.
- [34] V. Balasubramanian, J. Parsons, and S. F. Ross, "States of a chiral 2d CFT," *Class. Quant. Grav.* 28 (2011) 045004, 1011.1803.
- [35] V. de Alfaro, S. Fubini, and G. Furlan, "Conformal Invariance in Quantum Mechanics," *Nuovo Cim.* A34 (1976) 569.

- [36] A. Connes, "Trace formula in noncommutative geometry and the zeros of the Riemann zeta function," Selecta Math. (N.S.) 5 (1999), no. 1 29–106.
- [37] M. V. Berry and J. P. Keating, "H= xp and the riemann zeros," in Supersymmetry and Trace Formulae, pp. 355–367. Springer, 1999.
- [38] M. V. Berry and J. P. Keating, "The Riemann zeros and eigenvalue asymptotics," SIAM Rev. 41 (1999), no. 2 236–266 (electronic).
- [39] G. Sierra and P. K. Townsend, "Landau levels and Riemann zeros," *Phys. Rev. Lett.* **101** (2008), no. 11 110201, 4.
- [40] G. Sierra and J. Rodriguez-Laguna, "The H=xp model revisited and the Riemann zeros," *Phys.Rev.Lett.* **106** (2011) 200201, **1102.5356**.
- [41] M. V. Berry and J. P. Keating, "A compact Hamiltonian with the same asymptotic mean spectral density as the Riemann zeros," J. Phys. A 44 (2011), no. 28 285203, 14.
- [42] V. Bargmann, "Irreducible unitary representations of the Lorentz group," Annals Math. 48 (1947) 568–640.
- [43] H. Jacquet, "Fonctions de Whittaker associées aux groupes de Chevalley," Bull. Soc. Math. France 95 (1967) 243–309.
- [44] G. Schiffmann, "Intégrales d'entrelacement et fonctions de Whittaker," Bull. Soc. Math. France 99 (1971) 3–72.
- [45] B. Kostant, "On Whittaker vectors and representation theory," *Invent. Math.*48 (1978), no. 2 101–184.
- [46] P. Etingof, "Whittaker functions on quantum groups and q-deformed Toda operators," in *Differential topology, infinite-dimensional Lie algebras, and applications*, vol. 194 of *Amer. Math. Soc. Transl. Ser. 2*, pp. 9–25. Amer. Math. Soc., Providence, RI, 1999.
- [47] N. Seiberg, "Notes on quantum Liouville theory and quantum gravity," *Prog. Theor. Phys. Suppl.* **102** (1990) 319–349.
- [48] B. S. DeWitt, "Quantum Theory of Gravity. 1. The Canonical Theory," Phys. Rev. 160 (1967) 1113–1148.
- [49] C. W. Misner, "MINISUPERSPACE,".

- [50] G. W. Moore, N. Seiberg, and M. Staudacher, "From loops to states in 2-D quantum gravity," Nucl. Phys. B362 (1991) 665–709.
- [51] V. Fateev, A. B. Zamolodchikov, and A. B. Zamolodchikov, "Boundary Liouville field theory. 1. Boundary state and boundary two point function," hep-th/0001012.
- [52] M. Aganagic, R. Dijkgraaf, A. Klemm, M. Marino, and C. Vafa, "Topological strings and integrable hierarchies," *Commun.Math.Phys.* 261 (2006) 451–516, hep-th/0312085.
- [53] A.-K. Kashani-Poor, "The Wave Function Behavior of the Open Topological String Partition Function on the Conifold," JHEP 04 (2007) 004, hep-th/0606112.
- [54] M. C. N. Cheng, R. Dijkgraaf, and C. Vafa, "Non-Perturbative Topological Strings And Conformal Blocks," *JHEP* 09 (2011) 022, 1010.4573.
- [55] C. Beem, T. Dimofte, and S. Pasquetti, "Holomorphic Blocks in Three Dimensions," JHEP 1412 (2014) 177, 1211.1986.
- [56] J. D. Brown and M. Henneaux, "Central Charges in the Canonical Realization of Asymptotic Symmetries: An Example from Three-Dimensional Gravity," *Commun.Math.Phys.* **104** (1986) 207–226.
- [57] S. Gubser, I. R. Klebanov, and A. M. Polyakov, "Gauge theory correlators from noncritical string theory," *Phys.Lett.* B428 (1998) 105–114, hep-th/9802109.
- [58] E. Witten, "Anti-de Sitter space and holography," Adv. Theor. Math. Phys. 2 (1998) 253-291, hep-th/9802150.
- [59] I. S. Gradshteyn and I. Ryzhik, "Table of integrals, series, and products. translated from the russian. translation edited and with a preface by alan jeffrey and daniel zwillinger," 2000.
- [60] H. M. Edwards, *Riemann's zeta function*. Academic Press [A subsidiary of Harcourt Brace Jovanovich, Publishers], New York-London, 1974. Pure and Applied Mathematics, Vol. 58.