

Strong limitations on allowable gauge transformations in electrodynamics

H. R. Reiss

*Max Born Institute, 12489 Berlin, Germany and
American University, Washington, DC 20016-8058, USA**

(Dated: 27 October 2015)

Abstract

Conservation principles establish the primacy of potentials over fields in electrodynamics, both classical and quantum. The contrary conclusion that fields are primary is based on the Newtonian concept that forces completely determine dynamics, and electromagnetic forces depend directly on fields. However, physical conservation principles come from symmetries such as those following from Noether's theorem, and these require potentials for their statement. Examples are given of potentials that describe fields correctly but that violate conservation principles, demonstrating that the correct statement of potentials is necessary. An important consequence is that gauge transformations are severely limited when conservation conditions must be satisfied. When transverse and longitudinal fields are present concurrently, the only practical gauge is the radiation gauge.

*Electronic address: reiss@american.edu

It is widely accepted that the Aharonov-Bohm effect [1, 2] establishes the need for electromagnetic potentials for the proper description of electromagnetic phenomena, but this is a unique effect that is specifically quantum-mechanical. It is demonstrated here that physical conservation principles that exist in both classical and quantum mechanics are sufficient to establish the primacy of potentials over fields for the description of electromagnetic phenomena. Basic examples are shown in which one set of potentials satisfies conservation principles and another does not, even though both sets of potentials predict exactly the same fields. A direct consequence is that physically allowable gauge transformations impose severe limitations on physically valid gauge transformations. An example from strong-field physics shows that problems containing both longitudinal and transverse fields can be described only within the radiation gauge.

The first example is well-known, since it pertains to the description of a charged particle in a static electric field. This is an energy-conserving system, consistent with Noether's theorem that the Lagrangian should be independent of time. However, time-independence is true for only one set of potentials and not a second one that nevertheless correctly describes the fields. The second example to be presented is less well-known. It violates the conservation of the ponderomotive potential energy of a charged particle in a plane-wave field, and it is only recently that this has been demonstrated [3] to be a conserved quantity. Furthermore, potentials that violate the conservation rule about the ponderomotive potential are not widely known, although they have been in the literature [4] for many years.

Despite the familiarity of the example of the static electric field, its implications have not been fully appreciated. The static electric field \mathbf{E}_0 can be described by the scalar and vector potentials

$$\phi = -\mathbf{r} \cdot \mathbf{E}_0, \quad \mathbf{A} = 0, \quad (1)$$

or by the potentials

$$\tilde{\phi} = 0, \quad \tilde{\mathbf{A}} = -ct\mathbf{E}_0. \quad (2)$$

(Gaussian units are used throughout.) A charged particle in a constant electric field is a system that is conservative, which is consistent with the time-independent potentials (1), but the time-dependent potentials (2) are unphysical because Noether's Theorem show that they violate the energy conservation principle. Starting from the potentials (1), it has been shown [5] that there is no possible gauge transformation that can preserve energy

conservation. That is, no gauge freedom at all exists for this case.

The uniqueness of the potentials (1) is easily demonstrated. The general gauge transformation is given by

$$\tilde{A}^\mu = A^\mu + \partial^\mu \Lambda, \quad (3)$$

where Λ is a scalar function that satisfies the homogeneous wave equation

$$\partial^\mu \partial_\mu \Lambda = 0. \quad (4)$$

The covariant expression (3) can be devolved into scalar and vector potentials as

$$\tilde{\phi} = \phi + \frac{1}{c} \partial_t \Lambda, \quad (5)$$

$$\tilde{\mathbf{A}} = \mathbf{A} - \nabla \Lambda. \quad (6)$$

Starting with the potentials (1), Eq. (5) shows that the necessity to keep potentials independent of time means that Λ can, at most, have a dependence that is linear in t , leading to the form $\Lambda = tf(\mathbf{r})$. However, this would generate time dependence in $\tilde{\mathbf{A}}$. Therefore, dependence of Λ is limited to the form $\Lambda(\mathbf{r})$. In that case, Eq. (1) provides a complete description of the field, meaning that $\Lambda = \text{constant}$ is the only possibility. This is a trivial result, with the significance that the potentials (1) are the unique potentials for the description of a constant electric field.

The gauge transformation connecting Eqs. (1) and (2) is just the Göppert-Mayer gauge transformation [6] applied in the zero frequency limit. The unphysical nature of the potentials (2) has consequences for strong-field physics that are explained in Ref. [7].

The second example relates to the very important case of propagating fields that may, equivalently, be referred to as transverse fields or plane-wave (PW) fields. Their profound importance stems from the widespread use of lasers in laboratory experiments, and lasers produce PW fields. The unique ability of PW fields to propagate without sources means that the fields that impinge on a target in a laser experiment can only be a superposition of PW fields. Any “contamination” by fields other than pure PW fields damps out over short distances, of the order of wavelengths of the laser field.

A basic limitation on possible gauge transformations that preserve the identity of PW fields follows immediately from the premise of special relativity that all inertial frames of reference are equivalent. This is expressible by the constraint that the spacetime 4-vector

x^μ can occur only in the Lorentz-scalar combination [8, 9]

$$\varphi \equiv k^\mu x_\mu = \omega t - \mathbf{k} \cdot \mathbf{r}, \quad (7)$$

where k^μ is the propagation 4-vector of the PW field, and \mathbf{k} is the propagation 3-vector. The quantity φ is just the phase of a propagating field. The propagation 4-vector lies on the light cone, meaning that

$$k^\mu k_\mu = 0. \quad (8)$$

A gauge transformation that preserves the basic condition for a PW field is thus constrained by the limitation that the generator of the gauge transformation can depend on x^μ only in the form of φ [3, 5], meaning that

$$\tilde{A}^\mu = A^\mu + \partial^\mu \Lambda = A^\mu + (\partial^\mu \varphi) \frac{d}{d\varphi} \Lambda = A^\mu + k^\mu \Lambda', \quad (9)$$

where Λ' is the total derivative of Λ with respect to φ . A direct consequence of Eq. (9) is that

$$\tilde{A}^\mu \tilde{A}_\mu = A^\mu A_\mu, \quad (10)$$

following from Eq. (8) and from the transversality property

$$k^\mu A_\mu = 0. \quad (11)$$

This has the basic physical significance that the ponderomotive potential U_p of a particle of charge q and mass m immersed in a PW field, defined as

$$U_p \equiv \frac{q^2}{2mc^2} (-A^\mu A_\mu), \quad (12)$$

is gauge-invariant under any gauge transformation that preserves the PW property of the field. It is also, from its definition, Lorentz invariant. The minus sign in Eq. (12) is included to make U_p a positive quantity, since the 4-vector A^μ is a spacelike 4-vector, whose square is negative in terms of the time-favoring real metric employed here. Two important facts are that the period average generally employed in the definition of U_p is needlessly limiting, and is not necessary; and that U_p defined for a PW field is a true potential energy [3] and not the kinetic “quiver energy” associated with oscillatory electric fields.

One of the important properties, just demonstrated in Eq. (9), is that any gauge transformation applied to a PW field is limited to an alteration of the 4-potential by an additive

function lying on the light cone. This limitation is strong in the sense that adding a light-cone contribution does not confer any simplifications in a practical problem. If the vector potential A^μ for the PW field is expressed in the radiation gauge, where $A^0 = 0$, there is no practical advantage to departing from this simple choice. The radiation gauge (often called the Coulomb gauge) thus has the convenient property that transverse fields are represented by 3-vector potentials, and longitudinal fields are represented by scalar potentials.

There is another reason to limit the possible gauges for the expression of a PW field to the unique choice of the radiation gauge when a PW field is applied in the concurrent presence of a longitudinal field, such as a Coulomb field. When the PW field is very strong, its fundamentally relativistic nature can be viewed as requiring the entire problem of an atom or molecule in the strong PW field to be basically relativistic. If the atomic “electron” in such a problem is viewed as a spinless particle, as is usually done in AMO physics, then the Klein-Gordon equation is the appropriate quantum dynamical equation. The Klein-Gordon equation for an “electron” in a strong laser field was examined in Ref. [10] with the result that a cross-coupling term between longitudinal and transverse fields was found of the form $2VeA^0$, where V is a Coulomb potential and \mathbf{A}^0 is the time part of the 4-vector PW potential. This cross coupling would survive into the nonrelativistic limit, which is the Schrödinger equation. The Schrödinger equation does not have such a term, which constitutes a requirement that the radiation gauge must be selected for the PW field. Stated briefly: the concurrent existence of a longitudinal field (like the Coulomb potential) and a transverse field (a laser field) limits gauge selection to the sole physical possibility of the radiation gauge.

Now the second example will be examined of a gauge transformation that results in an unphysical set of potentials despite these potentials predicting the fields correctly and satisfying all of the normal requirements for carrying out a gauge transformation. To examine this problem, we temporarily set aside the requirement that x^μ can occur only in the combination φ of Eq. (7). The example concerns the fundamentally important propagating field (or transverse field or PW field). The generating function

$$\Lambda_K = -x^\nu A_\nu(\varphi) \tag{13}$$

is introduced. The subscript “K” is intended to stand for “Keldysh”, since the original introduction of the this transformation [4] was in an (unsuccessful) attempt to place the

well-known Keldysh approximation [11] on a foundation that referred to a laser field rather than to the oscillatory electric field treated by Keldysh. The 4-vector potential so obtained from the gauge transformation (3) produces the result

$$\tilde{A}_K^\mu = -k^\mu [x^\nu A'_\nu(\varphi)], \quad (14)$$

where A'_ν refers to the total derivative of A_ν with respect to φ . The significance of this result becomes more clear if the original 4-potential A^μ is in the radiation gauge, in which case Eq. (14) can be written as [4]

$$\tilde{A}_K^\mu = -\frac{k^\mu}{\omega/c} [\mathbf{r} \cdot \mathbf{E}(\varphi)], \quad (15)$$

which has the time component

$$\tilde{A}_K^0 \equiv \phi_K = -\mathbf{r} \cdot \mathbf{E}(\varphi). \quad (16)$$

The covariant expression for the 4-potential (14) then appears to be a relativistic generalization of the ordinary length-gauge scalar potential. That this is not the case will shortly become clear.

It can be shown readily [4] that Λ_K satisfies the homogeneous wave equation as specified in Eq. (4), the 4-vector potential obtained from the transformation (13) satisfies the transversality condition

$$k_\mu \tilde{A}_K^\mu = 0, \quad (17)$$

and it is also a *Lorentz gauge*, meaning that

$$\partial_\mu \tilde{A}_K^\mu = 0. \quad (18)$$

In other words, all of the requirements attached to a gauge transformation are satisfied. Furthermore, the potentials (14) produce exactly the correct PW electric and magnetic fields; perpendicular to each other and perpendicular to the propagation direction. That is, the gauge transformation is legitimate in that it satisfies all the usual conditions imposed on gauge transformations and it reproduces the desired electromagnetic fields exactly.

However, the gauge-transformed 4-vector potential of Eq. (14) is unphysical.

One defect is immediately clear: \tilde{A}_K^μ is lightlike rather than spacelike. In consequence, the importantly gauge-invariant ponderomotive potential is predicted to be exactly zero in the gauge of Eq. (14). The cause of these problems is directly traceable to the fact that the

generating function (13) introduces a dependence on the spacetime 4-vector x^μ that is not in the form of $k^\mu x_\mu$, thus violating a basic premise of special relativity for the propagation of a light wave.

A dependence on physical reasoning based entirely on the fields and not on potentials is consequential. An important example comes from Atomic, Molecular, and Optical (AMO) physics, where it is customary in the description of laser-produced phenomena to introduce what is generally called the “dipole approximation”, viewed in terms of fields as the statement that the electric field is a function of time only, with no spatial dependence, and where the magnetic field is neglected altogether:

$$\mathbf{E} = \mathbf{E}(t), \quad \mathbf{B} = 0. \quad (19)$$

This is seemingly plausible as long as the wavelength of the field is much longer than atomic dimensions. However, this criterion can be seriously misleading when very strong laser fields are present. For example, the fields (19) do not obey the same Maxwell equations as those for propagating fields [12], which means that there is no possible gauge connection to propagating fields (despite the widespread assumption that there is such a connection). The dipole fields (19) have led to the practice of subjecting both analytical and numerical calculations to the test that behavior should replicate that of a constant electric field in the zero-frequency limit. This is a basic premise, for example, in a textbook on strong-field physics [13], which reflects the frequently-encountered use of this criterion in the AMO community. However, from the point of view of potentials, it is clear that the fields of (19) are incompatible with the requirements for a propagating field. An electric field of zero frequency certainly does not have the propagation property, and the zero-frequency limit of a propagating field is simply an extremely long-wavelength radio field. Thus, the requirement that there should be a zero-frequency limit for laser effects that is a constant electric field is fundamentally unphysical.

Another widespread concept is that laser-caused atomic ionization is a tunneling process wherein the attractive Coulomb potential of an electron bound in an atom can be wholly or partially compromised by the addition of a scalar potential representing an oscillatory electric field. This is a process that is completely dependent on the superposition of two longitudinal fields and cannot be representative of the interaction of a transverse field with a Coulomb potential. The tunneling model has some success, but it is plainly confined to a

limited domain of frequencies and intensities of the laser field. In particular, the “tunneling limit” where the Keldysh parameter

$$\gamma_K = \sqrt{E_B/2U_p} \quad (20)$$

(where E_B is the field-free binding energy of the electron) becomes extremely small ($\gamma_K \ll 1$), is actually a domain where the magnetic field becomes important, the fields of Eq. (19) do not properly represent the laser environment, and conclusions based on the tunneling concept are inappropriate [14, 15].

The principal conclusions reached above can be summarized:

Electromagnetic fields are more fundamental than the fields that can be derived from them in both classical and quantum electrodynamics.

There exist potentials that predict fields correctly but violate physical conservation laws.

Gauge transformations that connect physically possible gauges are severely limited.

For each of the important special cases of constant electric fields and of laser fields (i.e., propagating fields) there exists only one physical gauge.

Methods based on the electric dipole approximation of AMO physics can support unphysical conclusions about strong laser fields.

-
- [1] W. Ehrenberg and R. E. Siday, Proc. Roy. Soc. B **62**, 8 (1949).
 - [2] Y. Aharonov and D. Bohm, Phys. Rev. **115**, 485 (1959).
 - [3] H. R. Reiss, Phys. Rev. A **89**, 022116 (2014).
 - [4] H. R. Reiss, Phys. Rev. A **19**, 1140 (1979).
 - [5] H. R. Reiss, J. Mod. Opt. **59**, 1371 (2012); **60**, 687 (2013).
 - [6] M. Göppert-Mayer, Ann. Phys. (Leipzig) **9**, 273 (1931).
 - [7] H. R. Reiss, arXiv:1509.02764v1 [quant-ph] 9 Sep 2015.
 - [8] J. Schwinger, Phys. Rev. **82**, 664 (1951).
 - [9] E. S. Sarachik and G. T. Schappert, Phys. Rev. D **1**, 2738 (1970).
 - [10] H. R. Reiss, Phys. Rev. A **42**, 1476 (1990).
 - [11] L. V. Keldysh, Zh. Eksp. Teor. Fiz. **47**, 1945 (1964) [Sov. Phys. JETP **20**, 1307 (1965)].
 - [12] H. R. Reiss, J. Phys. B **46**, 175601 (2013).

- [13] C. J. Joachain, N. J. Kylstra, and R. M. Potvliege, *Atoms in Intense Laser Fields* (Cambridge Univ. Press, Cambridge, 2012).
- [14] H. R. Reiss, Phys. Rev. Lett. **101**, 043002 (2008); **101**, 159901(E) (2008).
- [15] H. R. Reiss, J. Phys. B **47**, 204006 (2014).