

PARALLEL METHODS FOR REGULARIZING SYSTEMS OF EQUATIONS INVOLVING ACCRETIVE OPERATORS

Pham Ky Anh*

Department of Mathematics, Hanoi University of Science, Hanoi, Vietnam

Nguyen Buong

Institute of Information Technology, Vietnamese Academy of Science & Technology

Dang Van Hieu

Department of Mathematics, Hanoi University of Science, Hanoi, Vietnam

□ *In this paper, two parallel methods for solving systems of accretive operator equations in Banach spaces are studied. The convergence analysis of the methods in both free-noise and noisy data cases is provided.*

Keywords: Uniformly smooth and uniformly convex Banach space; Accretive and inverse uniformly accretive operators; Iterative regularization method; Newton-type method; Parallel computation.

AMS Subject Classifications 47J06; 47J25; 65J15; 65J20; 65Y05.

1. INTRODUCTION, PRELIMINARIES, AND NOTATIONS

Various problems of science and engineering, including a multi-parameter identification problem, the convex feasibility problem, a common fixed point problem, etc..., lead to a system of ill-posed operator equations

$$A_i(x) = 0, \quad x \in X, (i = 1, 2, \dots, N), \quad (1.1)$$

where X is a real Banach space and $A_i : D(A_i) = X \rightarrow X$ are possibly nonlinear operators on X .

Very recently, several sequential and parallel regularizing methods for solving system (1.1) have been proposed. The Kaczmarz method [1, 2], the Newton-Kaczmarz method [3], the steepest-descent-Kaczmarz method [4], parallel iterative regularization methods [5], parallel regularized Newton-type methods [6, 7], parallel hybrid methods [8], to name only few. However, most of the investigation of available methods was carried out in the framework of Hilbert spaces.

In this paper we study parallel methods extended to system (1.1) involving m -accretive operators in the setting of Banach spaces. In the sequel we always assume that system (1.1) is consistent, i.e., the solution set S of (1.1) is not empty. It is known that if $A_i (i = 1, \dots, N)$ are not strongly or uniformly accretive, then system (1.1) in general is ill-posed, i.e., the solution set S of (1.1) may not depend continuously on data. In that case, a process known as regularization should be applied for stable solution of (1.1).

In what follows, for the reader's convenience, we collect some definitions and results concerning the geometry of Banach spaces and accretive operators, which are used in this paper. We refer the reader to [9-13] for more details.

*Corresponding author: Pham Ky Anh, Department of Mathematics, Hanoi University of Science, 334 Nguyen Trai, Thanh Xuan, Hanoi, Vietnam. Email address: phamkyanh@hus.edu.vn.

Definition 1.1. A Banach space X is called

- 1) *strictly convex* if the unit sphere $S_1(0) = \{x \in X : \|x\| = 1\}$ is strictly convex, i.e., the inequality $\|x + y\| < 2$ holds for all $x, y \in S_1(0), x \neq y$;
- 2) *uniformly convex* if for any given $\epsilon > 0$ there exists $\delta = \delta(\epsilon) > 0$ such that for all $x, y \in X$ with $\|x\| \leq 1, \|y\| \leq 1, \|x - y\| = \epsilon$ the inequality $\|x + y\| \leq 2(1 - \delta)$ holds.

The modulus of convexity of X is defined by

$$\delta_X(\epsilon) = \inf \left\{ 1 - \frac{\|x - y\|}{2} : \|x\| = \|y\| = 1, \|x - y\| = \epsilon \right\}.$$

The modulus of smoothness of X is defined by

$$\rho_X(\tau) = \sup \left\{ \frac{\|x + y\| + \|x - y\|}{2} - 1 : \|x\| = 1, \|y\| = \tau \right\}.$$

Definition 1.2. A Banach space X is called *uniformly smooth* if

$$\lim_{\tau \rightarrow 0} h_X(\tau) := \lim_{\tau \rightarrow 0} \frac{\rho_X(\tau)}{\tau} = 0.$$

Observe that if X is a real uniformly convex and uniformly smooth Banach space, then the modulus of convexity δ_X is a continuous and strictly increasing function on the whole segment $[0, 2]$ (see, for example [14]).

Definition 1.3. A Banach space X possesses the *approximation* if there exists a directed family of finite dimensional subspaces X_n ordered by inclusion, and a corresponding family of projectors $P_n : X \rightarrow X_n$, such that $\|P_n\| = 1$ for all $n \geq 0$ and $\cup_n X_n$ is dense in X .

Throughout this paper we assume that the so-called normalized duality mapping $J : X \rightarrow X^*$, satisfying the relation

$$\langle x, J(x) \rangle = \|x\|^2 = \|J(x)\|^2, \quad \forall x \in X,$$

is single valued. This assumption will be fulfilled if X is smooth.

For the sake of simplicity, we will denote norms of both spaces X and X^* by the same symbol $\|\cdot\|$. The dual product of $f \in X^*$ and $x \in X$ will be denoted by $\langle x, f \rangle$ or $\langle f, x \rangle$. Besides, we put $\mathbb{R}^+ := (0, \infty), \mathbb{R}_*^+ := [0, \infty)$.

Definition 1.4. An operator $A : X \rightarrow X$ is called

- 1) *accretive*, if

$$\langle A(x) - A(y), J(x - y) \rangle \geq 0 \quad \forall x, y \in X;$$
- 2) *maximal accretive*, if it is accretive and its graph is not the right part of the graph of any other accretive operator;
- 3) *m-accretive*, if it is accretive and $R(A + \alpha I) = X$ for all $\alpha > 0$, where I is the identity operator in X ;
- 4) *uniformly accretive*, if there exists a strictly increasing function $\psi : \mathbb{R}_*^+ \rightarrow \mathbb{R}_*^+, \psi(0) = 0$, such that

$$\langle A(x) - A(y), J(x - y) \rangle \geq \psi(\|x - y\|) \quad \forall x, y \in X; \quad (1.2)$$

- 5) strongly accretive, if there exists a positive constant c , such that in (1.2), $\psi(t) = ct^2$;
- 6) inverse strongly accretive, if there exists a positive constant c , such that

$$\langle A(x) - A(y), J(x - y) \rangle \geq c \|A(x) - A(y)\|^2 \quad \forall x, y \in X.$$

If X is a Hilbert space then J is an identity operator and accretive operators are also called monotone.

Definition 1.5. A continuous operator A mapping a Banach space X into itself is called φ -inverse uniformly accretive (or simply, inverse uniformly accretive), if there exists a function $\varphi : \mathbb{R}^+ \times \mathbb{R}_*^+ \rightarrow \mathbb{R}_*^+$, which is continuous and strictly increasing in the second variable and $\varphi(s, t) = 0$ if and only if $t = 0$ for every fixed $s > 0$, such that

$$\langle A(x) - A(y), J(x - y) \rangle \geq \varphi(R, \|A(x) - A(y)\|) \quad \forall x, y \in X, \|x\|, \|y\| \leq R, \forall R > 0. \quad (1.3)$$

Example 1. Any inverse strongly accretive operator is inverse uniformly accretive, hence is accretive. Indeed, let A be a c -inverse strongly accretive operator. Then A is Lipschitz continuous with the Lipschitz constant c^{-1} and the inequality (1.3) holds for the function $\varphi(s, t) = ct^2$.

Example 2. Let T be a nonexpansive operator on a uniformly convex and uniformly smooth Banach space X . Then $A := I - T$ is a Lipschitz continuous operator. Moreover, according to Alber [15],

$$\langle A(x) - A(y), J(x - y) \rangle \geq L^{-1}R^2\delta_X \left(\frac{\|A(x) - A(y)\|}{4R} \right) \quad \forall x, y \in X, \|x\|, \|y\| \leq R,$$

where $L \in (1; 1.7)$ is the Figiel constant and $\delta_X(\epsilon)$ is the modulus of the convexity of X . Observe that $\epsilon := \frac{\|A(x) - A(y)\|}{4R} \leq 1$ for any $x, y \in X; \|x\|, \|y\| \leq R$ and inequality (1.3) holds for the function $\varphi(s, t) = L^{-1}s^2\delta_X\left(\frac{t}{4s}\right), s \in \mathbb{R}^+; t \in [0; 2s]$.

Example 3. Now let X in Example 2 be one of the following Banach spaces L^p, l^p, W_p^m , where $1 < p < \infty$. Then X is uniformly smooth and uniformly convex, and it is well-known that (see [9])

$$\begin{aligned} \delta_X(t) &\geq \frac{p-1}{16}t^2, \quad 1 < p < 2; \\ \delta_X(t) &\geq \frac{1}{p2^p}t^p, \quad p \geq 2. \end{aligned}$$

Thus, for all $x, y \in X, \|x\|, \|y\| \leq R$, one gets

$$\begin{aligned} \langle A(x) - A(y), J(x - y) \rangle &\geq \frac{p-1}{256L} \|A(x) - A(y)\|^2, \quad 1 < p < 2; \\ \langle A(x) - A(y), J(x - y) \rangle &\geq \frac{1}{Lp8^p} \frac{\|A(x) - A(y)\|^p}{R^{p-2}}, \quad p \geq 2. \end{aligned}$$

So, the operator $A = I - T$, where $T : l^p \rightarrow l^p$ is a nonexpansive operator, is inverse strongly accretive if $1 < p < 2$, and is inverse uniformly accretive with $\varphi(s, t) = \frac{t^p}{pL8^p s^{p-2}}$, if $p \geq 2$.

Definition 1.6. An operator, $B : D(B) \subset X \rightarrow X$ is called

- 1) *hemicontinuous at a point* $x_0 \in D(B)$, if $B(x_0 + t_n h) \rightarrow x_0$ as $t_n \rightarrow 0$ for any vector h such that $x_0 + t_n h \in D(B)$ and $0 \leq t_n \leq t(x_0)$;
- 2) *weakly continuous at* $x_0 \in D(B)$, if $D(B) \ni x \rightarrow x_0$ implies that $B(x) \rightarrow B(x_0)$.

If B is hemicontinuous (weakly continuous) at every point of $D(B)$, then B is said to be hemicontinuous (weakly continuous), respectively.

For regularizing accretive operator equations one needs the following fact [9].

Lemma 1.1. *Suppose that the Banach space X possesses the approximation, $A : X \rightarrow X$ is a hemicontinuous accretive operator with $D(A) = X$, and the normalized duality mapping $J : X \rightarrow X^*$ is sequentially weakly continuous and continuous. Then the problem*

$$A(x) + \alpha x = y, \quad (1.4)$$

where α is a fixed positive parameter and $y \in X$, is well-posed.

The unique solvability of (1.4) is established in [9]. The continuous dependence of the solution x_α of (1.4) on the right-hand side y follows from the inequality $\|x_{\alpha,1} - x_{\alpha,2}\| \leq \frac{\|y_1 - y_2\|}{\alpha}$, where $x_{\alpha,i}$ are the unique solution of (1.4) with respect to the right-hand side $y = y_i, i = 1, 2$.

The next five lemmas will be used in Section 2 for establishing the convergence of implicit and explicit parallel iterative regularization methods.

Lemma 1.2. [16] *Let X be a real uniformly smooth Banach space. Then for any $x, y \in X$ such that $\|x\| \leq R, \|y\| \leq R$, the following inequality holds:*

$$\|J(x) - J(y)\| \leq 8Rh_X \left(\frac{16L \|x - y\|}{R} \right),$$

where L is Figiel constant, $(1 < L < 1.7)$.

Lemma 1.3. [9] *If X is a real uniformly smooth Banach space, then the inequality*

$$\begin{aligned} \|x\|^2 &\leq \|y\|^2 + 2 \langle x - y, J(x) \rangle \\ &\leq \|y\|^2 + 2 \langle x - y, J(y) \rangle + 2 \langle x - y, J(x) - J(y) \rangle \end{aligned}$$

holds for every $x, y \in X$.

Lemma 1.4. [9] *Let X be a uniformly smooth Banach space. Then for $x, y \in X$*

$$\langle x - y, J(x) - J(y) \rangle \leq 8 \|x - y\|^2 + C(\|x\|, \|y\|) \rho_X(\|x - y\|),$$

where $C(\|x\|, \|y\|) \leq 4 \max\{2L, \|x\| + \|y\|\}$.

Lemma 1.5. [9] *In a uniformly smooth Banach space X , for $x, y \in X$,*

$$\langle x - y, J(x) - J(y) \rangle \leq R^2(\|x\|, \|y\|) \rho_X \left(\frac{4 \|x - y\|}{R(\|x\|, \|y\|)} \right),$$

where $R(\|x\|, \|y\|) = \sqrt{2^{-1}(\|x\|^2 + \|y\|^2)}$.

If $\|x\| \leq R, \|y\| \leq R$, then

$$\langle x - y, J(x) - J(y) \rangle \leq 2LR^2 \rho_X \left(\frac{4 \|x - y\|}{R} \right).$$

Lemma 1.6. [9, 17] Let $\{\lambda_n\}$ and $\{p_n\}$ be sequences of nonnegative numbers, $\{b_n\}$ be a sequence of positive numbers, satisfying the inequalities

$$\lambda_{n+1} \leq (1 - p_n) \lambda_n + b_n, \quad \forall n \geq 0,$$

where $p_n \in (0; 1)$, $\frac{b_n}{p_n} \rightarrow 0$ ($n \rightarrow +\infty$) and $\sum_{i=1}^{\infty} p_n = +\infty$. Then $\lambda_n \rightarrow 0$ ($n \rightarrow +\infty$).

In Section 3, when dealing with a parallel Newton-type regularization method, we need some more results.

Lemma 1.7. [18] Suppose $A : D(A) = X \rightarrow X$ is a continuously Fréchet differentiable accretive operator and let $L := A'(h)$, $h \in X$, and α be a real positive number. Then

$$\|(\alpha I + L)^{-1}\| \leq \frac{1}{\alpha}; \quad \|(\alpha I + L)^{-1} L\| \leq 2.$$

Lemma 1.8. [7, 19] Let $\{\omega_n\}$ be a sequence of nonnegative numbers satisfying the relations

$$\omega_{n+1} \leq a + b\omega_n + c\omega_n^2, \quad n \geq 0,$$

for some $a, b, c > 0$.

Let $M_+ := (1 - b + \sqrt{(1 - b)^2 - 4ac})/2c$, $M_- := (1 - b - \sqrt{(1 - b)^2 - 4ac})/2c$. If $b + 2\sqrt{ac} < 1$ and $\omega_0 \leq M_+$, then $\omega_n \leq l := \max\{\omega_0, M_-\}$ for all $n \geq 0$.

An outline of the remainder of the paper is as follows: In Section 2 we propose two parallel iterative regularizations methods (PIRMs) for system (1.1), namely implicit PIRM and explicit PIRM. The convergence of these PIRMs is established for both exact and noisy data cases. Section 3 studies a parallel regularizing Newton-type method for system (1.1). The convergence analysis of the proposed method in exact and noisy data cases is also studied.

2. EQUATIONS WITH INVERSE UNIFORMLY ACCRETIVE OPERATORS

In this section, we consider system (1.1) with inverse uniformly accretive operators. Clearly, if each operator A_i is φ_i -inverse uniformly accretive, then it is φ -inverse uniformly accretive with $\varphi(s, t) := \min_{i=1, \dots, N} \varphi_i(s, t)$. Thus, without loss of generality we can assume that all the operators $A_i, i = 1, \dots, N$ are φ -inverse uniformly accretive with the same function φ .

We begin with the following simple fact (cf. [6]).

Lemma 2.1. Suppose $A_i, i = 1, 2, \dots, N$, are inverse uniformly accretive operators. If system (1.1) is consistent, then it is equivalent to the operator equation

$$A(x) := \sum_{i=1}^N A_i(x) = 0. \quad (2.1)$$

Proof. Let the operators $A_i, i = 1, \dots, N$, be φ -inverse uniformly accretive with the same function φ . Obviously, any solution of (1.1) is a solution of (2.1). Conversely, let y be a solution of (2.1), i.e.,

$$\sum_{i=1}^N A_i(y) = 0.$$

Let z be a solution of system (1.1), i.e., $A_i(z) = 0, i = 1, 2, \dots, N$. Then, $\sum_{i=1}^N (A_i(y) - A_i(z)) = 0$, and since A_i are inverse uniformly accretive, one gets

$$\sum_{i=1}^N \varphi(R, \|A_i(y) - A_i(z)\|) \leq \sum_{i=1}^N \langle A_i(y) - A_i(z), J(y - z) \rangle = 0,$$

where $R = \max\{\|y\|, \|z\|\}$. Thus $\varphi(R, \|A_i(y) - A_i(z)\|) = 0$, hence $A_i(y) = A_i(z) = 0, i = 1, 2, \dots, N$. Therefore, y is a solution of system (1.1). \square

In this section we need the following result .

Lemma 2.2. ([20], Theorem 2.1). *Let X be a real, reflexive and strictly convex Banach space with a uniformly Gateaux differentiable norm and let A be an m -accretive mapping on X . Then for each $\alpha > 0$ and a fixed $y \in X$, equation (1.4) possesses a unique solution x_α , and in addition, if the solution set S_A of the equation $A(x) = y$ is nonempty, then the net $\{x_\alpha\}$ converges strongly to the unique element \hat{x}^* solving the following variational inequality*

$$\langle \hat{x}^*, J(\hat{x}^* - x^*) \rangle \leq 0, \forall x^* \in S_A.$$

Moreover we have $\|x_\alpha^\delta - x_\alpha\| \leq \delta/\alpha$, where x_α^δ is the unique solution of the equation $A(x) + \alpha x = y_\delta$, for any $\alpha > 0$ and $y_\delta \in X$ satisfying $\|y_\delta - y\| \leq \delta$.

In the remainder of Section 2, we impose two sets of conditions on the space X , the duality mapping J , and the operators $A_i, i = 1, 2, \dots, N$.

Conditions (AJX)

- A1. $A_i, i = 1, 2, \dots, N$, are φ -inverse uniformly accretive operators with $D(A_i) = X$;
- A2. The normalized duality mapping J is sequentially weakly continuous and continuous;
- A3. X is a smooth and reflexive Banach space, possessing the approximation.

Conditions (AX)

- B1. $A_i, i = 1, 2, \dots, N$, are m -accretive and φ -inverse uniformly accretive operators with $D(A_i) = X$;
- B2. X is a uniformly smooth and uniformly convex Banach space.

Together with equation (2.1) we consider the following regularized one

$$A(x) + \alpha_n x = \sum_{i=1}^N A_i(x) + \alpha_n x = 0. \quad (2.2)$$

Lemma 2.3. *Let conditions A1-A3 or B1-B2 be fulfilled. Then the following statements hold:*

- i) For every $\alpha_n > 0$, equation (2.2) has a unique solution x_n^* .
- ii) $\|x_n^*\| \leq 2\|\hat{x}\|$, where \hat{x} is an arbitrary element of S .
- iii) $x_n^* \rightarrow \hat{x}^*$ as $n \rightarrow +\infty$, where \hat{x}^* is an unique solution of the inequality $\langle \hat{x}^*, J(\hat{x}^* - x^*) \rangle \leq 0, \forall x^* \in S$.
- iv) $\|x_n^* - x_{n+1}^*\| \leq 2\|\hat{x}^*\| \frac{|\alpha_{n+1} - \alpha_n|}{\alpha_n}$.
- v) $\|A_i(x_n^*)\| \leq \varphi_R^{-1} \left(6\alpha_n \|\hat{x}^*\|^2 \right) \quad i = 1, 2, \dots, N$, where $R > 0$ is a fixed number satisfying an a-priori estimate $R \geq 2\|\hat{x}^*\|$ and φ_s^{-1} denotes the inverse function of $\varphi(s, t)$ with respect to the second variable t for fixed $s > 0$.

Proof. 1. Suppose that conditions A1-A3 hold. We perform the regularization process (2.2) for equation (2.1) with the accretive operator $A = \sum_{i=1}^N A_i$. For the proofs of statements i) – iv) we refer the reader to [9]. Concerning the last part v) we observe that

$$\sum_{i=1}^N (A_i(x_n^*) - A_i(\hat{x}^*)) + \alpha_n x_n^* = 0,$$

hence,

$$\sum_{i=1}^N \langle A_i(x_n^*) - A_i(\hat{x}^*), J(x_n^* - \hat{x}^*) \rangle + \alpha_n \langle x_n^*, J(x_n^* - \hat{x}^*) \rangle = 0.$$

Observing that by part ii) $\|x_n^*\| \leq 2\|\hat{x}^*\|$, hence, $\|x_n^* - \hat{x}^*\| \leq 3\|\hat{x}^*\|$ and using the inverse uniform accretiveness of A_i , from the last inequality we have

$$\sum_{i=1}^N \varphi(R, \|A_i(x_n^*) - A_i(\hat{x}^*)\|) \leq -\alpha_n \langle x_n^*, J(x_n^* - \hat{x}^*) \rangle \leq \alpha_n \|x_n^*\| \cdot \|x_n^* - \hat{x}^*\|,$$

where $R \geq 2\|\hat{x}^*\|$. The last inequality gives $\varphi(R, \|A_i(x_n^*)\|) \leq 6\alpha_n \|\hat{x}^*\|^2$. Thus $\|A_i(x_n^*)\| \leq \varphi_R^{-1}(6\alpha_n \|\hat{x}^*\|^2)$.

2. Now suppose that conditions B1-B2 hold. Since all A_i are inverse uniformly accretive, they are continuous, hence locally bounded. Besides, $A_i, i = 1, \dots, N$, are m -accretive, $D(A_i) = X$, and the spaces X and X^* are uniformly convex, then by Theorem 1.15.22 ([9]), the operator $A = \sum_{i=1}^N A_i$ is also m -accretive. Lemma 2.2 applied to equation (2.2) ensures the convergence of regularized solutions x_n^* to \hat{x}^* . The remaining statements can be argued similarly as in part 1. \square

Following [5] we consider an implicit PIRM consisting of solving simultaneously N regularized equations

$$A_i(x_n^i) + \left(\frac{\alpha_n}{N} + \gamma_n\right)x_n^i = \gamma_n x_n, i = 1, 2, \dots, N, \quad (2.3)$$

where $\alpha_n > 0$ and $\gamma_n > 0$ are regularization and parallel splitting up parameters, respectively, and defining the next approximation as an average of the regularized solutions x_n^i ,

$$x_{n+1} = \frac{1}{N} \sum_{i=1}^N x_n^i, \quad n = 0, 1, \dots, \quad x_0 \in X. \quad (2.4)$$

According to Lemmas 2.2, 2.3, all the problems (2.3) are well posed and independent from each other, hence the regularized solutions x_n^i can be found stably and simultaneously by parallel processors.

We first prove the boundedness of the sequence $\{x_n\}$ defined by the implicit PIRM (2.3)-(2.4).

Lemma 2.4. *Under conditions A1-A3 or B1-B2, the sequence $\{x_n\}$ generated by (2.3) and (2.4) is bounded.*

Proof. By Lemma 1.1 (Lemma 2.2), the regularized equation (2.3) has a unique solution denoted by x_n^i . Let $B_r(\hat{x}^*)$ be the closed ball with center \hat{x}^* and radius r . Choose $r > 0$ sufficiently large such that $r \geq \|\hat{x}^*\|$ and $x_0 \in B_r(\hat{x}^*)$. Supposing for some $n > 0$, $x_n \in B_r(\hat{x}^*)$, we will show that $x_{n+1} \in B_r(\hat{x}^*)$. Indeed, from (2.3) and $A_i(\hat{x}^*) = 0$, we get

$$(A_i(x_n^i) - A_i(\hat{x}^*)) + \left(\frac{\alpha_n}{N} + \gamma_n\right)(x_n^i - \hat{x}^*) = \gamma_n(x_n - \hat{x}^*) - \frac{\alpha_n}{N}\hat{x}^*.$$

Thus

$$\begin{aligned} & \left\langle A_i(x_n^i) - A_i(\widehat{x}^*), J(x_n^i - \widehat{x}^*) \right\rangle + \left(\frac{\alpha_n}{N} + \gamma_n \right) \left\langle x_n^i - \widehat{x}^*, J(x_n^i - \widehat{x}^*) \right\rangle \\ & = \gamma_n \left\langle x_n - \widehat{x}^*, J(x_n^i - \widehat{x}^*) \right\rangle - \frac{\alpha_n}{N} \left\langle \widehat{x}^*, J(x_n^i - \widehat{x}^*) \right\rangle. \end{aligned}$$

By the accretiveness of A_i , we get

$$\left(\frac{\alpha_n}{N} + \gamma_n \right) \left\| x_n^i - \widehat{x}^* \right\|^2 \leq \gamma_n \|x_n - \widehat{x}^*\| \left\| x_n^i - \widehat{x}^* \right\| + \frac{\alpha_n}{N} \|\widehat{x}^*\| \left\| x_n^i - \widehat{x}^* \right\|,$$

hence

$$\left(\frac{\alpha_n}{N} + \gamma_n \right) \left\| x_n^i - \widehat{x}^* \right\| \leq \gamma_n \|x_n - \widehat{x}^*\| + \frac{\alpha_n}{N} \|\widehat{x}^*\|.$$

Using the inequalities $\|x_n - \widehat{x}^*\| \leq r$ and $r \geq \|\widehat{x}^*\|$, we have

$$\left(\frac{\alpha_n}{N} + \gamma_n \right) \left\| x_n^i - \widehat{x}^* \right\| \leq \gamma_n r + \frac{\alpha_n}{N} r \leq \left(\frac{\alpha_n}{N} + \gamma_n \right) r,$$

which gives $\|x_n^i - \widehat{x}^*\| \leq r$. By (2.4), one gets

$$\|x_{n+1} - \widehat{x}^*\| \leq \frac{1}{N} \sum_{i=1}^N \left\| x_n^i - \widehat{x}^* \right\| \leq r.$$

Therefore, $x_{n+1} \in B_r(\widehat{x}^*)$. Thus, $\{x_n\}$ is bounded. \square

Theorem 2.1. *Suppose conditions A1-A3 or B1-B2 are fulfilled. Let $\{\alpha_n\}$ and $\{\gamma_n\}$ be real sequences, such that*

- i) $\alpha_n \rightarrow 0, \gamma_n \rightarrow +\infty$ as $n \rightarrow +\infty$,
- ii) $\frac{\gamma_n |\alpha_{n+1} - \alpha_n|}{\alpha_n^2} \rightarrow 0$ as $n \rightarrow +\infty$, $\sum_{n=1}^{\infty} \frac{\alpha_n}{\gamma_n} = +\infty$,
- iii) $\frac{h_X(\tau_n) \varphi_R^{-1}(R_1 \alpha_n)}{\alpha_n} \rightarrow 0$ as $n \rightarrow +\infty$, where $R \geq 2\|\widehat{x}^*\|$, $R_1 := \frac{3R^2}{2}$ and $\tau_n = \gamma_n^{-1}$.

If in addition, the function $\frac{\varphi(s,t)}{t}$ is coercive in t for any fixed $s > 0$, i.e., $\frac{\varphi(s,t)}{t} \rightarrow +\infty$ as $t \rightarrow +\infty$, then starting from arbitrary $x_0 \in X$, the sequence $\{x_n\}$ defined by (2.3) and (2.4) converges strongly to \widehat{x}^* .

Proof. Let x_n^* be the unique solution of (2.2). Setting $e_n^i = x_n^i - x_n^*$; $e_n = x_n - x_n^*$; $\epsilon_n = \frac{\alpha_n \tau_n}{N}$, we can rewrite (2.3) as

$$x_n^i + \tau_n A_i(x_n^i) + \epsilon_n x_n^i = x_n,$$

or

$$\left(e_n^i - e_n \right) + \tau_n \left[A_i(x_n^i) - A_i(x_n^*) \right] + \epsilon_n e_n^i = -\tau_n A_i(x_n^*) - \epsilon_n x_n^*, \quad i = 1, 2, \dots, N.$$

From the last relation, using the accretiveness of A_i we find

$$2 \left\langle e_n^i - e_n, J(e_n^i) \right\rangle + 2\epsilon_n \left\| e_n^i \right\|^2 \leq -2 \left\langle \tau_n A_i(x_n^*) + \epsilon_n x_n^*, J(e_n^i) \right\rangle. \quad (2.5)$$

From Lemma 1.3, we get

$$2 \left\langle e_n^i - e_n, J(e_n^i) \right\rangle \geq \left\| e_n^i \right\|^2 - \|e_n\|^2.$$

Combining this inequality with (2.5), we obtain

$$(1 + 2\epsilon_n) \left\| e_n^i \right\|^2 - \|e_n\|^2 \leq -2\tau_n \left\langle A_i(x_n^*) + \frac{\alpha_n}{N} x_n^*, J(e_n^i) \right\rangle,$$

hence,

$$(1 + 2\epsilon_n) \sum_{i=1}^N \left\| e_n^i \right\|^2 - N \|e_n\|^2 \leq -2\tau_n \sum_{i=1}^N \left\langle A_i(x_n^*) + \frac{\alpha_n}{N} x_n^*, J(e_n^i) \right\rangle. \quad (2.6)$$

Observing that x_n^* is the solution of (2.2) and using Lemma 2.3, we can estimate the right-hand side of (2.6) as follows

$$\begin{aligned} - \sum_{i=1}^N \left\langle A_i(x_n^*) + \frac{\alpha_n}{N} x_n^*, J(e_n^i) \right\rangle &= - \left\langle \sum_{i=1}^N A_i(x_n^*) + \alpha_n x_n^*, J(e_n) \right\rangle \\ &\quad - \sum_{i=1}^N \left\langle A_i(x_n^*) + \frac{\alpha_n}{N} x_n^*, J(e_n^i) - J(e_n) \right\rangle \\ &\leq \sum_{i=1}^N \left(\|A_i(x_n^*)\| + \frac{\alpha_n}{N} \|x_n^*\| \right) \|J(e_n^i) - J(e_n)\| \\ &\leq \sum_{i=1}^N \left(\varphi_R^{-1} \left(6\alpha_n \|\hat{x}^*\|^2 \right) + \frac{2\alpha_n}{N} \|\hat{x}^*\| \right) \|J(e_n^i) - J(e_n)\|. \end{aligned} \quad (2.7)$$

By Lemma 2.3 and Lemma 2.4, the sequences $\{x_n^*\}$, $\{x_n\}$ and $\{x_n^i\}$ are bounded, hence the sequences $\{e_n\}$ and $\{e_n^i\}$ are also bounded, i.e., there exists a positive constant $C > 0$ such that $\|e_n\| \leq C$; $\|e_n^i\| \leq C$; $\|x_n^i\| \leq C$.

From Lemma 1.2, we get

$$\|J(e_n^i) - J(e_n)\| \leq 8Ch_X \left(\frac{16L \|x_n^i - x_n\|}{C} \right), \quad (2.8)$$

where $L \in (1, 1.7)$ is Figiel constant.

We show that $\|A_i(z)\| \leq C_i < +\infty$ for all $\|z\| \leq R_0 := C + 2\|\hat{x}^*\|$ and $i = 1, \dots, N$. Indeed, suppose in contrary, that there exists a sequence $\{z_n\}$, such that $\|z_n\| \leq R_0$, and $\|A_i(z_n)\| \rightarrow \infty$ as $n \rightarrow \infty$. Then $t_n := \|A_i(z_n) - A_i(0)\| \geq \|A_i(z_n)\| - \|A_i(0)\| \rightarrow \infty$ as $n \rightarrow \infty$. Since $\varphi(R_0, t_n) = \varphi(R_0, \|A_i(z_n) - A_i(0)\|) \leq \langle A_i(z_n) - A_i(0), J(z_n - 0) \rangle \leq \|A_i(z_n) - A_i(0)\| \|z_n\| \leq R_0 t_n$, we get $R_0 \geq \frac{\varphi(R_0, t_n)}{t_n}$, which contradicts the coerciveness of $\frac{\varphi(R_0, t)}{t}$.

Thus, we can put

$$M = \sup \left\{ \left\| A_i(x) + \frac{\alpha_n}{N} x \right\| : \|x\| \leq R_0, n = 1, 2, \dots, i = 1, 2, \dots, N \right\}.$$

Relation (2.3) yields

$$A_i(x_n^i) + \frac{\alpha_n}{N} x_n^i = \gamma_n (x_n - x_n^i), i = 1, 2, \dots, N,$$

which gives $\gamma_n \|x_n - x_n^i\| \leq \|A_i(x_n^i) + \frac{\alpha_n}{N} x_n^i\| \leq M$, hence, $\|x_n - x_n^i\| \leq \frac{M}{\gamma_n} = M\tau_n$.

Combining the last inequality with (2.8), we obtain

$$\|J(e_n^i) - J(e_n)\| \leq c_2 h_X(k_0 \tau_n), \quad (2.9)$$

where $c_2 = 8C, k_0 = \frac{16LM}{C}$. By (2.7) and (2.9), we have

$$-\sum_{i=1}^N \left\langle A_i(x_n^*) + \frac{\alpha_n}{N} x_n^*, J(e_n^i) \right\rangle \leq Nc_2 \left(\varphi_R^{-1} \left(6\alpha_n \|\hat{x}^*\|^2 \right) + \frac{2\alpha_n}{N} \|\hat{x}^*\| \right) h_X(k_0\tau_n). \quad (2.10)$$

From (2.6), (2.10), we get

$$(1 + 2\epsilon_n) \sum_{i=1}^N \|e_n^i\|^2 \leq N \|e_n\|^2 + 2Nc_2\tau_n \left(\varphi_R^{-1} \left(6\alpha_n \|\hat{x}^*\|^2 \right) + \frac{2\alpha_n}{N} \|\hat{x}^*\| \right) h_X(k_0\tau_n). \quad (2.11)$$

Taking into account relation (2.4), Lemma 2.3, and the inequality $(a + b)^2 \leq (1 + \epsilon_n)(a^2 + \frac{b^2}{\epsilon_n})$, we find

$$\begin{aligned} \|e_{n+1}\|^2 &= \|x_{n+1} - x_{n+1}^*\|^2 \leq (\|x_{n+1} - x_n^*\| + \|x_n^* - x_{n+1}^*\|)^2 \\ &\leq \left(\|x_{n+1} - x_n^*\| + 2\|\hat{x}^*\| \frac{|\alpha_{n+1} - \alpha_n|}{\alpha_n} \right)^2 \\ &\leq \left(\frac{1}{N} \sum_{i=1}^N \|e_n^i\| + 2\|\hat{x}^*\| \frac{|\alpha_{n+1} - \alpha_n|}{\alpha_n} \right)^2 \\ &\leq \left(\frac{1}{\sqrt{N}} \left(\sum_{i=1}^N \|e_n^i\|^2 \right)^{1/2} + 2\|\hat{x}^*\| \frac{|\alpha_{n+1} - \alpha_n|}{\alpha_n} \right)^2 \\ &\leq (1 + \epsilon_n) \left(\frac{1}{N} \sum_{i=1}^N \|e_n^i\|^2 + 4\|\hat{x}^*\|^2 \frac{|\alpha_{n+1} - \alpha_n|^2}{\alpha_n^2 \epsilon_n} \right). \end{aligned}$$

Thus,

$$\frac{N}{(1 + \epsilon_n)} \|e_{n+1}\|^2 - 4N\epsilon_n \left(\frac{\alpha_{n+1} - \alpha_n}{\epsilon_n \alpha_n} \right)^2 \|\hat{x}^*\|^2 \leq \sum_{i=1}^N \|e_n^i\|^2. \quad (2.12)$$

From (2.12), (2.11), one gets

$$\begin{aligned} \|e_{n+1}\|^2 &\leq \frac{1 + \epsilon_n}{1 + 2\epsilon_n} \|e_n\|^2 + 4(1 + \epsilon_n)\epsilon_n \left(\frac{\alpha_{n+1} - \alpha_n}{\epsilon_n \alpha_n} \right)^2 \|\hat{x}^*\|^2 \\ &\quad + \frac{2c_2(1 + \epsilon_n)\tau_n}{1 + 2\epsilon_n} \left(\varphi_R^{-1} \left(6\alpha_n \|\hat{x}^*\|^2 \right) + \frac{2\alpha_n}{N} \|\hat{x}^*\| \right) h_X(k_0\tau_n). \end{aligned} \quad (2.13)$$

Setting $\lambda_n = \|e_n\|^2$; $p_n = \frac{\epsilon_n}{1+2\epsilon_n}$ and $b_n = b_{1n} + b_{2n} + b_{3n}$, where

$$\begin{aligned} b_{1n} &= \frac{4(1 + \epsilon_n)}{\epsilon_n} \left(\frac{\alpha_{n+1} - \alpha_n}{\alpha_n} \right)^2 \|\hat{x}^*\|^2; \\ b_{2n} &= \frac{2c_2(1 + \epsilon_n)\tau_n}{1 + 2\epsilon_n} \varphi_R^{-1} \left(6\alpha_n \|\hat{x}^*\|^2 \right) h_X(k_0\tau_n); \\ b_{3n} &= \frac{4c_2(1 + \epsilon_n)\tau_n\alpha_n}{N(1 + 2\epsilon_n)} \|\hat{x}^*\| h_X(k_0\tau_n). \end{aligned}$$

We can rewrite (2.13) as $\lambda_{n+1} \leq (1 - p_n)\lambda_n + b_n$. Clearly, $\lambda_n, b_n \geq 0$; $p_n \in (0; 1)$ and $p_n \rightarrow 0$ as $n \rightarrow +\infty$.

Since $p_n = \frac{\epsilon_n}{1+2\epsilon_n}$ and $\epsilon_n \rightarrow 0$ as $n \rightarrow +\infty$, the series $\sum_{n=1}^{\infty} p_n = +\infty$ if and only if $\sum_{n=1}^{\infty} \epsilon_n = +\infty$. The last fact is equivalent to the assumption $\sum_{n=1}^{\infty} \frac{\alpha_n}{\gamma_n} = +\infty$.

By the assumption ii), $\frac{b_{1n}}{p_n} = 4(1 + \epsilon_n)(1 + 2\epsilon_n) \left(\frac{\alpha_{n+1} - \alpha_n}{\epsilon_n \alpha_n} \right)^2 \|\hat{x}^*\|^2 \rightarrow 0$ as $n \rightarrow +\infty$.

Further, using assumption iii), we will show that the expression

$$\frac{b_{2n}}{p_n} = 2c_2 N (1 + \epsilon_n) \frac{\varphi_R^{-1} \left(6\alpha_n \|\hat{x}^*\|^2 \right) h_X(k_0 \tau_n)}{\alpha_n} \quad (2.14)$$

will tend to zero as $n \rightarrow +\infty$. We first prove that there exist positive integers m and n_0 , such that for all $n \geq n_0$, $h_X(k_0 \tau_n) \leq \frac{5^m}{k_0} h_X(\tau_n)$. Indeed, according to [10, Lemma 1, page 65], we have

$$2 \leq \limsup_{\tau \rightarrow 0^+} \frac{\rho_X(2\tau)}{\rho_X(\tau)} \leq 4.$$

Hence, there exists $\tau_0 > 0$, such that $\frac{\rho_X(2\tau)}{\rho_X(\tau)} \leq 5$ for all $\tau \leq \tau_0$. Since $\tau_n \rightarrow 0$ as $n \rightarrow +\infty$, we can find a number n_0 such that $k_0 \tau_n \leq \tau_0$ for all $n \geq n_0$. Let m be a sufficiently large positive integer, such that $2^m \geq k_0$. Then for all $n \geq n_0$ we have $\rho_X(k_0 \tau_n) = \rho_X(2^{\frac{k_0 \tau_n}{2^m}}) \leq 5 \rho_X(2^{\frac{k_0 \tau_n}{2^{m-1}}}) \leq 5^2 \rho_X(2^{\frac{k_0 \tau_n}{2^{m-2}}}) \leq \dots \leq 5^m \rho_X(2^{\frac{k_0 \tau_n}{2^m}})$.

Because of the convexity of ρ_X and $\frac{k_0}{2^m} \leq 1$, we get $\rho_X(k_0 \tau_n) \leq 5^m \rho_X(\frac{k_0 \tau_n}{2^m}) \leq 5^m \rho_X(\tau_n)$. Thus, we come to the relation $h_X(k_0 \tau_n) = \frac{\rho_X(k_0 \tau_n)}{k_0 \tau_n} \leq \frac{5^m \rho_X(\tau_n)}{k_0 \tau_n} = \frac{5^m}{k_0} h_X(\tau_n)$.

Now using the last inequality and taking into account the fact that $\varphi_R^{-1}(t)$ is an increasing function and $R_1 := \frac{3}{2}R^2 \geq 6\|\hat{x}^*\|^2$, we can estimate the expression (2.14) as $\frac{b_{2n}}{p_n} \leq \frac{2c_2 N 5^m (1 + \epsilon_n) \varphi_R^{-1}(R_1 \alpha_n) h_X(\tau_n)}{\alpha_n}$ for all $n \geq n_0$. The assumption iii) implies that $\frac{b_{2n}}{p_n} \rightarrow 0$ as $n \rightarrow +\infty$.

Finally, the uniform smoothness of X gives

$$\frac{b_{3n}}{p_n} = 4c_2 (1 + \epsilon_n) \|\hat{x}^*\| h_X(k_0 \tau_n) \rightarrow 0$$

as $n \rightarrow +\infty$. Thus, $\frac{b_n}{p_n} \rightarrow 0 (n \rightarrow +\infty)$. Lemma 1.6 ensures that $\lambda_n = \|e_n\|^2 = \|x_n - x_n^*\|^2 \rightarrow 0 (n \rightarrow +\infty)$. Besides, by Lemma 2.3, $x_n^* \rightarrow \hat{x}^* (n \rightarrow +\infty)$, hence $\|x_n - \hat{x}^*\| \leq \|x_n - x_n^*\| + \|x_n^* - \hat{x}^*\| \rightarrow 0 (n \rightarrow +\infty)$. The proof of Theorem 2.1 is complete. \square

Example 4. Let $A_i, i = 1, \dots, N$ be c -inverse strongly monotone operators on a real Hilbert space X . Then all the conditions A1-A3 and B1-B2 are satisfied. Further, since $\varphi(s, t) = ct^2$, the function $\frac{\varphi(s, t)}{t} = ct$ is coercive. Conditions i), ii) on the parameters α_n, γ_n have been already stated in [5, Theorem 2.1]. On the other hand, for a Hilbert space, $\rho_X(t) = \sqrt{1 + t^2} - 1 \leq \frac{t^2}{2}$, hence the assumption iii) of Theorem 2.1 leads to the additional constraint $\gamma_n \alpha_n^{1/2} \rightarrow +\infty (n \rightarrow +\infty)$.

An example of such a pair of parameters could be $\alpha_n = (n + 1)^{-p}$, where, $0 < p < 1/2$ and $\gamma_n = (n + 1)^{1/2}$.

In the next two examples we suppose that $X = l^p, 1 \leq p < +\infty$ and $A_i = I - T_i$, where $T_i : X \rightarrow X, i = 1, \dots, N$, are nonexpansive operators. In this case both sets of conditions A1-A3 and B1-B2 are fulfilled. Observe that for proving the m -accretiveness of A_i one should use the identity $A_i + \alpha I = (1 + \alpha)\{I - (1 + \alpha)^{-1}T_i\}$ and the fact that $(1 + \alpha)^{-1}T_i$ is a contraction for $i = 1, \dots, N$.

Example 5. Let $X = l^p$ with $p \geq 2$, then $\rho_X(t) \leq (p - 1)t^2$ and $h_X(t) \leq (p - 1)t$ (see [9], page 48). According to Example 3, all the operators $A_i := I - T_i, i = 1, 2, \dots, N$,

are inverse uniformly accretive with $\varphi(s, t) = \frac{1}{Lp8^p} \frac{t^p}{s^{p-2}}$. For any fixed $s > 0$, $\varphi_s^{-1}(t) = c(s)t^{\frac{1}{p}}$, where $c(s)$ is a positive constant, and the function $\frac{\varphi(s, t)}{t}$ is coercive in t . The assumption iii) of Theorem 2.1 becomes $\gamma_n \alpha_n^{\frac{p-1}{p}} \rightarrow \infty$ ($n \rightarrow +\infty$) and we can choose $\alpha_n = (n+1)^{-k}$, $\gamma_n = (n+1)^{1/2}$ with $0 < k < \frac{1}{2}$.

Example 6. Suppose $X = l^p$, $1 < p < 2$, then we have (see [9], page 48) $\rho_X(t) \leq \frac{t^p}{p}$, $h_X(t) \leq \frac{t^{p-1}}{p}$. Example 3 shows that $A_i, i = 1, \dots, N$ are c -inverse strongly accretive operators with $\varphi(s, t) = ct^2$, $\varphi_s^{-1}(t) = \frac{\sqrt{t}}{\sqrt{c}}$, $c = \frac{p-1}{256L}$ and the assumption iii) of Theorem 2.1 becomes $\alpha_n^{1/2} \gamma_n^{p-1} \rightarrow \infty$ ($n \rightarrow +\infty$). We can choose $\alpha_n = (n+1)^{-k}$, $\gamma_n = (n+1)^{1/2}$ with $0 < k < \min\{\frac{1}{2}, p-1\}$.

Next we turn to the noisy data case. Assume that $A_i(x) := F_i(x) - f_i$ and the exact operators $F_i(x), i = 1, \dots, N$, are inverse uniformly accretive. Suppose that instead of the exact data (F_i, f_i) , we have only noisy ones $(F_{n,i}, f_{n,i})$, where the perturbed operators $F_{n,i} : D(F_{n,i}) = X \rightarrow X$ are just accretive for all $n \geq 1$ and $i = 1, \dots, N$. Moreover, let

$$\|F_{n,i}(x) - F_i(x)\| \leq h_n g(\|x\|), \quad (2.15)$$

$$\|f_{n,i} - f_i\| \leq \delta_n, \quad i = 1, \dots, N, \quad (2.16)$$

where, $g(t)$ is a nonnegative continuous nondecreasing function, $h_n > 0, \delta_n > 0$ for all $n > 0$. Starting from arbitrary $z_0 \in X$, we perform the following implicit PIRM:

$$A_{n,i}(z_n^i) + \left(\frac{\alpha_n}{N} + \gamma_n\right) z_n^i = \gamma_n z_n, \quad i = 1, 2, \dots, N, \quad (2.17)$$

$$z_{n+1} = \frac{1}{N} \sum_{i=1}^N z_n^i, \quad n = 0, 1, \dots, \quad (2.18)$$

where $A_{n,i}(x) := F_{n,i}(x) - f_{n,i}, i = 1, 2, \dots, N$.

Theorem 2.2. Assume that all the conditions of Theorem 2.1 and relations (2.15), (2.16) are fulfilled. If in addition $\frac{h_n + \delta_n}{\alpha_n} \rightarrow 0$ as $n \rightarrow +\infty$, then the sequence $\{z_n\}$ generated by (2.17), (2.18) converges strongly to \hat{x}^* as $n \rightarrow +\infty$.

Proof. Let $\{x_n\}$ be the sequence of approximations defined by (2.3), (2.4). From (2.3) and (2.17), we have

$$A_{n,i}(z_n^i) - A_i(x_n^i) + \left(\frac{\alpha_n}{N} + \gamma_n\right) (z_n^i - x_n^i) = \gamma_n (z_n - x_n),$$

or

$$\begin{aligned} \left(F_{n,i}(z_n^i) - F_{n,i}(x_n^i)\right) + \left(F_{n,i}(x_n^i) - F_i(x_n^i)\right) + (f_{n,i} - f_i) + \left(\frac{\alpha_n}{N} + \gamma_n\right) (z_n^i - x_n^i) \\ = \gamma_n (z_n - x_n). \end{aligned}$$

Therefore

$$\begin{aligned} \langle F_{n,i}(z_n^i) - F_{n,i}(x_n^i), J(z_n^i - x_n^i) \rangle + \langle F_{n,i}(x_n^i) - F_i(x_n^i), J(z_n^i - x_n^i) \rangle \\ + \langle f_{n,i} - f_i, J(z_n^i - x_n^i) \rangle + \left(\frac{\alpha_n}{N} + \gamma_n\right) \|z_n^i - x_n^i\|^2 \\ = \gamma_n \langle z_n - x_n, J(z_n^i - x_n^i) \rangle \end{aligned} \quad (2.19)$$

By relations (2.15), (2.16) and (2.19), we get

$$\left(\frac{\alpha_n}{N} + \gamma_n\right) \|z_n^i - x_n^i\|^2 \leq h_n g(\|x_n^i\|) \|z_n^i - x_n^i\| + \delta_n \|z_n^i - x_n^i\| + \gamma_n \|z_n - x_n\| \|z_n^i - x_n^i\|$$

Lemma 2.4 ensures the boundedness of the sequence $\{x_n^i\}$. Thus, $\|x_n^i\| \leq R$ for some $R > 0$. Setting $\lambda_n = \|z_n - x_n\|$, from the last inequality we find

$$\|z_n^i - x_n^i\| \leq \frac{N\gamma_n}{\alpha_n + N\gamma_n} \lambda_n + \frac{Ng(R)h_n}{\alpha_n + N\gamma_n} + \frac{N\delta_n}{\alpha_n + N\gamma_n} \quad (2.20)$$

On account of (2.4), (2.18) and (2.20), we have

$$\lambda_{n+1} = \|z_{n+1} - x_{n+1}\| \leq \frac{1}{N} \sum_{i=1}^N \|z_n^i - x_n^i\| \leq \frac{N\gamma_n}{\alpha_n + N\gamma_n} \lambda_n + \frac{Ng(R)h_n}{\alpha_n + N\gamma_n} + \frac{N\delta_n}{\alpha_n + N\gamma_n}. \quad (2.21)$$

Putting $p_n = \frac{\alpha_n}{\alpha_n + N\gamma_n}$, $b_n = \frac{Ng(R)h_n}{\alpha_n + N\gamma_n} + \frac{N\delta_n}{\alpha_n + N\gamma_n}$, from (2.21) we get $\lambda_{n+1} = (1 - p_n)\lambda_n + b_n$. By virtue of the hypotheses of Theorem 2.2, we have $\sum_{n=1}^{\infty} p_n = +\infty$, $\frac{b_n}{p_n} \rightarrow 0$ as $n \rightarrow +\infty$. Lemma 1.6 implies that $\lambda_n = \|z_n - x_n\| \rightarrow 0$ as $n \rightarrow +\infty$. Finally, by Theorem 2.1, $x_n \rightarrow \hat{x}^*$ ($n \rightarrow +\infty$), hence $\|z_n - \hat{x}^*\| \leq \|z_n - x_n\| + \|x_n - \hat{x}^*\| \rightarrow 0$ ($n \rightarrow +\infty$). The proof of Theorem 2.2 is complete. \square

We now consider an explicit PIRM for solving system (1.1), consisting of synchronous computation of intermediate approximations z_{ni}

$$z_{ni} = z_n - \frac{1}{\gamma_n} \left\{ A_i(z_n) + \frac{\alpha_n}{N} z_n \right\} = z_n - \tau_n \left\{ A_i(z_n) + \frac{\alpha_n}{N} z_n \right\}, i = 1, 2, \dots, N, \quad (2.22)$$

and defining the next approximation z_{n+1} as an average of intermediate approximations z_{ni}

$$z_{n+1} = \frac{1}{N} \sum_{i=1}^N z_{ni}, n = 1, 2, \dots \quad (2.23)$$

Lemma 2.5. *Suppose conditions B1-B2 are satisfied. Assume in addition the function $\frac{\varphi(s,t)}{t}$ is coercive in t for every fixed $s > 0$. Let $\{\alpha_n\}$ and $\{\gamma_n\}$ be positive sequences such that for all $n \geq 0$, $\alpha_n \leq 1$, $\gamma_n \geq 1$, and*

$$\tau_n \leq d, \quad \frac{\rho_X(\tau_n)}{\tau_n \alpha_n} \leq d^2, \quad (2.24)$$

where $\tau_n := 1/\gamma_n$ and $d \in (0, 1)$ is a fixed number. Then starting from arbitrary $z_0 \in X$, the sequence $\{z_n\}$ generalized by (2.22), (2.23) is bounded.

Proof. A simple verification shows that the sequence $\{z_n\}$ defined by (2.22) and (2.23) satisfies the relation

$$z_{n+1} = z_n - \frac{1}{N\gamma_n} \{A(z_n) + \alpha_n z_n\}, \quad (2.25)$$

where $A(z) := \sum_{i=1}^N A_i(z)$. By our assumptions, all the operators $A_i, i = 1, \dots, N$, are continuous, m -accretive and φ -inverse uniformly accretive. Moreover, as it was shown in the proof of Theorem 2.1, A_i is bounded for every i , hence the operator $A : D(A) = X \rightarrow X$ is bounded, continuous and m -accretive. According to Lemma 2.1, equation $A(z) = 0$ is equivalent to the consistent system (1.1). By [21, Theorem 5.1] the sequence $\{z_n\}$ defined by (2.25) is bounded. \square

Theorem 2.3. Assume that all the conditions of Lemma 2.5 are fulfilled. In addition, let $\alpha_n \rightarrow 0, \tau_n := 1/\gamma_n \rightarrow 0$ as $n \rightarrow +\infty$, such that

$$\sum_{i=1}^{\infty} \alpha_n \tau_n = +\infty, \quad \frac{\tau_n}{\alpha_n} \rightarrow 0, \quad \frac{|\alpha_n - \alpha_{n+1}|}{\tau_n \alpha_n^2} \rightarrow 0, \quad \frac{\rho_X(\tau_n)}{\tau_n \alpha_n} \rightarrow 0. \quad (2.26)$$

Then the sequence $\{z_n\}$ generalized by (2.22) and (2.23) converges to \hat{x}^* as $n \rightarrow \infty$.

Proof. Let x_n^* be the unique solution of regularized equation (2.2). It follows from Lemma 2.3 that $\{x_n^*\}$ is bounded, hence there exists a constant $\tilde{d} > 0$ such that

$$\|x_n^* - x_{n+1}^*\| \leq \tilde{d}.$$

By Lemma 1.3, we have

$$\begin{aligned} \|z_{ni} - x_{n+1}^*\|^2 &\leq \|z_{ni} - x_n^*\|^2 + 2 \langle x_{n+1}^* - x_n^*, J(x_n^* - z_{ni}) \rangle \\ &\quad + 2 \langle x_{n+1}^* - x_n^*, J(x_{n+1}^* - z_{ni}) - J(x_n^* - z_{ni}) \rangle. \end{aligned} \quad (2.27)$$

Further, by Lemma 1.4, we get

$$\begin{aligned} \|z_{ni} - x_{n+1}^*\|^2 &\leq \|z_{ni} - x_n^*\|^2 + 2 \|z_{ni} - x_n^*\| \cdot \|x_{n+1}^* - x_n^*\| \\ &\quad + 16 \|x_{n+1}^* - x_n^*\|^2 + c_1(n) \rho_X(\|x_{n+1}^* - x_n^*\|), \end{aligned} \quad (2.28)$$

where $c_1(n) = 8 \max\{2L, \|z_{ni} - x_{n+1}^*\| + \|z_{ni} - x_n^*\|\}$. Taking into account Lemma 2.5 and the boundedness of the operators A_i we conclude that the sequence $\{z_{ni}\}$ is also bounded, therefore there exist positive numbers c_1, k_0 such that $c_1(n) \leq c_1$ and $\|z_{ni} - x_n^*\| \leq k_0$ for all $n \geq 0$. Note that, if H is a Hilbert space, then for all $0 < \tau < \hat{\tau}$

$$\rho_X(\tau) \geq \rho_H(\tau) = \sqrt{1 + \tau^2} - 1 \geq \hat{c}\tau^2, \quad (2.29)$$

where $\hat{c} = (\sqrt{1 + \hat{\tau}^2} + 1)^{-1}$.

Now, summing up both sides of (2.28) for $i = 1, 2, \dots, N$, and using Lemma 1.5, as well as inequality (2.29) with $\hat{\tau} := \tilde{d} \geq \tau := \|x_{n+1}^* - x_n^*\|$, we obtain

$$\begin{aligned} \sum_{i=1}^N \|z_{ni} - x_{n+1}^*\|^2 &\leq \sum_{i=1}^N \|z_{ni} - x_n^*\|^2 + 4Nk_0 \frac{|\alpha_{n+1} - \alpha_n|}{\alpha_n} \|\hat{x}^*\| \\ &\quad + N \left(16\hat{c}^{-1} + c_1\right) \rho_X \left(2 \frac{|\alpha_{n+1} - \alpha_n|}{\alpha_n} \|\hat{x}^*\|\right). \end{aligned}$$

From the last inequality and the fact that $\rho_X(\tau) \leq \tau$, one gets

$$\sum_{i=1}^N \|z_{ni} - x_{n+1}^*\|^2 \leq \sum_{i=1}^N \|z_{ni} - x_n^*\|^2 + c_3 \frac{|\alpha_{n+1} - \alpha_n|}{\alpha_n}, \quad (2.30)$$

where $c_3 = 2N \|\hat{x}^*\| (2k_0 + 16\hat{c}^{-1} + c_1)$.

Next, we shall estimate the expression $\sum_{i=1}^N \|z_{ni} - x_n^*\|^2$. By Lemma 1.3 and (2.22), we have

$$\begin{aligned} \|z_{ni} - x_n^*\|^2 &= \left\| z_n - x_n^* - \tau_n \left\{ A_i(z_n) + \frac{\alpha_n}{N} z_n \right\} \right\|^2 \\ &\leq \|z_n - x_n^*\|^2 - 2\tau_n \left\langle A_i(z_n) + \frac{\alpha_n}{N} z_n, J(z_n - x_n^*) \right\rangle \\ &\quad + 2 \langle z_{ni} - z_n, J(z_{ni} - x_n^*) - J(z_n - x_n^*) \rangle. \end{aligned} \quad (2.31)$$

Besides,

$$\|z_{ni} - z_n\| = \tau_n \left\| A_i(z_n) + \frac{\alpha_n}{N} z_n \right\| \leq M\tau_n, \quad (2.32)$$

where $M = \sup \{ \|A_i(z_n) + \frac{\alpha_n}{N} z_n\| : i = 1, 2, \dots, N; n = 1, 2, \dots \}$. Using (2.32) and Lemma 1.4, one obtains

$$\langle z_{ni} - z_n, J(z_{ni} - x_n^*) - J(z_n - x_n^*) \rangle \leq 8M^2\tau_n^2 + c_2(n)\rho_X(M\tau_n),$$

where $c_2(n) = 4\max \{ 2L, \|z_{ni} - x_n^*\| + \|z_n - x_n^*\| \} \leq c_2$ because of the boundedness of the sequences $\{z_{ni}\}$, $\{z_n\}$ and $\{x_n^*\}$. Therefore

$$\langle z_{ni} - z_n, J(z_{ni} - x_n^*) - J(z_n - x_n^*) \rangle \leq 8M^2\tau_n^2 + c_2\rho_X(M\tau_n). \quad (2.33)$$

On the other hand, since the operators A_i are accretive and x_n^* is the solution of (2.2), we have

$$\begin{aligned} \sum_{i=1}^N \langle A_i(z_n) + \frac{\alpha_n}{N} z_n, J(z_n - x_n^*) \rangle &= \sum_{i=1}^N \langle A_i(z_n) - A_i(x_n^*), J(z_n - x_n^*) \rangle \\ &\quad + \left\langle \sum_{i=1}^N A_i(x_n^*) + \alpha_n x_n^*, J(z_n - x_n^*) \right\rangle + \alpha_n \|z_n - x_n^*\|^2 \\ &\geq \alpha_n \|z_n - x_n^*\|^2. \end{aligned} \quad (2.34)$$

Now, summing the both sides of (2.31) for $i = 1, 2, \dots, N$ and taking account relations (2.33), (2.34), we get

$$\sum_{i=1}^N \|z_{ni} - x_n^*\|^2 \leq N \|z_n - x_n^*\|^2 - 2\tau_n\alpha_n \|z_n - x_n^*\|^2 + 16NM^2\tau_n^2 + 2Nc_2\rho_X(M\tau_n). \quad (2.35)$$

Note that

$$\|z_{n+1} - x_{n+1}^*\|^2 \leq \frac{1}{N^2} \left(\sum_{i=1}^N \|z_{ni} - x_{n+1}^*\| \right)^2 \leq \frac{1}{N} \sum_{i=1}^N \|z_{ni} - x_{n+1}^*\|^2. \quad (2.36)$$

From (2.30), (2.35), (2.36), we get

$$\begin{aligned} \|z_{n+1} - x_{n+1}^*\|^2 &\leq \|z_n - x_n^*\|^2 - \frac{2\tau_n\alpha_n}{N} \|z_n - x_n^*\|^2 \\ &\quad + 16M^2\tau_n^2 + 2c_2\rho_X(M\tau_n) + c_3 \frac{|\alpha_{n+1} - \alpha_n|}{\alpha_n}. \end{aligned} \quad (2.37)$$

Setting $\lambda_n = \|z_n - x_n^*\|^2$, $p_n = \frac{2\tau_n\alpha_n}{N}$, $b_n = 16M^2\tau_n^2 + 2c_2\rho_X(M\tau_n) + c_3 \frac{|\alpha_{n+1} - \alpha_n|}{\alpha_n}$, we can rewrite (2.37) as

$$\lambda_{n+1} \leq (1 - p_n)\lambda_n + b_n.$$

In the same manner as in the proof of Theorem 2.1, we can find positive integers n_0 and m , such that for all $n \geq n_0$, $\rho_X(M\tau_n) \leq 5^m\rho_X(\tau_n)$. By Lemma 1.6 and the hypothesis (2.26), we conclude that $\lambda_n = \|z_n - x_n^*\|^2 \rightarrow 0$ as $n \rightarrow +\infty$. Finally, by Lemma 2.3,

$$\|z_n - \hat{x}^*\| \leq \|z_n - x_n^*\| + \|x_n^* - \hat{x}^*\| \rightarrow 0,$$

which implies that $\{z_n\}$ converges to \hat{x}^* . The proof of Theorem 2.3 is complete. \square

3. EQUATIONS WITH SMOOTH ACCRETIVE OPERATORS

In this section, for solving system (1.1) with smooth accretive operators A_i , we consider a parallel regularized Newton-type method (cf. [6, 7])

$$A_i(x_n) + \frac{\alpha_n}{N}(x_n - x_i^0) + (A_i'(x_n) + \frac{\alpha_n}{N}I)(x_n^i - x_n) = 0, i = 1, 2, \dots, N, \quad (3.1)$$

$$x_{n+1} = \frac{1}{N} \sum_{i=1}^N x_n^i, n = 0, 1, \dots \quad (3.2)$$

The following assumptions will be needed throughout Section 3.

C1. System (1.1) possesses an exact solution \hat{x}^* . The operators A_i ($i = 1, \dots, N$) are accretive on a real Banach space X and Fréchet differentiable in a closed ball $B_r(\hat{x}^*) \subset X$ with center \hat{x}^* and radius $r > 0$. Moreover,

$$\|A_i'(x) - A_i'(y)\| \leq K \|x - y\| \quad \forall x, y \in B_r(\hat{x}^*), i = 1, \dots, N.$$

C2. The following componentwise source condition (see [3, 7]) holds

$$x_i^0 - \hat{x}^* = A_i'(\hat{x}^*)v_i,$$

where $x_i^0 \in B_r(\hat{x}^*)$, $v_i \in X$, $1 \leq i \leq N$.

C3. The parameters α_n are chosen such that

$$\alpha_n > 0, \alpha_n \rightarrow 0, 1 \leq \frac{\alpha_n}{\alpha_{n+1}} \leq \rho,$$

where the constant $\rho > 1$.

Theorem 3.1. *Let all the assumptions C1 – C3 be satisfied. If $\sum_{i=1}^N \|v_i\|$ is sufficiently small and x_0 is close enough to \hat{x}^* , then there holds the estimate*

$$\|x_n - \hat{x}^*\| = O(\alpha_n). \quad (3.3)$$

Proof. We suppose by induction that $x_n \in B_r(\hat{x}^*)$ for some $n \geq 0$. Setting $e_n^i = x_n^i - \hat{x}^*$ and $e_n = x_n - \hat{x}^*$, from equation (3.1) and assumption C2, we get

$$\begin{aligned} e_n^i &= e_n + x_n^i - x_n = e_n - \left(A_i'(x_n) + \frac{\alpha_n}{N}I \right)^{-1} \left(A_i(x_n) + \frac{\alpha_n}{N}(x_n - x_i^0) \right) \\ &= \left(A_i'(x_n) + \frac{\alpha_n}{N}I \right)^{-1} \left[\left(A_i'(x_n) + \frac{\alpha_n}{N}I \right) e_n - \left(A_i(x_n) + \frac{\alpha_n}{N}(x_n - x_i^0) \right) \right] \\ &= \left(A_i'(x_n) + \frac{\alpha_n}{N}I \right)^{-1} \left[\frac{\alpha_n}{N}(x_i^0 - \hat{x}^*) + A_i'(x_n)e_n - A_i(x_n) \right] \\ &= \frac{\alpha_n}{N} \left(A_i'(x_n) + \frac{\alpha_n}{N}I \right)^{-1} A_i'(\hat{x}^*)v_i + \left(A_i'(x_n) + \frac{\alpha_n}{N}I \right)^{-1} (A_i'(x_n)e_n - A_i(x_n)). \end{aligned}$$

Using Lemma 1.7, from the last inequality we obtain

$$\|e_n^i\| \leq \frac{\alpha_n}{N} \left\| \left(A_i'(x_n) + \frac{\alpha_n}{N}I \right)^{-1} A_i'(\hat{x}^*) \right\| \|v_i\| + \frac{N}{\alpha_n} \|A_i'(x_n)e_n - A_i(x_n)\|. \quad (3.4)$$

Obviously, if $x_t := x_n + t(\hat{x}^* - x_n)$, where $0 \leq t \leq 1$, then

$$\|x_t - \hat{x}^*\| = (1 - t) \|x_n - \hat{x}^*\| \leq r,$$

hence $x_t \in B_r(\hat{x}^*)$.

The second term of the right-hand side of (3.4) can be estimated as

$$\begin{aligned} \frac{N}{\alpha_n} \|A'_i(x_n)e_n - A_i(x_n)\| &= \frac{N}{\alpha_n} \|A_i(\hat{x}^*) - A_i(x_n) + A'_i(x_n)(e_n)\| \\ &= \frac{N}{\alpha_n} \left\| \int_0^1 (A'_i(x_n) - A'_i(x_t)) e_n dt \right\| \\ &\leq \frac{N}{\alpha_n} \int_0^1 Kt \|e_n\|^2 dt = \frac{KN}{2\alpha_n} \|e_n\|^2. \end{aligned} \quad (3.5)$$

On the other hand, we have

$$\begin{aligned} \left(A'_i(x_n) + \frac{\alpha_n}{N}I\right)^{-1} - \left(A'_i(\hat{x}^*) + \frac{\alpha_n}{N}I\right)^{-1} &= \\ &= \left(A'_i(x_n) + \frac{\alpha_n}{N}I\right)^{-1} [A'_i(\hat{x}^*) - A'_i(x_n)] \left(A'_i(\hat{x}^*) + \frac{\alpha_n}{N}I\right)^{-1}. \end{aligned}$$

Therefore,

$$\begin{aligned} \left\| \left(A'_i(x_n) + \frac{\alpha_n}{N}I\right)^{-1} A'_i(\hat{x}^*) \right\| &\leq \left\| \left(A'_i(\hat{x}^*) + \frac{\alpha_n}{N}I\right)^{-1} A'_i(\hat{x}^*) \right\| \\ &+ \left\| \left(A'_i(x_n) + \frac{\alpha_n}{N}I\right)^{-1} [A'_i(\hat{x}^*) - A'_i(x_n)] \left(A'_i(\hat{x}^*) + \frac{\alpha_n}{N}I\right)^{-1} A'_i(\hat{x}^*) \right\|. \end{aligned}$$

From the last inequality and using Lemma 1.7 as well as assumption C1, we get

$$\left\| \left(A'_i(x_n) + \frac{\alpha_n}{N}I\right)^{-1} A'_i(\hat{x}^*) \right\| \leq 2 + \frac{2KN}{\alpha_n} \|e_n\|. \quad (3.6)$$

Combining (3.4), (3.5) and (3.6), one has

$$\|e_n^i\| \leq \frac{2\alpha_n \|v_i\|}{N} + 2K \|v_i\| \|e_n\| + \frac{NK}{2\alpha_n} \|e_n\|^2. \quad (3.7)$$

By (3.2) and (3.7), we obtain

$$\begin{aligned} \|e_{n+1}\| = \|x_{n+1} - \hat{x}^*\| &\leq \frac{1}{N} \sum_{i=1}^N \|e_n^i\| \leq \frac{2\alpha_n \sum_{i=1}^N \|v_i\|}{N^2} \\ &+ \frac{2K \sum_{i=1}^N \|v_i\|}{N} \|e_n\| + \frac{NK}{2\alpha_n} \|e_n\|^2. \end{aligned} \quad (3.8)$$

Setting $\omega_n = \frac{N\|e_n\|}{\alpha_n}$ and using assumption C3, from (3.8) we find

$$\begin{aligned} \omega_{n+1} &\leq \frac{2 \sum_{i=1}^N \|v_i\|}{N} \left(\frac{\alpha_n}{\alpha_{n+1}}\right) + \frac{2K \sum_{i=1}^N \|v_i\|}{N} \left(\frac{\alpha_n}{\alpha_{n+1}}\right) \omega_n + \frac{K}{2} \left(\frac{\alpha_n}{\alpha_{n+1}}\right) \omega_n^2 \\ &\leq \frac{2\rho \sum_{i=1}^N \|v_i\|}{N} + \frac{2\rho K \sum_{i=1}^N \|v_i\|}{N} \omega_n + \frac{K\rho}{2} \omega_n^2 \\ &= a + b\omega_n + c\omega_n^2, \end{aligned} \quad (3.9)$$

where

$$a = \frac{2\rho \sum_{i=1}^N \|v_i\|}{N}, \quad b = \frac{2\rho K \sum_{i=1}^N \|v_i\|}{N}, \quad c = \frac{K\rho}{2}.$$

If $\sum_{i=1}^N \|v_i\|$ is small enough, then a, b will be small, hence

$$b + 2\sqrt{ac} < 1, \frac{2a\alpha_0}{N} \leq r \left(1 - b + \sqrt{(1-b)^2 - 4ac} \right). \quad (3.10)$$

Now if x_0 is sufficiently close to \hat{x}^* then

$$\omega_0 = \frac{N \|e_0\|}{\alpha_0} = \frac{N \|x_0 - \hat{x}^*\|}{\alpha_0} \leq M_+ := \frac{(1-b + \sqrt{(1-b)^2 - 4ac})}{2c}.$$

Lemma 1.8 applied to (3.9) ensures that

$$\omega_n := \frac{N \|e_n\|}{\alpha_n} \leq l := \max \{ \omega_0, M_- \}, \forall n \geq 0,$$

$$\text{where } M_- = \frac{(1-b - \sqrt{(1-b)^2 - 4ac})}{2c} = \frac{2a}{(1-b + \sqrt{(1-b)^2 - 4ac})/2c}.$$

In particular,

$$\|x_{n+1} - \hat{x}^*\| = \|e_{n+1}\| = \frac{\omega_{n+1}\alpha_{n+1}}{N} \leq \frac{l\alpha_0}{N}. \quad (3.11)$$

Observing that $\frac{\omega_0\alpha_0}{N} = \|x_0 - \hat{x}^*\| \leq r$. From (3.10), we have

$$\frac{M_- \alpha_0}{N} = \frac{2a\alpha_0}{N \left(1 - b + \sqrt{(1-b)^2 - 4ac} \right)} \leq r.$$

Therefore, $\frac{l\alpha_0}{N} \leq r$, hence $x_{n+1} \in B_r(\hat{x}^*)$. Thus, the estimate $\omega_n \leq l$ yields

$$\|e_n\| = \frac{\omega_n \alpha_n}{N} \leq \frac{l\alpha_n}{N} = O(\alpha_n).$$

The proof of Theorem 3.1 is complete. \square

Now, assume that $A_i(x) = F_i(x) - f_i$ and instead of (F_i, f_i) , we only have approximations (F_i^h, f_i^δ) , such that

$$\|f_i^\delta - f_i\| \leq \delta, \quad (3.12)$$

and the operators F_i^h , $(i = 1, \dots, N)$, are accretive on a real Banach space X and Fréchet differentiable in a closed ball $B_r(\hat{x}^*)$. Moreover, suppose that

$$\|F_i^{h'}(x) - F_i^{h'}(y)\| \leq K \|x - y\| \quad \forall x, y \in B_r(\hat{x}^*), i = 1, 2, \dots, N. \quad (3.13)$$

Also, assume that

$$\|F_i^h(\hat{x}^*) - F_i(\hat{x}^*)\| \leq h, \quad (3.14)$$

$$\|F_i^{h'}(\hat{x}^*) - F_i'(\hat{x}^*)\| \leq h. \quad (3.15)$$

Given a starting point $x_0 \in B_r(\hat{x}^*)$, we define a sequence $\{x_n\}$:

$$\tilde{A}_i(x_n) + \frac{\alpha_n}{N}(x_n - x_i^0) + (\tilde{A}_i'(x_n) + \frac{\alpha_n}{N}I)(x_n^i - x_n) = 0, i = 1, 2, \dots, N \quad (3.16)$$

$$x_{n+1} = \frac{1}{N} \sum_{i=1}^N x_n^i, n = 1, 2, \dots, \quad (3.17)$$

where $\tilde{A}_i(x) = F_i^h(x) - f_i^\delta$. We define the stopping index $N(\delta, h)$ as

$$N(\delta, h) = \max \left\{ n : \alpha_n^2 \geq \frac{\delta + h}{\eta} \right\}, \quad (3.18)$$

where η is a fixed parameter.

By the same argument as in the proof of Theorem 3.1, we come to the following estimate for iteration (3.16) and (3.17)

$$\|e_n^i\| \leq \frac{2\alpha_n \|v_i\|}{N} + 2 \|v_i\| \|F_i'(\hat{x}^*) - F_i'(x_n)\| + \frac{N}{\alpha_n} \|F_i^{h'}(x_n)e_n - F_i^h(x_n) + f^\delta\| \quad (3.19)$$

From conditions (3.13) and (3.15), we have

$$\|F_i'(\hat{x}^*) - F_i'(x_n)\| \leq \|F_i'(\hat{x}^*) - F_i'(\hat{x}^*)\| + \|F_i^{h'}(\hat{x}^*) - F_i'(x_n)\| \leq h + K \|e_n\|. \quad (3.20)$$

And using conditions (3.12) and (3.14), we have

$$\begin{aligned} \|F_i^{h'}(x_n)e_n - F_i^h(x_n) + f^\delta\| &\leq \|F_i^{h'}(x_n)e_n - F_i^h(x_n) + F_i^h(\hat{x}^*)\| + \\ &\quad + \|F_i^h(\hat{x}^*) - F_i^h(x_n)\| + \|f - f^\delta\| \leq \frac{K}{2} \|e_n\|^2 + h + \delta. \end{aligned} \quad (3.21)$$

Combining (3.19), (3.20), (3.21) and noting that $\|e_{n+1}\| \leq \frac{1}{N} \sum_{i=1}^N \|e_n^i\|$, we obtain

$$\|e_{n+1}\| \leq \left[\frac{2 \sum_{i=1}^N \|v_i\|}{N} (h + \alpha_n) + \frac{h + \delta}{\alpha_n} \right] + \frac{2L \sum_{i=1}^N \|v_i\|}{N} \|e_n\| + \frac{K}{2\alpha_n} \|e_n\|^2. \quad (3.22)$$

From (3.18) we get $\frac{h}{\alpha_n} \leq \alpha_n \eta \leq \alpha_0 \eta$. Setting $\omega_n = \frac{N \|e_n\|}{\alpha_n}$ we can rewrite (3.22) as

$$\begin{aligned} \omega_{n+1} &\leq \frac{\alpha_n}{\alpha_{n+1}} \left[\frac{2 \sum_{i=1}^N \|v_i\|}{N} \left(\frac{h}{\alpha_n} + 1 \right) + \frac{h + \delta}{\alpha_n^2} + \frac{2L \sum_{i=1}^N \|v_i\|}{N} \omega_n + \frac{K}{2} \omega_n^2 \right] \\ &\leq \rho \left[\left(\frac{2 \sum_{i=1}^N \|v_i\|}{N} (\alpha_0 \eta + 1) + \eta \right) + \frac{2K \sum_{i=1}^N \|v_i\|}{N} \omega_n + \frac{K}{2} \omega_n^2 \right] \\ &= a + b\omega_n + c\omega_n^2, \end{aligned}$$

where

$$a = \rho \left(\frac{2 \sum_{i=1}^N \|v_i\|}{N} (\alpha_0 \eta + 1) + \eta \right), b = \frac{2\rho K \sum_{i=1}^N \|v_i\|}{N}, c = \frac{K\rho}{2}.$$

Again, if $\sum_{i=1}^N \|v_i\|$ and η are small enough and x_0 is sufficiently close to \hat{x}^* , then arguing similarly as in the proof of Theorem 3.1, we can conclude that $x_{n+1} \in B_r(\hat{x}^*)$ and $\|x_n - \hat{x}^*\| = O(\alpha_n)$ for $n = 1, 2, \dots, N(\delta, h)$. Thus, we come to the following convergence result.

Theorem 3.2. *Assume that the assumptions C1-C3 hold for the exact operators $F_i, i = 1, \dots, N$, and conditions (3.12), (3.13), (3.14) and (3.15) are satisfied. If $\sum_{i=1}^N \|v_i\|$ is sufficiently small and x_0 is close enough to \hat{x}^* , then there holds the following estimate*

$$\|x_n - \hat{x}^*\| = O(\alpha_n), \quad n = 1, 2, \dots, N(\delta, h). \quad (3.23)$$

Finally, taking into account the stopping rule (3.18), from Theorem 3.2 we obtain the convergence rate for the parallel regularized Newton-type method (3.16)-(3.17) in noisy data cases.

Corollary 3.1. *Assume that all conditions of Theorem 3.2 are fulfilled. Then*

$$\|x_{n_*} - \hat{x}^*\| = O\left(\delta^{1/2} + h^{1/2}\right), \quad (3.24)$$

where $n_* = N(\delta, h) + 1$.

4. CONCLUSION

Most of existing solution methods for systems of ill posed operator equations deal with Hilbert spaces. In this paper we investigate two parallel iterative regularization methods and a parallel regularized Newton-type method for solving systems of equations involving m -accretive operators in Banach spaces. The convergence analysis of the proposed methods in both free-noise and noisy data cases is provided.

5. ACKNOWLEDGMENTS

The first author of this manuscript wishes to express his gratitude to Vietnam Institute for Advanced Study in Mathematics for financial support.

References

- [1] M. Haltmeier, R. Kowar, A. Leitao, O. Scherzer, Kaczmarz methods for regularizing nonlinear ill-posed equations, I. Convergence analysis, *Inverse Probl. Imaging* 1(2) (2007) 289-298.
- [2] M. Haltmeier, R. Kowar, A. Leitao, O. Scherzer, Kaczmarz methods for regularizing nonlinear ill-posed equation, II. Applications, *Inverse Probl. Imaging* 1(3) (2007) 507-523.
- [3] M. Burger, B. Kaltenbacher, Regularizing Newton-Kaczmarz methods for nonlinear ill-posed problems, *SIAM J. Numer. Anal.* 44 (2006) 153-182.
- [4] A. De Cezaro, M. Haltmeier, A. Leitao, O. Scherzer, On steepest-descent-Kaczmarz method for regularizing systems of nonlinear ill-posed equations, *Appl. Math. Comput.* 202 (2008) 596-607.
- [5] P.K. Anh, C.V. Chung, Parallel iterative regularization methods for solving systems of ill-posed equations, *Appl. Math. Comput.* 212 (2009) 542 - 550.
- [6] P.K. Anh, C.V. Chung, Parallel regularized Newton method for nonlinear ill-posed equations, *Numer. Algor.* 58(3) (2011) 379-398.
- [7] P.K. Anh, V.T. Dzung, Parallel iteratively regularized Gauss-Newton method for systems of nonlinear ill-posed equations, *Inter. J. Computer Math.* DOI: 10.1080/00207160.2013.782399.
- [8] P.K. Anh, C.V. Chung, Parallel hybrid methods for a finite family of relatively nonexpansive mappings, *Numer. Funct. Anal. Optim.* DOI: 10.1080/01630563.2013.830127.
- [9] Ya. I. Alber, I. Ryazantseva, *Nonlinear ill-posed problems of monotone type*, Springer, Dordrecht, 2006.

- [10] J. Diestel, *Geometry of Banach spaces - Selected topics*. Lecture Notes in Mathematics, 485, Springer-Verlag, Berlin, 1975.
- [11] I. Cioranescu, *Geometry of Banach spaces, Duality Mappings and Nonlinear Problems*, Kluwer, Dordrecht, 1990.
- [12] S. Reich, *Review of Geometry of Banach spaces, Duality Mappings and Nonlinear Problems by Ioana Cioranescu*, Kluwer Academic Publishers, Dordrecht, 1990, *Bull. Amer. Math. Soc.* 26 (1992) 367-370.
- [13] S. Reich, *Approximating zeros of accretive operators*, *Proc. Amer. Math. Soc.* 51 (1975) 381-384.
- [14] K. Ball, E. Carlen, E. Lieb, *Sharp uniform convexity and smoothness inequalities for trace norms*, *Invent. Math.* 115 (1994) 463-482.
- [15] Ya. I. Alber, *New results in fixed point theory*, Preprint, Technion, 2000.
- [16] Ya. I. Alber, *On the stability of iterative approximations to fixed point of nonexpansive mappings*, *J. Math. Anal. Appl.* 328 (2007) 958-971.
- [17] H.K. Xu, *Viscosity approximation methods for nonexpensive mappings*, *J. Math. Anal. Appl.* 289(1) (2004) 279-291.
- [18] J. Wang, J. Li, Z. Liu, *Regularization methods for nonlinear ill-posed problems with accretive operators*, *Acta Mathematica Scientia.* 28(1) (2008) 141-150.
- [19] B. Blaschke, A. Neubauer, O. Schezer, *On convergence rates for the iteratively regularized Gauss-Newton method*, *IMA J. Numer. Anal.* 17 (1997) 421-436.
- [20] N. Buong, N.T.H. Phuong, *Convergence Rates in Regularization for Nonlinear Ill-Posed Equations Involving m -Accretive Mappings in Banach Spaces*, *Appl. Math. Sci.* 63 (2012) 3109-3117.
- [21] Ya. I. Alber, C.E. Chidume, H. Zegeye, *Regularization of nonlinear ill-posed equations with accretive operators*, *Fixed Point Theory Appl.* 2005(1)(2005) 11- 33.