

HIGHER TODA BRACKETS AND THE ADAMS SPECTRAL SEQUENCE IN TRIANGULATED CATEGORIES

J. DANIEL CHRISTENSEN AND MARTIN FRANKLAND

ABSTRACT. The Adams spectral sequence is available in any triangulated category equipped with a projective or injective class. Higher Toda brackets can also be defined in a triangulated category, as observed by B. Shipley based on J. Cohen's approach for spectra. We provide a family of definitions of higher Toda brackets, show that they are equivalent to Shipley's, and show that they are self-dual. Our main result is that the Adams differential d_r in any Adams spectral sequence can be expressed as an $(r+1)$ -fold Toda bracket and as an r^{th} order cohomology operation. We also show how the result simplifies under a sparseness assumption, discuss several examples, and give an elementary proof of a result of Heller, which implies that the three-fold Toda brackets in principle determine the higher Toda brackets.

CONTENTS

1. Introduction	1
2. The Adams spectral sequence	4
3. 3-fold Toda brackets	8
4. Adams d_2 in terms of 3-fold Toda brackets	12
5. Higher Toda brackets	15
6. Higher order operations determine d_r	22
7. Sparse rings of operations	26
Appendix A. Computations in the stable module category of a group	32
Appendix B. 3-fold Toda brackets determine the triangulated structure	34
References	36

1. INTRODUCTION

The Adams spectral sequence is an important tool in stable homotopy theory. Given finite spectra X and Y , the classical Adams spectral sequence is

$$E_2^{s,t} = \text{Ext}_{\mathcal{A}}^{s,t}(H^*Y, H^*X) \Rightarrow [\Sigma^{t-s}X, Y_p^\wedge],$$

where $H^*X := H^*(X; \mathbb{F}_p)$ denotes mod p cohomology and $\mathcal{A} = H\mathbb{F}_p^*H\mathbb{F}_p$ denotes the mod p Steenrod algebra. Determining the differentials in the Adams spectral sequence generally requires a combination of techniques and much ingenuity. The approach that provides a basis for our work is found in [28], where Maunder showed that the differential d_r in this spectral sequence is determined by r^{th} order cohomology operations, which we now review.

Date: April 2, 2018.

2010 Mathematics Subject Classification. Primary 55T15; Secondary 18E30, 55S20.

Key words and phrases. triangulated category, Adams spectral sequence, Toda bracket, cohomology operation, differential, higher order operation, projective class.

A primary cohomology operation in this context is simply an element of the Steenrod algebra, and it is immediate from the construction of the Adams spectral sequence that the differential d_1 is given by primary cohomology operations. A secondary cohomology operation corresponds to a relation among primary operations, and is partially defined and multi-valued: it is defined on the kernel of a primary operation and takes values in the cokernel of another primary operation. Tertiary operations correspond to relations between relations, and have correspondingly more complicated domains and codomains. The pattern continues for higher operations. Using that cohomology classes are representable, secondary cohomology operations can also be expressed using 3-fold Toda brackets involving the cohomology class and two operations whose composite is null. However, what one obtains in general is a *subset* of the Toda bracket with less indeterminacy. This observation will be the key to our generalization of Maunder’s result to other Adams spectral sequences in other categories.

The starting point of this paper is the following observation. On the one hand, the Adams spectral sequence can be constructed in any triangulated category equipped with a projective class or an injective class [13]. For example, the classical Adams spectral sequence is constructed in the stable homotopy category with the injective class consisting of retracts of products $\prod_i \Sigma^{n_i} H\mathbb{F}_p$. On the other hand, higher Toda brackets can also be defined in an arbitrary triangulated category. This was done by Shipley in [40], based on Cohen’s construction for spaces and spectra [15], and was studied further in [36]. The goal of this paper is to describe precisely how the Adams d_r can be described as a particular subset of an $(r+1)$ -fold Toda bracket which can be viewed as an r^{th} order cohomology operation, all in the context of a general triangulated category.

Triangulated categories arise throughout mathematics, so our work applies in various situations. As an example, we give calculations involving the Adams spectral sequence in the stable module category of a group algebra. Even in stable homotopy theory, there are a variety of Adams spectral sequences, such as the Adams–Novikov spectral sequence or the motivic Adams spectral sequence, and our results apply to all of them. Moreover, by working with minimal structure, our approach gains a certain elegance.

Organization and main results. In Section 2, we review the construction of the Adams spectral sequence in a triangulated category equipped with a projective class or an injective class. In Section 3, we review the construction of 3-fold Toda brackets in a triangulated category and some of their basic properties. Section 4 describes how the Adams d_2 is given by 3-fold Toda brackets. This section serves as a warm-up for Section 6.

In Section 5, we recall the construction of higher Toda brackets in a triangulated category via filtered objects. We provide a family of alternate constructions, and prove that they are all equivalent. The main result is Theorem 5.11, which says roughly the following.

Theorem. *There is an inductive way of computing an n -fold Toda bracket $\langle f_n, \dots, f_1 \rangle \subseteq \mathcal{T}(\Sigma^{n-2}X_0, X_n)$, where the inductive step picks three consecutive maps and reduces the length by one. The $(n-2)!$ ways of doing this yield the same subset, up to an explicit sign.*

As a byproduct, we obtain Corollary 5.13, which would be tricky to prove directly from the filtered object definition.

Corollary. *Toda brackets are self-dual up to suspension: $\langle f_n, \dots, f_1 \rangle \subseteq \mathcal{T}(\Sigma^{n-2}X_0, X_n)$ corresponds to the Toda bracket computed in the opposite category*

$$\langle f_1, \dots, f_n \rangle \subseteq \mathcal{T}^{\text{op}}\left(\Sigma^{-(n-2)}X_n, X_0\right) = \mathcal{T}(X_0, \Sigma^{-(n-2)}X_n).$$

Section 6 establishes how the Adams d_r is given by $(r + 1)$ -fold Toda brackets. Our main results are Theorems 6.1 and 6.5, which say roughly the following.

Theorem. *Let $[x] \in E_r^{s,t}$ be a class in the E_r term of the Adams spectral sequence. As subsets of $E_1^{s+r,t+r-1}$, we have*

$$\begin{aligned} d_r[x] &= \langle \Sigma^{r-1}d_1, \dots, \Sigma^2d_1, \Sigma d_1, \Sigma p_{s+1}, \delta_s x \rangle \\ &= \langle \Sigma^{r-1}d_1!, \dots!, \Sigma d_1!, d_1, x \rangle. \end{aligned}$$

Here, d_1 , p_{s+1} , and δ_s are maps appearing in the Adams resolution of Y , where each d_1 is a primary cohomology operation. The first expression for $d_r[x]$ is an $(r + 1)$ -fold Toda bracket. The second expression denotes an appropriate subset of the bracket $\langle \Sigma^{r-1}d_1, \dots, \Sigma d_1, d_1, x \rangle$ with some choices dictated by the Adams resolution of Y . This description exhibits $d_r[x]$ as an r^{th} order cohomology operation applied to x .

In Section 7, we show that when certain sparseness assumptions are made, the subset $\langle \Sigma^{r-1}d_1!, \dots!, \Sigma d_1!, d_1, x \rangle$ coincides with the full Toda bracket, and give examples of this phenomenon. See Theorem 7.14, Proposition 7.15, and Example 7.17. The main application is to computing maps in the homotopy category of R -module spectra, for a ring spectrum R whose coefficient ring π_*R is sufficiently sparse, such as ku . See Example 7.21.

In Appendix A, we compute examples of Toda brackets in stable module categories. In particular, Proposition A.1 provides an example where the inclusion $d_2[x] \subseteq \langle \Sigma d_1, d_1, x \rangle$ is proper. Appendix B provides for the record a short, simple proof of a theorem due to Heller, that 3-fold Toda brackets determine the triangulated structure. As a corollary, we note that the 3-fold Toda brackets indirectly determine the higher Toda brackets.

Related work. Detailed treatments of secondary operations can be found in [1, §3.6], where Adams used secondary cohomology operations to solve the Hopf invariant one problem, [32, Chapter 16], and [19, Chapter 4].

There are various approaches to higher order cohomology operations and higher Toda brackets in the literature, many of which use some form of enrichment in spaces, chain complexes, or groupoids; see for instance [42], [27], [24], and [23]. In this paper, we work solely with the triangulated structure, without enhancement, and provide no comparison to those other approaches.

In [5] and [6], Baues and Jibladze express the Adams d_2 in terms of secondary cohomology operations, and this is generalized to higher differentials by Baues and Blanc in [7]. Their approach starts with an injective resolution as in Diagram (2.3), and witnesses the equations $d_1d_1 = 0$ by providing suitably coherent null-homotopies, described using mapping spaces. Using this coherence data, the authors express a representative of $d_r[x]$ as a specific element of the Toda bracket $\langle \Sigma^{r-1}d_1, \dots, \Sigma d_1, d_1, x \rangle$. While this approach makes use of an enrichment, we suspect that by translating the (higher dimensional) null-homotopies into lifts to fibers or extensions to cofibers, one could relate their expression for $d_r[x]$ to ours.

Acknowledgments. We thank Robert Bruner, Dan Isaksen, Peter Jorgensen, Fernando Muro, Irakli Patchkoria, Steffen Sagave, and Dylan Wilson for helpful conversations, as well as the referee for their useful comments. The second author also thanks the Max-Planck-Institut für Mathematik for its hospitality. The second author was partially funded by a grant of the DFG SPP 1786: Homotopy Theory and Algebraic Geometry.

2. THE ADAMS SPECTRAL SEQUENCE

In this section, we recall the construction of the Adams spectral sequence in a triangulated category, along with some of its features. We follow [13, §4], or rather its dual. Some references for the classical Adams spectral sequence are [2, §III.15], [26, Chapter 16], and [10]. Background material on triangulated categories can be found in [33, Chapter 1], [26, Appendix 2], and [44, Chapter 10]. We assume that the suspension functor Σ is an equivalence, with chosen inverse Σ^{-1} . Moreover, we assume we have chosen natural isomorphisms $\Sigma\Sigma^{-1} \cong \text{id}$ and $\Sigma^{-1}\Sigma \cong \text{id}$ making Σ and Σ^{-1} into an adjoint equivalence. We silently use these isomorphisms when needed, e.g., when we say that a triangle of the form $\Sigma^{-1}Z \rightarrow X \rightarrow Y \rightarrow Z$ is distinguished.

Definition 2.1 ([13, Proposition 2.6]). A **projective class** in a triangulated category \mathcal{T} is a pair $(\mathcal{P}, \mathcal{N})$ where \mathcal{P} is a class of objects and \mathcal{N} is a class of maps satisfying the following properties.

- (1) A map $f: X \rightarrow Y$ is in \mathcal{N} if and only if the induced map

$$f_*: \mathcal{T}(P, X) \rightarrow \mathcal{T}(P, Y)$$

is zero for all P in \mathcal{P} . In other words, \mathcal{N} consists of the **\mathcal{P} -null** maps.

- (2) An object P is in \mathcal{P} if and only if the induced map

$$f_*: \mathcal{T}(P, X) \rightarrow \mathcal{T}(P, Y)$$

is zero for all f in \mathcal{N} .

- (3) For every object X , there is a distinguished triangle $P \rightarrow X \xrightarrow{f} Y \rightarrow \Sigma P$, where P is in \mathcal{P} and f is in \mathcal{N} .

In particular, the class \mathcal{P} is closed under arbitrary coproducts and retracts. The objects in \mathcal{P} are called **projective**.

Definition 2.2. A projective class is **stable** if it is closed under shifts, i.e., $P \in \mathcal{P}$ implies $\Sigma^n P \in \mathcal{P}$ for all $n \in \mathbb{Z}$.

We will assume for convenience that our projective class is stable. We suspect that many of the results can be adapted to unstable projective classes, with a careful treatment of shifts.

Definition 2.3. Let \mathcal{P} be a projective class. A map $f: X \rightarrow Y$ is called

- (1) **\mathcal{P} -epic** if the map

$$f_*: \mathcal{T}(P, X) \rightarrow \mathcal{T}(P, Y)$$

is surjective for all $P \in \mathcal{P}$. Equivalently, the map to the cofiber $Y \rightarrow C_f$ is \mathcal{P} -null.

- (2) **\mathcal{P} -monic** if the map

$$f_*: \mathcal{T}(P, X) \rightarrow \mathcal{T}(P, Y)$$

is injective for all $P \in \mathcal{P}$. Equivalently, the map from the fiber $\Sigma^{-1}C_f \rightarrow X$ is \mathcal{P} -null.

Example 2.4. Let \mathcal{T} be the stable homotopy category and \mathcal{P} the projective class consisting of retracts of wedges of spheres $\bigvee_i S^{n_i}$. This is called the *ghost projective class*, studied for instance in [13, §7].

Now we dualize everything.

Definition 2.5. An **injective class** in a triangulated category \mathcal{T} is a projective class in the opposite category \mathcal{T}^{op} . Explicitly, it is a pair $(\mathcal{I}, \mathcal{N})$ where \mathcal{I} is a class of objects and \mathcal{N} is a class of maps satisfying the following properties.

- (1) A map $f: X \rightarrow Y$ is in \mathcal{N} if and only if the induced map

$$f^*: \mathcal{T}(Y, I) \rightarrow \mathcal{T}(X, I)$$

is zero for all I in \mathcal{I} .

- (2) An object I is in \mathcal{I} if and only if the induced map

$$f^*: \mathcal{T}(Y, I) \rightarrow \mathcal{T}(X, I)$$

is zero for all f in \mathcal{N} .

- (3) For every object X , there is a distinguished triangle $\Sigma^{-1}I \rightarrow W \xrightarrow{f} X \rightarrow I$, where I is in \mathcal{I} and f is in \mathcal{N} .

In particular, the class \mathcal{I} is closed under arbitrary products and retracts. The objects in \mathcal{I} are called **injective**. Just as for projective classes, we will assume for convenience that our injective class is stable.

Example 2.6. Let \mathcal{T} be the stable homotopy category. Take \mathcal{N} to be the class of maps inducing zero on mod p cohomology and \mathcal{I} to be the retracts of (arbitrary) products $\prod_i \Sigma^{n_i} H\mathbb{F}_p$ with $n_i \in \mathbb{Z}$. One can generalize this example to any cohomology theory (spectrum) E instead of $H\mathbb{F}_p$, letting \mathcal{I}_E denote the injective class consisting of retracts of products $\prod_i \Sigma^{n_i} E$.

Definition 2.7. Let \mathcal{I} be an injective class. A map $f: X \rightarrow Y$ is called

- (1) **\mathcal{I} -monic** if the map

$$f^*: \mathcal{T}(Y, I) \rightarrow \mathcal{T}(X, I)$$

is surjective for all $I \in \mathcal{I}$. Equivalently, the map from the fiber $\Sigma^{-1}C_f \rightarrow X$ is \mathcal{I} -null.

- (2) **\mathcal{I} -epic** if the map

$$f^*: \mathcal{T}(Y, I) \rightarrow \mathcal{T}(X, I)$$

is injective for all $I \in \mathcal{I}$. Equivalently, the map to the cofiber $Y \rightarrow C_f$ is \mathcal{I} -null.

Remark 2.8. The projectives and \mathcal{P} -epic maps determine each other via the lifting property

$$\begin{array}{ccc} & X & \\ & \nearrow & \downarrow f \\ P & \longrightarrow & Y. \end{array}$$

Dually, the injectives and \mathcal{I} -monic maps determine each other via the extension property

$$\begin{array}{ccc} X & \longrightarrow & I \\ \downarrow f & & \nearrow \\ Y & & \end{array}$$

This is part of the equivalent definition of a projective (resp. injective) class described in [13, Proposition 2.4].

Convention 2.9. We will implicitly use the natural isomorphism $\mathcal{T}(A, B) \cong \mathcal{T}(\Sigma^k A, \Sigma^k B)$ sending a map f to $\Sigma^k f$.

Definition 2.10. An **Adams resolution** of an object X in \mathcal{T} with respect to a projective class $(\mathcal{P}, \mathcal{N})$ is a diagram

$$\begin{array}{ccccccc} X = X_0 & \xrightarrow{i_0} & X_1 & \xrightarrow{i_1} & X_2 & \xrightarrow{i_2} & X_3 \longrightarrow \dots \\ & \swarrow p_0 & \swarrow \delta_0 & \swarrow p_1 & \swarrow \delta_1 & \swarrow p_2 & \swarrow \delta_2 \\ & & P_0 & & P_1 & & P_2 \end{array} \quad (2.1)$$

where every P_s is projective, every map i_s is in \mathcal{N} , and every triangle $P_s \xrightarrow{p_s} X_s \xrightarrow{i_s} X_{s+1} \xrightarrow{\delta_s} \Sigma P_s$ is distinguished. Here the arrows $\delta_s: X_{s+1} \rightarrow \Sigma P_s$ denote degree-shifting maps, namely, maps $\delta_s: X_{s+1} \rightarrow \Sigma P_s$.

Dually, an **Adams resolution** of an object Y in \mathcal{T} with respect to an injective class $(\mathcal{I}, \mathcal{N})$ is a diagram

$$\begin{array}{ccccccc}
 Y = Y_0 & \xleftarrow{i_0} & Y_1 & \xleftarrow{i_1} & Y_2 & \xleftarrow{i_2} & Y_3 \xleftarrow{\dots} \\
 & \searrow p_0 & \nearrow \delta_0 & \searrow p_1 & \nearrow \delta_1 & \searrow p_2 & \nearrow \delta_2 \\
 & & I_0 & & I_1 & & I_2
 \end{array} \tag{2.2}$$

where every I_s is injective, every map i_s is in \mathcal{N} , and every triangle $\Sigma^{-1}I_s \xrightarrow{\Sigma^{-1}\delta_s} Y_{s+1} \xrightarrow{i_s} Y_s \xrightarrow{p_s} I_s$ is distinguished.

From now on, fix a triangulated category \mathcal{T} and a (stable) injective class $(\mathcal{I}, \mathcal{N})$ in \mathcal{T} . By repeatedly using condition (3) in the definition of an injective class, we get the following lemma.

Lemma 2.11. *Every object Y of \mathcal{T} admits an Adams resolution.*

Given an object X and an Adams resolution of Y , applying $\mathcal{T}(X, -)$ yields an exact couple

$$\begin{array}{ccc}
 \bigoplus_{s,t} \mathcal{T}(\Sigma^{t-s} X, Y_s) & \xrightarrow{i = \bigoplus (i_s)_*} & \bigoplus_{s,t} \mathcal{T}(\Sigma^{t-s} X, Y_s) \\
 & \searrow \delta = \bigoplus (\delta_s)_* & \swarrow p = \bigoplus (p_s)_* \\
 & \bigoplus_{s,t} \mathcal{T}(\Sigma^{t-s} X, I_s) &
 \end{array}$$

and thus a spectral sequence with E_1 term

$$E_1^{s,t} = \mathcal{T}(\Sigma^{t-s} X, I_s) \cong \mathcal{T}(\Sigma^t X, \Sigma^s I_s)$$

and differentials

$$d_r: E_r^{s,t} \rightarrow E_r^{s+r, t+r-1}$$

given by $d_r = p \circ i^{-(r-1)} \circ \delta$, where i^{-1} means choosing an i -preimage. This is called the **Adams spectral sequence** with respect to the injective class \mathcal{I} abutting to $\mathcal{T}(\Sigma^{t-s} X, Y)$.

Lemma 2.12. *The E_2 term is given by*

$$E_2^{s,t} = \text{Ext}_{\mathcal{T}}^{s,t}(X, Y) := \text{Ext}_{\mathcal{T}}^s(\Sigma^t X, Y)$$

where $\text{Ext}_{\mathcal{T}}^s(X, Y)$ denotes the s^{th} derived functor of $\mathcal{T}(X, -)$ (relative to the injective class \mathcal{I}) applied to the object Y .

Proof. The Adams resolution of Y yields an \mathcal{I} -injective resolution of Y

$$0 \longrightarrow Y \xrightarrow{p_0} I_0 \xrightarrow{(\Sigma p_1)\delta_0} \Sigma I_1 \xrightarrow{(\Sigma^2 p_2)(\Sigma \delta_1)} \Sigma^2 I_2 \longrightarrow \dots \tag{2.3}$$

□

Remark 2.13. We do not assume that the injective class \mathcal{I} generates, i.e., that every non-zero object X admits a non-zero map $X \rightarrow I$ to an injective. Hence, we do not expect the Adams spectral sequence to be conditionally convergent in general; c.f. [13, Proposition 4.4].

Example 2.14. Let E be a commutative (homotopy) ring spectrum. A spectrum is called E -**injective** if it is a retract of $E \wedge W$ for some W [22, Definition 2.22]. A map of spectra $f: X \rightarrow Y$ is called E -**monic** if the map $E \wedge f: E \wedge X \rightarrow E \wedge Y$ is a split monomorphism. The E -injective objects and E -monic maps form an injective class in the stable homotopy category. The Adams spectral sequence associated to this injective class is the *Adams spectral sequence based on E -homology*, as described in [35, Definition 2.2.4], also called the *unmodified Adams spectral sequence* in [22, §2.2]. Further assumptions are needed in order to identify the E_2 term as Ext groups in E_*E -comodules.

Definition 2.15. The \mathcal{I} -**cohomology** of an object X is the family of abelian groups $H^I(X) := \mathcal{T}(X, I)$ indexed by the injective objects $I \in \mathcal{I}$.

A **primary operation** in \mathcal{I} -cohomology is a natural transformation $H^I(X) \rightarrow H^J(X)$ of functors $\mathcal{T}^{\text{op}} \rightarrow \text{Ab}$. Equivalently, by the (additive) Yoneda lemma, a primary operation is a map $I \rightarrow J$ in \mathcal{I} .

Example 2.16. The differential d_1 is given by primary operations. More precisely, let $x \in E_1^{s,t}$ be a map $x: \Sigma^{t-s}X \rightarrow I_s$. Then $d_1(x) \in E_1^{s+1,t}$ is the composite

$$\Sigma^{t-s}X \xrightarrow{x} I_s \xrightarrow{\delta_s} \Sigma Y_{s+1} \xrightarrow{\Sigma p_{s+1}} \Sigma I_{s+1}.$$

In other words, $d_1(x)$ is obtained by applying the primary operation $d_1 := (\Sigma p_{s+1})\delta_s: I_s \rightarrow \Sigma I_{s+1}$ to x .

Proposition 2.17. A primary operation $\theta: I \rightarrow J$ appears as $d_1: I_s \dashrightarrow I_{s+1}$ in some Adams resolution if and only if θ admits an \mathcal{I} -epi – \mathcal{I} -mono factorization.

Proof. The condition is necessary by construction. In the factorization $d_1 = (\Sigma p_{s+1})\delta_s$, the map δ_s is \mathcal{I} -epic while p_{s+1} is \mathcal{I} -monic.

To prove sufficiency, assume given a factorization $\theta = iq: I \rightarrow W \rightarrow J$, where $q: I \twoheadrightarrow W$ is \mathcal{I} -epic and $i: W \hookrightarrow J$ is \mathcal{I} -monic. Taking the fiber of q twice yields the distinguished triangle

$$\Sigma^{-1}W \longrightarrow Y_0 \twoheadrightarrow I \xrightarrow{q} W$$

which we relabel

$$Y_1 \xrightarrow{i_0} Y_0 \twoheadrightarrow I \xrightarrow{\delta_0} \Sigma Y_1.$$

Relabeling the given map $i: W \hookrightarrow J$ as $\Sigma p_1: \Sigma Y_1 \hookrightarrow \Sigma I_1$, we can continue the usual construction of an Adams resolution of Y_0 as illustrated in Diagram (2.2), in which $\theta = iq$ appears as the composite $(\Sigma p_1)\delta_0$. Note that by the same argument, for any $s \geq 0$, θ appears as $d_1: I_s \dashrightarrow I_{s+1}$ in some (other) Adams resolution. \square

Example 2.18. Not every primary operation appears as d_1 in an Adams resolution. For example, consider the stable homotopy category with the projective class \mathcal{P} generated by the sphere spectrum $S = S^0$, that is, \mathcal{P} consists of retracts of wedges of spheres. The \mathcal{P} -epis (resp. \mathcal{P} -monos) consist of the maps which are surjective (resp. injective) on homotopy groups. The primary operation $2: S \rightarrow S$ does *not* admit a \mathcal{P} -epi – \mathcal{P} -mono factorization.

Indeed, assume that $2 = iq: S \twoheadrightarrow W \hookrightarrow S$ is such a factorization. We will show that this implies $\pi_2(S/2) = \mathbb{Z}/2 \oplus \mathbb{Z}/2$, contradicting the known fact $\pi_2(S/2) = \mathbb{Z}/4$. Here $S/2$ denotes the mod 2 Moore spectrum, sitting in the cofiber sequence $S \xrightarrow{2} S \rightarrow S/2$.

By the octahedral axiom applied to the factorization $2 = iq$, there is a diagram

$$\begin{array}{ccccccc}
 S & \xrightarrow{q} & W & \longrightarrow & C_q & \xrightarrow{\delta'} & S^1 \\
 \parallel & & \downarrow i & & \downarrow \alpha & & \parallel \\
 S & \xrightarrow{2} & S & \longrightarrow & S/2 & \xrightarrow{\delta} & S^1 \\
 & & \downarrow j & & \downarrow \beta & & \\
 & & C_i & \xlongequal{\quad} & C_i & &
 \end{array}$$

with distinguished rows and columns. The long exact sequence in homotopy yields $\pi_n C_q = {}_2\pi_{n-1}S$, where the induced map $\pi_n(\delta'): \pi_n C_q \rightarrow \pi_n S^1$ corresponds to the inclusion ${}_2\pi_{n-1}S \hookrightarrow \pi_{n-1}S$. Likewise, we have $\pi_n C_i = (\pi_n S)/2$, where the induced map $\pi_n(j): \pi_n S \rightarrow \pi_n C_i$ corresponds to the quotient map $\pi_n S \twoheadrightarrow (\pi_n S)/2$. The defining cofiber sequence $S \xrightarrow{2} S \rightarrow S/2$ yields the exact sequence

$$\pi_n S \xrightarrow{2} \pi_n S \longrightarrow \pi_n(S/2) \xrightarrow{\pi_n \delta} \pi_{n-1} S \xrightarrow{2} \pi_{n-1} S$$

which in turn yields the short exact sequence

$$0 \longrightarrow (\pi_n S)/2 \longrightarrow \pi_n(S/2) \xrightarrow{\pi_n \delta} {}_2\pi_{n-1} S \longrightarrow 0.$$

The map $\pi_n(\alpha): {}_2\pi_{n-1}S \rightarrow \pi_n(S/2)$ is a splitting of this sequence, because of the equality $\pi_n(\delta)\pi_n(\alpha) = \pi_n(\delta\alpha) = \pi_n(\delta')$. However, the short exact sequence does not split in the case $n = 2$, by the isomorphism $\pi_2(S/2) = \mathbb{Z}/4$. For references, see [39, Proposition II.6.48], [38, Proposition 4], and [31].

3. 3-FOLD TODA BRACKETS

In this section, we review different constructions of 3-fold Toda brackets and some of their properties.

Definition 3.1. Let $X_0 \xrightarrow{f_1} X_1 \xrightarrow{f_2} X_2 \xrightarrow{f_3} X_3$ be a diagram in a triangulated category \mathcal{T} . We define subsets of $\mathcal{T}(\Sigma X_0, X_3)$ as follows.

- The **iterated cofiber Toda bracket** $\langle f_3, f_2, f_1 \rangle_{cc} \subseteq \mathcal{T}(\Sigma X_0, X_3)$ consists of all maps $\psi: \Sigma X_0 \rightarrow X_3$ that appear in a commutative diagram

$$\begin{array}{ccccccc}
 X_0 & \xrightarrow{f_1} & X_1 & \longrightarrow & C_{f_1} & \longrightarrow & \Sigma X_0 \\
 \parallel & & \parallel & & \downarrow \varphi & & \downarrow \psi \\
 X_0 & \xrightarrow{f_1} & X_1 & \xrightarrow{f_2} & X_2 & \xrightarrow{f_3} & X_3
 \end{array} \tag{3.1}$$

where the top row is distinguished.

- The **fiber-cofiber Toda bracket** $\langle f_3, f_2, f_1 \rangle_{fc} \subseteq \mathcal{T}(\Sigma X_0, X_3)$ consists of all composites $\beta \circ \Sigma\alpha: \Sigma X_0 \rightarrow X_3$, where α and β appear in a commutative diagram

$$\begin{array}{ccccccc}
 X_0 & \xrightarrow{f_1} & X_1 & & & & \\
 \alpha \downarrow & & \parallel & & & & \\
 \Sigma^{-1}C_{f_2} & \longrightarrow & X_1 & \xrightarrow{f_2} & X_2 & \longrightarrow & C_{f_2} \\
 & & & & \parallel & & \downarrow \beta \\
 & & & & X_2 & \xrightarrow{f_3} & X_3
 \end{array} \tag{3.2}$$

where the middle row is distinguished.

- The **iterated fiber Toda bracket** $\langle f_3, f_2, f_1 \rangle_{ff} \subseteq \mathcal{T}(\Sigma X_0, X_3)$ consists of all maps $\Sigma\delta: \Sigma X_0 \rightarrow X_3$ where δ appears in a commutative diagram

$$\begin{array}{ccccccc}
 X_0 & \xrightarrow{f_1} & X_1 & \xrightarrow{f_2} & X_2 & \xrightarrow{f_3} & X_3 \\
 \delta \downarrow & & \gamma \downarrow & & \parallel & & \parallel \\
 \Sigma^{-1}X_3 & \longrightarrow & \Sigma^{-1}C_{f_3} & \longrightarrow & X_2 & \xrightarrow{f_3} & X_3
 \end{array} \tag{3.3}$$

where the bottom row is distinguished.

Remark 3.2. In the literature, there are variations of these definitions, which sometimes differ by a sign. With the notion of cofiber sequence implicitly used in [43], our definitions agree with Toda's. The Toda bracket also depends on the choice of triangulation. Given a triangulation, there is an associated negative triangulation whose distinguished triangles are those triangles whose negatives are distinguished in the original triangulation (see [3]). Negating a triangulation negates the 3-fold Toda brackets. Dan Isaksen has pointed out to us that in the stable homotopy category there are 3-fold Toda brackets which are not equal to their own negatives. For example, Toda showed in [43, Section VI.v, and Theorems 7.4 and 14.1] that the Toda bracket $\langle 2\sigma, 8, \nu \rangle$ has no indeterminacy and contains an element ζ of order 8. We give another example in Example A.4.

The following proposition can be found in [36, Remark 4.5 and Figure 2] and was kindly pointed out by Fernando Muro. It is also proved in [30, §4.6]. We provide a different proof more in the spirit of this article. In the case of spaces, it was originally proved by Toda [43, Proposition 1.7].

Proposition 3.3. *The iterated cofiber, fiber-cofiber, and iterated fiber definitions of Toda brackets coincide. More precisely, for any diagram $X_0 \xrightarrow{f_1} X_1 \xrightarrow{f_2} X_2 \xrightarrow{f_3} X_3$ in \mathcal{T} , the following subsets of $\mathcal{T}(\Sigma X_0, X_3)$ are equal:*

$$\langle f_3, f_2, f_1 \rangle_{cc} = \langle f_3, f_2, f_1 \rangle_{fc} = \langle f_3, f_2, f_1 \rangle_{ff}.$$

Proof. We will prove the first equality; the second equality is dual.

(\supseteq) Let $\beta(\Sigma\alpha) \in \langle f_3, f_2, f_1 \rangle_{fc}$ be obtained from maps α and β as in Diagram (3.2). Now consider the diagram with distinguished rows

$$\begin{array}{ccccccc}
 X_0 & \xrightarrow{f_1} & X_1 & \longrightarrow & C_{f_1} & \longrightarrow & \Sigma X_0 \\
 \downarrow \alpha & & \parallel & & \downarrow \varphi & & \downarrow \Sigma\alpha \\
 \Sigma^{-1}C_{f_2} & \longrightarrow & X_1 & \xrightarrow{f_2} & X_2 & \longrightarrow & C_{f_2} \\
 & & & & \parallel & & \downarrow \beta \\
 & & & & X_2 & \xrightarrow{f_3} & X_3
 \end{array}$$

where there exists a filler $\varphi: C_{f_1} \rightarrow X_2$. The commutativity of the tall rectangle on the right exhibits the membership $\beta(\Sigma\alpha) \in \langle f_3, f_2, f_1 \rangle_{cc}$.

(\subseteq) Let $\psi \in \langle f_3, f_2, f_1 \rangle_{cc}$ be as in Diagram (3.1). The octahedral axiom comparing the cofibers of q_1 , φ , and $\varphi \circ q_1 = f_2$ yields a commutative diagram

$$\begin{array}{ccccccc}
 & & & & \Sigma^{-1}C_\varphi & \xlongequal{\quad} & \Sigma^{-1}C_\varphi \\
 & & & & \downarrow -\Sigma^{-1}\iota & & \downarrow -\Sigma^{-1}\eta \\
 X_0 & \xrightarrow{f_1} & X_1 & \xrightarrow{q_1} & C_{f_1} & \xrightarrow{\iota_1} & \Sigma X_0 & \xrightarrow{-\Sigma f_1} & \Sigma X_1 \\
 \downarrow \alpha & & \parallel & & \downarrow \varphi & & \downarrow \Sigma\alpha & & \parallel \\
 \Sigma^{-1}C_{f_2} & \xrightarrow{-\Sigma^{-1}\iota_2} & X_1 & \xrightarrow{f_2} & X_2 & \xrightarrow{q_2} & C_{f_2} & \xrightarrow{\iota_2} & \Sigma X_1 \\
 & & & & \downarrow q & & \downarrow \xi & & \\
 & & & & C_\varphi & \xlongequal{\quad} & C_\varphi, & & \\
 & & & & \swarrow f_3 & & \swarrow \beta & & \\
 & & & & X_3 & & & & \\
 & & & & \swarrow \theta & & & & \\
 & & & & C_\varphi & & & &
 \end{array}$$

where the rows and columns are distinguished. By exactness of the sequence

$$\mathcal{T}(C_{f_2}, X_3) \xrightarrow{(\Sigma\alpha)^*} \mathcal{T}(\Sigma X_0, X_3) \xrightarrow{(-\Sigma^{-1}\eta)^*} \mathcal{T}(\Sigma^{-1}C_\varphi, X_3)$$

there exists a map $\beta: C_{f_2} \rightarrow X_3$ satisfying $\psi = \beta(\Sigma\alpha)$ if and only if the restriction of ψ to the fiber $\Sigma^{-1}C_\varphi$ of $\Sigma\alpha$ is zero. That condition does hold: one readily checks the equality $\psi(-\Sigma^{-1}\eta) = 0$. The chosen map $\beta: C_{f_2} \rightarrow X_3$ might *not* satisfy the equation $\beta q_2 = f_3$, but we will correct it to another map β' which does. The error term $f_3 - \beta q_2$ is killed by restriction along φ , and therefore factors through the cofiber of φ , i.e., there exists a factorization

$$f_3 - \beta q_2 = \theta \iota$$

for some $\theta: C_\varphi \rightarrow X_3$. The corrected map $\beta' := \beta + \theta\xi: C_{f_2} \rightarrow X_3$ satisfies $\beta' q_2 = f_3$. Moreover, this corrected map β' still satisfies $\beta'(\Sigma\alpha) = \psi = \beta(\Sigma\alpha)$, since the correction term satisfies $\theta\xi(\Sigma\alpha) = 0$. \square

Thanks to the proposition, we can write $\langle f_3, f_2, f_1 \rangle$ if we do not need to specify a particular definition of the Toda bracket.

We also recall this well-known fact, and leave the proof as an exercise:

Lemma 3.4. For any diagram $X_0 \xrightarrow{f_1} X_1 \xrightarrow{f_2} X_2 \xrightarrow{f_3} X_3$ in \mathcal{T} , the subset $\langle f_3, f_2, f_1 \rangle$ of $\mathcal{T}(\Sigma X_0, X_3)$ is a coset of the subgroup

$$(f_3)_* \mathcal{T}(\Sigma X_0, X_2) + (\Sigma f_1)^* \mathcal{T}(\Sigma X_1, X_3). \quad \square$$

The displayed subgroup is called the **indeterminacy**, and when it is trivial, we say that the Toda bracket **has no indeterminacy**.

Lemma 3.5. Consider maps $X_0 \xrightarrow{f_1} X_1 \xrightarrow{f_2} X_2 \xrightarrow{f_3} X_3 \xrightarrow{f_4} X_4$. Then the following inclusions of subsets of $\mathcal{T}(\Sigma X_0, X_4)$ hold.

- (1) $f_4 \langle f_3, f_2, f_1 \rangle \subseteq \langle f_4 f_3, f_2, f_1 \rangle$
- (2) $\langle f_4, f_3, f_2 \rangle f_1 \subseteq \langle f_4, f_3, f_2 f_1 \rangle$
- (3) $\langle f_4 f_3, f_2, f_1 \rangle \subseteq \langle f_4, f_3 f_2, f_1 \rangle$
- (4) $\langle f_4, f_3, f_2 f_1 \rangle \subseteq \langle f_4, f_3 f_2, f_1 \rangle$.

Proof. (1)-(2) These inclusions are straightforward.

(3)-(4) Using the iterated cofiber definition, the subset $\langle f_4 f_3, f_2, f_1 \rangle_{cc}$ consists of the maps $\psi: \Sigma X_0 \rightarrow X_4$ appearing in a commutative diagram

$$\begin{array}{ccccccc} X_0 & \xrightarrow{f_1} & X_1 & \longrightarrow & C_{f_1} & \longrightarrow & \Sigma X_0 \\ \parallel & & \parallel & & \downarrow \varphi & & \downarrow \psi \\ X_0 & \xrightarrow{f_1} & X_1 & \xrightarrow{f_2} & X_2 & \xrightarrow{f_3} & X_3 & \xrightarrow{f_4} & X_4 \end{array}$$

where the top row is distinguished. Given such a diagram, the diagram

$$\begin{array}{ccccccc} X_0 & \xrightarrow{f_1} & X_1 & \longrightarrow & C_{f_1} & \longrightarrow & \Sigma X_0 \\ \parallel & & \parallel & & \searrow f_3 \varphi & & \downarrow \psi \\ X_0 & \xrightarrow{f_1} & X_1 & \xrightarrow{f_2} & X_2 & \xrightarrow{f_3} & X_3 & \xrightarrow{f_4} & X_4 \end{array}$$

exhibits the membership $\psi \in \langle f_4, f_3 f_2, f_1 \rangle_{cc}$. A similar argument can be used to prove the inclusion $\langle f_4, f_3, f_2 f_1 \rangle_{ff} \subseteq \langle f_4, f_3 f_2, f_1 \rangle_{ff}$. \square

Example 3.6. The inclusion $\langle f_4 f_3, f_2, f_1 \rangle \subseteq \langle f_4, f_3 f_2, f_1 \rangle$ need not be an equality. For example, consider the maps $X \xrightarrow{0} Y \xrightarrow{1} Y \xrightarrow{0} Z \xrightarrow{1} Z$. The Toda brackets being compared are

$$\begin{aligned} \langle 1_Z 0, 1_Y, 0 \rangle &= \langle 0, 1_Y, 0 \rangle = \{0\} \\ \langle 1_Z, 0 1_Y, 0 \rangle &= \langle 1_Z, 0, 0 \rangle = \mathcal{T}(\Sigma X, Z). \end{aligned}$$

Definition 3.7. In the setup of Definition 3.1, the **restricted Toda brackets** are the subsets of the Toda bracket

$$\begin{aligned} \left\langle f_3, f_2, f_1 \right\rangle_{fc}^\alpha &\subseteq \langle f_3, f_2, f_1 \rangle_{fc} \\ \left\langle f_3, f_2, f_1 \right\rangle_{fc}^\beta &\subseteq \langle f_3, f_2, f_1 \rangle_{fc} \end{aligned}$$

consisting of all composites $\beta(\Sigma\alpha): \Sigma X_0 \rightarrow X_3$, where α and β appear in a commutative diagram (3.2) where the middle row is distinguished, with the prescribed map $\alpha: X_0 \rightarrow \Sigma^{-1}C_{f_2}$ (resp. $\beta: C_{f_2} \rightarrow X_3$).

The lift to the fiber $\alpha: X_0 \rightarrow \Sigma^{-1}C_{f_2}$ is a witness of the equality $f_2 f_1 = 0$. Dually, the extension to the cofiber $\beta: C_{f_2} \rightarrow X_3$ is a witness of the equality $f_3 f_2 = 0$.

Remark 3.8. Let $X_1 \xrightarrow{f_2} X_2 \xrightarrow{q_2} C_{f_2} \xrightarrow{\iota_2} \Sigma X_1$ be a distinguished triangle. By definition, we have equalities of subsets

$$\begin{aligned} \left\langle f_3, f_2, f_1 \right\rangle_{fc}^\alpha &= \left\langle f_3, f_2, -\Sigma^{-1}\iota_2 \right\rangle_{fc}^1 (\Sigma\alpha) \\ \left\langle f_3, f_2, f_1 \right\rangle_{fc}^\beta &= \beta \left\langle q_2, f_2, f_1 \right\rangle_{fc}^1. \end{aligned}$$

4. ADAMS d_2 IN TERMS OF 3-FOLD TODA BRACKETS

In this section, we show that the Adams differential d_r can be expressed in several ways using 3-fold Toda brackets. One of these expressions is as a secondary cohomology operation.

Given an injective class \mathcal{I} , an Adams resolution of an object Y as in Diagram (2.2), and an object X , consider a class $[x] \in E_2^{s,t}$ represented by a cycle $x \in E_1^{s,t} = \mathcal{T}(\Sigma^{t-s}X, I_s)$. Recall that $d_2[x] \in E_2^{s+2,t+1}$ is obtained as illustrated in the diagram

Explicitly, since x satisfies $d_1(x) = (\Sigma p_{s+1})\delta_s x = 0$, we can choose a lift $\tilde{x}: \Sigma^{t-s}X \rightarrow \Sigma Y_{s+2}$ of $\delta_s x$ to the fiber of Σp_{s+1} . Then the differential d_2 is given by

$$d_2[x] = [(\Sigma p_{s+2})\tilde{x}].$$

From now on, we will unroll the distinguished triangles and keep track of the suspensions. Following Convention 2.9, we will use the identifications

$$E_1^{s+2,t+1} = \mathcal{T}(\Sigma^{t-s-1}X, I_{s+2}) \cong \mathcal{T}(\Sigma^{t-s}X, \Sigma I_{s+2}) \cong \mathcal{T}(\Sigma^{t-s+1}X, \Sigma^2 I_{s+2}).$$

Proposition 4.1. *Denote by $d_2[x] \subseteq E_1^{s+2,t+1}$ the subset of all representatives of the class $d_2[x] \in E_2^{s+2,t+1}$. Then the following equalities hold:*

(1)

$$\begin{aligned} d_2[x] &= \left\langle \begin{array}{c} \Sigma^2 p_{s+2} \\ \Sigma d_1, \Sigma p_{s+1}, \delta_s x \end{array} \right\rangle_{fc} \\ &= \langle \Sigma d_1, \Sigma p_{s+1}, \delta_s x \rangle \end{aligned}$$

(2)

$$\begin{aligned} d_2[x] &= (\Sigma^2 p_{s+2}) \left\langle \Sigma \delta_{s+1}, \Sigma p_{s+1}, \delta_s x \right\rangle_{fc} \\ &= (\Sigma^2 p_{s+2}) \langle \Sigma \delta_{s+1}, \Sigma p_{s+1}, \delta_s x \rangle \end{aligned}$$

(3)

$$d_2[x] = \left\langle \Sigma d_1, d_1, x \right\rangle_{fc},$$

where β is the composite $C \xrightarrow{\tilde{\beta}} \Sigma^2 Y_{s+2} \xrightarrow{\Sigma^2 p_{s+2}} \Sigma^2 I_{s+2}$ and $\tilde{\beta}$ is obtained from the octahedral axiom applied to the factorization $d_1 = (\Sigma p_{s+1}) \delta_s: I_s \rightarrow \Sigma Y_{s+1} \rightarrow \Sigma I_{s+1}$.

In (3), β is a witness to the fact that the composite $(\Sigma d_1) d_1$ of primary operations is zero, and so the restricted Toda bracket is a secondary operation.

Proof. Note that t plays no role in the statement, so we will assume without loss of generality that $t = s$ holds.

(1) The first equality holds by definition of $d_2[x]$, namely choosing a lift of $\delta_s x$ to the fiber of Σp_{s+1} . The second equality follows from the fact that $\Sigma^2 p_{s+2}$ is the *unique* extension of $\Sigma d_1 = (\Sigma^2 p_{s+2})(\Sigma \delta_{s+1})$ to the cofiber of Σp_{s+1} . Indeed, $\Sigma \delta_{s+1}$ is \mathcal{I} -epic and ΣI_{s+2} is injective, so that the restriction map

$$(\Sigma \delta_{s+1})^*: \mathcal{T}(\Sigma^2 Y_{s+2}, \Sigma^2 I_{s+2}) \rightarrow \mathcal{T}(\Sigma I_{s+1}, \Sigma^2 I_{s+2})$$

is injective.

(2) The first equality holds by Remark 3.8. The second equality holds because $\Sigma \delta_{s+1}$ is \mathcal{I} -epic and ΣI_{s+2} is injective, as in part (1).

(3) The map $d_1: I_s \rightarrow \Sigma I_{s+1}$ is the composite $I_s \xrightarrow{\delta_s} \Sigma Y_{s+1} \xrightarrow{\Sigma p_{s+1}} \Sigma I_{s+1}$. The octahedral axiom applied to this factorization yields the dotted arrows in a commutative diagram

$$\begin{array}{ccccccc} I_s & \xrightarrow{\delta_s} & \Sigma Y_{s+1} & \xrightarrow{\Sigma i_s} & \Sigma Y_s & \xrightarrow{-\Sigma p_s} & \Sigma I_s \\ \parallel & & \downarrow \Sigma p_{s+1} & & \downarrow \tilde{\alpha} & & \parallel \\ I_s & \xrightarrow{d_1} & \Sigma I_{s+1} & \xrightarrow{q} & C_{d_1} & \xrightarrow{\iota} & \Sigma I_s \\ & & \downarrow \Sigma \delta_{s+1} & & \downarrow \tilde{\beta} & & \\ & & \Sigma^2 Y_{s+2} & \xlongequal{\quad} & \Sigma^2 Y_{s+2} & & \\ & & \downarrow -\Sigma^2 i_{s+1} & & \downarrow & & \\ & & \Sigma^2 Y_{s+1} & \xrightarrow{\Sigma^2 i_s} & \Sigma^2 Y_s & & \end{array}$$

where the rows and columns are distinguished and the equation $(-\Sigma^2 i_{s+1})\tilde{\beta} = (\Sigma\delta_s)\iota$ holds. The restricted bracket $\left\langle \Sigma d_1, d_1, x \right\rangle_{fc}^\beta$ consists of the maps $\Sigma X \rightarrow \Sigma^2 I_{s+2}$ appearing as downward composites in the commutative diagram

$$\begin{array}{ccccc}
& & & \Sigma X & \xrightarrow{-\Sigma x} & \Sigma I_s \\
& & & \downarrow \Sigma\alpha & & \parallel \\
I_s & \xrightarrow{d_1} & \Sigma I_{s+1} & \xrightarrow{q} & C_{d_1} & \xrightarrow{\iota} & \Sigma I_s \\
& & \parallel & & \downarrow \beta & & \parallel \\
& & \Sigma I_{s+1} & \xrightarrow{\Sigma d_1} & \Sigma^2 I_{s+2} & & \\
& & \nearrow \Sigma\delta_{s+1} & & \searrow \Sigma^2 p_{s+2} & & \\
& & \Sigma^2 Y_{s+2} & & & &
\end{array}$$

(\supseteq) Let $\beta(\Sigma\alpha) \in \left\langle d_1, d_1, x \right\rangle_{fc}^\beta$. By definition of β , we have $\beta(\Sigma\alpha) = (\Sigma^2 p_{s+2})\tilde{\beta}(\Sigma\alpha)$. Then $\tilde{\beta}(\Sigma\alpha): \Sigma X \rightarrow \Sigma^2 Y_{s+2}$ is a valid choice of the lift \tilde{x} in the definition of $d_2[x]$:

$$\begin{aligned}
(\Sigma^2 i_{s+1})\tilde{\beta}(\Sigma\alpha) &= -(\Sigma\delta_s)\iota(\Sigma\alpha) \\
&= -(\Sigma\delta_s)(-\Sigma x) \\
&= \Sigma(\delta_s x).
\end{aligned}$$

(\subseteq) Given a representative $(\Sigma p_{s+2})\tilde{x} \in d_2[x]$, let us show that $\Sigma\tilde{x}: \Sigma X \rightarrow \Sigma^2 Y_{s+2}$ factors as $\Sigma X \xrightarrow{\Sigma\alpha} C_{d_1} \xrightarrow{\tilde{\beta}} \Sigma^2 Y_{s+2}$ for some $\Sigma\alpha$, yielding a factorization of the desired form:

$$\begin{aligned}
(\Sigma^2 p_{s+2})(\Sigma\tilde{x}) &= (\Sigma^2 p_{s+2})\tilde{\beta}(\Sigma\alpha) \\
&= \beta(\Sigma\alpha).
\end{aligned}$$

By construction, the map $(\Sigma^2 i_s)(-\Sigma^2 i_{s+1}): \Sigma^2 Y_{s+2} \rightarrow \Sigma^2 Y_s$ is a cofiber of $\tilde{\beta}$. The condition

$$(\Sigma^2 i_s)(\Sigma^2 i_{s+1})(\Sigma\tilde{x}) = (\Sigma^2 i_s)\Sigma(\delta_s x) = 0$$

guarantees the existence of some lift $\Sigma\alpha: \Sigma X \rightarrow C_{d_1}$ of $\Sigma\tilde{x}$. The chosen lift $\Sigma\alpha$ might *not* satisfy $\iota(\Sigma\alpha) = -\Sigma x$, but we will correct it to a lift $\Sigma\alpha'$ which does. The two sides of the equation become equal after applying $-\Sigma\delta_s$, i.e., $(-\Sigma\delta_s)(-\Sigma x) = (-\Sigma\delta_s)\iota(\Sigma\alpha)$ holds. Hence, the error term factors as

$$-\Sigma x - \iota\Sigma\alpha = (-\Sigma p_s)(\Sigma\theta)$$

for some $\Sigma\theta: \Sigma X \rightarrow \Sigma Y_s$, since $-\Sigma p_s$ is a fiber of $-\Sigma\delta_s$. The corrected map $\Sigma\alpha' := \Sigma\alpha + \tilde{\alpha}(\Sigma\theta): \Sigma X \rightarrow C_{d_1}$ satisfies $\iota(\Sigma\alpha') = -\Sigma x$ and still satisfies $\tilde{\beta}(\Sigma\alpha') = \tilde{\beta}(\Sigma\alpha) = \Sigma\tilde{x}$, since the correction term $\tilde{\alpha}(\Sigma\theta)$ satisfies $\tilde{\beta}\tilde{\alpha}(\Sigma\theta) = 0$. \square

Proposition 4.2. *The following inclusions of subsets hold in $E_1^{s+2, t+1}$:*

$$d_2[x] \subseteq (\Sigma^2 p_{s+2}) \langle \Sigma\delta_{s+1}, d_1, x \rangle \subseteq \langle \Sigma d_1, d_1, x \rangle.$$

Proof. The first inclusion is

$$d_2[x] = (\Sigma^2 p_{s+2}) \langle \Sigma\delta_{s+1}, \Sigma p_{s+1}, \delta_s x \rangle \subseteq (\Sigma^2 p_{s+2}) \langle \Sigma\delta_{s+1}, (\Sigma p_{s+1})\delta_s, x \rangle,$$

whereas the second inclusion is

$$(\Sigma^2 p_{s+2}) \langle \Sigma \delta_{s+1}, d_1, x \rangle \subseteq \langle (\Sigma^2 p_{s+2})(\Sigma \delta_{s+1}), d_1, x \rangle,$$

both using Lemma 3.5. \square

Proposition 4.3. *The inclusion $(\Sigma^2 p_{s+2}) \langle \Sigma \delta_{s+1}, d_1, x \rangle \subseteq \langle \Sigma d_1, d_1, x \rangle$ need not be an equality in general.*

It was pointed out to us by Robert Bruner that this can happen in principle. We give an explicit example in Proposition A.1.

5. HIGHER TODA BRACKETS

We saw in Section 3 that there are several equivalent ways to define 3-fold Toda brackets. Following the approach given in [29], we show that the fiber-cofiber definition generalizes nicely to n -fold Toda brackets. There are $(n-2)!$ ways to make this generalization, and we prove that they are all the same up to a specified sign. We also show that this Toda bracket is self-dual.

Other sources that discuss higher Toda brackets in triangulated categories are [40, Appendix A], [18, IV §2] and [36, §4], which all give definitions that follow Cohen's approach for spectra or spaces [15]. We show that our definition agrees with those of [40] and [36]. (We believe that it sometimes differs in sign from [15]. We have not compared carefully with [18].)

Definition 5.1. Let $X_0 \xrightarrow{f_1} X_1 \xrightarrow{f_2} X_2 \xrightarrow{f_3} X_3$ be a diagram in a triangulated category \mathcal{T} . We define the **Toda family** of this sequence to be the collection $T(f_3, f_2, f_1)$ consisting of all pairs $(\beta, \Sigma\alpha)$, where α and β appear in a commutative diagram

$$\begin{array}{ccccccc} X_0 & \xrightarrow{f_1} & X_1 & & & & \\ \alpha \downarrow & & \parallel & & & & \\ \Sigma^{-1}C_{f_2} & \longrightarrow & X_1 & \xrightarrow{f_2} & X_2 & \longrightarrow & C_{f_2} \\ & & & & \parallel & & \downarrow \beta \\ & & & & X_2 & \xrightarrow{f_3} & X_3 \end{array}$$

with distinguished middle row. Equivalently,

$$\begin{array}{ccccccc} & & & & \Sigma X_0 & \xrightarrow{-\Sigma f_1} & \Sigma X_1 \\ & & & & \Sigma \alpha \downarrow & & \parallel \\ X_1 & \xrightarrow{f_2} & X_2 & \longrightarrow & C_{f_2} & \longrightarrow & \Sigma X_1 \\ & & \parallel & & \downarrow \beta & & \\ & & X_2 & \xrightarrow{f_3} & X_3, & & \end{array}$$

where the middle row is again distinguished. (The negative of Σf_1 appears, since when a triangle is rotated, a sign is introduced.) Note that the maps in each pair form a composable sequence $\Sigma X_0 \xrightarrow{\Sigma\alpha} C_{f_2} \xrightarrow{\beta} X_3$, with varying intermediate object, and that the collection of composites $\beta \circ \Sigma\alpha$ is exactly the Toda bracket $\langle f_3, f_2, f_1 \rangle$, using the fiber-cofiber definition (see Diagram (3.2)). (Also note that the Toda family is generally a proper class, but this is

only because the intermediate object can be varied up to isomorphism, and so we will ignore this.)

More generally, if S is a set of composable triples of maps, starting at X_0 and ending at X_3 , we define $T(S)$ to be the union of $T(f_3, f_2, f_1)$ for each triple (f_3, f_2, f_1) in S .

Definition 5.2. Let $X_0 \xrightarrow{f_1} X_1 \xrightarrow{f_2} X_2 \xrightarrow{f_3} \dots \xrightarrow{f_n} X_n$ be a diagram in a triangulated category \mathcal{T} . We define the **Toda bracket** $\langle f_n, \dots, f_1 \rangle$ inductively as follows. If $n = 2$, it is the set consisting of just the composite $f_2 f_1$. If $n > 2$, it is the union of the sets $\langle \beta, \Sigma\alpha, \Sigma f_{n-3}, \dots, \Sigma f_1 \rangle$, where $(\beta, \Sigma\alpha)$ is in $T(f_n, f_{n-1}, f_{n-2})$.

In fact, there are $(n - 2)!$ such definitions, depending on a sequence of choices of which triple of consecutive maps to apply the Toda family construction to. In Theorem 5.11 we will enumerate these choices and show that they all agree up to sign.

Example 5.3. Let us describe 4-fold Toda brackets in more detail. We have

$$\langle f_4, f_3, f_2, f_1 \rangle = \bigcup_{\beta, \alpha} \langle \beta, \Sigma\alpha, \Sigma f_1 \rangle = \bigcup_{\beta, \alpha} \bigcup_{\beta', \alpha'} \{ \beta' \circ \Sigma\alpha' \}$$

with $(\beta, \Sigma\alpha) \in T(f_4, f_3, f_2)$ and $(\beta', \Sigma\alpha') \in T(\beta, \Sigma\alpha, \Sigma f_1)$. These maps fit into a commutative diagram

$$\begin{array}{ccccc} \Sigma^2 X_0 & \xrightarrow{\Sigma\alpha'} & C_{\Sigma\alpha} & \longrightarrow & \Sigma^2 X_1 & \text{row} = -\Sigma^2 f_1 \\ & & \uparrow & \searrow^{\beta'} & & \\ \Sigma X_1 & \xrightarrow{\Sigma\alpha} & C_{f_3} & \longrightarrow & \Sigma X_2 & \text{row} = -\Sigma f_2 \\ & & \uparrow & \searrow^{\beta} & & \\ X_2 & \xrightarrow{f_3} & X_3 & \xrightarrow{f_4} & X_4 & \\ & & \uparrow & & & \\ & & 0 & & & \end{array}$$

where the horizontal composites are specified as above, and each “snake”

$$\begin{array}{ccc} & \longrightarrow & \cdot \\ & \uparrow & \\ \cdot & \longrightarrow & \cdot \end{array}$$

is a distinguished triangle. The middle column is an example of a *3-filtered object* as defined below.

Next, we will show that Definition 5.2 coincides with the definitions of higher Toda brackets in [40, Appendix A] and [36, §4], which we recall here.

Definition 5.4. Let $n \geq 1$ and consider a diagram in \mathcal{T}

$$Y_0 \xrightarrow{\lambda_1} Y_1 \xrightarrow{\lambda_2} Y_2 \longrightarrow \dots \xrightarrow{\lambda_{n-1}} Y_{n-1}$$

consisting of $n - 1$ composable maps. An *n -filtered object* Y based on $(\lambda_{n-1}, \dots, \lambda_1)$ consists of a sequence of maps

$$0 = F_0 Y \xrightarrow{i_0} F_1 Y \xrightarrow{i_1} \dots \xrightarrow{i_{n-1}} F_n Y = Y$$

together with distinguished triangles

$$F_j Y \xrightarrow{i_j} F_{j+1} Y \xrightarrow{q_{j+1}} \Sigma^j Y_{n-1-j} \xrightarrow{e_j} \Sigma F_j Y$$

for $0 \leq j \leq n-1$, such that for $1 \leq j \leq n-1$, the composite

$$\Sigma^j Y_{n-1-j} \xrightarrow{e_j} \Sigma F_j Y \xrightarrow{\Sigma q_j} \Sigma^j Y_{n-j}$$

is equal to $\Sigma^j \lambda_{n-j}$. In particular, the n -filtered object Y comes equipped with maps

$$\sigma'_Y: Y_{n-1} \cong F_1 Y \rightarrow Y$$

$$\sigma_Y: Y = F_n Y \rightarrow \Sigma^{n-1} Y_0.$$

Definition 5.5. Let $X_0 \xrightarrow{f_1} X_1 \xrightarrow{f_2} X_2 \xrightarrow{f_3} \dots \xrightarrow{f_n} X_n$ be a diagram in a triangulated category \mathcal{T} . The **Toda bracket** in the sense of Shipley–Sagave $\langle f_n, \dots, f_1 \rangle_{SS} \subseteq \mathcal{T}(\Sigma^{n-2} X_0, X_n)$ is the set of all composites appearing in the middle row of a commutative diagram

$$\begin{array}{ccccc} & & X_{n-1} & & \\ & & \downarrow \sigma'_X & \searrow f_n & \\ \Sigma^{n-2} X_0 & \dashrightarrow & X & \dashrightarrow & X_n \\ & \searrow \Sigma^{n-2} f_1 & \downarrow \sigma_X & & \\ & & \Sigma^{n-2} X_1 & & \end{array}$$

where X is an $(n-1)$ -filtered object based on $(f_{n-1}, \dots, f_3, f_2)$.

Example 5.6. For a 3-fold Toda bracket $\langle f_3, f_2, f_1 \rangle_{SS}$, a 2-filtered object X based on f_2 amounts to a cofiber of $-f_2$, more precisely, a distinguished triangle

$$X_2 \xrightarrow{\sigma'_X} X \xrightarrow{\sigma_X} \Sigma X_1 \xrightarrow{\Sigma f_2} \Sigma X_2.$$

Using this, one readily checks the equality $\langle f_3, f_2, f_1 \rangle_{SS} = \langle f_3, f_2, f_1 \rangle_{fc}$, as noted in [36, Definition 4.5].

Example 5.7. For a 4-fold Toda bracket $\langle f_4, f_3, f_2, f_1 \rangle_{SS}$, a 3-filtered object X based on (f_3, f_2) consists of the data displayed in the diagram

$$\begin{array}{ccccc} & & F_3 X = X \xrightarrow{q_3 = \sigma_X} \Sigma^2 X_1 & & \\ & & \uparrow i_2 & & \\ \Sigma X_1 & \xrightarrow{-\Sigma^{-1} e_2} & F_2 X & \xrightarrow{q_2} & \Sigma X_2 & \text{row} = -\Sigma f_2 \\ & & \uparrow i_1 & & \\ X_2 & \xrightarrow{-\Sigma^{-1} e_1} & F_1 X & \xrightarrow[q_1]{\cong} & X_3 & \text{row} = -f_3 \\ & & \uparrow i_0 & & \\ & & F_0 X = 0, & & \end{array}$$

where the two snakes are distinguished. The bracket consists of the maps $\Sigma^2 X_0 \rightarrow X_4$ appearing as composites of the dotted arrows in a commutative diagram

$$\begin{array}{ccccc}
 \Sigma^2 X_0 & \dashrightarrow & X & \xrightarrow{\sigma_X} & \Sigma^2 X_1 & \text{row} = \Sigma^2 f_1 \\
 & & \uparrow & \dashrightarrow & & \\
 \Sigma X_1 & \xrightarrow{-\Sigma^{-1}e_2} & F_2 X & \xrightarrow{q_2} & \Sigma X_2 & \text{row} = -\Sigma f_2 \\
 & & \uparrow & \dashrightarrow & \downarrow & \\
 X_2 & \xrightarrow{-f_3} & X_3 & \xrightarrow{f_4} & X_4 & \\
 & & \uparrow & & & \\
 & & 0, & & &
 \end{array}$$

where the two snakes are distinguished. By negating the first and third map in each snake, this recovers the description in Example 5.3, thus proving the equality of subsets

$$\langle f_4, f_3, f_2, f_1 \rangle_{SS} = \langle f_4, f_3, f_2, f_1 \rangle.$$

Proposition 5.8. *Definitions 5.2 and 5.5 agree. In other words, we have the equality*

$$\langle f_n, \dots, f_1 \rangle_{SS} = \langle f_n, \dots, f_1 \rangle$$

of subsets of $\mathcal{T}(\Sigma^{n-2} X_0, X_n)$.

Proof. This is a straightforward generalization of Example 5.7. □

We define the **negative** of a Toda family $T(f_3, f_2, f_1)$ to consist of pairs $(\beta, -\Sigma\alpha)$ for $(\beta, \Sigma\alpha) \in T(f_3, f_2, f_1)$. (Since changing the sign of two maps in a triangle doesn't affect whether it is distinguished, it would be equivalent to put the minus sign with the β .)

Lemma 5.9. *Let $X_0 \xrightarrow{f_1} X_1 \xrightarrow{f_2} X_2 \xrightarrow{f_3} X_3 \xrightarrow{f_4} X_4$ be a diagram in a triangulated category \mathcal{T} . Then the two sets of pairs $T(T(f_4, f_3, f_2), \Sigma f_1)$ and $T(f_4, T(f_3, f_2, f_1))$ are negatives of each other.*

This is stronger than saying that the two ways of computing the Toda bracket $\langle f_4, f_3, f_2, f_1 \rangle$ are negatives, and the stronger statement will be used inductively to prove Theorem 5.11.

Proof. We will show that the negative of $T(T(f_4, f_3, f_2), \Sigma f_1)$ is contained in the family $T(f_4, T(f_3, f_2, f_1))$. The reverse inclusion is proved dually.

Suppose $(\beta, \Sigma\alpha)$ is in $T(T(f_4, f_3, f_2), \Sigma f_1)$, that is, $(\beta, \Sigma\alpha)$ is in $T(\beta', \Sigma\alpha', \Sigma f_1)$ for some $(\beta', \Sigma\alpha')$ in $T(f_4, f_3, f_2)$. This means that we have the following commutative diagram

$$\begin{array}{ccccc}
 & & \Sigma X_1 & \xrightarrow{-\Sigma f_2} & \Sigma X_2 \\
 & & \vdots & & \parallel \\
 & & \Sigma\alpha' & & \\
 & & \downarrow & & \\
 X_2 & \xrightarrow{f_3} & X_3 & \longrightarrow & C_{f_3} & \longrightarrow & \Sigma X_2 \\
 & & \searrow f_4 & & \swarrow \beta' & & \\
 & & & & X_4 & & \\
 & & & & \swarrow \beta & & \\
 & & & & C_{\Sigma\alpha'} & & \\
 & & \swarrow \Sigma\alpha & & \downarrow & & \\
 & & \Sigma^2 X_0 & & \downarrow & & \\
 & & \swarrow -\Sigma^2 f_1 & & \Sigma^2 X_1 & &
 \end{array}$$

in which the long row and column are distinguished triangles.

Using the octahedral axiom, there exists a map $\delta : C_{f_2} \rightarrow X_3$ in the following diagram making the two squares commute and such that the diagram can be extended as shown, with all rows and columns distinguished:

$$\begin{array}{ccccccc}
 & & & & \Sigma X_0 & & \\
 & & & & \swarrow \gamma & \searrow -\Sigma f_1 & \\
 X_2 & \longrightarrow & C_{f_2} & \longrightarrow & \Sigma X_1 & \xrightarrow{-\Sigma f_2} & \Sigma X_2 \\
 \parallel & & \vdots & & \vdots & & \parallel \\
 X_2 & \xrightarrow{f_3} & X_3 & \longrightarrow & C_{f_3} & \longrightarrow & \Sigma X_2 \\
 & & \searrow f_4 & & \swarrow \beta' & & \\
 & & & & X_4 & & \\
 & & & & \swarrow \beta & & \\
 & & & & C_{\Sigma\alpha'} & & \\
 & & \swarrow \Sigma\alpha & & \downarrow & & \\
 & & \Sigma^2 X_0 & & \downarrow & & \\
 & & \swarrow -\Sigma^2 f_1 & & \Sigma^2 X_1 & & \\
 & & \downarrow \Sigma\gamma & & \downarrow & & \\
 & & \Sigma C_{f_2} & \longrightarrow & \Sigma^2 X_1 & &
 \end{array}$$

Define $\Sigma\gamma$ to be the composite $\Sigma^2 X_0 \rightarrow C_{\Sigma\alpha'} = C_\delta \rightarrow \Sigma C_{f_2}$, where the first map is $\Sigma\alpha$. Then the small triangles at the top and bottom of the last diagram commute as well. Therefore, (δ, γ) is in $T(f_3, f_2, f_1)$. Moreover, this diagram shows that $(\beta, -\Sigma\alpha)$ is in $T(f_4, \delta, \gamma)$, completing the argument. \square

To concisely describe different ways of computing higher Toda brackets, we introduce the following notation. For $0 \leq j \leq n - 3$, write $T_j(f_n, f_{n-1}, \dots, f_1)$ for the set of tuples

$$\{(f_n, f_{n-1}, \dots, f_{n-j+1}, \beta, \Sigma\alpha, \Sigma f_{n-j-3}, \dots, \Sigma f_1)\},$$

where $(\beta, \Sigma\alpha)$ is in $T(f_{n-j}, f_{n-j-1}, f_{n-j-2})$. (There are j maps to the left of the three used for the Toda family.) If S is a set of n -tuples of composable maps, we define $T_j(S)$ to be the union of the sets $T_j(f_n, f_{n-1}, \dots, f_1)$ for $(f_n, f_{n-1}, \dots, f_1)$ in S . With this notation, the standard Toda bracket $\langle f_n, \dots, f_1 \rangle$ consists of the composites of all the pairs occurring in the iterated Toda family

$$T(f_n, \dots, f_1) := T_0(T_0(T_0(\dots T_0(f_n, \dots, f_1) \dots))).$$

A general Toda bracket can be written in the form $T_{j_1}(T_{j_2}(T_{j_3}(\dots T_{j_{n-2}}(f_n, \dots, f_1) \dots)))$, where j_1, j_2, \dots, j_{n-2} is a sequence of natural numbers with $0 \leq j_i < i$ for each i . There are $(n - 2)!$ such sequences.

Remark 5.10. There are six ways to compute the five-fold Toda bracket $\langle f_5, f_4, f_3, f_2, f_1 \rangle$, as the set of composites of the pairs of maps in one of the following sets:

$$\begin{aligned} T_0(T_0(T_0(f_5, f_4, f_3, f_2, f_1))) &= T(T(T(f_5, f_4, f_3), \Sigma f_2), \Sigma^2 f_1), \\ T_0(T_0(T_1(f_5, f_4, f_3, f_2, f_1))) &= T(T(f_5, T(f_4, f_3, f_2)), \Sigma^2 f_1), \\ T_0(T_1(T_1(f_5, f_4, f_3, f_2, f_1))) &= T(f_5, T(T(f_4, f_3, f_2), \Sigma f_1)), \\ T_0(T_1(T_2(f_5, f_4, f_3, f_2, f_1))) &= T(f_5, T(f_4, T(f_3, f_2, f_1))), \\ T_0(T_0(T_2(f_5, f_4, f_3, f_2, f_1))) &\text{, and} \\ T_0(T_1(T_0(f_5, f_4, f_3, f_2, f_1))) & . \end{aligned}$$

The last two cannot be expressed directly just using T .

Now we can prove the main result of this section.

Theorem 5.11. *The Toda bracket computed using the sequence j_1, j_2, \dots, j_{n-2} equals the standard Toda bracket up to the sign $(-1)^{\sum j_i}$.*

Proof. Let j_1, j_2, \dots, j_{n-2} be a sequence with $0 \leq j_i < i$ for each i . Lemma 5.9 tells us that if we replace consecutive entries $k, k + 1$ with k, k in any such sequence, the two Toda brackets agree up to a sign. To begin with, we ignore the signs. We will prove by induction on ℓ that the initial portion j_1, \dots, j_ℓ of such a sequence can be converted into any other sequence, using just the move allowed by Lemma 5.9 and its inverse, and without changing j_i for $i > \ell$. For $\ell = 1$, there is only one sequence 0. For $\ell = 2$, there are two sequences, 0, 0 and 0, 1, and Lemma 5.9 applies. For $\ell > 2$, suppose our goal is to produce the sequence j'_1, \dots, j'_ℓ . We break the argument into three cases:

Case 1: $j'_\ell = j_\ell$. We can directly use the induction hypothesis to adjust the entries in the first $\ell - 1$ positions.

Case 2: $j'_\ell > j_\ell$. By induction, we can change the first $\ell - 1$ entries in the sequence j so that the entry in position $\ell - 1$ is j_ℓ , since $j_\ell < j'_\ell \leq \ell - 1$. Then, using Lemma 5.9, we can change the entry in position ℓ to $j_\ell + 1$. Continuing in this way, we get j'_ℓ in position ℓ , and then we are in Case 1.

Case 3: $j'_\ell < j_\ell$. Since the moves are reversible, this is equivalent to Case 2.

To handle the sign, first note that signs propagate through the Toda family construction. More precisely, suppose S is a set of n -tuples of maps, and let S' be a set obtained by negating

the k^{th} map in each n -tuple for some fixed k . Then $T_j(S)$ has the same relationship to $T_j(S')$, possibly for a different value of k .

As a result, applying the move of Lemma 5.9 changes the resulting Toda bracket by a sign. That move also changes the parity of $\sum_i j_i$. Since we get a plus sign when each j_i is zero, it follows that the difference in sign in general is $(-1)^{\sum_i j_i}$. \square

An animation of this argument is available at [14]. It was pointed out by Dylan Wilson that the combinatorial part of the above proof is equivalent to the well-known fact that if a binary operation is associative on triples, then it is associative on n -tuples.

In order to compare our Toda brackets to the Toda brackets in the opposite category, we need one lemma.

Lemma 5.12. *Let $X_0 \xrightarrow{f_1} X_1 \xrightarrow{f_2} X_2 \xrightarrow{f_3} X_3$ be a diagram in a triangulated category \mathcal{T} . Then the Toda family $T(\Sigma f_3, \Sigma f_2, \Sigma f_1)$ is the negative of the suspension of $T(f_3, f_2, f_1)$. That is, it consists of $(\Sigma\beta, -\Sigma^2\alpha)$ for $(\beta, \Sigma\alpha)$ in $T(f_3, f_2, f_1)$.*

Proof. Given a distinguished triangle $\Sigma^{-1}C_{f_2} \xrightarrow{k} X_1 \xrightarrow{f_2} X_2 \xrightarrow{\iota} C_{f_2}$, a distinguished triangle involving Σf_2 is

$$C_{f_2} \xrightarrow{-\Sigma k} \Sigma X_1 \xrightarrow{\Sigma f_2} \Sigma X_2 \xrightarrow{\Sigma \iota} \Sigma C_{f_2}.$$

Because of the minus sign at the left, the maps that arise in the Toda family based on this triangle are $-\Sigma^2\alpha$ and $\Sigma\beta$, where $\Sigma\alpha$ and β arise in the Toda family based on the starting triangle. \square

Given a triangulated category \mathcal{T} , the opposite category \mathcal{T}^{op} is triangulated in a natural way. The suspension in \mathcal{T}^{op} is Σ^{-1} and a triangle

$$Y_0 \xrightarrow{g_1} Y_1 \xrightarrow{g_2} Y_2 \xrightarrow{g_3} \Sigma^{-1}Y_0$$

in \mathcal{T}^{op} is distinguished if and only if the triangle

$$\Sigma\Sigma^{-1}Y_0 \xleftarrow{g'_1} Y_1 \xleftarrow{g_2} Y_2 \xleftarrow{g_3} \Sigma^{-1}Y_0$$

in \mathcal{T} is distinguished, where g'_1 is the composite of g_1 with the natural isomorphism $Y_0 \cong \Sigma\Sigma^{-1}Y_0$.

Corollary 5.13. *The Toda bracket is self-dual up to suspension. More precisely, let $X_0 \xrightarrow{f_1} X_1 \xrightarrow{f_2} X_2 \xrightarrow{f_3} \dots \xrightarrow{f_n} X_n$ be a diagram in a triangulated category \mathcal{T} . Then the subset*

$$\langle f_1, \dots, f_n \rangle^{\mathcal{T}^{\text{op}}} \subseteq \mathcal{T}^{\text{op}}(\Sigma^{-(n-2)}X_n, X_0) = \mathcal{T}(X_0, \Sigma^{-(n-2)}X_n)$$

defined by taking the Toda bracket in \mathcal{T}^{op} is sent to the subset

$$\langle f_n, \dots, f_1 \rangle^{\mathcal{T}} \subseteq \mathcal{T}(\Sigma^{n-2}X_0, X_n)$$

defined by taking the Toda bracket in \mathcal{T} under the bijection $\Sigma^{n-2} : \mathcal{T}(X_0, \Sigma^{-(n-2)}X_n) \rightarrow \mathcal{T}(\Sigma^{n-2}X_0, X_n)$.

Proof. First we compare Toda families in \mathcal{T} and \mathcal{T}^{op} . It is easy to see that the Toda family $T^{\mathcal{T}^{\text{op}}}(f_1, f_2, f_3)$ computed in \mathcal{T}^{op} consists of the pairs $(\alpha, \Sigma^{-1}\beta)$ for $(\Sigma\alpha, \beta)$ in the Toda family $T^{\mathcal{T}}(f_3, f_2, f_1)$ computed in \mathcal{T} . In short, one has to desuspend and transpose the pairs.

Using this, one can see that the iterated Toda family

$$T^{\mathcal{T}^{\text{op}}}(T^{\mathcal{T}^{\text{op}}}\dots T^{\mathcal{T}^{\text{op}}}(f_1, f_2, f_3), \dots, \Sigma^{-(n-3)}f_n)$$

is equal to the transpose of

$$\Sigma^{-1}T^\mathcal{T}(\Sigma^{-(n-3)}f_n, \Sigma^{-1}T^\mathcal{T}(\Sigma^{-(n-4)}f_{n-1}, \Sigma^{-1}T^\mathcal{T} \cdots \Sigma^{-1}T^\mathcal{T}(f_3, f_2, f_1) \cdots))$$

By Lemma 5.12, the desuspensions pass through all of the Toda family constructions, introducing an overall sign of $(-1)^{1+2+3+\cdots+(n-3)}$, and producing

$$\Sigma^{-(n-2)}T^\mathcal{T}(f_n, T^\mathcal{T}(f_{n-1}, T^\mathcal{T} \cdots T^\mathcal{T}(f_3, f_2, f_1) \cdots))$$

By Theorem 5.11, composing the pairs gives the usual Toda bracket up to the sign $(-1)^{0+1+2+\cdots+(n-3)}$. The two signs cancel, yielding the result. \square

We do not know a direct proof of this corollary. To summarize, our insight is that by generalizing the corollary to all $(n-2)!$ methods of computing the Toda bracket, we were able to reduce the argument to the 4-fold case (Lemma 5.9) and some combinatorics.

Remark 5.14. As with the 3-fold Toda brackets (see Remark 3.2), the higher Toda brackets depend on the triangulation. If the triangulation is negated, the n -fold Toda brackets change by the sign $(-1)^n$.

6. HIGHER ORDER OPERATIONS DETERMINE d_r

In this section, we show that the higher Adams differentials can be expressed in terms of higher Toda brackets, in two ways. One of these expressions is as an r^{th} order cohomology operation.

Given an injective class \mathcal{I} , an Adams resolution of an object Y as in Diagram (2.2), and an object X , consider a class $[x] \in E_r^{s,t}$ represented by an element $x \in E_1^{s,t} = \mathcal{T}(\Sigma^{t-s}X, I_s)$. The class $d_r[x]$ is the set of all $(\Sigma p_{s+r})\tilde{x}$, where \tilde{x} runs over lifts of $\delta_s x$ through the $(r-1)$ -fold composite $\Sigma(i_{s+1} \cdots i_{s+r-1})$ which appears across the top edge of the Adams resolution.

Our first result will be a generalization of Proposition 4.1(1), expressing d_r in terms of an $(r+1)$ -fold Toda bracket.

Theorem 6.1. *As subsets of $E_1^{s+r, t+r-1}$, we have*

$$d_r[x] = \langle \Sigma^{r-1}d_1, \dots, \Sigma^2d_1, \Sigma d_1, \Sigma p_{s+1}, \delta_s x \rangle.$$

Proof. We compute the Toda bracket, applying the Toda family construction starting from the right, which introduces a sign of $(-1)^{1+2+\cdots+(r-2)}$, by Theorem 5.11. We begin with the Toda family $\mathbb{T}(\Sigma d_1, \Sigma p_{s+1}, \delta_s x)$. There is a distinguished triangle

$$\Sigma Y_{s+2} \xrightarrow{\Sigma i_{s+1}} \Sigma Y_{s+1} \xrightarrow{\Sigma p_{s+1}} \Sigma I_{s+1} \xrightarrow{\Sigma \delta_{s+1}} \Sigma^2 Y_{s+2},$$

with no needed signs. The map Σd_1 factors through $\Sigma \delta_{s+1}$ as $\Sigma^2 p_{s+2}$, and this factorization is unique because $\Sigma \delta_{s+1}$ is \mathcal{I} -epic and $\Sigma^2 I_{s+2}$ is injective. The other maps in the Toda family are Σx_1 , where x_1 is a lift of $\delta_s x$ through Σi_{s+1} . So

$$\mathbb{T}(\Sigma d_1, \Sigma p_{s+1}, \delta_s x) = \{(\Sigma^2 p_{s+2}, \Sigma x_1) \mid x_1 \text{ a lift of } \delta_s x \text{ through } \Sigma i_{s+1}\}.$$

(The Toda family also includes $(\Sigma^2 p_{s+2} \phi, \phi^{-1}(\Sigma x_1))$, where ϕ is any isomorphism, but these contribute nothing additional to the later computations.) The composites of such pairs give $d_2[x]$, up to suspension, recovering Proposition 4.1(1).

Continuing, for each such pair we compute

$$\begin{aligned} \mathbb{T}(\Sigma^2 d_1, \Sigma^2 p_{s+2}, \Sigma x_1) &= -\Sigma \mathbb{T}(\Sigma d_1, \Sigma p_{s+2}, x_1) \\ &= -\Sigma \{(\Sigma^2 p_{s+3}, \Sigma x_2) \mid x_2 \text{ a lift of } x_1 \text{ through } \Sigma i_{s+2}\}. \end{aligned}$$

The first equality is Lemma 5.12, and the second reuses the work done in the previous paragraph, with s increased by 1. Composing these pairs gives $-d_3[x]$. The sign which is needed to produce the standard Toda bracket is $(-1)^1$, and so the signs cancel.

At the next step, we compute

$$\begin{aligned} \mathrm{T}(\Sigma^3 d_1, \Sigma^3 p_{s+3}, -\Sigma^2 x_2) &= -\Sigma^2 \mathrm{T}(\Sigma d_1, \Sigma p_{s+3}, x_2) \\ &= -\Sigma^2 \{(\Sigma^2 p_{s+4}, \Sigma x_3) \mid x_3 \text{ a lift of } x_2 \text{ through } \Sigma i_{s+3}\}. \end{aligned}$$

Again, the composites give $-d_4[x]$. Since it was a double suspension that passed through the Toda family, no additional sign was introduced. Similarly, the sign to convert to the standard Toda bracket is $(-1)^{1+2}$, and since 2 is even, no additional sign was introduced. Therefore, the signs still cancel.

The pattern continues. In total, there are $1+2+\cdots+(r-2)$ suspensions that pass through the Toda family, and the sign to convert to the standard Toda bracket is also based on that number, so the signs cancel. \square

Remark 6.2. Theorem 6.1 can also be proved using the definition Toda brackets based on r -filtered objects, as in Definitions 5.4 and 5.5. However, one must work in the opposite category $\mathcal{T}^{\mathrm{op}}$. In that category, there is a unique r -filtered object, up to isomorphism, based on the maps in the Toda bracket. One of the dashed arrows in the diagram from Definition 5.5 is also unique, and the other corresponds naturally to the choice of lift in the Adams differential.

In the remainder of this section, we describe the analog of Proposition 4.1(3). We begin by defining restricted higher Toda brackets, in terms of restricted Toda families.

Consider a Toda family $\mathrm{T}(gh_1, g_1 h_0, g_0 h)$, where the maps factor as shown, there are distinguished triangles

$$Z_i \xrightarrow{g_i} J_i \xrightarrow{h_i} Z_{i+1} \xrightarrow{k_i} \Sigma Z_i \quad (6.1)$$

for $i = 0, 1$, and g and h are arbitrary maps $Z_2 \rightarrow A$ and $B \rightarrow Z_0$, respectively. This information determines an essentially unique element of the Toda family in the following way. The octahedral axiom applied to the factorization $g_1 h_0$ yields the dotted arrows in a commutative diagram

$$\begin{array}{ccccccc} J_0 & \xrightarrow{h_0} & Z_1 & \xrightarrow{k_0} & \Sigma Z_0 & \xrightarrow{-\Sigma g_0} & \Sigma J_0 \\ \parallel & & \downarrow g_1 & & \downarrow \alpha_2 & & \parallel \\ J_0 & \xrightarrow{g_1 h_0} & J_1 & \dashrightarrow^q & W_2 & \dashrightarrow^\iota & \Sigma J_0 \\ & & \downarrow h_1 & & \downarrow \beta_2 & & \\ & & Z_2 & \xlongequal{\quad} & Z_2 & & \\ & & \downarrow k_1 & & \downarrow \gamma_2 & & \\ & & \Sigma Z_1 & \xrightarrow{\Sigma k_0} & \Sigma^2 Z_0, & & \end{array}$$

where the rows and columns are distinguished and $\gamma_2 := (\Sigma k_0)k_1$. It is easy to see that $-\Sigma(g_0 h)$ lifts through ι as $\alpha_2(\Sigma h)$, and that gh_1 extends over q as $g\beta_2$. We define the **restricted Toda family** to be the set $\mathrm{T}(gh_1^!, g_1 h_0^!, g_0 h)$ consisting of the pairs $(g\beta_2, \alpha_2(\Sigma h))$ that arise in this way. Since α_2 and β_2 come from a distinguished triangle involving a fixed map γ_2 , such pairs are unique up to the usual ambiguity of replacing the pair with $(g\beta_2 \phi, \phi^{-1} \alpha_2(\Sigma h))$, where ϕ is an isomorphism. Similarly, given any map $x : B \rightarrow J_0$, we

define $T(gh_1!g_1h_0, x)$ to be the set consisting of the pairs $(g\beta_2, \Sigma\alpha)$, where β_2 arises as above and $\Sigma\alpha$ is any lift of $-\Sigma x$ through ι .

Definition 6.3. Given distinguished triangles as in Equation (6.1), for $i = 1, \dots, n-1$, and maps $g : Z_n \rightarrow A$ and $x : B \rightarrow J_1$, we define the **restricted Toda bracket**

$$\left\langle gh_{n-1}!, g_{n-1}h_{n-2}!, \dots!, g_3h_2!, g_2h_1, x \right\rangle$$

inductively as follows: If $n = 2$, it is the set consisting of just the composite gh_1x . If $n = 3$, it is the set of composites of the pairs in $T(gh_2!, g_2h_1, x)$. If $n > 3$, it is the union of the sets

$$\left\langle g\beta_2!, \alpha_2(\Sigma h_{n-3})!, \Sigma(g_{n-3}h_{n-4})!, \dots, \Sigma x \right\rangle,$$

where $(g\beta_2, \alpha_2(\Sigma h_{n-3}))$ is in $T(gh_{n-1}!, g_{n-1}h_{n-2}!, g_{n-2}h_{n-3})$.

Remark 6.4. Despite the notation, we want to make it clear that these restricted Toda families and restricted Toda brackets depend on the choice of factorizations and on the distinguished triangles in Equation (6.1). Moreover, the elements of the restricted Toda families are not simply pairs, but also include the factorizations of the maps in those pairs, and the distinguished triangle involving α_2 and β_2 . This information is used in the $(n-1)$ -fold restricted Toda bracket that is used to define the n -fold restricted Toda bracket.

Recall that the maps d_1 are defined to be $(\Sigma p_{s+1})\delta_s$, and that we have distinguished triangles

$$Y_s \xrightarrow{p_s} I_s \xrightarrow{\delta_s} \Sigma Y_{s+1} \xrightarrow{\Sigma i_s} \Sigma Y_s$$

for each s . The same holds for suspensions of d_1 , with the last map changing sign each time it is suspended. Thus for $x : \Sigma^{t-s}X \rightarrow I_s$ in the E_1 term, the $(r+1)$ -fold restricted Toda bracket $\left\langle \Sigma^{r-1}d_1!, \dots!, \Sigma d_1!, d_1, x \right\rangle$ makes sense for each r , where we are implicitly using the defining factorizations and the triangles from the Adams resolution.

Theorem 6.5. *As subsets of $E_1^{s+r, t+r-1}$, we have*

$$d_r[x] = \left\langle \Sigma^{r-1}d_1!, \dots!, \Sigma d_1!, d_1, x \right\rangle.$$

This is a generalization of Proposition 4.1(3). The data in the Adams resolution is the witness that the composites of the primary operations are zero in a sufficiently coherent way to permit an r^{th} order cohomology operation to be defined.

Proof. The restricted Toda bracket $\left\langle \Sigma^{r-1}d_1!, \dots!, \Sigma d_1!, d_1, x \right\rangle$ is defined recursively, working from the left. Each of the $r-2$ doubly restricted Toda families has essentially one element. The first one involves maps α_2, β_2 and γ_2 that form a distinguished triangle, and γ_2 is equal to $[(-1)^r \Sigma^r i_{s+r-2}] [-(-1)^r \Sigma^r i_{s+r-1}]$. We will denote the corresponding maps in the following octahedra α_k, β_k and γ_k , where each γ_k equals $[(-1)^r \Sigma^r i_{s+r-k}] \gamma_{k-1}$, and so $\gamma_k = -(-1)^{rk} \Sigma^r (i_{s+r-k} \cdots i_{s+r-1})$. One is left to compute the singly restricted Toda family $\left\langle \Sigma^r p_{s+r} \beta_{r-1}!, \alpha_{r-1} \Sigma^{r-2} \delta_s, \Sigma^{r-2} x \right\rangle$, where α_{r-1} and β_{r-1} fit into a distinguished triangle

$$\Sigma^{r-1} Y_{s+1} \xrightarrow{\alpha_{r-1}} W_{r-1} \xrightarrow{\beta_{r-1}} \Sigma^r Y_{s+r} \xrightarrow{\gamma_{r-1}} \Sigma^r Y_{s+1},$$

and $\gamma_{r-1} = -\Sigma^r(i_{s+1} \cdots i_{s+r-1})$. Thus, to compute the last restricted Toda bracket, one uses the following diagram, obtained as usual from the octahedral axiom:

$$\begin{array}{ccccccc}
 & & & & & & \Sigma^{t-s+r-1}X \\
 & & & & & & \downarrow -\Sigma^{r-1}x \\
 \Sigma^{r-2}I_s & \xrightarrow{\Sigma^{r-2}\delta_s} & \Sigma^{r-1}Y_{s+1} & \xrightarrow{(-1)^r \Sigma^{r-1}i_s} & \Sigma^{r-1}Y_s & \xrightarrow{-\Sigma^{r-1}p_s} & \Sigma^{r-1}I_s \\
 \parallel & & \downarrow \alpha_{r-1} & & \downarrow \alpha_r & & \parallel \\
 \Sigma^{r-2}I_s & \longrightarrow & W_{r-1} & \xrightarrow{q_{r-1}} & W_r & \xrightarrow{\iota_{r-1}} & \Sigma^{r-1}I_s \\
 & & \downarrow \beta_{r-1} & & \downarrow \beta_r & & \\
 \Sigma^r I_{s+r} & \xleftarrow{\Sigma^r p_{s+r}} & \Sigma^r Y_{s+r} & \xlongequal{\quad} & \Sigma^r Y_{s+r} & & \\
 & & \downarrow \gamma_{r-1} & & \downarrow \gamma_r & & \\
 & & \Sigma^r Y_{s+1} & \xrightarrow{(-1)^r \Sigma^r i_s} & \Sigma^r Y_s & &
 \end{array}$$

Up to suspension, both $d_r[x]$ and the last restricted Toda bracket are computed by composing certain maps $\tilde{x} : \Sigma^{t-s+r-2}X \rightarrow \Sigma^r Y_{s+r}$ with $\Sigma^r p_{s+r}$. For $d_r[x]$, the maps \tilde{x} must lift $\Sigma^{r-1}(\delta_s x)$ through $-\gamma_{r-1}$. For the last bracket, the maps \tilde{x} are of the form $\beta_r y$, where $y : \Sigma^{t-s+r-1}X \rightarrow W_r$ is a lift of $-\Sigma^{r-1}x$ through ι_{r-1} . As in the proof of Proposition 4.1(3), one can see that the possible choices of \tilde{x} coincide. \square

We next give a description of $d_r[x]$ using higher Toda brackets defined using filtered objects, as in Definitions 5.4 and 5.5. The computation of the restricted Toda bracket above produces a sequence

$$0 = W_0 \xrightarrow{q_0} W_1 \xrightarrow{q_1} \cdots \xrightarrow{q_{r-1}} W_r, \quad (6.2)$$

where W_k is the fibre of the k -fold composite $\Sigma^r(i_{s+r-k} \cdots i_{s+r-1})$. (The map γ_k may differ in sign from this composite, but that doesn't affect the fibre.) For each k , we have a distinguished triangle

$$W_k \xrightarrow{q_k} W_{k+1} \xrightarrow{\iota_k} \Sigma^{r-1}I_{s+r-k-1} \xrightarrow{-(\Sigma\alpha_k)(\Sigma^{r-1}\delta_{s+r-k-1})} \Sigma W_k,$$

where we extend downwards to $k = 0$ by defining $W_1 = \Sigma^{r-1}I_{s+r-1}$ and using the non-obvious triangle

$$W_0 \xrightarrow{q_0=0} W_1 \xrightarrow{\iota_0=-1} \Sigma^{r-1}I_{s+r-1} \xrightarrow{0} \Sigma W_0.$$

One can check that

$$(\Sigma\iota_{k-1})(-\Sigma\alpha_k)(\Sigma^{r-1}\delta_{s+r-k-1}) = (\Sigma^r p_{s+r-k})(\Sigma^{r-1}\delta_{s+r-k-1}) = \Sigma^{r-1}d_1 = \Sigma^k(\Sigma^{r-k-1}d_1),$$

where $\Sigma^{r-k-1}d_1$ is the map appearing in the $(k+1)$ st spot of the Toda bracket. In other words, the sequence (6.2) is an r -filtered object based on $(\Sigma^{r-2}d_1, \dots, d_1)$.

The natural map $\sigma_W : W_r \rightarrow \Sigma^{r-1}I_s$ is ι_{r-1} , and the natural map $\sigma'_W : \Sigma^{r-1}I_{s+r-1} \cong W_1 \rightarrow W_r$ is the composite $q_{r-1} \cdots q_1 \iota_0 = -q_{r-1} \cdots q_1$. The Toda bracket computed using the filtered object W consists of all composites appearing in the middle row of this commutative

diagram:

$$\begin{array}{ccccc}
 & & \Sigma^{r-1}I_{s+r-1} & & \\
 & & \downarrow \sigma'_W & \searrow \Sigma^{r-1}d_1 & \\
 \Sigma^{t-s+r-1}X & \dashrightarrow^a & W_r & \dashrightarrow^b & \Sigma^r I_{s+r} \\
 & \searrow \Sigma^{r-1}x & \downarrow \sigma_W & & \\
 & & \Sigma^{r-1}I_s & &
 \end{array} \tag{6.3}$$

We claim that there is a natural choice of extension b . Since $\Sigma^{r-1}d_1 = (\Sigma^r p_{s+r})(\Sigma^{r-1}\delta_{s+r-1})$, it suffices to extend $\Sigma^{r-1}\delta_{s+r-1}$ over σ'_W . Well, β_2 by definition is an extension of $\Sigma^{r-1}\delta_{s+r-1}$ over q_1 , and each subsequent β_k gives a further extension. Because $\iota_0 = -1$, $-(\Sigma^r p_{s+r})\beta_r$ is a valid choice for b .

On the other hand, as described at the end of the previous proof, the lifts a of $\Sigma^{r-1}x$ through $\sigma_W = \iota_{r-1}$, when composed with $-(\Sigma^r p_{s+r})\beta_r$, give exactly the Toda bracket computed there.

In summary, we have:

Theorem 6.6. *Given an Adams resolution of Y and $r \geq 2$, there is an associated r -filtered object W and a choice of a map b in Diagram (6.3), such that for any X and class $[x] \in E_r^{s,t}$, we have*

$$d_r[x] = \langle \Sigma^{r-1}d_1, \dots, \Sigma d_1, d_1, x \rangle,$$

where the Toda bracket is computed only using the r -filtered object W and the chosen extension b .

7. SPARSE RINGS OF OPERATIONS

In this section, we focus on injective and projective classes which are generated by an object with a “sparse” endomorphism ring. In this context, we can give conditions under which the restricted Toda bracket appearing in Theorem 6.5 is equal to the unrestricted Toda bracket, producing a cleaner correspondence between Adams differentials and Toda brackets. We begin in Subsection 7.1 by giving the results in the case of an injective class, and then briefly summarize the dual results in Subsection 7.2. Subsection 7.3 gives examples.

Let us fix some notation and terminology, also discussed in [36], [34], [37, §2], and [8].

Definition 7.1. Let N be a natural number. A graded abelian group R_* is N -**sparse** if R_* is concentrated in degrees which are multiples of N , i.e., $R_i = 0$ whenever $i \not\equiv 0 \pmod{N}$.

7.1. Injective case.

Notation 7.2. Let E be an object of the triangulated category \mathcal{T} . Define the E -**cohomology** of an object X to be the graded abelian group E^*X given by $E^n X := \mathcal{T}(X, \Sigma^n E)$. Postcomposition makes E^*X into a left module over the graded endomorphism ring E^*E .

Assumption 7.3. For the remainder of this subsection, we assume the following.

- (1) The triangulated category \mathcal{T} has infinite products.
- (2) The graded ring E^*E is N -sparse for some $N \geq 2$.

Let \mathcal{I}_E denote the injective class generated by E , as in Example 2.6. Explicitly, \mathcal{I}_E consists of retracts of (arbitrary) products $\prod_i \Sigma^{n_i} E$.

Lemma 7.4. *With this setup, we have:*

- (1) Let I be an injective object such that E^*I is N -sparse. Then I is a retract of a product $\prod_i \Sigma^{m_i N} E$.
- (2) If, moreover, W is an object such that E^*W is N -sparse, then we have $\mathcal{T}(W, \Sigma^t I) = 0$ for $t \not\equiv 0 \pmod{N}$.

Proof. (1) I is a retract of a product $P = \prod_i \Sigma^{n_i} E$, with a map $\iota: I \hookrightarrow P$ and retraction $\pi: P \rightarrow I$. Consider the subproduct $P' = \prod_{N|n_i} \Sigma^{n_i} E$, with inclusion $\iota': P' \hookrightarrow P$ (via the zero map into the missing factors) and projection $\pi': P \rightarrow P'$. Then the equality

$$\iota' \pi' \iota = \iota: I \rightarrow P$$

holds, using the fact that E^*I is N -sparse. Therefore, we obtain $\pi \iota' \pi' \iota = \pi \iota = 1_I$, so that I is a retract of P' .

- (2) By the first part, $\mathcal{T}(W, \Sigma^t I)$ is a retract of

$$\begin{aligned} \mathcal{T}(W, \Sigma^t \prod_i \Sigma^{m_i N} E) &= \mathcal{T}(W, \prod_i \Sigma^{m_i N+t} E) \\ &= \prod_i \mathcal{T}(W, \Sigma^{m_i N+t} E) \\ &= \prod_i E^{m_i N+t} W \\ &= 0, \end{aligned}$$

using the assumption that E^*W is N -sparse. \square

Lemma 7.5. Let $I_0 \xrightarrow{f_1} I_1 \xrightarrow{f_2} I_2 \rightarrow \dots \xrightarrow{f_r} I_r$ be a diagram in \mathcal{T} , with $r \leq N+1$. Assume that each object I_j is injective and that each $E^*(I_j)$ is N -sparse. Then the iterated Toda family $\mathbb{T}(f_r, f_{r-1}, \dots, f_1)$ is either empty or consists of a single composable pair $\Sigma^{r-2} I_0 \rightarrow C \rightarrow I_r$, up to automorphism of C .

Proof. In the case $r = 2$, there is nothing to prove, so we may assume $r \geq 3$. The iterated Toda family is obtained by $r-2$ iterations of the 3-fold Toda family construction. The first iteration computes the Toda family of the diagram

$$I_{r-3} \xrightarrow{f_{r-2}} I_{r-2} \xrightarrow{f_{r-1}} I_{r-1} \xrightarrow{f_r} I_r.$$

Choose a cofiber of f_{r-1} , i.e., a distinguished triangle $I_{r-2} \xrightarrow{f_{r-1}} I_{r-1} \rightarrow C_1 \rightarrow \Sigma I_{r-2}$. A lift of f_{r-2} to the fiber $\Sigma^{-1} C_1$, if it exists, is determined up to

$$\mathcal{T}(I_{r-3}, \Sigma^{-1} I_{r-1}) = \mathcal{T}(\Sigma I_{r-3}, I_{r-1}),$$

which is zero by Lemma 7.4(2). Likewise, an extension of f_r to the cofiber C_1 , if it exists, is determined up to

$$\mathcal{T}(\Sigma I_{r-2}, I_r) = 0.$$

Hence, $\mathbb{T}(f_r, f_{r-1}, f_{r-2})$ is either empty or consists of a single pair $(\beta_1, \Sigma \alpha_1)$, up to automorphisms of C_1 . It is easy to see that the object C_1 has the following property:

$$\text{If } E^*W = 0 \text{ for } * \equiv 0, 1 \pmod{N}, \text{ then } \mathcal{T}(W, C_1) = 0. \quad (7.1)$$

For $r \geq 4$, the next iteration computes the Toda family of the diagram

$$\Sigma I_{r-4} \xrightarrow{\Sigma f_{r-3}} \Sigma I_{r-3} \xrightarrow{\Sigma \alpha_1} C_1 \xrightarrow{\beta_1} I_r.$$

The respective indeterminacies are

$$\mathcal{T}(\Sigma^2 I_{r-4}, C_1),$$

which is zero by Property (7.1), and

$$\mathcal{T}(\Sigma^2 I_{r-3}, I_r),$$

which is zero by Lemma 7.4(2), since $N \geq 3$ in this case. Hence, $\mathcal{T}(\beta_1, \Sigma\alpha_1, \Sigma f_{r-3})$ is either empty or consists of a single pair $(\beta_2, \Sigma\alpha_2)$, up to automorphism of the cofiber C_2 of $\Sigma\alpha_1$. Repeating the argument inductively, the successive iterations compute the Toda family of a diagram

$$\Sigma^j I_{r-3-j} \xrightarrow{\Sigma^j f_{r-2-j}} \Sigma^j I_{r-2-j} \xrightarrow{\Sigma\alpha_j} C_j \xrightarrow{\beta_j} I_r$$

for $0 \leq j \leq r-3$, where C_j has the following property:

$$\text{If } E^*W = 0 \text{ for } * \equiv 0, 1, \dots, j \pmod{N}, \text{ then } \mathcal{T}(W, C_j) = 0. \quad (7.2)$$

The indeterminacy groups $\mathcal{T}(\Sigma^{j+1} I_{r-3-j}, C_j)$ and $\mathcal{T}(\Sigma^{j+1} I_{r-2-j}, I_r)$ are again zero. Hence, $\mathcal{T}(\beta_j, \Sigma\alpha_j, \Sigma^j f_{r-2-j})$ is either empty or consists of a single pair $(\beta_{j+1}, \Sigma\alpha_{j+1})$, up to automorphism of C_{j+1} . Note that the argument works until the last iteration $j = r-3$, by the assumption $r-2 < N$. \square

We will need the following condition on an object Y :

Condition 7.6. Y admits an \mathcal{I}_E -Adams resolution Y_\bullet (see (2.2)) such that for each injective I_j in the resolution, $E^*(\Sigma^j I_j)$ is N -sparse.

Remark 7.7.

- (1) Condition 7.6 implies that E^*Y is itself N -sparse, because of the surjection $E^*I_0 \twoheadrightarrow E^*Y$.
- (2) The condition can be generalized to: there is an integer m such that for each j , $E^*(\Sigma^j I_j)$ is concentrated in degrees $* \equiv m \pmod{N}$. We take $m = 0$ for notational convenience.
- (3) We will see in Propositions 7.9 and 7.10 situations in which Condition 7.6 holds.

Theorem 7.8. *Let X and Y be objects in \mathcal{T} and consider the Adams spectral sequence abutting to $\mathcal{T}(X, Y)$ with respect to the injective class \mathcal{I}_E . Assume that Y satisfies Condition 7.6. Then for all $r \leq N$, the Adams differential is given, as a subset of $E_1^{s+r, t+r-1}$, by*

$$d_r[x] = \langle \Sigma^{r-1} d_1, \dots, \Sigma d_1, d_1, x \rangle.$$

In other words, the restricted bracket appearing in Theorem 6.5 coincides with the full Toda bracket.

Proof. We will show that

$$\langle \Sigma^{r-1} d_1!, \dots, \Sigma d_1!, d_1, x \rangle = \langle \Sigma^{r-1} d_1, \dots, \Sigma d_1, d_1, x \rangle.$$

Consider the diagram

$$\begin{array}{ccccccc} I_s & \xrightarrow{d_1} & \Sigma I_{s+1} & \xrightarrow{\Sigma d_1} & \Sigma^2 I_{s+2} & \longrightarrow \dots \longrightarrow & \Sigma^{r-1} I_{r-1} & \xrightarrow{\Sigma^{r-1} d_1} & \Sigma^r I_{s+r} \\ & \uparrow x & & & & & & & \\ & X & & & & & & & \end{array}$$

whose Toda bracket is being computed. The corresponding Toda family is

$$\mathrm{T}(\Sigma^{r-1}d_1, \dots, \Sigma d_1, d_1, x) = \mathrm{T}(\mathrm{T}(\Sigma^{r-1}d_1, \dots, \Sigma d_1, d_1), \Sigma^{r-2}x).$$

We know that

$$\mathrm{T}(\Sigma^{r-1}d_1^!, \dots, \Sigma d_1^!, d_1) \subseteq \mathrm{T}(\Sigma^{r-1}d_1, \dots, \Sigma d_1, d_1).$$

By Lemma 7.5, the Toda family on the right has at most one element, up to automorphism. But fully-restricted Toda families are always non-empty, so the inclusion must be an equality.

Write $\Sigma^{r-2}I_s \xrightarrow{f} C \xrightarrow{g} \Sigma^r I_{s+r}$ for an element of these families. It remains to show that the inclusion

$$\langle g^!, f, \Sigma^{r-2}x \rangle \subseteq \langle g, f, \Sigma^{r-2}x \rangle$$

is an equality, i.e., that the extension of g to the cofiber of f is unique. This follows from the equality $\mathcal{T}(\Sigma^{r-1}I_s, \Sigma^r I_{s+r}) = 0$, which uses the assumption on the injective objects I_j and that $r-1 < N$. \square

Next, we describe situations in which Theorem 7.8 applies.

Proposition 7.9. *Assume that every product of the form $\prod_i \Sigma^{m_i N} E$ has cohomology E^* ($\prod_i \Sigma^{m_i N} E$) which is N -sparse. Then every object Y such that E^*Y is N -sparse also satisfies Condition 7.6.*

Proof. Let (y_i) be a set of non-zero generators of E^*Y as an E^*E -module. Then the corresponding map $Y \rightarrow \prod_i \Sigma^{|y_i|} E$ is \mathcal{I}_E -monic into an injective object; we take this map as the first step $p_0: Y_0 \rightarrow I_0$, with cofiber ΣY_1 . By our assumption on Y , each degree $|y_i|$ is a multiple of N , and thus E^*I_0 is N -sparse, by the assumption on E . The distinguished triangle $Y_1 \rightarrow Y_0 \xrightarrow{p_0} I_0 \rightarrow \Sigma Y_1$ induces a long exact sequence on E -cohomology which implies that the map $I_0 \rightarrow \Sigma Y_1$ is injective on E -cohomology. It follows that $E^*(\Sigma Y_1)$ is N -sparse as well. Repeating this process, we obtain an \mathcal{I}_E -Adams resolution of Y such that for every j , $E^*(\Sigma^j Y_j)$ and $E^*(\Sigma^j I_j)$ are N -sparse. \square

The condition on E is discussed in Example 7.17.

Proposition 7.10. *Assume that the ring E^*E is left coherent, and that E^*Y is N -sparse and finitely presented as a left E^*E -module. Then Y satisfies Condition 7.6.*

Proof. Since E^*Y is finitely generated over E^*E , the map $p_0: Y \rightarrow I_0$ can be chosen so that $I_0 = \prod_i \Sigma^{m_i N} E \cong \oplus_i \Sigma^{m_i N} E$ is a finite product. It follows that E^*I_0 is N -sparse and finitely presented. We have that $E^{*-1}Y_1 = \ker(p_0^*: E^*I_0 \rightarrow E^*Y)$. This is N -sparse, since E^*I_0 is, and is finitely presented over E^*E , since both E^*I_0 and E^*Y are, and E^*E is coherent [9, §I.2, Exercises 11–12]. Repeating this process, we obtain an \mathcal{I}_E -Adams resolution of Y such that for every j , $\Sigma^j I_j$ is a finite product of the form $\prod_i \Sigma^{m_i N} E$. \square

7.2. Projective case. The main applications of Theorem 7.8 are to projective classes instead of injective classes. For future reference, we state here the dual statements of the previous subsection and adopt a notation inspired from stable homotopy theory.

Notation 7.11. Let R be an object of the triangulated category \mathcal{T} . Define the **homotopy** (with respect to R) of an object X as the graded abelian group π_*X given by $\pi_n X := \mathcal{T}(\Sigma^n R, X)$. Precomposition makes π_*X into a right module over the graded endomorphism ring π_*R .

Assumption 7.12. For the remainder of this subsection, we assume the following.

- (1) The triangulated category \mathcal{T} has infinite coproducts.

(2) The graded ring π_*R is N -sparse for some $N \geq 2$.

Let \mathcal{P}_R denote the stable projective class spanned by R , as in Example 2.4. Explicitly, \mathcal{P}_R consists of retracts of (arbitrary) coproducts $\bigoplus_i \Sigma^{m_i} R$.

Condition 7.13. X admits a \mathcal{P}_R -Adams resolution X_\bullet as in Diagram (2.1) with the property that $\pi_*(\Sigma^{-j}P_j)$ is N -sparse for each projective P_j .

Theorem 7.14. *Let X and Y be objects in \mathcal{T} and consider the Adams spectral sequence abutting to $\mathcal{T}(X, Y)$ with respect to the projective class \mathcal{P}_R . Assume that X satisfies Condition 7.13. Let $[y] \in E_r^{s,t}$ be a class represented by $y \in E_1^{s,t} = \mathcal{T}(\Sigma^{t-s}P_s, Y)$. Then for all $r \leq N$, the Adams differential is given, as a subset of $E_1^{s+r, t+r-1}$, by*

$$d_r[y] = \left\langle y, d_1, \Sigma^{-1}d_1, \dots, \Sigma^{-(r-1)}d_1 \right\rangle.$$

Note that we used Corollary 5.13 to ensure that the equality holds as stated, not merely up to sign.

Proposition 7.15. *Assume that every coproduct of the form $\bigoplus_i \Sigma^{m_i} R$ has homotopy $\pi_*(\bigoplus_i \Sigma^{m_i} R)$ which is N -sparse. Then every object X such that π_*X is N -sparse also satisfies Condition 7.13.*

Recall the following terminology:

Definition 7.16. An object X of \mathcal{T} is **compact** if the functor $\mathcal{T}(X, -)$ preserves infinite coproducts.

Example 7.17. If R is compact in \mathcal{T} , then R satisfies the assumption of Proposition 7.15. This follows from the isomorphism

$$\pi_* \left(\bigoplus_i \Sigma^{m_i} R \right) \cong \bigoplus_i \pi_*(\Sigma^{m_i} R) = \bigoplus_i \Sigma^{m_i} \pi_* R$$

and the assumption that π_*R is N -sparse. The same argument works if R is a retract of a coproduct of compact objects.

Dually, if E is cocompact in \mathcal{T} , then E satisfies the assumption of Proposition 7.9. This holds more generally if E is a retract of a product of cocompact objects.

Remark 7.18. Some of the related literature deals with compactly generated triangulated categories. As noted in Remark 2.13, we do *not* assume that the object R is a generator, i.e., that the condition $\pi_*X = 0$ implies $X = 0$.

Proposition 7.19. *Assume that the ring π_*R is right coherent, and that π_*X is N -sparse and finitely presented as a right π_*R -module. Then X satisfies Condition 7.13.*

The following is a variant of [34, Lemma 2.2.2], where we do not assume that R is a generator. It identifies the E_2 term of the spectral sequence associated to the projective class \mathcal{P}_R . The proof is straightforward.

Proposition 7.20. *Assume that the object R is compact.*

(1) *Let P be in the projective class \mathcal{P}_R . Then the map of abelian groups*

$$\mathcal{T}(P, Y) \rightarrow \text{Hom}_{\pi_*R}(\pi_*P, \pi_*Y)$$

is an isomorphism for every object Y .

(2) *There is an isomorphism*

$$\text{Ext}_{\mathcal{P}_R}^s(X, Y) \cong \text{Ext}_{\pi_*R}^s(\pi_*X, \pi_*Y)$$

which is natural in X and Y .

7.3. Examples. Theorem 7.14 applies to modules over certain ring spectra. We describe some examples, along the lines of [34, Examples 2.4.6 and 2.4.7].

Example 7.21. Let R be an A_∞ ring spectrum, and let $h\text{Mod}_R$ denote the homotopy category of the stable model category of (right) R -modules [37, Example 2.3(ii)] [17, §III]. Then R itself, the free R -module of rank 1, is a compact generator for $h\text{Mod}_R$. The R -homotopy of an R -module spectrum X is the usual homotopy of X , as suggested by the notation:

$$h\text{Mod}_R(\Sigma^n R, X) \cong h\text{Mod}_S(S^n, X) = \pi_n X.$$

In particular, the graded endomorphism ring $\pi_* R$ is the usual coefficient ring of R .

The projective class \mathcal{P}_R is the ghost projective class [13, §7.3], generalizing Example 2.4, where R was the sphere spectrum S . The Adams spectral sequence relative to \mathcal{P}_R is the universal coefficient spectral sequence

$$\text{Ext}_{\pi_* R}^s(\Sigma^t \pi_* X, \pi_* Y) \Rightarrow h\text{Mod}_R(\Sigma^{t-s} X, Y)$$

as described in [17, §IV.4] and [13, Corollary 7.12]. We used Proposition 7.20 to identify the E_2 term.

Some A_∞ ring spectra R with sparse homotopy $\pi_* R$ are discussed in [34, §4.3, 5.3, 6.4]. In view of Proposition 7.20, the Adams spectral sequence in $h\text{Mod}_R$ collapses at the E_2 page if $\pi_* R$ has (right) global dimension less than 2.

The Johnson–Wilson spectrum $E(n)$ has coefficient ring

$$\pi_* E(n) = \mathbb{Z}_{(p)}[v_1, \dots, v_n, v_n^{-1}], \quad |v_i| = 2(p^i - 1),$$

which has global dimension n and is $2(p-1)$ -sparse. Hence, Theorem 7.14 applies in this case to the differentials d_r with $r \leq 2(p-1)$, while d_r is zero for $r > n$. Likewise, connective complex K -theory ku has coefficient ring

$$\pi_* ku = \mathbb{Z}[u], \quad |u| = 2,$$

which has global dimension 2 and is 2-sparse.

Example 7.22. Let R be a differential graded (dg for short) algebra over a commutative ring k , and consider the category of dg R -modules dgMod_R . The homology $H_* X$ of a dg R -module is a (graded) $H_* R$ -module. The derived category $D(R)$ is defined as the localization of dgMod_R with respect to quasi-isomorphisms. The free dg R -module R is a compact generator of $D(R)$. The R -homotopy of an object X of $D(R)$ is its homology $\pi_* X = H_* X$. In particular, the graded endomorphism ring of R in $D(R)$ is the graded k -algebra $H_* R$.

The Adams spectral sequence relative to \mathcal{P}_R is an Eilenberg–Moore spectral sequence

$$\text{Ext}_{H_* R}^s(\Sigma^t H_* X, H_* Y) \Rightarrow D(R)(\Sigma^{t-s} X, Y)$$

from ordinary Ext to differential Ext, as described in [4, §8, 10]. See also [25, §III.4], [21, Example 10.2(b)], and [16].

Remark 7.23. Example 7.22 can be viewed as a special case of Example 7.21. Letting HR denote the Eilenberg–MacLane spectrum associated to R , the categories Mod_{HR} and dgMod_R are Quillen equivalent, by [37, Example 2.4(i)] [41, Corollary 2.15], yielding a triangulated equivalence $h\text{Mod}_{HR} \cong D(R)$. The generator HR corresponds to the generator R via this equivalence.

Example 7.24. Let R be a ring, viewed as a dg algebra concentrated in degree 0. Then Example 7.22 yields the ordinary derived category $D(R)$. The graded endomorphism ring of R in $D(R)$ is $H_* R$, which is R concentrated in degree 0. This is N -sparse for any $N \geq 2$.

The Adams spectral sequence relative to \mathcal{P}_R is the hyperderived functor spectral sequence

$$\mathrm{Ext}_{H_*R}^s(\Sigma^t H_*X, H_*Y) = \prod_{i \in \mathbb{Z}} \mathrm{Ext}_R^s(H_{i-t}X, H_iY) \Rightarrow D(R)(\Sigma^{t-s}X, Y) = \mathbf{Ext}_R^{s-t}(X, Y)$$

from ordinary Ext to hyper-Ext, as described in [44, §5.7, 10.7].

APPENDIX A. COMPUTATIONS IN THE STABLE MODULE CATEGORY OF A GROUP

In this appendix, we give some computations in the stable module category of a group algebra kG , where k is a field and G is a finite group. These computations are used in Proposition 4.3.

Write R for the group algebra kG . We will work in the stable module category $\mathcal{T} := \mathrm{StMod}(R)$. This is the category whose objects are (left) R -modules, and whose morphisms from M to N consist of the R -module homomorphisms from M to N modulo those that factor through a projective module. An isomorphism in $\mathrm{StMod}(R)$ is called a **stable equivalence**, and two R -modules M and N are stably equivalent if and only if there are projectives P and Q such that $M \oplus P \cong N \oplus Q$. The category $\mathrm{StMod}(R)$ is triangulated. The suspension ΣM is defined by choosing an embedding of M into an injective module and taking the quotient, the desuspension ΩM is defined by choosing a surjection from a projective to M and taking the kernel, and these are inverse to each other because the projectives and injectives coincide. Given a short exact sequence

$$0 \longrightarrow M_1 \longrightarrow M_2 \longrightarrow M_3 \longrightarrow 0$$

and an embedding of M_1 into an injective module I , one can choose maps

$$\begin{array}{ccccccccc} 0 & \longrightarrow & M_1 & \longrightarrow & M_2 & \longrightarrow & M_3 & \longrightarrow & 0 \\ & & \parallel & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & M_1 & \longrightarrow & I & \longrightarrow & \Sigma M_1 & \longrightarrow & 0 \end{array} \quad (\text{A.1})$$

making the diagram commute in Mod_R . The distinguished triangles are defined to be those triangles isomorphic in $\mathrm{StMod}(R)$ to one of the form

$$M_1 \longrightarrow M_2 \longrightarrow M_3 \longrightarrow \Sigma M_1$$

constructed in this way.

Rather than working with respect to an injective class in \mathcal{T} , we will consider the **ghost projective class** \mathcal{P} , which is generated by the trivial module k . More precisely, \mathcal{P} consists of the retracts of coproducts $\bigoplus_i \Sigma^{n_i} k$, and the associated ideal consists of the maps which induce the zero map in Tate cohomology. See [12, §4.2] for details.

Proposition A.1. *Let G be the cyclic group $C_4 = \langle g \mid g^4 = 1 \rangle$, let $k = \mathbb{F}_2$, and write $R = kG$. There exists an R -module M , an Adams resolutions of M with respect to the ghost projective class, and a map $\kappa : M \rightarrow M$ such that the inclusion $\langle \kappa, d_1, \delta \rangle (\Sigma p) \subseteq \langle \kappa, d_1, d_1 \rangle$ from Proposition 4.2 (dualized) is proper.*

Proof. To produce our counterexample, we will consider the Adams spectral sequence abutting to $\mathrm{StMod}(M, \Omega^* M)$, where M is a two-dimensional module with basis vectors that are interchanged by g .

In order to make concrete computations, it will be helpful to observe that, as a k -algebra, R is the truncated polynomial algebra

$$R = k[g]/(g^4 - 1) = k[g]/(g - 1)^4 = k[x]/x^4,$$

where we define $x := g - 1 \in R$. In this notation, the trivial module k is R/x and the module M is R/x^2 .

We will need to compute their desuspensions, which are given, as R -modules, by:

$$\begin{aligned}\Omega k &= \ker(R \twoheadrightarrow k) = k\{x, x^2, x^3\} \cong R/x^3 \\ \Omega^2 k &= \ker(R \twoheadrightarrow R/x^3) = k\{x^3\} \cong R/x = k \\ \Omega M &= \ker(R \twoheadrightarrow R/x^2) = k\{x^2, x^3\} \cong R/x^2 = M,\end{aligned}$$

where curly brackets denote the k -vector space with the given generating set.

In order to produce a \mathcal{P} -epic map to M , we need to know the maps from suspensions of k to M . Since k is 2-periodic, the following calculations give us what we need:

$$\begin{aligned}\mathcal{T}(k, M) &= \text{Mod}_R(k, M)/\sim \cong \text{Mod}_R(R/x, R/x^2)/\sim = k\{\mu_x\}/\sim = k\{\mu_x\} \\ \mathcal{T}(\Omega k, M) &= \text{Mod}_R(\Omega k, M)/\sim = \text{Mod}_R(R/x^3, R/x^2)/\sim = k\{\mu_1, \mu_x\}/\sim = k\{\mu_1\},\end{aligned}$$

where $f \sim g$ if $f - g$ factors through a projective, and $\mu_r: R/x^m \rightarrow R/x^n$ denotes the R -module map given by multiplication by $r \in R$ (when this is well-defined). Here, we used the fact that $\mu_x: R/x^3 \rightarrow R/x^2$ is stably null, since it factors as

$$R/x^3 \xrightarrow{\mu_x} R \xrightarrow{\mu_1} R/x^2.$$

Using this, we obtain a \mathcal{P} -epic map $p := \mu_x + \mu_1: k \oplus \Omega k \rightarrow M$. Since p is surjective, its fiber is its kernel. This kernel is generated by $(1, x)$ and is readily seen to be isomorphic to M . Under the identification of ΩM with M , the natural map $\Omega M \rightarrow M$ (using the dual of Equation (A.1)) is μ_x . Since we are working at the prime 2, fibre sequences and cofiber sequences agree, so we obtain the following Adams resolution of M

$$\begin{array}{ccccccc} M & \xrightarrow{\mu_x} & M & \xrightarrow{\mu_x} & M & \xrightarrow{\mu_x} & M \rightarrow \dots \\ & \swarrow p & \searrow \delta & & \swarrow p & \searrow \delta & \\ & & k \oplus \Omega k & & k \oplus \Omega k & & k \oplus \Omega k, \end{array}$$

where $\delta = [\begin{smallmatrix} \mu_1 \\ \mu_x \end{smallmatrix}]$, and we have chosen to put the degree shifts on the horizontal arrows.

We will be considering the Adams spectral sequence formed by applying the functor $\mathcal{T}(-, M)$. The map $d_1 = \delta p: k \oplus \Omega k \rightarrow k \oplus \Omega k$ is $[\begin{smallmatrix} 0 & \mu_1 \\ \mu_{x^2} & \mu_x \end{smallmatrix}]$, which simplifies to $[\begin{smallmatrix} 0 & \mu_1 \\ \mu_{x^2} & 0 \end{smallmatrix}]$, using the fact that $\mu_x: \Omega k \rightarrow \Omega k$ is stably null, since it factors as $\Omega k \xrightarrow{\mu_x} R \xrightarrow{\mu_1} \Omega k$. The stable maps $k \oplus \Omega k \rightarrow M$ are of the form $[a\mu_x \ b\mu_1]$ for a and b in k , and all composites $[a\mu_x \ b\mu_1] \circ d_1$ are stably null. Using this twice for d_1 's in different positions, one sees that if $\kappa: k \oplus \Omega k \rightarrow M$ is any map, then $d_2[\kappa]$ is defined and has no indeterminacy.

Now consider $\langle \kappa, d_1, \delta \rangle (\Sigma p)$. One part of the indeterminacy here consists of maps of the form $f\Sigma(\delta)\Sigma(p) = f\Sigma(d_1)$, for $f: \Sigma(k \oplus \Omega k) \rightarrow M$. As above, all such composites are zero. The other part of the indeterminacy consists of maps of the form $\kappa f\Sigma(p)$, for $f: \Sigma M \rightarrow k \oplus \Omega k$, and again, one can show that all such composites are zero. So $\langle \kappa, d_1, \delta \rangle (\Sigma p)$ has no indeterminacy and therefore equals $d_2[\kappa]$.

Finally, consider $\langle \kappa, d_1, d_1 \rangle$. The part of the indeterminacy involving d_1 is again zero. The other part consists of all composites κf , for $f: \Sigma(k \oplus \Omega k) \rightarrow k \oplus \Omega k$. Since there is an isomorphism $\Sigma(k \oplus \Omega k) \rightarrow k \oplus \Omega k$, this indeterminacy is non-zero if and only if κ is non-zero.

Since non-zero maps $\kappa: k \oplus \Omega k \rightarrow M$ exist, we conclude that the containment

$$\langle \kappa, d_1, \delta \rangle (\Sigma p) \subseteq \langle \kappa, d_1, d_1 \rangle$$

can be proper. \square

Remark A.2. If in the proof above we take κ to be the map $[\mu_x \ 0] : k \oplus \Omega k \rightarrow M$, then using the same techniques one can show that

$$\begin{aligned} \langle \kappa, d_1, \delta \rangle &= \{1_M\}, \\ \langle \kappa, d_1, \delta \rangle (\Sigma p) &= \{\Sigma p\} = d_2[\kappa] = \{[\mu_1 \ \mu_x]\}, \end{aligned}$$

and

$$\langle \kappa, d_1, d_1 \rangle = \{[\mu_1 \ b\mu_x] \mid b \in \mathbb{F}_2\}$$

as subsets of $\mathcal{T}(\Omega k \oplus k, M) \cong \mathcal{T}(\Sigma(k \oplus \Omega k), M)$, where we identify M with ΩM and ΣM , as before.

Remark A.3. Theorem 7.14 does not apply to the example in Proposition A.1. Indeed, the graded endomorphism ring of k in $\text{StMod}(kG)$ is the Tate cohomology ring $\tilde{H}^n(G; k) = \text{StMod}(kG)(\Omega^n k, k)$ [11, §6]. This ring is not sparse, as we have $\tilde{H}^{-1}(G; k) \neq 0$.

Example A.4. The following example illustrates the fact that a Toda bracket need not be equal to its own negative, as noted in Remark 3.2.

Consider the ground field $k = \mathbb{F}_3$ and the group algebra $R = kC_3 \cong k[x]/x^3$, where we denote $x = g - 1 \in R$ for $g \in C_3$ a generator. Consider the R -modules $k = R/x$ and $M = R/x^2$. Let us compute the Toda bracket of the diagram

$$M \xrightarrow{\mu_1} k \xrightarrow{\mu_x} M \xrightarrow{\mu_1} k$$

in the triangulated category $\mathcal{T} = \text{StMod}(R)$. We will use appropriate isomorphisms $\Sigma k \cong M$ and $\Sigma M \cong k$, and in particular compute the Toda bracket as a subset of $\mathcal{T}(k, k) \cong \mathcal{T}(\Sigma M, k)$. Via these isomorphisms, the suspension $\Sigma\mu_1 : \Sigma M \rightarrow \Sigma k$ equals $\mu_x : k \rightarrow M$. Consider the commutative diagram in \mathcal{T}

$$\begin{array}{ccccc} & & & k & \xrightarrow{-\mu_x} & M \\ & & & \downarrow & & \parallel \\ & & & \Sigma\alpha \downarrow & & \\ k & \xrightarrow{\mu_x} & M & \xrightarrow{\mu_1} & k & \xrightarrow{\mu_x} & M \\ & & \parallel & & \downarrow & & \\ & & M & \xrightarrow{\mu_1} & k & & \\ & & & & \downarrow & & \\ & & & & \beta \downarrow & & \end{array}$$

where the middle row is distinguished. The only choices for the dotted arrows are $\Sigma\alpha = -1_k$ and $\beta = 1_k$, from which we conclude

$$\langle \mu_1, \mu_x, \mu_1 \rangle_{f_C} = \{-1_k\} \subset \mathcal{T}(k, k).$$

APPENDIX B. 3-FOLD TODA BRACKETS DETERMINE THE TRIANGULATED STRUCTURE

Heller proved the following theorem in [20, Theorem 13.2]. We present an arguably simpler proof here. The argument was kindly provided by Fernando Muro.

Theorem B.1. *Let \mathcal{T} be a triangulated category. Then the diagram $X \xrightarrow{f} Y \xrightarrow{g} Z \xrightarrow{h} \Sigma X$ in \mathcal{T} is a distinguished triangle if and only if the following two conditions hold.*

- (1) *The sequence of abelian groups*

$$\mathcal{T}(A, \Sigma^{-1}Z) \xrightarrow{(\Sigma^{-1}h)_*} \mathcal{T}(A, X) \xrightarrow{f_*} \mathcal{T}(A, Y) \xrightarrow{g_*} \mathcal{T}(A, Z) \xrightarrow{h_*} \mathcal{T}(A, \Sigma X)$$

is exact for every object A of \mathcal{T} .

- (2) *The Toda bracket $\langle h, g, f \rangle \subseteq \mathcal{T}(\Sigma X, \Sigma X)$ contains the identity map $1_{\Sigma X}$.*

Proof. (\Rightarrow) A distinguished triangle satisfies the first condition. For the second condition, consider the commutative diagram

$$\begin{array}{ccccccc} X & \xrightarrow{f} & Y & \xrightarrow{g} & Z & \xrightarrow{h} & \Sigma X \\ \parallel & & \parallel & & \downarrow 1_Z & & \downarrow 1_{\Sigma X} \\ X & \xrightarrow{f} & Y & \xrightarrow{g} & Z & \xrightarrow{h} & \Sigma X. \end{array}$$

Since the top row is distinguished, this diagram exhibits the membership $1_{\Sigma X} \in \langle h, g, f \rangle$.

(\Leftarrow) Assume that $1_{\Sigma X} \in \langle h, g, f \rangle$ holds. By definition of the Toda bracket, there exists a map $\varphi: C_f \rightarrow Z$ making the diagram

$$\begin{array}{ccccccc} X & \xrightarrow{f} & Y & \xrightarrow{q} & C_f & \xrightarrow{\iota} & \Sigma X \\ \parallel & & \parallel & & \downarrow \varphi & & \downarrow 1_{\Sigma X} \\ X & \xrightarrow{f} & Y & \xrightarrow{g} & Z & \xrightarrow{h} & \Sigma X \end{array}$$

commute, where the top row is distinguished. To show that the bottom row is distinguished, it suffices to show that $\varphi: C_f \rightarrow Z$ is an isomorphism. By the Yoneda lemma, it suffices to show that $\varphi_*: \mathcal{T}(A, C_f) \rightarrow \mathcal{T}(A, Z)$ is an isomorphism for every object A of \mathcal{T} .

Consider the diagram

$$\begin{array}{ccccccccc} X & \xrightarrow{f} & Y & \xrightarrow{q} & C_f & \xrightarrow{\iota} & \Sigma X & \xrightarrow{-\Sigma f} & \Sigma Y \\ \parallel & & \parallel & & \downarrow \varphi & & \downarrow 1_{\Sigma X} & & \downarrow 1_{\Sigma Y} \\ X & \xrightarrow{f} & Y & \xrightarrow{g} & Z & \xrightarrow{h} & \Sigma X & \xrightarrow{-\Sigma f} & \Sigma Y \end{array} \quad (\text{B.1})$$

Applying $\mathcal{T}(A, -)$ yields the diagram of abelian groups

$$\begin{array}{ccccccccc} \mathcal{T}(A, X) & \xrightarrow{f_*} & \mathcal{T}(A, Y) & \xrightarrow{q_*} & \mathcal{T}(A, C_f) & \xrightarrow{\iota_*} & \mathcal{T}(A, \Sigma X) & \xrightarrow{(-\Sigma f)_*} & \mathcal{T}(A, \Sigma Y) \\ \parallel & & \parallel & & \downarrow \varphi_* & & \downarrow 1 & & \downarrow 1 \\ \mathcal{T}(A, X) & \xrightarrow{f_*} & \mathcal{T}(A, Y) & \xrightarrow{g_*} & \mathcal{T}(A, Z) & \xrightarrow{h_*} & \mathcal{T}(A, \Sigma X) & \xrightarrow{(-\Sigma f)_*} & \mathcal{T}(A, \Sigma Y). \end{array}$$

The top row is exact, since the top row of (B.1) is a cofiber sequence, and the bottom row is exact, using the first condition. By the 5-lemma, φ_* is an isomorphism. \square

Remark B.2. Some remarks about the first condition.

- (1) It implies $gf = g_*f_*(1_X) = 0$ and $hg = h_*g_*(1_Y) = 0$.
- (2) It is equivalent to the exactness of the long sequence (infinite in both directions)

$$\cdots \longrightarrow \mathcal{T}(A, \Sigma^n X) \xrightarrow{(\Sigma^n f)_*} \mathcal{T}(A, \Sigma^n Y) \xrightarrow{(\Sigma^n g)_*} \mathcal{T}(A, \Sigma^n Z) \xrightarrow{(\Sigma^n h)_*} \mathcal{T}(A, \Sigma^{n+1} X) \longrightarrow \cdots$$

for every object A of \mathcal{T} .

- (3) It is a weaker version of what is sometimes called a *pre-triangle* [33, §1.1]. Indeed, the condition states that the sequence

$$H(\Sigma^{-1}Z) \xrightarrow{H(\Sigma^{-1}h)} H(X) \xrightarrow{H(f)} H(Y) \xrightarrow{H(g)} H(Z) \xrightarrow{H(h)} H(\Sigma X)$$

is exact for every decent homological functor $H: \mathcal{T} \rightarrow \text{Ab}$ of the form $H = \mathcal{T}(A, -)$.

Corollary B.3. *Given the suspension functor $\Sigma: \mathcal{T} \rightarrow \mathcal{T}$, 3-fold Toda brackets in \mathcal{T} determine the triangulated structure. In particular, 3-fold Toda brackets determine the higher Toda brackets, via the triangulation.*

Remark B.4. It is unclear to us if the higher Toda brackets can be expressed directly in terms of three-fold brackets.

REFERENCES

- [1] J. F. Adams, *On the non-existence of elements of Hopf invariant one*, Ann. of Math. (2) **72** (1960), 20–104. MR0141119 (25 #4530)
- [2] ———, *Stable homotopy and generalised homology*, Chicago Lectures in Mathematics, vol. 17, University of Chicago Press, Chicago, IL, 1974.
- [3] P. Balmer, *Triangulated categories with several triangulations* (2002), Preprint, available at <http://www.math.ucla.edu/~balmer/Pubfile/TriangulationS.pdf>.
- [4] T. Barthel, J. P. May, and E. Riehl, *Six model structures for DG-modules over DGAs: model category theory in homological action*, New York J. Math. **20** (2014), 1077–1159. MR3291613
- [5] H.-J. Baues and M. Jibladze, *Secondary derived functors and the Adams spectral sequence*, Topology **45** (2006), no. 2, 295–324, DOI 10.1016/j.top.2005.08.001. MR2193337 (2006k:55031)
- [6] ———, *Dualization of the Hopf algebra of secondary cohomology operations and the Adams spectral sequence*, J. K-Theory **7** (2011), no. 2, 203–347, DOI 10.1017/is010010029jkt133. MR2787297 (2012h:55023)
- [7] H.-J. Baues and D. Blanc, *Higher order derived functors and the Adams spectral sequence*, J. Pure Appl. Algebra **219** (2015), no. 2, 199–239, DOI 10.1016/j.jpaa.2014.04.018. MR3250522
- [8] D. Benson, H. Krause, and S. Schwede, *Realizability of modules over Tate cohomology*, Trans. Amer. Math. Soc. **356** (2004), no. 9, 3621–3668 (electronic), DOI 10.1090/S0002-9947-03-03373-7. MR2055748
- [9] N. Bourbaki, *Commutative algebra. Chapters 1–7*, Elements of Mathematics (Berlin), Springer-Verlag, Berlin, 1998. Translated from the French; Reprint of the 1989 English translation. MR1727221
- [10] R. R. Bruner, *An Adams spectral sequence primer*, 2009, Unpublished notes.
- [11] J. F. Carlson, *Modules and group algebras*, Lectures in Mathematics ETH Zürich, Birkhäuser Verlag, Basel, 1996. Notes by Ruedi Suter. MR1393196
- [12] S. K. Chebolu, J. D. Christensen, and J. Mináč, *Ghosts in modular representation theory*, Adv. Math. **217** (2008), no. 6, 2782–2799, DOI 10.1016/j.aim.2007.11.008. MR2397466 (2008m:20018)
- [13] J. D. Christensen, *Ideals in triangulated categories: phantoms, ghosts and skeleta*, Adv. Math. **136** (1998), no. 2, 284–339, DOI 10.1006/aima.1998.1735. MR1626856 (99g:18007)
- [14] ———, *Python code for comparing sequences* (2015), <https://trinket.io/python/8932cfbf2f>.
- [15] J. M. Cohen, *The decomposition of stable homotopy*, Ann. of Math. (2) **87** (1968), 305–320. MR0231377 (37 #6932)
- [16] S. Eilenberg and J. C. Moore, *Homology and fibrations. I. Coalgebras, cotensor product and its derived functors*, Comment. Math. Helv. **40** (1966), 199–236. MR0203730
- [17] A. D. Elmendorf, I. Kriz, M. A. Mandell, and J. P. May, *Rings, modules, and algebras in stable homotopy theory*, Mathematical Surveys and Monographs, vol. 47, American Mathematical Society, Providence, RI, 1997. With an appendix by M. Cole. MR1417719 (97h:55006)
- [18] S. I. Gelfand and Y. I. Manin, *Methods of homological algebra*, 2nd ed., Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2003. MR1950475 (2003m:18001)
- [19] J. R. Harper, *Secondary cohomology operations*, Graduate Studies in Mathematics, vol. 49, American Mathematical Society, Providence, RI, 2002. MR1913285 (2004c:55035)
- [20] A. Heller, *Stable homotopy categories*, Bull. Amer. Math. Soc. **74** (1968), 28–63. MR0224090 (36 #7137)
- [21] M. Hovey, J. H. Palmieri, and N. P. Strickland, *Axiomatic stable homotopy theory*, Mem. Amer. Math. Soc. **128** (1997), no. 610, x+114, DOI 10.1090/memo/0610. MR1388895
- [22] M. Hovey and N. P. Strickland, *Morava K-theories and localisation*, Mem. Amer. Math. Soc., vol. 139, AMS, 1999.
- [23] S. Klaus, *Towers and pyramids. I*, Forum Math. **13** (2001), no. 5, 663–683, DOI 10.1515/form.2001.028. MR1858494 (2002k:55044)
- [24] S. O. Kochman, *Uniqueness of Massey products on the stable homotopy of spheres*, Canad. J. Math. **32** (1980), no. 3, 576–589, DOI 10.4153/CJM-1980-044-9. MR586976 (81k:55032)
- [25] I. Kříž and J. P. May, *Operads, algebras, modules and motives*, Astérisque **233** (1995), iv+145pp (English, with English and French summaries). MR1361938

- [26] H. R. Margolis, *Spectra and the Steenrod algebra*, North-Holland Mathematical Library, vol. 29, North-Holland Publishing Co., Amsterdam, 1983. Modules over the Steenrod algebra and the stable homotopy category. MR738973 (86j:55001)
- [27] C. R. F. Maunder, *Cohomology operations of the Nth kind*, Proc. London Math. Soc. (3) **13** (1963), 125–154. MR0211398 (35 #2279)
- [28] ———, *On the differentials in the Adams spectral sequence*, Proc. Cambridge Philos. Soc. **60** (1964), 409–420. MR0167980 (29 #5245)
- [29] J. C. McKeown, *nLab: Toda bracket* (2012), available at <http://ncatlab.org/nlab/show/Toda+bracket>.
- [30] L. Meier, *United Elliptic Homology*, 2012. Thesis (Ph.D.)—Universität Bonn.
- [31] *MathOverflow: Second homotopy group of the mod 2 Moore spectrum*, <http://mathoverflow.net/questions/100272/>.
- [32] R. E. Mosher and M. C. Tangora, *Cohomology operations and applications in homotopy theory*, Harper & Row, Publishers, New York-London, 1968. MR0226634 (37 #2223)
- [33] A. Neeman, *Triangulated categories*, Annals of Mathematics Studies, vol. 148, Princeton University Press, Princeton, NJ, 2001. MR1812507 (2001k:18010)
- [34] I. Patchkoria, *On the algebraic classification of module spectra*, Algebr. Geom. Topol. **12** (2012), no. 4, 2329–2388, DOI 10.2140/agt.2012.12.2329. MR3020210
- [35] D. C. Ravenel, *Complex cobordism and stable homotopy groups of spheres*, Second, Vol. 347, AMS Chelsea Publishing, 2004.
- [36] S. Sagave, *Universal Toda brackets of ring spectra*, Trans. Amer. Math. Soc. **360** (2008), no. 5, 2767–2808, DOI 10.1090/S0002-9947-07-04487-X. MR2373333 (2008j:55009)
- [37] S. Schwede and B. Shipley, *Stable model categories are categories of modules*, Topology **42** (2003), no. 1, 103–153, DOI 10.1016/S0040-9383(02)00006-X. MR1928647
- [38] S. Schwede, *Algebraic versus topological triangulated categories*, Triangulated categories, London Math. Soc. Lecture Note Ser., vol. 375, Cambridge University Press, Cambridge, 2010, pp. 389–407, DOI 10.1017/CBO9781139107075.010. MR2681714 (2012i:18012)
- [39] ———, *Symmetric spectra*, Version 3.0, Apr. 12, 2012, Unpublished.
- [40] B. Shipley, *An algebraic model for rational S^1 -equivariant stable homotopy theory*, Q. J. Math. **53** (2002), no. 1, 87–110, DOI 10.1093/qjmath/53.1.87. MR1887672 (2003a:55026)
- [41] ———, *HZ-algebra spectra are differential graded algebras*, Amer. J. Math. **129** (2007), no. 2, 351–379, DOI 10.1353/ajm.2007.0014. MR2306038
- [42] E. Spanier, *Higher order operations*, Trans. Amer. Math. Soc. **109** (1963), 509–539. MR0158399 (28 #1622)
- [43] H. Toda, *Composition methods in homotopy groups of spheres*, Annals of Mathematics Studies, No. 49, Princeton University Press, Princeton, N.J., 1962. MR0143217 (26 #777)
- [44] C. Weibel, *An introduction to homological algebra*, Cambridge Studies in Advanced Mathematics, Cambridge University Press, 1994.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WESTERN ONTARIO, LONDON, ONTARIO, N6A 5B7,
CANADA

E-mail address: `jdc@uwo.ca`

UNIVERSITÄT OSNABRÜCK, INSTITUT FÜR MATHEMATIK, ALBRECHTSTR. 28A, 49076 OSNABRÜCK, GER-
MANY

E-mail address: `martin.frankland@uni-osnabrueck.de`