

Upper bounds for the achromatic and coloring numbers of a graph*

Baoyindureng Wu [†]

College of Mathematics and System Sciences, Xinjiang University

Urumqi, Xinjiang 830046, P.R.China

Clive Elphick [‡]

Abstract: Dvořák *et al.* introduced a variant of the Randić index of a graph G , denoted by $R'(G)$, where $R'(G) = \sum_{uv \in E(G)} \frac{1}{\max\{d(u), d(v)\}}$, and $d(u)$ denotes the degree of a vertex u in G . The coloring number $col(G)$ of a graph G is the smallest number k for which there exists a linear ordering of the vertices of G such that each vertex is preceded by fewer than k of its neighbors. It is well-known that $\chi(G) \leq col(G)$ for any graph G , where $\chi(G)$ denotes the chromatic number of G . In this note, we show that for any graph G without isolated vertices, $col(G) \leq 2R'(G)$, with equality if and only if G is obtained from identifying the center of a star with a vertex of a complete graph. This extends some known results. In addition, we present some new spectral bounds for the coloring and achromatic numbers of a graph.

Keywords: Chromatic number; Coloring number; Achromatic number; Randić index

1 Introduction

The Randić index $R(G)$ of a (molecular) graph G was introduced by Milan Randić [17] in 1975 as the sum of $1/\sqrt{d(u)d(v)}$ over all edges uv of G , where $d(u)$ denotes the degree of a vertex u in G . Formally,

$$R(G) = \sum_{uv \in E(G)} \frac{1}{\sqrt{d(u)d(v)}}.$$

*Research supported by NSFC (No. 11161046)

[†]Corresponding author : Email: wubaoyin@hotmail.com (B. Wu)

[‡]Email: clive.elphick@gmail.com (C. Elphick)

This index is useful in mathematical chemistry and has been extensively studied, see [12]. For some recent results on the Randić index, we refer to [3, 13, 14, 15].

A variation of the Randić index of a graph G is called the Harmonic index, denoted by $H(G)$, which was defined in [8] as follows:

$$H(G) = \sum_{uv \in E(G)} \frac{2}{d(u) + d(v)}.$$

In 2011 Dvořák *et al.* introduced another variant of the Randić index of a graph G , denoted by $R'(G)$, which has been further studied by Knor *et al.* [16]. Formally,

$$R'(G) = \sum_{uv \in E(G)} \frac{1}{\max\{d(u), d(v)\}}.$$

It is clear from the definitions that for a graph G ,

$$R'(G) \leq H(G) \leq R(G). \quad (1)$$

The chromatic number of G , denoted by $\chi(G)$, is the smallest number of colors needed to color all vertices of G such that no pair of adjacent vertices is colored the same. As usual, $\delta(G)$ and $\Delta(G)$ denote the minimum degree and the maximum degree of G , respectively. The coloring number $col(G)$ of a graph G is the least integer k such that G has a vertex ordering in which each vertex is preceded by fewer than k of its neighbors. The *degeneracy* of G , denoted by $deg(G)$, is defined as $deg(G) = \max\{\delta(H) : H \subseteq G\}$. It is well-known (see Page 8 in [11]) that for any graph G ,

$$col(G) = deg(G) + 1. \quad (2)$$

List coloring is an extension of coloring of graphs, introduced by Vizing [21] and independently, by Erdős *et al.* [6]. For each vertex v of a graph G , let $L(v)$ denote a list of colors assigned to v . A list coloring is a coloring l of vertices of G such that $l(v) \in L(v)$ and $l(x) \neq l(y)$ for any $xy \in E(G)$, where $v, x, y \in V(G)$. A graph G is k -choosable if for any list assignment L to each vertex $v \in V(G)$ with $|L(v)| \geq k$, there always exists a list coloring l of G . The list chromatic number $\chi_l(G)$ (or choice number) of G is the minimum k for which G is k -choosable.

It is well-known that for any graph G ,

$$\chi(G) \leq \chi_l(G) \leq col(G) \leq \Delta(G) + 1. \quad (3)$$

The detail of the inequalities in (3) can be found in a survey paper by Tuza [20] on list coloring.

In 2009, Hansen and Vukicević [9] established the following relation between the Randić index and the chromatic number of a graph.

Theorem 1.1. (Hansen and Vukicević [9]) *Let G be a simple graph with chromatic number $\chi(G)$ and Randić index $R(G)$. Then $\chi(G) \leq 2R(G)$ and equality holds if G is a complete graph, possibly with some additional isolated vertices.*

Some interesting extensions of Theorem 1.1 were recently obtained.

Theorem 1.2. (Deng et al [2]) *For a graph G , $\chi(G) \leq 2H(G)$ with equality if and only if G is a complete graph possibly with some additional isolated vertices.*

Theorem 1.3. (Wu, Yan and Yang [23]) *If G is a graph of order n without isolated vertices, then*

$$\text{col}(G) \leq 2R(G),$$

with equality if and only if $G \cong K_n$.

Let n and k be two integers such that $n \geq k \geq 1$. We denote the graph obtained from identifying the center of the star $K_{1,n-k}$ with a vertex of the complete graph K_k by $K_k \bullet K_{1,n-k}$. In particular, if $k \in \{1, 2\}$, $K_k \bullet K_{1,n-k} \cong K_{1,n-1}$; if $k = n$, $K_k \bullet K_{1,n-k} \cong K_n$. The primary aim of this note is to prove stronger versions of Theorems 1.1-1.3, noting the inequalities in (1).

Theorem 1.4. *For a graph G of order n without isolated vertices, $\text{col}(G) \leq 2R'(G)$, with equality if and only if $G \cong K_k \bullet K_{1,n-k}$ for some $k \in \{1, \dots, n\}$.*

Corollary 1.5. *For a graph G of order n without isolated vertices, $\chi(G) \leq 2R'(G)$, with equality if and only if $G \cong K_k \bullet K_{1,n-k}$ for some $k \in \{1, \dots, n\}$.*

Corollary 1.6. *For a graph G of order n without isolated vertices, $\chi_l(G) \leq 2R'(G)$, with equality if and only if $G \cong K_k \bullet K_{1,n-k}$ for some $k \in \{1, \dots, n\}$.*

Corollary 1.7. *For a graph G of order n without isolated vertices, $\text{col}(G) \leq 2H(G)$, with equality if and only if $G \cong K_n$.*

The proofs of these results will be given in the next section.

A *complete k -coloring* of a graph G is a k -coloring of the graph such that for each pair of different colors there are adjacent vertices with these colors. The *achromatic number* of G , denoted by $\psi(G)$, is the maximum number k for which the graph has a complete k -coloring. Clearly, $\chi(G) \leq \psi(G)$ for a graph G . In general, $\text{col}(G)$ and $\psi(G)$ are incomparable. Tang *et al.* [19] proved that for a graph G , $\psi(G) \leq 2R(G)$.

In Section 3, we prove new bounds for the coloring and achromatic numbers of a graph in terms of its spectrum, which strengthen $\text{col}(G) \leq 2R(G)$ and $\psi(G) \leq 2R(G)$. In Section 4, we provide an example and propose two related conjectures.

2 The proofs

For convenience, an edge e of a graph G may be viewed as a 2-element subset of $V(G)$ and if a vertex v is an end vertex of e , we denote the other end of e by $e \setminus v$. Moreover, $\partial_G(v)$ denotes the set of edges which are incident with v in G .

First we need the following theorem, which will play a key role in the proof of Theorem 1.4.

Theorem 2.1. *If v is a vertex of G with $d(v) = \delta(G)$, then*

$$R'(G) - R'(G - v) \geq 0,$$

with equality if and only if $N_G(v)$ is an independent set of G and $d(w) < d(v_i)$ for all $w \in N(v_i) \setminus \{v\}$.

Proof. Let $k = d_G(v)$. The result is trivial for $k = 0$. So, let $k > 0$ and let $N_G(v) = \{v_1, \dots, v_k\}$ and $d_i = d_G(v_i)$ for each i . Without loss of generality, we may assume that $d_1 \geq \dots \geq d_k$. Let

$$E_1 = \{e : e \in \partial_G(v_1) \setminus \{vv_1\} \text{ such that } d_G(e \setminus v_1) < d_G(v_1) \text{ and } e \not\subseteq N_G(v)\},$$

$$E'_1 = \{e : e \in \partial_G(v_1) \setminus \{vv_1\} \text{ such that } d_G(e \setminus v_1) = d_G(v_1) \text{ and } e \subseteq N_G(v)\},$$

and for an integer $i \geq 2$,

$$E_i = \{e : e \in \partial_G(v_i) \setminus \{vv_i\} \text{ such that } d_G(e \setminus v_i) < d_G(v_i) \text{ and } e \not\subseteq N_G(v)\},$$

$$E'_i = \{e : e \in \partial_G(v_i) \setminus \{vv_i\} \text{ such that } d_G(e \setminus v_i) = d_G(v_i) \text{ and } e \subseteq N_G(v)\} \setminus (\cup_{j=1}^{i-1} E'_j).$$

Let $a_i = |E_i|$ and $b_i = |E'_i|$ for any $i \geq 1$. Since $a_i + b_i \leq d_i - 1$,

$$R'(G) - R'(G - v) = \sum_{i=1}^k \frac{1}{d_i} - \sum_{i=1}^k \frac{a_i + b_i}{d_i(d_i - 1)} \geq 0,$$

with equality if and only if $a_i + b_i = d_i - 1$ for all i , i.e., $N_G(v)$ is an independent set of G and $d(w) < d(v_i)$ for all $w \in N(v_i) \setminus \{v\}$. □

Lemma 2.2. *If T is a tree of order $n \geq 2$, then $R'(T) \geq 1$, with equality if and only if $T = K_{1,n-1}$.*

Proof. By induction on n . It can be easily checked that $R'(K_{1,n-1}) = 1$. So, the assertion of the lemma is true for $n \in \{2, 3\}$. So, we assume that $n \geq 4$. Let v be a leaf of T . Then $T - v$ is a tree of order $n - 1$. By Theorem 2.1, $R'(T) \geq R'(T - v) \geq 1$. If $R'(T) = 1$, then $R'(T - v) = 1$, and by the induction hypothesis, $T - v \cong K_{1,n-2}$. Let u be the center of $T - v$. We claim that $vu \in E(T)$. If this is not, then v is adjacent to a leaf of $T - v$, say x , in T . Thus $d_T(x) = 2$. However, since $R'(T) = R'(T - v)$ and $u \in N_T(x)$, by Theorem 2.1 that $d_T(x) > d_T(u) = n - 2 \geq 2$, a contradiction. This shows that $vu \in E(T)$ and $T \cong K_{1,n-1}$. □

The proof of Theorem 1.4:

Let v_1, v_2, \dots, v_n be an ordering of all vertices of G such that v_i is a minimum degree vertex of $G_i = G - \{v_{i+1}, \dots, v_n\}$ for each $i \in \{1, \dots, n\}$, where $G_n = G$ and $d_G(v_n) = \delta(G)$. It is well known that $\deg(G) = \max\{d_{G_i}(v_i) : 1 \leq i \leq n\}$ (see Theorem 12 in [11]). Let k be the maximum number such that $\deg(G) = d_{G_k}(v_k)$, and n_k the order of G_k . By Theorem 2.1,

$$2R'(G) \geq 2R'(G_{n-1}) \geq \dots \geq 2R'(G_k). \quad (4)$$

Moreover,

$$\begin{aligned} 2R'(G_k) &= \sum_{uv \in E(G_k)} \frac{2}{\max\{d_{G_k}(u), d_{G_k}(v)\}} \\ &\geq \sum_{uv \in E(G_k)} \frac{2}{\Delta(G_k)} \\ &\geq \frac{\Delta(G_k) + (n_k - 1)\delta(G_k)}{\Delta(G_k)} \\ &\geq \delta(G_k) + 1 \\ &= \text{col}(G). \end{aligned}$$

It follows that

$$\text{if } R'(G_k) = \delta(G_k) + 1, \text{ then } G_k \cong K_k. \quad (5)$$

Now assume that $\text{col}(G) = 2R'(G)$. Observe that $\text{col}(G) = \max\{\text{col}(G_i) : G_i \text{ is a component of } G\}$ and $R'(G) = \sum_i R'(G_i)$. Thus, by the assumption that $\text{col}(G) = 2R'(G)$ and G has no isolated vertices, G is connected and $\text{col}(G) \geq 2$. If $\text{col}(G) = 2$, then G is a tree. By Corollary 2.1, $G \cong K_{1, n-1}$. Next we assume that $\text{col}(G) \geq 3$. By (4) and (5) we have $R'(G) = \dots = R'(G_k)$, $G_k \cong K_k$ and thus $\text{col}(G) = k$. We show $G \cong K_k \bullet K_{1, n-k}$ by induction on $n - k$. It is easy to check that $\text{col}(K_n) = n + 1 = 2R'(K_n)$ and thus the result is for $n - k = 0$. If $n - k = 1$, then $G_k = G - v_n$ and $R'(G_k) = R'(G)$, by Theorem 2.1, $N_G(v_n)$ is an independent set of G . Combining with $G_k \cong K_k$ ($k \geq 3$), $d_{G_n}(v_n) = 1$. Thus $G = K_k \bullet K_{1, 1}$.

Next assume that $n - k \geq 2$. We consider G_{n-1} . Since $\text{col}(G_{n-1}) = 2R'(G_{n-1})$, by the induction hypothesis $G_{n-1} \cong K_k \bullet K_{1, n-1-k}$. Without loss of generality, let $N_{G_{n-1}}(v_{k+1}) = \dots = N_{G_{n-1}}(v_{n-1}) = \{v_k\}$. So, it remains to show that $N_G(v_n) = \{v_k\}$.

Claim 1. $N_G(v_n) \cap \{v_{k+1}, \dots, v_{n-1}\} = \emptyset$.

If it is not, then $\{v_{k+1}, \dots, v_{n-1}\} \subseteq N_G(v_n)$, because $d_G(v_n) = \delta(G)$. By Theorem 2.1, $d_G(v_{n-1}) > d_G(v_k)$. But, in this case, $d_G(v_{n-1}) = 2 < d_G(v_k)$, a contradiction.

By Claim 1, $N_G(v_n) \subseteq \{v_1, \dots, v_k\}$. Since $N_G(v_n)$ is an independent set and $\{v_1, \dots, v_k\}$ is a clique of G , $|N_G(v_n) \cap \{v_1, \dots, v_k\}| = 1$. If $N_G(v_n) = \{v_i\}$, where $i \neq k$, then $d_G(v_i) = k$. By Theorem 2.1, $d_G(v_i) > d_G(v_k)$. However, $d_G(v_k) \geq n - 2 \geq k$, a contradiction. This shows $N_G(v_n) = \{v_k\}$ and hence $G \cong K_k \bullet K_{1, n-k}$.

It is straightforward to check that

$$2R'(K_k \bullet K_{1, n-k}) = 2 \times \left(\frac{k-2}{2} + 1 \right) = k = \text{col}(G).$$

The proofs of Corollaries 1.5 and 1.6:

By (3) and Theorem 1.4, we have $\chi(G) \leq \chi_l(G) \leq 2R'(G)$. If $\chi(G) = 2R'(G)$ (or $\chi_l(G) = 2R'(G)$), then $\text{col}(G) = 2R'(G)$. By Theorem 1.4, $G \cong K_k \bullet K_{1, n-k}$. On the other hand, it is easy to check that

$$\chi(K_k \bullet K_{1, n-k}) = \chi_l(K_k \bullet K_{1, n-k}) = k = 2R'(K_k \bullet K_{1, n-k}).$$

The proof of Corollary 1.7:

By (1) and Theorem 1.4, we have $\text{col}(G) \leq 2H(G)$. If $\text{col}(G) = 2H(G)$, then $\text{col}(G) = 2R'(G)$. By Theorem 1.4, $G \cong K_k \bullet K_{1, n-k}$. It can be checked that $k = 2H(K_k \bullet K_{1, n-k})$ if and only if $k = n$, i.e., $G \cong K_n$.

3 Spectral bounds

3.1 Definitions

Let $\mu = \mu_1 \geq \dots \geq \mu_n$ denote the eigenvalues of the adjacency matrix of G and let π, ν and γ denote the numbers (counting multiplicities) of positive, negative and zero eigenvalues respectively. Then let

$$s^+ = \sum_{i=1}^{\pi} \mu_i^2 \text{ and } s^- = \sum_{i=n-\nu+1}^n \mu_i^2.$$

Note that $\sum_{i=1}^n \mu_i^2 = s^+ + s^- = \text{tr}(A^2) = 2m$.

3.2 Bounds for $\psi(G)$ and $\text{col}(G)$

Theorem 3.1. *For a graph G , $\psi(G) \leq 2m/\sqrt{s^+} \leq 2m/\mu \leq 2R(G)$.*

Proof. Ando and Lin [1] proved a conjecture due to Wocjan and Elphick [22] that $1 + s^+/s^- \leq \chi$ and consequently $s^+ \leq 2m(\chi - 1)/\chi$. It is clear that $\psi(\psi - 1) \leq 2m$. Therefore:

$$s^+ \leq \frac{2m(\chi - 1)}{\chi} \leq \frac{2m(\psi - 1)}{\psi} \leq \frac{4m^2}{\psi^2}.$$

Taking square roots and re-arranging completes the first half of the proof. Favaron *et al* [7] proved that $R(G) \geq m/\mu$. Therefore $\psi(G) \leq 2m/\mu \leq 2R(G)$. \square

Note that for regular graphs, $2m/\mu = 2R' = 2H = 2R = n$ whereas for almost all regular graphs $2m/\sqrt{s^+} < n$.

Lemma 3.2. *For all graphs, $col(col - 1) \leq 2m$.*

Proof. As noted above, $s^+ \leq 2m(\chi - 1)/\chi$ and $\chi(\chi - 1) \leq 2m$. Therefore:

$$\sqrt{s^+}(\sqrt{s^+} + 1) = s^+ + \sqrt{s^+} \leq \frac{2m(\chi - 1)}{\chi} + \frac{2m}{\chi} = 2m.$$

We can show that $deg(G) \leq \mu(G)$ as follows.

$$deg(G) = \max(\delta(H) : H \subseteq G) \leq \max(\mu(H) : H \subseteq G) \leq \mu(G).$$

Therefore $col(G) \leq \mu + 1$, so:

$$col(col - 1) \leq \mu(\mu + 1) \leq \sqrt{s^+}(\sqrt{s^+} + 1) \leq 2m.$$

\square

We can now prove the following theorem, using the same proof as for Theorem 3.1.

Theorem 3.3. *For a graph G , $col(G) \leq 2m/\sqrt{s^+} \leq 2m/\mu \leq 2R(G)$.*

3.3 Bounds for s^+

The proof of Lemma 3.2 uses that $s^+ + \sqrt{s^+} \leq 2m$, from which it follows that:

$$\sqrt{s^+} \leq \frac{1}{2}(\sqrt{8m + 1} - 1).$$

This strengthens Stanley's inequality [18] that:

$$\mu \leq \frac{1}{2}(\sqrt{8m + 1} - 1).$$

However, $\sqrt{2m - n + 1} \leq (\sqrt{8m + 1} - 1)/2$, and Hong [10] proved for graphs with no isolated vertices that $\mu \leq \sqrt{2m - n + 1}$. Elphick *et al* [5] recently conjectured that for connected graphs $s^+ \leq 2m - n + 1$, or equivalently that $s^- \geq n - 1$.

4 Example and Conjectures

A *Grundy k -coloring* of G is a k -coloring of G such that each vertex is colored by the smallest integer which has not appeared as a color of any of its neighbors. The *Grundy number* $\Gamma(G)$ is the largest integer k , for which there exists a Grundy k -coloring for G . It is clear that for any graph G ,

$$\Gamma(G) \leq \psi(G) \text{ and } \chi(G) \leq \Gamma(G) \leq \Delta(G) + 1. \quad (6)$$

Note that each pair of $\text{col}(G)$ and $\psi(G)$, or $\text{col}(G)$ and $\Gamma(G)$, or $\psi(G)$ and $\Delta(G)$ is incomparable in general.

As an example of the bounds discussed in this paper, if $G = P_4$, then

$$\chi(G) = \text{col}(G) = 2$$

$$\Gamma(G) = \psi(G) = \Delta(G) + 1 = 2R'(G) = 3$$

$$2H(G) = 3.67 \text{ and } 2R(G) = 3.83$$

$$\mu_1(G) = 1.618 \text{ and } \mu_2 = 0.618$$

$$2m/\mu_1(G) = 3.71 \text{ and } 2m/\sqrt{s^+} = 3.46.$$

We believe the following conjectures to be true.

Conjecture 4.1. *For any graph G , $\psi(G) \leq 2R'(G)$.*

In view of (6), a more tractable conjecture than the above is as follows.

Conjecture 4.2. *For any graph G , $\Gamma(G) \leq 2R'(G)$.*

References

- [1] T. Ando and M. Lin, *Proof of a conjectured lower bound on the chromatic number of a graph*, Linear Algebra and Appl. 485 (2015) 480-484.
- [2] H. Deng, S. Balachandran, S.K. Ayyaswamy and Y.B. Venkatakrisnan, *On the harmonic index and the chromatic number of a graph*, Discrete Appl. Math. 16 (2013) 2740-2744.
- [3] T.R. Divnić and L.R. Pavlović, *Proof of the first part of the conjecture of Aouchiche and Hansen about the Randić*, Discrete Appl. Math. 161 (2013) 953-960.
- [4] Z. Dvořák, B. Lidický and R. Škrekovski, *Randić index and the diameter of a graph*, European J. Combin. 32 (2011) 434-342.

- [5] C. Elphick, M. Farber, F. Goldberg and P. Wocjan, *Conjectured bounds for the sum of squares of positive eigenvalues of a graph*, (2015), math arXiv : 1409.2079v2.
- [6] P. Erdős, A. Rubin, and H. Taylor, *Choosability in graphs*, Congr. Num. 26 (1979) 125-157.
- [7] O. Favaron, M. Maheo and J. -F. Sacle, *Some eigenvalue properties in Graphs (conjectures of Graffiti II)*, Discrete Maths. 111 (1993) 197-220.
- [8] S. Fajtlowicz, *On conjectures of Graffiti-II*, Conr. Numer. 60 (1987) 187-197.
- [9] P. Hansen and D. Vukicević, *On the Randić index and the chromatic number*, Discrete Math. 309 (2009) 4228-4234.
- [10] Y. Hong, *Bounds on eigenvalues of graphs*, Discrete Math., 123, (1993), 65 - 74.
- [11] T.R. Jensen and B. Toft, *Graph Coloring Problems*, A Wiley-Interscience Publication, Jhon Wiley and Sons, Inc, New York, 1995.
- [12] X. Li and I. Gutman, *Mathematical Aspects of Randić-Type Molecular Structure Descriptors*, Mathematical Chemistry Monographs No. 1, Kragujevac, 2006.
- [13] J. Liu, M. Liang, B. Cheng and B. Liu, *A proof for a conjecture on the Randić index of graphs with diameter*, Appl. Math. Lett. 24 (2011) 752-756.
- [14] B. Liu, L.R. Pavlović, T.R. Divnić, J. Liu and M.M. Stojanović, *On the conjecture of Aouchiche and Hansen about the Randić index*, Discrete Math. 313 (2013) 225-235.
- [15] X. Li and Y. Shi, *On a relation between the Randić index and the chromatic number*, Discrete Math. 310 (2010) 2448-2451.
- [16] M. Knor, B. Kuzar and R. Skrekovski, *Sandwiching the (generalized) Randić index*, Discrete Appl. Math. 181 (2015) 160-166.
- [17] M. Randić, *On characterization of molecular branching*, J. Amer. Chem. Soc. 97 (1975) 6609-6615.
- [18] R. P. Stanley, *A bound on the spectral radius of graphs with e edges*, Linear Algebra Appl., 87, (1987), 267 - 269.
- [19] Z. Tang, B. Wu and L. Hu, *The Grundy number of a graph*, math arXiv:1507.01080 (2015).
- [20] Z. Tuza, *Graph colorings with local constraints- a survey*, Discuss. Math. Graph Theory 17 (1997) 161-228.

- [21] V.G. Vizing, *Coloring the vertices of a graph in prescribed colors* (Russian), Diskret. Analiz. 29 (1976) 3-10.
- [22] P. Wocjan and C. Elphick, *New spectral bounds on the chromatic number encompassing all eigenvalues of the adjacency matrix*, Electron. J. Combin. 20(3) (2013), P39.
- [23] B. Wu, J. Yan and X. Yang, *Randic index and coloring number of a graph*, Discrete Appl. Math. 178 (2014) 163 - 165.