Needles and straw in a haystack: robust confidence for possibly sparse sequences

Eduard Belitser and Nurzhan Nurushev VU Amsterdam and University of Amsterdam

Abstract

In the general $signal + noise$ (allowing non-normal, non-independent observations) model we construct an empirical Bayes posterior which we then use for *uncertainty quantification* for the unknown, possibly sparse, signal. We introduce a novel excessive bias restriction (EBR) condition, which gives rise to a new slicing of the entire space that is suitable for uncertainty quantification. Under EBR and some mild exchangeable exponential moment condition on the noise, we establish the local (oracle) optimality of the proposed confidence ball. Without EBR, we derive the full coverage for confidence balls of at least $\sigma n^{1/4}$ -radius, implying the local optimality only for cases when the oracle rate is at least of the order $\sigma n^{1/4}$. In passing, we also get the local optimal results for estimation and posterior contraction problems. Adaptive minimax results (also for the estimation and posterior contraction problems) over various sparsity classes follow from our local results.

1 Introduction

The model and the main problem. Suppose we observe $X = X^{(\sigma,n)} = (X_1, \ldots, X_n)$:

$$
X_i = \theta_i + \sigma \xi_i, \quad i \in [n] = \{1, ..., n\},
$$
 (1)

where $\theta = (\theta_1, \dots, \theta_n) \in \mathbb{R}^n$ is an unknown high-dimensional parameter of interest, the ξ_i 's are random errors with $E\xi_i = 0$, $Var(\xi_i) \leq C_{\xi}$, $\sigma > 0$ is the known noise intensity. The goal is to make inference about the parameter θ based on the data X: recovery of θ and uncertainty quantification by constructing an optimal confidence set. We pursue robust inference in the sense that the distribution of the error vector $\xi = (\xi_1, \ldots, \xi_n)$ is unknown and can also depend on θ , but assumed to satisfy only certain mild *exchangeable exponential moment condition*; see Condition [\(A1\)](#page-7-0) in Section [2.](#page-4-0) For inference on θ , we exploit the empirical Bayes approach. We derive non-asymptotic results, which imply asymptotic assertions as well if needed. Possible asymptotic regimes are decreasing noise level $\sigma \to 0$, high-dimensional setup $n \to \infty$ (the leading case for high dimensional models), or their combination, e.g., $\sigma = n^{-1/2}$ and $n \to \infty$.

Useful inference is not possible without some structure on the parameter θ . Popular structural assumptions are smoothness and sparsity, in this paper we are concerned with the latter.

MSC2010 subject classification: primary 62G15, secondary 62C12.

Keywords and phrases: confidence set, empirical Bayes posterior, local radial rate.

The best studied problem in the sparsity context is that of estimating θ in the many normal means model, a variety of estimation methods and results are available in the literature: [\[16\]](#page-34-0), [\[6\]](#page-34-1), [\[17\]](#page-34-2), [\[1\]](#page-33-0), [\[13\]](#page-34-3), [\[27\]](#page-35-0). However, even an optimal estimator does not reveal how far it is from θ . It is of importance to quantify this uncertainty, which can be seen as the problem of constructing confidence sets for θ .

Bayesian approach and accompanying posterior contraction problem. Many inference methods have Bayesian connections. For example, even some seemingly non Bayesian estimators can be obtained as certain quantities (like posterior mode for penalized minimum contrast estimators) of the (empirical Bayes) posterior distributions resulting from imposing some specific priors on the parameter; cf. [\[17\]](#page-34-2) and [\[1\]](#page-33-0). Although the Bayesian methodology is used or can be related to in constructing many (frequentist) inference procedures, only recently the posterior distributions themselves have been studied in the sparsity context: [\[13\]](#page-34-3), [\[27\]](#page-35-0), [\[19\]](#page-35-1), [\[12\]](#page-34-4), [\[7\]](#page-34-5), [\[25\]](#page-35-2), [\[23\]](#page-35-3).

In this paper, for inference on θ we use an empirical Bayes approach. Since any Bayesian approach always delivers a posterior $\pi(\vartheta|X)$ (in the posteriors for θ , we will use the variable ϑ to distinguish it from the "true" θ), an accompanying problem of interest is the contraction of the resulting (empirical Bayes) posterior to the "true" θ from the frequentist perspective of the "true" measure P_{θ} , the distribution of X from [\(1\)](#page-0-0). The quality of posterior is characterized by the posterior contraction rate. We pursue a novel local approach by allowing the posterior contraction rate to be a local quantity, i.e., depending on the true θ , whereas global minimax rates are typically studied in the literature on Bayesian nonparametrics.

A common Bayesian way to model sparsity structure is by the so called two-groups priors. Such a prior puts positive mass on vectors θ with some exact zero coordinates (zero group) and the remaining coordinates (signal group) are drawn from a chosen distribution. So the marginal prior for each coordinate is a mixture of a continuous distribution and a point-mass at zero. In [\[13\]](#page-34-3) it is shown that for a suitably chosen two-groups prior, the posterior concentrates around the true θ at the minimax rate (as $n \to \infty$) for two sparsity classes, nearly black vectors $\ell[p_n]$ with p_n nonzero coordinates and weak ℓ_s -balls $m_s[p_n]$. As pointed out by [\[13\]](#page-34-3) (also by [\[17\]](#page-34-2)), the distributions of non-zero coordinates should not have too light tails, otherwise one gets suboptimal rates. The important Gaussian case is for example excluded. This has to do with the so called *over-shrinkage effect* of the normal prior with a fixed mean for nonzero coordinates, which pushes the posterior too much towards the prior mean, missing the true parameter that in general differs from the prior mean. That is why [\[17\]](#page-34-2) and [\[13\]](#page-34-3) discard normal priors on non-zero coordinates and use heavy tailed priors. A way to construct such a prior is to put a next level heavy-tailed prior, like half-Cauchy, on the variance in the normal prior, resulting in the so called (one-component) horseshoe prior on θ (cf. [\[11\]](#page-34-6) and [\[27\]](#page-35-0)). In the present paper we show that normal priors are still usable (cf. [\[19\]](#page-35-1)) and even lead to strong local results (and even for non-normal models) if combined with empirical Bayes approach.

Uncertainty quantification problem. The main aim in this paper is to construct confidence sets with optimal properties. The size of a confidence set is measured by the smallest radius of a ball containing this set, hence it suffices to consider confidence balls. For the usual norm

 $\|\cdot\|$ in \mathbb{R}^n , a random ball in \mathbb{R}^n is $B(\hat{\theta}, \hat{r}) = \{\theta \in \mathbb{R}^n : \|\hat{\theta} - \theta\| \leq \hat{r}\}\$, where the center $\hat{\theta} = \hat{\theta}(X) : \mathbb{R}^n \to \mathbb{R}^n$ and radius $\hat{r} = \hat{r}(X) : \mathbb{R}^n \to \mathbb{R}_+ = [0, +\infty]$ are measurable functions of the data X . Let us introduce the optimality framework for uncertainty quantification. The goal is to construct such a confidence ball $B(\hat{\theta}, C\hat{r})$ that for any $\alpha_1, \alpha_2 \in (0, 1]$ and some functional $r(\theta) = r_{\sigma,n}(\theta), r : \mathbb{R}^n \to \mathbb{R}_+$, there exist $C, c > 0$ such that

$$
\sup_{\theta \in \Theta_0} P_{\theta} \big(\theta \notin B(\hat{\theta}, C\hat{r}) \big) \le \alpha_1, \quad \sup_{\theta \in \Theta_1} P_{\theta} \big(\hat{r} \ge cr(\theta) \big) \le \alpha_2,
$$
\n(2)

for some $\Theta_0, \Theta_1 \subseteq \mathbb{R}^n$. The function $r(\theta)$, called the *radial rate*, is a benchmark for the effective radius of the confidence ball $B(\hat{\theta}, C\hat{r})$. The first expression in [\(2\)](#page-2-0) is called *coverage relation* and the second size relation. Notice that our approach is local (and hence genuinely adaptive) as the radial rate $r(\theta)$ is a function of the "true" parameter θ . Recall the common (global) minimax adaptive version of [\(2\)](#page-2-0): given a family of sets Θ_β with corresponding minimax estimation rates $r(\Theta_\beta)$ indexed by a structural parameter $\beta \in \mathcal{B}$ (e.g., smoothness or sparsity), the minimax adaptive version of [\(2\)](#page-2-0) would be obtained by taking $\Theta_0 = \Theta_1 = \Theta_\beta$ and the radial rate $r(\theta)$ $r(\Theta_{\beta})$ for all $\theta \in \Theta_{\beta}$ and all $\beta \in \mathcal{B}$.

Coming back to our local framework [\(2\)](#page-2-0), it is desirable to find the smallest $r(\theta)$ and the biggest Θ_0 , Θ_1 , for which [\(2\)](#page-2-0) holds. These are contrary requirements, so we have to trade them off against each other. There are different ways of doing this, leading to different optimality frameworks. For example, if we insist on overall uniformity $\Theta_0 = \mathbb{R}^n$, then the results in [\[18\]](#page-34-7) and [\[10\]](#page-34-8) (more refined versions are in [\[3\]](#page-34-9) and [\[20\]](#page-35-4)) say basically that the radial rate r cannot be of a faster order than $\sigma n^{1/4}$ for every θ and is at least of order $\sigma n^{1/2}$ for some θ . This means that any confidence ball that is optimal with respect to the optimality framework [\(2\)](#page-2-0) with $\Theta_0 = \mathbb{R}^n$ will necessarily have a big size, even if the true θ happens to lie in a very "good", smooth or sparse, class Θ_1 . Many good confidence sets cannot be optimal in this sense (called "honest" in some papers) and effectively excluded from the consideration. For minimax adaptive versions of [\(2\)](#page-2-0) this means that as soon as we require $\Theta_0 = \Theta_\beta$, $\beta \in \mathcal{B}$ in the coverage relation, the minimax rate $r(\Theta_\beta)$ in the size relation is unattainable even for $\beta \in \mathcal{B} = {\beta_1, \beta_2}$; cf. [\[20\]](#page-35-4) for two nearly black classes. Essentially, the overall uniform coverage and optimal size properties can not hold together, it is necessary to sacrifice at least one of these, preferably as little as possible. We argue that it is unreasonable to pursue an optimality framework with the entire space $\Theta_0 = \mathbb{R}^n$ in the coverage relation, because this leads to discarding many good procedures and optimality of uninteresting ones. Instead, it makes sense to sacrifice in the set $\Theta_0 = \mathbb{R}^n \backslash \Theta'$, by removing a preferably small portion of "deceptive parameters" Θ' from \mathbb{R}^n so that that the optimal radial rates become attainable in the size relation with interesting (preferably "massive") sets Θ_1 .

This "deceptiveness" phenomenon is well understood for some smoothness structures (e.g., Sobolev scale), especially in global minimax settings; see [\[22\]](#page-35-5), [\[9\]](#page-34-10), [\[4\]](#page-34-11) and [\[26\]](#page-35-6). If we now insist on the optimal size property in [\(2\)](#page-2-0) for all Θ_{β} , $\beta \in \mathcal{B}$, the coverage relation in (2) will not hold for all $\Theta_0 = \Theta_\beta$, but only for $\Theta_0 = \Theta_\beta \backslash \Theta'$, with some set of "deceptive parameters" Θ' removed from Θ_{β} . In [\[26\]](#page-35-6) such parameters are called "inconvenient truths" and an implicit construction of a $\theta' \in \Theta'$ is given. Examples of non-deceptive parameters are the set of selfsimilar parameters $\Theta_0 = \Theta_{ss}$ introduced by [\[21\]](#page-35-7) and studied by [\[8\]](#page-34-12), [\[9\]](#page-34-10), [\[26\]](#page-35-6), and the set of polished tail parameters $\Theta_0 = \Theta_{pt}$ considered by [\[26\]](#page-35-6). In all the above mentioned papers global minimax radial rates (i.e., $r(\theta) = r(\Theta_{\beta})$ for all $\theta \in \Theta_{\beta}$) for specific smoothness structures are

studied. A local approach, delivering also the adaptive minimax results for many smoothness structures simultaneously, is considered by [\[2\]](#page-34-13) for posterior contraction rates and by [\[4\]](#page-34-11) for constructing optimal confidence balls. In [\[4\]](#page-34-11), yet a more general (than Θ_{ss} and Θ_{pt}) set of nondeceptive parameters was introduced, $\Theta_0 = \Theta_{ebr}$, parameters satisfying the so called *excessive* bias restriction (EBR). More on this can be found in Section [4.2.](#page-22-0)

To the best of our knowledge, there are very few papers about adaptive results on uncertainty quantification [\(2\)](#page-2-0). The case of two nearly black classes is treated by [\[20\]](#page-35-4), the "general polished tail" condition was introduced in [\[24\]](#page-35-8) to describe non-deceptive parameters, a restricted scale of nearly black classes is treated in [\[28\]](#page-35-9), where a version of our EBR condition is used, more on relation to paper [\[28\]](#page-35-9) can be found in Section [4.1.](#page-18-0)

The scope of this paper. In this paper, we introduce a family of normal mixture priors and propose an empirical Bayes procedure (in fact, two procedures). We use the normal likelihood, whereas the true model [\(1\)](#page-0-0) does not have to be normal (and independence of ξ_i 's is not required either), but only satisfying some mild Condition (A1) (called *exchangeable exponential moment* condition). There are three distinctive features of our approach: *robust*, *local* and *refined*.

First, robust means that our results cover also misspecified models, as we allow the ξ_i 's to be not necessarily independent normals (a certain type of error misspecification was also mentioned in a remark of the supplement to $[12]$, but only satisfying Condition $(A1)$. It turned out that, although we use the normal likelihood in the Bayesian analysis, in the proof of the main results we can handle the frequentist behavior of the posterior from the perspective of the true measure only on the basis of Condition (A1).

Second, we develop the novel *local* approach, meaning that the radial rate $r(\theta)$ in [\(2\)](#page-2-0) is allowed to be a function of θ , which, in a way, measures the amount of sparsity for each $\theta \in \mathbb{R}^n$: the smaller $r(\theta)$, the more sparse θ . The local radial rate $r(\theta)$ is constructed as the best (smallest) rate over a certain family of local rates, therefore called *oracle rate*. We demonstrate that the local approach is more powerful than global in that we do not need to impose any specific sparsity structure, because the proposed local approach automatically exploits the "effective" sparsity of each underlying θ , and our local results imply a whole panorama of the global minimax results for many scales at once. More on this is in Section [3.5.](#page-16-0)

Third, we derive the local posterior contraction result for the resulting empirical Bayes posterior $\hat{\pi}(\vartheta|X)$ in the refined non-asymptotic formulation: $\sup_{\theta \in \mathbb{R}^n} E_{\theta} \hat{\pi}(\|\vartheta - \theta\|^2 \ge M_0 r^2(\theta) +$ $M\sigma^2|X) \leq H_0e^{-m_0M}$ for some fixed $M_0, H_0, m_0 > 0$ and arbitrary $M \geq 0$, as an exponential nonasymptotic concentration bound in terms of M, uniformly in $\theta \in \mathbb{R}^n$. This formulation provides a rather subtile characterization of the quality of the posterior (finer, than, e.g., asymptotically in terms of the dimension n), allowing subtle analysis for various asymptotic regimes. This result is of interest and importance on its own as it actually establishes the contraction of the empirical Bayes posterior with the local rate $r(\theta)$. Besides, we obtain the oracle estimation result (also in similar refined formulation, finer than traditional oracle inequalities) by constructing an estimator, the empirical Bayes posterior mean, which converges to θ with the local rate $r(\theta)$. This result, besides being an ingredient for the uncertainty quantification problem [\(2\)](#page-2-0), is also of interest and importance on its own as it delivers the same (oracle and minimax) estimation results as in [\[1\]](#page-33-0) and [\[17\]](#page-34-2) and posterior convergence results as in [\[13\]](#page-34-3), obtained for different priors.

Next, we construct a confidence ball by using the empirical Bayes posterior quantities. Since we want the size of our confidence sets to be of an oracle rate order, this comes with the price that the coverage property can hold uniformly only over some set of parameters satisfying the so called *excessive bias restriction* (EBR) $\Theta_0 = \Theta_{eb} \subseteq \mathbb{R}^n$. The main result consists in establishing the optimality [\(2\)](#page-2-0) of the constructed confidence ball for the optimality framework $\Theta_0 = \Theta_{eb}$, $\Theta_1 = \mathbb{R}^n$ and the local radial rate $r(\theta)$. The important consequence of our local approach is that a whole panorama of adaptive (global) minimax results (for all the three problems: estimation, posterior contraction rate and confidence sets) over all sparsity scales covered by $r(\theta)$ (see Section [3.5\)](#page-16-0) follow from our local results. In particular, our local results imply the same type of adaptive minimax estimation results over sparsity scales as in [\[17\]](#page-34-2), and the same type of global minimax results on contraction posterior rates as in [\[13\]](#page-34-3) (and actually more).

Also we treat the situation when $\Theta_0 = \mathbb{R}^n$ in [\(2\)](#page-2-0) by constructing a confidence ball such that its radius is of the order $\sigma n^{1/4} + r(\theta)$. As we already discussed, the term $\sigma n^{1/4}$ in the size relation is necessary for the uniform coverage to hold. Clearly, this confidence ball will have optimal size only for non-sparse parameters (for which $r(\theta) \geq c \sigma n^{1/4}$).

Although the original motivation of the EBR condition was to remove the deceptive parameters, it turned out to be a very useful notion in the context of uncertainty quantification. In effect, the EBR condition leads to a *new sparsity EBR-scale* which gives the slicing of the entire space that is very suitable for uncertainty quantification. This provides a new perspective at the above mentioned "deceptiveness" issue: basically, each parameter is deceptive (or non deceptive) to some extent. It is the structural parameter of the new EBR-scale that measures the deceptiveness amount, and the (mild and controllable) price for handling deceptive parameters is the effective amount of inflating of the confidence ball that matches the amount of deceptiveness needed to provide a high coverage. More on the EBR condition can be found in Section [4.2.](#page-22-0)

The paper is organized as follows. In Section [2](#page-4-0) we introduce the notation, the prior, describe the empirical Bayes procedure in detail, make a link with the penalization method, and provide some conditions. Section [3,](#page-8-0) where we also introduce the EBR, contains the main results of the paper. In Section [4,](#page-18-1) we discuss some variations of our results, present concluding remarks and discuss the EBR. The theoretical results are illustrated in Section [5](#page-23-0) by a small simulation study. The proofs of the lemmas and theorems are given in Sections [6](#page-24-0) and [7](#page-27-0) respectively.

2 Preliminaries

First we introduce some notation and a certain family of normal priors (similar to priors from [\[4\]](#page-34-11) but geard towards modeling sparsity rather than smoothness). Next, by applying the empirical Bayes approach to the the normal likelihood, we derive an empirical Bayes posterior which we will use in the construction of the estimator and the confidence ball. The empirical Bayes procedure is linked to the penalization method. We complete this section with some conditions on the errors ξ_i 's and the prior.

2.1 Notation

Denote the probability measure of X from the model [\(1\)](#page-0-0) by $P_{\theta} = P_{\theta}^{(\sigma,n)}$, and by E_{θ} the corresponding expectation. For notational simplicity, we often skip the dependence on σ and n.

Denote by $\mathbb{1}_E = \mathbb{1}{E}$ the indicator function of the event E, by |S| the cardinality of the set S, the difference of sets $S\setminus S_0 = \{s \in S : s \notin S_0\}$. Let $[k] = \{1, \ldots, k\}$ and $[k]_0 = \{0\} \cup [k]$ for $k \in \mathbb{N} = \{1, 2, \ldots\}$. For $I \subseteq [n]$, define $I^c = [n] \setminus I$. Let $\mathcal{I} = \mathcal{I}_n = \{I : I \subseteq [n]\}$ be the family of all subsets of $[n]$ including the empty set. If the summation range in \sum_I is not specified (for brevity), this means $\sum_{I\in\mathcal{I}}$. Throughout we assume the conventions that $\sum_{i\in\mathcal{I}} a_i = 0$, $\sum_{a}^{b} a_i = \sum_{a \leq i \leq b} a_i$ for any $a_i, a, b \in \mathbb{R}$ and $0 \log(c/0) = 0$ (hence $(c/0)^0 = 1$) for any $c > 0$. Let $\theta_{(1)}^2 \leq \theta_{(2)}^2 \leq \ldots \leq \theta_{(n)}^2$ and $\theta_{[1]}^2 \geq \theta_{[2]}^2 \geq \ldots \geq \theta_{[n]}^2$ be the ordered values of $\theta_1^2, \ldots, \theta_n^2$. To have some quantity well defined in the sequel, introduce also $0 = \theta_{(0)}^2 = \theta_{[n+1]}^2$ and $\theta_{[0]}^2 = \theta_{(n+1)}^2 = \infty$.

If random quantities appear in a relation, this relation should be understood in P_{θ} -almost sure sense. Throughout $\phi(x, \mu, \sigma^2)$ will be the density of $\mu + \sigma Z \sim N(\mu, \sigma^2)$ at point x, where $Z \sim N(0, 1)$. By convention, $N(\mu, 0) = \delta_{\mu}$ denotes a Dirac measure at point μ . The symbol \triangleq will refer to equality by definition, $(a \vee b) = \max\{a, b\}$ and $(a \wedge b) = \min\{a, b\}$. Finally, denote $X(I) = (X_i 1\{i \in I\}, i \in \mathbb{N}_n)$ for $I \in \mathcal{I}$, and let $\langle x, y \rangle = \sum_i x_i y_i$ denote the usual scalar product between $x, y \in \mathbb{R}^n$.

2.2 Multivariate normal prior

When deriving all the posterior quantities in the Bayesian analysis below, we will use the normal likelihood $\ell(\theta, X) = (2\pi\sigma^2)^{-n/2} \exp\{-\|X-\theta\|^2/2\sigma^2\}$, which is equivalent to imposing the classical high-dimensional normal model $X = (X_i, i \in \mathbb{N}_n) \sim \bigotimes_{i=1}^n \mathbb{N}(\theta_i, \sigma^2)$. Recall however that the "true" model $X \sim P_{\theta}$ is not assumed to be normal, but satisfying Condition [\(A1\)](#page-7-0).

To model possible sparsity in the parameter θ , the coordinates of θ can be split into two distinct groups of coordinates of θ : for some $I \in \mathcal{I}$, $\theta_I = (\theta_i, i \in I)$ and $\theta_{I^c} = (\theta_i, i \in I^c)$, so that $\theta = (\theta_I, \theta_{I^c})$. The group of coordinates $\theta_{I^c} = (\theta_i, i \notin I)$ consists of (almost) zeros and $\theta_I = (\theta_i, i \in I)$ is the group of non-zeros coordinates. For any $\theta \in \mathbb{R}^n$ (even "not sparse" one) there is the best (oracle) splitting into two groups, we will come back to this in Section [3.](#page-8-0) To model sparsity, we propose a prior on θ given I as follows:

$$
\pi_I = \bigotimes_{i=1}^n \mathcal{N}(\mu_i(I), \tau_i^2(I)), \quad \mu_i(I) = \mu_i \mathbb{1}\{i \in I\}, \quad \tau_i^2(I) = \sigma^2 K_n(I) \mathbb{1}\{i \in I\},\tag{3}
$$

and $K_n(I) = \left(\frac{en}{|I|} - 1\right) \mathbb{1}\left\{I \neq \emptyset\right\}$. The indicators in prior [\(3\)](#page-5-0) ensure the sparsity of the group I^c . The rather specific choice of $K_n(I)$ is made for the sake of concise expressions in later calculations, many other choices are actually possible. By using normal likelihood $\ell(\theta, X) = (2\pi\sigma^2)^{-n/2} \exp\{-\|X - \theta\|^2/2\sigma^2\},\$ the corresponding posterior distribution for θ is readily obtained:

$$
\pi_I(\vartheta|X) = \bigotimes_{i=1}^n \mathcal{N}\Big(\frac{\tau_i^2(I)X_i + \sigma^2 \mu_i(I)}{\tau_i^2(I) + \sigma^2}, \frac{\tau_i^2(I)\sigma^2}{\tau_i^2(I) + \sigma^2}\Big). \tag{4}
$$

Next, introduce the prior λ on I, discussed in Section [4.1](#page-18-0) below. For $\varkappa > 1$, draw a random set from $\mathcal I$ with probabilities

$$
\lambda_I = c_{\varkappa,n} \exp\left\{-\varkappa |I| \log\left(\frac{en}{|I|}\right)\right\} = c_{\varkappa,n} \left(\frac{en}{|I|}\right)^{-\varkappa |I|}, \quad I \in \mathcal{I},\tag{5}
$$

where $c_{\varkappa,n}$ is the normalizing constant. Since $(\frac{n}{k})^k \leq {n \choose k}$ $\binom{n}{k} \leq \left(\frac{en}{k}\right)$ $(\frac{2n}{k})^k$ and $(\frac{n}{0})^k$ $\binom{n}{0} = 1,$

$$
1 = \sum_{I \in \mathcal{I}} \lambda_I = c_{\varkappa, n} \sum_{k=0}^n \binom{n}{k} \left(\frac{en}{k}\right)^{-\varkappa k} \le c_{\varkappa, n} \sum_{k=0}^n \left(\frac{en}{k}\right)^{-(\varkappa - 1)k} \le c_{\varkappa, n} \sum_{k=0}^n e^{-(\varkappa - 1)k},\tag{6}
$$

so that $c_{\varkappa,n} \geq 1 - e^{1-\varkappa} > 0$, $n \in \mathbb{N}$. Combining [\(3\)](#page-5-0) and [\(5\)](#page-5-1) gives the mixture prior on θ : $\pi = \sum_{I \in \mathcal{I}} \lambda_I \pi_I$. This leads to the marginal distribution of X: $P_X = \sum_{I \in \mathcal{I}} \lambda_I P_{X,I}$, with $P_{X,I} = \overline{\bigotimes_{i=1}^{n}} \mathbb{N}(\mu_i(I), \sigma^2 + \tau_i^2(I)),$ and the posterior of θ is

$$
\pi(\vartheta|X) = \pi_{\varkappa}(\vartheta|X) = \sum_{I \in \mathcal{I}} \pi_I(\vartheta|X)\pi(I|X),\tag{7}
$$

where $\pi_I(\vartheta|X)$ is defined by [\(4\)](#page-5-2) and the posterior $\pi(I|X)$ for I is

$$
\pi(I|X) = \frac{\lambda_I P_{X,I}}{P_X} = \frac{\lambda_I \prod_{i=1}^n \phi(X_i, \mu_i(I), \sigma^2 + \tau_i^2(I))}{\sum_{J \in \mathcal{I}} \lambda_J \prod_{i=1}^n \phi(X_i, \mu_i(J), \sigma^2 + \tau_i^2(J))}.
$$
\n(8)

2.3 Empirical Bayes posterior

The parameters $\mu_{1,i}$ are yet to be chosen in the prior. We choose μ_i by using empirical Bayes approach. The marginal likelihood P_X is readily maximized with respect to μ_i : $\tilde{\mu}_i = X_i$, which we then substitute instead of μ_i in the expression [\(7\)](#page-6-0) for $\pi(\theta|X)$, obtaining the empirical Bayes posterior

$$
\tilde{\pi}(\vartheta|X) = \tilde{\pi}_{\varkappa}(\vartheta|X) = \sum_{I \in \mathcal{I}} \tilde{\pi}_{I}(\vartheta|X)\tilde{\pi}(I|X),\tag{9}
$$

where the empirical Bayes conditional posterior (recall that $N(0, 0) = \delta_0$)

$$
\tilde{\pi}_I(\vartheta|X) = \bigotimes_{i=1}^n \mathcal{N}\big(X_i \mathbb{1}\{i \in I\}, \frac{K_n(I)\sigma^2 \mathbb{1}\{i \in I\}}{K_n(I)+1}\big) \tag{10}
$$

is obtained from [\(4\)](#page-5-2) with $\mu_{1,i} = X_i$, and

$$
\tilde{\pi}(I|X) = \frac{\lambda_I P_{X,I}}{\sum_{J \in \mathcal{I}} \lambda_J P_{X,J}} = \frac{\lambda_I \prod_{i=1}^n \phi(X_i, X_i \mathbb{1}\{i \in I\}, \sigma^2 + \tau_i^2(I))}{\sum_{J \in \mathcal{I}} \lambda_J \prod_{i=1}^n \phi(X_i, X_i \mathbb{1}\{i \in J\}, \sigma^2 + \tau_i^2(J))}
$$
(11)

is the empirical Bayes posterior for $I \in \mathcal{I}$, obtained from [\(8\)](#page-6-1) with $\mu_i = X_i$. Let \tilde{E} and \tilde{E}_I be the expectations with respect to the measures $\tilde{\pi}(\vartheta|X)$ and $\tilde{\pi}_I(\vartheta|X)$ respectively. Then $\tilde{E}_I(\vartheta|X)$ = $X(I) = (X_i 1\{i \in I\}, i \in [n])$. Introduce the *empirical Bayes posterior mean* estimator

$$
\tilde{\theta} = \tilde{\mathcal{E}}(\theta|X) = \sum_{I \in \mathcal{I}} \tilde{\mathcal{E}}_I(\theta|X)\tilde{\pi}(I|X) = \sum_{I \in \mathcal{I}} X(I)\tilde{\pi}(I|X).
$$
\n(12)

Consider an alternative empirical Bayes posterior. First derive an empirical Bayes variable selector \hat{I} by maximizing $\tilde{\pi}(I|X)$ over $I \in \mathcal{I}$ (any maximizer will do) as follows:

$$
\hat{I} = \underset{I \in \mathcal{I}}{\operatorname{argmax}} \tilde{\pi}(I|X) = \underset{I \in \mathcal{I}}{\operatorname{argmax}} \lambda_I P_{X,I} = \underset{I \in \mathcal{I}}{\operatorname{argmax}} \left\{ -\sum_{i \in I^c} \frac{X_i^2}{2\sigma^2} - \frac{|I|}{2} \log(K_n(I) + 1) + \log \lambda_I \right\}
$$
\n
$$
= \underset{I \in \mathcal{I}}{\operatorname{argmin}} \left\{ \sum_{i \in I^c} X_i^2 + (2\kappa + 1)\sigma^2 |I| \log\left(\frac{en}{|I|}\right) \right\},\tag{13}
$$

which is reminiscent of the penalization procedure from [\[6\]](#page-34-1) (cf. also [\[1\]](#page-33-0)). Now plugging in \hat{I} into $\tilde{\pi}_I(\vartheta|X)$ defined by [\(10\)](#page-6-2) yields another empirical (now with respect to μ_i 's and I) Bayes posterior and the corresponding empirical Bayes mean estimator for θ :

$$
\check{\pi}(\vartheta|X) = \tilde{\pi}_{\hat{I}}(\vartheta|X), \quad \check{\theta} = \check{E}(\vartheta|X) = X(\hat{I}) = (X_i 1\{i \in \hat{I}\}, i \in [n]),\tag{14}
$$

where E denotes the expectation with respect to the measure $\check{\pi}(\vartheta|X)$.

2.4 Conditions

We provide some technical conditions and definitions. The following condition (called *exchange*able exponential moment condition) on the error vector $\xi = (\xi_1, \ldots, \xi_n)$ will be assumed throughout.

CONDITION (A1). The random variables ξ_i 's from [\(1\)](#page-0-0) satisfy: $E\xi_i = 0$, $Var(\xi_i) \le C_\xi$, $i \in [n]$; and for some $\beta, B > 0$ (without loss of generality assume $C_{\xi} = 1$ and $\beta \in (0, 1]$),

$$
\mathcal{E} \exp\left\{\beta \sum_{i \in I} \xi_i^2\right\} \le e^{B|I|} \quad \text{for all } I \in \mathcal{I}.\tag{A1}
$$

Notice that the unknown distribution of ξ may also depend on θ , in that case we assume Condition (A1) to be fulfilled for all $\theta \in \mathbb{R}^n$. The constants $\beta \in (0,1]$ and $B > 0$ will be fixed in the sequel and we omit the dependence on these constants in all further notation. A short discussion about this condition can be found in Section [4.1.](#page-18-0) There is no need to assume $Var(\xi_i) \leq C_{\xi}$ as this follows from [\(A1\)](#page-7-0), but we provide this just for reader's convenience. Note also that the ξ_i 's do not have to be even independent. For example, ξ_i 's can follow an autoregressive model, see Section [4.1.](#page-18-0)

Condition [\(A1\)](#page-7-0) is clearly satisfied for independent normals $\xi_i \stackrel{\text{ind}}{\sim} N(0, 1)$ and for bounded (arbitrarily dependent) ξ_i 's. In case of independent normal errors, some bounds in the proofs can be sharpened; we will mention possible refinements in Section [4.1.](#page-18-0)

In the proof of Theorem [1](#page-10-0) below, we will need a bound for $E\left(\sum_{i\in I}\xi_i^2\right)^2$, $I \in \mathcal{I}$. Actually, Condition [\(A1\)](#page-7-0) ensures such a bound. Indeed, since $x^2 \le e^{2x}$ for all $x \ge 0$, by using the Hölder inequality and [\(A1\)](#page-7-0), we derive that for any $t \in (0, \beta]$

$$
\mathcal{E}_{\theta}\left(\sum_{i\in I}\xi_i^2\right)^2 = \frac{4}{t^2}\mathcal{E}_{\theta}\left(\frac{t}{2}\sum_{i\in I}\xi_i^2\right)^2 \le \frac{4}{t^2}\mathcal{E}_{\theta}e^{t\sum_{i\in I}\xi_i^2} \le \frac{4}{t^2}\big[\mathcal{E}e^{\beta\sum_{i\in I}\xi_i^2}\big]^{t/\beta} \le \frac{4}{t^2}e^{Bt\beta^{-1}|I|}.
$$

To summarize, Condition [\(A1\)](#page-7-0) implies that for any $\rho \in (0, B/2]$ and any $I \in \mathcal{I}$,

$$
E\left(\sum_{i \in I} \xi_i^2\right)^2 \le \frac{B^2}{(\beta \rho)^2} e^{2\rho|I|}.
$$
\n(15)

In some proofs we need a technical condition on the parameter \varkappa appearing in [\(5\)](#page-5-1). CONDITION (A2). The parameter \varkappa of the prior λ defined by [\(5\)](#page-5-1) satisfies

$$
\varkappa > \bar{\varkappa} \triangleq (12 - \beta + 4B)/(4\beta), \tag{A2}
$$

where β , B are from Condition [\(A1\)](#page-7-0).

Finally, we give one more technical definition which we will need in the claims. For constants β , B from Condition [\(A1\)](#page-7-0), define

$$
\bar{\tau} = \bar{\tau}(\varkappa, \beta, B) \triangleq 3(\varkappa \beta + \frac{\beta}{2} + B)/\beta.
$$
 (16)

3 Main results

In this section we give the main results of the paper. From now on, by $\hat{\pi}(\vartheta|X)$ (with corresponding expectation $E(\cdot|X)$) we denote either $\tilde{\pi}(\vartheta|X)$ defined by [\(9\)](#page-6-3) or $\tilde{\pi}(\vartheta|X)$ defined by [\(14\)](#page-7-1), and $\hat{\theta}$ will stand either for $\hat{\theta}$ defined by [\(12\)](#page-6-4) or for $\hat{\theta}$ defined by [\(14\)](#page-7-1). Also, $\hat{\pi}(I \in \mathcal{G}|X)$ should be read as $\tilde{\pi}(I \in \mathcal{G}|X)$ in case $\hat{\pi} = \tilde{\pi}$, and as $\mathbb{1}\{\hat{I} \in \mathcal{G}\}\$ in case $\hat{\pi} = \tilde{\pi}$, for all $\mathcal{G} \subseteq \mathcal{I}$ that appear in what follows. Hence, $\hat{\pi}(I|X) = \tilde{\pi}(I|X)$ and $E_{\theta}\hat{\pi}(I \in \mathcal{G}|X) = E_{\theta}\tilde{\pi}(I \in \mathcal{G}|X)$ in the former case, and $\hat{\pi}(I|X) = \mathbb{1}\{\hat{I} = I\}$ and $E_{\theta}\hat{\pi}(I \in \mathcal{G}|X) = P_{\theta}(\hat{I} \in \mathcal{G})$ in the latter case.

3.1 Oracle rate

The empirical Bayes posterior $\hat{\pi}(\vartheta|X)$ is a random mixture over $\tilde{\pi}_I(\vartheta|X)$, $I \in \mathcal{I}$. From the P_θ-perspective, each posterior $\tilde{\pi}_I(\vartheta|X)$ (and the corresponding estimator $\tilde{E}_I(\vartheta|X) = X(I)$) contracts to the true parameter θ with the local rate $R^2(I,\theta) = \sum_{i\in I^c} \theta_i^2 + \sigma^2|I|$. Indeed, since $\tilde{\mathrm{E}}_I(\vartheta|X) = X(I) = (X_i \mathbb{1}\{i \in I\}, i \in \mathbb{N}_n),$ [\(10\)](#page-6-2) and the Markov inequality yields

$$
\mathbf{E}_{\theta}\tilde{\pi}_I(\|\vartheta-\theta\|^2 \geq M^2R^2(I,\theta)|X) \leq \frac{\mathbf{E}_{\theta}\|X(I)-\theta\|^2 + \frac{K_n(I)\sigma^2|I|}{K_n(I)+1}}{M^2R^2(I,\theta)} \leq \frac{2}{M^2}.
$$

For each $\theta \in \mathbb{R}^n$, among $I \in \mathcal{I}$ there exists the best choice $I_o = I_o(\theta) = I_o(\theta, \sigma)$ (called the R-oracle) corresponding to the fastest local rate $R^2(\theta) = R^2(\theta, \mathcal{I}) = \min_{I \in \mathcal{I}} R^2(I, \theta) =$ $\sum_{i\in I_c^c} \theta_i^2 + \sigma^2 |I_o|$. Ideally, we would like to *mimic* the *R*-oracle, i.e., to construct an empirical Bayesian procedure (e.g., $\hat{\pi}(\vartheta|X)$) which performs as good as the oracle empirical Bayes posterior $\tilde{\pi}_{I_o}(\vartheta|X)$ without knowing I_o , uniformly in $\theta \in \mathbb{R}^n$. However, the lower bounds for the estimation problem (hence, also for the posterior contraction problem), obtained by [\[15\]](#page-34-14) and later by [\[6\]](#page-34-1), show that it is impossible to mimic the R -oracle and a logarithmic factor is the unavoidable price for the uniformity over \mathbb{R}^n (otherwise this would contradict to the minimax lower bound over the scale of sparsity classes, cf. [\[6\]](#page-34-1)). Therefore only a modification of the risk R-oracle where the variance term $\sigma^2|I_o|$ is inflated with the factor $\log(en/|I_o|)$ (thought of as payment for not knowing I_o) is "mimicable".

The above discussion motivates the following definition. Introduce the family of local rates

$$
r^{2}(I,\theta) = r_{\sigma}^{2}(I,\theta) = \sum_{i \in I^{c}} \theta_{i}^{2} + \sigma^{2}|I| \log(\frac{en}{|I|}) = B(I,\theta) + V(I), \quad I \in \mathcal{I},
$$
\n
$$
(17)
$$

where $B(I,\theta) = \sum_{i\in I^c} \theta_i^2$ is the bias part of the rate and $V(I) = V(I,\sigma,n) = \sigma^2|I|\log(\frac{en}{|I|})$ is the adjusted variance part, the variance term $\sigma^2|I|$ of the rate $R(I,\theta)$ multiplied by the logarithmic factor $\log(\frac{en}{|I|})$. There exists the best choice $I_o = I_o(\theta) = I_o(\theta, \sigma^2) = I_o(\theta, \sigma^2, n) \in \mathcal{I}$ (called oracle) at which the rate [\(17\)](#page-8-1) is minimal:

$$
r^{2}(\theta) = r^{2}(\theta, \mathcal{I}) = \min_{I \in \mathcal{I}} r^{2}(I, \theta) = r^{2}(I_{o}(\theta), \theta) = B(I_{o}(\theta), \theta) + V(I_{o}(\theta)),
$$
\n(18)

called the *oracle rate*. Note that the oracle I_0 may not be unique (but $|I_0|$ is unique) as some coordinates of θ can coincide, in that case take the one with the earliest coordinates. Clearly, $I_o = \{i \in [n] : \theta_i^2 \geq \theta_{[i_o]}^2\}$, where $i_o = |I_o| = \operatorname{argmin}_{k \in [n]_0} {\sum_{i=1}^{n-k} \theta_{(i)}^2 + \sigma^2 k \log(\frac{en}{k})}$. Thus the oracle I_o classifies the coordinates $(\theta_i, i \in I_o)$ as *significant* and the coordinates $(\theta_i, i \in I_o)$ I_o^c) as insignificant. The bias related term $B(I_o(\theta), \theta) = \sum_{i \in I_o^c} \theta_i^2 = \sum_{i=1}^{n-i_o} \theta_{(i)}^2$ of the oracle rate is called the *excessive bias*. This is the error the oracle makes when setting insignificant coordinates of θ to zero. The variance related term $\sigma^2 |I_o| \log(\frac{en}{|I_o|})$ is the error the oracle makes when recovering the significant coordinates (the log factor is the payment for not knowing the locations). The definition [\(18\)](#page-9-0) of the oracle I_0 implies the following characterization of the significant coordinates $\{\theta_{[i]}, i = 1, \ldots, i_o\}$:

$$
\theta_{[i_o]}^2 \ge \sigma^2 \Big[\log(\frac{en}{i_o}) - (i_o - 1) \log(\frac{i_o}{i_o - 1}) \Big],
$$

\n
$$
\theta_{[i_o]}^2 + \theta_{[i_o + 1]}^2 \ge \sigma^2 \Big[2 \log(\frac{en}{i_o}) - (i_o - 2) \log(\frac{i_o}{i_o - 2}) \Big],
$$

\n...,
\n
$$
\sum_{i=1}^{i_o} \theta_{[i]}^2 \ge \sigma^2 i_o \log(\frac{en}{i_o}).
$$
\n(19)

The insignificant coordinates $\{\theta_{(i)}, i = 1, \ldots, n - i_o\}$ can be characterized in a similar manner.

Introduce a family of the so called τ -oracles $I_o^{\tau} = I_o^{\tau}(\theta) = I_o(\theta, \tau \sigma^2), \tau \ge 0$ and let $i_{\tau} = |I_o^{\tau}(\theta)|$ be the corresponding cardinalities. A τ -oracle $I_o^{\tau}(\theta)$ is just the usual oracle defined by [\(18\)](#page-9-0) with σ² substituted by $\tau \sigma^2$, the oracle itself is the τ-oracle with $\tau = 1$: $I_o(\theta) = I_o^1(\theta)$. Notice that $I_{o}^{\tau_1} \subseteq I_{o}^{\tau_2}$ if $\tau_1 \geq \tau_2$. For $\tau \downarrow 0$, $r^2(\theta, I_{o}^{\tau}) \downarrow 0$ and the "limiting" τ -oracle recovers the active index set $I^* = I^*(\theta) = \{i \in [n]: \theta_i \neq 0\}$ in the sense that $I_o^{\tau} \uparrow I^*$ as $\tau \downarrow 0$. Informally, since the τ -oracle is defined by substituting $\tau \sigma^2$ instead of σ^2 in the oracle rate, one can think of the τ -oracle with $\tau \in [0, 1)$ as if X is observed with a "magnifying glass" since the error variance is reduced by the factor τ so that the τ -oracle can distinguish more coordinates from zero. In the case $\tau > 1$, the error variance in [\(1\)](#page-0-0) increases by the factor τ (as if the observations X_i 's get blurred), resulting in a smaller set of significant coordinates recovered by the τ -oracle. However, all τ -oracle rates $r^2(\theta, I_o^{\tau})$, $\tau > 0$, are related to the oracle rate $r^2(\theta) = r^2(\theta, I_o^1)$ by the trivial relations

$$
r^2(\theta) \le r^2(\theta, I_o^{\tau}) \le \tau r^2(\theta) \quad \text{for } \tau \ge 1, \quad r^2(\theta, I_o^{\tau}) \le r^2(\theta) \le \tau^{-1} r^2(\theta, I_o^{\tau}) \quad \text{for } 0 < \tau < 1.
$$

So, in principle we can obtain the result for any τ -oracle rate $r^2(\theta, I_o^{\tau})$ via the result for the oracle rate $r^2(\theta)$ and vice versa, but at the price of some multiplicative constant.

Actually, we can look at all τ -oracles I_o^{τ} , $\tau \geq 0$, from the following general perspective. Introduce a family of $n + 1$ sets

$$
\mathcal{I}_o = \{I_o(k), \, k \in [n]_0\}, \quad \text{where} \quad I_o(k) = \{i \in [n] : \theta_i^2 \ge \theta_{[k]}^2\}.
$$
 (20)

Clearly, these are embedded sets $\emptyset \triangleq I_o(0) \subseteq I_o(1) \subseteq I_o(2) \subseteq \ldots \subseteq I_o(n) = [n]$. Now notice that the oracle set $I_o(\theta)$ and actually all τ -oracles $I_o^{\tau}(\theta)$, $\tau \geq 0$, are all from this family, in fact, $I_o = I_o(i_o)$ and $I_o^{\tau} = I_o(i_{\tau}).$

3.2 Contraction results with oracle rate

The following theorem establishes that the empirical Bayes posterior $\hat{\pi}(\vartheta|X)$ (which is either $\tilde{\pi}(\vartheta|X)$ defined by [\(9\)](#page-6-3) or $\tilde{\pi}(\vartheta|X)$ defined by [\(14\)](#page-7-1)) contracts to θ with the oracle rate $r(\theta)$ from the frequentist P_θ-perspective, and the empirical Bayes posterior mean $\hat{\theta}$ which is either $\hat{\theta}$ defined by [\(12\)](#page-6-4) or $\check{\theta}$ defined by [\(14\)](#page-7-1)) converges to θ with the oracle rate $r(\theta)$, uniformly over the entire parameter space.

Theorem 1. Let Conditions $(A1)$ and $(A2)$ be fulfilled. Then there exist positive constants $M_0, M_1, H_0, H_1, m_0, m_1$ such that for any $\theta \in \mathbb{R}^n$ and any $M \geq 0$,

$$
\mathcal{E}_{\theta}\hat{\pi}\left(\|\vartheta - \theta\|^2 \ge M_0 r^2(\theta) + M\sigma^2|X\right) \le H_0 e^{-m_0 M},\tag{i}
$$

$$
\mathbf{P}_{\theta}(\|\hat{\theta} - \theta\|^2 \ge M_1 r^2(\theta) + M\sigma^2) \le H_1 e^{-m_1 M}.
$$
 (ii)

Remark 1. Notice that already claim [\(i\)](#page-10-1) of the theorem contains an oracle bound for the estimator $\hat{\theta}$. Indeed, by Jensen's inequality, we derive the oracle inequality

$$
\mathcal{E}_{\theta} \|\hat{\theta} - \theta\|^2 \le \mathcal{E}_{\theta} \hat{\mathcal{E}}(\|\theta - \theta\|^2 | X) \le M_0 r^2(\theta) + H_0 \int_0^{+\infty} e^{-m_0 u/\sigma^2} du = M_0 r^2(\theta) + \frac{H_0 \sigma^2}{m_0}.
$$
 (21)

However, comparing claim [\(ii\)](#page-10-2) with the oracle inequality [\(21\)](#page-10-3), we see that claim [\(ii\)](#page-10-2) is actually stronger (and more refined) than [\(21\)](#page-10-3) and therefore requires a separate proof.

Remark 2. A few more remarks on the theorem are in order.

- (i) The above local result implies the minimax optimality over various sparsity scales, see Section [3.5](#page-16-0) for more detail on this.
- (ii) The constants $M_0, M_1, H_0, H_1, m_0, m_1 > 0$ in the theorem depend only on β , B and some also on \varkappa , the exact expressions can be found in the proof.
- (iii) The non-asymptotic exponential bounds in terms of the constant M from the expression $M'r^2(\theta) + M\sigma^2$ (with some fixed M') in claims (i) and (ii) of the theorem provide a very refined characterization of the quality of the posterior $\hat{\pi}(\vartheta|X)$ and estimator $\hat{\theta}$, finer than, e.g., the traditional oracle inequalities like [\(21\)](#page-10-3). This refined formulation allows for subtle analysis in various asymptotic regimes $(n \to \infty, \sigma \to 0)$, or their combination) as we can let M depend in any way on n, σ , or both.

The next theorem describes the frequentist behavior of the selector \hat{I} and the empirical Bayes posterior for I, saying basically that \hat{I} and $\tilde{\pi}(I|X)$ "live" on a certain set that is, in a sense, almost as good as the oracle $I_o = I_o(\theta)$ defined by [\(18\)](#page-9-0). For any $\theta \in \mathbb{R}^n$, introduce

$$
I_* = I_*(\theta) \triangleq I_o^{\tau_0}(\theta) = I_o(\theta, \tau_0 \sigma^2), \quad i_* = |I_*|,
$$
\n(22)

where we fix some $\rho \in (0,1)$ and $\tau_0 > \frac{1+\rho}{1-\rho}$ $\frac{1+\rho}{1-\rho}\bar{\tau}$, $\bar{\tau}$ is defined by [\(16\)](#page-8-3). For example, we can take $\varrho = 0.1$ and $\tau_0 = \frac{11}{9}$ $rac{11}{9}\bar{\tau} + 0.1.$

Theorem 2. Let Condition [\(A1\)](#page-7-0) be fulfilled. The following relations hold for any $\theta \in \mathbb{R}^n$, $M \geq 0$.

(i) Let
$$
\kappa > \frac{4+\beta+2B}{2\beta}
$$
 (Condition (A2) implies this). There exist $M'_0, H'_0 > 0$ such that

$$
E_{\theta}\hat{\pi}(I \in \mathcal{I}: |I|\log(\frac{en}{|I|}) \geq M'_0|I \cap I_o|\log(\frac{en}{|I \cap I_o|}) + M|X) \leq H'_0 e^{-M}.
$$

(*ii*) Let $\varkappa > \beta^{-1} - \frac{1}{2}$ $\frac{1}{2}$ (Condition [\(A2\)](#page-8-2) implies this), $\overline{\tau}$ be defined by [\(16\)](#page-8-3). Fix any $I' \in \mathcal{I}$. Then there exist $H'_1, m'_0 > 0$ (independent of θ and I') such that

$$
\mathcal{E}_{\theta}\hat{\pi}\left(I\in\mathcal{I}:\sum_{i\in I'\setminus I}\frac{\theta_i^2}{\sigma^2}\geq \bar{\tau}|I\cup I'|\log(\frac{en}{|I\cup I'|})+M|X\right)\leq H_1'e^{-m_0'M}.
$$

In particular, let I_* be defined by [\(22\)](#page-10-4), then there exist $\alpha'_1, m'_1 > 0$ such that

$$
\mathcal{E}_{\theta}\hat{\pi}\big(I \in \mathcal{I}:|I|\log(\frac{en}{|I|}) \leq \varrho|I_{*}|\log(\frac{en}{|I_{*}|}) - M|X\big) \leq H'_{1}(\frac{en}{|I_{*}|})^{-\alpha'_{1}|I_{*}|}e^{-m'_{1}M}.\tag{23}
$$

(iii) Let Condition $(A2)$ be fulfilled, c_1, c_2, c_3 be the constants defined in Lemma [2.](#page-25-0) Then

$$
E_{\theta}\hat{\pi}(I \in \mathcal{I}: r^2(I,\theta) \ge c_3 r^2(\theta) + M\sigma^2|X) \le C_0 e^{-c_2 M}, \text{ where } C_0 = (1 - e^{1-c_1})^{-1}.
$$

Remark 3. The assertion [\(23\)](#page-11-0) holds also for I_* defined differently:

$$
I_* = I_*(\theta) = I_*(\theta, \tau) = I_*(\theta, \tau, \varrho) = \{ i \in [n] : \theta_i^2 \ge \theta_{[i_*]}^2 \}
$$
\n(24)

and $i_* = i_*(\tau, \varrho, \theta) = \max\{k \in [n]_0 : \sum_{\varrho k}^k \theta_{[i]}^2 \ge \tau(1-\varrho)\sigma^2 k \log(\frac{en}{k})\}\.$ Indeed, the only difference in the proof of [\(23\)](#page-11-0) for I_* defined by [\(24\)](#page-11-1) is that, instead of [\(43\)](#page-31-0), we have the bound

$$
\sum_{i \in I_* \setminus I} \frac{\theta_i^2}{\sigma^2} \ge \sum_{\varrho |I_*|}^{|I_*|} \frac{\theta_{[i]}^2}{\sigma^2} \ge \tau(1-\varrho)|I_*| \log(\frac{\epsilon n}{|I_*|}) \ge \tau' |I \cup I_*| \log(\frac{\epsilon n}{|I \cup I_*|}) + \tau' M,
$$

so that $m'_1 = \tau' m'_0$ in this case and the rest of the proof is the same.

For any $\theta \in \mathbb{R}^n$, the set $I_*(\theta)$ is the representative from the family \mathcal{I}_o (defined by [\(20\)](#page-9-1)) that consists of "distinctly significant" coordinates of θ such that $\hat{\pi}(I|X)$ makes (almost) no mistake for selecting a big proportion of this set. Recall that for $\tau_1 \ge \tau_2$, $I_o(i_{\tau_1}) = I_o^{\tau_1} \subseteq I_o^{\tau_2} = I_o(i_{\tau_2})$ so that $i_{\tau_1} \leq i_{\tau_2}$. Since $I_* = I_0^{\tau_0}$ for $\tau_0 > 1$, we have $I_* \subseteq I_0$. So, the claims of the above theorem roughly mean that $\hat{\pi}(I|X)$ (i.e., the selector \hat{I} and the posterior $\tilde{\pi}(I|X)$) lives in the "shell" $\{I: I_o(K^{-1}i_*)\subseteq I \subseteq I_o(Ki_o)\}\$ for some sufficiently large K, where the sets $I_o(k)$ are defined by [\(20\)](#page-9-1). So, if I_* and I_o are "close" to each other (i.e., $i_o \le Ci_*$ for some $C > 0$), then $\hat{\pi}(I|X)$ recovers well the oracle structure I_o . The case $i_* \ll i_o$ is problematic (the corresponding θ is "deceptive") as the living shell for $\hat{\pi}(I|X)$ is then too wide. Property (i) of Theorem [2](#page-11-2) claims good "over-dimensionality" control of $\hat{\pi}(I|X)$ in terms of the oracle I_o . In other words, the method does a good job in assigning insignificant coordinates to zeros, for any $\theta \in \mathbb{R}^n$. On the other hand, there is no full "under-dimensionality" control for $\hat{\pi}(I|X)$, as property (ii) is only in terms of the set I_* (which may be much "smaller" than I_o): basically for deceptive θ 's, $\hat{\pi}(I|X)$ can make relatively many errors by assigning many significantly non-zero coordinates to zeros.

This is reminiscent of the same asymmetric situation for adaption to smoothness where it is also possible to control under-smoothing (e.g., by penalization procedures or Lepski's method), but not over-smoothing. In view of the lower bound results mentioned in the introduction, this is not an artefact of the method, it is a fundamental, unavoidable problem. It occurs for the so called deceptive parameters θ that have many smallish coordinates, just slightly under the noise level. Interestingly, controlling over-dimensionality (or under-smoothing for smoothness structures) is enough for solving adaptive estimation problem, but not for uncertainty quantification where we need both over-dimensionality control (for the optimal size) and under-dimensionality control (for the coverage). This is possible for the non-deceptive parameters described by the so called EBR condition and introduced in the next section.

3.3 Confidence ball under excessive bias restriction

Theorem [1](#page-10-0) establishes the strong local optimal properties of the empirical Bayes posterior $\hat{\pi}(\vartheta|X)$ and the empirical Bayes posterior mean $\hat{\theta}$, but these do not solve the uncertainty quantification problem yet. Let us construct a confidence ball by using the empirical Bayes posterior $\check{\pi}(\vartheta|X)$ defined by [\(14\)](#page-7-1). Since $\check{\pi}(\vartheta|X) = \bigotimes_{i=1}^{n} N(\check{\theta}_i, \check{\sigma}_i^2)$ with $\check{\theta}_i = X_i \mathbb{1}\{i \in \hat{I}\}\$ and $\check{\sigma}_i^2 = (1 - |\hat{I}|/en)\sigma^2 \mathbb{1}\{i \in \hat{I}\},\$ denoting by $\chi^2_{k,\alpha}$ the $(1 - \alpha)$ -quantile of χ^2_k -distribution we have

$$
\check{\pi}\big(\|\vartheta-\check{\theta}\|^2\leq \sigma^2\chi^2_{|\hat{I}|,\alpha}|X\big)\geq \check{\pi}\big(\|\vartheta-\check{\theta}\|^2\leq (1-|\hat{I}|/en)\sigma^2\chi^2_{|\hat{I}|,\alpha}|X\big)=1-\alpha.
$$

But $\chi^2_{|\hat{I}|,\alpha}$ is bounded by a constant multiple of $|\hat{I}|$, and hence for simplicity the latter can replace the former to obtain a credible ball. This leads to $B(\check{\theta}, M\sigma|\hat{I}|^{1/2})$ as a credible ball for θ , which can be guaranteed to have at least a given level of credibility by choosing a sufficiently large constant M. From (i) of Theorem [2](#page-11-2) it follows that $|\hat{I}|$ is of the order $|I_o|$. However, it is clear that $B(\check{\theta}, M\sigma|\hat{I}|^{1/2})$ cannot have a guaranteed coverage, since otherwise the center $\check{\theta}$ would be an estimator that mimics the R-oracle uniformly in $\theta \in \mathbb{R}^n$, which is impossible as we discussed earlier. Hence to obtain coverage, the order of the radius of any confidence ball must contain a logarithmic factor. This leads us to the inflated credible ball $B(\check{\theta}, M\hat{r})$, where

$$
\hat{r}^2 = \hat{r}^2(X) = \sigma^2 + \sigma^2 |\hat{I}| \log(en/|\hat{I}|).
$$
\n(25)

The empirical Bayes posterior $\check{\pi}(\vartheta|X)$ is well concentrated (in fact, in a ball of the size $M\sigma^2|I_o|$, but not around the truth, rather around its mean $\check{\theta}$ which in general is away from the truth by the distance at most of the order of the oracle rate $r(\theta)$. We can also construct a confidence ball by using the posterior $\tilde{\pi}(\vartheta|X)$ defined by [\(9\)](#page-6-3) with the same resulting properties, but with more involved mathematical derivations. Property (i) of Theorem [2](#page-11-2) means that \hat{r}^2 is at most of the order of the variance part of the oracle rate $r^2(\theta)$, so the size property holds uniformly over $\theta \in \mathbb{R}^n$. But this goes at the expense of the coverage, namely, the coverage property does not hold uniformly.

Indeed, according to Theorem [2,](#page-11-2) $\varrho \sigma^2 |I_*| \log(en/|I_*|) - M \sigma^2 \leq \hat{r}^2 \leq M_0' \sigma^2 |I_o| \log(en/|I_o|) +$ $M\sigma^2$ with large probability. But this shell can be wide if $\sigma^2|I_*| \log(en/|I_*|) \ll \sigma^2|I_o| \log(en/|I_o|)$.

If this happens (for deceptive θ 's), then the coverage property of the ball $B(\check{\theta}, M\hat{r})$ cannot be guaranteed because its radius can be of a smaller order than the oracle rate $r^2(\theta)$. This problem will not occur for those (non-deceptive) θ 's for which the bias part of the rate $r^2(I_*,\theta)$ (see definition [\(17\)](#page-8-1)) is within a multiple of its variance part $\sigma^2|I_*| \log(en/|I_*|)$. Indeed, then $\sigma^2|I_*| \log(en/|I_*|)$ must be at least some multiple of $r^2(I_*,\theta)$ which is in turn bigger than the oracle rate $r^2(\theta)$ by the definition [\(18\)](#page-9-0) of the oracle. This means that $\sigma^2|I_*| \log(en/|I_*|)$ is at least of the oracle rate order, which, together with (ii) of Theorem [2,](#page-11-2) imply that \hat{r} is also at least of the oracle rate order, resulting in a good coverage of the confidence ball $B(\check{\theta}, M_2\hat{r} + M\sigma^2)$ for some M_2 and sufficiently large M . This discussion motivates introducing the following condition.

CONDITION EBR. We say that $\theta \in \mathbb{R}^n$ satisfies the *excessive bias restriction* (EBR) condition with structural parameter $t \geq 0$ if $\theta \in \Theta_{\text{eb}}(t)$, where the corresponding set (called the *EBR* class) is

$$
\Theta_{\text{eb}}(t) = \Theta_{\text{eb}}(t, \tau_0) = \left\{ \theta \in \mathbb{R}^n : \sum_{i \in I_*^c} \theta_i^2 \le t\sigma^2 \left(1 + |I_*| \log(\frac{en}{|I_*|}) \right) \right\},\tag{26}
$$

where the set $I_* = I_o^{\tau_0}$ is defined by [\(22\)](#page-10-4).

The condition EBR essentially requires that the bias part of the rate $r^2(I_*,\theta)$ is dominated by a multiple of its variance part (additional σ^2 is needed to handle the case $I_* = \emptyset$). This is obviously satisfied also for the rate $r^2(I', \theta)$ for all $I' \in \mathcal{I}_o$ such that $I_* \subseteq I'$ (hence also for the oracle I_o since $I_* \subseteq I_o$, where the family \mathcal{I}_o is defined by [\(20\)](#page-9-1). Besides, $\Theta_{eb}(t_1) \subseteq \Theta_{eb}(t_2)$ for $t_1 \leq t_2$, and, by the definition of $I_*, \mathbb{R}^n = \Theta_{\rm eb}(\tau_0 n)$.

Now we can use the center $\ddot{\theta}$ and the radius \hat{r} in constructing a confidence ball for θ . The following theorem, which is the main result in the paper, describes the coverage and size properties of the confidence ball based on $\hat{\theta}$ and \hat{r} .

Theorem 3. Let Conditions [\(A1\)](#page-7-0) and [\(A2\)](#page-8-2) be fulfilled. Then there exist constants $M_2, H_2, m_2 >$ 0 such that for any $t, M \ge 0$, and with $\hat{R}_{M}^{2} = \hat{R}_{M}^{2}(M_{2}) = (t + 1)M_{2}\hat{r}^{2} + (t + 2)M\sigma^{2}$,

$$
\sup_{\theta \in \Theta_{\text{eb}}(t)} \mathbf{P}_{\theta} \big(\theta \notin B(\hat{\theta}, \hat{R}_M) \big) \leq H_2 e^{-m_2 M}, \sup_{\theta \in \mathbb{R}^n} \mathbf{P}_{\theta} \big(\hat{r}^2 \geq M'_0 \sigma^2 |I_o| \log(\frac{en}{|I_o|}) + (M+1) \sigma^2 \big) \leq H'_0 e^{-M},
$$

where $\Theta_{eb}(t)$ is defined by [\(26\)](#page-13-0), the constants M'_0, H'_0 are defined in Theorem [2.](#page-11-2)

Remark 4. Let the quantity $b(\theta)$ (called *excessive bias ratio*) be defined by

$$
b(\theta) = b(\theta, \tau_0) = \frac{\sum_{i \in I_*^c} \theta_i^2}{\sigma^2 + \sigma^2 |I_*| \log(en/|I_*|)} = \frac{\sum_{i=i_*+1}^n \theta_{[i]}^2}{\sigma^2 + \sigma^2 i_* \log(en/i_*)} = \frac{B(I_*, \theta)}{\sigma^2 + V(I_*, \theta)}.
$$
(27)

Note that, when proving Theorem [3,](#page-13-1) we actually established the following local assertions: there exist constants $M_2, \alpha_1, m''_1, H_2, m_2 > 0$ such that for any $\theta \in \mathbb{R}^n$ and any $\alpha, M \ge 0$

$$
P_{\theta}(\theta \notin B(\hat{\theta}, [(b(\theta) + 1)M_2\hat{r}^2 + (b(\theta) + 2)M\sigma^2]^{1/2})
$$

\n
$$
\leq H_1(\frac{en}{|I_o|})^{-\alpha_1|I_o|}e^{-m_1M} + H'_1(\frac{en}{|I_*|})^{-\alpha'_1|I_*|}e^{-m''_1M} \leq H_2e^{-m_2M},
$$

\n
$$
P_{\theta}(\hat{r}^2 \geq \sigma^2(M'_0 + \alpha)|I_o|\log(\frac{en}{|I_o|}) + (M + 1)\sigma^2) \leq H'_0(\frac{ne}{|I_o|})^{-\alpha|I_o|}e^{-M},
$$

where all the other constants $(H_1, m_1, H'_1, \alpha'_1, M'_0, H'_0)$ are defined in Theorems [1](#page-10-0) and [2.](#page-11-2) Notice that the above size relation holds uniformly in $\theta \in \mathbb{R}^n$. Although the coverage relation is also uniform in $\theta \in \mathbb{R}^n$, the main (unavoidable) problem is the dependence of the coverage relation on $b(\theta)$. That is why we introduced the EBR condition which essentially provides control over the quantity $b(\theta)$.

Remark 5. The smaller constant τ_0 (involved in the definition of the EBR condition) is, the less restrictive the EBR condition is, the limiting case $\tau_0 \downarrow 0$ corresponds basically to no condition. However, the main message here is that for any specific distribution of error vector ξ there is always some value of the constant τ_0 in the EBR condition (bounded away from zero, depending on how "bad" ξ is). We treat a general situation and are not concerned with the most exact (smallest) value for τ_0 , our bound for τ_0 is in terms of β , B and possibly too conservative for each specific distribution of ξ .

The idea of the set $I_* = I_0^{\tau_0}$ introduced by [\(22\)](#page-10-4) is that it contains $i_* = |I_*|$ most significant coordinates of vector θ that are (essentially) not missed by the procedure \hat{I} . The bias term of the rate $r^2(I_*,\theta)$ is the error that is made when setting significant coordinates to zero (whereas they may not be zero). Large ratio $b(\theta)$ defined by [\(27\)](#page-13-2) means that this error is relatively large as compared to the variance part of the rate $r^2(I_*,\theta)$. In a way, such θ 's "trick" the procedure $\hat{\theta}$ and can therefore be regarded as deceptive. For each $\theta \in \mathbb{R}^n$, $b(\theta)$ measures the amount of deceptiveness of θ : the bigger $b(\theta)$, the more deceptive θ . The EBR condition says that the deceptiveness has to be restricted: $\Theta_{eb}(t) = \{ \theta \in \mathbb{R}^n : b(\theta) \leq t \}.$ An explicit example of EBR parameters is the set of self-similar parameters introduced in [\[28\]](#page-35-9) which is in our terms $\Theta_{ss}(p,c,\tau_0) = \{ \theta \in \ell_0[p] : |I_*(\theta)| \geq cp \}$ for $p \in [n], c \in (0,1]$. If $\theta \in \Theta_{ss}(p,c,\tau_0)$, then $p \leq c^{-1} |I_*|$ and $\sum_{i \in I_*^c} \theta_i^2 \leq \sum_{|I_*|}^{c^{-1} |I_*|}$ $\int_{|I_{*}|}^{c^{-1}|I_{*}|} \theta_{[i]}^{2} \leq (c^{-1} - 1)\tau_{0}\sigma^{2}|I_{*}| \log(\frac{en}{|I_{*}|}),$ where the second inequality follows by the oracle definition. Hence, $\Theta_{\rm ss}(p,c,\tau_0) \subseteq \Theta_{\rm eb}((c^{-1}-1)\tau_0)$. Notice that $\Theta_{\text{ss}}(p,c,\tau) \subseteq \Theta_{\text{eb}}((c^{-1}-1)\tau)$ for any $\tau > 0$.

In particular, for $\theta \in \Theta_{ss}(p,1,\tau_0) = {\theta \in \ell_0[p] : |I_*(\theta)| = p}$, the insignificant coordinates I^c_* of such θ 's are the true zeros and the significant coordinates I_* are sufficiently distinct from zero. Then the set $I_*(\theta)$ coincides with the support supp $(\theta) \triangleq \{i \in [n] : \theta_i \neq 0\}$, i.e., $I_*(\theta)$ supp (θ) , so that $B(I_*, \theta) = 0$, implying $\Theta_{ss}(p, 1, \tau_0) \subseteq \Theta_{eb}(0)$. This class consists of the "nicest" (least deceptive) parameters satisfying the EBR condition with zero deceptiveness $t = 0$. The uncertainty quantification result is the strongest for this class because the inflating factor is the smallest as $t = 0$. More about the EBR condition is in Section [4.2.](#page-22-0)

3.4 Confidence ball of $n^{1/4}$ -radius without EBR

By analyzing the previous results, we see that the resulting radius \hat{r} of our constructed confidence ball is of the oracle rate only under the EBR condition. In general, \hat{r}^2 underestimates the oracle rate $r^2(\theta)$. The difference is the bias term which may in general be of a bigger order than the variance part, leading to a bad coverage. Suppose we want to construct a confidence ball of a full coverage uniformly over the whole space \mathbb{R}^n . Recall however that, in view of the above mentioned negative results of [\[18\]](#page-34-7), [\[10\]](#page-34-8), [\[3\]](#page-34-9) and [\[20\]](#page-35-4), no data dependent ball can provide full coverage and adaptive size simultaneously. Insisting on the full coverage, one can at best adapt to sparsity levels only in the range $s \geq C \sqrt{n}$ (i.e., actually for non-sparse parameters) and the

term of order $\sigma^2 \sqrt{n}$ should be in the radius. Let us give a heuristics behind this. An idea is to mimic the quantity $\|\theta - \hat{\theta}\|^2$ by $\hat{R}^2 = \|X - \hat{\theta}\|^2$. Clearly, there is a lot of bias in \hat{R}^2 , the biggest part of which is due to the term $\sigma^2 ||\xi||^2$ contained in \hat{R} . To de-bias for that part, we need to subtract its expectation $\sigma^2 E ||\xi||^2 = n\sigma^2$, where we assumed $Var(\xi_i) = 1$ in the model [\(1\)](#page-0-0) for simplicity. However, even de-biased quantity \hat{R}^2 can only be controlled up to a margin of the order $\sigma^2 \sqrt{n}$. That is why a term of the order $\sigma n^{1/4}$ is necessary in the radius of the confidence ball to provide coverage uniformly over the whole space \mathbb{R}^n .

To handle some technical issues in this case, we impose the following additional condition.

CONDITION (A3). Besides X given by [\(1\)](#page-0-0), we also observe $X' \in \mathbb{R}^n$ independent of X, where $X' = \theta + \sigma \xi'$, the random vector ξ' satisfies the following relations: $E\xi_i' = 0$, $Var(\xi_i') \leq C_{\xi}$, $i \in [n];$

$$
P(|\langle v, \xi' \rangle| \ge \sqrt{M}) \le \psi_1(M) \quad \forall v \in \mathbb{R}^n : ||v|| = 1; \ P(||\xi'||^2 - E||\xi'||^2| \ge M\sqrt{n}) \le \psi_2(M). \tag{A3}
$$

Here $\psi_1(M), \psi_2(M)$ are some positive monotonically decreasing functions such that $\psi_1(M) \downarrow 0$ and $\psi_2(M) \downarrow 0$ as $M \uparrow \infty$.

Condition (A4) is satisfied for independent normals $\xi_i \stackrel{\text{ind}}{\sim} N(0, 1)$ even if we do not have the sample X' at our disposal. Indeed, in this case we can "duplicate" the observations by randomization at the cost of doubling the variance in the following manner: create samples $X'=X+\sigma Z$ and $X'' = X - \sigma Z$, for a $Z = (Z_1, \ldots, Z_n)$ (independent of X) such that $Z_i \stackrel{\text{ind}}{\sim} N(0, 1)$. Relations [\(A3\)](#page-15-0) are then fulfilled with exponential functions $\psi_l(M) = Ce^{-cM}$, $l = 1, 2$, for some $C, c > 0$. If the sub-gaussianity condition [\(32\)](#page-20-0) is fulfilled for ξ' (which is the same as Condition (A1) in case of independent ξ_i 's), then $\psi_1(M) = e^{-\rho M}$. By Chebyshev's inequality, we see that the second relation in [\(A3\)](#page-15-0) is fulfilled with function $\psi_2(M) = cM^{-2}$ for any zero mean independent ξ_i 's with $E\xi_i^{\prime 4} \leq C$.

Coming back to the problem of constructing a confidence ball of full coverage uniformly over \mathbb{R}^n , let $\hat{\theta}$ and \hat{I} be defined as before and based on the sample X. We propose to mimic $\|\theta - \hat{\theta}\|^2$ by the de-biased quantity $\|X' - \hat{\theta}\|^2 - n\sigma^2$ plus additional $\sigma^2 \sqrt{n}$ -order term to control its oscillations, leading us to the following data dependent radius

$$
\tilde{R}_M^2 = (||X' - \hat{\theta}||^2 - n\sigma^2 + 2\sigma^2 G_M \sqrt{n})_+, \text{ where } G_M = \sqrt{M(M + M_1)}, \quad (28)
$$

 $x_+ = x \vee 0$ and the constant M_1 is from Theorem [1.](#page-10-0) The next theorem establishes the coverage and size properties of the confidence ball $B(\hat{\theta}, \tilde{R}_M)$.

Theorem 4. Let Conditions [\(A1\)](#page-7-0), [\(A2\)](#page-8-2) and [\(A3\)](#page-15-0) be fulfilled and \tilde{R}_{M}^{2} be defined by [\(28\)](#page-15-1). Then for any $M \geq 0$

$$
\sup_{\theta \in \mathbb{R}^n} \mathcal{P}_{\theta}(\theta \notin B(\hat{\theta}, \tilde{R}_M)) \leq \psi_1(M/4) + \psi_2(M) + H_1 e^{-m_1 M},
$$

\n
$$
\sup_{\theta \in \mathbb{R}^n} \mathcal{P}_{\theta}(\tilde{R}_M^2 \geq g_M(\theta, n)) \leq \psi_1(M/4) + \psi_2(M) + 2H_1 e^{-m_1 M},
$$

 $g_M(\theta,n) = M_1 r^2(\theta) + M\sigma^2 + 4\sigma^2 G_M \sqrt{n}$ and the constants H_1, m_1, M_1 are defined in Theorem [1.](#page-10-0)

By taking large enough M we can ensure the coverage and size relations uniformly over the entire space \mathbb{R}^n . However, notice the price for this overall uniformity: the radius of the constructed confidence ball is essentially of the order $\sigma n^{1/4} + r(\theta)$. So, it is always of the order at least $\sigma n^{1/4}$ even for very sparse parameters θ , and it is of the oracle rate order only for non-sparse parameters, when $r(\theta) \geq C \sigma n^{1/4}$. This is a fundamental problem for uncertainty quantification, which typically occurs when the parameter θ has many smallish coordinates θ_i , say, with θ_i^2 just under the noise level σ^2 . Clearly, in this case no method can reliably assign those coordinates to the significant set. As demonstrated in [\[18\]](#page-34-7), [\[10\]](#page-34-8), [\[3\]](#page-34-9) and [\[20\]](#page-35-4), the above mentioned price in the form of a big radius is absolutely unavoidable (even in the case of just two sparsity classes as is shown in [\[20\]](#page-35-4)), as soon as we require uniform coverage.

3.5 Implications: the minimax results over sparsity classes

In this subsection we elucidate the potential strength of the local approach. In particular, we demonstrate how the global adaptive minimax results over certain scales can be derived from the local results. Note that the oracle rate $r(\theta)$ is a local quantity in that it quantifies the level of accuracy of inference about specific θ and originally it is not linked to any particular scale of classes. However, it is always possible to relate the oracle rate to various scales. Precisely, if we want to establish global adaptive minimax results over certain scale, say, $\{\Theta_\beta, \beta \in \mathcal{B}\},\$ with corresponding minimax rates $\{r(\Theta_\beta), \beta \in \mathcal{B}\}\$ (the minimax rate over Θ_β is $r^2(\Theta_\beta) \triangleq$ $\inf_{\hat{\theta}} \sup_{\theta \in \Theta_{\beta}} \mathbb{E}_{\theta} \|\hat{\theta} - \theta\|^2$, where the infimum is taken over all estimators), the only thing we need to show is

$$
\sup_{\theta \in \Theta_{\beta}} r^2(\theta) \leq cr^2(\Theta_{\beta}), \quad \text{for all } \beta \in \mathcal{B}.
$$

If the above property holds, we say the oracle rate $r(\theta)$ covers the scale $\{\Theta_\beta, \beta \in \mathcal{B}\}\$. In this case, the local results on the estimation, the posterior contraction and the size relation of the confidence ball will immediately imply the corresponding global adaptive minimax results over the covered scale, (actually, simultaneously for all scales that are covered by the oracle rate $r(\theta)$). As to the coverage property, according to Theorem [3,](#page-13-1) it holds uniformly only over the EBR class $\Theta_{eb}(t)$, whichever scale we consider. Thus, specializing the coverage property to a particular scale boils down to intersecting this scale with the EBR class $\Theta_{eb}(t)$ in the coverage property.

Next, we consider two sparsity scales $\{\Theta_{\beta}, \beta \in \mathcal{B}\}\$ for which the adaptive minimax results (on the estimation problem, the contraction rate of the empirical Bayes posterior, and the size property of the confidence ball $B(\hat{\theta}, (M_2\hat{r}^2 + M)^{1/2})$ follow from our local results Theorems [1](#page-10-0) and [3.](#page-13-1) The results for other (covered) scales can also be readily derived.

Nearly black vectors. For $p_n \in [n]$ such that $p_n = o(n)$ as $n \to \infty$ (we use the usual o, O notation to describe the asymptotic behavior of certain quantities as $n \to \infty$), introduce the sparsity class $\ell_0[p_n] = \{\theta \in \mathbb{R}^n : s(\theta) = |I^*(\theta)| \leq p_n\}$, where by $I^*(\theta)$ and $s(\theta)$ we denote the active index set and the sparsity of $\theta \in \mathbb{R}^n$:

$$
I^*(\theta) = \{ i \in [n] : \theta_i \neq 0 \}, \qquad s(\theta) = |I^*(\theta)|. \tag{29}
$$

The minimax estimation rate over the class of nearly black vectors $\ell_0[p_n]$ with the sparsity parameter p_n is known to be $r^2(\ell_0[p_n]) = O(\sigma^2 p_n \log(\frac{n}{p_n}))$ as $n \to \infty$; see [\[14\]](#page-34-15). By the definition [\(18\)](#page-9-0) of the oracle rate $r^2(\theta)$, we have that $r^2(\theta) \le r^2(I^*(\theta), \theta)$. Then we obtain trivially that

$$
\sup_{\theta \in \ell_0[p_n]} r^2(\theta) \le \sup_{\theta \in \ell_0[p_n]} r^2(I^*(\theta), \theta) \le \sigma^2 p_n \log\left(\frac{en}{p_n}\right) = O\big(r^2(\ell_0[p_n])\big).
$$

The last relation, Theorems [1](#page-10-0) and [3](#page-13-1) immediately imply the adaptive minimax results for the scale $\ell_0[p_n]$. We summarize these results in the following corollary.

Corollary 1. Under the conditions of Theorem [3,](#page-13-1) we have for any $M \geq 0$

$$
\sup_{\theta \in \ell_0[p_n]} \mathbf{E}_{\theta} \hat{\pi} \left(\|\vartheta - \theta\|^2 \ge M_0 \sigma^2 p_n \log(\frac{en}{p_n}) + M \sigma^2 | X \right) \le H_0 e^{-m_0 M},
$$

\n
$$
\sup_{\theta \in \ell_0[p_n]} \mathbf{P}_{\theta} \left(\|\hat{\theta} - \theta\|^2 \ge M_1 \sigma^2 p_n \log(\frac{en}{p_n}) + M \sigma^2 \right) \le H_1 e^{-m_1 M},
$$

\n
$$
\sup_{\theta \in \ell_0[p_n]} \mathbf{P}_{\theta} \left(\hat{r}^2 \ge M_0' \sigma^2 p_n \log(\frac{en}{p_n}) + (M+1) \sigma^2 \right) \le H_0' e^{-M}.
$$

The next assertion describes some "over-dimensionality" (or "undersmoothing") control of the empirical Bayes posterior $\hat{\pi}(I|X)$ from the P_θ-perspective.

Theorem 5. Let $s(\theta)$ be defined by [\(29\)](#page-16-1). Under the conditions of Theorem [2,](#page-11-2) there exist $M_4, m_4 > 0$ such that for any $M > M_4$ and $\theta \in \mathbb{R}^n$

$$
\mathbb{E}_{\theta}\hat{\pi}(I:|I|>Ms(\theta)|X)\leq C_0\exp\big\{-m_4s(\theta)\big[(M-M_4)\log(\frac{en}{s(\theta)})-M\log M\big]\big\}.
$$

In particular, there exist constants $M'_4, m'_4 > 0$ such that

$$
\mathbf{E}_{\theta}\hat{\pi}(I:|I|>M_4's(\theta)|X)\leq C_0\exp\big\{-m_4's(\theta)\log(\tfrac{en}{s(\theta)})\big\}.
$$

The above assertion is a local type result, but can readily be specialized to the sparsity class $\theta \in \ell_0[s_n]$ in the minimax sense. If $s(\theta) \geq 1$, the probability bound goes to 0 as $n \to \infty$.

Weak ℓ_s -balls. For $s \in (0, 2)$, the weak ℓ_s -ball with the sparsity parameter p_n is defined by

$$
m_s[p_n] = \left\{ \theta \in \mathbb{R}^n : \theta_{[i]}^2 \le (p_n/n)^2 (n/i)^{2/s}, \ i \in \mathbb{N}_n \right\}, \quad p_n = o(\sigma n) \text{ as } n \to \infty,
$$

where $\theta_{[1]}^2 \geq \ldots \geq \theta_{[n]}^2$ are the ordered $\theta_1^2, \ldots, \theta_n^2$. This scale can be thought of as Sobolev hyper-rectangle for ordered (with unknown locations) coordinates: $m_s[p_n] = \mathcal{H}(\beta, \delta_n) = \{ \theta \in$ $\mathbb{R}^n: |\theta_{[i]}| \leq \delta_n i^{-\beta},$ with $\delta_n = n^{1/s} \frac{p_n}{n}$ and $\beta = 1/s > 1/2$.

Denote $j = O_{\theta}(i)$ if $\theta_i^2 = \theta_{[j]}^2$, with the convention that in the case $\theta_{i_1}^2 = \ldots = \theta_{i_k}^2$ for $i_1 < \ldots < i_k$ we let $O_\theta(i_{l+1}) = O_\theta(i_l) + 1, l = 1, \ldots, k-1$. The minimax estimation rate over this class is $r^2(m_s[p_n]) = n(\frac{p_n}{n})$ $(\frac{2n}{n})^s [\sigma^2 \log(\frac{n\sigma}{p_n})]^{1-s/2}$ when $n^{2/s}(\frac{p_n}{n})$ $(\frac{p_n}{n})^2 \ge \sigma^2 \log n$, and $r^2(m_s[p_n]) =$ $n^{2/s}(\frac{p_n}{n}$ $(\frac{p_n}{n})^2 + \sigma^2$ when $n^{2/s}(\frac{p_n}{n})$ $(\frac{2n}{n})^2 < \sigma^2 \log n$, as $n \to \infty$; see [\[16\]](#page-34-0) and [\[6\]](#page-34-1). Then take $I^*(\theta) = \{i \in \mathbb{R}^n : |f_i| = 1\}$ \mathbb{N}_n : $\ddot{O}_{\theta}(i) \leq p_n^* \},$ with $p_n^* = en(\frac{p_n}{n\sigma})$ $\frac{p_n}{n\sigma}$)^s $\left[\log(\frac{n\sigma}{p_n})\right]^{-s/2}$ in the case $n^{2/s}(\frac{p_n}{n})$ $(\frac{\partial n}{n})^2 \ge \sigma^2 \log n$, to derive

$$
\sup_{\theta \in m_s[p_n]} r^2(\theta) \le \sup_{\theta \in m_s[p_n]} r^2(I^*(\theta), \theta) \le \sigma^2 p_n^* \log(\frac{en}{p_n^*}) + n^{2/s} (\frac{p_n}{n})^2 \sum_{i > p_n^*} i^{-2/s}
$$
\n
$$
\le K_1 \sigma^2 p_n^* \log(\frac{n\sigma}{p_n}) + K_2 n^{2/s} (\frac{p_n}{n})^2 (p_n^*)^{1-2/s} \le K n (\frac{p_n}{n})^s \left[\sigma^2 \log(\frac{n\sigma}{p_n})\right]^{1-s/2} = O\left(r^2(m_s[p_n])\right),\tag{30}
$$

for some $K = K(s)$. The case $n^{2/s}(\frac{p_n}{n})$ $(\frac{2n}{n})^2 < \sigma^2 \log n$ is treated similarly by taking $p_n^* = 0$. Theorems [1](#page-10-0) and [3](#page-13-1) imply the minimax adaptive results for the scale m_s [p_n].

Corollary 2. Under the conditions of Theorem [3,](#page-13-1) we have for any $M \geq 0$

$$
\sup_{\theta \in m_s[p_n]} E_{\theta} \hat{\pi} \left(\| \vartheta - \theta \|^2 \ge M_0 K r^2(m_s[p_n]) + M \sigma^2 | X \right) \le H_0 e^{-m_0 M},
$$

\n
$$
\sup_{\theta \in m_s[p_n]} P_{\theta} \| (\hat{\theta} - \theta \|^2 \ge M_1 K r^2(m_s[p_n]) + M \sigma^2) \le H_1 e^{-m_1 M},
$$

\n
$$
\sup_{\theta \in m_s[p_n]} P_{\theta} (\hat{r}^2 \ge M_0' K r^2(m_s[p_n]) + (M+1) \sigma^2) \le H_0' e^{-M}.
$$

The following claim concerns the "over-dimensionality" control for the class m_s [p_n].

Theorem 6. Let the conditions of Theorem [2](#page-11-2) be fulfilled, $p_n^* = en(\frac{p_n}{n\sigma})$ $(\frac{p_n}{n\sigma})^s \left[\log(\frac{n\sigma}{p_n})\right]^{-s/2}$ and $n^{2/s}(\frac{p_n}{n})$ $(n\geq n^2) \geq \sigma^2 \log n$. Then there exist constants $M_5, m_5 > 0$ such that for any $M > M_5$ there exists $n_0 = n_0(M, s)$ such that, for all $n \geq n_0$,

$$
\sup_{\theta \in m_s[p_n]} \mathbb{E}_{\theta} \hat{\pi}(I : |I| > Mp_n^*|X) \le C_0 \exp\big\{-m_5(M-M_5)p_n^* \log(\frac{n\sigma}{p_n})\big\}.
$$

Notice that the exponential upper bound from the last relation converges to zero as $n\to\infty$ because $p_n^* \log(\frac{n\sigma}{p_n}) \geq e(\sigma^2 \log n)^{s/2} (\log(\frac{n\sigma}{p_n})^{1-s/2})$.

Remark 6. The same minimax results hold over the so called *strong* ℓ_s -ball $\ell_s[p_n] = \{ \theta \in \mathbb{R}^n :$ 1 $\frac{1}{n}\sum_{i=1}^n |\theta_i|^s \leq (\frac{p_n}{n})$ $\{\{e_n\}_{n=0}^{\infty}$, $s \in (0, 2)$, since $\ell_s[p_n] \subseteq m_s[p_n] \subseteq \ell_{s'}[p_n]$ for any $s' > s$.

4 Concluding remarks and EBR

4.1 Concluding remarks

Improving constants. Since our approach applies to a very general situation, many constants involved in the conditions and proofs may be rather conservative. Indeed, we do not specify any distribution of ξ and even dot not assume independence of its coordinates. For the problem to be at all solvable, the vector ξ has to have some minimal structure which is in our case provided by Condition [\(A1\)](#page-7-0). The constants β , B reflect in a generic way how bad (or how good) the vector ξ is, implying that almost all the constants in the proofs and conditions depend on β , B. Clearly, if a distribution of ξ is specified, many bounds can made more precise and many constants can

be improved, including the constants $\bar{\varkappa}$ and τ_0 from Conditions [\(A2\)](#page-8-2) and (A3), see Remark [5](#page-14-0) for more on constant τ_0 . Besides, some constants can be improved by using more precise inequalities at some steps of the proof. But this would make the presentation significantly lengthier without adding anything new conceptually.

For example, in case $\xi_i \stackrel{\text{ind}}{\sim} N(0, 1)$, we can sharpen up many constants in the proofs and conditions. In the proof of Lemma [1,](#page-24-1) we can compute exactly the right hand side of [\(33\)](#page-25-1) by using the elementary identity: for $Y \sim N(\mu_y, \sigma_y^2)$,

$$
\text{E} \exp\left\{\frac{aY^2}{2}\right\} = \exp\left\{\frac{a\mu_y^2}{2(1 - a\sigma_y^2)} - \frac{1}{2}\log(1 - a\sigma_y^2)\right\}, \quad \text{for any } a < \sigma_y^{-2}.\tag{31}
$$

By some tedious but straightforward calculations, we obtain the claim of Lemma [1](#page-24-1) for any $h \in [0,1)$ with the constants $A_h = \frac{h}{2(1+h)}$ $\frac{h}{2(1+h)}, B_h = \frac{h}{2(1-h)}$ $\frac{h}{2(1-h)},\ C_h=\frac{h}{2}$ $\frac{h}{2}$ and $D_h = \frac{h}{2} + \frac{1}{2}$ $\frac{1}{2} \log(1-h)$. If $I \setminus I_0 = \emptyset$, the bound holds also for $h = 1$ with $A_1 = \frac{1}{4}$ $\frac{1}{4}$, $B_1 = 0$, $C_1 = D_1 = \frac{1}{2}$ $\frac{1}{2}$. Next, since Lemma [1](#page-24-1) now holds for any $h \in [0, 1)$, we can try to optimize the choice of h in Lemma [2.](#page-25-0) We can also relax the requirement $c_1 > 2$ to $c_1 > 1$ in Lemma [2,](#page-25-0) leading to the bound for $\varkappa > \bar{\varkappa} = 2.04.$

The constants in the proof of Theorem [2](#page-11-2) can also be improved in the normal case and we can use the bound $E\left(\sum_{i\in I} \xi_i^2\right)^2 = |I|^2 + 2|I| \leq 3|I|^2$ instead of [\(15\)](#page-7-2) in the proof of Theorem [1.](#page-10-0)

Product prior. If, instead of the prior π , we take a prior $\bar{\pi} = \bar{\pi}_{K,\varkappa} = \sum_{I \in \mathcal{I}} \lambda_I \pi_I$ with $\tau_i^2(I) = K\sigma^2 \mathbb{1}\{i \in I\}$ for any fixed $K > 0$ (we can even allow $K = K_n \to \infty$, but $K_n = O(n)$, as $n \to \infty$) in [\(3\)](#page-5-0) and $\lambda_I = c_{\varkappa,n} \exp\{-\varkappa|I|\log n\}$ (with $\varkappa > \varkappa_0$ for some $\varkappa_0 > 0$) in [\(5\)](#page-5-1), then all the results will hold with $\log n$ instead of $\log(\frac{en}{|I|})$ in the oracle rate [\(18\)](#page-9-0). This case was studied in the first version of the arXiv-preprint of this paper. Thus, the results for the prior $\bar{\pi}$ are weaker than the results obtained in this paper. For example, the minimax rates for the sparsity classes (Corollaries [1,](#page-17-0) [2\)](#page-18-2) follow from these weaker results only if the sparsity parameter $p_n = O(n^{\gamma})$ for $\gamma \in [0,1)$ as $n \to \infty$, otherwise we obtain only the *near-minimax* rates, with the factor log n instead of $\log(\frac{n}{p_n})$.

However, there is an advantageous feature of the prior $\bar{\pi}$ as compared with π . Namely, it is of the product structure: for $\lambda_I = c_{\lambda} \prod_{i \in I} \lambda_i$ with $c_{\lambda} = \prod_{i=1}^n (1 + \lambda_i)^{-1}$, we compute $\bar{\pi} = \sum_{I \in \mathcal{I}} \lambda_I \pi_I = \bigotimes_{i=1}^n \left[\omega_i \mathcal{N}(\mu_{1,i}, K\sigma^2) + (1 - \omega_i) \delta_0 \right], \ \omega_i = \frac{\lambda_i}{1 + \lambda_i}$ $\frac{\lambda_i}{1+\lambda_i}$ $(\omega_i = \lambda (i \in I)$ is the prior probability that the random set I contains i). This leads to the product structure of the empirical Bayes posterior, so that the computation of the corresponding empirical Bayes estimator can easily be done in the coordinatewise fashion. Indeed, in our case $\lambda_i = \lambda = n^{-\varkappa}$ and some computations give the following empirical Bayes posterior

$$
\bar{\pi}(\vartheta|X) = \bigotimes_{i=1}^{n} \left[p_i \mathcal{N}(X_i, \frac{K\sigma^2}{K+1}) + (1-p_i)\delta_0 \right], \qquad p_i = 1/\left[1 + h \exp\{-\frac{X_i^2}{2\sigma^2}\} \right],
$$

where $p_i = \bar{\pi}(\theta_i \neq 0|\mathbf{X})$ and $h = h_{\varkappa,K} = \frac{\sqrt{K+1}}{\lambda} = n^{\varkappa}(K+1)^{1/2}$. The mean with respect to $\bar{\pi}(\vartheta|X)$ is readily obtained: $\bar{\theta} = \mathbb{E}_{\bar{\pi}}(\vartheta|X) = \hat{p}_i X_i, i \in [n],$ a shrinkage estimator with easily computable shrinkage factors p_i . Coordinatewise empirical Bayes medians can also be easily computed.

Cardinality dependent prior λ . Notice that the prior $\lambda = (\lambda_I, I \in \mathcal{I})$ defined by [\(5\)](#page-5-1) depends on the set $I \in \mathcal{I}$ only via its cardinality |I|, i.e., $\lambda_I = g(|I|)$ for some nonnegative function $g(k)$, $k = 0, 1, ..., n$. It is easy to see that in this case $\pi_n(k) = g(k) \binom{n}{k}$ $\binom{n}{k}, k = 0, 1, \ldots, n,$ determines the prior on the cardinality of I. Hence, the prior λ_I can always be modeled in two steps: first draw the random cardinality K according to the prior $\pi_n(k)$, and then given $K = k$, draw a random set $\mathcal I$ uniformly from the family of all subsets of $\mathcal I$ of cardinality k. Such priors λ are used in [\[13\]](#page-34-3), where the cardinality prior $\pi_n(k)$ can be taken to be a so called "complexity prior" $\pi_n(k) = \exp\{-ak \log(bn/k)\}\$ for some $a, b > 0$. Since $e^{k \log(n/k)} \leq {n \choose k}$ $\binom{n}{k}$ \leq $e^{k \log(ne/k)}$, the resulting prior mass λ_I on I is bounded below and above by expressions of the type $\exp\{-a_1|I|\log(b_1n/|I|)\}$, resembling the prior [\(5\)](#page-5-1). The condition on the complexity prior from [\[13\]](#page-34-3) essentially corresponds to our condition $\varkappa > \bar{\varkappa}$ for some $\bar{\kappa} > 0$ (Condition [\(A2\)](#page-8-2)).

Computing the estimators. Note that the estimator [\(12\)](#page-6-4) is a shrinkage estimator, and the estimator [\(14\)](#page-7-1) is a hard thresholding procedure. Indeed, the estimator [\(12\)](#page-6-4) is $\theta_i = p_i X_i$ where $p_i = \sum_{I: i \in I} \tilde{\pi}(I|X)$, and the estimator [\(14\)](#page-7-1) is $\check{\theta}_i = X_i 1\{|X_i| \geq \check{t}\}$, where $\check{t} = |X_{[\check{k}]}|$, $|X_{[1]}| \geq \ldots \geq |X_{[n]}|$, and \check{k} is the minimizer of $\sum_{i=k+1}^{n} X_{[i]}^2 + (2\varkappa + 1)\sigma^2 k \log(en/k)$.

The thresholding procedure is easy to implement, whereas the values p_i in the shrinkage procedure are more difficult to compute. It is demonstrated in [\[13\]](#page-34-3) how one can use the partial product structure (in the model and in π_I , but not in λ_I) to facilitate the computation of p_i 's. Other estimators can be considered, for example, the coordinatewise median with respect to $\tilde{\pi}$, which is going to be something in between shrinkage and thresholding.

Condition [\(A1\)](#page-7-0). First we mention that all the results still hold, if, instead of Condition [\(A1\)](#page-7-0), we assume the weaker condition: $\mathbb{E} \exp \{ \beta \sum_{i \in I} \xi_i^2 \} \leq C_{\beta} e^{B|I| \log(en/|I|)}$ for all $I \in \mathcal{I}$ and some $\beta \in (0,1], B, C_{\beta} > 0.$ However, we leave Condition [\(A1\)](#page-7-0) in its present form to provide a cleaner mathematical exposition.

It is interesting to relate Condition [\(A1\)](#page-7-0) to the so called sub-gaussianity condition on the error vector $\xi = (\xi_i, i \in [n])$. The random vector ξ is called sub-gaussian with parameter $\rho > 0$ if

$$
P(|\langle v,\xi\rangle| > t) \le e^{-\rho t^2} \quad \text{for all } t \ge 0 \text{ and all } v \in \mathbb{R}^n \text{ such that } ||v|| = 1. \tag{32}
$$

The sub-gaussianity condition [\(32\)](#page-20-0) and Condition [\(A1\)](#page-7-0) are close, but in general incomparable. For example, let $\xi_i = \xi_0, i \in [n]$, for some bounded random variable ξ_0 (say, uniform on $[-1, 1]$), then Condition (A1) trivially holds whereas the sub-gaussianity condition is not fulfilled. If the ξ_i 's are independent, then the sub-gaussianity condition is equivalent to Condition [\(A1\)](#page-7-0).

In a way, Condition [\(A1\)](#page-7-0) prevents too much dependence, but it still allows some interesting cases of dependent ξ_i 's. Suppose that the ξ_i 's follow an autoregressive model AR(1) with normal white noise:

$$
\xi_k = \alpha \xi_{k-1} + \epsilon_k, \quad \epsilon_k \stackrel{\text{ind}}{\sim} \mathcal{N}(0, 1), \quad k \in [n]; \quad \xi_0 = 0, \quad |\alpha| < 1.
$$

Let us show that Condition [\(A1\)](#page-7-0) is fulfilled for the vector $\xi = (\xi_i, i \in [n])$. We have that for any $k > l$, $\xi_k = \alpha^{k-l} \xi_l + \alpha^{k-l-1} \epsilon_{l+1} + \ldots + \epsilon_k = \alpha^{k-l} \xi_l + Z_{k-l}$, where $Z_{k'} \sim N(0, \sigma_{k'}^2)$ with $\sigma_{k'}^2 = 1 + \alpha^2 + \ldots + \alpha^{2(k'-1)} \leq \frac{1}{1-\alpha^2} \triangleq \sigma_0^2$. Clearly, for any $I \in \mathcal{I}$, there are $1 \leq k_1 < k_2 < \ldots <$

 $k_{|I|} \leq n$ such that $\sum_{i \in I} \xi_i^2 = \sum_{i=1}^{|I|} \xi_{k_i}^2$. Denote $\mathcal{F}_m = \sigma(\xi_{k_i}, 1 \leq i \leq m)$, $m \in [|I|]$, the σ -algebra generated by $\{\xi_{k_i}, 1 \leq i \leq m\}$. Choose β and α in such a way that $0 < \frac{2\beta\alpha^2}{1-4\beta\sigma^2}$ $\frac{2\rho\alpha^2}{1-4\beta\sigma_0^2} \leq \beta$. By using the elementary identity [\(31\)](#page-19-0), we first evaluate the conditional expectation

$$
E\left(e^{\beta(\xi_{k_{m-1}}^2+\xi_{k_m}^2)}|\mathcal{F}_{m-1}\right) \leq e^{\beta\xi_{k_{m-1}}^2}E\left(e^{2\beta\xi_{k_m}^2}|\mathcal{F}_{m-1}\right)
$$

= $\exp\left\{\left(\beta + \frac{2\beta\alpha^{2(k_m-k_{m-1})}}{1-4\beta\sigma_{k_m,k_{m-1}}^2}\right)\xi_{k_{m-1}}^2 - \frac{1}{2}\log(1-4\beta\sigma_{k_m,k_{m-1}}^2)\right\} \leq (1-4\beta\sigma_0^2)^{-1/2}e^{2\beta\xi_{k_{m-1}}^2}.$

Iterating the above conditional expectation argument, we establish Condition [\(A1\)](#page-7-0):

$$
\operatorname{E} \exp \left\{ \beta \sum_{i \in I} \xi_i^2 \right\} = \operatorname{E} \mathbb{E} \left[\exp \left\{ \beta \sum_{i \in I} \xi_i^2 \right\} \middle| \mathcal{F}_{|I|-1} \right] = (1 - 4\beta \sigma_0^2)^{-1/2} \operatorname{E} \left[\exp \left\{ \beta \sum_{i=1}^{|I|-2} \xi_{k_i}^2 \right\} e^{2\beta \xi_{k|I|-1}^2} \right]
$$

$$
\leq \dots \leq (1 - 4\beta \sigma_0^2)^{-|I|/2} = e^{B|I|}, \qquad \text{with} \quad B = \log(1 - 4\beta \sigma_0^2)^{-1/2}.
$$

Relation to paper [\[28\]](#page-35-9). When the present paper was under review, paper [\[28\]](#page-35-9) on the same topic appeared (with discussion, see also our contribution [\[5\]](#page-34-16) to this discussion). The main result of [\[28\]](#page-35-9) is the adaptivity of the confidence set constructed by the Bayesian approach over the sparsity scale of nearly black vectors (introduced in Section [3.5\)](#page-16-0) within a grand space $\ell_0[p_n]$ for some $p_n \to \infty$, $p_n = o(n)$ as $n \to \infty$, under the EBR condition. The EBR condition introduced in [\[28\]](#page-35-9) is essentially a version of our EBR condition adopted to the sparsity scale within the grand space $\ell_0[p_n]$. It is not difficult to see that, within the asymptotic framework $n \to \infty$ and restricting the values of θ to some grand space $\ell_0[p_n]$ with $p_n = o(n)$, the EBR condition introduced in [\[28\]](#page-35-9) is actually equivalent to our EBR condition specified to that embedded sparsity scale with appropriate choices of the constants involved.

Restricting the values of θ to some grand space $\ell_0[p_n]$ excludes some "almost sparse" parameters that are formally non-sparse (with many very small, but nonzero, entries), but this is in fact necessary to ensure the asymptotic regime $n \to \infty$ considered in [\[28\]](#page-35-9). The main differences of our approach and that of [\[28\]](#page-35-9) are the following. We obtain local results without relating to any sparsity scale, e.g., the true parameter θ may be not $\ell_0[p_n]$ -sparse at all. For example, as a consequence we derive the results not only for $\ell_0[p_n]$, but also for other sparsity scales, such as weak ℓ_s -balls $m_s[p_n]$. Next, we allow the error vector ξ to be non-normal and even not necessarily independent (but just satisfying Condition (A1)). Some of our constants in the proofs and conditions may be more conservative, which is not surprising since we pursue a more general situation. Finally, we derive non-asymptotic exponential concentration bounds, which give a refined characterization of the quality of coverage and size relation results (finer, than, e.g., Theorem 5 from [\[28\]](#page-35-9), which is asymptotic in $n \to \infty$) and allow subtle analysis for various asymptotic regimes.

We should mention that the derivation of our somewhat stronger results relies on certain explicit posterior expressions resulting from our choice of prior (mixture of normals, although the model is not assumed to be normal), whereas the horseshoe prior studied in [\[28\]](#page-35-9) leads to only implicit posterior quantities so that the authors had to overcome difficult technical issues in the proofs.

4.2 The EBR condition

A new perspective on EBR – the EBR scale. As mentioned in the introduction, it is impossible to construct optimal (fully) adaptive confidence set in the minimax sense over traditional smoothness and sparsity scales with a prescribed high coverage probability. Namely, there exist "deceptive" parameters $\theta \in \Theta'_0 = \mathbb{R}^n \backslash \Theta_0$ for which the coverage property in [\(2\)](#page-2-0) may not hold for arbitrarily small α_1 . Removing deceptive parameters Θ'_0 and restricting to the remaining set Θ_0 of non-deceptive parameters resolves this issue. This was the original motivation of introducing the EBR condition.

An interesting additional feature of the EBR condition is that it leads to *slicing of the* entire parameter space \mathbb{R}^n . This opens up a new perspective on the EBR and its role in the deceptiveness issue, which we explain next.

Note that the EBR condition $\theta \in \Theta_{\rm eb}(t,\tau)$ (see [\(26\)](#page-13-0)) is actually a family of embedded conditions parametrized by $t \geq 0$: $\Theta_{eb}(t_1, \tau) \subseteq \Theta_{eb}(t_2, \tau)$ for $t_1 \leq t_2$ and any $\tau \geq 0$. Note that, by the oracle definition, $\Theta_{\rm eb}(\tau n, \tau) = \mathbb{R}^n$ for any $\tau > 0$. An important observation is that this family of conditions effectively introduces a new scale $\cup_{t>0}\Theta_{\rm eb}(t,\tau)=\cup_{0\leq t\leq\tau n}\Theta_{\rm eb}(t,\tau)$ (for any fixed $\tau > 0$, to be called the *EBR scale*, with the structural parameter $t > 0$ measuring the allowed amount of deceptiveness for parameters $\theta \in \Theta_{eb}(t, \tau)$. Indeed, this scale "slices" \mathbb{R}^n in the sense that $\mathbb{R}^n = \bigcup_{0 \leq t \leq \tau_n} \Theta_{\rm eb}(t, \tau)$. The main benefit of introducing the EBR scale is that it gives the slicing of the entire space that is very suitable for uncertainty quantification. Indeed, the dictum "removing deceptive parameters" becomes a very natural notion in terms of the scale $\bigcup_{0\leq t\leq\tau n}\Theta_{\rm eb}(t,\tau)$ as it is nothing else but restricting the amount of deceptiveness t. This provides a new perspective at the above mentioned "deceptiveness" issue: basically, each parameter $\theta \in \mathbb{R}^n$ has a certain amount of "deceptiveness" that is measured by the excessive bias ratio $b_\tau(\theta)$ defined by [\(27\)](#page-13-2), or the smallest t for which $\theta \in \Theta_{eb}(t, \tau)$. The larger t, the more deceptive parameters are allowed in $\Theta_{\rm eh}(t, \tau)$. A mild and controllable price for the uniformity over $\Theta_{\rm eb}(t,\tau)$ in the coverage relation is the amount of inflating of the confidence ball needed to provide a guaranteed high coverage for the parameters of deceptiveness at most t. We should mention that the EBR scale is intrinsically tied to our Bayesian procedure as it depends on the proposed family of priors $\{\pi_I, I \in \mathcal{I}\}\$. A different family may lead to a different EBR scale. Note however that a version of our EBR condition (adapted to the sparsity scale) is still used for a different (horseshoe) prior in paper [\[28\]](#page-35-9) .

Interestingly, slicing is also possible by the parameter $\tau > 0$: $\mathbb{R}^n = \cup_{\tau \geq 0} \Theta_{\rm eb}(t, \tau)$ (for any $t > 0$, the embedding goes in the opposite direction: the smaller the τ , the weaker the EBR. Namely, $\Theta_{eb}(t, \tau_2) \subseteq \Theta_{eb}(t, \tau_1)$ for any $0 \leq \tau_1 \leq \tau_2$, $t > 0$, and the "limiting" EBR set $\lim_{\tau \downarrow 0} \Theta_{\rm eb}(t,\tau)$ expands to the entire space: $\Theta_{\rm eb}(t,0) = \mathbb{R}^n$. Besides, notice that the inflating factor in the confidence ball from Theorem [3](#page-13-1) will not not increase as $\tau \downarrow 0$ (in fact, it will decrease). A paradox seems to have emerged: by considering very small τ 's, we can have less deceptiveness without any price in the coverage relation. However, this paradox is resolved by reminding that the coverage relation from Theorem [3](#page-13-1) does not hold for arbitrarily small τ_0 because in [\(22\)](#page-10-4) $\tau_0 > \frac{1+\varrho}{1-\varrho}$ $\frac{1+\varrho}{1-\varrho}\bar{\tau}$, showing that "there is no free lunch". The lower bound for τ_0 can be relaxed (made smaller) for specific distribution of ξ , but it will always be some positive threshold reflecting the complexity of ξ .

The EBR does not affect the minimaxity over the sparsity scale $\ell_0[p]$. The EBR condition is mild from the minimax point of view in the following sense: if we take the traditional sparsity class $\ell_0[p] = \{ \theta \in \mathbb{R}^n : |I^*(\theta)| = ||\theta||_0 \leq p \}$ for $p \in \mathbb{N}_n$ and remove non-EBR parameters, then the minimax rate over the remaining part will not change (up to a constant). We outline the argument below. The minimax estimation rate was established by Birgé and Massart $[6]$ (Theorem 4 from [\[6\]](#page-34-1), formulated in our notation): for some universal constant $c > 0$,

$$
r^{2}(\ell_0[p]) \triangleq \inf_{\hat{\theta}} \sup_{\theta \in \ell_0[p]} \mathbb{E}_{\theta} \|\hat{\theta} - \theta\|^{2} \ge c\sigma^{2} p \log(en/p).
$$

The proof is based on considering the subset $\mathcal{B}_1(p) = \{ \theta \in \mathbb{R}^n : |I^*(\theta)| \leq p, |\theta_i| \leq p \}$ $\sigma^2 \log(en/p)$ $\subset \ell_0[p]$ and establishing the required lower bound for the minimax risk $R(\mathcal{B}_1(p))$ over the set $\mathcal{B}_1(p)$, thus obtaining $r^2(\ell_0[p]) \ge r^2(\mathcal{B}_1(p)) \ge c\sigma^2 p \log(en/p)$. Inspecting all the steps in the proof, we see that essentially the same lower bound (with a different constant c) holds for another subset of $\ell_0[p]$: $\mathcal{B}_2(p,\tau_0) = \{ \theta \in \mathbb{R}^n : |I^*(\theta)| = p, 2\tau_0\sigma^2 \log(en/p) \leq |\theta_i| \leq$ $(2\tau_0+1)\sigma^2\log(en/p)$ for all $i \in I^*(\theta)$, for τ_0 defined in [\(22\)](#page-10-4). For each $\theta \in \mathcal{B}_2(p,\tau_0)$, we have $I_*(\theta) = I_0^{\tau_0}(\theta) = I^*(\theta)$ so that I_*^c $\mathcal{L}_{*}^{c}(\theta) = \emptyset$, $|I_{*}(\theta)| = p$, and the EBR condition is trivially satisfied for any $t \geq 0$:

$$
\frac{\sum_{i\in I_*^c} \theta_i^2}{\sigma^2 (1+|I_*|\log(en/|I_*|))} = 0 \le t.
$$

This means that $r^2(\ell_0[p] \cap \Theta_{\text{eb}}(t)) \ge r^2(\mathcal{B}_2(p, \tau_0)) \ge c\sigma^2 p \log(en/p).$

5 Simulations

Here we present a small simulation study according to the model [\(1\)](#page-0-0) with $\xi_i \stackrel{\text{ind}}{\sim} N(0, 1)$, $\sigma = 1$ and $n = 500$. We used signals $\theta = (\theta_1, \ldots, \theta_n)$ of the form $\theta = (0, \ldots, 0, A, \ldots, A)$, where $p = #(\theta_{0,i} \neq 0)$ last coordinates of θ are equal to a fixed number A. Different sparsity levels $p_n \in \{25, 50, 100\}$ and "signal strengths" $A \in \{3, 4, 5\}$ are considered. In case $\xi_i \stackrel{\text{ind}}{\sim} N(0, 1)$, the Conditions [\(A1\)](#page-7-0) and [\(A2\)](#page-8-2) are fulfilled with $\beta = 0.4$, $B = 1$ and $\varkappa > 3.24$. The idea is to construct an empirical counterpart of the ball $B(\hat{\theta}, [b(\theta) + 1)M_2\hat{r}^2]^{1/2})$ which appears in Remark [4](#page-13-3) (for $M = 0$). We consider \hat{I} and $\check{\theta}$ defined respectively by [\(13\)](#page-7-3) and [\(14\)](#page-7-1), and take $\hat{b} = \frac{\sum_{i \notin I} (X_i^2 - 1)}{|\hat{I}|\log(|x_i| + |\hat{I}|)}$ $\frac{\sum_{i \notin I} (1 - i)^{-1}}{|\hat{I}| \log(en/|\hat{I}|)}$ as the estimator of $b(\theta)$ defined by [\(27\)](#page-13-2).

When computing \hat{I} given by [\(13\)](#page-7-3), an important choice is that of parameter $\varkappa > 0$. In our simulation study, we choose \varkappa via a *cross-validation* procedure. For that, we create two independent normal samples $X'_i = X_i + \eta_i$ and $X''_i = X_i - \eta_i$, where simulated independent standard normal η_i 's are assumed to be independent of ξ_i , $i \in [n]$. Then X'_i and X''_i are independent random variables with means θ_i and variances 2. In words, the observation sample can be duplicated at the cost of multiplying the variance by 2. Using these samples, we estimate $\varkappa > 0$ as follows: let $\check{\theta}' = \check{\theta}'(\hat{I}') = (X_i'1\{i \in \hat{I}'\}, i \in [n]),$ where $\hat{I}'(\varkappa) = \operatorname{argmin}_{I \in \mathcal{I}} \left\{-\frac{1}{\varkappa}\right\}$ $\sum_{i \in I} (X'_i)^2 + 2(2\varkappa + 1)|I| \log \left(\frac{en}{|I|} \right)$ $\frac{en}{|I|}\Big\}$, then $\hat{\varkappa} = \operatorname{argmin}_{\varkappa \in (0, \log n]} \| \hat{I}'(\varkappa) - X'' \|^{2}$. Now, let $\hat{I} = \hat{I}(\hat{\varkappa})$ be defined by [\(13\)](#page-7-3) with $\varkappa = \hat{\varkappa}$. Finally, consider the confidence ball $B(\check{\theta}, [\check{M}(\hat{b}+1)\hat{r}^2]^{1/2})$ around $\check{\theta}$ defined by [\(14\)](#page-7-1) where $\hat{b} = \frac{\sum_{i \in I^c} (X_i^2 - 1)}{\|\hat{t}\|_{\text{Lip}} \|\hat{t}\| \|\hat{t}\|}$ $\frac{\sum_{i \in \hat{I}^c} (X_i^z - 1)}{|\hat{I}| \log(en/|\hat{I}|)}$ and $\hat{r}^2 = |\hat{I}| \log(en/|\hat{I}|)$ given by [\(25\)](#page-12-0). Recall

that this confidence ball construction is inspired by local result formulation from Remark [4.](#page-13-3) The multiplicative factor \tilde{M} is intended to trade-off the size of the ball against its coverage probability. We take $\tilde{M}=2$, an intuitive justification for this choice would be the fact that we doubled the variance in the randomization procedure when duplicating the sample. For each sparsity level $p \in \{25, 50, 100\}$ and signal strength $A \in \{3, 4, 5\}$, we simulated 100 data vectors X of dimension $n = 500$ from the model [\(1\)](#page-0-0) and computed the average squared radius by \hat{R}^2 . Table [1](#page-24-2) shows the ratio of \hat{R}^2 to the oracle radial rate $r^2(\theta) = |I_o| \log(en/|I_o|) = p \log(en/p)$, where the oracle I_0 is defined by [\(18\)](#page-9-0), and the frequency $\bar{\alpha}$ of the event that confidence ball $B(\check{\theta},[\check{M}(\hat{b}+1)\hat{r}^2]^{1/2})$ contains the signal θ , respectively. The former quantity estimates the average inflating factor with respect to the oracle rate, and the latter estimates the coverage of the constructed confidence ball. We see that the most difficult case is $p = 25$ and $A = 3$. An informal interpretation of the simulation results for this case: we need to blow up the confidence ball of the oracle rate radius approximately by the factor 3.24 in order to cover a few small needles in a haystack with coverage probability approximately 0.92.

Table 1: The ratio $\hat{R}^2/r^2(\theta)$ and the frequency $\bar{\alpha}$ of the event that the confidence ball $B(\check{\theta},[\check{M}(\hat{b}+\check{b}(\theta))])$ $(1)\hat{r}^{2}\hat{z}^{1/2}$ contains the signal $\theta = (0,\ldots,0,A,\ldots,A)$ (where p last coordinates are equal to A) computed for 100 vectors X simulated from [\(1\)](#page-0-0) with $n = 500$, $\sigma = 1$.

	25				50			100		
		$3\quad 4\quad 5$		3 4 5			3 4 5			
$\hat{R}^2/r^2(\theta)$ 3.24 2.73 2.21 3.24 2.14 2.07 2.26 1.83 1.85										
α		$0.92 \quad 0.97 \quad 0.98$			$0.99 \quad 0.99 \quad 1$			$0.96 \qquad 1$		

6 Technical lemmas

First we provide a couple of technical lemmas used in the proofs of the main results.

Remark 7. Notice that in the below lemma we established the same bound for the both quantities $E_{\theta}\hat{\pi}(I|X) = E_{\theta}\tilde{\pi}(I|X)$ and $E_{\theta}\mathbb{1}\{\hat{I} = I\} = P_{\theta}(\hat{I} = I)$. The proofs of the properties of $\check{\pi}(\vartheta|X)$ and $\check{\theta}$ are exactly in the same as for $\tilde{\pi}(\vartheta|X)$ and $\tilde{\theta}$, with the only difference that everywhere (in the claims and in the proofs) $\hat{\pi}(I \in \mathcal{G}|X)$ should be read as $\tilde{\pi}(I \in \mathcal{G}|X)$ in case $\hat{\pi} = \tilde{\pi}$; and as $\mathbb{1}\{\hat{I} \in \mathcal{G}\}\$ in case $\hat{\pi} = \tilde{\pi}$, for all $\mathcal{G} \subseteq \mathcal{I}$ that appear in the proof. Hence, $E_{\theta}\hat{\pi}(I \in \mathcal{G}|X) = E_{\theta}\tilde{\pi}(I \in \mathcal{G}|X)$ in the former case, and $E_{\theta}\hat{\pi}(I \in \mathcal{G}|X) = P_{\theta}(\hat{I} \in \mathcal{G})$ in the latter case.

Lemma 1. Let Condition [\(A1\)](#page-7-0) be fulfilled. Then for any $\theta \in \mathbb{R}^n$ and any $I, I_0 \in \mathcal{I}$,

$$
\mathcal{E}_{\theta}\hat{\pi}(I|X) \leq \left[\frac{\lambda_I}{\lambda_{I_0}}\right]^h \exp\bigg\{B_h \sum_{i \in I \setminus I_0} \frac{\theta_i^2}{\sigma^2} - A_h \sum_{i \in I_0 \setminus I} \frac{\theta_i^2}{\sigma^2} + C_h|I_0|\log(\frac{en}{|I_0|}) - D_h|I|\log(\frac{en}{|I|})\bigg\},\
$$

where $h=\frac{2\beta}{3}$ $\frac{2\beta}{3}, A_h = \frac{\beta}{6}$ $\frac{\beta}{6}$, $B_h = \frac{2\beta}{3}$ $\frac{2\beta}{3}$, $C_h = \frac{\beta+B}{3}$ $\frac{1}{3}$ and $D_h = \frac{\beta - 2B}{3}$. If $I \setminus I_0 = \varnothing$, the bound holds also for $h = \beta$ with $A_h = \frac{\beta}{3}$ $\frac{\beta}{3},\ B_h=0,\ C_h=\frac{\beta}{2}+B,\ D_h=\frac{\beta}{2}$ $\frac{\beta}{2}$. If $I_0 \backslash I = \varnothing$, the bound holds also for $h = \beta$ with $A_h = 0, B_h = \beta, C_h = \frac{\beta}{2}$ $\frac{\beta}{2}$, $D_h = \frac{\beta}{2} - B$.

Proof of Lemma [1.](#page-24-1) Recall that $P_{X,I} = \phi(X_i 1\{i \notin I\}, 0, \sigma^2 + K_n(I)\sigma^2 1\{i \in I\})$. In case $\hat{\pi}(I|X) = \tilde{\pi}(I|X)$, we get by [\(11\)](#page-6-5) that, for any $I, I_0 \in \mathcal{I}$ and any $h \in [0, 1]$,

$$
E_{\theta}\hat{\pi}(I|X) = E_{\theta}\tilde{\pi}(I|X) = E_{\theta}\frac{\lambda_I P_{X,I}}{\sum_{J\in\mathcal{I}}\lambda_J P_{X,J}} \leq E_{\theta}\Big(\frac{\lambda_I P_{X,I}}{\lambda_{I_0}P_{X,I_0}}\Big)^h
$$
(33)

$$
= E_{\theta}\Big[\frac{\lambda_I \prod_{i=1}^n \phi(X_i \mathbb{1}\{i \notin I\}, 0, \sigma^2 + K_n(I)\sigma^2 \mathbb{1}\{i \in I\})}{\lambda_{I_0} \prod_{i=1}^n \phi(X_i \mathbb{1}\{i \notin I_0\}, 0, \sigma^2 + K_n(I_0)\sigma^2 \mathbb{1}\{i \in I_0\})}\Big]^h
$$

$$
= \Big[\frac{\lambda_I}{\lambda_{I_0}}\Big]^h E_{\theta} \exp\Big\{\frac{h}{2}\Big[\sum_{i\in I\setminus I_0} \frac{X_i^2}{\sigma^2} - \sum_{i\in I_0\setminus I} \frac{X_i^2}{\sigma^2} + |I_0|\log(\frac{en}{|I_0|}) - |I|\log(\frac{en}{|I|})\Big]\Big\}. \tag{34}
$$

In case $\hat{\pi}(I|X) = \mathbb{1}\{\hat{I} = I\}$, by the definition [\(13\)](#page-7-3) of \hat{I} and the Markov inequality, we derive that, for any $I, I_0 \in \mathcal{I}$ and any $h \geq 0$

$$
\mathrm{E}_{\theta}\hat{\pi}(I|X) = \mathrm{P}_{\theta}(\hat{I}=I) \leq \mathrm{P}_{\theta}\Big(\frac{\tilde{\pi}(I|X)}{\tilde{\pi}(I_0|X)} \geq 1\Big) \leq \mathrm{E}_{\theta}\Big[\frac{\tilde{\pi}(I|X)}{\tilde{\pi}(I_0|X)}\Big]^{h} = \mathrm{E}_{\theta}\Big(\frac{\lambda_I \mathrm{P}_{X,I}}{\lambda_{I_0} \mathrm{P}_{X,I_0}}\Big)^{h},
$$

which yields exactly the bound [\(33\)](#page-25-1), and hence the bound [\(34\)](#page-25-2) again.

Using Hölder's inequality, Condition [\(A1\)](#page-7-0) and the two elementary facts $X_i^2 \leq 2\theta_i^2 + 2\sigma^2 \xi_i^2$ and $-X_i^2 \le -\frac{\theta_i^2}{2} + \sigma^2 \xi_i^2$, we obtain

$$
\begin{split} \mathcal{E}_{\theta} \exp\Big\{ \t\frac{\beta}{3} \Big[\sum_{i \in I \setminus I_0} \tfrac{X_i^2}{\sigma^2} - \sum_{i \in I_0 \setminus I} \tfrac{X_i^2}{\sigma^2} \Big] \Big\} & \leq \Big(\mathcal{E}_{\theta} e^{-\beta \sum_{i \in I \setminus I_0} \tfrac{X_i^2}{\sigma^2}} \Big)^{2/3} \Big(\mathcal{E}_{\theta} e^{-\beta \sum_{i \in I_0 \setminus I} \tfrac{X_i^2}{\sigma^2}} \Big)^{1/3} \\ & \leq \exp\Big\{ \tfrac{2\beta}{3} \sum_{i \in I \setminus I_0} \tfrac{\theta_i^2}{\sigma^2} + \tfrac{2B}{3} |I \setminus I_0| - \tfrac{\beta}{6} \sum_{i \in I_0 \setminus I} \tfrac{\theta_i^2}{\sigma^2} + \tfrac{B}{3} |I_0 \setminus I| \Big\}. \end{split}
$$

Since $|I\setminus I_0| \leq |I| \leq |I| \log(\frac{en}{|I|})$ and $|I_0\setminus I| \leq |I_0| \leq |I_0| \log(\frac{en}{|I_0|})$, the lemma follows for $h = \frac{2\beta}{3}$ 3 from the last display and [\(34\)](#page-25-2).

 $\frac{X_i^2}{\sigma^2}$ } \leq If $I \setminus I_0 = \emptyset$, we take $h = \beta$ in [\(34\)](#page-25-2) and combine this with E_{θ} exp $\{-\frac{\beta}{2}$ $\frac{\beta}{2} \sum_{i \in I_0 \setminus I}$ $\frac{\theta_i^2}{\sigma^2} + B|I_0\setminus I|\},\$ which holds in view of Condition [\(A1\)](#page-7-0) and $-\frac{X_i^2}{\sigma^2} \leq -\frac{2\theta_i^2}{3\sigma^2} +$ $\exp\big\{-\frac{\beta}{3}$ $\frac{\beta}{3}\sum_{i\in I_0\setminus I}$ $2\xi_i^2$, as $(a + b)^2 \geq 2a^2/3 - 2b^2$. If $I_0 \setminus I = \emptyset$, we take $h = \beta$ in [\(34\)](#page-25-2) and combine this with $\left\{\frac{X_i^2}{\sigma^2}\right\} \le \exp\left\{\beta \sum_{i \in I \setminus I_0}$ $\frac{\theta_i^2}{\sigma^2} + B|I\setminus I_0|\}$ which holds in view of Condition [\(A1\)](#page-7-0) $E_{\theta} \exp \left\{ \frac{\beta}{2} \sum_{i \in I \setminus I_0}$ and $\frac{X_i^2}{\sigma^2} \le \frac{2\theta_i^2}{\sigma^2} + 2\xi_i^2$. 2 \Box

Note that above lemma holds for any set $I_0 \in \mathcal{I}$. By taking $I_0 = I_0$ defined by [\(18\)](#page-9-0), we obtain the following lemma.

Lemma 2. Let Conditions [\(A1\)](#page-7-0) and [\(A2\)](#page-8-2) be fulfilled. Then there exist positive constants $c_1 = c_1(\varkappa) > 2$, c_2 and $c_3 = c_3(\varkappa)$ such that for any $\theta \in \mathbb{R}^n$

$$
\mathrm{E}_{\theta}\hat{\pi}(I|X) \leq \left(\frac{ne}{|I|}\right)^{-c_1|I|} \exp\big\{-c_2\sigma^{-2}\big[r^2(I,\theta)-c_3r^2(\theta)\big]\big\}.
$$

Proof of Lemma [2.](#page-25-0) With constants h , A_h , B_h , C_h , D_h given in Lemma [1,](#page-24-1) define the constant $c_1 = c_1(\varkappa) = \varkappa h + D_h - A_h = \frac{2\beta\varkappa}{3} + \frac{\beta-2B}{3} - \frac{\beta}{6} > 2$ as $\varkappa > \bar{\varkappa}$ by Condition (A2). Since $xh + D_h = c_1 + A_h$, the definition [\(5\)](#page-5-1) of λ_I entails that

$$
\begin{aligned} &\left(\frac{\lambda_I}{\lambda_{I_0}}\right)^h \exp\left\{C_h|I_0|\log(\frac{en}{|I_0|}) - D_h|I|\log(\frac{en}{|I|})\right\} \\ &= (\frac{ne}{|I|})^{-c_1|I|} \exp\left\{(\varkappa h + C_h)|I_0|\log(\frac{en}{|I_0|}) - A_h|I|\log(\frac{en}{|I|})\right\}. \end{aligned}
$$

Using the last relation and Lemma [1](#page-24-1) with $I_0 = I_o$, we bound

$$
\mathcal{E}_{\theta}\hat{\pi}(I|X) \leq \left[\frac{\lambda_I}{\lambda_{I_o}}\right]^h \exp\left\{B_h \sum_{i\in I\backslash I_o} \frac{\theta_i^2}{\sigma^2} - A_h \sum_{i\in I_o\backslash I} \frac{\theta_i^2}{\sigma^2} + C_h|I_o|\log(\frac{en}{|I_o|}) - D_h|I|\log(\frac{en}{|I|})\right\}
$$

$$
= \left(\frac{ne}{|I|}\right)^{-c_1|I|} \exp\left\{-A_h \sum_{i\in I_o\backslash I} \frac{\theta_i^2}{\sigma^2} - A_h|I|\log(\frac{en}{|I|}) + B_h \sum_{i\in I\backslash I_o} \frac{\theta_i^2}{\sigma^2} + (\varkappa h + C_h)|I_o|\log(\frac{en}{|I_o|})\right\}.
$$

The claim of the lemma follows with the constants $c_1 = (4\beta \varkappa + \beta - 4B)/6 > 2$, $c_2 = A_h = \beta/6$
and $c_3 = c_3(\varkappa) = \max\{B_h, \varkappa h + C_h\}/A_h = (\varkappa h + C_h)/A_h = 4\varkappa + 2(\beta + B)/\beta$. and $c_3 = c_3(\kappa) = \max\{B_h, \kappa h + C_h\}/A_h = (\kappa h + C_h)/A_h = 4\kappa + 2(\beta + B)/\beta.$

Lemma 3. Let Y_1, \ldots, Y_n be some random variables such that, for any $I \in \mathcal{I}$, $E e^{t \sum_{i \in I} Y_i} \le$ $A_{|I|}(t)$ for some $t > 0$ and $A_k(t)$. Let $Y_{[1]} \geq Y_{[2]} \geq \ldots \geq Y_{[n]}$. Then, for any $k \in \mathbb{N}_n$ and $C, c \geq 0,$

$$
P\Big(\sum_{i=1}^k Y_{[i]} \ge Ck \log(\frac{en}{k}) + c\Big) \le A_k(t) \exp\{-(Ct - 1)k \log(\frac{en}{k}) - ct\},
$$

$$
E\sum_{i=1}^k Y_{[i]} \le t^{-1} [k \log(\frac{en}{k}) + \log(A_k(t))].
$$

In particular, if $\xi_1, \ldots, \xi_n \stackrel{\text{ind}}{\sim} N(0, 1)$, then for any $k \in \mathbb{N}_n$, $C, c \ge 0$

$$
\mathbb{P}\Big(\sum_{i=1}^k \xi_{[i]}^2 \ge Ck \log\left(\frac{en}{k}\right) + c\Big) \le \left(\frac{en}{k}\right)^{-(0.4C-2)k} e^{-0.4c}, \quad \mathbb{E}\sum_{i=1}^k \xi_{[i]}^2 \le 6k \log\left(\frac{en}{k}\right).
$$

Proof. By Jensen's inequality, we derive

$$
\exp\left\{t\mathbf{E}\sum_{i=1}^k Y_{[i]}\right\} \le \mathbf{E}\exp\left\{t\sum_{i=1}^k Y_{[i]}\right\} \le \sum_{I:|I|=k} \mathbf{E}\exp\left\{t\sum_{i\in I} Y_i\right\} \le {n \choose k} A_k(t).
$$

Then E exp $\{t\sum_{i=1}^k Y_{[i]}\}\leq {n \choose k}$ $\binom{n}{k} A_k(t) \leq e^{k \log(\frac{en}{k}) + \log(A_k(t))}$, where we used $\binom{n}{k}$ $\binom{n}{k} \leq \left(\frac{en}{k}\right)$ $(\frac{en}{k})^k$. This and the (exponential) Markov inequality yield the first relation:

$$
P\left(\sum_{i=1}^k Y_{[i]} \ge Ck \log\left(\frac{en}{k}\right) + c\right) \le A_k(t) \exp\left\{-(Ct-1)k \log\left(\frac{en}{k}\right) - ct\right\}.
$$

The first display implies also the second relation: $E\sum_{i=1}^{k} Y_{[i]} \leq t^{-1} [\log {n \choose k}$ ${k \choose k} + \log(A_k(t))$.

As to the normal case, for any $I \in \mathcal{I}$ and any $t < \frac{1}{2}$ we have that $E \exp\left\{t \sum_{i \in I} \xi_i^2\right\}$ $(1-2t)^{-|I|/2} = A_{|I|}(t)$. Since $A_k(t) \le e^k \le e^{k \log(\frac{en}{k})}$ for any $t \le (1-e^{-2})/2 < 0.43$, the first assertion for the normal case follows by taking $t = 0.4$. By taking $t = \frac{1}{4}$ $\frac{1}{4}$, the second assertion follows since $E \sum_{i=1}^{k} \xi_{[i]}^2 \leq 4k \log(\frac{en}{k}) + 2k \log 2 \leq 6k \log(\frac{en}{k})$.

This lemma is useful if $A_k(t) \leq C_1(\frac{en}{k})$ $(\frac{kn}{k})^{C_2 k}$ for some $t, C_1, C_2 > 0$; in particular, for $Y_i = \xi_i^2$, where the ξ_i 's satisfy Condition [\(A1\)](#page-7-0). Then Lemma [3](#page-26-0) applies with $t = \beta$ and $A_k(\beta) = e^{Bk}$:

$$
\mathbb{P}\Big(\sum_{i=1}^k \xi_{[i]}^2 \ge \frac{(1+B)}{\beta} k \log(\frac{en}{k}) + M\Big) \le \exp\{-\beta M\}, \quad k \in \mathbb{N}_n, M \ge 0. \tag{35}
$$

7 Proofs of the theorems

Here we gather the proofs of the theorems. By C_0, C_1, C_2 etc., denote constants which are different in different proofs. Recall that $Y_{[1]} \geq Y_{[2]} \geq \ldots \geq Y_{[n]}$ denote the ordered Y_1, \ldots, Y_n .

Proof of Theorem [1.](#page-10-0) Recall the constants c_1, c_2, c_3 defined in the proof of Lemma [2.](#page-25-0) Let $M_0 =$ $2c_3(6+\frac{1+B}{\beta})$. Introduce the subfamily of index sets $\mathcal{S}_M = \mathcal{S}_M(\theta) = \{I \in \mathcal{I} : r^2(I,\theta) \leq$ $c_3r^2(\theta) + \frac{\beta}{40(1+B)}M\sigma^2$, $m = m_M(\theta) = \max\{|I| : I \in \mathcal{S}_M\}$, and the event $A_M = A(\theta) =$ $\left\{ \sum_{i=1}^{m} \xi_{[i]}^2 \leq \frac{(1+B)}{\beta} m \log(\frac{en}{m}) + \frac{M}{8} \right\}$. We have

$$
\hat{\pi}(\|\vartheta - \theta\|^2 \ge M_0 r^2(\theta) + M\sigma^2|X) \le \mathbb{1}_{A_M^c} + \hat{\pi}(I \in \mathcal{S}_M^c|X) \n+ \sum_{I \in \mathcal{S}_M} \mathbb{1}_{A_M} \hat{\pi}_I(\|\vartheta - \theta\|^2 \ge M_0 r^2(\theta) + M\sigma^2|X) \hat{\pi}(I|X) = T_1 + T_2 + T_3.
$$

Now we bound the quantities $E_{\theta}T_1$, $E_{\theta}T_2$ and $E_{\theta}T_3$.

First, we bound $E_{\theta}T_1$ by using Lemma [3](#page-26-0) (see also [\(35\)](#page-27-1)):

$$
E_{\theta}T_1 = P_{\theta}(A_M^c) = P\left(\sum_{i=1}^m \xi_{[i]}^2 > \frac{(1+B)}{\beta} m \log(\frac{en}{m}) + \frac{M}{8}\right) \le \exp\left\{-\beta M/8\right\}.
$$
 (36)

Let us bound $E_{\theta}T_2$. Since $\binom{n}{k}$ $\binom{n}{k} \leq \left(\frac{en}{k}\right)$ $(\frac{k}{k})^k$ and $c_1 > 2$, the following relation holds:

$$
\sum_{I \in \mathcal{I}} \left(\frac{ne}{|I|} \right)^{-c_1 |I|} = \sum_{k=0}^{n} {n \choose k} \left(\frac{en}{k} \right)^{-c_1 k} \le \sum_{k=0}^{n} \left(\frac{en}{k} \right)^{-k(c_1 - 1)} \le (1 - e^{1 - c_1})^{-1} \triangleq C_0.
$$
 (37)

If $I \in \mathcal{S}_M^c$, then $r^2(I,\theta) > c_3 r^2(\theta) + \frac{\beta}{40(1+B)} M \sigma^2$. Using this, Lemma [2](#page-25-0) and [\(37\)](#page-27-2), we bound $E_{\theta}T_2$:

$$
E_{\theta}T_2 = \sum_{I \in S_M^c} E_{\theta}\hat{\pi}(I|X) \le \sum_{I \in S_M^c} \left(\frac{ne}{|I|}\right)^{-c_1|I|} \exp\left\{-c_2\sigma^{-2}\left[r^2(I,\theta) - c_3r^2(\theta)\right]\right\}
$$

$$
\le \sum_{I \in \mathcal{I}} \left(\frac{ne}{|I|}\right)^{-c_1|I|} \exp\{-c_2\beta M/(40(1+B))\} \le C_0 \exp\{-c_2\beta M/(40(1+B))\}. \tag{38}
$$

It remains to bound $E_{\theta}T_3$. For each $I \in S_M$, $\sigma^2|I|\log(en/|I|) \le r^2(I,\theta) \le c_3r^2(\theta) + \frac{\beta}{40(I+B)}M\sigma^2$. Since $m = \max\{|I| : I \in S_M\}$, then $\sigma^2m \log(\frac{en}{m}) \le c_3r^2(\theta) + \frac{\beta}{40(I+B)}M\sigma^2$. Thus, for any $I \in \mathcal{S}_M$, the event A_M implies that $\sum_{i \in I} \xi_i^2 \leq \sum_{i=1}^m \xi_{[i]}^2 \leq \frac{(1+B)}{\beta} m \log(\frac{en}{m}) + \frac{M}{8} \leq$ $(1 + B)$ $\frac{1}{\beta}B_{c3}\sigma^{-2}r^2(\theta)+\frac{3M}{20}$. Denote for brevity $\Delta_M(\theta)=M_0r^2(\theta)+M\sigma^2$ and recall that $\sum_{i\in I^c}\theta_i^2\leq$ $r^2(I,\theta) \leq c_3 r^2(\theta) + \frac{\beta}{40(I+B)} M\sigma^2 \leq c_3 r^2(\theta) + \frac{M}{40}\sigma^2$ for any $I \in \mathcal{S}_M$. Then for any $I \in \mathcal{S}_M$

$$
A_M \subseteq \left\{ \frac{\Delta_M(\theta)}{2} - \sigma^2 \sum_{i \in I} \xi_i^2 - \sum_{i \in I^c} \theta_i^2 \ge \left[\frac{M_0}{2} - \frac{1 + B + \beta}{\beta} c_3 \right] r^2(\theta) + \frac{13M\sigma^2}{40} \right\}.
$$
 (39)

According to [\(10\)](#page-6-2), $\hat{\pi}_I(\vartheta|X) = \bigotimes_{i=1}^n N(X_i(I), \sigma_i^2(I)),$ with $X_i(I) = X_i \mathbb{1}\{i \in I\}$ and $\sigma_i^2(I) =$ $i(I)$), with $\Lambda_i(I) - \Lambda_i \mathbb{1} \setminus \mathbb{1}$ and $\mathbb{0} \setminus \mathbb{1}$ $\frac{K_n(I)\sigma^2 \mathbb{1}\{i\in I\}}{K_n(I)+1}$. Let P_Z be the measure of $Z=(Z_1,\ldots,Z_n)$, with $Z_i \stackrel{\text{ind}}{\sim} N(0,1)$. By using [\(39\)](#page-28-0), the fact that $\frac{r^2(\theta)}{\sigma^2} \ge c_3^{-1}(m \log(\frac{en}{m}) - \frac{\beta}{40(1+B)}M)$ and Lemma [3](#page-26-0) (now applied to the Gaussian case), we obtain that, for any $I \in \mathcal{S}_M$,

$$
\hat{\pi}_{I}(\|\vartheta - \theta\|^{2} \ge M_{0}r^{2}(\theta) + M\sigma^{2}|X) \mathbb{1}_{A_{M}} = \mathbb{P}_{Z}\Big(\sum_{i=1}^{n}(\sigma_{i}(I)Z_{i} + X_{i}(I) - \theta_{i})^{2} \ge \Delta_{M}(\theta)\Big) \mathbb{1}_{A_{M}}
$$
\n
$$
\le \mathbb{P}_{Z}\Big(\sum_{i=1}^{n}\sigma_{i}^{2}(I)Z_{i}^{2} \ge \frac{\Delta_{M}(\theta)}{2} - \sum_{i=1}^{n}(X_{i}(I) - \theta_{i})^{2}\Big) \mathbb{1}_{A_{M}}
$$
\n
$$
\le \mathbb{P}_{Z}\Big(\sum_{i\in I}\sigma^{2}Z_{i}^{2} \ge \frac{\Delta_{M}(\theta)}{2} - \sum_{i\in I}\sigma^{2}\xi_{i}^{2} - \sum_{i\in I}\theta_{i}^{2}\Big) \mathbb{1}_{A_{M}}
$$
\n
$$
\le \mathbb{P}_{Z}\Big(\sum_{i\in I}Z_{i}^{2} \ge \left[\frac{M_{0}}{2} - \left(\frac{1+B}{\beta} + 1\right)c_{3}\right] \frac{r^{2}(\theta)}{\sigma^{2}} + \frac{13M}{40}\Big)
$$
\n
$$
\le \mathbb{P}_{Z}\Big(\sum_{i=1}^{m}Z_{[i]}^{2} \ge \left(\frac{M_{0}}{2c_{3}} - \frac{1+B}{\beta} - 1\right)\left[m\log(\frac{en}{m}) - \frac{\beta}{40(1+B)}M\right] + \frac{13M}{40}\Big)
$$
\n
$$
\le \mathbb{P}_{Z}\Big(\sum_{i=1}^{m}Z_{[i]}^{2} \ge 5m\log(\frac{en}{m}) + \frac{M}{5}\Big) \le \exp\{-2M/25\},
$$

where we also used in the last step that $\frac{M_0}{2c_3} - \frac{1+B}{\beta} - 1 = 5$. Hence,

$$
\begin{aligned} \mathcal{E}_{\theta}T_3 &= \mathcal{E}_{\theta} \sum_{I \in \mathcal{S}_M} \mathbb{1}_{A_M} \hat{\pi}_I \big(\| \vartheta - \theta \|^2 \ge M_0 r^2(\theta) + M \sigma^2 |X \big) \hat{\pi}(I|X) \\ &\le \exp\{-2M/25\} \mathcal{E}_{\theta} \sum_{I \in \mathcal{I}} \hat{\pi}(I|X) \le \exp\{-2M/25\}. \end{aligned}
$$

This completes the proof of assertion [\(i\)](#page-10-1) since, in view of [\(36\)](#page-27-3), [\(38\)](#page-27-4) and the last display, we established that $E_{\theta}\hat{\pi}(\|\theta-\theta\|^2 \geq M_0r^2(\theta) + M\sigma^2|X) \leq E_{\theta}(T_1 + T_2 + T_3) \leq (2 + C_0)e^{-m_0M}$, with constants $M_0 = 2c_3(6 + \frac{1+B}{\beta}), H_0 = 2 + C_0, m_0 = \min\{\frac{\beta}{8}$ $\frac{\beta}{8}, \frac{c_2\beta}{40(1+}$ $\frac{c_2\beta}{40(1+B)}, \frac{2}{25}$ and C_0 defined in [\(37\)](#page-27-2).

The proof of assertion [\(ii\)](#page-10-2) proceeds along similar lines. Recall the constants $c_1 > 2$, c_2 , c_3 from Lemma [2](#page-25-0) and define $M_1 = 4c_3(1 + B + \beta)/\beta$. Introduce the subfamily of sets

$$
\bar{S}_M = \bar{S}_M(\theta) = \left\{ I \in \mathcal{I} : r^2(I, \theta) \leq 2c_3 r^2(\theta) + \frac{\beta}{6(1+B)} M \sigma^2 \right\},\
$$

and the event $\bar{A}_M = \bar{A}_M(\theta) = \left\{ \sum_{i=1}^{\bar{m}} \xi_{[i]}^2 \le \frac{(1+B)}{\beta} \bar{m} \log(\frac{en}{\bar{m}}) + \frac{M}{6} \right\}$, where $\bar{m} = \bar{m}_M(\theta) =$ $\max\{|I| : I \in \bar{S}_M\}.$ Introduce the notation $\bar{\Delta}_M(\theta) = M_1 r^2(\theta) + M\sigma^2$ for brevity. By the definition of $\hat{\theta}$ and the Cauchy-Schwartz inequality, we have that $\|\hat{\theta} - \theta\|^2 \leq \sum_{I \in \mathcal{I}} \|X(I) - \theta\|^2$ $\theta\|^2\hat{\pi}(I|X)$, where $\|X(I) - \theta\|^2 = \sigma^2 \sum_{i \in I} \xi_i^2 + \sum_{i \in I^c} \theta_i^2$. Using this, we derive

$$
\begin{split} \mathbf{P}_{\theta} \big(\| \hat{\theta} - \theta \|^{2} \geq \bar{\Delta}_{M}(\theta) \big) &\leq \mathbf{P}_{\theta} \Big(\sum_{I \in \mathcal{I}} \| X(I) - \theta \|^{2} \hat{\pi}(I|X) \geq \bar{\Delta}_{M}(\theta) \Big) \\ &\leq \mathbf{P}_{\theta}(\bar{A}_{M}^{\epsilon}) + \mathbf{P}_{\theta} \Big(\Big\{ \sum_{I \in \bar{\mathcal{S}}_{M}} \Big[\sigma^{2} \sum_{i \in I} \xi_{i}^{2} + \sum_{i \in I^{c}} \theta_{i}^{2} \Big] \hat{\pi}(I|X) \geq \bar{\Delta}_{M}(\theta)/2 \Big\} \cap \bar{A}_{M} \Big) \\ &+ \mathbf{P}_{\theta} \Big(\sum_{I \in \bar{\mathcal{S}}_{M}^{\epsilon}} \Big[\sigma^{2} \sum_{i \in I} \xi_{i}^{2} + \sum_{i \in I^{c}} \theta_{i}^{2} \Big] \hat{\pi}(I|X) \geq \bar{\Delta}_{M}(\theta)/2 \Big) = \bar{T}_{1} + \bar{T}_{2} + \bar{T}_{3}. \end{split}
$$

Similar to [\(36\)](#page-27-3), we bound the term \bar{T}_1 by Lemma [3](#page-26-0) (see also [\(35\)](#page-27-1)):

$$
\bar{T}_1 = P_{\theta}(\bar{A}_M^c) = P\Big(\sum_{i=1}^{\bar{m}} \xi_{[i]}^2 > \frac{(1+B)}{\beta} \bar{m} \log(\frac{en}{\bar{m}}) + \frac{M}{6}\Big) \le \exp\Big\{-M\beta/6\Big\}.
$$

Now we evaluate the term \bar{T}_2 . Since $\bar{m} = \max\{|I| : I \in \bar{S}_M\}, \sigma^2 \bar{m} \log(\frac{en}{\bar{m}}) \leq 2c_3 r^2(\theta) +$ $\frac{\beta}{6(1+B)}M\sigma^2$. Then for any $I \in \bar{\mathcal{S}}_M$, the event \bar{A}_M implies that $\sum_{i\in I}\xi_i^2 \leq \sum_{i=1}^{\bar{m}}\xi_{[i]}^2 \leq$ $\frac{(1+B)}{\beta} \bar{m} \log(\frac{en}{\bar{m}}) + \frac{M}{6} \leq \frac{2c_3(1+B)}{\beta}$ β $\frac{r^2(\theta)}{\sigma^2}+\frac{M}{3}$ $\frac{M}{3}$. Also $\sum_{i \in I^c} \theta_i^2 \leq r^2(I,\theta) \leq 2c_3 r^2(\theta) + \frac{\beta}{6(1+B)} M \sigma^2$ for any $I \in \overline{S}_M$. Hence, for any $I \in \overline{S}_M$, we obtain the implication

$$
\bar{A}_M \subseteq \Big\{ \sigma^2 \sum_{i \in I} \xi_i^2 + \sum_{i \in I^c} \theta_i^2 \le \frac{2c_3(1 + B + \beta)}{\beta} r^2(\theta) + \left(\frac{1}{3} + \frac{\beta}{6(1 + B)}\right) M \sigma^2 \Big\}.
$$

As $M_1 = 4c_3(1 + B + \beta)/\beta$, $\beta \in (0, 1]$ and $B > 0$, the last relation entails

$$
\overline{T}_2 = P_\theta \Big(\Big\{ \sum_{I \in \overline{S}_M} \Big(\sigma^2 \sum_{i \in I} \xi_i^2 + \sum_{i \in I^c} \theta_i^2 \Big) \hat{\pi}(I|X) \ge \frac{\overline{\Delta}_M}{2} \Big\} \cap \overline{A}_M \Big) \le P_\theta \Big(\frac{2c_3(1+B+\beta)}{\beta} r^2(\theta) + \left(\frac{1}{3} + \frac{\beta}{6(1+B)}\right) M \sigma^2 \ge \frac{M_1}{2} r^2(\theta) + \frac{M}{2} \sigma^2 \Big) = 0.
$$

It remains to handle the term \bar{T}_3 . Applying first the Markov inequality and then the Cauchy-Schwarz inequality, we obtain

$$
\bar{T}_3 \leq \frac{\mathrm{E}_{\theta}\left(\sum_{I \in \bar{\mathcal{S}}_M^c} \left[\sigma^2 \sum_{i \in I} \xi_i^2 + \sum_{i \in I^c} \theta_i^2\right] \hat{\pi}(I|X)\right)}{\bar{\Delta}_M(\theta)/2} \n\leq \frac{\sum_{I \in \bar{\mathcal{S}}_M^c} \left(\sigma^2 \left[\mathrm{E}_{\theta}\left(\sum_{i \in I} \xi_i^2\right)^2\right]^{1/2} \left[\mathrm{E}_{\theta}(\hat{\pi}(I|X))^2\right]^{1/2} + r^2(I,\theta)\mathrm{E}_{\theta}\hat{\pi}(I|X)\right)}{\bar{\Delta}_M(\theta)/2} = T_{31} + T_{32}.
$$

For any $I \in \bar{\mathcal{S}}_M^c$, we have $c_3 r^2(\theta) \leq \frac{r^2(I,\theta)}{2} - \frac{\beta}{12(I+B)} M \sigma^2$, yielding the bound

$$
\frac{c_2}{2} \left(r^2(I,\theta) - c_3 r^2(\theta) \right) \ge C_1 r^2(I,\theta) + C_2 M \sigma^2 \quad \text{for any } I \in \bar{\mathcal{S}}_M^c,
$$
\n
$$
\tag{40}
$$

where $C_1 = c_2/4$ and $C_2 = c_2\beta/[24(1+B)]$. By [\(40\)](#page-29-0) and Lemma [2,](#page-25-0)

$$
\left[\mathcal{E}_{\theta}\hat{\pi}(I|X)\right]^{1/2} \leq \left(\frac{ne}{|I|}\right)^{-c_1|I|/2} \exp\left\{-C_1\sigma^{-2}r^2(I,\theta) - C_2M\right\} \text{ for any } I \in \bar{\mathcal{S}}_M^c. \tag{41}
$$

Since $c_1 > 2$, [\(37\)](#page-27-2) gives $\sum_{I \in \mathcal{I}} \left(\frac{ne}{|I|} \right)$ $\frac{ne}{|I|}$ ^{-c₁|I|/2} $\leq (1 - e^{-c_1/2})^{-1} \triangleq C_3$. According to [\(15\)](#page-7-2) with $\rho = \min\{C_1, B/2\}, \left[\mathbb{E}\left(\sum_{i \in I} \xi_i^2\right)^2\right]^{1/2} \leq \frac{B}{\beta \rho} \exp\{\rho |I|\}.$ If $M \in [0, 1]$, the claim (ii) holds for any $H_1 \geq e^{m_1}$. Let $M \geq 1$, then $\sigma^2/\overline{\Delta}_M(\theta) \leq M^{-1} \leq 1$. Besides, $\sigma^{-2}r^2(I,\theta) \geq |I| \log(en/|I|) \geq |I|$. Piecing all these relations together with [\(41\)](#page-30-0), we derive

$$
T_{31} \leq \frac{2B}{\beta \rho} \sum_{I \in \bar{\mathcal{S}}_M^c} \exp\{\rho |I|\} \left(\frac{ne}{|I|}\right)^{-c_1 |I|/2} \exp\left\{-C_1 \sigma^{-2} r^2 (I,\theta) - C_2 M\right\} \leq C_4 \exp\{-C_2 M\},\
$$

where $C_4 = 2BC_3/(\beta \rho) = 2BC_3/(\beta \min\{C_1, B\})$. Finally, by [\(37\)](#page-27-2), [\(41\)](#page-30-0) and the facts that $\max_{x\geq 0} \{xe^{-cx}\}\leq (ce)^{-1}$ (for any $c>0$) and $\sigma^2/\Delta_M(\theta)\leq 1$, we bound the term T_{32} :

$$
T_{32} = \frac{2}{\Delta_M(\theta)} \sum_{I \in \bar{\mathcal{S}}_M^c} r^2(I, \theta) \mathbb{E}_{\theta} \hat{\pi}(I|X)
$$

\$\leq \frac{2}{\Delta_M(\theta)} \sum_{I \in \bar{\mathcal{S}}_M^c} r^2(I, \theta) \left(\frac{ne}{|I|}\right)^{-c_1|I|} \exp \left\{-2C_1 \sigma^{-2} r^2(I, \theta) - 2C_2 M\right\} \leq C_5 \exp\{-2C_2 M\},

where $C_5 = C_0/(C_1e)$. The assertion [\(ii\)](#page-10-2) is proved since we showed that $P_\theta(\|\hat{\theta}-\theta\|^2 \ge M_1r^2(\theta) +$ $(M\sigma^2) \leq H_1 e^{-m_1 M}$ with $M_1 = 4c_3(1+B+\beta)/\beta$, $H_1 = (1+C_4+C_5) \vee e^{m_1}$, $m_1 = \min\{\frac{\beta}{6}$ $\frac{6}{6}, C_2\}.$

Proof of Theorem [2.](#page-11-2) First we prove (i). If the inequality $|I \setminus I_o| \log(\frac{en}{|I|}) < \sum_{i \in I \setminus I_o}$ $\frac{\theta_i^2}{\sigma^2}$ would hold for some $I \in \mathcal{I}$, then

$$
r^2(I \cup I_o, \theta) = \sum_{i \notin I \cup I_o} \theta_i^2 + \sigma^2 |I \cup I_o| \log(\frac{en}{|I \cup I_o|}) \le \sum_{i \notin I \cup I_o} \theta_i^2 + \sigma^2 |I \setminus I_o| \log(\frac{en}{|I|}) + \sigma^2 |I_o| \log(\frac{en}{|I_o|})
$$

$$
< \sum_{i \notin I \cup I_o} \theta_i^2 + \sum_{i \in I \setminus I_o} \frac{\theta_i^2}{\sigma^2} + \sigma^2 |I_o| \log(\frac{en}{|I_o|}) = \sum_{i \notin I_o} \theta_i^2 + \sigma^2 |I_o| \log(\frac{en}{|I_o|}) = r^2(\theta),
$$

which contradicts the definition of the oracle. Hence, $\sum_{i \in I \setminus I_o}$ $\frac{\theta_i^2}{\sigma^2} \leq |I \setminus I_o| \log(\frac{en}{|I|})$ for any $I \in \mathcal{I}$. $|I|$ Define $c_4 = \varkappa \beta - \frac{\beta}{2} - B - 1$ and note that $c_4 > 1$ by the condition of the theorem. Using the relation $\sum_{i\in I\setminus I_o}$ $\frac{\theta_i^2}{\sigma^2} \leq |I \setminus I_o| \log(\frac{en}{|I|}) \leq |I| \log(\frac{en}{|I|})$ and Lemma [1](#page-24-1) with $h = \beta$ and $I_0 = I_o \cap I$ (so that $I\setminus I_0 = I\setminus I_o$, we obtain for each $I \in \mathcal{G}_1 = \{I \in \mathcal{I} : |I|\log(\frac{en}{|I|}) \geq M'_0 |I_0|\log(\frac{en}{|I_0|}) + M\}$ with $M_0'=\varkappa\beta+\frac{\beta}{2}$ $\frac{5}{2}$

$$
\begin{split} \mathcal{E}_{\theta}\hat{\pi}(I|X) &\leq \left[\frac{\lambda_{I}}{\lambda_{I_{0}}}\right]^{\beta} \exp\left\{\beta\sum_{i\in I\backslash I_{0}}\frac{\theta_{i}^{2}}{\sigma^{2}}+\frac{\beta}{2}|I_{0}|\log(\frac{en}{|I_{0}|})-(\frac{\beta}{2}-B)|I|\log(\frac{en}{|I|})\right\} \\ &\leq (\frac{ne}{|I|})^{-c_{4}|I|} \exp\left\{-\left(\varkappa\beta-\frac{\beta}{2}-B-c_{4}\right)|I|\log(\frac{en}{|I|})+(\beta\varkappa+\frac{\beta}{2})|I_{0}|\log(\frac{en}{|I_{0}|})\right\} \\ &=\left(\frac{ne}{|I|}\right)^{-c_{4}|I|} \exp\left\{-|I|\log(\frac{en}{|I|})+(\beta\varkappa+\frac{\beta}{2})|I_{0}|\log(\frac{en}{|I_{0}|})\right\} \\ &\leq (\frac{ne}{|I|})^{-c_{4}|I|} \exp\left\{-\left(M_{0}'-\varkappa\beta-\frac{\beta}{2}\right)|I_{0}|\log(\frac{en}{|I_{0}|})-M\right\}=(\frac{ne}{|I|})^{-c_{4}|I|}e^{-M} .\end{split}
$$

Since $c_4 > 1$, by the same reasoning as in [\(6\)](#page-6-6) we bound $\sum_{I \in \mathcal{I}} \left(\frac{ne}{|I|} \right)$ $\frac{ne}{|I|}$ $\big)^{-c_4|I|} \le (1 - e^{1 - c_4})^{-1} \triangleq H'_0.$ Using this and the last display, we finish the proof of (i):

$$
\mathcal{E}_{\theta}\hat{\pi}(I \in \mathcal{G}_1|X) = \sum_{I \in \mathcal{G}_1} \mathcal{E}_{\theta}\hat{\pi}(I|X) \leq e^{-M} \sum_{I \in \mathcal{I}} \left(\frac{ne}{|I|}\right)^{-c_4|I|} \leq H'_0 e^{-M}.
$$

Next we prove (ii). Define $\mathcal{G}_2 = \mathcal{G}_2(I') = \{I \in \mathcal{I} : \sum_{i \in I' \setminus I}$ $\frac{\theta_i^2}{\sigma^2} \geq \overline{\tau} |I \cup I'| \log(\frac{en}{|I \cup I'|}) + M$. Using [\(5\)](#page-5-1) and Lemma [1](#page-24-1) with $h = \beta$ and $I_0 = I_0(I, \theta) = I \cup I'$, we evaluate for each $I \in \mathcal{G}_2$

$$
\begin{split} \mathcal{E}_{\theta}\hat{\pi}(I|X) &\leq \left[\frac{\lambda_{I}}{\lambda_{I_{0}}}\right]^{\beta}\exp\left\{-\frac{\beta}{3}\sum_{i\in I_{0}\setminus I}\frac{\theta_{i}^{2}}{\sigma^{2}}+(\frac{\beta}{2}+B)|I_{0}|\log\left(\frac{en}{|I_{0}|}\right)-\frac{\beta}{2}|I|\log\left(\frac{en}{|I|}\right)\right\} \\ &=\left[\frac{\lambda_{I}}{c_{\varkappa,n}}\right]^{\beta}\exp\left\{-\frac{\beta}{3}\sum_{i\in I'\setminus I}\frac{\theta_{i}^{2}}{\sigma^{2}}+(\varkappa\beta+\frac{\beta}{2}+B)|I\cup I'|\log\left(\frac{en}{|I\cup I'|}\right)-\frac{\beta}{2}|I|\log\left(\frac{en}{|I|}\right)\right\} \\ &\leq \left[\frac{\lambda_{I}}{c_{\varkappa,n}}\right]^{\beta+\frac{\beta}{2\varkappa}}\exp\left\{\left(-\frac{\beta}{3}\bar{\tau}+\varkappa\beta+\frac{\beta}{2}+B\right)|I\cup I'|\log\left(\frac{en}{|I\cup I'|}\right)-\frac{\beta}{3}M\right\}\leq \left[\frac{\lambda_{I}}{c_{\varkappa,n}}\right]^{\beta+\frac{\beta}{2\varkappa}}e^{-\frac{\beta}{3}M}. \end{split}
$$

Since $\varkappa > \beta^{-1} - \frac{1}{2}$ $\frac{1}{2}$, by the same reasoning as in [\(6\)](#page-6-6) we bound $\sum_I \left[\frac{\lambda_I}{c_{\kappa}} \right]$ $\frac{\lambda_I}{c_{\varkappa,n}}\big]^{\beta(1+1/2\varkappa)} \leq (1-\varkappa)$ $e^{1-\varkappa\beta-\beta/2}$ ⁻¹ $\triangleq H'_1$. This relation and the last display imply claim (ii): with $m'_0=\frac{\beta}{3}$ $\frac{p}{3}$,

$$
\mathcal{E}_{\theta}\hat{\pi}\big(I \in \mathcal{G}_2|X\big) = \sum_{I \in \mathcal{G}_2} \mathcal{E}_{\theta}\hat{\pi}(I|X) \le H'_1 \exp\{-m'_0 M\}.
$$
 (42)

Let us derive the second claim of (ii). If $|I| \log(\frac{en}{|I|}) \le \varrho |I_*| \log(\frac{en}{|I_*|}) - M$, then $|I \cup I_*| \log(\frac{en}{|I \cup I_*|}) \le$ $|I|\log(\frac{en}{|I|})+|I_*|\log(\frac{en}{|I_*|})\leq (1+\varrho)|I_*|\log(\frac{en}{|I_*|})-M.$ Hence, $|I_*|\log(\frac{en}{|I_*|})\geq \frac{1}{1+\varrho}$ $\frac{1}{1+\varrho}|I\cup I_*|\log(\frac{en}{|I\cup I_*|})+$ M $\frac{M}{1+\varrho}$, which, together with the definition of the τ -oracle, imply

$$
\sum_{i \in I_* \setminus I} \frac{\theta_i^2}{\sigma^2} \ge \left(\sum_{i \in I^c} \frac{\theta_i^2}{\sigma^2} - \sum_{i \in I_*^c} \frac{\theta_i^2}{\sigma^2} \right) \ge \tau_0 \left(|I_*| \log(\frac{en}{|I_*|}) - |I| \log(\frac{en}{|I|}) \right)
$$
\n
$$
\ge \tau_0 (1 - \varrho) |I_*| \log(\frac{en}{|I_*|}) + \tau M \ge \tau' |I \cup I_*| \log(\frac{en}{|I \cup I_*|}) + \frac{2\tau_0}{1 + \varrho} M,\tag{43}
$$

where $\tau' = \frac{1-\varrho}{1+\varrho}\tau_0 > \bar{\tau}$ by the condition of the theorem. Thus, we obtain

$$
\mathcal{E}_{\theta}\hat{\pi}\big(I:|I|\log(\tfrac{en}{|I|})\leq \varrho|I_*|\log(\tfrac{en}{|I_*|})-M\big|X\big)\leq \mathcal{E}_{\theta}\hat{\pi}\bigg(\sum_{i\in I_*\setminus I}\frac{\theta_i^2}{\sigma^2}\geq \tau'|I\cup I_*|\log(\tfrac{en}{|I\cup I_*|})+\tfrac{2\tau_0}{1+\varrho}M|X\bigg).
$$

By this and [\(42\)](#page-31-1) with $I' = I_*$, the second claim of (ii) follows with $\alpha'_1 = \tau' - \overline{\tau} > 0$ and $m'_1 = \frac{2\tau_0 m'_0}{1+\varrho}.$

Finally, let us prove (iii). Denote $\mathcal{G}_3 = \mathcal{G}_3(\theta, M) = \{I : r^2(I, \theta) \ge c_3 r^2(\theta) + M\sigma^2\}$, where the constants $c_1 > 2$ $c_1 > 2$, c_2 , c_3 are defined in Lemma [2.](#page-25-0) Applying Lemma 2 and using the fact [\(37\)](#page-27-2), we complete the proof of (iii):

$$
\mathrm{E}_{\theta}\hat{\pi}\big(I\in\mathcal{G}_3\big|X\big)=\sum_{I\in\mathcal{G}_3}\mathrm{E}_{\theta}\hat{\pi}(I\big|X)\leq e^{-c_2M}\sum_{I\in\mathcal{I}}\big(\frac{ne}{|I|}\big)^{-c_1|I|}\leq C_0e^{-c_2M}.\quad\Box
$$

Proof of Theorem [3.](#page-13-1) The biggest part of the proof is already contained in Theorem [2.](#page-11-2) We first establish the coverage property. The constants M_1 , H_1 and m_1 are defined in Theorem [1,](#page-10-0) the constant ϱ is from [\(22\)](#page-10-4). Take some $M_2 > \frac{M_1}{\varrho}$, for example $M_2 = \frac{M_1}{\varrho} + 1$. From [\(18\)](#page-9-0) and [\(27\)](#page-13-2), it follows that $r^2(\theta) \le r^2(I_*, \theta) = (b(\theta) + 1)\sigma^2 |I_*| \log(\frac{en}{|I_*|}) + b(\theta)\sigma^2 \le (b(\theta) + 1)\sigma^2(|I_*| \log(\frac{en}{|I_*|}) + 1).$ Combining this with claims (ii) from Theorems [1](#page-10-0) and [2](#page-11-2) and the definition [\(25\)](#page-12-0) of \hat{r} yields the coverage property:

$$
P_{\theta}(\theta \notin B(\hat{\theta}, [(b(\theta) + 1)M_2\hat{r}^2 + (b(\theta) + 2)M\sigma^2]^{1/2})
$$

\n
$$
\leq P_{\theta}(|\hat{\theta} - \theta|^2 > (b(\theta) + 1)M_2\hat{r}^2 + (b(\theta) + 2)M\sigma^2, \hat{r}^2 \geq \varrho\sigma^2|I_*|\log(\frac{en}{|I_*|}) + \sigma^2 - \frac{M\sigma^2}{M_2})
$$

\n
$$
+ P_{\theta}(\hat{r}^2 < \varrho\sigma^2|I_*|\log(\frac{en}{|I_*|}) + \sigma^2 - \frac{M\sigma^2}{M_2})
$$

\n
$$
\leq P_{\theta}(|\hat{\theta} - \theta|^2 > \varrho M_2r^2(\theta) + M\sigma^2) + P_{\theta}(|\hat{I}|\log(\frac{en}{|\hat{I}|}) < \varrho|I_*|\log(\frac{en}{|I_*|}) - \frac{M}{M_2})
$$

\n
$$
\leq H_1(\frac{en}{|I_o|})^{-\alpha_1|I_o|}e^{-m_1M} + H'_1(\frac{en}{|I_*|})^{-\alpha'_1|I_*|}e^{-m''_1M} \leq H_2e^{-m_2M},
$$

where $\alpha_1 = \varrho M_2 - M_1$, $m_1'' = m_1'/M_2$, $H_2 = H_1 + H_1'$, $m_2 = m_1 \wedge m_1''$; H_1', α_1', m_1' are defined in Theorem [2](#page-11-2) and the constant ρ is from [\(22\)](#page-10-4). The first claim of the theorem follows.

The size property follows from the definition (25) of \hat{r} , Remark [7](#page-24-3) and property (i) of Theorem [2.](#page-11-2) Indeed, $P_{\theta}(\hat{r}^2 \geq \sigma^2(M'_0+\alpha)|I_o|\log(\frac{en}{|I_o|})+(M+1)\sigma^2) = P_{\theta}(|\hat{I}|\log(\frac{en}{|\hat{I}|}) \geq (M'_0+\alpha)|I_o|\log(\frac{en}{|I_o|})+$ M) $\leq P_{\theta}(|\hat{I}| \log(\frac{en}{|\hat{I}|}) \geq M'_{0}|I \cap I_{o}| \log(\frac{en}{|I \cap I_{o}|}) + \alpha |I_{o}| \log(\frac{en}{|I_{o}|}) + M$) $\leq H'_{0}(\frac{ne}{|I_{o}|})$ $\frac{ne}{|I_o|}$ $\rho^{-\alpha|I_o|}e^{-M}.$ \Box

Proof of Theorem [4.](#page-15-2) Since $X' = \theta + \xi'$, we rewrite [\(28\)](#page-15-1) as

$$
\tilde{R}_M^2 = (||X' - \hat{\theta}||^2 - \sigma^2 E ||\xi'||^2 + 2\sigma^2 G_M \sqrt{n})_+ \n= (||\theta - \hat{\theta}||^2 + \sigma^2 (||\xi'||^2 - E||\xi'||^2) + 2\sigma \langle \xi', (\theta - \hat{\theta}) \rangle + 2\sigma^2 G_M \sqrt{n})_+.
$$
\n(44)

Introduce the events $\mathcal{C}_M = \left\{2|\langle \xi',(\theta - \hat{\theta}) \rangle| < \sqrt{M(M_1 r^2(\theta) + M \sigma^2)} \right\}$ and $\mathcal{D}_M = \left\{\|\hat{\theta} - \theta\|^2 < \infty\right\}$ $M_1r^2(\theta) + M\sigma^2$. According to [\(A3\)](#page-15-0), $\hat{\theta}$ and \hat{I} are independent of ξ' . Using this fact, the first relation from [\(A3\)](#page-15-0) and Theorem [1,](#page-10-0) we obtain that

$$
P_{\theta}(C_M^c) = E_{\theta} P_{\theta}(C_M^c \cap \mathcal{D}_M | X) + P_{\theta}(C_M^c \cap \mathcal{D}_M^c)
$$

\$\leq E_{\theta} \left[\psi_1 \left(\frac{M(M_1 r^2(\theta) + M \sigma^2)}{4 \|\hat{\theta} - \theta\|^2} \right) \mathbb{1}_{\mathcal{D}_M} \right] + P_{\theta}(\mathcal{D}_M^c) \leq \psi_1(M/4) + H_1 e^{-m_1 M}. \tag{45}

Since $r^2(\theta) \leq r^2([n], \theta) = \sigma^2 n$ by the oracle definition [\(18\)](#page-9-0), the event \mathcal{C}_M implies that $2\sigma \langle \xi', (\theta - \xi') \rangle$ $\langle \hat{\theta} \rangle$ > $-\sigma \sqrt{\overline{M(M_1\sigma^2n + M\sigma^2)}} \geq -\sigma^2 G_M \sqrt{n}$. Combining this with [\(44\)](#page-32-0), [\(45\)](#page-32-1) and the second relation from [\(A3\)](#page-15-0) yields the coverage relation:

$$
P_{\theta}(\theta \notin B(\hat{\theta}, \tilde{R}_M)) = P_{\theta}(\theta \notin B(\hat{\theta}, \tilde{R}_M), C_M) + P_{\theta}(\theta \notin B(\hat{\theta}, \tilde{R}_M), C_M^c)
$$

\n
$$
\leq P_{\theta}(\|\theta - \hat{\theta}\|^2 \geq \tilde{R}_M^2, C_M) + P_{\theta}(C_M^c) \leq P_{\theta}(0 \geq \sigma^2(\|\xi'\|^2 - E\|\xi'\|^2) + \sigma^2 G_M \sqrt{n}) + P_{\theta}(C_M^c)
$$

\n
$$
\leq P_{\theta}(\|\xi'\|^2 - E\|\xi'\|^2 \leq -M\sqrt{n}) + \psi_1(M/4) + H_1 e^{-m_1 M} \leq \psi_2(M) + \psi_1(M/4) + H_1 e^{-m_1 M}.
$$

Let us show the size property. By [\(45\)](#page-32-1), $P_{\theta}(2\sigma \langle \xi', (\theta - \hat{\theta}) \rangle \geq \sigma^2 G_M \sqrt{n}) \leq P_{\theta}(2\langle \xi', (\theta - \hat{\theta}) \rangle >$ $\sqrt{M(M_1r^2(\theta) + M\sigma^2)} \le P_\theta(\mathcal{C}_M^c) \le \psi_1(M/4) + H_1e^{-m_1M}$. This, Theorem [1](#page-10-0) and [\(44\)](#page-32-0) imply

$$
P_{\theta}(\tilde{R}_{M}^{2} \ge g_{M}(\theta, n)) \le P_{\theta}(\|\theta - \hat{\theta}\|^{2} \ge M_{1}r^{2}(\theta) + M\sigma^{2}) + P_{\theta}(\sigma^{2}(\|\xi'\|^{2} - \mathbb{E}\|\xi'\|^{2}) \ge \sigma^{2}G_{M}\sqrt{n}) + P_{\theta}(2\sigma\langle\xi', (\theta - \hat{\theta})\rangle \ge \sigma^{2}G_{M}\sqrt{n}) \le H_{1}e^{-m_{1}M} + \psi_{2}(M) + \psi_{1}(M/4) + H_{1}e^{-m_{1}M}.
$$

Proof of Theorem [5.](#page-17-1) Observe that $r^2(\theta) \le r^2(I^*(\theta), \theta) \le \sigma^2 s(\theta) \log(\frac{en}{s(\theta)})$. Since the function $x \mapsto x \log(en/x)$ is increasing over $(0, n]$, $|I| \geq Ms(\theta)$ implies that $r^2(I, \theta) \geq \sigma^2|I| \log(\frac{en}{|I|}) \geq$ $\sigma^2 M s(\theta) \log(\frac{en}{Ms(\theta)})$. Thus, if $|I| \geq M s(\theta)$, then

$$
r^2(I,\theta) \ge M\sigma^2 s(\theta) \log(\frac{en}{Ms(\theta)}) \ge M_4 r^2(\theta) - M_4 \sigma^2 s(\theta) \log(\frac{en}{s(\theta)}) + M \sigma^2 s(\theta) \log(\frac{en}{Ms(\theta)}).
$$

The first claim follows from Theorem [2](#page-11-2) with $M_4 = c_3$ and $m_4 = c_2$.

To prove the second claim, note that for any $M' > 2M_4$, $|I| \ge M' s(\theta)$ implies that

$$
r^2(I,\theta) \ge \sigma^2|I|\log(en/|I|) \ge M'\sigma^2s(\theta)\left[\log(en/s(\theta)) - \log M'\right] \ge \frac{M'}{2}\sigma^2s(\theta)\log(en/s(\theta)),
$$

provided that $s(\theta) < en/(M')^2$. Since $r^2(\theta) \le r^2(I^*(\theta), \theta) \le \sigma^2 s(\theta) \log(en/s(\theta))$, the relation above implies that $r^2(I,\theta) \ge M_4 r^2(\theta) + M\sigma^2$, where $M = (M'/2 - M_4)s(\theta) \log(en/s(\theta))$. Hence by Theorem [2,](#page-11-2) the assertion holds for $M'_4 = M'$ whenever $s(\theta) < en/(M')^2$. If $s(\theta) \geq en/(M')^2$, the result trivially holds by choosing $M'_4 = (M')^2/e$. Hence the choice $M'_4 \ge \max\{M', (M')^2/e\}$ ensures the result with $m'_4 = m_4(M'/2 - M_4)$ for any $\theta \in \mathbb{R}^n$.

Proof of Theorem [6.](#page-18-3) Recall [\(30\)](#page-18-4): $\sigma^{-2} r^2(\theta) \leq Kn(\frac{p_n}{n\sigma})$ $\frac{p_n}{n\sigma}$ ^s $\left[\log(\frac{n\sigma}{p_n})\right]^{1-s/2}$ for each $\theta \in m_s[p_n]$ with some $K = K(s)$. On the other hand, if $|I| > Mp_n^* = Men(\frac{p_n}{n\sigma})$ $\frac{p_n}{n\sigma}$ ^s $\left[\log(\frac{n\sigma}{p_n})\right]^{-s/2}$, then $\sigma^{-2}r^2(I,\theta) \ge$ $|I|\log(\frac{en}{|I|}) \geq M p_n^* \log(\frac{en}{Mp_n^*}) = M p_n^* \left[s \log(\frac{nc}{p_n}) + \frac{s}{2} \log \log(\frac{nc}{p_n}) - \log(M)\right] \geq M s p_n^* \log(\frac{nc}{p_n})$ for sufficiently large *n* as $p_n = o(n)$. Then, for any $\theta \in m_s[p_n]$, $M > c_3K/(se)$ and $|I| > Mp_n^*$, we have that, for sufficiently large n ,

$$
\sigma^{-2}(r^2(I,\theta) - c_3r^2(\theta)) \ge M p_n^* \log\left(\frac{en}{Mp_n^*}\right) - c_3 K n \left(\frac{p_n}{n\sigma}\right)^s \left[\log\left(\frac{nc}{p_n}\right)\right]^{1-s/2}
$$

\n
$$
\ge M s p_n^* \log\left(\frac{nc}{p_n}\right) - c_3 K e^{-1} p_n^* \log\left(\frac{nc}{p_n}\right) = (Ms - c_3 K e^{-1}) p_n^* \log\left(\frac{nc}{p_n}\right).
$$

Finally, applying Theorem [2,](#page-11-2) we obtain

$$
\sup_{\theta \in m_s[p_n]} \mathbb{E}_{\theta} \hat{\pi}(I : |I| > M p_n^* | X) \le C_0 \exp \big\{ -c_2 s \big(M - c_3 K (se)^{-1} \big) p_n^* \log(\frac{n\sigma}{p_n}) \big\},
$$

 \Box

which gives the claim with $m_5 = c_2 s$ and $M_5 = c_3 K(se)^{-1}$.

References

[1] Abramovich, F., Grinshtein, V. and Pensky, M. (2007). On optimality of Bayesian testimation in the normal means problem. Ann. Statist. 35, 2261–2286.

- [2] Babenko, A. and Belitser, E. (2010). Oracle projection convergence rate of posterior. Math. Meth. Statist. 19, 219–245.
- [3] Baraud, Y. (2004). Confidence balls in Gaussian regression. Ann. Statist. 32, 528–551.
- [4] Belitser, E. (2017). On coverage and local radial rates of credible sets. Ann. Statist. 45, 1124–1151.
- [5] Belitser, E. and Nurushev, N. (2017). Discussion on article by van der Pas, Szabo, and van der Vaart. Bayesian Analysis 12, 1267–1269.
- [6] Birg´e, L. and Massart, P. (2001). Gaussian model selection. J. Eur. Math. Soc. 3, 203– 268.
- [7] BHATTACHARYA, A., DANSON D.B., PATI D. and PILLAI, N.S. (2016). Sub-optimality of some continuous shrinkage priors. Stoch. Proc. and their Appl. 126, 3828–3842.
- [8] BULL, A. (2012). Honest adaptive confidence bands and self-similar functions. *Electron. J.* Statist. 6, 1490–1516.
- [9] BULL, A. and NICKL, R. (2013). Adaptive confidence sets in ℓ_2 . Probab. Theory and Rel. Fields. 156, 889–919.
- [10] CAI, T.T. and LOW, M.G. (2004). An adaptation theory for nonparametric confidence intervals. Ann. Statist. 32, 1805–1840.
- [11] CARVALHO, C.M., POLSON N.G. and SCOTT J.G. (2010). The horseshoe estimator for sparse signals. Biometrika 97, 465–480.
- [12] Castillo, I., Schmidt-Hieber, J. and van der Vaart, A. (2015). Bayesian linear regression with sparse priors. Ann. Statist. 43, 1986–2018.
- [13] Castillo, I. and van der Vaart, A. (2012). Needles and straw in a haystack: posterior concentration for possibly sparse sequences. Ann. Statist. 40, 2069–2101.
- [14] DONOHO, D.L., JOHNSTONE, I.M., HOCH, J.C. and STERN, A.S. (1992). Maximum entropy and the nearly black object(with Discussion). J. Roy. Statist. Soc. Ser. B 54, 41–81.
- [15] DONOHO, D.L. and JOHNSTONE, I.M. (1994). Ideal spatial adaptation by wavelet shrinkage. Biometrika 81, 425–455.
- [16] DONOHO, D.L. and JOHNSTONE, I.M. (1994). Minimax risk over ℓ_p -balls for ℓ_q -error. Probab. Theory Related Fields. 99, 277–303.
- [17] Johnstone, I. and Silverman, B. (2004). Needles and straw in haystacks: empirical Bayes estimates of possibly sparse sequences. Ann. Statist. 32, 1594–1649.
- [18] Li, K.-C. (1989). Honest confidence regions for nonparametric regression. Ann. Statist. 17, 1001–1008.
- [19] Martin, R. and Walker, S.G. (2014). Asymptotically minimax empirical Bayes estimation of a sparse normal mean vector. Electron. J. Statist. 8, 2188–2206.
- [20] Nickl, R. and van de Geer, S. (2013). Confidence sets in sparse regression. Ann. Statist. 41, 2852–2876.
- [21] PICARD, D. and TRIBOULEY, K. (2000). Adaptive confidence interval for pointwise curve estimation. Ann. Statist. 28, 298–335.
- [22] ROBINS, J. and VAN DER VAART, A.W. (2006). Adaptive nonparametric confidence sets. Ann. Statist. 34, 229–253.
- [23] Rockova, V. (2018). Bayesian estimation of sparse signals with a continuous spike-andslab prior. Ann. Statist. 46, 401–437.
- [24] ROUSSEAU, J. and SZABÓ, B. (2016). Asymptotic frequentist coverage properties of Bayesian credible sets for sieve priors. [arXiv:1609.05067.](http://arxiv.org/abs/1609.05067)
- [25] ROUSSEAU, J. and SZABÓ, B. (2017) . Asymptotic behaviour of the empirical Bayes posteriors associated to maximum marginal likelihood estimator. Ann. Statist. 45, 833–865.
- [26] SZABÓ, B. T., VAN DER VAART, A.W. and VAN ZANTEN, J.H. (2015). Frequentist coverage of adaptive nonparametric Bayesian credible sets. Ann. Statist. 43, 1391–1428.
- [27] van der Pas, S.L., Kleijn, B.J.K. and van der Vaart, A.W. (2014). The horseshoe estimator: Posterior concentration around nearly black vectors. Electron. J. Stat. 8, 2585– 2618.
- [28] VAN DER PAS, S.L., SZABÓ, B. T. and VAN DER VAART, A.W. (2017). Uncertainty quantification for the horseshoe (with discussion). Bayesian Anal. 12, 1221–1274.