# Global boundedness of solutions in a parabolic-parabolic chemotaxis system with singular sensitivity

Xiangdong Zhao Sining Zheng<sup>\*</sup>

School of Mathematical Sciences, Dalian University of Technology, Dalian 116024, P. R. China

October 13, 2018

#### Abstract

We consider a parabolic-parabolic Keller-Segel system of chemotaxis model with singular sensitivity  $u_t = \Delta u - \chi \nabla \cdot (\frac{u}{v} \nabla v)$ ,  $v_t = k \Delta v - v + u$  under homogeneous Neumann boundary conditions in a smooth bounded domain  $\Omega \subset \mathbb{R}^n$   $(n \geq 2)$ , with  $\chi, k > 0$ . It is proved that for any k > 0, the problem admits global classical solutions, whenever  $\chi \in (0, -\frac{k-1}{2} + \frac{1}{2}\sqrt{(k-1)^2 + \frac{8k}{n}})$ . The global solutions are moreover globally bounded if  $n \leq 8$ . This shows an exact way the size of the diffusion constant k of the chemicals v effects the behavior of solutions.

2010MSC: 35B40; 92C17; 35K55

Keywords: Chemotaxis; Logarithmic sensitivity; Boundedness

# 1 Introduction

In this paper, we consider the parabolic-parabolic chemotaxis system with singular sensitivity

$$\begin{aligned} u_t &= \Delta u - \chi \nabla \cdot \left(\frac{u}{v} \nabla v\right), & x \in \Omega, \quad t \in (0, T), \\ v_t &= k \Delta v - v + u, & x \in \Omega, \quad t \in (0, T), \\ \frac{\partial u}{\partial \nu} &= \frac{\partial v}{\partial \nu} = 0, & x \in \partial \Omega, \quad t \in (0, T), \\ (u(x, 0), v(x, 0)) &= (u_0(x), v_0(x)), \quad x \in \Omega, \end{aligned}$$

$$(1.1)$$

where  $\chi, k > 0$ ,  $\Omega$  is a smooth bounded domain in  $\mathbb{R}^n$   $(n \ge 2)$ ,  $\frac{\partial}{\partial \nu}$  denotes the derivation with respect to the outer normal of  $\partial\Omega$ , and the initial datum  $u_0 \in C^0(\overline{\Omega}), u_0(x) \ge 0$  on  $\overline{\Omega},$  $v_0 \in W^{1,q}(\Omega)$   $(q > n), v_0(x) > 0$  on  $\overline{\Omega}$ .

The classical Keller-Segel system of chemotaxis model was introduced by Keller and Segel [9] in 1970 to describe the cells (with density u) move towards the concentration gradient of a chemical substance v produced by the cells themselves. Various forms of sensitivity

<sup>\*</sup>Corresponding author. E-mail: snzheng@dlut.edu.cn (S. N. Zheng), 704456001@qq.com (X. D. Zhao)

functions can be chosen to model different types of chemotaxis mechanisms. Among them  $\phi(v) = \frac{\chi}{v}$  was selected in (1.1) largely due to the Weber-Fechner's law for cellular behaviors, where the subjective sensation is proportional to the logarithm of the stimulus intensity [6]. With  $\phi(v) = \frac{\chi}{v}$ , the cellular movements are governed by the taxis flux  $\frac{\chi \nabla v}{v}$ , which may be unbounded when  $v \approx 0$ . In the model (1.1), the values of the chemotactic sensitivity coefficient  $\chi$  and the chemicals diffusion constant k play significant roles to determine the behavior of solutions. Obviously, either small  $\chi > 0$  or large k > 0 benefits the global boundedness of solutions.

Recall the known results in the field with k = 1. At first consider the parabolic-elliptic analogue of (1.1), namely, the second parabolic equation in (1.1) is replaced by the elliptic equation  $0 = \Delta v - v + u$ . It was known that all radial classical solutions are global-in-time if either  $n \ge 3$  with  $\chi < \frac{2}{n-2}$ , or n = 2 with  $\chi > 0$  arbitrary [11]. When  $0 < \phi(v) < \frac{\chi}{v^l}$  with  $l \ge 1$ ,  $\chi > 0$ , there is a unique and globally bounded classical solution if  $\chi < \frac{2}{n}$  (l = 1) or  $\chi < \frac{2}{n} \cdot \frac{l^l}{(l-1)^{(l-1)}} \gamma^{(l-1)}$  (l > 1), with  $\gamma > 0$  depending on  $\Omega$  and  $u_0$  [3]. Next consider the parabolic-parabolic case. All solutions of (1.1) are global in time when either n = 1 [13], or n = 2 and  $\chi < \frac{5}{2}$  under the radial assumption, while  $\chi < 1$  under the non-radial assumption [1, 12]. For  $n \ge 2$ , (1.1) possesses global classical solutions if  $0 < \chi < \sqrt{\frac{2}{n}}$ , and moreover,  $\chi < \sqrt{\frac{n+2}{3n-4}}$  ensures the global existence of weak solutions [19]. Certain radial weak solutions have been constructed when  $\chi < \sqrt{\frac{n}{n-2}}$  [15]. Following, the global boundedness of solutions has been obtained for  $\chi < \sqrt{\frac{2}{n}}$  [2]. Refer to [4, 10, 21] for the results on more general chemotaxis models.

Recently, under somewhat complicated conditions, Wang [16] established classical global solutions to the problem, a similar model to (1.1),

$$\begin{cases} u_t = \nabla \cdot (\nabla u - \frac{\chi u}{v+c} \nabla v), & x \in \Omega, \ t \in (0,T), \\ v_t = k \Delta v - \alpha v + \beta u, & x \in \Omega, \ t \in (0,T), \\ \frac{\partial u}{\partial \nu} = \frac{\partial v}{\partial \nu} = 0, & x \in \partial \Omega, \ t \in (0,T), \\ (u(x,0), v(x,0)) = (u_0(x), v_0(x)), \ x \in \Omega, \end{cases}$$

with  $\chi, c, k, \alpha, \beta > 0$ . In the present paper, motivated by Winkler [19] and Fujie [2], we will prove the global existence-boundedness of classical solutions to (1.1), with simplified conditions. That is the following theorem.

**Theorem 1** Let  $n \geq 2$ ,  $u_0 \in C^0(\overline{\Omega})$ ,  $v_0 \in W^{1,q}(\Omega)$  (q > n) with  $u_0 \geq 0$ ,  $v_0 > 0$  on  $\overline{\Omega}$ . Then, for any k > 0, there exists a global classical solution to (1.1), provided  $\chi \in (0, -\frac{k-1}{2} + \frac{1}{2}\sqrt{(k-1)^2 + \frac{8k}{n}})$ . Moreover, the solution is globally bounded under  $n \leq 8$ .

**Remark 1** Theorem 1 shows in what way the size of k > 0 (the diffusion strength of the chemicals v) effects the behavior of solutions to (1.1). It is interesting to observe that when

n = 2 the global existence-boundedness of solutions is independent of the size of k > 0, since  $-\frac{k-1}{2} + \frac{1}{2}\sqrt{(k-1)^2 + \frac{8k}{n}} \equiv 1$  with n = 2. Quite differently, when  $n \ge 3$  the contribution of k > 0 is significant that the range of  $\chi$  for global existence-boundedness of solutions will be enlarged (shrunk) as k > 0 is degreasing (increasing). The arbitrariness of k > 0 yields the "maximal" range with  $\chi \in (0, 1)$  or the "minimal" range with  $\chi \in (0, \frac{2}{n})$ . That is to say for any  $\chi \in (0, 1)$  (close to 1), there is k > 0 (small) such that the classical solution of (1.1) is globally bounded. On the other hand, for any k > 0 (large), there is  $\chi \in (0, \frac{2}{n})$  to ensure the global boundedness. Finally, it is pointed out that if k = 1, the required range of  $\chi$  in Theorem 1 becomes  $0 < \chi < \sqrt{\frac{2}{n}}$ , which coincides with those in [2, 19].

**Remark 2** Now compare our results for the parabolic-parabolic chemotaxis model (1.1) with those for the corresponding parabiotic-elliptic model, which can be thought as a special case of (1.1) with the diffusion constant of the chemicals v sufficiently large [3]. Just as mentioned in Remark 1, letting k > 0 be arbitrarily large results in the "minimal" permitted range with  $0 < \chi < \frac{2}{n}$ . This does agree those obtained for the parabolic-elliptic model in [11].

### 2 Preliminaries

In this section we introduce the local existence of classical solutions to (1.1) with required estimates involving  $\chi$  and k, as well as some technical lemmas for the global boundedness as preliminaries.

Lemma 2.1 Let  $n \geq 2$ ,  $u_0 \in C^0(\overline{\Omega})$ ,  $v_0 \in W^{1,q}(\Omega)$  (q > n) with  $u_0 \geq 0$ ,  $v_0 > 0$  on  $\overline{\Omega}$ . Then, for any  $k, \chi > 0$ , there exists  $T_{\max} \in (0, \infty]$ , such that (1.1) has a unique nonnegative solution  $u \in C^0([0, T_{\max}); C^0(\Omega)) \times C^{2,1}(\overline{\Omega} \times (0, T_{\max}))$  and  $v \in C^0([0, T_{\max}); C^0(\Omega)) \times C^{2,1}(\overline{\Omega} \times (0, T_{\max})) \times L^\infty_{\text{loc}}([0, T); W^{1,q}(\Omega))$ , where either  $T_{\max} = \infty$ , or  $T_{\max} < \infty$  with  $\lim_{t \to T_{\max}} \|u(\cdot, t)\|_{L^\infty(\Omega)} + \|v(\cdot, t)\|_{W^{1,q}(\Omega)} = \infty$ .

**Proof.** For k > 0, it is known from Lemma 2.2 in [2] that there is  $\eta > 0$ , such that  $\inf_{x \in \Omega} v(x,t) \ge \eta > 0$  for all t > 0. Consequently, the local existence lemma can be obtained by the classical parabolic theory, refer to Theorem 3.1 in [17].

The following lemma is crucial to establish the global existence-boundedness conclusions of the paper. Denote  $r_{\pm}(p) = (p-1)\left[\frac{p\chi(1-k)+2k}{p(k-1)^2+4k} \pm \frac{2\sqrt{k^2-p\chi k(k-1)-p\chi^2 k}}{p(k-1)^2+4k}\right]$ . Throughout the paper, for simplicity, denote  $T = T_{\text{max}}$ .

**Lemma 2.2** Let (u, v) solve (1.1) with k > 0 and  $\chi \in \left(0, -\frac{k-1}{2} + \frac{1}{2}\sqrt{(k-1)^2 + \frac{8k}{n}}\right)$ . If

$$\|v(\cdot,t)\|_{L^{p-r}(\Omega)} \le c, \quad t \in (0,T)$$
(2.1)

with  $p < \frac{k}{[\chi^2 - \chi(1-k)]_+}$ ,  $r \in (r_-(p), r_+(p))$ , and c > 0, then  $\int_{\Omega} u^p v^{-r} dx \le \tilde{c}, \quad t \in (0,T)$ (2.2) with some  $\tilde{c} > 0$ .

**Proof.** It is known via a simple computation with (1.1) that

$$\begin{split} \frac{d}{dt} \int_{\Omega} u^p v^{-r} dx &= p \int_{\Omega} u^{p-1} v^{-r} [\Delta u - \chi \nabla (\frac{u}{v} \nabla v)] dx - r \int_{\Omega} u^p v^{-r-1} (k \Delta v - v + u) dx \\ &= -p \int_{\Omega} \nabla (u^{p-1} v^{-r}) \cdot (\nabla u - \chi \frac{u}{v} \nabla v) dx + rk \int_{\Omega} \nabla (u^p v^{-r-1}) \cdot \nabla v dx \\ &+ r \int_{\Omega} u^p v^{-r} dx - r \int_{\Omega} u^{p+1} v^{-r-1} dx \\ &= -p(p-1) \int_{\Omega} u^{p-2} v^{-r} |\nabla u|^2 dx + [pr + prk + p(p-1)\chi] \int_{\Omega} u^{p-1} v^{-r-1} \nabla u \cdot \nabla v dx \\ &- [r(r+1)k + pr\chi] \int_{\Omega} u^p v^{-r-2} |\nabla v|^2 dx + r \int_{\Omega} u^p v^{-r} dx - r \int_{\Omega} u^{p+1} v^{-r-1} dx \\ &\leq \int_{\Omega} \Big[ \frac{p[(p-1)\chi + r + rk]^2}{4(p-1)} - pr\chi - r(r+1)k \Big] u^p v^{-r-2} |\nabla v|^2 dx \\ &+ r \int_{\Omega} u^p v^{-r} dx - r \int_{\Omega} u^{p+1} v^{-r-1} dx \end{split}$$

by Young's inequality. Denote

$$f(r; p, \chi, k) = \frac{p[(p-1)\chi + r + rk]^2}{4(p-1)} - pr\chi - r(r+1)k,$$

and rewrite as the quadric expression in r

$$4(p-1)f(r;p,\chi,k) = [p(k-1)^2 + 4k]r^2 + [2p(p-1)\chi(k-1) - 4(p-1)k]r + p(p-1)^2\chi^2.$$

We know

$$\Delta_r = 4(p-1)^2 [p\chi(k-1) - 2k]^2 - 4(p-1)^2 p\chi^2 [p(k-1)^2 + 4k]$$
  
= 16(p-1)^2 [k^2 - p\chi k(k-1) - p\chi^2 k] > 0

whenever  $p < \frac{k}{[\chi^2 + \chi(k-1)]_+}$ . Consequently,  $f(r; p, \chi, k) < 0$  for any  $r \in (r_-(p), r_+(p))$ . This yields

$$\frac{d}{dt} \int_{\Omega} u^{p} v^{-r} dx \le r \int_{\Omega} u^{p} v^{-r} dx - r \int_{\Omega} u^{p+1} v^{-r-1} dx, \quad t \in (0,T).$$
(2.3)

Due to  $\int_{\Omega} u^{p} v^{-r} \leq \left(\int_{\Omega} u^{p+1} v^{-r-1}\right)^{\frac{p}{p+1}} \left(\int_{\Omega} v^{p-r}\right)^{\frac{1}{p+1}}$ , we obtain (2.2) from (2.3) and (2.1).

**Lemma 2.3** Let (u, v) satisfy the second equation of (1.1) with k > 0,  $1 \le q \le p \le \infty$ ,  $\frac{n}{2}(\frac{1}{q}-\frac{1}{p}) < 1$ . Then there exists C > 0, such that

$$\|v(\cdot,t)\|_{L^{p}(\Omega)} \leq C(1 + \sup_{s \in (0,t)} \|u(\cdot,s)\|_{L^{q}(\Omega)}), \quad t \in (0,T)$$
(2.4)

**Proof.** Noticing  $v(\cdot, t) = e^{t(k\Delta-1)}v_0 + \int_0^t e^{(t-s)(k\Delta-1)}u(\cdot, s)ds$  for t > 0, by the standard smoothing estimates for the heat semigroup under homogeneous Neumann boundary conditions [20], we can obtain for  $q \leq p$  that

$$\begin{aligned} \|v(\cdot,t)\|_{L^{p}(\Omega)} &\leq \|\mathbf{e}^{t(k\Delta-1)}v_{0}\|_{L^{p}(\Omega)} + \int_{0}^{t} \|\mathbf{e}^{(t-s)(k\Delta-1)}u(\cdot,s)\|_{L^{p}(\Omega)}ds \\ &\leq C_{1}\|v_{0}(x)\|_{L^{\infty}(\Omega)} + C_{2}\sup_{s\in(0,t)}\|u(\cdot,s)\|_{L^{q}(\Omega)}\int_{0}^{t} (1+[k(t-s)])^{-\frac{n}{2}(\frac{1}{q}-\frac{1}{p})}\mathbf{e}^{-(\lambda_{1}+\frac{1}{k})[k(t-s)]}ds \\ &\leq C_{1}\|v_{0}(x)\|_{L^{\infty}(\Omega)} + \frac{C_{2}}{k}\sup_{s\in(0,t)}\|u(\cdot,s)\|_{L^{q}(\Omega)}\int_{0}^{\infty} (1+\alpha)^{-\frac{n}{2}(\frac{1}{q}-\frac{1}{p})}\mathbf{e}^{-\lambda_{1}\alpha}d\alpha, \quad t\in(0,T), \end{aligned}$$

where  $C_1, C_2 > 0$ ,  $\lambda_1$  is the first nonzero eigenvalue of  $-\Delta$  under the Neumann boundary condition. This proves (2.3).

Instead of the choice of  $r = \frac{p-1}{2}$  in [2], we deal with the more complicated form  $r = (p-1)\frac{[p\chi(1-k)+2k]}{p(1-k)^2+4k}$  to describe the effect of the chemicals diffusion rate k, and denote  $h(p) \triangleq h(p;\chi,k) = \frac{p\chi(1-k)+2k}{p(1-k)^2+4k}$ . For such h(p), we have the following lemma.

**Lemma 2.4** Let k > 0,  $\chi \in \left(0, -\frac{k-1}{2} + \frac{1}{2}\sqrt{(k-1)^2 + \frac{8k}{n}}\right)$ , and  $p \in \left(1, \frac{k}{[\chi^2 + \chi(k-1)]_+}\right)$ . Then  $h(p) \in (0, 1)$ .

**Proof.** We have  $h'(p) = \frac{2k(1-k)[2\chi-(1-k)]}{[p(1-k)^2+4k]^2}$ .

If 
$$k = 1$$
, then  $h(p) \equiv \frac{1}{2}$ .  
If  $k > 1$ , then  $h'(p) < 0$ , and so  
 $0 < \frac{\chi}{2\chi + k - 1} = h(\frac{k}{\chi^2 - \chi(1 - k)} - 0) = h(\frac{k}{[\chi^2 - \chi(1 - k)]_+} - 0) < h(p) < h(1 + 0) = \frac{2k + \chi(1 - k)}{(1 - k)^2 + 4k} < \frac{2k}{(1 + k)^2} < 1.$ 

Now suppose 0 < k < 1.

 $\begin{aligned} &\text{If } 0 < \chi < \frac{1-k}{2}, \text{ then } h'(p) < 0, \text{ and so} \\ &0 < \frac{\chi}{1-k} = \lim_{p \to \infty} \frac{p\chi(1-k)+2k}{p(1-k)^2+4k} < h(p) < h(1+0) = \frac{2k+\chi(1-k)}{(1-k)^2+4k} < \frac{1}{2}; \\ &\text{If } \chi = \frac{1-k}{2}, \text{ then } h(p) \equiv \frac{1}{2}; \\ &\text{If } \frac{1-k}{2} < \chi \leq 1-k, \text{ then } h'(p) > 0, \text{ and so} \\ &\frac{1}{2} < \frac{2k+\chi(1-k)}{(1+k)^2} = h(1+0) < h(p) < \lim_{p \to \infty} \frac{p\chi(1-k)+2k}{p(1-k)^2+4k} = \frac{\chi}{1-k} \leq 1; \\ &\text{If } 1-k < \chi < \frac{-(k-1)+\sqrt{(k-1)^2+\frac{8k}{n}}}{2}, \text{ then } h'(p) > 0, \text{ and so} \\ &0 < \frac{1-k^2}{(1+k)^2} < \frac{2k+\chi(1-k)}{(1+k)^2} = h(1+0) < h(p) < h(p) < h(\frac{k}{\chi^2-\chi(1-k)}-0) = \frac{\chi}{2\chi+k-1} < 1. \\ &\text{The proof is complete.} \qquad \Box \end{aligned}$ 

Denote  $c_0 = \inf_{p \in (1, \frac{n}{2}]} h(p)$  and  $c^0 = \sup_{p \in (1, \frac{n}{2}]} h(p)$  with  $n \ge 3$ . By Lemma 2.4 and its proof,  $c_0, c^0 \in (0, 1)$ . The following lemma with  $c_0$  and  $c^0$  will play an important role for estimating the bound of u by the involved iteration in the next section.

**Lemma 2.5** Let k > 0,  $\chi \in \left(0, -\frac{k-1}{2} + \frac{1}{2}\sqrt{(k-1)^2 + \frac{8k}{n}}\right)$ , then  $f(x) \triangleq \frac{n[(1-c^0)x + c^0 - c_0] + 2c_0x}{(n-2x)(1-c_0)} - x > 0$  for  $x \in (1, \frac{n}{2}]$ , provided  $n \le 8$ .

**Proof.** It suffices to show  $g(x) \triangleq 2(1-c_0)x^2 + [2c_0 - n(c^0 - c_0]x + n(c^0 - c_0) > 0$  in  $(1, \frac{n}{2}]$ . The case of  $c_0 = c^0 \in (0, 1)$  is trivial.

Now suppose  $0 < c_0 < c^0 < 1$ . We have  $\Delta_g = [2c_0 - n(c^0 - c_0)]^2 - 8n(1 - c_0)(c^0 - c_0) = [n(c^0 - c_0)]^2 + (4c_0 - 8)n(c^0 - c_0) + 4c_0^2 < 0$ , and hence g(x) > 0, whenever  $4 - 2c_0 - 4\sqrt{1 - c_0} < n(c^0 - c_0) < 4 - 2c_0 + 4\sqrt{1 - c_0}$ .

If  $n(c^0 - c_0) \le 4 - 2c_0 - 4\sqrt{1 - c_0}$ , then  $\Delta_g > 0$ . Due  $n(c^0 - c_0) - 2c_0 < 4(1 - c_0) - 4\sqrt{1 - c_0} < 0$ , we know that both the two roots of g(x) must be negative. With g(1 + 0) = 2, we obtain g(x) > 0.

If  $n(c^0 - c_0) \ge 4 - 2c_0 + 4\sqrt{1 - c_0}$ , then  $\Delta_g > 0$ , and the two roots of g(x) satisfy  $x_2 \ge x_1 > 0$ . Together with g(1+0) = 2 and  $g(\frac{n}{2}) = \frac{n^2}{2}(1-c^0) + nc_0 > 0$ , the positivity of g(x) for  $x \in (1, \frac{n}{2}]$  requires that the minimal point of g(x) satisfies  $\frac{n(c^0 - c_0) - 2c_0}{4(1 - c_0)} < 1$ , i.e.,  $n(c^0 - c_0) + 2c_0 < 4$ , by Vieta's formulas. This contradicts the case  $n(c^0 - c_0) \ge 4 - 2c_0 + 4\sqrt{1 - c_0}$ . So, the case itself should be excluded to ensure g(x) > 0 in  $(1, \frac{n}{2}]$ . Rewrite the case as  $c^0 \ge \frac{(n-2)c_0+4+4\sqrt{1-c_0}}{n} \triangleq \alpha(c_0)$  with  $c_0 \in (0, 1)$ . We have  $c^0 \ge \alpha(c_0) > \min\{\alpha(0), \alpha(1)\} = \min\{\frac{n+2}{n}, \frac{8}{n}\}$ . Thus, we would get a contradiction that  $c^0 > 1$ , whenever  $n \le 8$ .

#### **3** Proof of main result

We deal with the proof of the main result of the paper in this section.

#### Proof of Theorem 1.

We at first show that the local solutions ensured by Lemma 2.1 should be global.

Assume p > q. By the Hölder inequality, we have

$$\int_{\Omega} u^{q} dx = \int_{\Omega} (u^{p} v^{-r})^{\frac{q}{p}} v^{\frac{rq}{p}} dx \le \left(\int_{\Omega} u^{p} v^{-r} dx\right)^{\frac{q}{p}} \left(\int_{\Omega} v^{\frac{rq}{p-q}} dx\right)^{\frac{p-q}{p}}, \quad 0 < t < T.$$
(3.1)

Let  $p < \frac{k}{[\chi^2 + \chi(k-1)]_+}$ . We know from (2.3) in the proof of Lemma 2.2 with  $r \in (r_-(p), r_+(p))$  that

$$\frac{d}{dt}\int_{\Omega}u^{p}v^{-r}dx \le r\int_{\Omega}u^{p}v^{-r}dx, \ t \in (0,T),$$

and hence

$$\int_{\Omega} u^p v^{-r} dx \le C, \quad t \in (0, T)$$
(3.2)

with C = C(t) > 0. Notice  $\chi \in \left(0, -\frac{k-1}{2} + \frac{1}{2}\sqrt{(k-1)^2 + \frac{8k}{n}}\right)$  with  $p < \frac{k}{[\chi^2 + \chi(k-1)]_+}$  admits  $p > \frac{n}{2}$ .

Take  $q \in (\frac{n}{2}, p) \subset (1, \min\{p, \frac{n(p-r)}{[n-2r]_+}\})$ . Then  $\frac{n}{2}(\frac{1}{q} - \frac{p-q}{rq}) < 1$ . By Lemma 2.3, we have with  $C_1 > 0$  that

$$\|v\|_{\frac{rq}{p-q}} \le C_1(1 + \sup_{s \in (0,t)} \|u\|_q), \quad 0 < t < T.$$
(3.3)

Combining (3.1)–(3.3) yields

$$\int_{\Omega} u^q dx \le C_2 \left( 1 + \left( \sup_{s \in (0,t)} \int_{\Omega} u^q dx \right)^{\frac{r}{p}} \right), \quad t \in (0,T),$$

and hence  $\int_{\Omega} u^q dx \leq C_3$  for  $t \in (0,T)$ , where  $C_2 = C_2(t) = \tilde{C}_2 e^{rt}$  with  $\tilde{C}_2 > 0$ ,  $C_3 = C_3(t) > 0$ , and  $\frac{r}{p} < \frac{p-1}{p} < 1$  by Lemma 2.4. Now we can follow the proof of Lemma 3.4 in [19] to obtain the global existence of solutions for (1.1).

Next, we prove the solutions established above are also globally bounded if  $n \leq 8$ . We should verify that  $\int_{\Omega} u^q dx < C$  with some  $q > \frac{n}{2}$  and C > 0 [19].

The case of n = 2 is simple. Take

$$\begin{cases} p \in \left(1, \frac{k}{[\chi^2 + \chi(k-1)]_+}\right), \\ r = (p-1)h(p). \end{cases}$$

Then  $1 - \frac{1}{p-r} < 1$ . By Lemma 2.3 with the L<sup>1</sup>-conservation of u, we have

$$||v||_{L^{p-r}} \le C_4, \ t \in (0,T)$$

with time-independent  $C_4 > 0$ . By Lemma 2.2,

$$\int_{\Omega} u^p v^{-r} dx \le C_5, \quad t \in (0,T)$$

$$(3.4)$$

with some  $C_5 > 0$ . We deduce from (3.1), (3.3) and (3.4) that there exists q > 1, such that  $||u||_q \leq C_6$  for  $t \in (0,T)$  with  $C_6 = C_6(k) > 0$ .

Now consider the case of  $3 \le n \le 8$ . We should fix the assumption (2.1) with some  $p > \frac{n}{2}$  and the constant independent of t there. We will do it via an iteration procedure based of Lemma 2.5.

 $1^\circ$  Take

$$\begin{cases} p_0 \in \left(1, \min\left\{\frac{k}{[\chi^2 + \chi(k-1)]_+}, \frac{n(1-c_0) + 2c_0}{(n-2)(1-c_0)}\right\}\right),\\ r_0 = (p_0 - 1)h(p_0). \end{cases}$$
(3.5)

Then  $p_0 - r_0 \le p_0 - (p_0 - 1)c_0 = (1 - c_0)p_0 + c_0 < \frac{n}{n-2}$ , i.e.,  $\frac{n}{2}(1 - \frac{1}{p_0 - r_0}) < 1$ . By Lemma 2.3 with the  $L^1$ -conservation of u, we have

$$\|v\|_{L^{p_0-r_0}} \le C_7, \ t \in (0,T),$$

and hence

$$\int_{\Omega} u^{p_0} v^{-r_0} dx \le C_8, \quad t \in (0,T)$$
(3.6)

by Lemma 2.2, with some  $C_7, C_8 > 0$ . We know from (3.1), (3.3) and (3.6) that  $||u||_{q_0} \leq C_9$  for  $t \in (0,T)$  with  $C_9 > 0$ . If  $p_0 > \frac{n}{2}$ , take  $q_0 \in (\frac{n}{2}, p_0) \subset (1, \min\{p_0, \frac{n(p_0 - r_0)}{[n - 2r_0]_+}\})$ .

 $2^{\circ}$  Assume  $p_0 \leq \frac{n}{2}$ . Take

$$\begin{cases} p_1 \in \left(p_0, \min\left\{\frac{k}{[\chi^2 + \chi(k-1)]_+}, \frac{n[(1-c^0)p_0 + c^0 - c_0] + 2c_0p_0}{(n-2p_0)(1-c_0)}\right\}\right),\\ r_1 = (p_1 - 1)h(p_1), \end{cases}$$

where the well-definedness of the interval for  $p_1$  is ensured by Lemma 2.5. A simple calculation shows

$$p_1 - r_1 \le (1 - c_0)p_1 + c_0 < \frac{n[(1 - c^0)p_0 + c^0]}{n - 2p_0} \le \frac{n(p_0 - r_0)}{n - 2p_0} = \frac{n\frac{n(p_0 - r_0)}{n - 2r_0}}{n - \frac{2n(p_0 - r_0)}{n - 2r_0}}$$

i.e.,  $\frac{n}{2}(\frac{1}{\frac{n(p_0-r_0)}{n-2r_0}}-\frac{1}{p_1-r_1}) < 1$ . By Lemma 2.3 with  $p_0 \le \frac{n}{2}$ , there is  $q_0 \in (1, \frac{n(p_0-r_0)}{n-2r_0})$  such that

$$||v||_{p_1-r_1} < C_{10}, \quad t \in (0,T),$$

and thus

$$\int_{\Omega} u^{p_1} v^{-r_1} dx \le C_{11}, \quad t \in (0,T)$$
(3.7)

by Lemma 2.2, with some  $C_{10}, C_{11} > 0$ . It follows from (3.1), (3.3) and (3.7) that  $||u||_{q_1} \le C_{12}$  for  $t \in (0, T)$  with  $C_{12} > 0$ . If  $p_1 > \frac{n}{2}$ , take  $q_1 \in (\frac{n}{2}, p_1) \subset (1, \min\{p_1, \frac{n(p_1 - r_1)}{[n - 2r_1]_+}\})$ .

3° Assume  $p_{l-1} \leq \frac{n}{2}, l = 1, 2, 3, \dots$  Take

$$\begin{cases} p_l \in \left( p_{l-1}, \min\left\{ \frac{k}{[\chi^2 + \chi(k-1)]_+}, \frac{n[(1-c^0)p_{l-1} + c^0 - c_0] + 2c_0 p_{l-1}}{(n-2p_{l-1})(1-c_0)} \right\} \right), \\ r_l = (p_{l-1} - 1)h(p_{l-1}), \end{cases}$$
(3.8)

where the interval for  $p_l$  is well defined due to Lemma 2.5. Repeat the procedure in 2°, we deduce with  $q_l \in (1, \min\{p_l, \frac{n(p_l-r_l)}{[n-2r_l]+}\})$  that  $||u||_{q_l} \leq C_{13}$ , 0 < t < T, with some  $C_{13} > 0$ . Noticing  $\frac{n[(1-c^0)p_{l-1}+c^0-c_0]+2c_0p_{l-1}}{(n-2p_{l-1})(1-c_0)} \to \infty$  as  $l \to \infty$  by Lemma 2.5, we can realize  $p_l > \frac{n}{2}$  after finite steps. Let  $q = q_l \in (\frac{n}{2}, p_l)$  to get  $\int_{\Omega} u^q dx \leq C_{14}$  for all  $t \in (0,T)$  with  $C_{14} > 0$ . It is mentioned that for any fixed k > 0, the constants  $C_i$ ,  $i = 5, \ldots, 14$ , are all independent of  $t \in (0,T)$  here.

Based on the above estimate  $||u(\cdot,t)||_{L^q(\Omega)}$  with  $q > \frac{n}{2}$ , uniform for  $t \in (0,T)$ , we conclude the global boundedness of u by coping the related arguments in [19] for that to the case of k = 1.  $\Box$  **Remark 3** Notice  $p_0 < \frac{k}{[\chi^2 + \chi(k-1)]_+}$  in (3.5) with  $\chi \in \left(0, -\frac{k-1}{2} + \frac{1}{2}\sqrt{(k-1)^2 + \frac{8k}{n}}\right)$  admits  $p_0 > \frac{n}{2}$ . Moreover, a simple computation shows  $p_0 < \frac{n(1-c_0)+2c_0}{(n-2)(1-c_0)}$  in (3.5) ensures  $\frac{n(1-c_0)+2c_0}{(n-2)(1-c_0)} > \frac{n}{2}$  whenever n = 3, 4. Therefore, the Steps 2° and 3° in the proof of Theorem 1 are unnecessary for n = 3, 4 there. In addition, it should be pointed out that  $c^0 = c_0 = \frac{1}{2}$  if k = 1, and thus the requirement  $n \leq 8$  itself can be removed away for the global boundedness of solutions with k = 1.

# References

- P. Biler, Global solutions to some parabolic-elliptic systems of chemotaxis, Adv. Math. Sci. Appl. 9 (1999) 347–359.
- [2] K. Fujie, Boundedness in a fully parabolic chemotaxis system with singular sensitivity, J. Math. Anal. Appl. 424 (2015) 675–684.
- [3] K. Fujie, M. Winkler, T. Yokota, Boundedness of solutions to parabolic-elliptic Keller-Segel systems with signal-dependent sensitivity, Math. Methods. Appl. Sci. 38 (2015) 1212–1224.
- [4] K. Fujie, T. Yokota, Boundedness in a fully parabolic chemotaxis system with strongly singular sensitivity, Appl. Math. Letters 38 (2014) 140–143.
- [5] D. Henry, Germetric Theory of Semilinear Parabolic Equations, Springer, Berlin, Heidelberg, 1981.
- [6] T. Hillen, K.J. Painter, A use'sguidence to PDE models for chemotaxis, J. Math. Biol. 58 (2009) 183–217.
- [7] D. Horstann, M. Winkler, Boundedness vs. blow-up in a chemotaxis system, J. Differential Equations 215 (2005) 52–107.
- [8] W. Jäger, S. Luchhaus, On explosions of solutions to a system of partial differential equations modelling chemotaxis, Trans. Amer. Math. Soc. 329 (1992) 817–824.
- [9] E.F. Keller, L.A. Segel, Initiation of slime mold aggregation viewed as an instability, J. Theoret. Biol. 26 (1970) 399–415.
- [10] J. Lankeit A new approach toward boundedness in a two-dimensional parabolic chemotaxis system with singular sensitivity, Math. Methods Appl. Sci. 2015.
- [11] T. Nagai, T. Senba, Behavior of radially symmetric solutions of a system related to chemotaxis, Nonlinear Anal. 30 (1997) 3837–3842.
- [12] T. Nagai, T. Senba, K. Yoshida, Global existence of solutions to the parabolic systems of chemotaxis, RIMS Kokyuroku, 1009 (1997) 22–28.
- [13] K. Osaki, A. Yagi, Finite dimensional attractors for one-dimensional Keller-Segel equations, Funkcial. Ekvac. 44 (2001) 441–469.
- [14] B. D. Sleeman, H. A. Levine, Partial differential equations of chemotaixs and angiogenesis, Math. Methods. Appl. Sci. 24 (2001) 405–426.
- [15] C. Stinner, M. Winkler, Global weak solutions in a chemotaxis system with large singular sensitivity, Nonliear Anal. Real World Appl. 12 (2011) 3727–3740.
- [16] Q. Wang, Global solutions of a Keller-Segel system with saturated logarithmic sensitivity function, Commun. Pure Appl. Anal. 14 (2015) 383–396.
- [17] D. Horstmann, M. Winkler, Boundedness vs. blow-up in a chemotaxis system, J. Differential Equations 215 (2005) 52–107.

- [18] M. Winkler, Absence of collapse in a parabolic chemotaxis system with signal-dependent sensitivity, Math. Nachr. 283 (2010) 1664–1673.
- [19] M. Winkler, Global solutions in a fully parabolic chemotaxis system with singular sensitivity, Math. Methods. Appl. Sci. 34 (2011) 176–190.
- [20] M. Winkler, Aggregation vs. global diffusive behavior in the higher-dimensional Keller-Segel model, J. Differential Equations 248 (2011) 3728–3740.
- [21] P. Zheng, C. Mu, X. Hu, Q. Zheng, Global boundedness in q quasilinear chemotaxis system with signal-dependent sensitivity, J. Math. Anal. Appl. 428 (2015) 508–524.