MEASURE-PRESERVING SEMIFLOWS AND ONE-PARAMETER KOOPMAN SEMIGROUPS

NIKOLAI EDEKO, MORITZ GERLACH AND VIKTORIA KÜHNER

Dedicated to Rainer Nagel on the occasion of his 75th birthday.

ABSTRACT. For a finite measure space X, we characterize strongly continuous Markov lattice semigroups on $L^p(X)$ by showing that their generator A acts as a derivation on the dense subspace $D(A) \cap L^{\infty}(X)$. We then use this to characterize Koopman semigroups on $L^p(X)$ if X is a standard probability space. In addition, we show that every measurable and measure-preserving flow on a standard probability space is isomorphic to a continuous flow on a compact Borel probability space.

In this article we address mainly the following two issues. First, we characterize strongly continuous Markov lattice semigroups $(T(t))_{t\geq 0}$ on $L^p(X)$ by properties of their generators for a finite measure space $X = (X, \Sigma, \mu)$. We will show that a strongly continuous semigroup on $L^p(X)$ is a Markov lattice semigroup if and only if its generator A acts as a derivation on $D(A) \cap L^{\infty}(X)$, $\mathbb{1} \in D(A)$ and the semigroup is locally bounded on $L^{\infty}(X)$. Similar results have been established by R. Nagel and R. Derndinger in [2, Satz 2.5] for semigroups on C(K), see also [8, Section B-II.3], and recently by T. ter Elst and M. Lemańczyk in [4] for unitary groups on $L^2(X)$.

Second, we show that such semigroups are always similar to a semigroup of Koopman operators. More precisely, we construct a compact space K and a Borel measure ν such that $L^1(X, \Sigma, \mu)$ is isometrically Banach lattice isomorphic to $L^1(K, \nu)$ and, via this isomorphism, the semigroup $(T(t))_{t\geq 0}$ is similar to a semigroup of Koopman operators on $L^1(K, \nu)$ induced by a continuous semiflow $(\varphi_t)_{t\geq 0}$ on K. Furthermore, in case that the space $L^1(X, \Sigma, \mu)$ is separable, we show that K can be chosen to be metrizable. Similar results have been already obtained for strongly continuous representations of locally compact groups on $L^p(X)$ as bi-Markov embeddings, see [7, Theorem 5.14].

The article is organized as follows. In the second part of the introduction, we specify our notation and recall some basic facts we use throughout the article. In Section 1, we prove our main result, Theorem 1.1, that characterizes strongly Markov semigroups of lattice homomorphisms by the condition that their generator acts as a derivation, followed by a version for semigroups that are not necessarily Markov. In Section 2 we then use these results to characterize Koopman semigroups and in particular obtain [4, Theorem 1.1] in which ter Elst and Lemańczyk proved a corresponding result for unitary operator groups as Corollary 2.6. In Section 3, we turn to the construction of topological models. Finally, in Section 4, we consider ergodic, measure-preserving flows and give a new proof for the fact that they contain at most countably many non-ergodic mappings, provided that their induced group of Koopman operators is strongly continuous on $L^2(X)$. This has previously been proven in [9, Theorem 1] for \mathbb{R}^k -actions.

Let us recall some concepts and fix the notation used in this article. For a measure space $X = (X, \Sigma, \mu)$ and $1 \le p \le \infty$ we denote by

$$\mathrm{L}^p(\mathrm{X}) \coloneqq \mathrm{L}^p(\mathrm{X};\mathbb{C}) = \mathrm{L}^p(\mathrm{X};\mathbb{R}) \otimes \mathrm{i}\mathrm{L}^p(\mathrm{X};\mathbb{R})$$

the corresponding complex L^p -space. This is a complex Banach lattice in the sense of [3, Definition 7.2]. The lattice operations in $L^p(X; \mathbb{R})$ are denoted by \vee and \wedge and we write f_+ and f_- for the positive part $f \vee 0$ and negative part $(-f) \vee 0$ of a function $f \in L^p(X; \mathbb{R})$. We call X *separable* if $L^p(X)$ is separable for one (and hence all) $p \in [1,\infty)$. When considering more than one measure on X, we may distinguish between a measurable function $f: X \to \mathbb{C}$ and its equivalence class with respect to a measure μ on X by writing $[f]_{\mu}$ for the equivalence class of f. If $f \in L^{\infty}(X)$ is an essentially bounded function, M_f will denote its associated multiplication operator on $L^p(X)$, $p \in [1,\infty]$. Let $X = (X, \Sigma, \mu)$ and $Y = (Y, \Sigma', \mu')$ be finite measure spaces and $T: L^p(X) \to L^p(Y)$ a linear operator. The operator T is called *positive* if it is real, i.e., if $TL^p(X; \mathbb{R}) \subseteq L^p(Y; \mathbb{R})$, and its restriction to the Banach lattice $L^p(X; \mathbb{R})$ is positive. The operator T is said to be a *lattice homomorphism* if |Tf| = T|f| for each $f \in L^p(X)$. In particular, every lattice homomorphism is positive and fulfills $T(f_+) = (Tf)_+$ and $T(f_-) = (Tf)_-$ for each $f \in L^p(X; \mathbb{R})$. The operator T is called a *Markov operator* if it is positive and $T\mathbb{1}_X = \mathbb{1}_Y$, and a *bi-Markov* operator if, additionally, $T'\mathbb{1}_Y = \mathbb{1}_X$. It is called a *Koopman operator* if there is a measurable map $\varphi: Y \to X$ such that φ^{-1} maps null-sets into null-sets and $Tf = f \circ \varphi$ for all $f \in L^p(X)$. In this case, we denote the operator by T_{φ} . Note that every Koopman operator is a Markov lattice homomorphism.

Let $(T(t))_{t\geq 0}$ be a C_0 -semigroup on $L^p(X)$ for some $1 \leq p < \infty$. We denote its generator by A and the domain of A by D(A). The semigroup $(T(t))_{t\geq 0}$ is called a *lattice semigroup* if each operator T(t) is a lattice homomorphism. If, additionally, for each $t \geq 0$ the operator T(t) is a (bi-)Markov operator, $(T(t))_{t\geq 0}$ is called a *(bi-)Markov lattice semigroup*. It is called a *Koopman semigroup*, if for each $t \geq 0$ the operator T(t) is a Koopman operator.

Consider the equivalence relation

$M \sim N$ if $\mathbb{1}_M = \mathbb{1}_N \mu$ -almost everywhere

on Σ . Then the set of equivalence classes $\Sigma(X) := \Sigma/\sim$ is called the *measure algebra* of the measure space X and is a Boolean algebra with respect to the set operations union, intersection and complementation. For the sake of simplicity, we do not distinguish notationally between elements of Σ and $\Sigma(X)$. A mapping $\theta \colon \Sigma(X) \to \Sigma(X)$ is called a *Boolean algebra homomorphism* if $\theta(\emptyset) = \emptyset$, $\theta(X) = X$ as well as $\theta(A \cup B) = \theta(A) \cup \theta(B)$ and $\theta(A \cap B) = \theta(A) \cap \theta(B)$ for all $A, B \in \Sigma(X)$. If, in addition, a Boolean algebra homomorphism $\theta \colon \Sigma(X) \to \Sigma(X)$ satisfies $\mu(\theta(A)) = \mu(A)$ for all $A \in \Sigma(X)$, then θ is called a *measure algebra homomorphism*. Every measurable map $\varphi \colon X \to X$ such that φ^{-1} maps null-sets into null-sets induces a measure algebra homomorphism $\varphi^* \colon \Sigma(X) \to \Sigma(X)$ via $A \mapsto \varphi^{-1}(A)$. For further information on $\Sigma(X)$ we refer to [3, Section 6.1]. We call a measure space (X, Σ, μ) a *Borel probability space* if X can be equipped with a Polish topology such that Σ is the corresponding Borel σ -algebra. A measure space (X, Σ, μ) is called a *standard probability space* if there is a measurable, measure-preserving and essentially invertible map to a Borel probability space.

Finally, we call a linear operator δ on $L^p(X)$ with domain $D(\delta)$ a derivation on $D(\delta) \cap L^{\infty}(X)$ if $D(\delta) \cap L^{\infty}(X)$ is an algebra (with respect to the pointwise multiplication) and $\delta(f \cdot g) = \delta f \cdot g + f \cdot \delta g$ for all $f, g \in D(\delta) \cap L^{\infty}(X)$.

MEASURE-PRESERVING SEMIFLOWS AND ONE-PARAMETER KOOPMAN SEMIGROUPS 3

1. CHARACTERIZATION OF MARKOV LATTICE SEMIGROUPS ON L^p-spaces

In this section we characterize strongly continuous Markov lattice semigroups on $L^{p}(X)$ -spaces, where $X = (X, \Sigma, \mu)$ is a finite measure space, by means of their generators. The following theorem is our main result.

Theorem 1.1. Let A be the generator of a C_0 -semigroup $(T(t))_{t\geq 0}$ on a space $L^p(X)$, where X is a finite measure space and $1 \leq p < \infty$. Then the following assertions are equivalent.

- (i) $(T(t))_{t\geq 0}$ is a Markov lattice semigroup.
- (ii) For every $t \ge 0$ there exists a Boolean algebra homomorphism $\theta_t \colon \Sigma(X) \to \Sigma(X)$ such that $T(t)\mathbb{1}_M = \mathbb{1}_{\theta_t(M)}$ for all $M \in \Sigma(X)$.
- (iii) The space $L^{\infty}(X)$ is invariant under $(T(t))_{t\geq 0}$, the map $t \mapsto ||T(t)||_{\mathscr{L}(L^{\infty}(X))}$ is locally bounded, $\mathbb{1} \in D(A)$ and A is a derivation on $D(A) \cap L^{\infty}(X)$.
- **Remark 1.2.** (i) Given a finite measure space X and a bounded operator S on $L^p(X)$ such that $L^{\infty}(X)$ is invariant under S, it follows from the closed graph theorem that the restriction of S to $L^{\infty}(X)$ is a bounded operator. Therefore, the map $t \mapsto ||T(t)||_{\mathscr{L}(L^{\infty}(X))}$ is well-defined in (iii). As will be shown in Lemma 2.5, the local boundedness condition is automatically fulfilled if T is an operator group.
 - (ii) A semigroup $(T(t))_{t\geq 0}$ satisfying (i)-(iii) in Theorem 1.1 uniquely extends to a strongly continuous Markov lattice semigroup on $L^q(X)$ for each $1 \leq q < \infty$ with

$$||T(t)||_{\mathscr{L}(\mathrm{L}^{q}(\mathrm{X}))} = ||T(t)||_{\mathscr{L}(\mathrm{L}^{1}(\mathrm{X}))}^{\frac{1}{q}},$$

use [3, Theorem 7.23]. In particular, it extends to the biggest L^q -space, $L^1(X)$, and we will therefore only consider semigroups on $L^1(X)$ in Sections 3 and 4. At this point, however, the dependence on p in Theorem 1.1 cannot be eliminated as easily since it is only clear a posteriori that the implication (iii) \implies (i) can be reduced to the case p = 1.

As a preparation for the proof of Theorem 1.1, recall the following lemma relating the algebra and the lattice structure of $L^{\infty}(X)$.

Lemma 1.3. Let X be a finite measure space and $T: L^{\infty}(X) \to L^{\infty}(X)$ be a bounded linear operator satisfying $T\mathbb{1} = \mathbb{1}$. Then the following assertions are equivalent.

- (i) T is multiplicative.
- (ii) T is a C^{*}-homomorphism.
- (iii) T is a lattice homomorphism.

Proof. Obviously, (ii) implies (i). The equivalence of (ii) and (iii) can be found in [3, Theorem 7.23]. (There, the operator is assumed to be conjugation-preserving but this assumption is trivially superfluous.) The implication (i) \implies (ii) follows from [3, Theorem 4.13], the analogous statement for spaces of continuous functions, by applying the Gelfand-Naimark theorem [3, Theorem 4.23].

The following continuity property will be essential for the proof of Theorem 1.1.

Lemma 1.4. Let X be a measure space and $B \subseteq L^{\infty}(X)$ be bounded. Then the multiplication $L^{p}(X) \times B \to L^{p}(X), (f,g) \mapsto fg$ is $\|\cdot\|_{p}$ -continuous.

Proof. Let M be a bound for B. For $f, u \in L^p(X), g, v \in B$ and c > 0

$$fg - uv = (f - u)g + u(g - v)$$

= $(f - u)g + u\mathbb{1}_{[|u| \le c]}(g - v) + u\mathbb{1}_{[|u| > c]}(g - v)$

and so

$$\limsup_{(f,g)\to(u,v)} \|fg - uv\|_p \le 2M \|u\mathbb{1}_{[|u|>c]}\|_p \to 0 \quad (c \to \infty).$$

Next, we apply Lemma 1.4 to a C_0 -semigroup as in Theorem 1.1.

Corollary 1.5. Let A be the generator of a C_0 -semigroup $(T(t))_{t>0}$ on a space $L^{p}(X)$, where X is a finite measure space and $1 \leq p < \infty$. In addition, suppose that the space $L^{\infty}(X)$ is invariant under $(T(t))_{t\geq 0}$ and that the map $t \mapsto ||T(t)||_{\mathscr{L}(L^{\infty}(X))}$ is locally bounded. Then, for all $f \in L^{\infty}(\overline{X})$ and all $g \in L^{p}(X)$ the function

$$[0,\infty) \to t \mapsto L^p(X), \quad t \mapsto T(t)f \cdot T(t)g$$

is continuous. Moreover, for $f, g \in D(A) \cap L^{\infty}(X)$ this function is differentiable and the product rule

$$\frac{\mathrm{d}}{\mathrm{d}t}\left(T(t)f\cdot T(t)g\right) = T(t)Af\cdot T(t)g + T(t)f\cdot T(t)Ag$$

holds.

Proof. It suffices to prove the second part since the first is a consequence of Lemma 1.4. Let $f, g \in D(A) \cap L^{\infty}(X)$ and $t \geq 0$. Use Lemma 1.4 and differentiate to obtain

$$\frac{\mathrm{d}}{\mathrm{d}t} \left(T(t)f \cdot T(t)g \right) = \lim_{h \to 0} \frac{1}{h} \left(\left[T(t+h)f - T(t)f \right] T(t)g + T(t+h)f \left[T(t+h)g - T(t)g \right] \right)$$

$$= \lim_{h \to 0} \frac{1}{h} \left(\left[T(t+h)f - T(t)f \right] \right) T(t)g + \lim_{h \to 0} T(t+h)f \cdot \lim_{h \to 0} \frac{1}{h} \left[T(t+h)g - T(t)g \right]$$

$$= \left(\frac{\mathrm{d}}{\mathrm{d}t} T(t)f \right) \cdot T(t)g + T(t)f \cdot \left(\frac{\mathrm{d}}{\mathrm{d}t} T(t)g \right)$$
roves the assertion.

which proves the assertion.

We are now able to prove Theorem 1.1.

Proof of Theorem 1.1. The equivalence (i) \Leftrightarrow (ii) is proved almost exactly as in the case of bi-Markov operators, see [3, Theorem 12.10].

To prove the implication (i) \implies (iii), first note that every operator T(t) is positive. Since $T(t)\mathbb{1} = \mathbb{1}$ for each t > 0, this already implies that the semigroup preserves the subspace $L^{\infty}(X)$ and the restriction of each T(t) to $L^{\infty}(X)$ is a contraction. In particular, it follows from Lemma 1.3 that every operator T(t) is multiplicative on $L^{\infty}(X)$. By Corollary 1.5, for every $f, g \in D(A) \cap L^{\infty}(X)$

$$\frac{\mathrm{d}}{\mathrm{d}t}T(t)(f \cdot g) = \frac{\mathrm{d}}{\mathrm{d}t} (T(t)f \cdot T(t)g)$$
$$= T(t)Af \cdot T(t)g + T(t)f \cdot T(t)Ag$$
$$= T(t) [Af \cdot g + f \cdot Ag] .$$

In particular, this shows $f \cdot g \in D(A)$ and $A(f \cdot g) = Af \cdot g + f \cdot Ag$ for t = 0. This proves that A is a derivation and clearly $\mathbb{1} \in D(A)$.

We now prove that (iii) implies (i). Because of the local boundedness of $t \mapsto$ $||T(t)||_{\mathscr{L}(L^{\infty}(X))}$, there exists a constant C > 0 such that

$$\left\|\frac{1}{t}\int_0^t T(s)f\,\mathrm{d}s\right\|_{\infty} \le C\|f\|_{\infty}$$

for $0 < t \leq 1$ and $f \in L^{\infty}(X)$. This implies that $D \coloneqq D(A) \cap L^{\infty}(X)$ is a dense subspace of $L^{p}(X)$. We use this fact to show that each T(t) is multiplicative on $L^{\infty}(X)$. For fixed $f, g \in D$ and t > 0 consider the mapping

$$s \mapsto \beta(s) \coloneqq T(t-s)[T(s)f \cdot T(s)g]$$

on [0, t]. Since $\beta(0) = T(t)(f \cdot g)$ and $\beta(t) = T(t)f \cdot T(t)g$, it suffices to show that β is constant. To this end, consider the operator valued mappings $P, Q: [0, t] \rightarrow \mathscr{L}(L^p(X))$ given by P(s) = T(t-s) and $Q(s) = M_{T(s)f} \circ T(s)$, where $M_{T(s)f}$ denotes the multiplication with the bounded function T(s)f. It follows from Corollary 1.5 that Q is strongly continuous and that for each $h \in D$, $s \mapsto Q(s)h$ is differentiable with derivative

$$\frac{\mathrm{d}}{\mathrm{d}s}Q(s)h = T(s)Af \cdot T(s)h + T(s)f \cdot T(s)Ah = A(T(s)f \cdot T(s)h).$$

Here, the second equality follows from the fact that, by assumption, A is a derivation and D is invariant under each T(t). In particular, D is invariant under Q. Since P is also strongly continuous and $s \mapsto P(s)h$ is differentiable for all $h \in D$, it follows from [5, Lemma B.16] that

$$\beta'(s) = -AT(t-s)[T(s)f \cdot T(s)g] + T(t-s)A[T(s)f \cdot T(s)g] = 0$$

for all $s \in [0, t]$. This shows that β is constant and thus that every T(t) is multiplicative on D.

Since the multiplication with a fixed bounded function induces a bounded operator on $L^p(X)$ and D is $\|\cdot\|_p$ -dense in $L^{\infty}(X)$, fixing a function $g \in D$ and using a standard approximation argument shows that $T(f \cdot g) = T(t)f \cdot T(t)g$ for all $f \in L^{\infty}(X)$ and $g \in D$. Fixing $f \in L^{\infty}(X)$ and repeating the argument shows that $T(t)(f \cdot g) = T(t)f \cdot T(t)g$ for all $f, g \in L^{\infty}(X)$, so T(t) is multiplicative on all of $L^{\infty}(X)$. Furthermore, $A\mathbb{1} = 0$ since A is a derivation, i.e., $T(t)\mathbb{1} = \mathbb{1}$ for all $t \geq 0$. Now Lemma 1.3 yields that every T(t) is a lattice homomorphism on $L^{\infty}(X)$ and hence, by density and continuity, also on $L^p(X)$.

Remark 1.6. The assumption $\mathbb{1} \in D(A)$ in (iii) of Theorem 1.1 is automatically satisfied if T(t) is an isometry for each $t \ge 0$. To see this, note that in the proof of Theorem 1.1 the assumption $\mathbb{1} \in D(A)$ was only important for the implication (iii) \implies (i). There, we showed that all the operators of the semigroup are multiplicative on $L^{\infty}(X)$ and therefore map characteristic functions to characteristic functions. If T(t) is an isometry, it follows that $T(t)\mathbb{1} = \mathbb{1}$ for each $t \ge 0$ and hence $\mathbb{1} \in D(A)$. This assumption is also fulfilled if $T(t)'\mathbb{1} = \mathbb{1}$ for each $t \ge 0$, since then $\langle T(t)\mathbb{1},\mathbb{1} \rangle = \langle \mathbb{1},\mathbb{1} \rangle = \mu(X)$ and so, $T(t)\mathbb{1}$ being a characteristic function, $T(t)\mathbb{1} = \mathbb{1}$ for each $t \ge 0$.

As a corollary of Theorem 1.1, we also obtain the following characterization of bi-Markov lattice semigroups.

Corollary 1.7. Let A be the generator of a C_0 -semigroup $(T(t))_{t\geq 0}$ on a space $L^p(X)$, where X is a finite measure space and $1 \leq p < \infty$. Then the following assertions are equivalent.

- (i) $(T(t))_{t\geq 0}$ is a bi-Markov lattice semigroup.
- (ii) For every $t \ge 0$ there exists a measure algebra homomorphism $\theta_t \colon \Sigma(X) \to \Sigma(X)$ such that $T(t)\mathbb{1}_M = \mathbb{1}_{\theta_t(M)}$ for all $M \in \Sigma(X)$.
- (iii) The space $L^{\infty}(X)$ is invariant under $(T(t))_{t\geq 0}$, the map $t \mapsto ||T(t)||_{\mathscr{L}(L^{\infty}(X))}$ is locally bounded, A is a derivation on $D(A) \cap L^{\infty}(X)$ and $A'\mathbb{1} = 0$.

Proof. For the equivalence of (i) and (ii), the reader is again referred to [3, Theorem 12.10]. The equivalence of (i) and (iii) follows from Theorem 1.1 and Remark 1.6 since $T(t)'\mathbb{1} = \mathbb{1}$ for all $t \ge 0$ is equivalent to $A'\mathbb{1} = 0$.

In the following we discuss C_0 -semigroups of lattice homomorphisms on $L^p(X)$ that are not necessarily Markov. We show that their generator is a derivation perturbed by a bounded multiplication operator.

Theorem 1.8. Let A be the generator of a C_0 -semigroup $(T(t))_{t\geq 0}$ on a space $L^p(X)$, where X is a finite measure space and $1 \leq p < \infty$. Assume that $\mathbb{1} \in D(A)$ and $q := A\mathbb{1} \in L^{\infty}(X)$. Then the following assertions are equivalent.

- (i) $(S(t))_{t>0}$ is a lattice semigroup.
- (ii) The function q is real-valued and A = B + q where B is the generator of a Markov lattice C_0 -semigroup $(T(t))_{t\geq 0}$ on $L^p(X)$.

If (ii) holds, then

(1)
$$S(t)f = \exp\left(\int_0^t T(s)q\,\mathrm{d}s\right) \cdot T(t)f$$

for all $t \ge 0$ and $f \in L^p(X)$.

Proof. To show the equivalence of (i) and (ii), we first recall from [5, Theorem III.1.3] that $B \coloneqq A-q$ is a generator of a C_0 -semigroup $(T(t))_{t\geq 0}$ on $L^p(X)$ because B is a bounded perturbation of A. Since $B\mathbb{1} = 0$, $T(t)\mathbb{1} = \mathbb{1}$ for all $t \geq 0$. Now it follows from [8, Corollary C-II.5.8] (Kato's identity) that $(S(t))_{t\geq 0}$ is a lattice semigroup if and only if D(A) is a sublattice of $L^p(X)$ and

$$A|f| = \operatorname{Re}(\operatorname{sign}(f)Af)$$

for all $f \in D(A)$. Since (i) implies that q is real-valued, A satisfies this condition if and only if B does, which proves the equivalence of the assertions (i) and (ii). Now assume that (ii) holds. Since each T(t) is multiplicative on $L^{\infty}(X)$ by Lemma 1.3, $T(t) \exp(g) = T(t) \sum_{n=0}^{\infty} \frac{g^n}{n!} = \exp(T(t)g)$ for each $g \in L^{\infty}(X)$. Using this, one proves by induction that

$$\left(T(t)\mathrm{e}^{t\mathrm{M}_{q}}\right)^{n} = \mathrm{M}_{\exp\left(t\sum_{j=1}^{n}T(jt)q\right)}T(nt)$$

for each $n \geq 1$. Replace t by $\frac{t}{n}$ and note that

$$\sum_{j=1}^{n} \frac{t}{n} T\left(\frac{jt}{n}\right) q \xrightarrow[n \to \infty]{} \int_{0}^{t} T(s) q \, \mathrm{d}s.$$

By Lemma 1.4, one obtains the convergence

$$\operatorname{M}_{\exp\left(\sum_{j=1}^{n} \frac{t}{n} T\left(\frac{jt}{n}\right) q\right)} \xrightarrow[n \to \infty]{} \operatorname{M}_{\exp\left(\int_{0}^{t} T(s) q \, \mathrm{d}s\right)}$$

of multiplication operators in the strong operator topology on $\mathscr{L}(L^p(X))$. By the Trotter product formula

$$S(t)f = \lim_{n \to \infty} \left[T\left(\frac{t}{n}\right) \exp\left(\frac{t}{n} \mathbf{M}_q\right) \right]^n f = \exp\left(\int_0^t T(s)q \,\mathrm{d}s\right) \cdot T(t)f$$

$$f \in \mathbf{L}^p(\mathbf{X}).$$

for all $f \in L^p(X)$.

2. KOOPMAN SEMIGROUPS ON L^p -spaces

Every Koopman semigroup on an $L^{p}(X)$ -space is a Markov lattice semigroup but the converse is, in general, not true. However, it does hold if X is a standard probability space: In the case of bi-Markov lattice homomorphisms, this is a classical theorem by von Neumann, cf. [3, Theorem 7.20]. Below, we give an operator-theoretic proof extending this theorem to Markov lattice homomorphisms on L^{p} -spaces. We then relate this to the results on semigroups from the previous section.

Theorem 2.1. Let $X = (X, \Sigma_X, \mu_X)$ and $Y = (Y, \Sigma_Y, \mu_Y)$ be standard probability spaces and $T: L^p(X) \to L^p(Y), 1 \le p \le \infty$, a Markov lattice homomorphism (not necessarily bi-Markov). Then there is a measurable map $\varphi: Y \to X$ such that $T = T_{\varphi}$. If $\vartheta: Y \to X$ is another such map, then $\varphi = \vartheta \mid \mu_Y$ -almost everyhwere.

Proof. We will factorize T in order to apply von Neumann's theorem for bi-Markov lattice homomorphisms. Since $L^{\infty}(X)$ is dense in $L^{p}(X)$ for any $p \in [1, \infty]$, we only need to prove the case $p = \infty$. Moreover, assume without loss of generality that $X = (K, \mathcal{B}(K), \mu_{K})$ and $Y = (L, \mathcal{B}(L), \mu_{L})$ are Borel probability spaces.

Set $\nu := (T'\mathbb{1}_Y)\mu$ and note that $(K, \mathcal{B}(K), \nu)$ is again a Borel probability space. Since $\nu \ll \mu$, the map $P : L^p(K, \mathcal{B}(K), \mu_K) \to L^p(K, \mathcal{B}(K), \nu), [f]_{\mu_K} \mapsto [f]_{\nu}$ is a well-defined, surjective and bounded operator. Moreover, for $f \in L^{\infty}(K, \mathcal{B}(K), \mu_K)$,

(2)
$$\int_{K} |f| \,\mathrm{d}\nu = \int_{L} T|f| \,\mathrm{d}\mu_{L} = \int_{L} |Tf| \,\mathrm{d}\mu_{L}$$

and so ker $T \subseteq$ ker P. Therefore, there is an operator $\hat{T} : L^{\infty}(K, \mathcal{B}(K), \nu) \to L^{\infty}(L, \mathcal{B}(L), \mu_L)$ such that $T = \hat{T}P$. Since P is surjective, \hat{T} also is a Markov lattice homomorphism and it follows from (2) that \hat{T} is, in fact, bi-Markov.

Now, von Neumann's theorem shows that $\hat{T}[f]_{\nu} = [f \circ \varphi]_{\mu_L}$ for a measurable and measure-preserving map $\varphi \colon (L, \mathcal{B}(L), \mu_L) \to (K, \mathcal{B}(K), \nu)$ and so $T = T_{\varphi}$. The proof that φ is unique almost everywhere is the same as for measure-preserving maps, so we refer the reader to, e.g., [3, Lemma 6.9].

Corollary 2.2. Let A be the generator of a C_0 -semigroup $(T(t))_{t\geq 0}$ on a space $L^p(X)$, where $X = (X, \Sigma, \mu)$ is a standard probability space and $1 \leq p < \infty$. Then the equivalent assertions (i), (ii) and (iii) of Theorem 1.1 are also equivalent to

(iv) There exists a family $(\varphi_t)_{t\geq 0}$ of measurable maps on X such that $T(t)f = f \circ \varphi_t$ for all $f \in L^p(X)$ and $t \geq 0$.

Proof. If assertion (i) of Theorem 1.1 holds, we obtain assertion (iv) by Theorem 2.1 below. Conversely, if (iv) holds, then every T(t) is a Markov lattice homomorphism, thus assertion (i) holds.

Remark 2.3. Given a Koopman semigroup $(T_{\varphi_t})_{t\geq 0}$ on $L^p(X)$ as in Corollary 2.2, it is immediate from the semigroup property and Theorem 2.1 that $\varphi_0 = \operatorname{id}_X \mu$ -almost everywhere and $\varphi_t \circ \varphi_s = \varphi_{t+s} \mu$ -almost everywhere for all $s, t \geq 0$. Therefore, the family $(\varphi_t)_{t\geq 0}$ forms a *semiflow* modulo null-sets, see Section 3. Note, however, that it can in general not be made into a semiflow by simply discarding a null-set since the identity $\varphi_t \circ \varphi_s = \varphi_{t+s}$ might hold outside of a null-set depending on sand t.

Corollary 2.4. Let A be the generator of a C_0 -semigroup $(T(t))_{t\geq 0}$ on a space $L^p(X)$, where $X = (X, \Sigma, \mu)$ is a standard probability space and $1 \leq p < \infty$. Then the equivalent assertions (i), (ii) and (iii) of Corollary 1.7 are also equivalent to

(iv) There exists a family $(\varphi_t)_{t\geq 0}$ of measurable and measure-preserving maps on X such that $T(t)f = f \circ \varphi_t$ for all $f \in L^p(X)$ and $t \geq 0$.

Proof. Assume (i) of Corollary 1.7. Then Corollary 2.2 shows that there are measurable maps $\varphi_t \colon X \to X$ such that $T(t)f = f \circ \varphi_t$. Moreover, for $M \in \Sigma$

$$\mu(\varphi_t^{-1}(M)) = \langle \mathbb{1}_{\varphi_*^{-1}(M)}, \mathbb{1}_X \rangle = \langle T(t)\mathbb{1}_M, \mathbb{1}_X \rangle = \langle \mathbb{1}_M, \mathbb{1}_X \rangle = \mu(M)$$

and so each φ_t is measure-preserving. On the other hand, (iv) implies (ii) with $\theta_t = \varphi_t^*$.

The next lemma shows that in the case of groups, the boundedness assumption in Theorem 1.1 (iii) is superfluous. This allows us to recover [4, Theorem 1.1] as Corollary 2.6. The idea of the following proof is based on the proof of [4, Theorem 2.5].

Lemma 2.5. Let X be a finite measure space and $(T(t))_{t>0}$ be a semigroup on $L^{\infty}(X)$, strongly continuous with respect to $\|\cdot\|_p$ where $1 \leq p < \infty$. Then the mapping $t \mapsto \|T(t)\|_{\mathscr{L}(L^{\infty}(X))}$ is locally bounded.

Proof. For $f \in L^{\infty}(X)$ and t > 0, setting $q := 1 - \frac{1}{n}$

$$||T(t)f||_{\infty} = \sup \{ |\langle T(t)f, g \rangle| \colon g \in L^{1}(\mathbf{X}) \text{ and } ||g||_{1} = 1 \}$$

= sup {|\langle T(t)f, g \rangle| \vert g \in L^{q}(\mathbf{X}) and ||g||_{1} = 1 }

Since $(T(t))_{t>0}$ is strongly continuous with respect to $\|\cdot\|_p$, $\langle T(\cdot)f, g \rangle$ is continuous for $g \in L^q(X)$. Hence, $\|T(\cdot)f\|_{\infty}$ is lower semicontinuous, being the supremum of continuous functions. Therefore, the sets $A_n := \{t > 0 : \|T(t)f\|_{\infty} \le n\}$ are closed. Since $\mathbb{R}_{>0} = \bigcup_n A_n$, Baire's category theorem yields that for every $k \in \mathbb{N}$, there are $0 < a_k < b_k \le \frac{1}{k}$ and $M_k \in \mathbb{N}$ such that $[a_k, b_k] \subseteq A_{M_k}$. By the semigroup property,

$$\sup_{s \in [a_k+t,b_k+t]} ||T(s)f||_{\infty} = \sup_{s \in [a_k+t,b_k+t]} ||T(t)T(s-t)f||_{\infty}$$
$$\leq \sup_{s \in [a_k,b_k]} ||T(t)||_{\infty} ||T(s)f||_{\infty} \leq M_k ||T(t)||_{\infty}$$

for each t > 0. This shows that $||T(\cdot)f||_{\infty}$ is locally bounded for each $f \in L^{\infty}(X)$ and the claim now follows by the principle of uniform boundedness. \Box

Corollary 2.6 ([4, Theorem 1.1]). Let A be the generator of a unitary C_0 -group $(T(t))_{t \in \mathbb{R}}$ on $L^2(X)$ where $X = (X, \Sigma, \mu)$ is a standard probability space. Then the following assertions are equivalent.

- (i) For every $t \in \mathbb{R}$ there exists an essentially invertible measurable and measurepreserving map $\varphi_t \colon X \to X$ such that $T(t)f = f \circ \varphi_t$ for all $f \in L^2(X)$.
- (ii) The space L[∞](X) is invariant under (T(t))_{t≥0} and A is a derivation on D(A) ∩ L[∞](X).

Proof. The implication (i) \implies (ii) is a consequence of Corollary 1.7. In order to prove the converse implication, we observe that it follows from Lemma 2.5 and Remark 1.6 that A and $(T(t))_{t\geq 0}$ as well as -A and $(T(-t))_{t\geq 0}$ fulfill condition (iii) in Theorem 1.1. Corollary 2.2 therefore shows that $T(t) = T_{\varphi_t}$ for measurable maps $\varphi_t \colon X \to X$ and $t \ge 0$. The essential invertibility of the maps φ_t follows from [3, Proposition 7.12] and [3, Corollary 7.21]. Also, since each T(t) is unitary and a Markov operator, one shows as in Corollary 2.4 that each φ_t is measurepreserving.

Corollary 2.7. Let A be the generator of a C_0 -semigroup $(S(t))_{t\geq 0}$ on a space $L^p(X)$, where X is a standard probability space and $1 \leq p < \infty$, such that $\mathbb{1} \in D(A)$ and $q \coloneqq A\mathbb{1} \in L^{\infty}(X)$. Then $(S(t))_{t\geq 0}$ is a lattice semigroup if and only if $q \in L^{\infty}(X; \mathbb{R})$ and there exists a family $(\varphi_t)_{t\geq 0}$ of measurable maps on X corresponding to a strongly continuous Koopman semigroup on $L^p(X)$ such that

(3)
$$S(t)f = \exp\left(\int_0^t q \circ \varphi_s \,\mathrm{d}s\right) \cdot (f \circ \varphi_t)$$

for all $f \in L^p(X)$ and $t \ge 0$.

Proof. If $(S(t))_{t\geq 0}$ is a lattice semigroup, it follows from Theorem 1.8 that there exists a Markov lattice semigroup $(T(t))_{t\geq 0}$ on $L^p(X)$ with generator (A-q, D(A)) such that (1) holds. The representation (3) hence follows from Corollary 2.2. Conversely, every semigroup of the form (3) with a real-valued q is a lattice semigroup.

3. TOPOLOGICAL MODEL

We have seen in Corollary 2.2 that on a standard probability space $X = (X, \Sigma, \mu)$ every strongly continuous Markov lattice semigroup $(T(t))_{t\geq 0}$ on $L^1(X)$ is induced by a family $(\varphi_t)_{t\geq 0}$ of measurable maps on X. Since $(T(t))_{t\geq 0}$ is a semigroup, one has $\varphi_0 = \mathrm{id}_X$ and $\varphi_s \circ \varphi_t = \varphi_{s+t}$ almost everywhere, using the uniqueness in Theorem 2.1. Recall that a family $(\varphi_t)_{t>0}$ of maps on a set X is called a *semiflow* if $\varphi_0 = \mathrm{id}_X$ and $\varphi_t \circ \varphi_s = \varphi_{s+t}$ for $s, t \ge 0$. A semiflow on a topological space X is called *continuous* if the map $\Phi: X \times \mathbb{R}_+ \to X$, $(x,t) \mapsto \varphi_t(x)$ is continuous. Similarly, a semiflow on a measure space is called *measurable* if Φ is measurable. It is called measure-preserving if for each t > 0 the map φ_t is measurable and measurepreserving. In that case, there is an induced semigroup of operators $(T_{\varphi_t})_{t>0}$ on $L^{1}(X)$ called the *Koopman semigroup* induced by the semiflow. This semigroup is weakly measurable if the semiflow is measurable but even then it need not be strongly continuous: Consider, e.g., the semiflow on $(\{0,1\}, \mathcal{P}(\{0,1\}), \frac{1}{2}(\delta_0 + \delta_1))$ with $\varphi_0 = \mathrm{id}_{\{0,1\}}$ and $\varphi_t \equiv 0$ for each t > 0. For measurable and measurepreserving flows $(\varphi_t)_{t\in\mathbb{R}}$ (defined analogously to semiflows) on separable measure spaces, however, the induced Koopman group is in fact strongly continuous.

Proposition 3.1. Let $X = (X, \Sigma, \mu)$ be a separable, finite measure space and $(\varphi_t)_{t\geq 0}$ a measurable and measure-preserving semiflow on X. Then the induced Koopman semigroup $(T(t))_{t\geq 0}$ on $L^p(X)$ is strongly continuous on $(0, \infty)$ for $1 \leq p < \infty$. If $(\varphi_t)_{t\in\mathbb{R}}$ is a measurable and measure-preserving flow on X, the induced Koopman group on $L^p(X)$ is strongly continuous on all of \mathbb{R} for $1 \leq p < \infty$.

Proof. Take $f \in L^p(X)$. Then the function $f \circ \Phi$ is measurable on $X \times \mathbb{R}$ and for each $g \in L^q(X)$ with $\frac{1}{q} = 1 - \frac{1}{p}$ the map

$$t \mapsto \langle T(t)f,g \rangle = \int_X f(\Phi(x,t))g(x) \,\mathrm{d}\mu(x)$$

is measurable (cf. [10, Theorem 1.7.15]). Therefore, the semigroup $(T(t))_{t\geq 0}$ is weakly measurable and even strongly measurable by [6, Theorem 3.5.5] since $L^p(X)$ is separable. Since strongly measurable semigroups are strongly continuous on $(0, \infty)$ by [6, Theorem 10.2.3], $(T(t))_{t\geq 0}$ is indeed strongly continuous. The case of flows follows immediately.

We now show that for such measurable flows, one can construct a continuous flow on a compact metric space with an invariant probability measure such that the two flows are isomorphic. This will be done by first proving that every strongly continuous Markov lattice semigroup is similar to a Koopman semigroup induced by a continuous semiflow. An analogous result was recently proved for bi-Markov lattice embedding representations of locally compact groups by de Jeu and Rozendaal, see [7, Theorem 5.14].

Definition 3.2. Let $X = (X, \Sigma, \mu)$ and $Y = (Y, \Sigma', \mu')$ be finite measure spaces. We say that two measurable semiflows $(\varphi_t)_{t\geq 0}$ and $(\psi_t)_{t\geq 0}$ on X and Y are *isomorphic* if there is a measure-preserving and essentially invertible map $\rho: X \to Y$ such that $\psi_t \circ \rho = \rho \circ \varphi_t$ almost everywhere for each $t \geq 0$. We say that two Markov lattice semigroups $(T(t))_{t\geq 0}$ and $(S(t))_{t\geq 0}$ on $L^1(X)$ and $L^1(Y)$ are *Markov similar* if there is an invertible bi-Markov lattice homomorphism $\Phi: L^1(X) \to L^1(Y)$ such

that $S(t) \circ \Phi = \Phi \circ T(t)$ for each $t \ge 0$. The same notions are defined for flows and operator groups analogously.

We call continuous flows (resp. Koopman semigroups) as described above topological models for measurable flows (resp. Markov lattice semigroups). See also [3, Section 12.3] for this terminology and similar results in the time-discrete case. For the sake of simplicity, we restrict ourselves to the case p = 1. The idea of the proof of the following result was kindly provided to us by Markus Haase. The employed technique of topological models, however, dates back much further.

Theorem 3.3 (Topological Model). Let A be the generator of a strongly continuous Markov lattice semigroup $(T(t))_{t\geq 0}$ on a space $L^1(X)$, where $X = (X, \Sigma, \mu)$ is a finite measure space. Then there exist a compact space K, a continuous semiflow $(\psi_t)_{t\geq 0}$ on K and a strictly positive Borel probability measure ν such that the semiflow $(\psi_t)_{t\geq 0}$ induces a Koopman-semigroup on $L^1(K, \nu)$ which is Markov similar to the semigroup $(T(t))_{t\geq 0}$ on $L^1(X)$. The measure ν is $(\psi_t)_{t\geq 0}$ -invariant if and only if $(T(t))_{t\geq 0}$ is a bi-Markov lattice semigroup.

Proof. Consider $\mathcal{A} := \{f \in L^{\infty}(\mathbf{X}) : s \mapsto T(s)f \text{ is } \|\cdot\|_{\infty}\text{-continuous}\}$. Since each operator T(t) is contractive on $L^{\infty}(\mathbf{X})$ and multiplicative by Lemma 1.3, \mathcal{A} is an algebra and clearly $\mathbb{1} \in \mathcal{A}$. Furthermore, \mathcal{A} is closed with respect to $\|\cdot\|_{\infty}$ and closed under conjugation. Therefore, \mathcal{A} is a commutative C*-algebra invariant under $(T(t))_{t>0}$.

We show that \mathcal{A} is dense in $L^1(X)$. The strong continuity of $(T(t))_{t\geq 0}$ on $L^1(X)$ implies that $\|\cdot\|_1-\lim_{t\searrow 0}\frac{1}{t}\int_0^t T(r)f\,dr = f$ for each $f \in L^\infty(X)$. Therefore, it suffices to show that $\int_0^t T(r)f\,dr \in \mathcal{A}$ for $f \in L^\infty(X)$. For all $0 \leq s \leq t$ and $f \in L^\infty(X)$

$$\begin{aligned} \left| T(s) \int_0^t T(r) f \, \mathrm{d}r - \int_0^t T(r) f \, \mathrm{d}r \right| &= \left| \int_0^t T(s+r) f \, \mathrm{d}r - \int_0^t T(r) f \, \mathrm{d}r \right| \\ &= \left| \int_s^{t+s} T(r) f \, \mathrm{d}r - \int_0^t T(r) f \, \mathrm{d}r \right| \\ &\leq \left| \int_t^{t+s} T(r) f \, \mathrm{d}r \right| + \left| \int_0^s T(r) f \, \mathrm{d}r \right| \\ &\leq 2s \|f\|_\infty \mathbb{1} \end{aligned}$$

since each T(t) is $\|\cdot\|_{\infty}$ -contractive. This shows that $s \mapsto T(s) \int_0^t T(r) f \, dr$ is continuous at zero and hence on $[0,\infty)$ with respect to $\|\cdot\|_{\infty}$. Therefore, \mathcal{A} is dense in $L^1(X)$.

By a combination of the Gelfand-Naimark theorem and the Riesz representation theorem as in [3, Section 12.3] or [7, Theorem 5.14] one obtains a compact space K, a *-isomorphism $\Phi: \mathcal{A} \to C(K)$ with $\Phi \mathbb{1} = \mathbb{1}$, a unique probability measure ν on K such that

$$\int_X \Phi^{-1} g \,\mathrm{d}\mu = \int_K g \,\mathrm{d}\nu$$

for all $g \in C(K)$ and a semiflow semiflow $(\psi_t)_{t\geq 0}$ on K such that $T(t)|_{\mathcal{A}} = \Phi^{-1} \circ T_{\psi_t} \circ \Phi$. By [3, Theorem 4.17], the semiflow ψ is continuous, cf. also [8, Theorem B-II.3.4].

Moreover, Φ extends to a bi-Markov lattice homomorphism $\Phi: L^1(X) \to L^1(K, \nu)$. Let $(S(t))_{t>0}$ denote the semigroup $(T(t))_{t>0}$ induces on $L^1(K, \nu)$ via Φ . Then

(4)
$$S(t)[f]_{\nu} = [f \circ \psi_t]_{\nu}$$

for all continuous functions $f \in C(K)$. By a standard approximation argument, this holds for all bounded, Baire-measurable functions, cf. [3, Theorem E.1]. Via monotone approximation, (4) extends to all positive integrable functions and is hence valid for all $[f]_{\nu} \in L^1(K, \nu)$. Finally, $(T(t))_{t\geq 0}$ is bi-Markov if and only if $(S(t))_{t\geq 0}$ is, if and only if

$$\int_{K} f \, \mathrm{d}\nu = \int_{K} S(t) f \, \mathrm{d}\nu = \int_{K} f \circ \psi_{t} \, \mathrm{d}\nu$$

for all $f \in L^1(K, \nu)$. This is given if and only if $(\psi_t)_{t \ge 0}$ preserves ν .

Proposition 3.4. If, in the situation of Theorem 3.3, the measure space is separable, then the compact space K can be chosen to be metrizable.

Proof. Let \mathcal{A} be the algebra

$$\mathcal{A} := \{ f \in \mathcal{L}^{\infty}(\mathcal{X}) \colon s \mapsto T(s)f \text{ is } \| \cdot \|_{\infty} \text{-continuous} \}$$

from the proof of Theorem 3.3. Since $L^1(X)$ is separable and \mathcal{A} is dense in the former, there is a countable dense subset D_0 of \mathcal{A} . We set

$$D := \{T(t)f \colon f \in D_0, t \in \mathbb{Q}_+\} \subseteq \mathcal{A}$$

and denote by \mathcal{A}_0 the C*-subalgebra of \mathcal{A} generated by D. The algebra \mathcal{A}_0 is then separable, dense in $L^1(X)$ and since $\mathcal{A}_0 \subseteq \mathcal{A}$, $T(t)\mathcal{A}_0 \subseteq \mathcal{A}_0$ for not only $t \in \mathbb{Q}_+$ but $t \in \mathbb{R}_+$. To complete the proof, one can now proceed as in the proof of Theorem 3.3 with \mathcal{A} replaced by \mathcal{A}_0 , obtaining a compact representation space which is metrizable because C(K) is separable.

Remark 3.5. With slight notational adjustments, the proofs of the previous two results also work for Markov lattice groups and continuous flows.

Corollary 3.6. Let $X = (X, \Sigma, \mu)$ be a standard probability space and $(\varphi_t)_{t \in \mathbb{R}}$ a measurable and measure-preserving flow on X. Then there are a compact metric space K, a continuous flow $(\psi_t)_{t \in \mathbb{R}}$ on K and a strictly positive $(\psi_t)_{t \in \mathbb{R}}$ -invariant Borel probability measure ν on K so that the flows $(\varphi_t)_{t \in \mathbb{R}}$ and $(\psi_t)_{t \in \mathbb{R}}$ are isomorphic.

Proof. By Proposition 3.1 and Theorem 1.1, the flow $(\varphi_t)_{t\in\mathbb{R}}$ induces a bi-Markov group on $L^1(X)$ and so Remark 3.5 shows that there are a compact metric space K, a continuous flow $(\psi_t)_{t\in\mathbb{R}}$ and a strictly positive $(\psi_t)_{t\in\mathbb{R}}$ -invariant probability measure ν on K such that the groups $(T(t))_{t\in\mathbb{R}}$ and $(S(t))_{t\in\mathbb{R}}$ induced by the flows $(\varphi_t)_{t\in\mathbb{R}}$ and $(\psi_t)_{t\in\mathbb{R}}$ are Markov similar via an invertible bi-Markov lattice homomorphism Φ . Applying von Neumann's theorem shows that there is a measurable and measure-preserving map $\rho: Y \to X$ such that $\Phi = T_{\rho}$ and ρ is essentially invertible because Φ is invertible, see [3, Corollary 7.21]. The identity $\Phi \circ T(t) = S(t) \circ \Phi$ now shows that $\varphi_t \circ \rho = \rho \circ \psi_t \nu$ -almost everywhere, see [3, Proposition 7.19]. \Box

Remark 3.7. Corollary 3.6 is similar to [1, Theorem 5] but for two important differences: On the one hand, the authors of [1] work with a slightly stronger notion of isomorphism of flows. On the other hand, the models considered in [1] need not be compact.

4. Ergodic flows

In this section, we give an operator-theoretic proof for the fact that a measurepreserving ergodic flow on a separable measure space comprises at most countably many non-ergodic maps if it induces a strongly continuous group on $L^2(X)$. This is a special case of [9, Theorem 1] where \mathbb{R}^k -actions where considered. First, recall the following two properties. **Definition 4.1.** Let $X = (X, \Sigma, \mu)$ be a finite measure space.

- (a) A measure-preserving semiflow $(\varphi_t)_{t\geq 0}$ on X is called *ergodic*, if for every $A \in \Sigma$ one has $A \subseteq \varphi_t^{-1}(A)$ modulo null-sets for each $t \geq 0$ if and only if $A = \emptyset$ or A = X modulo null-sets.
- (b) A Markov lattice semigroup $(T(t))_{t\geq 0}$ on $L^1(X)$ is called *irreducible* if there are no nontrivial closed $(T(t))_{t\geq 0}$ -invariant ideals in $L^1(X)$.

Similar notions can be defined for measure-preserving flows and Markov lattice groups in a straightforward way and it is not difficult to see that a flow $(\varphi_t)_{t\in\mathbb{R}}$ is ergodic if and only if one/both of the associated semiflows $(\varphi_t)_{\pm t\geq 0}$ is/are ergodic. As a consequence of [3, Theorem 7.10], the two properties in Definition 4.1 are equivalent for a (semi)flow and its corresponding Koopman (semi)group. Obviously, if some φ_s is ergodic (meaning that $A \subseteq \varphi_s^{-1}(A)$ implies $\mu(A) \in \{0,1\}$), so is the semiflow $(\varphi_t)_{t\geq 0}$, while the converse is not true in general. However, it follows from [9, Theorem 1] that for an ergodic flow $(\varphi_t)_{t\in\mathbb{R}}$ all but at most countably many maps φ_t are ergodic. In the following we give a short proof of this fact using the Perron-Frobenius spectral theory. We shall prove this for semiflows for which the induced Koopman semigroup is strongly continuous on $L^2(X)$. In light of the previous remarks, this implies [9, Theorem 1].

Theorem 4.2. Let $(\varphi_t)_{t\geq 0}$ be a measure-preserving semiflow on a separable finite measure space $X = (X, \Sigma, \mu)$ such that the corresponding Koopman semigroup $(T(t))_{t\geq 0}$ is strongly continuous on $L^2(X)$. If $(\varphi_t)_{t\geq 0}$ is ergodic, then at most countably many maps φ_t are not ergodic.

Proof. Let $(\varphi_t)_{t\geq 0}$ be measure-preserving and ergodic and its induced Koopman semigroup $(T(t))_{t\geq 0}$ strongly continuous on $L^2(X)$ with generator A. Since the eigenspaces of an isometry on $L^2(X)$ are pairwise orthogonal, the point-spectrum of T(t) is countable for each $t \geq 0$ because $L^2(X)$ is separable. The spectral inclusion theorem [5, Theorem IV.3.6] implies that the boundary point spectrum $G \coloneqq P\sigma(A) \cap i\mathbb{R}$ is countable.

For a fixed s > 0 the map φ_s is not ergodic if and only if dim fix $(T(s)) \ge 2$, see [3, Proposition 7.15]. By the spectral mapping theorem for the point spectrum [5, Corollary IV.3.8]

 $\operatorname{fix}(T(s)) = \overline{\operatorname{span}} \{ f \in L^2(X) \colon Af = i\lambda f \text{ for some } i\lambda \in G \text{ with } e^{i\lambda s} = 1 \}.$

Moreover, for an irreducible Markov lattice semigroup the common fixed space

$$\bigcap_{t \ge 0} \operatorname{fix}(T(t)) = \ker A$$

is one-dimensional since it is a Banach sublattice of $L^2(X)$ and for real-valued $f \in \bigcap_{t \ge 0} \operatorname{fix}(T(t))$ all characteristic functions $\mathbb{1}_{[f>c]}, c \in \mathbb{R}$, are contained in the fixed space (use [3, Exercise 7.13]). Thus, dim $\operatorname{fix}(T(s)) \ge 2$ if and only if there exists $i\lambda \in G$ such that $\lambda s = 2\pi k$ for some $k \in \mathbb{Z} \setminus \{0\}$. Since G is countable, this can only be the case for at most countably many s > 0.

As a consequence of Theorem 4.2, Proposition 3.1 and the remarks made above, we note the following corollary.

Corollary 4.3. Let $(\varphi_t)_{t \in \mathbb{R}}$ be a measurable and measure-preserving flow on a separable finite measure space X. If $(\varphi_t)_{t \in \mathbb{R}}$ is ergodic, then at most countably many maps φ_t are not ergodic.

The assertion in Theorem 4.2 does not hold for non-separable spaces. Take for example the Bohr compactification $\mathbb{b}\mathbb{R}$ of the additive group of real numbers and define for $g \in \mathbb{b}\mathbb{R}$ the translation $\varphi_q \colon \mathbb{b}\mathbb{R} \to \mathbb{b}\mathbb{R}$, $h \mapsto h + g$. Then $(\varphi_t)_{t \in \mathbb{R}}$ is a

REFERENCES

continuous flow on b \mathbb{R} preserving the Haar measure m. Denote the corresponding Koopman group on $L^2(b\mathbb{R}, m)$ by $(T(t))_{t \in \mathbb{R}}$. If $f \in L^2(b\mathbb{R}, m)$ is such that $T(t)f = f \circ \varphi_t = f$ for all $t \in \mathbb{R}$, then $f \circ \varphi_g = f$ for all $g \in b\mathbb{R}$ since \mathbb{R} is dense in b \mathbb{R} . It follows that f is constant almost everywhere. Because T(t)f = f is equivalent to T(-t)f = f for each $t \geq 0$, this shows that the flow $(\varphi_t)_{t \geq 0}$ is ergodic. However, φ_t is not ergodic since T(t)f = f for all periodic functions with period t.

Acknowledgement. We express our sincere gratitude towards the anonymous referee for his insightful observations and detailed comments.

References

- [1] Warren Ambrose and Shizuo Kakutani. "Structure and continuity of measurable flows". *Duke Math. J.* 9 no. 1 (1942), pp. 25–42.
- [2] Roland Derndinger and Rainer Nagel. "Der Generator stark stetiger Verbandshalbgruppen auf C(X) und dessen Spektrum". Math. Ann. 245 no. 2 (1979), pp. 159–174.
- [3] Tanja Eisner, Bálint Farkas, Markus Haase, and Rainer Nagel. Operator Theoretic Aspects of Ergodic Theory. Vol. 272. Graduate Texts in Mathematics. Springer International Publishing, 2015.
- [4] Tom ter Elst and Mariusz Lemańczyk. "On one-parameter Koopman groups". *Ergodic Theory Dynam. Systems* 37 no. 5 (2015), pp. 1635–1656.
- [5] Klaus-Jochen Engel and Rainer Nagel. One-Parameter Semigroups for Linear Evolution Equations. Vol. 194. Graduate Texts in Mathematics. Springer-Verlag, New York, 2000.
- [6] Einar Hille and Ralph S. Phillips. Functional Analysis and Semi-Groups. Vol. 31. Colloquium Publications. American Mathematical Society, Providence, 1957.
- [7] Marcel de Jeu and Jan Rozendaal. "Disintegration of positive isometric group representations on L^p-spaces". *Positivity* 21 no. 2 (2017), pp. 673–710.
- [8] R. Nagel, ed. One-parameter Semigroups of Positive Operators. Vol. 1184. Lecture Notes in Mathematics. Springer-Verlag, Berlin, 1986.
- Charles Pugh and Michael Shub. "Ergodic elements of ergodic actions". Compositio Math. 23 no. 1 (1971), pp. 115–122.
- [10] Terence Tao. An Introduction to Measure Theory. Vol. 126. Graduate Studies in Mathematics. American Mathematical Society, Providence, 2011.

NIKOLAI EDEKO, MATHEMATISCHES INSTITUT, UNIVERSITÄT TÜBINGEN, AUF DER MORGENSTELLE 10, D-72076 TÜBINGEN, GERMANY

Email address: nied@fa.uni-tuebingen.de

Moritz Gerlach, Institut für Mathematik, Universität Potsdam, Karl-Liebknecht-Strasse 24–25, 14476 Potsdam

 $Email \ address: \ \tt{moritz.gerlach} @uni-potsdam.de \\$

VIKTORIA KÜHNER, MATHEMATISCHES INSTITUT, UNIVERSITÄT TÜBINGEN, AUF DER MORGEN-STELLE 10, D-72076 TÜBINGEN, GERMANY *Email address*: viku@fa.uni-tuebingen.de