# On *s*-extremal singly even self-dual [24k+8, 12k+4, 4k+2] codes<sup>\*</sup>

Masaaki Harada<sup>†</sup>and Akihiro Munemasa<sup>‡</sup>

June 13, 2022

#### Abstract

A relationship between s-extremal singly even self-dual [24k + 8, 12k + 4, 4k + 2] codes and extremal doubly even self-dual [24k + 8, 12k + 4, 4k + 4] codes with covering radius meeting the Delsarte bound, is established. As an example of the relationship, s-extremal singly even self-dual [56, 28, 10] codes are constructed for the first time. In addition, we show that there is no extremal doubly even self-dual code of length 24k+8 with covering radius meeting the Delsarte bound for  $k \ge 137$ . Similarly, we show that there is no extremal doubly even self-dual code of length 24k+8 with covering radius meeting the Delsarte bound for  $k \ge 137$ . Similarly, we show that there is no extremal doubly even self-dual code of length 24k + 16 with covering radius meeting the Delsarte bound for  $k \ge 148$ . We also determine the covering radii of some extremal doubly even self-dual codes of length 80.

#### **1** Introduction

A (binary) [n, k] code C is a k-dimensional vector subspace of  $\mathbb{F}_2^n$ , where  $\mathbb{F}_2$ denotes the finite field of order 2. All codes in this paper are binary. The parameter n is called the *length* of C. The *weight* wt(x) of a vector  $x \in \mathbb{F}_2^n$  is

<sup>\*</sup>Keywords: extremal doubly even self-dual code, *s*-extremal singly even self-dual code, covering radius. Mathematics Subject Classification: 94B05, 94B75.

<sup>&</sup>lt;sup>†</sup>Research Center for Pure and Applied Mathematics, Graduate School of Information Sciences, Tohoku University, Sendai 980–8579, Japan.

<sup>&</sup>lt;sup>‡</sup>Research Center for Pure and Applied Mathematics, Graduate School of Information Sciences, Tohoku University, Sendai 980–8579, Japan.

the number of non-zero components of x. A vector of C is called a *codeword* of C. The minimum non-zero weight of all codewords in C is called the *minimum weight* of C and an [n, k] code with minimum weight d is called an [n, k, d] code. The *automorphism group* of C consists of all permutations of the coordinates of C which preserve C. Two codes are *equivalent* if one can be obtained from the other by permuting the coordinates.

The dual code  $C^{\perp}$  of a code C of length n is defined as  $C^{\perp} = \{x \in \mathbb{F}_2^n \mid x \cdot y = 0 \text{ for all } y \in C\}$ , where  $x \cdot y$  is the standard inner product. A code C is called *self-dual* if  $C = C^{\perp}$ . The minimum weight d(C) of a self-dual code C of length n is bounded by  $d(C) \leq 4\lfloor n/24 \rfloor + 4$  unless  $n \equiv 22 \pmod{24}$  when  $d(C) \leq 4\lfloor n/24 \rfloor + 6 \pmod{19}$ . We say that a self-dual code meeting the upper bound is *extremal*.

Self-dual codes are divided into two classes, namely, doubly even selfdual codes and singly even self-dual codes. A self-dual code is called *doubly* even and singly even if all codewords have weight  $\equiv 0 \pmod{4}$  and if some codeword has weight  $\equiv 2 \pmod{4}$ , respectively. A doubly even self-dual code of length n exists if and only if  $n \equiv 0 \pmod{8}$ , while a singly even self-dual code of length n exists if and only if n is even.

Let C be a singly even self-dual code and let  $C_0$  denote the subcode of codewords having weight  $\equiv 0 \pmod{4}$ . Then  $C_0$  is a subcode of codimension 1. The shadow S of C is defined to be  $C_0^{\perp} \setminus C$ . Shadows were introduced by Conway and Sloane [8], in order to provide restrictions on the weight enumerators of singly even self-dual codes. The minimum weight d(D) of a coset D is the minimum non-zero weight of all vectors in the coset. Let C and S be a singly even self-dual code of length n and its shadow, respectively. Bachoc and Gaborit [2] showed that

$$d(S) \le \frac{n}{2} + 4 - 2d(C),\tag{1}$$

unless  $n \equiv 22 \pmod{24}$  and  $d(C) = 4\lfloor n/24 \rfloor + 6$ , in which case  $d(S) = \frac{n}{2} + 8 - 2d(C)$ . A singly even self-dual code C meeting the bound is called *s-extremal*. The existence of many *s*-extremal singly even self-dual codes was mentioned in [2]. For minimum weight 10, it is known that there is an *s*-extremal singly even self-dual code of lengths 46, 50, 52, 54, 58 (see [2, Section 7]).

The covering radius R(C) of a code C of length n is the smallest integer R such that spheres of radius R around codewords of C cover the space  $\mathbb{F}_2^n$ . It is known that the covering radius R(C) is the same as the largest value among the minimum weights of nontrivial cosets (see [18, Theorem 13 (i)]). The covering radius is a basic and important geometric parameter of a code. Let C be a code of length n and let  $A_i^{\perp}$  be the number of vectors of weight i in  $C^{\perp}$ . It was shown in [9] that  $R(C) \leq |\{i \mid A_i^{\perp} \neq 0, 0 < i \leq n\}|$ . The bound is called the Delsarte bound (see [1]). Much work has been done concerning covering radii of extremal doubly even self-dual codes (see e.g., [1], [14], [17], [20]).

In this paper, a relationship between s-extremal singly even self-dual [24k+8, 12k+4, 4k+2] codes and extremal doubly even self-dual [24k+2][8, 12k + 4, 4k + 4] codes with covering radius meeting the Delsarte bound, is established. As an example of the relationship, we actually construct sextremal singly even self-dual [56, 28, 10] codes from some extremal doubly even self-dual [56, 28, 12] codes with covering radius 10. The existence of such self-dual codes was not previously known. We also determine the covering radii of some known extremal doubly even self-dual codes of length 80, along with two new extremal doubly even self-dual codes of the same length. It is shown that there is no s-extremal singly even self-dual [24k+8, 12k+4, 4k+2]code for  $k \geq 137$ . By the above relationship, there is no extremal doubly even self-dual code of length 24k + 8 with covering radius meeting the Delsarte bound for k > 137. Finally, we show that there is no s-extremal singly even self-dual [24k + 16, 12k + 8, 4k + 4] code for  $k \ge 148$ . By Proposition 4 in [14], there is no extremal doubly even self-dual code of length 24k + 16with covering radius meeting the Delsarte bound for  $k \ge 148$ .

All computer calculations in this paper were done by MAGMA [4].

# 2 A relationship between *s*-extremal self-dual [24k+8, 12k+4, 4k+2] codes and certain extremal doubly even self-dual codes

In this section, a relationship between *s*-extremal singly even self-dual [24k + 8, 12k + 4, 4k + 2] codes and extremal doubly even self-dual [24k + 8, 12k + 4, 4k + 4] codes with covering radius 4k + 2, is established.

Let C be a singly even self-dual code and let  $C_0$  denote the subcode of codewords having weight  $\equiv 0 \pmod{4}$ . Then there are cosets  $C_1, C_2, C_3$ of  $C_0$  such that  $C_0^{\perp} = C_0 \cup C_1 \cup C_2 \cup C_3$ , where  $C = C_0 \cup C_2$  and  $S = C_1 \cup C_3$ . Two self-dual codes C and C' of length n are said to be *neighbors* if  $\dim(C \cap C') = n/2 - 1$ . If C is a singly even self-dual code of length divisible by 8, then C has two doubly even self-dual neighbors, namely  $C_0 \cup C_1$  and  $C_0 \cup C_3$  (see [5]).

**Theorem 1.** If C is an s-extremal singly even self-dual [24k+8, 12k+4, 4k+2] code, then the two doubly even self-dual neighbors  $C_0 \cup C_1$  and  $C_0 \cup C_3$  are extremal and their covering radii are 4k+2, meeting the Delsarte bound. Conversely, if D is an extremal doubly even self-dual [24k+8, 12k+4, 4k+4] code with covering radius 4k+2, then D has an s-extremal singly even self-dual [24k+8, 12k+4, 4k+4] dual [24k+8, 12k+4, 4k+2] neighbor.

*Proof.* Suppose that C is an s-extremal singly even self-dual [24k + 8, 12k + 4, 4k + 2] code. Since  $d(C_i) \ge 4k + 4$  (i = 0, 1, 3),  $C_0 \cup C_1$  and  $C_0 \cup C_3$  have minimum weight at least 4k + 4. By the bound on the minimum weight, both  $C_0 \cup C_1$  and  $C_0 \cup C_3$  have minimum weight 4k + 4, that is, the codes are extremal. The sets  $C_2 \cup C_3$  and  $C_1 \cup C_2$  are cosets of  $C_0 \cup C_1$  and  $C_0 \cup C_3$ , respectively. Since  $d(C_2) = 4k + 2$ ,  $d(C_2 \cup C_3) = d(C_1 \cup C_2) = 4k + 2$ . Hence, both  $C_0 \cup C_1$  and  $C_0 \cup C_3$  have covering radius at least 4k + 2. By the Delsarte bound, both  $C_0 \cup C_1$  and  $C_0 \cup C_3$  have covering radius 4k + 2.

Conversely, suppose that D is an extremal doubly even self-dual [24k + 8, 12k + 4, 4k + 4] code with covering radius 4k + 2. Then there is a coset v + D of minimum weight 4k + 2. Set  $D_0 = \{x \in D \mid v \cdot x = 0\}$ , which is a subcode of index 2. Then there are cosets  $D_1, D_2, D_3$  of  $D_0$  such that  $D_0^{\perp} = D_0 \cup D_1 \cup D_2 \cup D_3$ , where  $D = D_0 \cup D_2, v + D = D_1 \cup D_3, D_1$  consists of the vectors whose weight is congruent to 2 (mod 4) and  $D_3$  consists of the vectors whose weight is congruent to 0 (mod 4) [5, Lemma 3]. In addition,  $D_0 \cup D_1$  is a singly even self-dual code with shadow  $D_2 \cup D_3$ . Since  $d(D_1) = 4k + 2, d(D_2) \ge 4k + 4$  and  $d(D_3) \ge 4k + 4, D_0 \cup D_1$  is a singly even self-dual [24k + 8, 12k + 4, 4k + 2] code with shadow of minimum weight at least 4k + 4. By the upper bound (1), the minimum weight of the shadow is exactly 4k + 4. Of course,  $D_0 \cup D_1$  is a neighbor of D.

*Remark* 2. A similar result can be found in [14, Proposition 4] for *s*-extremal singly even self-dual [24k + 16, 12k + 8, 4k + 4] codes (see also Lemma 9).

Let  $A_i$  and  $B_i$  denote the number of codewords of weight i in a singly even self-dual code C of length n and the number of vectors of weight i in its shadow S, respectively. The weight enumerators of C and S are given by  $\sum_{i=0}^{n} A_i y^i$  and  $\sum_{i=0}^{n} B_i y^i$ , respectively. The weight enumerators of an s-extremal singly even self-dual [n, n/2, d] code and its shadow are uniquely determined [2].

**Corollary 3.** If C is an s-extremal singly even self-dual [24k+8, 12k+4, 4k+2] code, then the weight enumerator of a coset of minimum weight 4k+2 in the two doubly even self-dual neighbors  $C_0 \cup C_1$  and  $C_0 \cup C_3$  is given by

$$\sum_{i\equiv 2 \pmod{4}} A_i y^i + \sum_i \frac{B_i}{2} y^i.$$
 (2)

Proof. By Theorem 1,  $C_2 \cup C_3$  (resp.  $C_2 \cup C_1$ ) is a coset of minimum weight 4k + 2 in  $C_0 \cup C_1$  (resp.  $C_0 \cup C_3$ ). The weight enumerator of an extremal doubly even self-dual code of length n is uniquely determined. Since  $C_0 \cup C_1$  and  $C_0 \cup C_3$  are extremal doubly even self-dual codes,  $C_1$  and  $C_3$  have the identical weight enumerators. Hence, the weight enumerators of  $C_2 \cup C_3$  and  $C_2 \cup C_1$  are given by (2). By Corollary 1 to Theorem 1 in [1], the cosets of minimum weight 4k+2, if there are any, have unique weight enumerators.  $\Box$ 

As an example of Theorem 1, we consider the case k = 1. There are five inequivalent extremal doubly even self-dual codes of length 32 [7]. These codes are denoted by C81, C82, C83, C84 and C85 in [7, Table A]. Every extremal doubly even self-dual code of length 32 has covering radius 6 [1, Theorem 3]. These codes contain 11253, 14756, 12236, 11354 and 11321 cosets of minimum weight 6, respectively [1, Table II] and [6, Tables 1, 2, 6, 8]. By Theorem 1, 11253, 14756, 12236, 11354 and 11321 *s*-extremal singly even selfdual [32, 16, 6] codes are obtained, respectively. We verified by MAGMA [4] that the 11253, 14756, 12236, 11354 and 11321 codes are divided into 6, 2, 3, 7 and 10 inequivalent codes, respectively. By determining equivalence of the 28 codes, there are 19 inequivalent *s*-extremal singly even self-dual [32, 16, 6] codes. This classification coincides with the result listed in [2, p. 29].

# 3 Construction of s-extremal singly even selfdual [56, 28, 10] codes

In this section, we consider the case k = 2. Some *s*-extremal singly even self-dual [56, 28, 10] codes are constructed. The existence of such self-dual codes was not previously known (see [2]).

By using [8, (10), (11)], the weight enumerators of an *s*-extremal singly even self-dual [56, 28, 10] code and its shadow are determined as follows:

$$\begin{split} 1 + 308y^{10} + 3990y^{12} + 42900y^{14} + 311850y^{16} + 1583120y^{18} + 5847688y^{20} \\ + 15961680y^{22} + 32458965y^{24} + 49520856y^{26} + 56972740y^{28} + \cdots, \\ 8400y^{12} + 620928y^{16} + 11704000y^{20} + 64901760y^{24} + 113965280y^{28} + \cdots, \end{split}$$

respectively.

The existence of extremal doubly even self-dual codes of length 56 with covering radius 10 is known [14], [17] and [20]. Theorem 1 gives the existence of an s-extremal singly even self-dual [56, 28, 10] code. As an example, we actually construct s-extremal singly even self-dual [56, 28, 10] codes from the extremal double circulant doubly even self-dual [56, 28, 12] code  $C_{56.1}$  in [12] and the extremal doubly even self-dual [56, 28, 12] code in [3], which have covering radius 10. We denote the code in [3] by BYB. Ozeki [17] determined the complete coset weight distributions of  $C_{56,1}$  and BYB. The codes contain 2925 and 6552 cosets of minimum weight 10, respectively [17]. By Theorem 1, 2925 and 6552 s-extremal singly even self-dual [56, 28, 10] codes are obtained, respectively. We verified by MAGMA [4] that the 2925 codes are divided into 70 inequivalent codes. We denote the new codes by  $D_1, D_2, \ldots, D_{70}$ . These codes are constructed as  $(C_{56,1})_0 \cup (v + (C_{56,1})_0)$ , where  $(C_{56,1})_0 = \{x \in C_{56,1} \mid x \in C_{56,1} \mid x \in C_{56,1}\}$  $v \cdot x = 0$  and the supports supp(v) of v are listed in Table 1. The orders |Aut| of the automorphism groups calculated by MAGMA [4] are listed in Table 2. We verified by MAGMA [4] that the automorphism group of the code BYB acts on the 6552 cosets of minimum weight 10 transitively. This means that the 6552 s-extremal singly even self-dual [56, 28, 10] codes are equivalent. The unique s-extremal code E is constructed as  $(BYB)_0 \cup (v + (BYB)_0)$ , where  $(BYB)_0 = \{x \in BYB \mid v \cdot x = 0\}$  and

 $\operatorname{supp}(v) = \{1, 2, 3, 17, 21, 22, 28, 37, 51, 55\}.$ 

We verified by MAGMA [4] that the code E has automorphism group of order 9.

By comparing the automorphism groups of  $D_i$  (i = 1, 2, ..., 70) and E, we see that these 71 codes are pairwise inequivalent. Therefore, we have the following:

**Proposition 4.** There are at least 71 s-extremal singly even self-dual [56, 28, 10] codes.

The existence of such self-dual codes was not previously known (see [2]).

i	$\operatorname{supp}(v)$	i	$\operatorname{supp}(v)$
1	$\{1, 2, 3, 4, 5, 9, 10, 24, 54, 55\}$	2	$\{1, 2, 3, 6, 9, 10, 21, 23, 36, 45\}$
3	$\{1, 2, 3, 7, 11, 16, 32, 33, 46, 51\}$	4	$\{1, 2, 3, 7, 12, 13, 14, 31, 37, 46\}$
5	$\{1, 2, 3, 4, 5, 15, 49, 50, 53, 55\}$	6	$\{1, 2, 3, 19, 24, 26, 39, 40, 44, 48\}$
7	$\{1, 2, 3, 18, 25, 36, 38, 39, 42, 53\}$	8	$\{1, 2, 3, 6, 15, 27, 36, 40, 45, 46\}$
9	$\{1, 2, 3, 6, 27, 42, 43, 46, 55, 56\}$	10	$\{1, 2, 3, 9, 11, 33, 40, 42, 47, 50\}$
11	$\{1, 2, 4, 6, 10, 12, 24, 31, 46, 52\}$	12	$\{1, 2, 3, 7, 15, 25, 30, 33, 43, 51\}$
13	$\{1, 2, 3, 22, 23, 25, 31, 41, 45, 53\}$	14	$\{1, 2, 3, 14, 20, 28, 30, 35, 41, 53\}$
15	$\{1, 2, 4, 7, 17, 23, 28, 42, 54, 55\}$	16	$\{1, 2, 3, 20, 24, 26, 29, 34, 38, 50\}$
17	$\{1, 2, 4, 10, 11, 25, 27, 30, 37, 52\}$	18	$\{1, 2, 4, 7, 18, 20, 25, 30, 32, 36\}$
19	$\{1, 2, 3, 12, 16, 21, 24, 30, 31, 48\}$	20	$\{1, 2, 3, 12, 16, 24, 29, 35, 36, 41\}$
21	$\{1, 2, 3, 12, 19, 24, 30, 34, 39, 56\}$	22	$\{1, 2, 3, 5, 10, 25, 30, 36, 45, 55\}$
23	$\{1, 2, 3, 6, 15, 20, 23, 27, 33, 42\}$	24	$\{1, 2, 3, 23, 25, 29, 31, 39, 47, 53\}$
25	$\{1, 2, 3, 18, 21, 24, 44, 47, 52, 53\}$	26	$\{1, 2, 3, 4, 9, 27, 29, 32, 36, 37\}$
27	$\{1, 2, 3, 7, 15, 18, 24, 31, 37, 52\}$	28	$\{1, 2, 3, 18, 22, 26, 31, 34, 41, 43\}$
29	$\{1, 2, 3, 4, 6, 16, 42, 45, 50, 52\}$	30	$\{1, 2, 3, 4, 11, 26, 27, 29, 32, 47\}$
31	$\{1, 2, 3, 4, 11, 20, 25, 38, 48, 54\}$	32	$\{1, 2, 3, 12, 14, 15, 18, 23, 36, 54\}$
33	$\{1, 2, 3, 4, 24, 28, 31, 39, 47, 52\}$	34	$\{1, 2, 3, 4, 14, 19, 35, 39, 48, 49\}$
35	$\{1, 2, 3, 7, 17, 25, 41, 50, 55, 56\}$	36	$\{1, 2, 3, 5, 7, 13, 20, 34, 48, 52\}$
37	$\{1, 2, 3, 7, 15, 25, 38, 40, 42, 43\}$	38	$\{1, 2, 3, 4, 9, 20, 46, 48, 53, 54\}$
39	$\{1, 2, 3, 9, 11, 19, 24, 30, 32, 45\}$	40	$\{1, 2, 3, 4, 13, 26, 39, 48, 52, 55\}$
41	$\{1, 2, 3, 4, 11, 17, 28, 30, 46, 55\}$	42	$\{1, 2, 3, 8, 13, 20, 38, 41, 44, 56\}$
43	$\{1, 2, 4, 5, 28, 33, 34, 47, 51, 55\}$	44	$\{1, 2, 3, 23, 26, 29, 34, 37, 40, 49\}$
45	$\{1, 2, 3, 6, 12, 22, 28, 31, 41, 51\}$	46	$\{1, 2, 4, 7, 18, 20, 26, 29, 43, 53\}$
47	$\{1, 2, 3, 21, 32, 38, 42, 48, 52, 53\}$	48	$\{1, 2, 3, 10, 11, 12, 14, 17, 20, 50\}$
49	$\{1, 2, 3, 4, 6, 15, 19, 36, 45, 55\}$	50	$\{1, 2, 3, 5, 8, 11, 32, 34, 37, 42\}$
51	$\{1, 2, 3, 5, 9, 27, 32, 45, 46, 56\}$	52	$\{1, 2, 4, 6, 11, 13, 15, 16, 37, 47\}$
53	$\{1, 2, 3, 4, 8, 16, 17, 19, 37, 55\}$	54	$\{1, 2, 3, 18, 23, 34, 36, 44, 47, 53\}$
55	$\{1, 2, 3, 5, 6, 21, 25, 32, 33, 49\}$	56	$\{1, 2, 3, 23, 35, 37, 40, 45, 46, 53\}$
57	$\{1, 2, 3, 4, 24, 28, 37, 41, 45, 48\}$	58	$\{1, 2, 4, 7, 23, 29, 32, 37, 46, 50\}$
59	$\{1, 2, 4, 11, 16, 22, 43, 46, 52, 53\}$	60	$\{1, 2, 3, 6, 12, 14, 20, 37, 46, 47\}$
61	$\{1, 2, 4, 5, 22, 29, 36, 46, 49, 54\}$	62	$\{1, 2, 3, 22, 23, 24, 35, 40, 50, 53\}$
63	$\{1, 2, 3, 5, 12, 14, 26, 32, 42, 48\}$	64	$\{1, 2, 3, 7, 15, 23, 31, 38, 39, 42\}$
65	$\{1, 2, 4, 12, 27, 28, 31, 34, 42, 53\}$	66	$\{1, 2, 3, 7, 8, 14, 34, 36, 42, 52\}$
67	$\{1, 2, 3, 4, 28, 29, 32, 37, 38, 46\}$	68	$\{1, 2, 3, 17, 24, 34, 45, 46, 48, 49\}$
69	$\{1, 2, 3, 8, 9, 12, 14, 20, 42, 46\}$	70	$\{1, 2, 3, 4, 13, 18, 36, 37, 52, 55\}$

Table 1: New s-extremal singly even self-dual [56, 28, 10] codes

Table 2: Automorphism groups of  $D_i$  (i = 1, 2, ..., 70)

Aut	i
1	2, 3, 4, 7, 8, 9, 10, 11, 12, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 25, 27,
	31, 34, 37, 41, 43, 44, 45, 47, 48, 49, 55, 57, 58, 61, 63, 67, 68, 70
2	1, 5, 6, 13, 24, 26, 28, 29, 30, 32, 33, 35, 36, 39, 40, 42, 46, 50, 51, 52,
	53, 54, 56, 59, 60, 62, 64, 65, 66, 69
6	38

# 4 Covering radii of some extremal doubly even self-dual codes of length 80

In this section, we consider the case k = 3. The covering radii of some extremal doubly even self-dual codes of length 80 are determined.

The extended quadratic residue code  $QR_{80}$  of length 80 is an extremal doubly even self-dual code. It was shown in [10] that there are 11 extremal doubly even self-dual [80, 40, 16] codes with an automorphism of order 19, up to equivalence. The 11 codes are denoted by  $C_{80,1}, C_{80,2}, \ldots, C_{80,11}$  in [10]. Also, none of them is equivalent to  $QR_{80}$ . There are six (resp. five) inequivalent pure (resp. bordered) double circulant extremal doubly even self-dual codes of length 80 [11]. These codes are denoted by  $P_{80,i}$   $(i = 1, 2, \ldots, 6)$ (resp.  $B_{80,i}$   $(i = 1, 2, \ldots, 5)$ ) in [11]. Note that  $P_{80,1}$  and  $B_{80,5}$  are equivalent. Also, we verified by MAGMA [4] that  $P_{80,1}$  is equivalent to  $QR_{80}$ . Very recently, 14 more inequivalent extremal doubly even self-dual codes of length 80 have been found in [15]. These codes are denoted by  $\mathcal{L}_{80,i}$   $(i = 1, 2, \ldots, 14)$ in [15, Table 11]. Hence, at least 35 inequivalent extremal doubly even selfdual codes of length 80 exist<sup>1</sup>.

By the method given in [13], we found two extremal doubly even self-dual codes  $N_{80,1}$  and  $N_{80,2}$  of length 80. The code  $N_{80,i}$  (i = 1, 2) has generator matrix ( $I_{40}$   $M_{80,i}$ ), where  $I_{40}$  denotes the identity matrix of order 40 and  $M_{80,1}$  (resp.  $M_{80,2}$ ) is listed in Figure 1 (resp. Figure 2). In the figures, each row of  $M_{80,i}$  is written in octal using  $0 = (000), 1 = (001), \ldots, 6 = (110)$  and 7 = (111), together with a = (0) and b = (1), where, for example, the (1, 2)-entry indicates the second row and the (2, 1)-entry indicates the 5th row.

<sup>&</sup>lt;sup>1</sup>The number of known extremal doubly even self-dual codes of length 80 is incorrectly reported as 36 in [15, Theorem 6.1].

We verified by MAGMA [4] that  $N_{80,1}$  and  $N_{80,2}$  have automorphism groups of orders 32 and 64, respectively. Since none of the 35 known inequivalent extremal doubly even self-dual codes of length 80 has automorphism group of order 32, 64, we have the following:

**Proposition 5.** There are at least 37 inequivalent extremal doubly even selfdual codes of length 80.

By the Assmus–Mattson theorem, the supports of the codewords of minimum weight in an extremal doubly even self-dual code of length 80 form a self-orthogonal 3-(80, 16, 665) design. We verified that each of the 37 extremal doubly even self-dual codes of length 80 is generated by the codewords of minimum weight. Hence, we have the following:

**Corollary 6.** There are at least 37 non-isomorphic self-orthogonal 3-(80, 16, 665) designs having 2-rank 40.

 $\begin{array}{l} 0040734771561a\ 4434543163675b\ 2602474666733b\ 1715423124350b\\ 0352004645161a\ 3206160642302a\ 2660112641633b\ 6053127640067a\\ 7431246537036b\ 4577441737365b\ 2663035540577b\ 1725203477272b\\ 0346314420530a\ 4577753007251a\ 1114607163256a\ 4052516666522a\\ 1346325653523b\ 7600030245123a\ 0463176442323a\ 4625262036154b\\ 2706324200063a\ 5757747317034b\ 2373176750413a\ 5561262173200b\\ 5513453515272b\ 2251030451530a\ 2247576744126b\ 6240715602721b\\ 4243224261422a\ 6535307327614a\ 7642356744703a\ 7335772175344b\\ 3142160261567a\ 5075665327676b\ 2022127744732a\ 2372131202427b\\ 6216136061561a\ 0264135550102a\ 4526643473044a\ 1130243355350b \end{array}$ 

Figure 1: Matrix  $M_{80,1}$  (in octal)

We determine the covering radii of the codes  $C_{80,i}$  (i = 1, 2, ..., 11) in [10],  $P_{80,i}$  (i = 1, 2, ..., 6),  $B_{80,i}$  (i = 1, 2, 3, 4) in [11],  $\mathcal{L}_{80,i}$  (i = 1, 2, ..., 14) in [15] and the new codes  $N_{80,i}$  (i = 1, 2) as follows. By the Delsarte bound, the covering radius of an extremal doubly even self-dual code of length 80 is at most 14. By modifying the method in [16], we verified that there is no coset of minimum weight 13 and there is a coset of minimum weight 12 for the codes  $C_{80,i}$  (i = 1, 2, ..., 11),  $P_{80,i}$  (i = 1, 2, ..., 6),  $B_{80,i}$  (i = 1, 2, 3),  $\mathcal{L}_{80,i}$  (i = 1, 2, ..., 14) and  $N_{80,i}$  (i = 1, 2). In this calculation, the automorphism groups are useful to reduce the number of cosets which must be considered. Also, we verified that there is no coset of minimum weight 14 and there is a coset of minimum weight 13 for the code  $B_{80,4}$ . Hence, we have the following:  $\begin{array}{l} 2017612646116b \ 1314312723575b \ 7122225026433b \ 4035272764050b \\ 5472655005261a \ 6526731002002a \ 7140743001533b \ 4004001777410b \\ 5466360400441a \ 6520567600712a \ 7143664300677b \ 4005452237172b \\ 5466545260630a \ 1257102647151a \ 3143721054621b \ 6005430751155b \\ 3311203764154a \ 2120661405223a \ 5343727202023a \ 6672344101523a \\ 0751202337414b \ 0077116557334b \ 7453727110713a \ 7536344044677a \\ 0233202355172b \ 0206116566147b \ 7567327104226b \ 3560144042421b \\ 6214302356055b \ 3215556567514a \ 2162507104403a \ 2415123735044b \\ 6662731421667a \ 7022743210201a \ 0075001673345b \ 7452760442727b \\ 4241010156116b \ 2233013467575b \ 1206012233344a \ 4610412515050b \\ \end{array}$ 

Figure 2: Matrix  $M_{80,2}$  (in octal)

**Proposition 7.** The covering radii of the codes  $C_{80,i}$  (i = 1, 2, ..., 11),  $P_{80,i}$  (i = 1, 2, ..., 6),  $B_{80,i}$  (i = 1, 2, 3),  $\mathcal{L}_{80,i}$  (i = 1, 2, ..., 14) and  $N_{80,i}$  (i = 1, 2) are 12. The covering radius of the code  $B_{80,4}$  is 13.

A vector a of a coset U is called a coset leader of U if the weight of a is minimal in U. We give the support of a coset leader of a coset of minimum weight 13 in  $B_{80,4}$ :

 $\{2, 5, 8, 11, 14, 17, 20, 23, 26, 29, 32, 35, 38\}.$ 

The covering radii of extremal doubly even self-dual codes of length 80 are determined for the first time. The above proposition indicates that there are many extremal doubly even self-dual codes with covering radius not meeting the Delsarte bound. It is an open problem to determine whether there is an extremal doubly even self-dual [80, 40, 16] code with covering radius meeting the Delsarte bound or not. Equivalently, it is an open problem to determine whether there is an extremal singly even self-dual [80, 40, 14] code or not.

By Corollary 2 to Theorem 1 in [1], the cosets of minimum weights 13 and 14, if there are any, have unique weight enumerators. We verified by MAGMA [4] that the unique weight enumerator of a coset of minimum weight 13 in  $B_{80.4}$  is as follows:

$$\begin{split} & 560y^{13} + 12624y^{15} + 184080y^{17} + 2103920y^{19} + 18380960y^{21} \\ & + 124273760y^{23} + 661003616y^{25} + 2796701600y^{27} + 9492488720y^{29} \\ & + 26027430000y^{31} + 57969919280y^{33} + 105320725328y^{35} \\ & + 156558001600y^{37} + 190784587840y^{39} + \cdots . \end{split}$$

By using [8, (10), (11)], the weight enumerators of an *s*-extremal singly even self-dual [80, 40, 14] code and its shadow are determined as follows:

$$\begin{split} 1 + 3200y^{14} + 47645y^{16} + 640640y^{18} + 6452992y^{20} + 49304320y^{22} \\ + 294979360y^{24} + 1398270720y^{26} + 5294263040y^{28} + 16137190784y^{30} \\ + 39853463650y^{32} + 80135036800y^{34} + 131652451840y^{36} \\ + 177157460480y^{38} + 195552496832y^{40} + \cdots , \\ 99840y^{16} + 12859392y^{20} + 590187520y^{24} + 10587822080y^{28} \\ + 79708428800y^{32} + 263302574080y^{36} + 391107684352y^{40} + \cdots . \end{split}$$

respectively. By Corollary 3, a coset of minimum weight 14 of an extremal doubly even self-dual code of length 80, if there is any, has the following weight enumerator:

$$\begin{split} & 3200y^{14} + 49920y^{16} + 640640y^{18} + 6429696y^{20} + 49304320y^{22} \\ & + 295093760y^{24} + 1398270720y^{26} + 5293911040y^{28} + 16137190784y^{30} \\ & + 39854214400y^{32} + 80135036800y^{34} + 131651287040y^{36} \\ & + 177157460480y^{38} + 195553842176y^{40} + \cdots . \end{split}$$

### 5 Non-existence of certain extremal doubly even self-dual codes of length 24k + 8

**Theorem 8.** There is no s-extremal singly even self-dual [24k + 8, 12k + 4, 4k + 2] code for  $k \ge 137$ . Equivalently, there is no extremal doubly even self-dual [24k + 8, 12k + 4, 4k + 4] code with covering radius 4k + 2, meeting the Delsarte bound for  $k \ge 137$ .

*Proof.* Zhang [21] showed the non-existence of an extremal doubly even selfdual code of length 24k + 8 for  $k \ge 159$ . Hence, by Theorem 1, there is no *s*-extremal singly even self-dual [24k + 8, 12k + 4, 4k + 2] code for  $k \ge 159$ .

By using [8, (10), (11)], we numerically determined the weight enumerators of an *s*-extremal singly even self-dual [24k+8, 12k+4, 4k+2] code *C* and its shadow for each  $k \leq 158$ . This calculation was done by the program written in MAGMA [4] which is listed in Appendix A. Then we find that the weight enumerator of *C* contains a negative coefficient for  $k = 137, 138, \ldots, 158$ . The first assertion follows. The second assertion follows from Theorem 1 immediately.

# 6 Non-existence of certain extremal doubly even self-dual codes of length 24k + 16

The argument in the previous section can be applied to s-extremal singly even self-dual [24k + 16, 12k + 8, 4k + 4] codes and extremal doubly even self-dual [24k + 16, 12k + 8, 4k + 4] codes with covering radius 4k + 4, meeting the Delsarte bound.

**Lemma 9** (Harada and Munemasa [14]). There is an s-extremal singly even self-dual [24k + 16, 12k + 8, 4k + 4] code if and only if there is an extremal doubly even self-dual [24k + 16, 12k + 8, 4k + 4] code with covering radius 4k + 4, meeting the Delsarte bound.

**Theorem 10.** There is no s-extremal singly even self-dual [24k + 16, 12k + 8, 4k + 4] code for  $k \ge 148$ . Equivalently, there is no extremal doubly even self-dual [24k + 16, 12k + 8, 4k + 4] code with covering radius 4k + 4, meeting the Delsarte bound for  $k \ge 148$ .

*Proof.* Zhang [21] showed the non-existence of an extremal doubly even selfdual code of length 24k + 16 for  $k \ge 164$ . Hence, by Lemma 9, there is no *s*-extremal singly even self-dual [24k + 16, 12k + 8, 4k + 4] code for  $k \ge 164$ .

By using [8, (10), (11)], we numerically determined the weight enumerators of an s-extremal singly even self-dual [24k + 16, 12k + 8, 4k + 4] code C and its shadow S for each  $k \leq 163$ . This calculation was done by slightly modifying the program given in Appendix A to this case. Then we find that the weight enumerator of S contains a negative coefficient for  $k = 148, 149, \ldots, 163$ . The first assertion follows. The second assertion follows from by Lemma 9 immediately.

Acknowledgment. This work is supported by JSPS KAKENHI Grant Number 15H03633.

#### References

 E.F. Assmus, Jr. and V. Pless, On the covering radius of extremal selfdual codes, *IEEE Trans. Inform. Theory* 29 (1983), 359–363.

- [2] C. Bachoc and P. Gaborit, Designs and self-dual codes with long shadows, J. Combin. Theory Ser. A 105 (2004), 15–34.
- [3] V.K. Bhargava, G. Young and A.K. Bhargava, A characterization of a (56, 28) extremal self-dual code, *IEEE Trans. Inform. Theory* 27 (1981), 258–260.
- [4] W. Bosma, J. Cannon and C. Playoust, The Magma algebra system I: The user language, J. Symbolic Comput. 24 (1997), 235–265.
- [5] R. Brualdi and V. Pless, Weight enumerators of self-dual codes, *IEEE Trans. Inform. Theory* 37 (1991), 1222–1225.
- [6] P. Camion, B. Courteau and A. Monpetit, Coset weight enumerators of the extremal self-dual binary codes of length 32, Eurocode '92 (Udine, 1992), 17–29, CISM Courses and Lectures, 339, Springer, Vienna, 1993.
- [7] J.H. Conway, V. Pless and N.J.A. Sloane, The binary self-dual codes of length up to 32: a revised enumeration, J. Combin. Theory Ser. A 60 (1992), 183–195.
- [8] J.H. Conway and N.J.A. Sloane, A new upper bound on the minimal distance of self-dual codes, *IEEE Trans. Inform. Theory* 36 (1990), 1319– 1333.
- [9] P. Delsarte, Four fundamental parameters of a code and their combinatorial significance, *Inform. Control* **23** (1973), 407–438.
- [10] R. Dontcheva and M. Harada, Extremal doubly-even [80, 40, 16] codes with an automorphism of order 19, *Finite Fields Appl.* 9 (2003), 157– 167.
- [11] T.A. Gulliver and M. Harada, Classification of extremal double circulant self-dual codes of lengths 74–88, Discrete Math. 306 (2006), 2064–2072.
- [12] M. Harada, T.A. Gulliver and H. Kaneta, Classification of extremal double-circulant self-dual codes of length up to 62, *Discrete Math.* 188 (1998), 127–136.
- [13] M. Harada and H. Kimura, New extremal doubly-even [64, 32, 12] codes, Des. Codes Cryptogr. 6 (1995), 91–96.

- [14] M. Harada and A. Munemasa, On the covering radii of extremal doubly even self-dual codes, Adv. Math. Commun. 1 (2007), 251–256.
- [15] A. Kaya and B. Yildiz, Various constructions for self-dual codes over rings and new binary self-dual codes, *Discrete Math.* **339** (2016), 460– 469.
- [16] A. Munemasa, On the enumeration of self-dual codes, Proceedings of the Fourth Conference on Algebraic Geometry, Number Theory, Coding Theory and Cryptography, University of Tokyo, (2000), pp. 69–77.
- [17] M. Ozeki, On covering radii and coset weight distributions of extremal binary self-dual codes of length 56, *IEEE Trans. Inform. Theory* 46 (2000), 2359–2372.
- [18] V. Pless, Introduction to the Theory of Error-Correcting Codes (Third edition), A Wiley-Interscience Publication, John Wiley & Sons, Inc., New York, 1998.
- [19] E.M. Rains, Shadow bounds for self-dual codes, IEEE Trans. Inform. Theory 44 (1998), 134–139.
- [20] H.-P. Tsai, The covering radius of extremal self-dual code D11 and its application, IEEE Trans. Inform. Theory 43 (1997), 316–319.
- [21] S. Zhang, On the nonexistence of extremal self-dual codes, Discrete Appl. Math. 91 (1999), 277–286.

#### Appendix A

```
Rxy<x,y>:=PolynomialRing(Rationals(),2);
```

```
S:=function(k)
n:=24*k+8;
n2:=n div 2;
n8:=n div 8;
d2:=2*k+1;
basis:=[ (x<sup>2</sup>+y<sup>2</sup>)<sup>(n2-4*j)*(x<sup>2</sup>*y<sup>2</sup>*(x<sup>2</sup>-y<sup>2</sup>)<sup>2</sup>)<sup>j</sup>
: j in [0..n8] ];</sup>
```

```
basisCoeff:=Matrix(Rationals(),n8+1,d2,
  [ [ MonomialCoefficient(basis[i],x^(n-2*j)*y^(2*j))
  : j in [0..d2-1] ] : i in [1..n8+1] ] );
sbasis:=[ (-1)^j*2^(n2-6*j)*(x*y)^(n2-4*j)*(x^4-y^4)^(2*j)
  : j in [0..n8] ];
sbasisCoeff:=Matrix(Rationals(),n8+1,d2+1,
  [ [ MonomialCoefficient(sbasis[i],x^(n-2*j)*y^(2*j))
  : j in [0..d2] ] : i in [1..n8+1] ] );
mat:=HorizontalJoin(basisCoeff,sbasisCoeff);
rhsv:=Vector(Rationals(),Ncols(mat),
    [1] cat [ 0: i in [1..d2-1] ] cat [ 0: i in [0..d2] ]);
sol,nulsp:=Solution(mat,rhsv);
print Dimension(nulsp) eq 0;
return [ &+[ sol[i]*basis[i] : i in [1..n8+1] ],
 &+[ sol[i]*sbasis[i] : i in [1..n8+1] ] ];
end function;
for k in [1..158] do
WS:=S(k);
print k,Minimum(Coefficients(WS[1]));
end for;
```