# A NOTE ON THE COLORFUL FRACTIONAL HELLY THEOREM

#### MINKI KIM

ABSTRACT. Helly's theorem is a classical result concerning the intersection patterns of convex sets in  $\mathbb{R}^d$ . Two important generalizations are the colorful version and the fractional version. Recently, Bárány et al. combined the two, obtaining a colorful fractional Helly theorem. In this paper, we give an improved version of their result.

# 1. Introduction

Helly's theorem is one of the most well-known and fundamental results in combinatorial geometry, which has various generalizations and applications. It was first proved by Helly [10] in 1913, but his proof was not published until 1923, after alternative proofs by Radon [15] and König [13]. Recall that a family is *intersecting* if the intersection of all members is non-empty. The following is the original version of Helly's theorem.

**Theorem 1.1** (Helly's theorem, Helly [10]). Let  $\mathcal{F}$  be a finite family of convex sets in  $\mathbb{R}^d$  with  $|\mathcal{F}| \geq d+1$ . Suppose every (d+1)-tuple of  $\mathcal{F}$  is intersecting. Then the whole family  $\mathcal{F}$  is intersecting.

The following variant of Helly's theorem was found by Lovász, whose proof appeared first in Bárány's paper [4]. Note that the original Helly's theorem is obtained by setting  $\mathcal{F}_1 = \mathcal{F}_2 = \cdots = \mathcal{F}_{d+1}$ .

**Theorem 1.2** (Colorful Helly theorem, Lovász [4]). Let  $\mathcal{F}_1, \mathcal{F}_2, \ldots, \mathcal{F}_{d+1}$  be finite, non-empty families (color classes) of convex sets in  $\mathbb{R}^d$  such that every colorful (d+1)-tuple is intersecting. Then for some  $1 \le i \le d+1$ , the whole family  $\mathcal{F}_i$  is intersecting.

One way to generalize Helly's theorem is by weakening the assumption: not necessarily all but only a positive fraction of (d+1)-tuples are intersecting. The following theorem shows how the conclusion changes.

**Theorem 1.3** (Fractional Helly theorem, Katchalski and Liu [12]). For every  $\alpha \in (0,1]$ , there exists  $\beta = \beta(\alpha,d) \in (0,1]$  such that the following holds: Let  $\mathcal{F}$  be a finite family of convex sets in  $\mathbb{R}^d$  with  $|\mathcal{F}| \geq d+1$ . If at least  $\alpha\binom{|\mathcal{F}|}{d+1}$  of the (d+1)-tuples in  $\mathcal{F}$  are intersecting, then  $\mathcal{F}$  contains an intersecting subfamily of size at least  $\beta|\mathcal{F}|$ .

Date: August 13, 2018.

The fractional variant of Helly's theorem first appeared as a conjecture on interval graphs, i.e. intersection graphs of families of intervals on  $\mathbb{R}$ . Abbott and Katchalski [1] proved that  $\beta = 1 - \sqrt{1 - \alpha}$  is optimal for every family whose intersection graph is a chordal graph. Note that, by the result of Gavril [8], interval graphs are chordal graphs.

The fractional Helly theorem for arbitrary dimensions was proved by Katchalski and Liu [12]. Their proof gives a lower bound  $\beta \geq \alpha/(d+1)$ . However, it seems natural that  $\beta$  tends to 1 as  $\alpha$  tends to 1, since the original Helly's theorem implies that  $\beta = 1$  when  $\alpha = 1$ . Later, the quantitatively sharp value  $\beta(\alpha, d) = 1 - (1 - \alpha)^{1/(d+1)}$  was found by Kalai [11] and Eckhoff [6]. The result follows from the upper bound theorem for families of convex sets.

The (p,q)-theorem, another important generalization of Helly's theorem, deals with a weaker version of the assumption, the so-called (p,q)-condition: for every p members in a given family, there are some q members of the family that are intersecting. For instance, the (d+1,d+1)-condition in  $\mathbb{R}^d$  is the hypothesis of Helly's theorem. The (p,q)-theorem was proved by Alon and Kleitman [2], settling a conjecture by Hadwiger and Debrunner [9].

The proof of the (p, q)-theorem is quite long and involved, using various techniques. However, one of the most crucial ingredients is the fractional Helly theorem. See the survey paper by Eckhoff [7] and the textbook by Matousek [14] for an overview and further knowledge of this field.

Recently, Bárány et al. [5] established the colorful and fractional versions of the (p,q)-theorem. A key ingredient in their proof was a colorful variant of the fractional Helly theorem.

**Theorem 1.4** (Bárány-Fodor-Montejano-Oliveros-Pór [5]). Let  $\mathcal{F}_1, \mathcal{F}_2, \ldots, \mathcal{F}_{d+1}$  be finite, non-empty families (color classes) of convex sets in  $\mathbb{R}^d$ , and assume that  $\alpha \in (0,1]$ . If at least  $\alpha |\mathcal{F}_1| \cdots |\mathcal{F}_{d+1}|$  of the colorful (d+1)-tuples are intersecting, then some  $\mathcal{F}_i$  contains an intersecting subfamily of size  $\frac{\alpha}{d+1} |\mathcal{F}_i|$ .

Note that for  $\alpha=1$  we recover the hypothesis of the colorful Helly theorem, and we should therefore expect the value  $\beta=1$  (rather than  $\beta=\frac{1}{d+1}$ ). Bárány et al. therefore proposed the problem of showing that the function  $\beta$  in Theorem 1.4 tends to 1 as  $\alpha$  tends to 1.

In this paper, we solve the problem of Bárány et al.

**Theorem 1.5.** For every  $\alpha \in (0, 1]$ , there exists  $\beta = \beta(\alpha, d) \in (0, 1]$  tending to 1 as  $\alpha$  tends to 1 such that the following holds: Let  $\mathcal{F}_1, \mathcal{F}_2, \ldots, \mathcal{F}_{d+1}$  be finite, non-empty families (color classes) of convex sets in  $\mathbb{R}^d$ . If at least  $\alpha |\mathcal{F}_1| \cdots |\mathcal{F}_{d+1}|$  of the colorful (d+1)-tuples are intersecting, then for some  $1 \le i \le d+1$ ,  $\mathcal{F}_i$  contains an intersecting subfamily of size  $\beta |\mathcal{F}_i|$ .

See the survey paper by Amenta, Loera, and Soberón [3] for an overview of recent results and open problems related to Helly's theorem.

## 2. Proof of Theorem 1.5

2.1. **The Matching number of hypergraphs.** Let  $\mathcal{H}$  be an r-uniform hypergraph on a vertex set X. A subset  $S \subseteq X$  is said to be an *independent set* in  $\mathcal{H}$  if the induced sub-hypergraph  $\mathcal{H}[S]$  contains no hyperedge. The *independence number*  $\alpha(\mathcal{H})$  of  $\mathcal{H}$  is the cardinality of a maximum independent set in  $\mathcal{H}$ . A *matching* of  $\mathcal{H}$  is a set of pairwise disjoint edges in  $\mathcal{H}$ . The *matching number*  $\nu(\mathcal{H})$  of  $\mathcal{H}$  is the cardinality of a maximum matching in  $\mathcal{H}$ . For our result, we need the following observation.

**Observation 2.1.** Let  $\mathcal{H} = (X, E)$  be an r-uniform hypergraph with |X| = n. Suppose  $\alpha(\mathcal{H}) < cn$  for some  $c \in (0, 1]$ . Let M be a maximum matching in  $\mathcal{H}$ . Note that  $X \setminus M$  is an independent set it  $\mathcal{H}$ . If not, assume that there is an edge e contained in  $X \setminus M$ . Then  $M \cup \{e\}$  is a matching in  $\mathcal{H}$ , which is a contradiction to the maximality of M. Thus  $|X \setminus M| = n - rv(\mathcal{H}) \le \alpha(\mathcal{H}) < cn$ , so  $v(\mathcal{H}) > \frac{n-cn}{r}$ .

2.2. **Proof of Theorem 1.5.** Theorem 1.6 is implied by the following more explicit result.

**Theorem 2.2.** For every  $\alpha \in (0,1]$ , the following holds: Let  $\mathcal{F}_1, \mathcal{F}_2, \ldots, \mathcal{F}_{d+1}$  be finite families (color classes) of convex sets in  $\mathbb{R}^d$ . If at least  $\alpha |\mathcal{F}_1| \cdots |\mathcal{F}_{d+1}|$  of the colorful (d+1)-tuples are intersecting, then for some  $1 \le i \le d+1$ ,  $\mathcal{F}_i$  contains an intersecting subfamily of size at least

$$\max\left\{\frac{\alpha}{d+1}, 1 - (d+1)(1-\alpha)^{\frac{1}{d+1}}\right\} |\mathcal{F}_i|.$$

The following is a key lemma of the proof of Theorem 2.2.

**Lemma 2.3.** Choose any (d+1) members from each color class, say  $\mathcal{F}'_1, \ldots, \mathcal{F}'_{d+1}$ . If each of  $\mathcal{F}'_i$  is not intersecting, then at least one of colorful (d+1)-tuple is not intersecting.

*Proof.* This follows directly from the colorful Helly theorem.

*Proof of Theorem* 2.2. It is sufficient to show that for every  $\alpha \in [1 - \frac{1}{(d+1)^{(d+1)}}, 1]$ , if at least  $\alpha |\mathcal{F}_1| \cdots |\mathcal{F}_{d+1}|$  of the colorful (d+1)-tuples are intersecting, then some  $\mathcal{F}_i$  contains an intersecting subfamily of size at least  $(1 - (d+1)(1-\alpha)^{\frac{1}{d+1}})|\mathcal{F}_i|$ .

Let  $\mathcal{F}$  be the disjoint union of  $\mathcal{F}_1, \mathcal{F}_2, \dots, \mathcal{F}_{d+1}$ . For each  $1 \le i \le d+1$ , denote  $n_i = |\mathcal{F}_i|$  and define a (d+1)-uniform hypergraph  $H_i := (\mathcal{F}_i, E_i)$  whose vertices are the members in  $\mathcal{F}_i$  and hyperedges are non-intersecting (d+1)-tuples in  $\mathcal{F}_i$ . Let  $v_i = v(H_i)$  for each  $1 \le j \le d+1$ .

Also define a (d + 1)-uniform hypergraph  $H := (\mathcal{F}, E)$  whose vertices are the members in  $\mathcal{F}$  and hyperedges are intersecting colorful (d + 1)-tuples in  $\mathcal{F}$ .

Given  $\alpha \in [1 - \frac{1}{(d+1)^{(d+1)}}, 1]$ , let  $\gamma = \gamma(\alpha, d) = 1 - (d+1)(1-\alpha)^{\frac{1}{d+1}}$ . For contraction, assume that in each family  $\mathcal{F}_j$ , every subfamily of size at least  $\gamma n_j$  has an empty intersection.

By Lemma 2.3, we have  $\alpha n_1 \cdots n_{d+1} \leq |E| \leq n_1 \cdots n_d + 1 - \nu_1 \cdots \nu_{d+1}$ . Recall that by Observation 2.1,  $\nu_j > \frac{n_j - \gamma n_j}{d+1} = \left(\frac{1-\gamma}{d+1}\right) n_j$  for each  $1 \leq j \leq d+1$ . Then we obtain

$$\alpha n_{1} \cdots n_{d+1} \leq n_{1} \cdots n_{d+1} - \nu_{1} \cdots \nu_{d+1} 
< n_{1} \cdots n_{d+1} - \left(\frac{1-\gamma}{d+1}\right)^{d+1} n_{1} \cdots n_{d+1} 
= \left(1 - \left(\frac{1-\gamma}{d+1}\right)^{d+1}\right) n_{1} \cdots n_{d+1},$$

hence  $\alpha < 1 - (\frac{1-\gamma}{d+1})^{d+1} = \alpha$ , which is a contradiction.

Thus, there should exist  $1 \le i \le d+1$  such that  $\mathcal{F}_i$  contains an intersecting subfamily of size  $(1-(d+1)(1-\alpha)^{\frac{1}{d+1}})n_i$ .

## 3. The Upper bound

First recall that in the fractional Helly theorem, the upper bound is given by

$$\beta = \beta(\alpha, d) \le (1 - (1 - \alpha)^{\frac{1}{d+1}}).$$

This can be seen by the following well-known construction, which also shows the exactness of upper bound theorem for convex sets [6][11].

**Example 3.1.** Let  $\mathcal{F}$  consist of  $\lfloor \beta n \rfloor - (d+1)$  copies of  $\mathbb{R}^d$  and  $n - \lfloor \beta n \rfloor + (d+1)$  hyperplanes in general position. Denote by  $f_d(\mathcal{F})$  the number of intersecting (d+1)-tuples in  $\mathcal{F}$ . Note that

$$\alpha \binom{n}{d+1} = f_d(\mathcal{F}) = \binom{n}{d+1} - \binom{n - (\lfloor \beta n \rfloor - (d+1))}{d+1}$$

$$< \binom{n}{d+1} - \binom{n - \lfloor \beta n \rfloor}{d+1}$$

$$\leq \binom{n}{d+1} - (1-\beta)^{d+1} \binom{n}{d+1}.$$

The colorful version of this example gives an upper bound for the colorful fractional Helly theorem.

**Theorem 3.2.** For every  $\alpha \in (0,1]$ , there exist finite families (color classes)  $\mathcal{F}_1, \ldots, \mathcal{F}_{d+1}$  of convex sets in  $\mathbb{R}^d$  such that the following holds.  $\alpha |\mathcal{F}_1| \cdots |\mathcal{F}_{d+1}|$  of the colorful (d+1)-tuples are intersecting, but in each color class  $\mathcal{F}_i$ , the maximum cardinality of an intersecting subfamily is at most  $(1-(1-\alpha)^{\frac{1}{d+1}})|\mathcal{F}_i|$ .

*Proof.* It follows from the following construction. Let  $\mathcal{F}_i$  consist of  $\lfloor \beta n \rfloor - d$  copies of  $\mathbb{R}^d$  and  $n - \lfloor \beta n \rfloor + d$  hyperplanes in general position. Moreover, let all hyperplanes in  $\mathcal{F}_1 \cup \cdots \cup \mathcal{F}_{d+1}$  be in general position. Note that each of  $\mathcal{F}_i$  has an intersecting subfamily of size at most  $\beta n$ . The number of colorful (d+1)-tuples is given by

$$\alpha n^{d+1} = n^{d+1} - (n - \lfloor \beta n \rfloor + d)^{d+1} < n^{d+1} - (1 - \beta)^{d+1} n^{d+1}.$$

As *n* tends to infinity, one may have that  $\alpha = 1 - (1 - \beta)^{d+1}$ , i.e.  $\beta = 1 - (1 - \alpha)^{\frac{1}{d+1}}$ .  $\square$ 

#### 4. Remarks

In this note, we found upper and lower bounds on the function  $\beta(\alpha, d)$  in the colorful fractional Helly theorem, however, there remains a large gap between them. It would be interesting to determine the exact value of  $\beta(\alpha, d)$ .

**Problem 4.1.** What is the exact value of  $\beta = \beta(\alpha, d)$  in Theorem 1.5?

It is easy to see that  $\beta(\alpha, 1) = 1 - \sqrt{1 - \alpha}$  is the optimal bound for d = 1. We conjecture that  $\beta(\alpha, d) = 1 - (1 - \alpha)^{\frac{1}{d+1}}$  is the optimal bound for d > 1.

## ACKNOWLEDGMENT

I would like to thank my advisor Prof.Andreas Holmsen for introducing the problem and providing insightful comments. I also would like to thank Dr.Ilkyoo Choi for his advice to improve the readability of the paper.

#### REFERENCES

- [1] H. Abbott and M. Katchalski. A Turán type problem for interval graphs. *Discrete Math.*, 25(1):85–88, 1979.
- [2] N. Alon and D. J. Kleitman. Piercing convex sets and the Hadwiger-Debrunner (p, q)-problem. *Adv. Math.*, 96(1):103–112, 1992.
- [3] N. Amenta, J. A. D. Loera, and P. Soberón. Helly's theorem: New variations and applications, 2015.
- [4] I. Bárány. A generalization of Carathéodory's theorem. *Discrete Math.*, 40(2-3):141–152, 1982.
- [5] I. Bárány, F. Fodor, L. Montejano, D. Oliveros, and A. Pór. Colourful and fractional (*p*, *q*)-theorems. *Discrete Comput. Geom.*, 51(3):628–642, 2014.
- [6] J. Eckhoff. An upper-bound theorem for families of convex sets. *Geom. Dedicata*, 19(2):217–227, 1985.
- [7] J. Eckhoff. Helly, Radon, and Carathéodory type theorems. In *Handbook of convex geometry, Vol. A, B*, pages 389–448. North-Holland, Amsterdam, 1993.
- [8] F. Gavril. The intersection graphs of subtrees in trees are exactly the chordal graphs. *J. Combinatorial Theory Ser. B*, 16:47–56, 1974.
- [9] H. Hadwiger and H. Debrunner. Über eine Variante zum Hellyschen Satz. *Arch. Math. (Basel)*, 8:309–313, 1957.
- [10] E. Helly. Über mengen konvexer körper mit gemeinschaftlichen punkte. *Jahresber. Deutsch. Math.-Verein.*, 32:175–176, 1923.

- [11] G. Kalai. Intersection patterns of convex sets. Israel J. Math., 48(2-3):161–174, 1984.
- [12] M. Katchalski and A. Liu. A problem of geometry in  $\mathbb{R}^n$ . *Proc. Amer. Math. Soc.*, 75(2):284–288, 1979.
- [13] D. König. Über konvexe körper. Math. Z., 14:208–210, 1922.
- [14] J. Matoušek. *Lectures on discrete geometry*, volume 212 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2002.
- [15] J. Radon. Mengen konvexer Körper, die einen gemeinsamen Punkt enthalten. *Math. Ann.*, 83(1-2):113–115, 1921.

Minki Kim, Department of Mathematical Sciences, KAIST, Daejeon, South Korea.

E-mail address: kmk90@kaist.ac.kr