Separable local fractional differential equations

Kiran M. Kolwankar*

Department of Physics, Ramniranjan Jhunjhunwala College, Ghatkopar(W) Mumbai 400086 (Dated: July 18, 2018)

Abstract

The concept of local fractional derivative was introduced in order to be able to study the local scaling behavior of functions. However it has turned out to be much more useful. It was found that simple equations involving these operators naturally incorporate the fractal sets into the equations. Here, the scope of these equations has been extended further by considering different possibilities for the known function. We have also studied a separable local fractional differential equation along with its method of solution.

INTRODUCTION AND DEFINITIONS

The derivatives and integrals of non-integer order [2–4] have been found to be useful in successfully describing scaling processes. The realm of applications of such a fractional calculus is fast expanding with ever new developments rapidly taking place in the field of statistical and nonlinear physics over last few years [5–11]. The most often used definition of the fractional derivative is through the so called Riemann-Liouville fractional derivative [2–4]. For q, the order of the derivative, between zero and one it is given by:

Definition .1 The Riemann-Liouville fractional derivative of a function f of order q (0 < q < 1) is defined as:

$$D_x^q f(x') = \begin{cases} D_{x+}^q f(x'), & x' > x, \\ D_{x-}^q f(x'), & x' < x. \end{cases}$$

$$= \frac{1}{\Gamma(1-q)} \begin{cases} \frac{d}{dx'} \int_x^{x'} f(t)(x'-t)^{-q} dt, & x' > x, \\ -\frac{d}{dx'} \int_{x'}^{x} f(t)(t-x')^{-q} dt, & x' < x. \end{cases}$$
(1)

Clearly this is nonlocal and depends on the limit x. The value of x is usually dictated by the problem one is investigating. In some cases it is appropriate to put it equal to $\pm \infty$. In which case it is called Weyl derivative. Another choice of the limit is made in the following definition called the Local fractional derivative (LFD) defined as follows [12]:

Definition .2 The local fractional derivative of order q (0 < q < 1) of a function $f \in C^0$: $\mathbb{R} \to \mathbb{R}$ is defined as

$$\mathcal{D}^q f(x) = \lim_{x' \to x} D_x^q (f(x') - f(x))$$

if the limit exists in \mathbb{R} .

Such a local definition of fractional derivative is useful in studying the local scaling behavior in fractal and multifractal functions. This definition naturally appears in the local fractional Taylor expansion [12] giving it a geometrical interpretation which is a simple extension of that of ordinary derivatives. That is, just as the first order derivative is a coefficient of the local linear approximation, the LFD is the coefficient of the local power law approximation. It should be noted that the extra limit in the definition of the LFD makes it very different from other definitions of fractional derivatives and some of its properties are very different from the non-local versions of fractional derivatives (see [17, 22, 23] for more mathematical

properties). One important difference is that though it reduces to the usual derivative when q=1, the LFD is not an analytic function of q for a given function. For example, if the function is smooth then the LFD of any order q less than one is zero. In fact, in general for any continuous nondifferentiable function there exists a critical order of differentiability between zero and one below which all the derivatives are zero and above which they do not exist. This also means that with this definition one can not take an LFD of order greater than the critical order. In [33] we have introduced a recursive definition of LFD which allows us to overcome this problem. Also, as of now, we do not know any example of a continuous everywhere but nowhere differentiable function for which the LFD exists at the critical order. But here we take a point of view that such functions nevertheless exist and hope that this aspect will receive rigorous treatment in future, possibly by using nonstandard analysis. This critical order of differentiability is equivalent to the local Hölder exponent or the local power law exponent [27], and for this formalism to yield meaningful results we should work at the order which is equal to this exponent. Several authors [15, 16, 18–21, 26] have attempted to take this work further

A SIMPLE LOCAL FRACTIONAL DIFFERENTIAL EQUATIONS

Once we have the definition of LFD, the next natural step is to consider equations involving LFD, the local fractional differential equation (LFDE). The simplest such equation is

$$\mathcal{D}^q f(x) = g(x) \tag{2}$$

where g(x) is a known function and f(x) an unknown. Using the local fractional Taylor expansion [12] its solution can be written as a generalised Riemann sum giving

$$f(x) = \int g(x)d^{q}x = \lim_{N \to \infty} \sum_{i=0}^{N-1} g(x_{i}^{*}) \frac{(x_{i+1} - x_{i})^{q}}{\Gamma(q+1)}$$
(3)

where $g(x_i^*)$ is an appropriately chosen point in the interval $[x_i, x_{i+1}]$. In [32], it was shown that this integral does not exist if g(x) is a continuous function in any interval.

Yet, two classes of functions g(x) can be identified which will yield a nontrivial function f(x) as a solution. The first class corresponds to the functions which have fractal support. An example $g(x) = 1_C(x)$, where $1_C(x)$ is an indicator or characteristic function of a Cantor set C, was considered in [29]. Then, if $q = \alpha$, the dimension of the support of the function, the sum above converges since only a few terms contribute to the sum giving rise to a finite solution. This function, called the "devil's staircase", changes only on the points of the Cantor set C and is constant everywhere else. We denote this solution by $P_C(x)$. The second class consists of rapidly oscillating functions which oscillate around zero in any small interval. These oscillations then result in cancellations in the sum in (3) again giving rise to a finite solution. A realization of the white noise is one example in this class of functions. This example has been discussed in [30, 32].

In this work, we are going to restrict to the first class discussed above, that is, the one involving functions having fractal support. The treatment so far is far from being rigorous and we have relied only on the handwaving arguments. There are several issues involved here. It is easy to convince that the sum in (3) will lead to a devil's staircase like functions when g(x) has a fractal support but, for a rigorous treatment, the limit has to be established by considering upper and lower generalized Riemann sums. Furthermore, proving that the inverse also holds is even more difficult. It is clear that the staircase like solution will have local scaling (Hölder) exponent equal to α , the dimension, at the points of the support. So the critical order of the solution will be α which is consistent with the order of the original differential equation. But as we have discussed before, it is difficult to show that the LFD at the critical order exists. This difficulty arises owing to the oscillations around the local power law which exist in fractal functions. But this is not a serious point as these oscillations, as shown in [28], are small and do not affect the power law exponent. But it becomes a hurdle in generalizing the fundamental theorem of calculus to LFDs. A lot of efforts would be needed, possibly with minor modification of the definition of LFD or a treatment using nonstandard analysis, to develope the theory rigorously. In fact, Mandelbrot [1] has already advocated studying such integrals using nonstandard analysis. In this work, we are not going to get into these technical difficulties but instead develop the theory, albeit symbolically, assuming that the quantities defined exist.

Carpinteri and Cornetti [18] used the concept of LFD to introduce a relation between the strain and the displacement when the strain is localized on the fractal set. They proposed

$$\epsilon^*(x) = \mathcal{D}^{\alpha} u(x)$$

where $\epsilon^*(x)$ is a renormalised strain and u(x) is the displacement. This also offers a physical

interpretation to LFD. This formalism is useful in studying the structural properties of concrete-like or disordered materials. In [17], we calculated resultant of a stress distribution and its moment when the stress is distributed over a fractal set. Further use of LFD and the fractal integral for the principle of virtual work and to generalize the constitutive equations of elasticity has been made in [24, 31].

SEPARABLE LOCAL FRACTIONAL DIFFERENTIAL EQUATIONS

In this section we develop the theory of LFDEs further. First we consider more general forms of the function g(x) and then we treat more general right hand side in (2) which can depend even on the unknown function.

Now consider $g(x) = \bar{g}(x)1_C(x)$ where $\bar{g}(x)$ is some sufficiently well-behaved function. The multiplication by $1_C(x)$ makes g(x) a function with fractal support. In such a situation the solution f(x) in Eq. (3) can in principle be computed. The exact result will also depend on the fractal set supporting the function. Some explicit calculations have been performed in [17] when $\bar{g}(x) = x$ or x^2 and when the support is either a middle 1/3 Cantor set or a middle 1/p Cantor set. A similar method can be used for any power x^n . And if $\bar{g}(x)$ is real analytic then one can use its power series representation and an approximating scheme for the solution (3) can be set up which can at least be evaluated numerically.

Next nontrivial step is to consider $g(x) = g_1(x)1_{C_1}(x) + g_2(x)1_{C_2}(x)$ where C_1 and C_2 are different fractal sets with the same dimension α and g_1 , g_2 are different functions. Clearly, the integral operator in (3) is a linear operator and the result of considering such a g(x) in the solution is an addition of two devil's staircase like functions coming from two individual terms in above expression for g(x). It is important to realise that this is an interesting new step since it allows to incorporate effect of two different processes on two different fractal sets in one single equation and is not just another devil's staircase on a simple union of two sets. That is, the g(x) can not be written as $(g_1(x) + g_2(x))1_{C_1 \cup C_2}(x)$ even when C_1 and C_2 are disjoint. It is not difficult to imagine physical situations where such a function is needed. For example, we can generalise the situation discussed in [29] in which a local fractional diffusion equation was solved for a diffusion process taking place in fractal time. Now suppose that there are two different processes happening simultaneously but on two different fractal time sets of the same dimension but possibly with different lacunarity and

with different constants of diffusion. This could be a result of two different kind of traps. This choice of the known function will allow us to incorporate such situations into equations.

Now, in order to incorporate further complex situations, we rewrite the Eq. (2) using a different notation and also generalize it. We write

$$\frac{dy}{dx^q} = g(x, y). (4)$$

Here the LHS is a mere change in notation where LFD is written in more suggestive form. First of all y is now our unknown function to be found out. The order of the LFD q appears only in the denominator since the numerator is only a first order difference. The RHS is now a generalization wherein the known function g depends also on g. Now we assume that the equation is separable, that is, $g(x,y) = \bar{g}(x)1_C(x)h(y)$ a product of functions which depend only on g or g. This also means that the fractal set g does not change with g. Therefore we have

$$\frac{dy}{dx^q} = \bar{g}(x)1_C(x)h(y). \tag{5}$$

We can formally rewrite this equations as

$$\frac{dy}{h(y)} = \bar{g}(x)1_C(x)dx^q \tag{6}$$

and integrate to obtain the solution apart from the constant of integration

$$\int \frac{dy}{h(y)} = \int \bar{g}(x) 1_C(x) dx^q \tag{7}$$

where the RHS is now the integral as in Eq. (3). These steps can be justified using the chain rule for LFD, proved in [22], as in the following. Owing to the chain rule we can write two auxiliary equations in place of Eq. (5):

$$\frac{dy}{du} = h(y)$$
 and $\frac{du}{dx^q} = \bar{g}(x)1_C(x)$. (8)

These equations can be solved separately and then the solutions can be combined to obtain the solution (7).

As an example, consider h(y) = y and $\bar{g}(x) = 1$. In this case, the solution becomes $y = A \exp(P_C(x))$. A similar case had arisen in the solution of the local fractional Fokker-Planck equation in [29] though, over there, it was solved using different steps. As another example, if we consider $h(y) = y^2$ keeping the $\bar{g}(x) = 1$ then the solution is $y = -1/P_C(x)$. This

demonstrates that complex situations involving fractals can be incorporated in a differential equation and solutions can be obtained which includes the information of the fractal set in the form of the function $P_C(x)$.

CONCLUDING DISCUSSIONS

In this paper we have briefly reviewed the introduction to LFD and related developments regarding the equations involving the LFDs. The simplest example of the so called LFDE was also recalled. The main purpose of this paper was to take these developments further. To this end, we have step by step increased the complexity in the RHS of the simple LFDE. From the indicator function of the Cantor set we first considered the case of a function multiplied by the indicator function and discussed its solution. Then we took a next nontrivial step consisting of addition of two functions with different fractal sets of the same dimension as the support. Lastly, we studied a more general function which depended on both x and y but was separable, that is, could be written as a product of functions only of x and only of y. We discussed and rationalized the method of its solution. The examples considered demonstrate the utility of this approach in studying the process taking place on a fractal set. One can also add a white noise to the known function g on the RHS of Eqs. (2) or (4) which will yield a random process with additional log-periodic oscillations. Such stochastic differential equations will be considered elsewhere.

Taking this line of exploration further will be interesting and also challenging. Also, it is necessary to establish these results rigorously. These developments will pave a way for fractal dynamical systems.

ACKNOWLEDGEMENT

The author would like to thank Council for Scientific and Industrial Research, India for financial assistance.

[1] B. B. Mandelbrot, The Fractal Geometry of Nature (Freeman, San Franscisco, 1982).

^{*} Kiran.Kolwankar@gmail.com

- [2] K. B. Oldham and J. Spanier *The Fractional Calculus* (Academic Press, New York, 1974).
- [3] I. Podlubny, Fractional Differential Equations, Academic Press, San Diego, 1999.
- [4] S. G. Samko, A. A. Kilbas and O. I. Marichev, Fractional Integrals and Derivatives: Theory and Applications, Gordon and Breach, Yverdon, 1993.
- [5] B. J. West and M. Bologna and P. Grigolini, *Physics of Fractal Operators*, Springer, New York, 2003.
- [6] R. Hilfer(Ed.), Applications of Fractional Calculus in Physics, World Scientific, 2000.
- [7] R. Metzler and J. Klafter, Phys. Rep. 339, 1 (2000).
- [8] G. M. Zaslavsky, Phys. Rep. 371, 461 (2002).
- [9] R. Metzler and J. Klafter, J. Phys. A: Math. Gen. 7, R161 (2004).
- [10] I. Grigorenko and E. Grigorenko, Phys. Rev. Lett., 91, 034101 (2003).
- [11] A. Carpinteri and F. Mainardi (Eds.), Fractals and Fractional Calculus in Cotinuum Mechanics (Springer, Wien, 1997).
- [12] K. M. Kolwankar and A. D. Gangal, Chaos 6 505 (1996).
- [13] K. M. Kolwankar and A. D. Gangal, Pramana J. Phys. 48 49 (1997).
- [14] K. M. Kolwankar and A. D. Gangal, in the proceedings of the Conference 'Fractals in Engineering', Archanon, 1997.
- [15] S. Krishnamurthy and A. Tanguy and P. Abry and S. Roux, Europhys. Lett., (2000) 51, 1.
- [16] X. Li, M. Davison and C. Essex, On the Concept of Local Fractional Differentiation, Preprint.
- [17] A. Carpinteri and P. Cornetti and K. M. Kolwankar, Chaos, Solitons and Fractals, (2004) 21, 623.
- [18] A. Carpinteri and P. Cornetti, Chaos, Solitons and Fractals, (2002) 13, 85.
- [19] Chen, Y., Yan, Y., Zhang, K., Journal of Mathematical Analysis and Applications (2010) 362, 17-33.
- [20] Wu, Guo-Cheng, Int. Rev. Chem. Eng 4.5 (2012): 505-510.
- [21] X. J. Yang, Advanced Local fractional Calculus and its Applications, 2012.
- [22] F. B. Adda and J. Cresson, J. Math. Anal. Appl., (2001) 263, 721.
- [23] A. Babakhani and V. Daftardar-Gejji, J. Math. Anal. Appl., (2002) 270, 66.
- [24] A. Carpinteri, B. Chiaia and P. Cornetti, Comput. Methods Appl. Mech. Engrg. 191 (2001) 3-19.
- [25] K. M. Kolwankar and A. D. Gangal, Local fractional calculus: A calculus for fractal space-

- time, in *Fractals: Theory and Applications in Engineering*, M. Dekking, J. Lévy Véhel, E. Lutton and C. Tricot, Eds. (Springer, 1999).
- [26] C. P. Li and Z. G. Zhao, Eur. Phys. J. Special Topics 193 5-26 (2011).
- [27] K. M. Kolwankar and J. Lévy Véhel, J. of Fractional Calculus and Applied Analysis, (2001) 4, 285.
- [28] K. M. Kolwankar, Fractals 12.04 (2004): 375-380.
- [29] K. M. Kolwankar and A. D. Gangal, Phys. Rev. Lett. 80, 214 (1998).
- [30] K. M. Kolwankar, Brownian motion of fractal particles: Levy flights from white noise Arxiv preprint cond-mat/0511307, 2005
- [31] A. Carpinteri, B. Chiaia and P. Cornetti, Computers and Structures 82 (2004) 499-508.
- [32] K. M. Kolwankar, Local fractional calculus: a review, in *Fractional Calculus: Theory and Applications*, V. Daftardar-Gejji, Ed. (Narosa, 2013) (arXiv:1307.0739).
- [33] K. M. Kolwankar, Commun. Frac. Cal. 4 105 (2013) (arXiv:1312.7675).